Seasonal and long-term consequences of esca on grapevine stem xylem

integrity

Running title: Impact of the trunk disease esca on stem hydraulic integrity

Highlight: Our study reveals that esca can critically affect xylem water movement in

grapevine perennial organs, by the presence of plant-derived tyloses.

G. Bortolami[a], E. Farolfi[a], E. Badel[b], R. Burlett[c], H. Cochard[b], N. Ferrer[a], A. King[d], L.J.

Lamarque[c,e], P. Lecomte[a], M. Marchesseau-Marchal[a], J. Pouzoulet[f], J.M. Torres-Ruiz[b], S.

Trueba[c,g], S. Delzon[c], G.A. Gambetta[f], C.E.L. Delmas[a*]

[a]INRAE, BSA, ISVV, SAVE, 33882 Villenave d'Ornon, France

[b]Université Clermont-Auvergne, INRAE, PIAF, 63000 Clermont-Ferrand, France

[c]Univ. Bordeaux, INRAE, BIOGECO, 33615 Pessac, France

[d]Synchrotron SOLEIL, L'Orme des Merisiers, Gif-sur-Yvette, 91192, France

[e]Département des Sciences de l'Environnement, Université du Québec à Trois-Rivières,

Trois-Rivières, Québec, G9A 5H7, Canada

[f]EGFV, Bordeaux-Sciences Agro, INRAE, Université de Bordeaux, ISVV, 210 chemin de

Leysotte, 33882 Villenave d'Ornon, France

[g]School of Forestry and Environmental Studies, Yale University, New Haven, CT 06511,

USA

*Author for correspondence

Chloé E. L. DELMAS

Tel: +33557122636

chloe.delmas@inrae.fr

ORCID ID: 0000-0003-3568-605X

Number of tables: 4; Number of figures: 4; Word count: 6337 words

ABSTRACT

Hydraulic failure has been extensively studied during drought-induced plant dieback, but its role in plant-pathogen interactions is under debate. During esca, a grapevine (*Vitis vinifera*) disease, symptomatic leaves are prone to irreversible hydraulic dysfunctions but little is known about the hydraulic integrity of perennial organs over the short- and long-term. We investigated the effects of esca on stem hydraulic integrity in naturally infected plants within a single season and across season(s). We coupled *in vivo* X-ray microtomography visualizations with direct (k_s) and indirect (k_{th}) hydraulic conductivity measurements, and tylose and vascular pathogen detection. Although no loss of hydraulic conductivity (PLC) was observed in asymptomatic stems, 60% of symptomatic stems presented xylem occlusions with subsequent PLC, which could reach critical levels (over 50% PLC). A loss of stem k_s was observed simultaneously, or after, the occurrence of leaf symptoms in the presence of tyloses. The impact of esca on xylem integrity was only seasonal and no long-term impact of disease history was recorded. Our study demonstrated how and to what extent a vascular disease such as esca, affecting xylem integrity, could amplify plant mortality by hydraulic failure.

Key words: Esca, hydraulic failure, plant dieback, tyloses, vascular pathogens, *Vitis vinifera* L., X-ray microCT, xylem anatomy

INTRODUCTION

In agricultural and forest ecosystems, perennial plant dieback causes decreases in plant productivity and longevity (Aleemullah and Walsh, 1996; Eskalen *et al.*, 2013; Urbez-Torres *et al.*, 2013; Alvindia and Gallema, 2017). Plant dieback is a complex process where different biotic and/or abiotic stress factors interact and contribute to leaf and crown wilting and ultimately plant death (Desprez-Lostau *et al.*, 2006; Anderegg *et al.*, 2013, Cailleret *et al.*, 2017; Bettenfeld *et al.*, 2020). Drought-mediated plant dieback has been extensively studied, and in this case hydraulic failure has been identified as the primary cause of plant death (Anderegg *et al.*, 2016). Hydraulic failure results from an interruption of the ascendant water flow by air embolism or xylem occlusion (Zimmermann, 1979; Tyree and Sperry, 1989). Vascular pathogens, which infect the xylem network (Yadeta and Thomma, 2013), are also important drivers of pathogen-mediated plant dieback (Goberville *et al.*, 2016; Pandey *et al.*, 2018; Fallon *et al.*, 2020).

Vascular pathogens induce wood necrosis, leaf symptoms, and crown defoliation (Beckmann and Roberts, 1995; Pearce, 1996). Their biology and toxic metabolite production has been well studied, in particular using controlled phytotoxicity assays (Andolfi *et al.*, 2011; Akpaninyang and Opara, 2017). However, the possible role of hydraulic failure during pathogen-mediated plant dieback has been poorly investigated, and the underlying physiological mechanisms inducing leaf symptoms are not clear yet (Fradin and Thomma, 2006; McDowell *et al.*, 2008). Moreover, the long-term impact (over seasons) and relationships between pathogens, leaf symptom presence, and the hydraulic functioning of the plant are still unknown. During vascular pathogenesis, both air (Pérez-Donoso *et al.*, 2016) and non-gaseous (Sun *et al.*, 2013; Czemmel *et al.,* 2015, Pouzoulet *et al.*, 2019) embolism have been observed. For example, air embolism is thought to accelerate pathogen progression during Pierce's disease (Pérez-Donoso *et al.*, 2016), and non-gaseous embolism is associated with occlusion of the xylem conduits by the plant that could slow the disease process while interfering with xylem water transport (Sun *et al.*, 2013; Pouzoulet *et al.*, 2019).

Xylem occlusion, usually through the production of tyloses and gels, is one of the first plant defense mechanisms against vascular pathogens (Pearce, 1996). Xylem parenchyma cells secrete gels and expand into the vessel lumen, forming tyloses, physically blocking pathogen progression (Zimmermann, 1979). Xylem anatomy plays an important role, both for vascular pathogen development (Martin *et al.*, 2009; Martín *et al.*, 2013; Venturas *et al.*, 2014;

Pouzoulet *et al*., 2017; 2020) and for tylose formation (Bonsen and Kucera, 1990; De Micco
et al., 2016; Pouzoulet *et al*., 2019). If effective, this occlusion mechanism allows the plant to
compartmentalize the infected zone and to generate new tissue around it (CODIT model,
Pearce, 1996). Because tyloses can potentially interfere with the hydraulic functioning of the
plant, they could exacerbate disease symptoms (Talboys, 1972). Tyloses are usually observed
in close proximity to pathogens, as shown in artificial inoculation studies (Czemmel *et al*.,
2015; Rioux *et al*., 2018, among others). However, pathogens frequently proliferate in
perennial organs without physically reaching the leaves, thus leaf symptoms are often
induced at a distance (Beckmann and Roberts, 1995). A recent study shows that tyloses can
be present in symptomatic leaves at a distance from the pathogen niches resulting in
decreased leaf hydraulic conductivity (Bortolami *et al*., 2019).

Over the last decades, grapevine (*Vitis vinifera* L.) mortality and yield loss have been
reported in European, American, and South African vineyards due to esca trunk disease
(Cloete *et al*., 2015; Guerin-Dubrana *et al*., 2019). Esca, a vascular disease caused by the
infection of multiple fungal pathogens, affects mostly mature grapevines (more than seven-
years-old), and symptoms include trunk necrosis and leaf symptoms, consisting of "tiger-
stripe" necrosis and leaf wilting (Lecomte *et al*., 2012; Claverie *et al*., 2020), which are not
regularly expressed season-to-season even within individual vines (Guerin-Dubrana *et al*.,
2013; Li *et al*., 2017). While the pathogens responsible for esca-induced trunk necrosis have
been identified (Morales-Cruz *et al*., 2018; Brown *et al*., 2020), the underlying mechanisms
of leaf and fruit symptoms, and plant death are still poorly understood. Bortolami *et al*.
(2019) demonstrated that the two vascular pathogens related to esca (*Phaeomoniella*
chlamydospora and *Phaeoacremonium minimum*) were never detected in leaves or in one-
year old stems, but always in the trunk (independently from leaf symptom presence). They
further showed that esca symptomatic leaves presented significant losses in hydraulic
conductivity due to the occlusion of the xylem conduits by tyloses. Together, these results
reveal that esca impacts leaf hydraulic functioning, but whether or not there is a
corresponding failure in perennial organs, and the exact timing of this phenomenon, are still
unknown. As stems and branches are the direct connections between the pathogen niche in
the trunk and the observed symptoms in the leaves, the study of stem xylem integrity is
crucial in the understanding of esca impact on grapevine physiology in the current year and
across seasons.

In this study, we investigated stem xylem integrity in grapevine during esca leaf symptom formation asking the following questions: (i) Can esca lead to hydraulic failure in perennial organs? (ii) Does stem hydraulic failure occur prior to or after leaf symptom expression, and does it depend on xylem anatomy? (iii) Do long-term symptomatic plants present different xylem anatomy and levels of hydraulic failure from long-term asymptomatic plants? To answer these questions, we transplanted 28-years-old grapevines (*Vitis vinifera* L. cv Sauvignon blanc) from the field into pots to transport, manipulate, and study naturally esca-infected vines. We coupled *in vivo* visualizations of stem xylem functionality (using synchrotron-based X-ray microcomputed tomography) with stem specific hydraulic conductivity measurements (k_s), theoretical hydraulic conductivity estimates (k_{th}), optical observations of vessel occlusions, and pathogen detection during symptom appearance, while comparing plants with different symptom history record.

MATERIALS AND METHODS

Plant material

Vitis vinifera cv. Sauvignon blanc grafted onto 101-14 MGt were uprooted in winter 2017, 2018, and 2019 from a vineyard planted in 1992 located at INRAE Bordeaux-Nouvelle Aquitaine (44°47'24.8"N, 0°34'35.1"W) and transferred into pots. Each of these plants has been surveyed each year in the field since 2012 for esca leaf symptom expression following Lecomte *et al*. (2012), and has been classified yearly as leaf-symptomatic or asymptomatic. Plants were then classified by their long-term symptomatology record: plants asymptomatic from 2012 to 2018 (pA, previously asymptomatic), and plants that have expressed symptoms at least once between 2012 and 2018 (pS, previously symptomatic). The uprooting method and greenhouse growth conditions are detailed in Bortolami *et al*. (2019).

Esca symptom notation

The evolution of esca leaf symptoms was surveyed twice a week from June to October 2019 on every plant (n=84, Fig. 1). As presented in Fig. 1A, esca symptoms were scored at the stem and whole plant scales. Single stems collected for analyses (both hydraulic measurements or microCT observations) could be noted as: asymptomatic (green leaves and apparently healthy), pre-symptomatic (leaves presenting yellowing or small yellow spots

between the veins), tiger-stripe (typical pattern of esca leaf symptoms), or apoplectic (leaves

passing from green to wilted in a couple of days). Along the experimentation, entire plants

could be noted as asymptomatic (control) or symptomatic (when at least 25% of the canopy

was presenting tiger-stripe leaf symptoms). At the end of the experiment (week 40, October

2019) each plant was classified as symptomatic or asymptomatic (control). We were then

able to group each stem measured into six different groups (Fig. 1A): one group of stems

from control plants (asymptomatic from June to October) and five groups of stems from

symptomatic plants: two before symptom appearance (asymptomatic and pre-symptomatic

stems); and three after symptom appearance (asymptomatic, tiger-stripe, and apoplectic

stems). To clearly differentiate asymptomatic stems collected from symptomatic plants and

asymptomatic stems collected from asymptomatic plants, we considered plants (and their

stems) that didn't show leaf symptoms during the experiment as control plants (or stems). We

investigated whether symptom expression (final symptom notation in October 2019, see Fig.

1) differed between plants with contrasted long-term symptom history (previously

asymptomatic vs previously symptomatic, Table 1) using a Chi-square test of independence.

X-ray microCT observation

Synchrotron-based microCT was used to visualize the content of vessels and their

functionality in esca tiger-stripe and control stems. Three symptomatic plants (presenting

tiger-stripe symptoms for 8, 7, and 3 weeks), and one asymptomatic-control plant were

brought to the PSICHE beamline (King *et al*., 2016) at SOLEIL synchrotron facility in

September 2019. Shoots (ca. 2 m long) were cut under water and transferred into a solution

containing 75mM of contrasting agent iohexol. The iohexol solution absorbs X-rays very

strongly and appears bright white in X-ray scans above the iodine K-edge at 33.2 keV, and,

once it has been taken up by the transpiration stream, the effective functionality of each

vessel can be confirmed (Pratt and Jacobsen, 2018; Bortolami *et al.*, 2019). These stems were

moved and left outdoor to transpire the solution for at least half a day. The stems were then

transferred to the beamline stage and scanned twice in less than 5 minutes using two different

energies of a high-flux (3×10^{11} photons mm^{-2}) monochromatic X-ray beam: 33.1 keV and

33.3 keV. The projections were recorded with a sCMOS camera equipped with a 250-mm-

thick LuAG scintillator (Orca Flash, Hamamatsu, Japan). The complete tomographic scan

included 1500 projections, and each projection lasted 50 ms. Tomographic reconstructions

were performed using PyHST2 software (Mirone *et al*., 2014) using the Paganin method

(Paganin *et al*., 2002), resulting in 32-bit volume reconstructions of 2048 x 2048 x 1024

voxels. The final spatial resolution was 2.8769 μm^3 voxel^{-1}.

Image analysis of microCT scans

The contrast agent iohexol allowed us to distinguish in intact scans the effective functionality

of each vessel. In the absence of iohexol, X-ray microCT scans are used to distinguish air-

filled vessels (appearing black, corresponding to native PLC) from sap-filled vessels

(appearing grey). The addition of iohexol in the xylem sap allows to distinguish the

functional vessels (they appear bright white when they transport the sap), from the non-

functional ones (i.e. occluded vessels remaining grey, corresponding to occlusion PLC). We

could also observe partially occluded vessels (i.e. vessels with simultaneous presence of air

and occlusions, or sap and occlusions). This specific case was observed by checking the

presence of any occlusion in at least 200 slices in each volume. Partially occluded vessels

were considered as occluded, some examples are presented in Fig. S1. The equivalent-circle

diameter of air-filled, occluded, and functional (iohexol-filled) vessels was measured on the

cross sections from the central slice of the microCT scanned volume using ImageJ software

(Schneider et al., 2012). Native PLC, occlusion PLC, and total PLC (i.e. summing occlusion

with native PLC) were calculated with the Hagen-Poiseuille equation, as in Bortolami *et al*.

2019, (see Method S1). We investigated whether native PLC, and occlusion PLC differed

between control and esca tiger-stripe plants, using two independent generalized mixed linear

models where plants were treated as a random effect. Proportional data (ranging from 0 to 1,

dividing all PLC values by 100) was analyzed to fit a logit link function and binomial

distribution as appropriate.

Monitoring stem hydraulic properties over time

Stem xylem integrity was monitored over time by measuring hydraulic properties in control

and symptomatic plants along the season and during esca development. Stem specific

hydraulic conductivity (k_s) was measured by the gravity method (Sperry et al., 1988), and

compared to its theoretical analog (k_{th}) calculated from xylem anatomical observations on the

same stem (see below). When there are observed differences in k_s among stems, comparisons

with theoretical maximums (k_{th}) can show if lower k_s values result from anatomical

differences (i.e. different vessel size distributions) or by hydraulic failure (in the case of

similar vessel size and density). If k_s varies in unity with k_{th}, differences in k_s might result

from anatomical differences (e.g. smaller k_s are related to smaller vessels), otherwise k_s

variations are the consequence of hydraulic failure. Each method to measure k_s, k_{th}, and to

observe tyloses is described below.

Sampling started on June 19[th] and finished on September 13[th] 2019 for a total of 10 sampling

dates, 39 stems from 23 control-asymptomatic plants, and 49 stems from 17 symptomatic

plants. We randomly sampled control plants and esca symptomatic plants all along the season

through the evolution of esca symptoms, obtaining measurements from 14 weeks before until

10 weeks after symptom appearance. To explore the contribution of the experimental design

to data analysis, we tested the effect of the year of uprooting (2018 and 2019), the position of

the analyzed internode, and the week of the measurement (i.e. evolution during the season)

on k_s and k_{th} in control plants using separate generalized linear mixed model with normal

distributions and the plant treated as a random variable (Table S1). A significant impact of

the year of uproot was found for k_s and k_{th} values in control plants (Table S1). This could

have resulted from the more favorable conditions (i.e. climatic stability and nutrient

availability) for the greenhouse grown vines (note that plants uprooted in 2017 were only

esca symptomatic and were not included in this analysis). However, once k_s and k_{th} are

plotted together (Fig. S2), all the values lie on the same regression line without generating

outlier values (smaller k_s values correspond to smaller k_{th} values independently of the

uprooting year).

Stem specific hydraulic conductivity (k_s)

k_s measurements were performed on one internode of >1.5m long one-year old stems,

following Torres-Ruiz *et al.* (2012) gravity method (see Method S2 for details). A flow of 20

mM KCl solution passed through the sample from a reservoir to a precision electronic

balance (AS220.R2, RADWAG, Radom, PL) recording the weight every 5 seconds using the

WinWedge v3 5.0 software (TAL Technologies, Philadelphia, PA, USA). Hydraulic

conductance, k [kg s^{-1} MPa^{-1}] was obtained by the slope generated by the flow and the

corresponding pressure. The linear relationship between flow and pressure obtained were

always characterized by $R^2>0.97$. Stem specific hydraulic conductivity, k_s [kg s^{-1} MPa^{-1} m^{-1}],

was calculated as follows:

$$237 \qquad k_s = (k \text{ x } l) / A$$

$$238 \qquad A = ((d_1/2)^2 \text{ x } \pi) - ((d_2/2)^2 \text{ x } \pi)$$

Where: k is the hydraulic conductance, l is the length of the sample, A is the xylem area, d_1 is

the external diameter of the debarked stem, and d_2 is the diameter of the central pith.

Stem theoretical hydraulic conductivity (k_{th}), vessel anatomy, and tylose observation

Just before hydraulic conductivity measurements, the lower internode was stored at 4 °C in

80% ethanol for analysis of xylem anatomy. When possible, the same internode of k_s

measurements was used for anatomical analysis and k_{th} estimations, otherwise the stored

internode was used for the following protocol. 50 µm thick slices were obtained using a GSL-

1 microtome (Gärtner *et al*., 2014). Slices were stained using a 0.5% safranin solution during

5 minutes, and then washed three to four times in ethanol (100%). They were quickly soaked

in xylene and mounted on microscope slides with Permount Mounting Medium (Electron

Microscopy Science, Hatfield, PA, USA). Images were captured with a stereo microscope

SMZ1270 (Nikon, France) mounted with a DS-Fi3 camera (Nikon, France). The theoretical

conductivity of each vessel (k_{vessel}) [kg m Mpa^{-1} s^{-1}] was calculated using the Hagen-

Poiseuille equation:

$$253 \qquad k_{vessel} = (\pi \text{ x } \varnothing^4 \text{ x } \rho) / (128 \text{ x } \eta)$$

254   Where $\varnothing$ is the equivalent circle diameter [m] (measured with ImageJ software), ρ the density

255   of water [998.2 kg m^{-3} at 20 °C], and η the viscosity of water [1.002 x 10^{-9} MPa s at 20 °C].

256   k_{th} of the stem [kg s^{-1} m^{-1} Mpa^{-1}] was then calculated by summing every k_{vessel} in the xylem

257   area (A) [m²]:

$$258 \qquad k_{th} = \Sigma \, k_{vessel} / A$$

In the entire cross section of each sample, the physical presence (or absence) of tyloses in

vessel lumina was visually assessed.

Regarding the statistical analysis, stems were grouped in six different categories following

their esca symptomatology (as presented in Fig. 1A). We investigated whether k_s, k_{th}, and

total vessel density differed among these different categories, and how k_s, k_{th}, and total vessel

density differed between stems with and without tyloses (independently from leaf symptom

presence), using independent mixed linear general models. The symptom / tylose category
and the year of uprooting (since it had a significant impact on k_s and k_{th} in control plants,
Table S1) were entered as fixed effects, with the plant treated as a random effect since
different stems were sometimes analyzed from the same plant (88 analyzed stems on 40
different plants). Total density and densities for each vessel diameter class were log-
transformed prior to analysis to fit normality requirements. For the classes with no vessels
(e.g. samples without vessel diameters above 160 µm), a minimal density of 0.0001 was
assigned prior to log transformation. We investigated whether the frequency of symptomatic
stems presenting tyloses changed with the symptom age (i.e. weeks between first symptom
detection and k_s measurements on the same plant) with a Chi-square test. The relationships
between stem k_s and k_{th} were tested using linear regression models. Finally, we investigated
whether k_s and k_{th} in control stems differed between plants with different symptom history
records using independent mixed linear general models with the plant treated as a random
effect.

Fungal detection

Detection and quantification of *Phaeomoniella chlamydospora* and *Phaeoacremonium*
minimum were performed using qPCR in a subsample of stems and trunks from the same
symptomatic and control plants used for hydraulic and anatomical measurements. All along
the season, basal internodes, from the same stems sampled for k_s and k_{th} measurements, were
directly placed in liquid nitrogen and stored at -80 °C. At the end of the experiment, a subset
of plants was cut at the base for trunk sampling. A 2 cm high section was cut with a sterilized
hand saw. The bark was removed and the different tissues of each section (necrotic and
apparently healthy wood) were separately collected using ethyl alcohol-sterilized shears in a
sterile environment, and immediately placed in liquid nitrogen. All samples were ground in
liquid nitrogen using a tissue lyser (Tissuelyser II, Qiagen, Germantown, MD, USA). DNA
extraction and qPCR analysis were conducted as described in Pouzoulet *et al.* (2013) using
the primer sets PchQF/R and PalQF/R. Detection and quantification of *P. chlamydospora* and
P. minimum DNA by qPCR (SYBR Green assays) was conducted as described by Pouzoulet
et al. (2017). After qPCR analysis, the results from each trunk sample (i.e. necrotic or
apparently healthy wood) were averaged together in order to obtain one quantification per
plant. Pathogen DNA quantity (average value of three technical replicates, fg/µl) was

normalized by the amount of total DNA (ng/µl), detected using a Qubit fluorometer. We

investigated whether the amount of fungal DNA (both for *P. chlamydospora* and for *P.*

minimum) in trunks differed between symptomatic and control plants, and between control

plants with different symptom history records, using generalized linear mixed model with a

poisson distribution and a log likelihood function.

Statistical analysis

All data management and statistical tests were done in SAS software (SAS 9.4; SAS

Institute). We used PROC GLIMMIX for generalized linear mixed models, PROC GLM for

generalized linear models, PROC REG for regression analyses and PROC FREQ for

frequency analyses (Chi-square test of independence). The normality of the response

variables was tested using a Kolmogorov-Smirnov test (PROC UNIVARIATE) prior to

analyses. Data were log-transformed (total density) or appropriate distributions (binomial,

poisson) were fitted when appropriate.

RESULTS

Esca leaf symptom expression within and across seasons

Esca leaf symptoms were recorded in 20 out of the 58 plants followed in this study (35%,

Fig. 1, Table 1). The number of symptomatic plants increased gradually with time, from the

first symptom appearance in early June to the last in late September (Fig. 1). There was no

effect of the plant history (previously asymptomatic pA, or previously symptomatic pS) on

2019 symptom expression (n=58, X^2=0.27, P=0.60). On 20 pA plants, six (30%) expressed

leaf symptoms in 2019 (Table 1). On 38 pS plants, fourteen (37%) showed symptoms in 2019

(Table 1). However, pS plants expressed symptoms from June to the end of September, while

pA plants showed leaf symptoms only in September.

***In vivo* observations of esca symptomatic stems**

Xylem vessels of control and tiger-stripe stems were observed using three dimensional X-ray
microCT scans in iohexol-fed samples (Fig. 2, 3, Table S2). As shown in Fig. 2, functional
and non-functional vessels can be discriminated through the use of iohexol (functional
vessels appear bright white, non functional vessels appear either black if air-filled or grey if
occluded). We observed almost totally functional stems in all asymptomatic stems (<20%
total PLC, Fig. 2A-C), and 40% of tiger-stripe stems (e.g. Fig. 2D-G). Higher levels of PLC
(>20% total PLC, Fig H-M) were observed in the remaining tiger-stripe stems, with 40% of
tiger-stripe stems exhibiting over 50% total PLC (Fig. 2J-M). When the two components of
PLC were disentangled, we observed that the level of native PLC remained low both in
control (6.5 ± 2.6%) and in tiger-stripe (12.2 ± 2.9%) stems (Fig. 3A). Occlusion PLC values
were virtually zero in control stems (0.7 ± 0.02%) while in tiger-stripe stems the mean
occlusion PLC values was 27.5 ± 8.2% (Fig. 3B). Nevertheless, the variability of occlusion
PLC across tiger-stripe stems was very high, the values ranging from 0.3% to 72.9% (Fig.
2D-M, and 3B), and occlusion PLC was not correlated to symptom age (n=10, $F_{2,7}$=0.19,
P=0.83). Consequently, no statistical differences in native or occlusion PLC were found
between control and tiger-stripe stems (Fig. 3). When higher occlusion PLC was measured
(Fig. 2H-M), occluded vessels could be organized either on one side of the stem (Fig 2J-L) or
randomly distributed across the section (Fig 2H, 2I, 2M). In 90% of symptomatic stems, we
observed that the most external vessels were functional. Occlusions were present equally in
all vessel diameter classes (Fig. S3).

Tylose development, stem specific (k_s) and theoretical (k_{th}) hydraulic conductivity
during esca leaf symptom formation

Tyloses were identified in the xylem vessels of certain tiger-stripe stems and throughout the
temporal development of esca leaf symptoms, from the appearance of symptoms to 11 weeks
after. All apoplectic stems and 62.5% (15 of 24 analyzed stems) of esca tiger-stripe stems
presented tyloses, while all other stems (control, asymptomatic or pre-symptomatic) did not
contain these occlusions, even until one week before symptom development. In esca tiger-
stripe stems, tyloses were not related to specific plants, or to symptom age (i.e. on the same
plant at the same moment, different symptomatic stems could present tyloses, or not, n=24,
X^2=7.47, P=0.38).

Overall, no significant impact of esca symptoms was observed on k_s (Fig. 4A), even if tiger-

stripe stems were divided between those with and without tyloses. Control stems presented a

mean (± SE) k_s of 24.97 ± 1.72 kg s^{-1} MPa^{-1} m^{-1}; all the stems without tyloses measured on

symptomatic plants showed the same range of values as control stems (Fig 4A, Table 2):

26.04 ± 4.71 for asymptomatic before symptoms appearance, 30.32 ± 4.26 for pre-

symptomatic stems, 19.80 ± 5.18 for asymptomatic stems after symptom appearance on the

plant, and 21.29 ± 5.40 for tiger-stripe stems without tyloses. Stems with tyloses (tiger-stripe

and apoplectic stems) presented the lowest average k_s values (11.27 ± 2.86 and 2.47 ± 1.45 kg

s^{-1} MPa^{-1} m^{-1} for tiger-stripe and apoplectic, respectively). Regarding k_{th}, no significant

impact of esca symptoms was found (Fig. 4C, Table 2), all the values were in the same range,

with average values ranging from 70.44 (for tiger-stripe stems with tyloses) to 87.88 (for pre-

symptomatic stems) kg s^{-1} MPa^{-1} m^{-1}.

In order to further investigate the impact of esca on stem hydraulics, we explored the

relationship between individual stem k_s and k_{th} in each symptom category (Fig. 4B, S4, Table

2). Significant relationships were found between k_s and k_{th} in all groups except in

asymptomatic stems after symptom appearance and symptomatic stems with the physical

presence of tyloses (Fig. S4, Table 2). The slopes of regression curves between k_s and k_{th} did

not vary among groups in the absence of tyloses (slope values ranged between 0.3 and 0.4,

Table 2) while it was close to 0 in the presence of tyloses (0.17 for tiger-stripe and 0.04 for

apoplectic stems). When k_s and k_{th} are compared in the presence or absence of tyloses, we

observed that k_s was significantly lower when tyloses were present (9.81 ± 2.51 kg s^{-1} MPa^{-1}

m^{-1} in the presence of tyloses vs 25.06 ± 1.46 kg s^{-1} MPa^{-1} m^{-1} in the absence of tyloses, Table

2, n=88, $F_{1,49}$=7.11, P=0.01) while k_{th} did not significantly differ. Stems without tyloses

presented a strong correlation between k_s and k_{th}, while in the presence of tyloses this

relationship was not significant (Table 2, Fig. 4B).

Total vessel density did not significantly differ between stem symptomatology (comparing all

the seven categories presented in Table 2), even when vessel density was partitioned by

vessel diameter classes (Fig. 4D).

Finally, we tested the impact of disease history (comparing pA and pS plants) on the

hydraulic conductivity and xylem anatomy in control plants. There were no differences

between long-term symptomatic (pS) and long-term asymptomatic (pA) plants in stem k_s,

stem k_{th}, or total vessel density (Table 3).

Fungal detection

The two vascular pathogens associated with esca (*Phaeomoniella chlamydospora* and
Phaeoacremonium minimum) were never detected in one-year old stems while they were
systematically detected in the trunk of both control and symptomatic plants (Table 4). In
trunks, a significantly higher quantity of fungal DNA was detected in tiger-stripe
symptomatic plants than in controls (Table 4). We found 2.14- and 1.64-fold more of *P.*
chlamydospora and *P. minimum* DNA in symptomatic trunks relative to controls. In control
plants, different symptom history record impacted the quantity of fungal DNA detected by
qPCR, for *Phaeomoniella chlamydospora*, and for *Phaeoacremonium minimum*. We found
1.65- and 2.84-fold more *P. chlamydospora* and *P. minimum* DNA in previously
symptomatic trunks relative to previously asymptomatic trunks (Table 3).

DISCUSSION

Our results regarding the xylem integrity during esca show that the presence of plant-derived
tyloses induced hydraulic failure in 60% of symptomatic stems. Tyloses were produced
simultaneously with, or after, the occurrence of leaf symptoms, and resulted in more than
50% PLC in 40% of symptomatic stems, unrelated to symptom age. We demonstrated that
the presence of leaf symptoms during previous seasons had no impact on the likelihood of
symptom appearance in the current year, or on stem hydraulic conductivity and xylem
anatomy. Vascular fungi were never detected in the same organs as the tyloses (one-year old
stems), and although they were present in trunks of both tiger-stripe and control plants, tiger-
stripe plants showed higher quantities of fungal DNA. Among control plants that did not
express symptoms in the year of the study, we found higher quantities of fungal DNA in
those plants with a long-term history of symptom formation. Albeit xylem occlusions were
not observed in the totality of tiger-stripe stems, they could amplify yield loss plant mortality,
especially in the context of climate change as they impair water transport in a majority of
symptomatic stems.

In vivo **xylem integrity observations and hydraulic vulnerability segmentation**

Using direct X-ray microCT imaging in esca symptomatic stems, we found that hydraulic conductivity loss was almost entirely associated with the presence of tyloses. Different studies have investigated the link between vascular pathogen development and hydraulic conductivity in stems (Collins *et al.*, 2009; Lachenbruch and Zhao, 2019; Mensah *et al.*, 2020). During biotic stresses, air embolisms have been shown to decrease hydraulic conductivity during bacterial leaf scorch disease (McElrone *et al.*, 2003; 2008), Pierce's disease (Pérez-Donoso *et al.*, 2016), and Pine wilt disease (Yazaki *et al.*, 2018). In the case of fungal wilt diseases, the hydraulic conductivity loss was associated with non-gaseous embolism (i.e. tyloses) at the point of pathogen inoculation (Guerard *et al.*, 2000; Sallé *et al.*, 2008; Beier *et al.*, 2017; Mensah *et al.*, 2020), or with canker presence in naturally infected stems (Lachenbruch and Zhao, 2019).

Using iohexol we were able to visually observe the exact spatial organization of functional vessels. Interestingly, in some symptomatic samples we found functional vessels surrounding the non-functional xylem (Fig. 2J-L), suggesting that the plant was able to preserve the more external vessels from occlusions or to form new functional vessels after the loss of conductivity. Moreover, the sectoriality of the occlusions observed in Fig. 2J-L was reminiscent of the sectoriality observed in the distributions of trunk necrosis, especially on the brown stripe necrosis appearing along the vasculature (Lecomte *et al.*, 2012).

Comparing these results with our precedent study using the same technique in leaves, we showed that esca symptomatic leaves presented higher levels of occlusion PLC ($60.7 \pm 6.80\%$ in midribs, and $54.02 \pm 8.72\%$ in petioles, data from Bortolami *et al.*, 2019) compared to stems (27.53 ± 8.24 %, occlusion PLC), suggesting hydraulic vulnerability segmentation (although PLC in leaves and stems were measured in different plants and years). The hydraulic segmentation theory relies on the fact that annual organs (i.e. leaves) are more vulnerable than perennial organs (i.e. stems) to drought induced air embolism (Tyree and Ewers, 1991). Grapevine is well known for exhibiting strong hydraulic vulnerability segmentation (Charrier *et al.*, 2016; Hochberg *et al.*, 2016; 2017). This is thought to be adaptive, where the higher vulnerability in leaves and petioles favors embolism formation and leaf shedding prior to embolism formation in stems, thus protecting the perennial organs. Our observations during esca pathogenesis demonstrate that, analogous to the hydraulic vulnerability segmentation theory, leaves appear more vulnerable to the formation of non-

gaseous embolism as well, which could mitigate the risk of hydraulic failure in perennial

organs. From another perspective, the difference may not be a direct effect of the specific

organ's vulnerability to non-gaseous embolism, but a consequence of a difference in the

accumulation of putative toxins and/or elicitors. Indeed, we confirmed here that esca leaf

symptoms occur at a distance from the pathogen niche because vascular pathogens were

never detected in one-year old stems, suggesting that the plant may transport a signal (i.e.

toxins or elicitors) from the infected trunk up to the leaves. If the signal accumulates in leaves

in a higher amount than it does in the stems, and stimulates occlusion formation, stems would

then be secondarily affected.

458   Hydraulic conductivity, tyloses, and vessel anatomy

Tyloses could have different impacts, both positive and negative, during wilt disease

pathogenesis: (i) tyloses contribute to pathogen resistance as they aim to seal off vessel

lumens and impede pathogens spread throughout the host (CODIT model, Pearce, 1996).

This is the case regarding the susceptibility of different species or varieties to specific

pathogens (Jacobi and MacDonald, 1980; Ouellette *et al*., 1999; Clérivet *et al.,* 2000; Et-

Touil *et al*., 2005; Venturas *et al*., 2014; Park and Juzwik 2014; Rioux *et al*., 2018), in

particular to *Phaeomoniella chlamydospora*, one of the pathogen associated with esca

(Pouzoulet *et al*., 2017; 2020). (ii) In other studies, it has been shown that tyloses can

exacerbate symptoms (Talboys, 1972): they cause a reduction in stem hydraulic conductivity,

sometimes associated with a reduction in stomatal conductance in leaves and, in the most

severe cases, wilting (Parke *et al*., 2007; Beier *et al*., 2017; Lachenbruch and Zhao, 2019,

Mensah *et al*., 2020 during fungi development; Sun *et al*., 2013; Deyett *et al*., 2019 during

Pierce's disease). Our results suggest that during esca (i) the development of tyloses in stems

cannot be interpreted as a systematic trait of pathogen resistance because visual symptoms

were observed despite the presence of tyloses and they were produced at a distance from

pathogens. (ii) When present, tyloses might lead to symptom exacerbation. Esca has also

been suggested to lead to a general reduction in xylem water transport and stomatal

conductance (Ouadi *et al*., 2019), and tyloses could be a major contributor to these

phenomena as during winter senescence (Salleo *et al*., 2002; Sun *et al*., 2008). However,

when stems have no tyloses, esca leaf symptom formation seems to arise from within the leaf

itself, and does not result from upstream hydraulic failure.

Xylem is the battleground between vascular pathogens and the plant's defense response

(Yadeta and Thomma, 2013). Even if xylem vessel anatomy is less investigated, it could have

a crucial role in plant resistance and response to vascular pathogens. For example, during

Dutch elm wilt disease (due to *Ophiostoma spp.*) the most sensitive species and varieties

present wider xylem vessels (Elgersma, 1970; Mcnabb *et al.*, 1970; Solla and Gil 2002; Pita

et al., 2018). Smaller vessels could occlude faster, sustaining a more efficient pathogen

restriction (Venturas *et al.*, 2014). Our results on xylem vessel anatomy suggest that stems

with tyloses tend to present higher densities of small vessels, even if we did not observe any

differences in total k_{th} values. Moreover, microCT scans showed that occlusions appear

randomly in every vessel size class (Fig. S3). In contrast, artificial inoculations showed that

xylem vessel diameter had a strong impact on esca-related vascular pathogen development

(Pouzoulet *et al.*, 2017; 2020), and in the kinetic of vessel occlusion in grapevine stems

(Pouzoulet *et al.*, 2019). The relationships between esca leaf symptoms, xylem anatomy, and

tylose presence should be studied in detail in trunks, where vascular pathogens are present,

and among different grapevine varieties and rootstocks as they are known to show different

susceptibility to symptom expression.

497   Long-term consequences of esca on leaf symptom expression and stem hydraulic

498   integrity

In field surveys, esca leaf symptoms are often randomly distributed spatially throughout

vineyards and are not consistent from season to season in individual vines (Mugnai *et al.*,

1999; Surico *et al.*, 2000; Marchi *et al.*, 2006; Guerin-Dubrana *et al.*, 2013; Li *et al.*, 2017).

However, esca-related vine death is strongly related to leaf symptoms as death is usually

observed following a year with symptom expression (Guerin-Dubrana *et al.*, 2013). In

agreement with these field studies, we observed similar percentages of symptomatic plants

between those that had already expressed esca symptoms in the past (from one to seven

consecutive years, pS plants), and those that had never expressed symptoms over the past

seven years (pA plants). However, we also found that pS plants expressed symptoms earlier

in the season than pA plants, suggesting that symptoms might require more time to develop in

pA plants. We did not find any significant differences in k_s and k_{th} values between plants with

contrasted long-term symptom history. This result suggests that esca leaf symptoms may

have xylem anatomical consequences within the year of expression by the production of

tyloses, but not across seasons. Moreover, we showed that DNA pathogen amount

(*Phaeoacremonium minimum* and *Phaeomoniella chlamydospora*) depends on the symptom

expression in the season of sampling, and on the long-term symptom history. Altogether,

these results suggest that a higher amount of vascular fungi in the trunk represents a higher

risk in reproducing leaf symptoms, and consequently, a higher risk of plant death.

Hydraulic failure and esca leaf symptom pathogenesis

Our results showed that, even if esca-related stem occlusion was extremely variable, 40% of

the microCT analyzed stems presented a total PLC greater than 50%. Under drought

conditions alone, studies suggest that grapevines are not able to recover in the current season

from PLC greater than 50% in stems (Charrier *et al*., 2018). Thus, to what extent these levels

of esca-induced hydraulic failure compromise future vine performance, and/or increase the

likelihood of developing esca leaf symptoms in the future remains an open question.

We showed that, similarly to visual leaf symptoms, tyloses in stems were generated at a

distance from the pathogen niche in the trunk. Comparing our results with Bortolami *et al*.

(2019), we show that hydraulic failure affected leaves, on average, two times more than

stems. We could hypothesize that, following pathogen activities in the trunk, a signal passing

through the xylem network and stimulating tyloses, first accumulates in leaves and then

affects the stems. However, the exact signal and action remain unknown, as we showed that

the presence of tyloses depended upon given symptomatic stems rather than symptomatic

plants (i.e. two stems in the same plant, with same tiger-stripe symptoms, sampled at the

same moment, could or could not present tyloses).

We showed that there were no differences in symptom expression, nor in the stem hydraulic

properties, regarding the long-term symptom history. We can conclude that the processes that

generate tiger-stripe symptoms are largely restricted to the current year of the symptom

expression. However, in plants expressing symptoms for the first time according to our

disease record, these processes could require more time, as they showed symptoms only late

in the season. The presence of occlusion, leading to hydraulic failure in stems, could

exacerbate leaf symptom expression in the following seasons, possibly contributing to death.

We could speculate that a stem expressing extensive hydraulic failure could be more prone to

express symptoms in the following year or, in the worst cases, to die. If the level of hydraulic

failure could affect the stem mortality in the following year, the choice of stems with a
complete absence of failure during the winter pruning could reduce the impact of esca in
vineyards. In addition, the presence of occlusions could also amplify plant susceptibility to
drought-induced hydraulic failure, enhancing the risk of plant mortality in the field as
suggested by McDowell et al. (2008). It could be speculated that a decrease in soil water
potential or a high evaporative demand, concomitant to esca-induced hydraulic failure, could
embolize the remaining functional xylem vessels stopping the water flow and desiccating
plant tissues (this could be the case in apoplectic plants for example). In perspective, future
studies should investigate the link between pathogen activities and occlusion development,
especially in trunks, and the subsequent hydraulic failure consequences on whole plant
physiology.

SUPPORTING INFORMATION
The following Supporting Information is available for this article:
Method S1: Image analysis of microCT scans.
Method S2. Specific stem hydraulic conductivity (k_s) measurements.
Fig. S1. Two-dimensional reconstruction of longitudinal cross sections from X-ray microCT
volumes of grapevine stems.
Fig. S2. Relationship between k_s and k_{th} in control plants.
Fig. S3. Vessel density and percentage of occluded vessels in tiger-stripe stems for different
vessel diameter classes.
Fig. S4. Relationships between k_s and k_{th} in each stem symptom category.
Table S1. Effect of year of uprooting, internode analyzed, and sampling date on k_s and k_{th} in
control stems.
Table S2. Calculated theoretical hydraulic conductivity (k_{th} %), and hydraulic conductivity
loss (PLC %) from X-ray microCT volumes of intact grapevine stems.
568

ACKNOWLEDGMENTS
We thank the experimental teams of UMR SAVE and UMR EGFV (Bord'O platform,
INRAE, Bordeaux, France) and the SOLEIL synchrotron facility (beamline PSICHE) for
providing the materials and logistics. Specifically, we thank Jérôme Jolivet and Sebastien
Gambier (UMR SAVE) for providing technical knowledge and support for plant
transplantation and maintenance. This work was supported by the French Ministry of

AUTHOR CONTRIBUTIONS

C.E.L.D., G.A.G., G.B., and S.D. designed the experiments;

E.F., G.A.G., S.D., E.B., R.B., H.C., A.K., L.J.L., J.M.T.-R., S.T. participated in synchrotron

campaigns;

G.B., C.E.L.D., E.F., and N.F. conducted the esca symptom notations;

G.B., M.M.-M.,and N.F. conducted the histological observations;

E.F. conducted the hydraulic conductivity measurements and participated to data analyses;

N.F., and J.P., conducted the pathogen detection;

G.B. analyzed the microCT, optical images, and analyzed the data;

P.L. provided data on disease history of the plants;

G.B., C.E.L.D., and G.A.G. wrote the article;

all authors edited and agreed on the last version of the article.

Data Availability Statement

594 We agree to archive the data associated with this manuscript should the manuscript be

595 accepted.

REFERENCES

Akpaninyang FE, Opara EU. 2017. The Influence of Toxins in Disease Symptom Initiation in Plants: A Review. Journal of Agriculture and Sustainability 10, 29-52.

Aleemullah M., Walsh K. 1996. Australian papaya dieback: Evidence against the calcium deficiency hypothesis and observations on the significance of laticifer autofluorescence. Australian Journal of Agricultural Research 47, 371-385.

Alvindia DG, Gallema FLM. 2017. *Lasiodiplodia theobromae* causes vascular streak dieback (VSD)–like symptoms of cacao in Davao Region, Philippines. Australasian Plant Disease Notes 12, 54.

Anderegg WRL, Kane JM, Anderegg LDL. 2013. Consequences of widespread tree mortality triggered by drought and temperature stress. Nature Climate Change 3, 30-36.

Anderegg WRL, Klein T, Bartlett M, Sack L, Pellegrini AFA, Choat B, Jansen S. 2016. Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. Proceedings of the National Academy of Sciences 113, 5024-5029.

Andolfi A, Mugnai L, Luque J, Surico G, Cimmino A, Evidente A. 2011. Phytotoxins Produced by Fungi Associated with Grapevine Trunk Diseases. Toxins 3, 1569-1605.

Beckman CH, Roberts EM. 1995. On the Nature and Genetic Basis for Resistance and Tolerance to Fungal Wilt Diseases of Plants. Advances in Botanical Research 21, 35-77.

Beier GL, Held BW, Giblin CP, Cavender-Bares J, Blanchette RA. 2017. American elm cultivars: Variation in compartmentalization of infection by *Ophiostoma novo-ulmi* and its effects on hydraulic conductivity. Forest Pathology 47, 1-11.

Bettenfeld P, Fontaine F, Trouvelot S, Fernandez O, Courty P-E. 2020. Woody Plant Declines. What's Wrong with the Microbiome? Trends in Plant Science 25, 381-394.

Bonsen KJM, Kučera LJ. (1990). Vessel Occlusions in Plants: Morphological, Functional and Evolutionary Aspects. IAWA Journal 11, 393-399.

Bortolami G, Gambetta GA, Delzon S, *et al*. (2019). Exploring the Hydraulic Failure Hypothesis of Esca Leaf Symptom Formation. Plant Physiology 181, 1163-1174.

Brown AA, Lawrence DP, Baumgartner K. 2020. Role of basidiomycete fungi in the grapevine trunk disease esca. Plant Pathology 69, 205-220.

Cailleret M, Jansen S, Robert EMR, *et al*. 2017. A synthesis of radial growth patterns preceding tree mortality. Global Change Biology 23, 1675-1690.

Charrier G, Delzon S, Domec J-C, *et al*. 2018. Drought will not leave your glass empty: Low risk of hydraulic failure revealed by long-term drought observations in world's top wine regions. Science Advances 4, 1-9.

Charrier G, Torres-Ruiz JM, Badel E, *et al*. 2016. Evidence for Hydraulic Vulnerability Segmentation and Lack of Xylem Refilling under Tension. Plant Physiology 172, 1657-1668.

Claverie M, Notaro M, Fontaine F, Wery J. 2020. Current knowledge on Grapevine Trunk Diseases with complex etiology□: A systemic approach. Phytopathologia Mediterranea 59, 29-53.

Clérivet A, Déon V, Alami I, Lopez F, Geiger J-P, Nicole M. 2000. Tyloses and gels associated with cellulose accumulation in vessels are responses of plane tree seedlings (*Platanus × acerifolia*) to the vascular fungus *Ceratocystis fimbriata* f. Sp *platani*. Trees 15, 25-31.

Cloete M, Mostert L, Fischer M, Halleen F. 2015. Pathogenicity of South African Hymenochaetales taxa isolated from esca-infected grapevines. Phytopathologia Mediterranea 54, 368-379.

Collins BR, Parke JL, Lachenbruch B, Hansen EM. 2009. The effects of *Phytophthora ramorum* infection on hydraulic conductivity and tylosis formation in tanoak sapwood. Canadian Journal of Forest Research 39, 1766-1776.

Czemmel S, Galarneau ER, Travadon R, McElrone AJ, Cramer GR, Baumgartner K. 2015. Genes expressed in grapevine leaves reveal latent wood infection by the fungal pathogen *Neofusicoccum parvum*. PloS one 10, e0121828.

Desprez-Loustau M-L, Marçais B, Nageleisen L-M, Piou D, Vannini A. 2006. Interactive effects of drought and pathogens in forest trees. Annals of Forest Science 63, 597-612.

Deyett E, Pouzoulet J, Yang J□I, Ashworth VE, Castro C, Roper MC, Rolshausen PE. 2019. Assessment of Pierce's disease susceptibility in *Vitis vinifera* cultivars with different pedigrees. Plant Pathology 68, 1079-1087.

De Micco V, Balzano A, Wheeler EA, Baas P. 2016. Tyloses and gums□: A review of structure, function and occurrence of vessel occlusions. IAWA Journal 37, 186-205.

Elgersma DM. 1970. Length and diameter of xylem vessels as factors in resistance of elms to Ceratocystis ulmi. Netherlands Journal of Plant Pathology 76, 179-182.

Eskalen A, Stouthamer R, Lynch SC, Rugman-Jones PF, Twizeyimana M, Gonzalez A, Thibault T. 2013. Host Range of Fusarium Dieback and Its Ambrosia Beetle (Coleoptera□: Scolytinae) Vector in Southern California. Plant Disease 97, 938-951.

Et-Touil A, Rioux D, Mathieu FM, Bernier L. 2005. External symptoms and histopathological changes following inoculation of elms putatively resistant to Dutch elm disease with genetically close strains of *Ophiostoma*. Canadian Journal of Botany 83, 656–667.

Fallon B, Yang A, Lapadat C, Armour I, Juzwik J, Montgomery RA, Cavender-Bares J. 2020. Spectral differentiation of oak wilt from foliar fungal disease and drought is correlated with physiological changes. Tree Physiology 40, 377-390.

Fradin EF, Thomma BPHJ. 2006. Physiology and molecular aspects of *Verticillium* wilt diseases caused by *V. dahliae* and *V. albo-atrum*. Molecular Plant Pathology 7, 71-86.

Gärtner H, Lucchinetti S, Schweingruber FH. 2014. New perspectives for wood anatomical analysis in dendrosciences: the GSL1-microtome. Dendrochronologia 32, 47-51.

Goberville E, Hautekèete N-C, Kirby RR, Piquot Y, Luczak C, Beaugrand G. 2016. Climate change and the ash dieback crisis. Scientific Reports 6 doi:10.1038/srep35303.

Guérard N, Maillard P, Bréchet C, Lieutier F, Dreyer E. 2007. Do trees use reserve or newly assimilated carbon for their defense reactions? A 13C labeling approach with young

Scots pines inoculated with a bark-beetle-associated fungus (*Ophiostoma brunneo ciliatum*). Annals of Forest Science 64, 601-608.

Guerin-Dubrana L, Labenne A, Labrousse JC, Bastien S, Rey P, Gegout-Petit A. 2013. Statistical analysis of grapevine mortality associated with esca or Eutypa dieback foliar expression. Phytopathologia Mediterranea 52, 276-288.

Guerin-Dubrana L, Fontaine F, Mugnai L. 2019. Grapevine trunk disease in European and Mediterranean vineyards : Occurrence, distribution and associated disease-affecting cultural factors. Phytopathologia Mediterranea 58, 49-71.

Hochberg U, Albuquerque C, Rachmilevitch S, Cochard H, David-Schwartz R, Brodersen CR, McElrone A, Windt CW. 2016. Grapevine petioles are more sensitive to drought induced embolism than stems : Evidence from *in vivo* MRI and microcomputed tomography observations of hydraulic vulnerability segmentation: Hydraulic vulnerability segmentation in grapevine. Plant, Cell & Environment 39, 1886-1894.

Hochberg U, Windt CW, Ponomarenko A, Zhang Y-J, Gersony J, Rockwell FE, Holbrook NM. 2017. Stomatal Closure, Basal Leaf Embolism, and Shedding Protect the Hydraulic Integrity of Grape Stems. Plant Physiology 174, 764-775.

Jacobi WR, MacDonald WL. 1980. Colonization of resistant and susceptible oaks by *Ceratocystis fagacearum*. Phytopathology 70, 618-623.

King A, Guignot N, Zerbino P, *et al*. 2016. Tomography and imaging at the PSICHE beam line of the SOLEIL synchrotron. Review of Scientific Instruments 87, 093704.

Lachenbruch B, Zhao J-P. 2019. Effects of phloem on canopy dieback, tested with manipulations and a canker pathogen in the *Corylus avellana/Anisogramma anomala* host/pathogen system. Tree Physiology 39, 1086-1098.

Lecomte P, Darrieutort G, Liminana J-M, Comont G, Muruamendiaraz A, Legorburu F-J, Choueiri E, Jreijiri F, El Amil R, Fermaud M. 2012. New Insights into Esca of Grapevine: The Development of Foliar Symptoms and Their Association with Xylem Discoloration. Plant Disease 96, 924-934.

Li S, Bonneu F, Chadoeuf J, Picart D, Gégout-Petit A, Guérin-Dubrana L. 2017. Spatial and Temporal Pattern Analyses of Esca Grapevine Disease in Vineyards in France. Phytopathology 107, 59-69.

Martin N, Vesentini D, Rego C, Monteiro S, Oliveira H, Ferreira RB. 2009. *Phaeomoniella chlamydospora* infection induces changes in phenolic compounds content in *Vitis vinifera*. Phytopathologia Mediterranea 48, 101-116.

Martín JA, Solla A, Ruiz-Villar M, Gil L. 2013. Vessel length and conductivity of *Ulmus* branches : Ontogenetic changes and relation to resistance to Dutch elm disease. Trees 27, 1239-1248.

McDowell N, Pockman WT, Allen CD, *et al*. 2008. Mechanisms of plant survival and mortality during drought : Why do some plants survive while others succumb to drought? New Phytologist 178, 719-739.

McElrone AJ, Sherald JL, Forseth IN. 2003. Interactive effects of water stress and xylem-limited bacterial infection on the water relations of a host vine. Journal of Experimental Botany 54, 419-430.

McElrone AJ, Jackson S, Habdas P. 2008. Hydraulic disruption and passive migration by a bacterial pathogen in oak tree xylem. Journal of Experimental Botany 59, 2649-2657.

McNabb HS, Heybroek HM, Macdonald WL. 1970. Anatomical factors in resistance to Dutch elm disease. Netherlands Journal of Plant Pathology 76, 196-205.

Mensah JK, Sayer MAS, Nadel RL, Matusick G, Eckhardt LG. 2020. Physiological response of *Pinus taeda* L. trees to stem inoculation with *Leptographium terebrantis*. Trees 34, 869–880.

Mirone A, Brun E, Gouillart E, Tafforeau P, Kieffer J. 2014. The PyHST2 hybrid distributed code for high speed tomographic reconstruction with iterative reconstruction and a priori knowledge capabilities. Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms 324, 41-48.

Morales-Cruz A, Allenbeck G, Figueroa-Balderas R, Ashworth VE, Lawrence DP, Travadon R, Smith RJ, Baumgartner K, Rolshausen PE, Cantu D. 2018. Closed-

reference metatranscriptomics enables *in planta* profiling of putative virulence activities in the grapevine trunk disease complex: Transcriptomics of pathogen communities. Molecular Plant Pathology 19, 490-503.

Mugnai L, Graniti A, Surico G. 1999. Esca (Black Measles) and Brown Wood-Streaking : Two Old and Elusive Diseases of Grapevines. Plant Disease 83, 404-418.

Ouadi L, Bruez E, Bastien S, Vallance J, Lecomte P, Domec J-C, Rey P. 2019. Ecophysiological impacts of Esca, a devastating grapevine trunk disease, on *Vitis vinifera* L. PloS one 14, 1-20.

Ouellette GB, Baayen RP, Simard M, Rioux D. 1999. Ultrastructural and cytochemical study of colonization of xylem vessel elements of susceptible and resistant *Dianthus caryophyllus* by *Fusarium oxysporum* f.sp. Dianthi. Canadian Journal of Botany 77, 644-663.

Paganin D, Mayo SC, Gureyev TE, Miller PR, Wilkins SW. 2002. Simultaneous phase and amplitude extraction from a single defocused image of a homogeneous object. Journal of Microscopy 206, 33-40.

Pandey S, Rishi RR, Jayaraj R, Giri K, Kumar R, Pandey A, Juwantha R, Madaan S, Bhandari MS. 2019. *Fusarium equiseti* is associated with the wilt and dieback of *Aquilaria malaccensis* in Northeast India. Forest Pathology 49, e12489.

Park J-H, Juzwik J. 2014. *Ceratocystis smalleyi* colonization of bitternut hickory and host responses in the xylem. Forest Pathology 44, 282-292.

Parke JL, Oh E, Voelker S, Hansen EM, Buckles G, Lachenbruch B. 2007. *Phytophthora ramorum* Colonizes Tanoak Xylem and Is Associated with Reduced Stem Water Transport. Phytopathology 97, 1558-1567.

Pearce RB. 1996. Antimicrobial defences in the wood of living trees. New Phytologist 132, 203-233.

Pérez-Donoso AG, Lenhof JJ, Pinney K, Labavitch JM. 2016. Vessel embolism and tyloses in early stages of Pierce's disease. Australian Journal of Grape and Wine Research 22, 81-86.

Pita P, Rodríguez-Calcerrada J, Medel D, Gil L. 2018. Further insights into the components of resistance to *Ophiostoma novo-ulmi* in *Ulmus minor*: Hydraulic conductance, stomatal sensitivity and bark dehydration. Tree Physiology 38, 252-262.

Pouzoulet J, Mailhac N, Couderc C, Besson X, Daydé J, Lummerzheim M, Jacques A. 2013. A method to detect and quantify *Phaeomoniella chlamydospora* and *Phaeoacremonium aleophilum* DNA in grapevine-wood samples. Applied Microbiology and Biotechnology 97, 10163-10175.

Pouzoulet J, Scudiero E, Schiavon M, Rolshausen PE. 2017. Xylem Vessel Diameter Affects the Compartmentalization of the Vascular Pathogen *Phaeomoniella chlamydospora* in Grapevine. Frontiers in Plant Science 8, 1-13.

Pouzoulet J, Scudiero E, Schiavon M, Santiago LS, Rolshausen PE. 2019. Modeling of xylem vessel occlusion in grapevine. Tree Physiology 39, 1438-1445.

Pouzoulet J, Rolshausen PE, Charbois R, Chen J, Guillaumie S, Ollat N, Gambetta GA, Delmas CEL. 2020. Behind the Curtain of the Compartmentalization Process: Exploring How Xylem Vessel Diameter Impacts Vascular Pathogen Resistance. Plant, Cell & Environment doi: 10.1111/pce.13848.

Pratt RB, Jacobsen AL. 2018. Identifying which conduits are moving water in woody plants: A new HRCT-based method. Tree Physiology 38, 1200-1212.

Rioux D, Blais M, Nadeau-Thibodeau N, Lagacé M, DesRochers P, Klimaszewska K, Bernier L. 2018. First Extensive Microscopic Study of Butternut Defense Mechanisms Following Inoculation with the Canker Pathogen *Ophiognomonia clavigignenti-juglandacearum* Reveals Compartmentalization of Tissue Damage. Phytopathology 108, 1237-1252.

Salle A, Ye H, Yart A, Lieutier F. 2008. Seasonal water stress and the resistance of *Pinus yunnanensis* to a bark-beetle-associated fungus. Tree Physiology 28, 679-687.

Salleo S, Nardini A, Lo Gullo MA, Ghirardelli LA. 2002. Changes in stem and leaf hydraulics preceding leaf shedding in *Castanea sativa* L. Biologia plantarum 45, 227-234.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. Nature Methods 9, 671-675.

Sperry JS, Donnelly JR, Tyree MT. 1988. A method for measuring hydraulic conductivity and embolism in xylem. Plant, Cell & Environment 11, 35-40.

Solla A, Gil L. 2002. Xylem vessel diameter as a factor in resistance of *Ulmus minor* to *Ophiostoma novo-ulmi*. Forest Pathology 32, 123-134.

Sun Q, Rost TL, Matthews MA. 2008. Wound-induced vascular occlusions in *Vitis vinifera* (Vitaceae): Tyloses in summer and gels in winter. American Journal of Botany 95, 1498-1505.

Sun Q, Sun Y, Walker MA, Labavitch JM. 2013. Vascular Occlusions in Grapevines with Pierce's Disease Make Disease Symptom Development Worse. Plant Physiology 161, 1529-1541.

Surico G, Marchi G, Ferrandino A, Braccini P, Mugnai L. 2000. Analysis of the spatial spread of esca in some Tuscan vineyards (Italy). Phytopathologia Mediterranea 39, 211-224.

Talboys PW. 1972. Resistance to Vascular Wilt Fungi. Proceedings of the Royal Society of London. Series B, Biological Science 181, 319-332.

Torres-Ruiz JM, Sperry JS, Fernández JE. 2012. Improving xylem hydraulic conductivity measurements by correcting the error caused by passive water uptake. Physiologia Plantarum 146, 129-135.

Tyree MT, Sperry JS. 1989. Vulnerability of Xylem to Cavitation and Embolism. Annual Review of Plant Physiology and Plant Molecular Biology 40, 19-38.

Tyree MT, Ewers FW. 1991. The hydraulic architecture of trees and other woody plants. New Phytologist 119, 345-360.

Úrbez-Torres JR, Peduto F, Vossen PM, Krueger WH, Gubler WD. 2013. Olive Twig and Branch Dieback: Etiology, Incidence, and Distribution in California. Plant Disease 97, 231-244.

Venturas M, López R, Martín JA, Gascó A, Gil L. 2014. Heritability of *Ulmus minor* resistance to Dutch elm disease and its relationship to vessel size, but not to xylem vulnerability to drought. Plant Pathology 63, 500-509.

Yadeta KA, Thomma BPHJ. 2013. The xylem as battleground for plant hosts and vascular wilt pathogens. Frontiers in Plant Science 4 doi:10.3389/fpls.2013.00097.

Yazaki K, Takanashi T, Kanzaki N, Komatsu M, Levia DF, Kabeya D, Tobita H, Kitao M, Ishida A. 2018. Pine wilt disease causes cavitation around the resin canals and irrecoverable xylem conduit dysfunction. Journal of Experimental Botany 69, 589-602.

Zimmermann MH. 1979. The Discovery of Tylose Formation by a Viennese lady in 1845. IAWA Bulletin 2, 51-56.

TABLES

Table 1. Esca leaf symptom observations over the experimental season on *Vitis vinifera* cv Sauvignon blanc.

Symptom notation before 2019	All plants	Previously asymptomatic (pA)	Previously symptomatic (pS)
Symptom notation in 2019			
Esca-symptomatic	35 % (20/58)	30 % (6/20)	37 % (14/38)
Control-asymptomatic	65 % (38/58)	70 % (14/20)	63 % (24/38)

Plants are grouped by their symptom history: previously asymptomatic (pA, plants that have never expressed leaf symptoms between 2012 and 2018) and previously symptomatic (pS, plants that have expressed leaf symptoms at least once since 2012). Ratios present the number of plants in each symptom category (esca-symptomatic or control-asymptomatic) over the total number of plants of the category.

Table 2. Values for specific stem hydraulic conductivity (k_s), theoretical stem hydraulic conductivity (k_{th}) and equations of regression lines between k_s and k_{th} for control and esca symptomatic stems.

Tyloses	Esca	k_s [kg s^{-1} m^{-1} Mpa^{-1}]	k_{th} [kg s^{-1} m^{-1} Mpa^{-1}]	n (stem - plant)	Regression
Absence	Control	24.97 ± 1.72	78.36 ± 4.51	39 - 23	$k_s = 0.3$ x $k_{th} + 1.6$ R²=0.61 P<0.0001
	Asymptomatic before symptoms	26.04 ± 4.71	74.75 ± 11.80	6 - 6	$k_s = 0.36$ x $k_{th} - 0.94$ R²=0.82 P=0.013
	Pre-symptomatic	30.32 ± 4.26	87.88 ± 8.54	11 - 7	$k_s = 0.37$ x $k_{th} - 1.9$ R²=0.54 P=0.010
	Asymptomatic after symptoms	19.80 ± 5.18	72.58 ± 12.64	5 - 2	$k_s = 0.33$ x $k_{th} - 4$ R²=0.64 P=0.104
	Esca (tiger-stripe)	21.29 ± 5.40	72.85 ± 10.41	9 - 5	$k_s = 0.45$ x $k_{th} - 11.26$ R²=0.74 P=0.003
Presence	Esca (tiger-stripe)	11.27 ± 2.86	70.44 ± 7.81	15 - 5	$k_s = 0.17$ x $k_{th} -0.84$ R²=0.22 P=0.077
	Esca (apoplectic)	2.47 ± 1.45	74.80 ± 33.48	3 - 2	$k_s = 0.04$ x $k_{th} - 0.2$ R²=0.68 P=0.385
Absence	All	25.06 ± 1.46	78.42 ± 3.37	70 - 37	$k_s = 0.34$ x $k_{th} - 1.90$ R²=0.63 P<0.0001
Presence	All	9.81 ± 2.51	71.16 ± 8.00	18 - 7	$k_s = 0.12$ x $k_{th} - 1.28$ R²=0.15 P=0.117

Values represent mean ± SE. n = sample size, (including the number of analyzed stems and number of analyzed plants, respectively). See text and Fig. 4 for statistical analysis. A detailed esca symptom notation is provided in Fig. 1A. Bivariate plots of each regression are presented in Fig. S4.

Table 3. Long-term impact of symptom presence (i.e. comparing plants with different disease history record) in control plants on specific stem hydraulic conductivity (k_s), theoretical stem hydraulic conductivity (k_{th}), stem total vessel density, and amount of *Phaeomoniella chlamydospora* and *Phaeoacremonium minimum* DNA in trunks.

	Previously asymptomatic (pA)	Previously asymptomatic (pS)	Type III Tests of Fixed Effects (pA vs pS)
k_s [kg s^{-1} m^{-1} Mpa^{-1}]	23.76 ± 2.30	26.54 ± 2.61	n=39, $F_{1,16}$=1.19, P=0.29
k_{th} [kg s^{-1} m^{-1} Mpa^{-1}]	72.22 ± 4.85	86.30 ± 7.98	n=39, $F_{1,16}$=3.01, P=0.10
total vessel density [count mm^{-2}]	57.28 ± 4.03	52.61 ± 3.25	n=39, $F_{1,16}$=0.72, P=0.41
P. chlamydospora [pg ng^{-1}]	6.14 ± 1.90	10.15 ± 3.41	**n=13, $F_{1,11}$=5900.06, P<0.0001**
P. minimum [pg ng^{-1}]	9.27 ± 6.97	26.40 ± 13.83	**n=13, $F_{1,11}$=51014, P<0.0001**

Values represent means $\pm$ SE. Pathogen quantification was estimated as: pg fungal DNA ng^{-1} total DNA. Statistical tests used are individual generalized linear mixed models to compare pA vs pS plants (fixed effect) with the individual plants entered as a random effect in the models and the year of uprooting as a co-variable (fixed effect). Statistically significant results (P<0.05) are shown in bold.

Table 4. Quantification by qPCR of *Phaeomoniella chlamydospora* and *Phaeoacremonium minimum* DNA in stems and trunks of different esca symptomatology.

Organ	Esca	n	*P. chlamydospora* [pg ng^{-1}]	*P. minimum* [pg ng^{-1}]
Stem	Control	8	0	0
	Pre-symptomatic	3	0	0
	Asymptomatic (after symptoms)	3	0	0
	Tiger-stripe (without tyloses)	4	0	0
	Tiger-stripe (with tyloses)	8	0	0
	Apoplectic	2	0	0
Trunk	Control	13	7.37 ± 1.67 (12/13)[*]	14.54 ± 6.51 (12/13)[*]
	Symptomatic	7	15.80 ± 3.12 (7/7)[*]	23.90 ± 8.82 (7/7)[*]

[*] Number of samples positive for the pathogen over the total number of analyzed samples.

Pathogen quantification was estimated as: pg fungal DNA per ng total DNA. Values represent means $\pm$ SE, n = sample size. Trunks of symptomatic plants presented higher amount of both *P. chlamydospora* and *P. minimum*, compared to control (n=20, $F_{1,18}$=29806.11.25, P<0.0001 and n=20, $F_{1,18}$=21925.4, P<0.0001, respectively). See text for statistical methods.

FIGURE LEGENDS

Fig. 1. Representation of esca symptom notation during the experimental season. (**A**) Single stems could be noted as esca asymptomatic, pre-symptomatic, tiger-stripe, or apoplectic. Whole plants have been noted as control (asymptomatic from June to October) or symptomatic (with tiger-stripe symptoms at the end of the season). (**B**) Proportion of plants in each symptom category over the experimental season (n=58). The blue area corresponds to control plants, green area to esca symptomatic plants before symptom appearance, and red area to esca symptomatic plants.

Fig. 2. Two-dimensional reconstruction of cross-sections from X-ray microCT volumes of grapevine stems. Each panel represents a cross section of different stems for control (**A-C**) and esca symptomatic (**D-M**) plants. Iohexol appears white bright in functional vessels; air-filled vessels (i.e. native PLC) appear black; occluded vessels (i.e. occlusion PLC) appear grey. Total PLC (i.e. native PLC plus occlusion PLC) values are given for the presented samples. Scale bars = 1000μm.

Fig. 3. (**A**) Mean values of native PLC in control (blue) and esca tiger-stripe (red) stems of grapevine plants using X-ray microCT imaging. Differences were not significant (n=13, $F_{1,9}$=0.07, P=0.79). (**B**) Mean values of occlusion PLC in control (blue) and esca tiger-stripe (red) stems of grapevine plants using X-ray microCT imaging. Differences were not significant (n=13, $F_{1,9}$=0.33, P=0.58). Boxes and bars show the median, quartiles and extreme values, circles show mean values. N represents the sample size (number of analyzed stems) for each group.

Fig. 4. Relationships between specific stem hydraulic conductivity (k_s), theoretical stem hydraulic conductivity (k_{th}), and vessel density in control and esca symptomatic grapevine plants. (**A**) k_s values for control (blue); asymptomatic (dark green) and pre-symptomatic (yellow) stems in plants before symptom appearance; asymptomatic (light green), tiger-stripe (red), and apoplectic (grey) stems in plants after symptom appearance, differences were not significant (n=88, $F_{5,45}$=1.30 P=0.28). Boxes and bars show the median, quartiles and extreme values, circles within boxes correspond to means, and circles outside boxes to outlier values. (**B**) Relationships between k_s and k_{th}. Symbols represent the absence (circles) or presence

(triangles) of tyloses in xylem vessels. Colors represent esca symptomatology (as in panel A). The dashed line represents the regression for stems in which no tyloses were observed in xylem vessels, and the solid line represents the regression for samples with tyloses. R^2 for the regression lines are indicated (see Table 2 and Fig. S4 for detailed analyses). (**C**) k_{th} values for the different stem categories as presented in panel a. Differences were not significant (n=88, $F_{5,45}$=0.58, P=0.71). (**D**) Relationships between mean values of xylem vessel density and their diameters. Differences in total vessel density and in vessel size distributions were not significant (n=88, $F_{6,45}$=0.77, P=0.60; n=792 (88 samples for 9 vessel classes), $F_{48,693}$=1.19, P=0.18). Colors and markers are the same as panel B.

A

Esca symptom notation

Stem

Asymptomatic   Asymptomatic   Asymptomatic
                    Pre-symptomatic   Tiger-stripe
                                            Apoplectic

Plant

Control   Before symptoms   After symptoms

Asymptomatic   Symptomatic

B