Nutrient Export and Periphyton Biomass in a Stream-Lake Basin from the Patagonian Andean Region

Alejandro Sosnovsky[1], Mailén Elizabeth Lallement[1], Magalí Rechencq[1], María Valeria Fernández[1], Eduardo Enrique Zattara[1] & Claudia Silvina Feijoó[2]

[1]Grupo de Evaluación y Manejo de Recursos Ícticos, INIBIOMA, CONICET-Universidad Nacional del Comahue, Quintral 1250, R8400FRF San Carlos de Bariloche, Argentina

[2]Programa Biogeoquímica de Ecosistemas Dulceacuíclas, INEDES-CONICET, Departamento de Ciencias Básicas, Universidad Nacional de Luján, Ruta Nacional N° 5 y Av. Constitución B6700ZBA Luján, Argentina

Corresponding author: Alejandro Sosnovsky

e-mail: alejandro.sosnovsky@comahue-conicet.gob.ar

Telephone number: +54 294 423374 (452), Fax number: +54-2944-422111

Nutrient Export and Periphyton Biomass in a Stream-Lake Basin from the Patagonian Andean Region

Abstract: We characterized land use, dissolved nutrients, and periphytic biomass in Patagonian Andean streams. Nutrient export, especially dissolved nitrogen increased with human population density. However, no correlation between nutrient concentration and algal biomass was found, which could instead be limited by riparian vegetation cover. Our results suggest that local N-limited ecosystems are liable to eutrophication by increased demographic pressure and that alternative wastewater treatment strategies are necessary for sustainable growth.

Keywords: Ecosystem Functioning; Water Quality; Eutrophication; Baseflow; Spatial studies

Rainfall water infiltrates or runs off drainage basins until it reaches a stream. Physicochemical characteristics of stream water thus reflect the results of the interactions between precipitated water and the basin it flows through. For example, a stream's N:P ratio depends on the flow paths of each individual watershed (Green *et al.*, 2007). Eutrophication, the enrichment of surface waters with nutrients, is a key human-induced stressor in streams and often a consequence of intensified land use (Dodds & Smith, 2016). Stream eutrophication results in higher export and concentration of phosphorus (P) and nitrogen (N), usually followed by increased algal biomass and Cyanobacteria blooms (Álvarez *et al.*, 2017). Since algal intracellular nutrient concentrations and stream nutrient concentration are directly linked, ambient N:P ratio can indicate if any of these elements acts as a limiting factor to primary productivity of the ecosystem. A review of field studies found that the optimal N:P molar ratio of freshwater benthic algae is 18:1, suggesting that higher ratios indicate P limitation, while lower ratios indicate N limitation (Kahlert, 1998). Thus, N:P ratios are a simple metric to evaluate the impact of land use changes in a basin on stream ecosystems.

Ecological characteristics of the Northwest Patagonian region are driven by the Andean range. This region has a temperate-cold climate with marked seasonality in precipitation (Paruelo *et al.*, 1998). Its lakes and forests are generally N limited (Díaz *et al.*, 2007; Diehl *et al.*, 2008), while its Andisols-type soils are characterized by high

capacity to retain P (Satti *et al.*, 2003). The region harbours a wide network of stream

ecosystems subject to a yearly cycle formed by three contrasting hydrological periods:

stormflow, meltflow and baseflow (Sosnovsky *et al.*, 2020). Although large part of the

region has been suffering increasing impact from human activities related to settlement

since the early 20[th] century, studies linking this impact with eutrophication-driving

nutrients like N and P are still scarce (Baffico, 2001; Miserendino *et al.*, 2011;

Sosnovsky *et al.*, 2019). We expect that more intensive land use will translate into

higher nutrient export and concentration of nutrients in fluvial ecosystems, as well as an

increased periphyton biomass. Thus, we aimed to determine, for a stream-lake basin

during the baseflow period, the existing relationships i) between human population

density and dissolved nutrient export (i.e., N and P) and ii) between nutrient

concentration and periphyton algal biomass.

Research was conducted in a suburban area within the city of San Carlos de

Bariloche, Argentina. This city has approximately 110 000 inhabitants, distributed in

several drainage basins. Stream Gutiérrez drainage basin *(41° 09' 36.18" S 71° 24'*

37.19"W) is large (162 km^2) and includes a large and deep lake (17 km^2, 111 m

maximum depth) (Fig. 1). The lake has 7 inflow perennial streams; its seasonal average

concentrations of soluble reactive phosphorus (Srp) and dissolved inorganic nitrogen

(Din) are 3.4 mg m^{-3} and 4.6 mg m^{-3}, respectively (Díaz *et al.*, 2007). Gutiérrez stream,

the outflow of the lake, flows 9 km through a wide gently sloped valley and has only a

perennial affluent, the Cascada stream. Gutiérrez's riverbanks are extensively colonized

by the exotic crack willow, *Salix fragilis*, and pass through a series of populated areas.

Most settlements in the basin are not connected to the city's sewer system, using instead

household sanitation systems to treat and dispose domestic wastewater. Newer or more

affluent households use septic tanks and nitrifying gravel beds, while older ones usually

employ cesspits. Thus, sewage input is directly related to local population density. In

addition, a trout farm with a total production of ~10 tons year^{-1} pours its sewage into

Gutiérrez stream.

Fig. 1. a) Gutiérrez stream basin with the sampled sub-basins highlighted. Sampling sites at tributaries and mainstem of Gutiérrez stream are shown: 1) Torrontegui stream, 2) Melgarejo stream, 3) Pescadero stream, 4) Loncochinoco stream, 5) Cascada stream. b) 6-10) Gutiérrez stream. Population zones are shown in grey. c) Export of nutrients and concentration of Chlorophyll a (Chl a). The vertical dotted line represents the lake, to its left are its tributary streams, to its right are the Gutiérrez stream and its tributary. The Y axes corresponding to dissolved nitrogen (NO_3 + NO_2 + NH_4) and Chl a has two segments for a better visualization of the data.

We surveyed water quality along the Gutiérrez stream basin in April 2016. Six points

along the stream, the Cascada tributary stream plus four other tributary streams flowing

into the headwater lake were sampled (Table 1, Fig. 1). We determined the specific

drainage area for each site and used demographic density data (inhabitants· km^{-2}) from

2010's National Census. We also estimated canopy cover from riparian vegetation using

a concave gridded mirror; and measured discharge (Q), electrical conductivity (CE),

temperature (T) and turbidity. We filtered 0.5 liters of stream water through a MGF

Munktell filter to determine the concentration of dissolved nutrients, Din and soluble

inorganic phosphorus. We calculated Din as the sum of ammonium (NH_4), nitrates

(NO_3) and nitrites (NO_2). We analysed NH_4 by the phenol method and $NO_3 + NO_2$ with

a FUTURA Autoanalyzer (Alliance Instruments,Frepillon, France) through a reaction

with sulfanilamide with a previous Cu-Cd-reduction for NO_3 (A.P.H.A., 2005). We

considered soluble inorganic phosphorus as Srp (A.P.H.A., 2005). The export nutrients

were calculated by the following formula:

$$Nutrient\ export = \frac{[N]\ Q\ t}{A}$$

Where $[N]$ is nutrient concentration, Q is discharge, t is time and A is drainage basin

area, and export was standardized to kg of nutrient per km^2 and year.

We also determined chlorophyll *a* (Chl *a*) concentration, corrected for

pheopigments (Lorenzen, 1967), as a proxy for periphyton algal biomass. We choose

two representative boulders and sampled periphyton from each one of the boulders,

using a piece of cylindrical plastic pipe measuring 16 cm^2 in cross-section and a brush to

standardize the sampled area and remove the algae. We used linear regressions to

explore the relationship between i) population density and nutrient export, and ii)

nutrient concentration and Chl *a* concentration. Variables for which data failed tests for

normal distribution were log-transformed.

Number/Site	Drainage basin (km^2)	Discharge (m^3 s^{-1})	EC (µS cm^{-1})	Temperature (°c)	Turbidity (NTU)	Nutrient concentration (mg m^{-3})				N:P (molar)	Canopy (%)
						Nh$_4$	NO$_2$ + NO$_3$	Din	Srp		
1) Str. Torrontegui	17.0	0.158	92	6.7	0.32	17.1	0.9	18.0	3.7	8.1	7
2) Str. Melgarejo	2.6	0.016	198	8.1	1.47	nd	1.1	1.1	6.5	0.2	0
3) Str. Pescadero	11.3	0.006	74	9.6	0.41	nd	1.7	1.7	4.6	0.2	66
4) Str. Loncochinoco	9.4	0.099	146	9.3	1.14	4.4	488.2	492.6	12.0	21.0	0
5) Str. Cascada	13.0	0.003	87	7.6	0.07	nd	27.9	27.9	8.3	1.7	15
6) Gutiérrez rise	133.1	1.051	70	14.8	0.17	nd	0.5	0.5	3.7	0.1	3
7) Gutiérrez	144.2	1.132	76	12.6	0.24	1.0	14.8	15.8	7.4	1.3	0
8) Gutiérrez	158.3	0.231	88	11.5	0.83	30.2	78.3	108.5	12.0	7.6	100
9) Gutiérrez	160.9	0.938	80	11.7	0.32	12.0	326.0	338.1	11.0	16.6	44
10) Gutiérrez mouth	161.6	0.937	80	11.7	0.35	5.3	452.3	457.6	5.5	42.4	0

Table 1 Morphometric and limnological characteristics of the streams (Str.). Electrical Conductivity (EC); Ammonium (Nh$_4$); Nitrates + Nitrites (NO$_3$ + NO$_2$); Soluble reactive phosphorus (Srp); Dissolved inorganic nitrogen (Din), Phosphorus:Nitrogen ratio (N:P). Nd: not detectable

Tributary streams to Gutiérrez lake and stream show a highly heterogeneous distribution

of variable values (Fig. 1a, Table 1, supplementary table ST1). These streams had

considerably lower temperature and discharge values relative to those found along

stream Gutiérrez. In contrast, their nutrient concentrations were higher than those

measured at the stream's headwater (site 6) (Table 1). Torrontegui (site 1) and

Melgarejo (site 2) streams, draining sparsely populated basins, showed the lowest Chl *a*

concentrations (Fig. 1a). Despite the low nutrient export at Gutiérrez stream headwater,

we detected an increasing trend in Din export and N:P ratio towards the stream mouth

(Fig. 1b-c, Table 1). This trend was mainly driven by increased $NO_3 + NO_2$ and not

NH_4; ammonium export was highest at the two sites immediately downstream of the

trout farm. This increase in Din was not accompanied by increases in either Srp exports

or Chl *a* concentration (Fig. 1c, Table 1). Lack of response to increased nitrogen load

could be due to light limitations; for example, sites 8 and 9, which showed high Din

values and low Chl *a* concentration were shaded by a very dense riparian canopy cover

(Table 1).

Integrating our results across the whole study basin, we observe that the stream

ecosystem seems to be responding to an increase in land use, given that Din and Srp

exports were higher at sites draining more densely populated areas (Fig. 2a-b). This

relationship was more noticeable for Din: the linear regression between Din and

population density was significantly higher ($P = 0.002$. compared to $P = 0.018$ for Srp)

and explained a larger amount of variance ($R^2 = 0.72$, compared to 0.52 for Srp).

Similarly, the increase in population density was reflected in an increase in the N:P

ratio, which surpasses 18: 1 once population reaches ~ 35 inhabitants per km^2 (Fig. 2c).

This potential effect of land use on nutrient load was not followed by a response from

periphyton (Din-Chl *a* R^2=0.02, n=10, F=0.17, *P*=0.69; Srp-Chl *a* R^2=0.01, n=10,

F=0.05, *P*=0.82).

Fig. 2. Linear regressions Human density - Nutrients export. a) Dissolved inorganic nitrogen export (Din_{exp}) b) Soluble reactive phosphorus export (Srp_{exp}) c) Nitrogen:Phosphorus ratio in a molar basis. Dotted line indicates N:P = 18:1.

Our results reflect a relationship between population density and exports of dissolved

nitrogen and phosphorus. On the one hand, the steepest response shown by Din values

indicates that nitrogen was more readily incorporated into the stream ecosystem. This

may be partially explained by leeching of nitrates from household-associated nitrifying

beds (Beal *et al.*, 2005; De & Toor, 2017) and cesspits towards groundwater. In contrast

to this pervasive and diffuse source of nitrates to the stream, ammonium increase is

better explained by single point sources, likely excretion from fishes at the trout farm.

On the other hand, septic chambers and Andisol soils seem to be better at retaining

phosphorus derived from human activities. Thus, the specific interactions between

human settlement, waste treatment strategies and natural environmental characteristics

in this area skew nutrient export ratios towards nitrogen enrichment. In mountain areas

as the study region, this enrichment becomes most relevant during the baseflow period,

when groundwater flow path makes the largest contribution to total stream discharge

(Ahearn *et al.*, 2004).

The range of ambient N:P ratio has been used to define the transition between N

and P limitation in benthic algae (Schanz & Juon, 1983). If the N:P ratio is greater than

20 then P can be assumed to be in limiting supply. If N:P is less than 10 then N can be

assumed to be in limiting supply. The distinction of the limiting nutrient when ambient

N:P ratios are between 10 and 20 is not precise. Since the range of N:P values recorded

on our study site spans a wide range, lack of response from periphyton to changes in

N:P ratio might be due to different sites being subject to different constraints to algal

growth. For example, stream ecosystems found within a forest matrix can be limited by

light availability (Hill *et al.*, 2001); indeed, we found that sites with low Chl *a*

concentration despite high nutrient loads were usually shadier. Thus, we can postulate

that in stream ecosystems as the one we studied, the response of primary producers to

increased nutrient availability is modulated by the nature and structure of the riparian

vegetation. Recently, a significant fraction of the invasive *S. fragilis* population has

been removed presenting a great opportunity to experimentally test this hypothesis; and

more generally, to measure the impact of riparian vegetation on stream productivity.

This work presents some points which deserve further study. NH_4 concentration

in Torrontegui stream was unusually high and deserves further investigation. Moreover,

since we have made our measurements only during the baseflow period, we still do not

know if the same patterns relating population density and nutrient exports hold for

stormflow and meltflow periods of the local hydrologic cycle. Furthermore, our

understanding of the relationships between periphytic algae and nutrient availability

would benefit from studying species and chemical composition of this primary

producers (Carrillo *et al.*, 2018). Finally, the study of the environmental C:N:P

relationships will give us a better comprehension of the auto-heterotrophic dynamics of

Andean fluvial ecosystems of Patagonia.

Our research yields insights into stream eutrophication, as it links land uses with

physicochemical characteristics of a N-limited drainage basin. The headwater lake acted

as an efficient nutrient trap, while residential land use caused increased nutrient release.

We can conclude that in the Northwest Patagonian region, household wastewater

treatment strategies as those currently used within Gutiérrez stream basin are likely

contributing to eutrophication of streams during the baseflow period. Thus, it would be

necessary to consider and evaluate the use of alternative wastewater treatment systems

capable of reducing the release of nitrogen into groundwater. Since this basin is part of a

touristic region that can seasonally increment its population density to almost twice its

year-round density, devising mitigation and remediation measurements for human

impact is imperative to achieve environmental sustainability.

Acknowledgements

192 We acknowledge technician Pablo Alvear for field assistance. Sosnovsky, A.,

193 Rechencq, M., Férnández, M.V. and Zattara, E.E. are CONICET researchers.

References

A.P.H.A. 2005. *Standard Methods for the Examination of Water and Wastewater*.
American Public Health Association. Washington DC.
AHEARN, D. S., R. W. SHEIBLEY, R. A. DAHLGREN & K. E. KELLER. 2004.
Temporal dynamics of stream water chemistry in the last free-flowing river
draining the western Sierra Nevada, California. *Journal of Hydrology,* 295: 47-
63. doi:10.1016/j.jhydrol.2004.02.016
ÁLVAREZ, X., E. VALERO, R. M. B. SANTOS, S. G. P. VARANDAS, L. F.
SANCHES FERNANDES & F. A. L. PACHECO. 2017. Anthropogenic
nutrients and eutrophication in multiple land use watersheds: Best management
practices and policies for the protection of water resources. *Land Use Policy,* 69:
1-11. doi.org/10.1016/j.landusepol.2017.08.028
BAFFICO, G. 2001. Variations in the periphytic community structure and dynamics of
Lake Nahuel Huapi (Patagonia, Argentina). *Hydrobiologia,* 455 (1-3): 79-85.
doi:10.1023/A:1011991402794
BEAL, C. D., E. A. GARDNER & N. W. MENZIES. 2005. Process, performance, and
pollution potential: A review of septic tank-soil absorption systems. *Australian*
Journal of Soil Research, 43: 781-802.
CARRILLO, U., V. DÍAZ VILLANUEVA & B. E. MODENUTTI. 2018. Sustained
effects of volcanic ash on biofilm stoichiometry, enzyme activity and
community composition in North-Patagonia streams. *Science of the Total*
Environment, 621: 235-244. doi.org/10.1016/j.scitotenv.2017.11.270
DE, M. & G. S. TOOR. 2017. Nitrogen transformation in the mounded drainfields of
drip dispersal and gravel trench septic systems. *Ecological Engineering,* 102:
352-360.
DÍAZ, M. M., F. L. PEDROZO, C. S. REYNOLDS & P. F. TEMPORETTI. 2007.
Chemical composition and the nitrogen-regulated trophic state of Patagonian
lakes. *Limnologica,* 37: 17-27. doi:10.1016/j.limno.2006.08.006
DIEHL, P., M. J. MAZZARINO & S. FONTENLA. 2008. Plant limiting nutrients in
Andean-Patagonian woody species: effects of interannual rainfall variation, soil
fertility and mycorrhizal infection. *Forest Ecolgoy and Management,* 225 (7):
2973-2980. doi.org/10.1016/j.foreco.2008.02.003

DODDS, W. K. & V. H. SMITH. 2016. Nitrogen, phosphorus, and eutrophication in
streams. *Inland Waters,* 6: 155-164. DOI: 10.5268/IW-6.2.909
GREEN, M. B., J. L. NIEBER, G. JOHNSON, J. MAGNER & B. SCHAEFER. 2007.
Flow path influence on an N:P ratio in two headwater streams: a paired
watershed study. *Journal of Geophysical Research,* 112.
doi:10.1029/2007JG000403
HILL, W. R., P. J. MULHOLLAND & E. R. MARZOLF. 2001. Stream ecosystem
response to forest leaf emergence in spring. *Ecology,* 82: 2306-2319.
KAHLERT, M. 1998. C:N:P ratios of freshwater benthic algae. *Archiv für*
Hydrobiologie, Special Issues, Advances in Limnology, 51: 105-114.
LORENZEN, C. J. 1967. Determination of chlorophyll and pheopigments:
spectrophotometric equations. *Limnology and Oceanography,* 12: 343-346.
MISERENDINO, M. L., R. CASAUX, M. ARCHANGELSKY, Y. C. DI PRINZIO, C.
BRAND & M. A. KUTSCHKER. 2011. Assessing land-use effects on water
quality, in-stream habitat, riparian ecosystems and biodiversity in Patagonian
northwest streams. *Science of the Total Environment,* 409: 612-624.
Doi:10.1016/j.scitotenv.2010.10.034
PARUELO, J. M., A. BELTRAN, E. JOBBÁGY, O. SALA & R. GOLLUSCIO. 1998.
The climate of Patagonia: general patterns and controls on biotic processes.
Ecología Austral, 8 (2): 85-101.
SATTI, P., M. J. MAZZARINO & M. GOBBI. 2003. Soil N dynamics in relation to
leaf litter quality and soil fertility in north-western Patagonian forests. *Journal of*
Ecology, 91: 173-181.
SCHANZ, F. & H. JUON. 1983. Two different methods of evaluating nutrient
limitantions of periphyton bioassays, using water from the River Rhine and
eights of its tributaries. *Hydrobiologia,* 102: 187-195.
https://doi.org/10.1007/BF00006346
SOSNOVSKY, A., M. E. LALLEMENT, M. RECHENCQ, E. E. ZATTARA, M. V.
FERNÁNDEZ, R. DAGA, M. J. SUAREZ, S. LEIVA & R. J. C. CANTET.
2019. Topography and Land Use Modulate Hydrological and Nutrient Export
Dynamics in two Andean Streams. *Bioriv.* doi: https://doi.org/10.1101/742411
SOSNOVSKY, A., M. RECHENCQ, V. FERNÁNDEZ, M. J. SUAREZ & R. J. C.
CANTET. 2020. Hydrological and Physico-chemical dynamics in two Andean
streams. *Limnetica,* 39 (1): 17-33. DOI: 10.23818/limn.39.02
262
263

Number/Site	Population (inhabitants km^{-2})	Nutrient export (kg km^{-2} y^{-1})				Chlorophyll *a* (mg m^{-2})
		Nh_4	NO_2 + NO_3	Din	Srp	
1) Str. Torrontegui	0	5.0	0.3	5.3	1.08	7
2) Str. Melgarejo	0	nd	0.2	0.2	1.27	7
3) Str. Pescadero	4	nd	nd	nd	0.08	117
4) Str. Loncochinoco	45	1.5	162.2	163.7	3.97	52
5) Str. Cascada	13	nd	0.2	0.2	0.06	77
6) Gutiérrez rise	7	nd	0.1	0.1	0.92	7
7) Gutiérrez 7	19	0.3	3.7	4.0	1.83	66
8) Gutiérrez 8	18	1.4	3.6	5.0	0.55	12
9) Gutiérrez 9	23	2.2	59.9	62.1	2.03	4
10) Gutiérrez mouth	28	1.0	83.1	84.1	1.02	29

Table 1 annexed Data supporting Figures 1 and 2. Population density and limnological characteristics of the study sites (Str.) Streams. Ammonium (Nh_4); Nitrates + Nitrites (NO_3 + NO_2); Soluble reactive phosphorus (Srp). Nd: not detectable.