Nutrient Export and Periphyton Biomass in a Stream-Lake Basin from

the Patagonian Andean Region

Alejandro Sosnovsky[1], Mailén Elizabeth Lallement[1], Magalí Rechencq[1], María Valeria

Fernández[1], Eduardo Enrique Zattara[1] & Claudia Silvina Feijoó[2]

[1]Grupo de Evaluación y Manejo de Recursos Ícticos, INIBIOMA, CONICET-

Universidad Nacional del Comahue, Quintral 1250, R8400FRF San Carlos de

Bariloche, Argentina

[2]Programa Biogeoquímica de Ecosistemas Dulceacuícolas, INEDES-CONICET,

Departamento de Ciencias Básicas, Universidad Nacional de Luján, Ruta Nacional N° 5

y Av. Constitución B6700ZBA Luján, Argentina

Corresponding author: Alejandro Sosnovsky

e-mail: alejandro.sosnovsky@comahue-conicet.gob.ar

Telephone number: +54 294 423374 (452), Fax number: +54-2944-422111

Nutrient Export and Periphyton Biomass in a Stream-Lake Basin from

the Patagonian Andean Region

Abstract: We characterized how land use influenced dissolved nutrients and periphytic

algal biomass in an Andean basin from Northwest Patagonia. Nutrient export, especially

dissolved inorganic nitrogen increased with human population density. However, no

correlation between nutrient concentration and algal biomass was found, which could

instead be limited by light availability. Our results suggest that local N-limited

ecosystems are liable to eutrophication by increased demographic pressure and that

alternative wastewater treatment strategies are necessary for sustainable growth.

Keywords: Ecosystem Functioning; Water Quality; Eutrophication; Baseflow period;

Spatial studies, On-site wastewater treatment systems

Physicochemical characteristics of a stream reflect the interactions between water and

the basin it flows through. For example, a stream's N:P ratio depends on water flow

paths of each individual watershed (Green *et al*., 2007). Eutrophication, the enrichment

of surface waters with nutrients, is a key human-induced stressor in streams and often a

consequence of intensified land use (Dodds & Smith, 2016). Stream eutrophication

results in higher export and concentration of phosphorus (P) and nitrogen (N), usually

followed by increased algal biomass and Cyanobacteria blooms (Álvarez *et al.*, 2017).

Since algal intracellular nutrient concentration and stream nutrient concentration are

directly linked, ambient N:P ratio indicate which of these elements could act as a

limiting factor to primary productivity of the ecosystem. A review of field studies found

that the optimal C:N:P molar ratio of freshwater benthic algae is 158:18:1 (Kahlert,

1998), suggesting that P is limiting when environmental concentration of N:P exceed

18:1, and N is limiting when ratio is < 18:1. Thus, N:P ratio is a simple metric to

evaluate the impact of land use changes on nutrient limitation in stream ecosystems.

The ecological characteristics of Northwest Patagonia region are driven by the

Andean range. This region has a temperate-cold climate with markedly seasonal

precipitation regimes (Paruelo *et al.*, 1998). Its lakes and forests are generally N-limited

ecosystems (Díaz *et al.*, 2007; Diehl *et al.*, 2008), while its Andisols-type soils are

characterized by high capacity to retain P (Satti *et al.*, 2003). The region harbours a

wide network of stream ecosystems subject to a yearly cycle formed by three
contrasting hydrological periods: i) stormflow, when chemistry is controlled by lateral
flow through the landscape, ii) meltflow, when chemistry is largely influenced by
melting snow in the uplands and iii) baseflow, when chemistry is controlled by
groundwater inflow (Ahearn *et al.*, 2004; Sosnovsky *et al.*, 2020). Although a
considerable fraction of the region has been under an increasing influence from human
activities since urban settlement started by the early 1900's, it is startling that studies
linking this influence to eutrophication-driving nutrients like N and P are still scarce
(Baffico, 2001; Miserendino *et al.*, 2011; Sosnovsky *et al.*, 2019). We hypothesize that
more intensive land use will translate into higher nutrient export and concentration of
nutrients in fluvial ecosystems, resulting in increased periphyton biomass on an
otherwise nutrient-limited ecosystem. Thus, we aimed to determine during the baseflow
period if i) dissolved inorganic nutrient export (i.e., N and P) varied with human
population density and ii) changes in nutrient concentration were reflected by changes
in periphytic algal biomass.

Research was conducted in a suburban area within the city of San Carlos de
Bariloche, Northwest Patagonia, Argentina. This city has approximately 110 000
inhabitants distributed across several drainage basins. One of them, Gutiérrez stream's
drainage basin *(41°09'36.18"S 71°24'37.19"W)*, is large (162 km^2) and includes a large
and deep lake (Gutiérrez Lake, 17 km^2, 111 m maximum depth) (Fig. 1). Settlements in
this basin are not connected to the city's main sewer system, using instead economic on-
site wastewater treatment systems (OWTS) to treat and dispose domestic wastewater.
Thus, we can assume that OWTS densities are directly related to local population
density within Gutiérrez's basin. Wastewater undergoes primary treatment in a septic
tank via sedimentation and anaerobic digestion in OWTS. Secondary treatment of the
septic tank effluents occurs within a rubble bed. This bed has a large aerobic zone,
favouring nitrification processes over denitrification ones. The lake of this basin has 7
inflow perennial streams; its seasonal average concentrations of soluble reactive
phosphorus (SRP) and dissolved inorganic nitrogen (DIN) are 3.4 mg m^{-3} and 4.6 mg m^{-3}
^{3}, respectively (Díaz *et al.*, 2007). Gutiérrez stream, the outflow of this lake, flows 9 km
through a wide gently sloped valley and has a single perennial affluent, the Cascada
stream. Gutiérrez's riparian zone is extensively colonized by the exotic crack willow,
Salix fragilis, and runs through a series of populated areas. In addition, a trout farm with
a total production of ~10 tons year^{-1} pours its sewage into Gutiérrez stream.

Fig. 1. a) Gutiérrez stream basin with the sampled sub-basins highlighted. Sampling sites at tributaries and mainstem of Gutiérrez stream are shown: 1) Torrontegui stream; 2) Melgarejo stream; 3) Pescadero stream; 4) Loncochinoco stream; 5) Cascada stream. b) 6-10) Gutiérrez stream. Population zones are shown in grey. c) Export of nutrients and concentration of Chlorophyll *a* (Chl *a*). The vertical dotted line represents the lake, to its left are its tributary streams, to its right are the Gutiérrez stream and its tributary. The Y axes corresponding to dissolved inorganic nitrogen ($NO_3 + NO_2 + NH_4$) and Chl *a* have two segments for a better visualization of the data.

We surveyed water quality along the Gutiérrez stream basin during year 2016's

baseflow period. Demographic density (inhabitants· km^{-2}) was calculated from 2010's

National Census data (https://www.indec.gob.ar). Water samples were taken at four of

the headwater lake's tributary streams (1-4), at the Cascada stream (5) and at five points

along the main stream (6-10; Table 1, Fig. 1), measured *in-situ* discharge (Q), electrical

conductivity (CE), temperature (T) and turbidity, and estimated canopy cover from

riparian vegetation using a concave gridded mirror. We filtered 0.5 liters of water

through a MGF Munktell filter to determine the concentration of DIN and SRP. We

calculated DIN as the sum of ammonium (NH_4), nitrates (NO_3) and nitrites (NO_2). We

analysed NH_4 by the phenol method, $NO_3 + NO_2$ with a FUTURA Autoanalyzer

(Alliance Instruments, Frepillon, France) through a reaction with sulfanilamide with a

previous Cu-Cd-reduction for NO_3 and SRP by the ascorbic acid method (A.P.H.A.,

2005). The export nutrients were calculated by the following formula:

$$Nutrient\ export = \frac{[N]\ Q\ t}{A}$$

Where $[N]$ is nutrient concentration, Q is discharge, t is time and A is drainage basin

area; nutrient export was standardized to kg of nutrient per km^2 and year ($kg\ km^{-2}\ yr^{-1}$).

We also determined chlorophyll *a* (Chl *a*) concentration, corrected for

pheopigments (Lorenzen, 1967), as a proxy for periphytic algal biomass. For periphyton

sampling, two representative boulders were chosen and scraped, using a piece of

cylindrical plastic pipe measuring $16\ cm^2$ in cross-section and a brush to standardize the

sampled area and remove the algae. Relationships between i) population density and

nutrient export, and ii) nutrient concentration and Chl *a* concentration were modelled

using standard linear regression, log-transforming variables that failed tests for normal

distribution.

Number/Site	Drainage basin (km^2)	Discharge (m^3 s^{-1})	EC (µS cm^{-1})	Temperature (°c)	Turbidity (NTU)	Nutrient concentration (mg m^{-3})				N:P (molar)	Canopy (%)
						Nh$_4$	NO$_2$ + NO$_3$	DIN	SRP		
1) Str. Torrontegui	17.0	0.158	92	6.7	0.32	17.1	0.9	18.0	3.7	8.1	7
2) Str. Melgarejo	2.6	0.016	198	8.1	1.47	nd	1.1	1.1	6.5	0.2	0
3) Str. Pescadero	11.3	0.006	74	9.6	0.41	nd	1.7	1.7	4.6	0.2	66
4) Str. Loncochinoco	9.4	0.099	146	9.3	1.14	4.4	488.2	492.6	12.0	21.0	0
5) Str. Cascada	13.0	0.003	87	7.6	0.07	nd	27.9	27.9	8.3	1.7	15
6) Gutiérrez rise	133.1	1.051	70	14.8	0.17	nd	0.5	0.5	3.7	0.1	3
7) Gutiérrez	144.2	1.132	76	12.6	0.24	1.0	14.8	15.8	7.4	1.3	0
8) Gutiérrez	158.3	0.231	88	11.5	0.83	30.2	78.3	108.5	12.0	7.6	100
9) Gutiérrez	160.9	0.938	80	11.7	0.32	12.0	326.0	338.1	11.0	16.6	44
10) Gutiérrez mouth	161.6	0.937	80	11.7	0.35	5.3	452.3	457.6	5.5	42.4	0

Table 1 Morphometric and limnological characteristics of the streams (Str.). Electrical Conductivity (EC); Ammonium (Nh$_4$); Nitrates + Nitrites (NO$_3$ + NO$_2$); Soluble reactive phosphorus (SRP); Dissolved inorganic nitrogen (DIN), Phosphorus:Nitrogen ratio (N:P). Nd: not detectable

Tributary streams to Gutiérrez Lake and stream show a highly heterogeneous distribution of

variable values (Fig. 1a, Table 1, supplementary table ST1). These streams had

considerably lower temperature and discharge values relative to those found along stream

Gutiérrez. In contrast, their nutrient concentrations were higher than those measured at the

stream's headwater (site 6) (Table 1). Torrontegui (site 1) and Melgarejo (site 2) streams,

draining sparsely populated basins, showed the lowest concentration of $NO_3 + NO_2$ and Chl

a (Fig. 1a, Table 1). Despite the low nutrient export at Gutiérrez stream headwater, we

detected an increasing trend in DIN export and N:P ratio towards the stream mouth (Fig.

1c, Table 1). This trend was mainly driven by increased $NO_3 + NO_2$ and not NH_4;

ammonium export was highest at the two sites immediately downstream of the trout farm.

The increase in DIN in stream G was not accompanied by increases in either SRP exports

or Chl *a* concentration (Fig. 1c, Table 1). Instead, Chl *a* concentration was low in sites

shaded by a dense riparian cover (sites 8 and 9, Figure 1c, Table 1).

Integrating our results across the whole study basin, we observed that the stream

ecosystem responded to an increase in land use intensity, given that DIN and SRP exports

were higher at sites draining more densely populated areas (Fig. 2a-b). This relationship

was more noticeable for DIN: the linear regression between DIN and population density

was significantly higher ($P = 0.002$. compared to $P = 0.018$ for SRP) and explained a larger

amount of variance ($R^2 = 0.72$, compared to $R^2 = 0.52$ for SRP). Similarly, the increase in

population density was reflected in an increase in the N:P ratio, which surpasses 18: 1 once

population reaches ~ 35 inhabitants per km^2 (Fig. 2c). This potential effect of land use on

nutrient load was not followed by a response from periphytic algal biomass (DIN-Chl *a* R^2

$= 0.02$, n $= 10$, F $= 0.17$, $P = 0.69$; SRP-Chl *a* $R^2 = 0.01$, n $= 10$, F $= 0.05$, $P = 0.82$).

Fig. 2. Linear regressions Human density - Nutrients export. a) Dissolved inorganic nitrogen export (DIN$_{exp}$) b) Soluble reactive phosphorus export (SRP$_{exp}$) c) Nitrogen:Phosphorus ratio in a molar basis. Dotted line indicates N:P = 18:1.

Our results reflect a positive relationship between population density and exports of DIN

and SRP. On the one hand, the steepest response shown by DIN values indicates that

nitrogen was more readily incorporated into the stream ecosystem. This may be partially

explained by leeching of nitrates from OWTS towards groundwater (Beal *et al.*, 2005; De

& Toor, 2017). In contrast to this pervasive and diffuse source of nitrates to the stream,

ammonium increase is better explained by single point sources, likely excretion from fishes

at the trout farm. On the other hand, OWTS and Andisol soils seem to be better at retaining

phosphorus derived from human activities. Thus, the specific interactions between human

settlement, waste treatment strategies and natural environmental characteristics in this area

skew nutrient export ratios towards nitrogen enrichment. In mountain areas as the study

region, this enrichment becomes most relevant during the baseflow, when groundwater

flow path makes the largest contribution to total stream discharge (Ahearn *et al.*, 2004).

The range of ambient N:P ratio has been used to define the transition between N and

P limitation in benthic algae (Schanz & Juon, 1983). If the N:P ratio is greater than 20 then

P can be assumed to be in limiting supply. If N:P is less than 10 then N can be assumed to

be in limiting supply. The distinction of the limiting nutrient when ambient N:P ratios are

between 10 and 20 is not precise. The N:P ratio recorded on our study spanned a wide

range, from <1 to 42. Thus, the observed lack of response from periphytic algal biomass to

the increase in nutrient concentration could be because the limiting nutrient differed

between sites, being phosphorous, nitrogen or a colimitation by both them. However, N:P

ratio as an indicator of nutrient limitation should be viewed with caution because nutrient

requirements for different algal species can vary widely (Allan & Castillo, 2007). In

addition, stream ecosystems found within a forest matrix can be limited by light availability

(Hill *et al.*, 2001); indeed, we found that sites with low Chl *a* concentration despite high

nutrient loads were usually shadier. Thus, we can postulate that in stream ecosystems as the

one we studied, the response of primary producers to increased nutrient availability is

modulated by the nature and structure of the riparian vegetation. Recently, a significant

fraction of the invasive *S. fragilis* population has been removed presenting a great

opportunity to test this hypothesis; and more generally, to measure the impact of riparian

vegetation on stream productivity.

This work presents some points which deserve further study. NH_4 concentration in

Torrontegui stream was unusually high and merits deeper investigation. Moreover,

measurements were limited to the baseflow period, which might have different dynamics

relating population density and nutrient exports than the stormflow and meltflow periods of

the local hydrologic cycle. Furthermore, our understanding about how periphytic algae

respond to nutrient availability would benefit from studying species and chemical

composition of these primary producers (Carrillo *et al.*, 2018). Finally, the study of the

environmental C:N:P relationships will give us a better comprehension of the auto-

heterotrophic dynamics of Andean fluvial ecosystems of Patagonia.

Our research yields insights into stream eutrophication, as it links land uses with

physicochemical characteristics of a N-limited drainage basin. The headwater lake acted as

an efficient nutrient trap, while residential land use caused increased nutrient load to the
basin. Whether this nutrient increase results in algal growth is modulated by the degree of
canopy shading. We can conclude that in the Northwest Patagonian region, OWTS as those
currently used within Gutiérrez stream basin are likely contributing to eutrophication of
streams during the baseflow period. Thus, it would be necessary to consider and evaluate
the use of alternative OWTS, with higher denitrification capacity, in order to reduce release
of nitrates into groundwater. Furthermore, inhibition of algal response to nitrogen increases
by canopy shading could allow build-up of nutrient concentrations to go unnoticed,
resulting in potentially harmful blooms if the canopy is removed due to changes in land
use. Thus, special care should be taken in preserving the riparian zone of these streams,
especially since nitrate concentration of the groundwater might be reduced by
denitrification in this zone (Hill, 1996). Given that the Gutiérrez basin is part of a touristic
region that can seasonally double its population density, devising mitigation and
remediation measurements for human impact is imperative to achieve environmental
sustainability.
198

Acknowledgements

200 We acknowledge technician Pablo Alvear for field assistance. Sosnovsky, A., Rechencq,
201 M., Férnández, M.V. and Zattara, E.E. are CONICET researchers. This work was funded
202 by Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET resol. N°
203 3646/13).

References

206
A.P.H.A. 2005. *Standard Methods for the Examination of Water and Wastewater.*
American Public Health Association. Washington DC.
AHEARN, D. S., R. W. SHEIBLEY, R. A. DAHLGREN & K. E. KELLER. 2004.
Temporal dynamics of stream water chemistry in the last free-flowing river draining
the western Sierra Nevada, California. *Journal of Hydrology,* 295: 47-63.
DOI:10.1016/j.jhydrol.2004.02.016

ÁLVAREZ, X., E. VALERO, R. M. B. SANTOS, S. G. P. VARANDAS, L. F. SANCHES
FERNANDES & F. A. L. PACHECO. 2017. Anthropogenic nutrients and
eutrophication in multiple land use watersheds: Best management practices and
policies for the protection of water resources. *Land Use Policy,* 69: 1-11.
DOI.org/10.1016/j.landusepol.2017.08.028
ALLAN, J. D. & M. M. CASTILLO. 2007. *Stream Ecology. Structure and Function of*
Running Waters. Springer. Dordrecht, The Netherlands.
BAFFICO, G. 2001. Variations in the periphytic community structure and dynamics of
Lake Nahuel Huapi (Patagonia, Argentina). *Hydrobiologia,* 455 (1-3): 79-85.
DOI:10.1023/A:1011991402794
BEAL, C. D., E. A. GARDNER & N. W. MENZIES. 2005. Process, performance, and
pollution potential: A review of septic tank-soil absorption systems. *Australian*
Journal of Soil Research, 43: 781-802. DOI.org/10.1071/SR05018
CARRILLO, U., V. DÍAZ VILLANUEVA & B. E. MODENUTTI. 2018. Sustained effects
of volcanic ash on biofilm stoichiometry, enzyme activity and community
composition in North-Patagonia streams. *Science of the Total Environment,* 621:
235-244. DOI.org/10.1016/j.scitotenv.2017.11.270
DE, M. & G. S. TOOR. 2017. Nitrogen transformation in the mounded drainfields of drip
dispersal and gravel trench septic systems. *Ecological Engineering,* 102: 352-360.
DOI.org/10.1016/j.ecoleng.2017.02.039
DÍAZ, M. M., F. L. PEDROZO, C. S. REYNOLDS & P. F. TEMPORETTI. 2007.
Chemical composition and the nitrogen-regulated trophic state of Patagonian lakes.
Limnologica, 37: 17-27. DOI:10.1016/j.limno.2006.08.006
DIEHL, P., M. J. MAZZARINO & S. FONTENLA. 2008. Plant limiting nutrients in
Andean-Patagonian woody species: effects of interannual rainfall variation, soil
fertility and mycorrhizal infection. *Forest Ecolgoy and Management,* 225 (7): 2973-
2980. DOI.org/10.1016/j.foreco.2008.02.003
DODDS, W. K. & V. H. SMITH. 2016. Nitrogen, phosphorus, and eutrophication in
streams. *Inland Waters,* 6: 155-164. DOI: 10.5268/IW-6.2.909

GREEN, M. B., J. L. NIEBER, G. JOHNSON, J. MAGNER & B. SCHAEFER. 2007.

Flow path influence on an N:P ratio in two headwater streams: a paired watershed

study. *Journal of Geophysical Research,* 112. DOI:10.1029/2007JG000403

HILL, A. R. 1996. Nitrate removal in stream riparian zones. *Journal of Environmental*

Quality, 25 (4): 743-755. DOI.org/10.2134/jeq1996.00472425002500040014x

HILL, W. R., P. J. MULHOLLAND & E. R. MARZOLF. 2001. Stream ecosystem

response to forest leaf emergence in spring. *Ecology,* 82 (8): 2306-2319.

KAHLERT, M. 1998. C:N:P ratios of freshwater benthic algae. *Archiv für Hydrobiologie,*

Special Issues, Advances in Limnology, 51: 105-114.

LORENZEN, C. J. 1967. Determination of chlorophyll and pheopigments:

spectrophotometric equations. *Limnology and Oceanography,* 12: 343-346.

MISERENDINO, M. L., R. CASAUX, M. ARCHANGELSKY, Y. C. DI PRINZIO, C.

BRAND & M. A. KUTSCHKER. 2011. Assessing land-use effects on water

quality, in-stream habitat, riparian ecosystems and biodiversity in Patagonian

northwest streams. *Science of the Total Environment,* 409: 612-624.

DOI:10.1016/j.scitotenv.2010.10.034

PARUELO, J. M., A. BELTRAN, E. JOBBÁGY, O. SALA & R. GOLLUSCIO. 1998. The

climate of Patagonia: general patterns and controls on biotic processes. *Ecología*

Austral, 8 (2): 85-101.

SATTI, P., M. J. MAZZARINO & M. GOBBI. 2003. Soil N dynamics in relation to leaf

litter quality and soil fertility in north-western Patagonian forests. *Journal of*

Ecology, 91 (2): 173-181. DOI.org/10.1046/j.1365-2745.2003.00756.x

SCHANZ, F. & H. JUON. 1983. Two different methods of evaluating nutrient limitantions

of periphyton bioassays, using water from the River Rhine and eights of its

tributaries. *Hydrobiologia,* 102: 187-195. DOI.org/10.1007/BF00006346

SOSNOVSKY, A., M. E. LALLEMENT, M. RECHENCQ, E. E. ZATTARA, M. V.

FERNÁNDEZ, R. DAGA, M. J. SUAREZ, S. LEIVA & R. J. C. CANTET. 2019.

Topography and Land Use Modulate Hydrological and Nutrient Export Dynamics

in two Andean Streams. *BioRxiv*. DOI: https://doi.org/10.1101/742411

SOSNOVSKY, A., M. RECHENCQ, V. FERNÁNDEZ, M. J. SUAREZ & R. J. C.

CANTET. 2020. Hydrological and Physico-chemical dynamics in two Andean

streams. *Limnetica,* 39 (1): 17-33. DOI: 10.23818/limn.39.02

274
275

Number/Site	Population (inhabitants km^{-2})	Nutrient export ($kg\ km^{-2}\ y^{-1}$)				Chlorophyll ($mg\ m^{-2}$)
		Nh_4	$NO_2 + NO_3$	DIN	SRP	
1) Str. Torrontegui	0	5.0	0.3	5.3	1.08	7
2) Str. Melgarejo	0	nd	0.2	0.2	1.27	7
3) Str. Pescadero	4	nd	nd	nd	0.08	117
4) Str. Loncochinoco	45	1.5	162.2	163.7	3.97	52
5) Str. Cascada	13	nd	0.2	0.2	0.06	77
6) Gutiérrez rise	7	nd	0.1	0.1	0.92	7
7) Gutiérrez 7	19	0.3	3.7	4.0	1.83	66
8) Gutiérrez 8	18	1.4	3.6	5.0	0.55	12
9) Gutiérrez 9	23	2.2	59.9	62.1	2.03	4
10) Gutiérrez mouth	28	1.0	83.1	84.1	1.02	29

276 **Table 1 annexed** Data supporting Figures 1 and 2. Population density and limnological
277 characteristics of the study sites (Str.) Streams.
278 Ammonium (Nh_4); Nitrates + Nitrites ($NO_3 + NO_2$); Soluble reactive phosphorus (SRP). Nd: not
279 detectable.
280