1   A shift to shorter cuticular hydrocarbons causes sexual isolation in the *Drosophila americana* species
2   group

4   Jeremy S. Davis[1, 2]*, Matthew A. Pearcy[1], Joanne Y. Yew[3], Leonie C. Moyle[1]

5   [1]Dept. of Biology, Indiana University, Bloomington, IN, USA 47405

6   [2]Dept. of Biology, University of Kentucky, Lexington, KY, USA 40508

7   [3]Pacific Biosciences Research Center, University of Hawaiʻi at Mānoa, Honolulu, HI 96822

9   *Corresponding author: J.Davis@uky.edu

Abstract

Because sensory signals often evolve rapidly, they could be instrumental in the emergence of

reproductive isolation between species. However, pinpointing their specific contribution to isolating

barriers, and the mechanisms underlying their divergence, remains challenging. Here we demonstrate

sexual isolation due to divergence in chemical signals between members of the *Drosophila americana*

group, dissect its underlying phenotypic and genetic mechanisms, and propose a model of its

evolutionary history. Mating trials revealed complete sexual isolation between *Drosophila*

novamexicana males and *Drosophila americana texana* females*,* as well as more moderate bi-directional

isolation between *D. americana* subspecies*.* Mating behavior recordings indicate *D. a. texana* males

have the highest courtship efficiency and, unlike males of the other two species, are accepted by

females of all species. Quantification of cuticular hydrocarbon (CHC) profiles, chemosensory signals that

are used for species recognition and mate finding in Drosophila, shows that *D. a. texana* differs from the

other two species primarily on the basis of compound carbon chain-length. Moreover, manipulation of

male CHC composition via heterospecific perfuming—specifically perfuming *D. novamexicana* males

with *D. a. texana* males—abolishes their sexual isolation from *D. a. texana* females. Evaluation of

candidate genes for patterns of gene expression reveals a single gene—elongase CG17821—with

patterns consistent with a causal role in CHC differences between species. Sequence comparisons

indicate *D. novamexicana* and *D. a. americana* share a derived CG17821 truncation mutation that could

also contribute to their shared "short" CHC phenotype. We present an evolutionary model for the origin

and spread of this allele and the consequences for CHC divergence and sexual isolation in this group.

Introduction

Sensory signals can act as sexual cues that are critical for intraspecific mate evaluation and

reproductive success. Moreover, because they are frequently among the most rapidly evolving species

differences (Smadja and Butlin 2009, Seddon et al 2013, Wilkins et al 2013), divergence in sensory sexual

signals might often contribute to the earliest stages of reproductive isolation, in the form of prezygotic

barriers between lineages (Butlin et al 2012, Ritchie 2007). Among all such sexual signals, nowhere is the

diversity more evident than those acting as premating traits, including coloration and other visual

signals, acoustic signals, and complex pheromone blends. However, convincingly demonstrating the

connection between such sensory signal divergence and emerging reproductive isolation can be

challenging because it requires, first, identification and demonstration of the direct role of specific

signals mediating sexual isolation between species and, second, understanding the specific mechanistic

changes that have given rise to lineage differences in this signal.

Among insects, sexual signals draw on three primary sensory modalities—visual, auditory, and

chemosensory—and most species likely use a combination of all three modalities to identify potential

mates. Of these, many insect chemosensory signals primarily consist of cuticular hydrocarbons (CHCs)—

a broad group of carbon-chain compounds important for many essential functions including sex

pheromone signaling, but also environmental adaptation (Blomquist and Bagneres 2010, Chung and

Carrol 2015, Yew and Chung 2017). Unlike many auditory and visual cues, CHCs are produced and

received by both males and females—making them potentially important for both male and female

mate choice or species recognition. Indeed, studies on *Drosophila* chemical communication—

particularly in the *melanogaster* subgroup—have detected clear evidence for sexual isolation based on

male choice of female CHCs (Coyne et al. 1995, Billeter et al. 2009), while work in other species has

demonstrated female choice of species-specific male CHC profiles (Coyne et al. 2002, Mas and Jallon

2005, Curtis et al. 2013, Dyer et al. 2014). Elements of the biochemical pathways of CHC production and

the underlying genes are also relatively well understood in *D. melanogaster* (Pardy et al 2018).

Accordingly, single genes contributing to species-specific CHC differences have been implicated as

causes of reproductive isolation in several cases, including between *D. melanogaster* and *D. simulans*

(desatF: Legendre et al. 2008) and *D. serrata* and *D. birchii* (mFAS: Chung et al. 2014). This framework

provides an excellent resource for identifying candidate genes for sensory sexual signal variation in

other *Drosophila* systems.

62   Here we use the *Drosophila americana* group to investigate how species variation in CHC

profiles contributes to reproductive isolation and to identify the underlying chemical and genetic

changes that may comprise this variation. This group—consisting of two subspecies *Drosophila*

americana americana and *Drosophila americana texana* and their sister species *Drosophila*

novamexicana—is a member of the *virilis* clade found across North America. These species are closely

related (MRCA ~0.5 mya, Morales-Hojas et al. 2008), and have been shown to occupy distinct

geographic and environmental habitats (Davis and Moyle 2019). Classic studies have shown some

evidence for premating isolation (Spieth 1951) as well as differences in CHC composition between males

(Bartelt at al. 1986), however the contribution of hydrocarbons to premating isolation has not been

assessed. With lines from each of the three lineages *D. novamexicana*, *D. a. americana*, and *D. a. texana*

(referred to here as species, for simplicity), here we use a combination of mating and behavioral studies,

chemical analysis and manipulation, and gene expression and molecular sequence variation analyses to

assess the role of CHCs in reproductive isolation. Together, our data suggest that evolution of a male

sexual signal—an overall shift in the relative abundance of longer versus shorter cuticular hydrocarbons,

due to novel mutation in an elongase gene—has produced complete premating isolation between

derived males and females from species that retain the ancestral trait and preference, as proposed in

classical models (Kaneshiro 1976, 1980) of the evolution of asymmetric sexual isolation.

Results

D. a. texana *females discriminate against heterospecific males in mating trials*

We found clear evidence of moderate to strong sexual isolation between *D. a. texana* and the

other two species (Table 1). Mating rate (average proportion of females mated, in 4x4 mating trials)

ranged from very high in conspecific crosses and some heterospecific crosses, to <50% in heterospecific

combinations of female *D. a. texana* with each of the other species. Notably, *D. novamexicana* males

were never successful in mating with *D. a. texana* females, indicating complete sexual isolation in this

cross direction between these two species. In contrast, mating rate in the reciprocal cross was 70%.

Other pairings with mating rates at or below 50% were both reciprocal directions between *D. a.*

americana and *D. a. texana*. Across all cross combinations in the 4x4 mating trials, we found that female

identity (Kruskal-Wallis test: $\chi^2(2)$ = 8.99, *P* = 0.012), but not male identity (Kruskal-Wallis test: $\chi^2(2)$ =

0.80, *P* = 0.67), significantly affected success. Post-hoc comparisons confirmed no pairwise differences in

mean mating rate between any male species (not shown). In contrast, *D. a. texana* female copulation

rate was significantly lower on average than both *D. a. americana* (*P-adj* = 0.025) and *D. novamexicana*

(*P-adj* = 0.042), whereas *D. a. americana* and *D. novamexicana* females did not differ (*P-adj* = 0.15).

Together, these results indicate that *D. a. texana* females discriminate against heterospecific males

more strongly than do females of the other species, with the greatest discrimination against *D.*

novamexicana males. In addition, while we observe clear asymmetric sexual isolation between *D. a.*

texana and *D. novamexicana*, premating isolation between the two *D. americana* subspecies is bi-

directional and more moderate, indicating the specific mechanisms of isolation might differ among pairs

of species within this group (see discussion).

Table 1: Copulation rates and latency for each cross in unperfumed 4x4 and 1x1 mating trials.

Female species	Male species	4x4 mating trials	1x1 mating trials	
		Mating rate[a] (mean ± SE)	Mating rate[a] (mean)	Copulation latency (mean ± SE)[b]
D. novamexicana	*D. novamexicana*	0.90 ± 0.06	1.00	69.89 ± 22.73
	D. a. americana	0.69 ± 0.12	0.60	147.48 ± 27.00
	D. a. texana	0.71 ± 0.18	0.60	83.63 ± 39.70
D. a. americana	*D. novamexicana*	1.00 ± 0	0.20	158.33 ± 21.67
	D. a. americana	1.00 ± 0	0.60	123.46 ± 29.01
	D. a. texana	0.50 ± 0.18	0.60	99.42 ± 34.74
D. a. texana	*D. novamexicana*	0.00 ± 0	0.00	180 ± 0
	D. a. americana	0.40 ± 0.14	0.60	99.42 ± 34.74
	D. a. texana	0.85 ± 0.06	0.80	50.32 ± 33.07

[a] n = 5, rate is a proportion out of maximum possible matings (4 or 1)

[b] n = 5, units are in minutes. Unmated trials are scored as a latency of 180 min—the maximum time observed.

D. a. texana *males have greater courtship efficiency and mating success*

We video-recorded single-pair matings for each cross combination to evaluate species differences in courtship strategies and whether these differed in heterospecific pairings. Three male courtship behaviors were quantified from these 1x1 matings—display rate, tapping rate, and licking rate. All three behaviors showed similar patterns of among-species variation, when each was evaluated for the effect of male species, female species, or their interaction, using 2-way ANOVAs. Males significantly differed with respect to display-rate ($F(2) = 3.76$, $P = 0.03$) and tap-rate ($F(2) = 4.76$, $P = 0.017$), but only marginally for lick-rate ($F(2) = 2.67$, $P = 0.083$). For all three behaviors, this difference is due to *D. a. texana* males exhibiting higher rates (Tukey HSD post-hoc tests; Figure 1, supplementary Table S1). In contrast, we detected no female identity effects on male behavioral rate (display-rate: $F(2) = 1.83$, $P = 0.18$; tap-rate $F(2) = 0.58$, $P = 0.56$; lick-rate: $F(2) = 0.43$, $P = 0.65$), or any female x male interactions (display-rate: $F(2) = 0.72$, $P = 0.58$; tap-rate $F(2) = 0.45$, $P = 0.77$; lick-rate: $F(2) = 0.65$, $P = 0.63$). Because behavior rates are dependent on copulation latency, and not simply the total number of behavioral events, these rates approximate the efficiency with which each courtship behavior results in a mating. Taken together, our results indicate that *D. a. texana* males have greater courtship efficiency (in terms

of display and tap rates), and that males do not significantly vary these behavioral rates depending on

female species identity.

The single pair mating assays broadly reiterated the patterns of moderate to strong sexual

isolation we observed for *D. a. texana* in 4x4 trials. As with our 4x4 mating assay, individual pairs of *D.*

novamexicana males never mated with *D. a. texana* females, while mating rate between the two *D.*

americana subspecies was moderately reduced in both directions. A logistic regression assessing how

mating rate varied based on male identity, female identity, and their interaction showed a significant

interaction effect only (males: $\chi^2(2)$ = 2.97, *P* = 0.28; females: $\chi^2(2)$ = 2.52, *P* = 0.23; males*females: $\chi^2(4)$

= 12.67, *P* = 0.013)—a difference from 4x4-mating trials where female species identity was the primary

predictor of mating rate. This variation between the assays appears to be driven by female *D. a.*

americana accepting fewer *D. novamexicana* males in 1x1 trials (20%) compared to 4x4 trials (100%,

Table 1), an observation that suggests male density within mating trials might affect copulation success

in this particular species combination. Finally, copulation latency differed marginally based on both male

identity (ANOVA: *F*(2) = 3.14, *P* = 0.055) and the male by female interaction (*F*(4) = 2.48, *P* = 0.062), but

did not differ based on female species identity (*F*(2) = 0.68, *P* = 0.5147). Post-hoc tests suggest that the

marginal male species effect is likely due to lower copulation latency in male *D. a. texana* relative to *D.*

novamexicana males (Tukey HSD; Nov-Tex: *P-adjust* = 0.07; Ame-Tex: *P-adjust* = 0.23; Nov-Ame: *P-adjust*

= 0.81).

Figure 1: Male courtship behavior rates in 1x1 mating assays across all cross types. Males (columns) show significant differences in display rate and tap rate, and marginal differences in lick rate. Rates did not differ based on female identity (rows) or male x female interaction (see results). Bar and whiskers indicate mean and standard error (SE).

Species and sexes differ in CHC composition

We found that both species and sexes within species showed different and distinctive CHC

profiles, with the largest differences detected between *D. a. texana* and the other two species (Figure

2). Across all samples (*n*=5 replicate samples for each identity) we detected 8 alkene compounds (2 of

which are sex-specific) and 4 methyl-branched alkane compounds present in at least one of the species.

A principal component analysis of profiles from unmanipulated flies (unmanipulated-PCA or 'U-PCA')

summarizing the primary axes of CHC profile composition found that 95.3% of compound variation

across all samples was explained by the first three principal components (U-PC1 to 3). Of these, CHC

composition varied significantly for both species and sex for U-PC1 (species: $F(2, 429.29)$, $P < 0.0001$;

sex: $F(1, 46.97)$, $P < 0.0001$) and U-PC2 (species: $F(2, 10.54)$, $P = 0.00043$; sex: $F(1, 181.69)$, $P < 0.0001$),

but had no sex or species effect for U-PC3 (species: $F(2, 0.31)$, $P = 0.737$; sex: $F(1, 0.57)$, $P = 0.458$)(Figure

2). Notably, *D. a. texana* of both sexes differed from the other two species along the U-PC1 axis (Tukey

HSD; Nov-Tex: *P-adjust* < 0.0001; Ame-Tex: *P-adjust* < 0.0001; Nov-Ame: *P-adjust* = 0.28), which also

explains most of the CHC variation between samples (69.9%). U-PC1 was positively loaded for most of

the shorter carbon chain length compounds in both alkene and methyl-branched alkane classes of

compounds (Table S2); consequently, negative values were strongly loaded for the compounds with

longer carbon chain length, and this axis can be interpreted as a composite of average compound length

across the CHC profile as a whole (Figure 2). Accordingly, both sexes of *D. a. texana* had a higher

abundance of longer-chain alkenes and methyl-branched alkanes than did either *D. a. americana* or *D.*

novamexicana.

In contrast to U-PC1, U-PC2 appeared to primarily differentiate sexes within species, but also *D.*

a. americana males from males of the other two species. This axis was most heavily loaded for two

compounds: C21:1 and Me-C28, followed by C27:1 and Me-C26, with much smaller loadings for all other

compounds (Table S2). C21:1 is a male-specific compound that is not detected in females; Me-C28, Me-

26, and C27:1 abundance was also consistently different between sexes, with males always having more

of these CHCs than females for each species (Figure 2). Therefore, negative U-PC2 values can be

interpreted as primarily representing more male-specific profiles, while positive U-PC2 values

correspond to more female-like profiles. With respect to the detected species difference in U-PC2, post-

hoc tests reveal that this was driven by *D. a. americana* (Tukey HSD; Nov-Tex: *P-adjust* = 0.99; Ame-Tex:

P-adjust = 0.0062; Nov-Ame: *P-adjust* = 0.0088). Furthermore, a post-hoc test specifically comparing the

abundance of the male C21:1 compound confirms a difference between *D. a. americana* and males of

the other two species, but not between *D. novamexicana* and *D. a. texana* (Tukey HSD; Nov-Tex: *P-*

adjust = 0.98; Ame-Tex: *P-adjust* = 0.016; Nov-Ame: *P-adjust* = 0.022).

Figure 2: Cuticular hydrocarbon composition of unmanipulated males and females of each species
stock. A) Relative log-scale abundance of compounds for males (upper) and females (lower), for each
major compound type (line = mean, box = central quartile, whiskers = S.E.). *n* = 5 samples for each sex
and line. Test statistics for species differences are in Table S3. B) First two principal components (U-
PC1 and U-PC2) of composite CHC variation among males (circle) and females (triangle). Ellipses
indicate 90% bivariate normal density for each species-sex group. Species and sex significantly
influenced both U-PC1 and U-PC2.

Heterospecific perfuming influences D. a. texana *female acceptance of con- and hetero-specific males*

We found that patterns of sexual isolation could be modified by specifically changing the CHC

profiles of males of different species. To do this, we used perfuming assays to manipulate the

pheromone profile of males by co-housing them with either con- or hetero-specific males, and then

evaluated how this manipulation influenced mating rates among species. For these analyses, we focused

on two species pairs in two separate, analogous experiments—*D. novamexicana* and *D. a. texana* ("Nov-

Tex pair"), and *D. a. americana* and *D. a. texana* ("Ame-Tex pair")— because *D. a. texana* shows the

strongest sexual isolation from the two other species, and the largest differences in CHC composition.

Within each perfuming experiment, we generated four combinations of target and donor male

identities: each species perfumed by conspecific males (control) and each species perfumed by

heterospecific males. Each class of perfumed males was evaluated for mating rate with *D. a. texana*

females, because these females were the most discriminating against heterospecific males in our

previous mating assays (Table 2). The observed mating rate of males perfumed with conspecific males

(the control manipulation) was similar to that observed in unperfumed mating trials: *D. novamexicana*

and *D. a. americana* males perfumed with their own species performed poorly with *D. a. texana*

females, relative to conspecific *D. a. texana* perfumed with their own males, which had a 100% mating

rate (Table 2). In strong contrast, heterospecific perfumed males showed altered mating rates relative to

control perfumed males. For Nov-Tex pairings, *D. novamexicana* males perfumed with *D. a. texana*

males successfully mated with *D. a. texana* females 94% of the time, a significantly higher mating rate

than *D. novamexicana* males perfumed with their own males (19%; Mann-Whitney U-test: U = 0, *P* =

0.025). Likewise, the reciprocal treatment—male *D. a. texana* perfumed with *D. novamexicana* males—

showed significantly reduced copulation success (19%) relative to conspecific-perfumed *D. a. texana*

(Mann-Whitney U-test: U = 0, *P* = 0.018). For Ame-Tex pairings, *D. a. texana* male mating rate was also

significantly reduced when perfumed with *D. a. americana* males (Mann-Whitney U-test: U = 0, *P* =

0.018), compared to *D. a. texana* males perfumed with conspecifics. In contrast, *D. a. americana* males

perfumed with *D. a. texana* males did not significantly differ from the mating rate of control

(conspecific-perfumed) males (Mann-Whitney U-test: U = 5.5, *P* = 0.54).

Overall, these results indicate that perfuming males with heterospecific CHCs influences the

frequency of successful copulation with *D. a. texana* females. *D. a. texana* males perfumed with

heterospecific profiles (of either *D. novamexicana* or *D. a. americana*) had significantly reduced mating

rates with their own females. Conversely, perfuming *D. novamexicana* males with heterospecific *D. a.*

texana males significantly increased their mating rate with *D. texana* females, almost completely

reversing the pattern of sexual isolation observed for unmanipulated males. This large shift in mating

rate for heterospecifically-perfumed *D. novamexicana* indicates that differences in male CHC profile play

a critical role in sexual isolation in the Nov-Tex pair. In contrast with the three other classes of

heterospecifically-perfumed males, perfuming *D. a. americana* with *D. a. texana* males did not

significantly change their mating rate. It possible that this manipulation might have been less effective at

changing the CHC profile of *D. a. americana* males (see next section), and/or that mating isolation in the

Ame-Tex pair might depend on more complex factors than a simple shift in CHC composition (see

Discussion).

Table 2: Mean copulation success (number of matings out of 4 per trial) with *D. a. texana* females for each perfuming identity in manipulated (perfumed) 4x4 mating assays.

Target male	Donor male	Copulation success (Mean ± SE)	Pairwise Mann-Whitney U test	
			U-score	*P*-value
Nov-Tex pair				
D. novamexicana	*D. novamexicana*	0.25 ± 0.15	} 0	0.0247
D. novamexicana	*D. a. texana*	0.94 ± 0.063		
D. a. texana	*D. a. texana*	1.00 ± 0	} 0	0.018
D. a. texana	*D. novamexicana*	0.19 ± 0.063		
Ame-Tex pair				
D. a. americana	*D. a. americana*	0.44 ± 0.12	} 5.5	0.54
D. a. americana	*D. a. texana*	0.56 ± 0.12		
D. a. texana	*D. a. texana*	1.00 ± 0	} 0	0.018
D. a. texana	*D. a. americana*	0.50 ± 0.063		

Note: *n* = 5 for all crosses

Perfuming shifts CHC composition towards the donor male profile

Using CHCs extracted from an additional set of perfumed males (see Methods), we confirmed that our heterospecific perfuming manipulation produced quantitative changes in composite male CHC profiles in both the Nov-Tex and Ame-Tex pairs, by shifting male CHCs closer to the donor male profile. PCAs were performed separately for each species pair (N-PCA for the Nov-Tex pair, and A-PCA for the Ame-Tex pair, hereafter), as were analyses of differences among classes of perfumed males. In both cases, of the first three PCs of CHC composition, PC1 (i.e. N-PC1 or A-PC1) varied by both donor and target male identity (Table S4), indicating that perfuming significantly shifted CHC profiles along the primary axis of variation in both perfuming experiments. Nonetheless, the specific donor-target manipulation that was most successful in this regard differed between the Nov-Tex and Ame-Tex pairs. In the Nov-Tex pair, *D. novamexicana* males perfumed with *D. a. texana* males significantly differed from control (*D. novamexicana*) perfumed samples in CHC composition (N-PC1) (t(5.9) = -2.8, *P* = 0.031); however, heterospecific-perfumed *D. a. texana* did not differ from conspecific *D. a. texana*-perfumed samples along the same axis (t(3.53) = 1.91, *P* = 0.14). In contrast, in the Am-Tex pair, *D. a. texana* perfumed with *D. a. americana* showed a significant shift in A-PC1 (t(4.85) = 4.23, *P* = 0.0088), but there

was no significant shift for *D. a. texana*-perfumed *D. a. americana* compared to conspecific-perfumed

controls (t(4.68) = 0.055, *P* = 0.96). The difference in pattern between these pairs might be attributed to

high variation seen among samples within a given group (in particular conspecific perfumed *D. a.*

texana). Regardless, it is clear from these results that this perfuming assay can produce significant,

detectable differences in CHC profiles after heterospecific perfuming, most notably in *D. novamexicana*

males.

Patterns of gene expression and sequence variation implicate an elongase gene that contributes to CHC

variation between species

From among a set of 23 candidate genes whose orthologs have functions related to cuticular

hydrocarbon variation, both transcriptome and sequence comparisons implicated one specific elongase

locus as potentially causal in CHC composition differences between our species. Using whole-

transcriptome RNA-seq data from the same three populations (previously generated in Davis and Moyle

2020), we found that 3 of our candidate genes in males and 2 candidates in females were within the top

10[th] percentile of genes most differentially expressed between species across the whole transcriptome

(i.e. they showed a stronger species effect than 90% of all 11301 expressed genes) (Table 3). Of these

candidate genes, only CG17821 showed elevated gene expression specifically in *D. a. texana* compared

to the other two species; this pattern was observed in both sexes but is more significant in females

(Figure 3).

Closer inspection of our CG17821 sequences revealed that alleles in *D. a. americana* and *D.*

novamexicana share a thymine insertion mutation that causes a premature stop codon 4 amino acids

upstream of the end of the gene, compared to the annotated gene model in outgroup *D. virilis* and the

allele in our *D. a. texana* stock (Figure 3); these four terminal amino acids are not present in RNA

transcripts for either *D. a. americana* and *D. novamexicana*. Among our three lines, this suggests an

association between the truncated protein product of CG17821 and the "short" CHC phenotype we

observe. This association is further supported by additional CHC and sequence data from another *D.*

americana population, originally collected in Anderson, IN (species stock center line 15010-0951.00).

Lamb et al. 2020 characterized CHC divergence between females from this *D. a. americana* stock and

another *D. novamexicana* line (species stock center 15010-1031.04). The profile for the *D.*

novamexicana line had absence of long (>C30) compounds, low abundance of C29 compounds, and

presence of multiple short alkenes (<C27), consistent with the "short" CHC profile we observe for our *D.*

novamexicana line. Likewise, the Anderson *D. a. americana* profile they observed is broadly consistent

with the "long" profile we observed here in *D. a. texana*, as both show presence of compounds longer

than 30 carbons, a high abundance of C29 length alkenes, and absence of compounds shorter than 27

carbons. In addition, using a genome assembly from the same Anderson *D. americana* stock (Kim et al.

unpubl.), we found that this population has the non-truncated allele of CG17821. These data are

consistent with our inference that truncation of this allele could contribute to the difference between

"long" and "short" CHC phenotypes in these species.

Evolutionary history of elongase CG17821

Several lines of evidence indicate that CG17821 has a complex evolutionary history within the *D.*

americana subgroup, that likely includes post-speciation introgression. First, the presence of the non-

truncated allele in outgroup *D. virilis* (Figure 3) indicate that the truncated allele is derived from a

thymine insertion mutation event that took place within the *D. americana* subgroup. Second, the

observation that *D. novamexicana* and our western *D. a. americana* line share an identical derived allele

at CG17821 indicates an evolutionary history at this locus that disagrees with expected phylogenetic

relationships among these three taxa (i.e. where lines from the two *D. americana* subspecies are

expected to be most closely related). Gene trees for each of our 23 candidate genes confirmed that

CG17821 has a topology that is discordant with the expected species relationships, placing our western

D. a. americana as sister to *D. novamexicana* rather than grouping it with *D. a. texana* (Figure 3). Four

other CHC candidate genes also show this discordant topology, most notably CG18609 which is located

immediately downstream of CG17821 (within 1kB) according to the annotated genome of *D. virilis*

(Figure 3). For 4 out of 5 of these loci—including both CG17821 and CG18609—using the allele from the

eastern (Anderson, IN) *D. a. americana* stock instead produces a genealogy where *D. a. americana*

groups with *D. a. texana*, as expected for the species tree (Figure 3).

The observation of phylogenetically discordant sites can be due to several factors, notably

incomplete lineage sorting (ILS) or post-speciation introgression. However, evidence from SNP variants

across the whole transcriptome and specifically at CG17821 is more consistent with introgression

between *D. novamexicana* and our western *D. a. americana* population. Genome-wide, of 34441 total

SNPs detected in 11301 loci within the transcriptome dataset (Davis and Moyle 2020), 14902 supported

D. americana subspecies as sister (in accordance with the species tree), 13372 SNPs grouped *D.*

novamexicana and *D. a. americana*, and 6167 grouped *D. novamexicana* with *D. a. texana*. These data

indicate a significant excess of shared variants between *D. novamexicana* and our western *D. a.*

americana line (D = 0.369, P < 0.0001), consistent with a history of introgression between the

progenitors of these two populations. Moreover, gene tree topologies at and around CG17821 also

indicate this specific genomic region shares recent ancestry between *D. novamexicana* and western *D. a.*

americana due to introgression. CG17821, the adjacent downstream CHC candidate CG18609, and the

next nearest gene (List, a neurotransmitter located ~14 kB downstream) all display shared derived SNPs

between *D. novamexicana* and our western *D. a. americana* line, and discordant gene tree topologies

that group these two lines as sister. In contrast, topologies for the next two nearest up/downstream

genes in this region reflect the species tree, indicating that recent shared ancestry between *D.*

novamexicana and our western *D. a. americana* extends across a genomic window between 18 and 68

kB long around the specific region containing CG17821 (Figure 3C). Estimates of linkage disequilibrium

(LD) from *D. melanogaster* autosomes indicate that most LD decays within 200 bp, and r^2 decreases to

around 0.1 within a 1 kB window (Franssen et al. 2015); therefore the size of the window observed here

is well outside the range expected to show discordant ancestry due to sorting from ancestral variation

(i.e. ILS). Coupled with genetic and phenotypic evidence above, these results indicate introgression of

these genes between *D. novamexicana* and western *D. a. americana*—and not ILS—is most likely

responsible for their shared allele at CG17821 and therefore potentially for their similarity in "short"

CHC phenotypes.

Table 3: Species differences in quantitative gene expression for 23 candidate genes associated with
CHC function, analyzed separately by sex, with *P*-value and rank compared to all genes in dataset.

Candidate gene	motif	Males		Females	
		P-value	Rank (%)	*P*-value	Rank (%)
Cyp4g1	Cyp4g	0.87	10780 (95%)	0.86	9834 (87%)
Cyp4g15	Cyp4g	0.22	3702 (33%)	0.41	4286 (38%)
Cyt-b5-r	desat	0.19	3244 (29%)	0.51	5946 (53%)
ifc	desat	0.89	10862 (96%)	0.52	5906 (52%)
CG17928	desat	0.40	6375 (56%)	0.98	10883 (96%)
CG8630	desat	0.11	2110 (19%)	0.30	3086 (27%)
CG9743	desat	0.15	2660 (24%)	0.25	2443 (22%)
desat2	desat	0.81	10465 (93%)	0.45	5138 (45%)
Baldspot	elongase	0.49	7711 (68%)	0.45	5078 (45%)
bond	elongase	0.38	5988 (53%)	0.76	8901 (79%)
ELOVL	**elongase**	**0.019**	**646 (6%)**	0.15	1535 (14%)
CG5326	elongase	0.087	1763 (16%)	0.87	10008 (89%)
CG17821	**elongase**	0.11	2099 (19%)	**0.0033**	**99 (0.8%)**
CG18609	**elongase**	**0.026**	**781 (7%)**	0.24	2325 (21%)
CG30008	**elongase**	0.21	3460 (31%)	**0.069**	**807 (7%)**
CG31522	elongase	0.52	8052 (71%)	0.70	8052 (73%)
CG31523	**elongase**	**0.040**	**1024 (9%)**	0.95	10710 (95%)
CG33110	elongase	0.65	9332 (83%)	0.67	7863 (70%)
CG6660	**elongase**	0.39	6150 (54%)	0.36	3752 (33%)
FASN1	fatty-acid-synthase	0.32	5197 (46%)	0.62	7194 (64%)
FASN2	fatty-acid-synthase	0.25	4136 (37%)	0.41	4385 (39%)
ebony	pigment	0.22	3686 (33%)	0.38	3948 (35%)
tan	pigment	0.31	5029 (45%)	0.72	8460 (75%)

Note: **Bold** genes indicate gene expression difference in the top 10th percentile

Figure 3: CHC candidate gene expression, genealogical relationships, and sequence variation among *D. a. americana* group lines. A) Gene expression variation showing mean and standard error between focal species lines for candidate genes with the largest differences in either males (upper, circles) or

females (lower, triangles). Gene expression is measured in transcripts per million (TPM, n = 3 for each
sample). B) Gene tree topologies for all 23 candidate genes using either western (Nebraska) or eastern
(Indiana) populations for *D. a. americana*. C) Schematic of loci (green/blue boxes) in the genomic
window surrounding candidates CG17821 and CG18609. Blue boxes indicate a genealogy that groups
the western *D. a. americana* population with *D. novamexicana*; gene topology of shaded blue region
displayed below line. Green boxes indicate loci with genealogies matching the expected species tree.
D) 3' terminal nucleotides and occurrence of inferred ancestral and derived (truncated) allele in
CG17821 among lineages with sequence data.

Discussion

344
Identifying the genetic and evolutionary mechanisms underlying the first steps of reproductive

isolation between species is essential for understanding the factors that drive speciation. Evolutionary

divergence in sexual signals may be an especially potent contributor to this process (Schemske 2000,

Coyne and Orr 2004, Ritchie 2007, Schluter et al. 2009, van Doorn et al. 2009). However, demonstrating

the connection between sensory signal divergence and emerging reproductive isolation can be

challenging, as it requires identification and demonstration of the direct role of specific signals

mediating sexual isolation between species and knowledge of the specific mechanistic changes that

have given rise to lineage differences in this signal. Here we have demonstrated that sexual isolation

between species in the *D. americana* group is based on female choice of male chemical signals,

identified both the specific phenotypic shift between species in pheromone chemistry and a genetic

variant likely contributing to this phenotypic change, and uncovered evidence of a complex evolutionary

history for this hypothesized causal locus. Together these data support a clear role for sensory signal

divergence in the evolution of premating species barriers in this diverse wild group and provide data on

the likely genetic and evolutionary mechanisms involved in its evolutionary emergence.

Female choice of male CHC variation is responsible for isolation between D. novamexicana *and* D. a.

texana

Differences in copulation success between species are the product of variation in male choice

(via differences in courtship intensity and copulation attempts, depending upon female identity), female

choice (via differences in acceptance rates of males, depending upon their identity), or a combination of

these factors. Disentangling these alternatives is critical for identifying the specific trait and

corresponding preference variation responsible for the emergence of prezygotic species barriers. Here

we showed that female choice of a male sensory signal results in strong sexual isolation within the *D.*

americana subgroup, specifically *D. a texana* female preference for CHC profiles of their own males and

rejection of male with heterospecific profiles. The role of CHCs is exceptionally clear in the case of

premating isolation between male *D. novamexicana* and female *D. a. texana*: otherwise complete sexual

isolation in this cross can be almost entirely reversed by perfuming *D. novamexicana* males with *D. a.*

texana male CHCs. In contrast, we find little evidence for differential male preference of females, even

though female CHC profiles are also divergent between species. Instead, our behavioral data indicates

that males did not alter their courtship behavior in response to female species identity. Interestingly,

these observations also suggest evidence for female choice of male courtship behaviors, whereby *D. a.*

texana courtship consistently induced females to copulate sooner than courtship behaviors of other

species males.

These findings fit within a body of studies that have identified either male or female choice of

sensory signals as critical for sexual isolation among *Drosophila* species (patterns reviewed Yukilevich

and Patterson 2019). In *Drosophila melanogaster,* many studies examining CHC divergence have focused

on male choice based on female compounds, and found that specific female compounds play a role in

isolation between closely related heterospecifics (Coyne et al 1995, Billeter et al. 2009) as well as

between intraspecific populations (Wu et al. 1995, Hollocher et al. 1997, Yukilevich and True 2008).

Female choice of male CHC variation has received comparably less mechanistic attention, however has

been demonstrated as a prezygotic barrier in several diverse *Drosophila* groups, most notably between

D. santomea and *D. yakuba* (Coyne et al. 2002, Mas and Jallon 2005), between the mycophagous *D.*

subquinaria and *D. recens* (Curtis et al. 2013, Dyer et al. 2013), and between *D. mojavensis* males reared

on different cactus substrates (Havens and Etges 2013). Our results therefore complement a growing

body of evidence that shows female choice can be an important factor dictating patterns of sexual

isolation among diverse wild *Drosophila* species.

CHC divergence and the role of Elongases

Dissecting the finer details of phenotypic divergence in sensory signals can help pinpoint

underlying mechanisms and associated genes responsible, and more clearly demonstrate how these

signal changes contribute to species isolation. Here we found that CHC divergence between our lines

occurs primarily on the basis of compound length, with *D. novamexicana* and *D. a. americana* profiles

both having similar enrichment of shorter compounds ("short" phenotype) compared to an enrichment

of longer compounds for *D. a. texana* ("long" phenotype). These differences in abundance of shorter-

versus longer-chain compounds were observed across both sexes, and across both alkenes and methyl-

branched alkanes (Figure 2, Table S3). In contrast, we find no evidence for variation in other features

such as double bond or methyl branch location or number. These features themselves suggest that the

striking difference between 'shorter' and 'longer' profile phenotypes could be due to variation in fatty-

acid elongase activity, which globally influences the carbon chain length of CHC precursors, and

therefore can have consistent downstream effects on both alkenes and methyl-branched alkanes,

because both are modified after the elongation step (Pardy et al. 2018).

Strikingly, analyses of gene expression and/or sequence variation revealed two putative fatty-

acid elongases (Szafer-Glusman et al. 2008, Gaudet et al. 2011)—CG17821 and CG18609—with patterns

of variation associated with species differences. Both could functionally contribute to our observed

variation between longer and shorter CHC profiles, as alleles at both loci are shared between our

western *D. a. americana* and *D. novamexicana*, differ from our *D. a. texana* line and the eastern *D. a.*

americana Anderson stock, and are therefore associated with the "short" versus "long" CHC

phenotypes. The variation we observed at CG17821 is particularly interesting, because we observe a

thymine insertion mutation that results in premature truncation of CG17821 specifically in *D.*

novamexicana and our western *D. a. americana* (Figure 3) and because our gene expression data also

indicates reduced expression of the CG17821 allele specifically in these lines. In the latter case, while we

might expect gene expression differences to be tissue-specific (that is, observed specifically in the

oenocytes: the CHC-producing organs), our data indicate that the species-specific expression signal at

this locus is strong enough to detect from whole-body transcriptome data. Both observed sequence and

expression changes at CG17821 could individually produce a greater abundance of short CHC products,

by reducing elongase functionality: gene expression changes could result in a lower abundance of

elongase proteins, while the truncation insertion mutation could reduce the efficiency or activity of the

resulting elongase enzymes. Therefore, although our current data cannot differentiate which change

may have occurred first (or whether they are pleiotropic), either change could produce the pattern of

phenotypic difference observed between species. Overall, these data suggest allelic variation in

CG17821 could be responsible for the major axis of CHC divergence observed between *D. a. texana* and

the other two species, consistent with a hypothesis of a simple underlying basis for chain length

variation. Moreover, because this axis of CHC divergence appears primarily responsible for sexual

isolation between *D. novamexicana* and *D. a. texana*, this points to a large role for this simple allelic

change in CG17821 in the emergence of a strong isolating barrier between these two species.

Complex patterns of isolation and multimodality of sexual signals

While our analysis indicates that sexual isolation between *D. novamexicana* males and *D. a*

texana females could be due to a single trait change of large effect, it also reveals that isolation in other

species combinations is likely influenced by more complex factors. This is particularly evident for cross

combinations involving our *D. a. americana* and *D. a. texana* lines. Despite having a similar "short" CHC

profile to *D. novamexicana*, western *D. a. americana* has more modest and bi-directional isolation from

D. a. texana, including more frequent acceptance of *D. a. americana* males by *D. a. texana* females

(compared to the complete rejection of *D. novamexicana* males). Furthermore, perfuming of *D. a.*

americana males did not alter mating rate with *D. a. texana* females. It is possible that these more

complex patterns are due to the additional contribution of other sensory signals to sexual interactions

between *D. a. texana* and *D. a. americana*, including via the two other main sensory modalities used in

insect courtship—visual and auditory signals (reviewed Bontonou and Wicker-Thomas 2014, Jezovit et

al. 2017). Nonetheless, it is unlikely that auditory signals make a major contribution here, as song

differences between the *D. americana* group species are much smaller than those between other

subgroups in the virilis clade, and these differences have been shown not to influence female choice

(Hoikkala and Lumme 1987). Similarly, unlike other *Drosophila* groups (e.g., Fuyama 1979, Massey et al.

2020), *D. americana* group males spend relatively little time during courtship on visual cues such as wing

displays (Spieth 1951, personal observation), and visual display intensity did not appear to influence

mating success in our trials (Table S5). Overall, these data suggest that chemical communication is the

primary means of sexual signaling in this subgroup. This would explain why a shift in CHCs has such a

large effect on mating in the *D. novamexicana-D. a. texana* pair, but also suggests that more subtle

variations in chemical signals might also be responsible for more complex isolation patterns involving *D.*

a. americana, especially between *D. a. americana* and *D. a. texana*. This possibility is supported by our

findings that, in addition to the primary axis of CHC variation between species, *D. a. americana* also

shows male specific divergence on the U-PC2 axis (Figure 2). This axis is loaded heavily for the male-

specific compound C21:1—the only major compound with species-specificity that differentiates *D. a.*

americana from both *D. novamexicana* and *D. a. texana*—and *D. a. americana* has significantly less

C21:1 than the other two species (Results). Lower C21:1 abundance might be critical for the persistence

of sexual isolation even after *D. a. americana* males are perfumed by *D. a. texana*, if *D. a. texana*

females require high abundance of C21:1 in males as a stimulus for mating, but transfer of C21:1 during

perfuming is insufficient. Indeed, CHC quantification indicates that heterospecific-perfumed *D. a.*

americana males do not differ from conspecific-perfumed males, suggesting incomplete transfer of

compounds may play a role. This preference would not impact mating patterns with *D. novamexicana*

males—who have a high abundance of C21:1 regardless of perfuming—but could explain continued

isolation from *D. a. americana* males even after perfuming manipulation. Therefore, in addition to *D. a.*

texana females discriminating against "short" CHC phenotypes, our data also indicate they might use

C21:1 as a threshold trait for assessment of male quality or identity.

Lastly, the expression of sexual isolation in some of our other cross combinations might also be

modulated by additional complex effects on the order and timing of courtship behaviors (Ruedi and

Hughes 2008, Gaertner et al. 2015). For instance, we observed that female *D. a. americana* accept fewer

D. novamexicana males in 1x1 trials (20%) compared to 4x4 (100%), suggesting the density of flies

during these mating opportunities might influence behaviors of males or females, with consequences

for mating rates. Male-male interactions may subtly alter courtship patterns in *Drosophila* (Baxter et al.

2018), for example, so that courtship intensity differences between trials in which single or multiple

males were present might explain these more complex *D. a. americana* mating patterns. Regardless, it is

clear that the primary signal modality—and major cause of sexual isolation—between *D. a. texana*

females and *D. novamexicana* males is female choice on the basis of male pheromone chemistry.

The evolutionary history of CHC divergence and consequences for past and future sexual isolation in this

group

Together, our data demonstrate sexual isolation between *D. a. texana* and *D. novamexicana* is

due to CHC divergence in compound length*,* and suggest that variation between truncated and non-

truncated alleles of the elongase CG17821 contribute to this phenotypic variation. Moreover, both

genome-wide SNP variation, as well as localized variation specifically around this this putative elongase

gene (Figure 3C), indicate that this phenotypic variation involves a history of introgression. These data

point to a model for the evolutionary history of transitions involved in the change in CHC profiles

between species and, potentially, in the emergence and expression of sexual isolation that depends

upon this phenotype.

First, the distribution of both "long" versus "short" CHC phenotypes (shown here and in Lamb et

al. 2020), and allelic variation in CG17821, indicate that the "short" phenotype and the CG17821

truncated allele are derived states that arose within the *D. americana* group. This shift most likely

occurred in western lineages that gave rise to contemporary *D. novamexicana*. The evolutionary forces

responsible for the persistence and spread of this phenotype are not yet known. The relationship

between variation in environmental factors, insect stress physiology, and features of CHC length and

branching is known to be complex. Prior work has associated aspects of CHC divergence with abiotic

variation such as latitude (Frentiu and Chenoweth 2010, Rajpurohit et al. 2017, but see Gibbs et al.

2003), and physiological traits such as desiccation resistance (reviewed Chung and Carroll 2015), that

suggest a role for natural selection in shaping CHC composition, but the specific sources(s) of selection

can be challenging to pinpoint. In the *D. americana* group, species habitats are differentiated primarily

on the basis of water availability, and species differ in key physiological traits such as desiccation

resistance (Davis and Moyle 2019). These ecological and physiological differences, combined with

evidence of elevated nonsynonymous molecular evolution at CG17821 across the whole *D. virilis*

subclade (Table S6), suggest there could be strong selection acting on CHC composition in this group;

this selection may have contributed to the emergence and persistence of the novel "short" phenotype in

the western distribution of the *D. americana* species group.

Based on our data associating phenotypic divergence with sexual isolation, the appearance of

this new CHC phenotype would have reduced sexual compatibility between derived "short" males and

females with strong preferences for the "long" ancestral CHC profile. Persistence of this phenotype

would therefore have required a broadening of female preference to accommodate males with "short"

CHC phenotypes, perhaps via selection of rare females that are already responsive to the new

pheromone blend (e.g., as observed in male *Ostrinia* moths; Roelofs et al. 2002). Our data support this

expectation, as *D. novamexicana* females are more accepting of both (putatively derived) *D.*

novamexicana and (ancestral) *D. a. texana* male CHC phenotypes, while *D. a. texana* females

discriminate against derived "short" CHC phenotypes. Interestingly, this model for the evolution of

asymmetric sexual isolation is broadly consistent with Kaneshiro's (1976, 1980) model for peripatric

speciation. Kaneshiro observed that females from derived populations frequently have broad

preferences for both derived and ancestral male phenotypes; he proposed that this was due to relaxed

selection on narrow female preferences in genetically bottlenecked island populations, where founder

effects have led to the loss of elements of male courtship. Although Kaneshiro's model explicitly invokes

genetic drift in male trait evolution, our observations indicate his explanation of sexual isolation

asymmetry could extend more generally to all cases where evolutionary change in male sexual traits

precipitates broadening of female preferences for them in derived populations. Regardless of the

evolutionary force that might have driven the initial trait change, this process results in the emergence

of strong premating asymmetry specifically between females with ancestral trait preferences and males

with derived trait values.

526   Intriguingly, our data also support the subsequent movement of this CHC phenotype from *D.*

novamexicana into western *D. a. americana* lineages. Although shared variation in CG17821 and CHC

phenotypes could have been due to segregating ancestral variation present in both *D. a. americana* and

D. novamexicana (that is, ILS)*,* our data strongly support the alternative hypothesis of more recent

introgression of this trait from *D. novamexicana* into western *D. a. americana* populations. Both their

significant excess of shared genome-wide variation and a genomic region of at least 18 kB of shared

recent ancestry surrounding CG17821, is consistent with introgression between *D. novamexicana* and

western *D. a. americana*, that was accompanied by a similar shift to the "short" CHC phenotype in

western *D. a. americana*.

Given the effect of the "short" CHC phenotype on sexual isolation between *D. novamexicana*

and *D. a texana*, this introgression is expected to have consequences for reproductive isolation involving

recipient *D. a. americana* lineages. Interestingly, our data indicate that current patterns of sexual

isolation differ from those seen in *D. novamexicana*, even though western *D. a. americana* shows the

general shift to shorter chain CHC length associated with the introgressed region. One explanation might

be that this introgression-mediated shift occurred on a novel *D. a. americana* genomic background that

has additional variation in other aspects of CHC composition (such as the abundance of C21:1, see

above) and/or different patterns and strengths of female preference. A concrete example of these

background effects can be seen for (putatively) introgressed locus CG18609, where our western *D. a.*

americana shares the *D. novamexicana* allele at this locus but not its patterns of gene expression.

Instead, this locus has significantly reduced expression in *D. a. americana* compared to the other two

lineages (Figure 3A). These and similar effects of genomic background could be responsible for more

complex mating relationships seen in the *D. a. americana* line examined here, that might also extend to

patterns more broadly among *D. a. americana* populations depending on their individual histories of

divergence and introgression. Recent work (Sramkowski et al. 2020) has described rare *D.*

novamexicana-like pigmentation alleles in geographically disparate *D. a. americana* populations,

similarly suggesting past gene exchange into diverse genomic backgrounds. If alleles for specific sexual

signals, behaviors, and/or preferences are being differentially exchanged throughout the range of *D.*

americana, we could expect to observe variable levels of mating success between disparate populations.

Indeed, 4x4 mating trials performed by Spieth (1951) support this expectation, by demonstrating up to

10-fold variation in mating success among seven populations of *D. a. americana* from disparate

geographic origins. These lines of evidence suggest that trait variation arising from introgression of a

locus with major effects on a signaling phenotype—such as CG17821—could initiate the further

emergence of reproductive isolating barriers between populations within the broad and diverse *D. a.*

americana subspecies. This expectation is testable with work examining mating success between

geographically diverse *D. americana* populations—particularly between eastern and western

populations—and its covariation with CHC phenotypic and genotypic variation.

562   Regardless of the collateral consequences for isolation within *D. a. americana*, our data clearly

support the role of divergent cuticular hydrocarbon profiles—specifically a general shift in carbon chain

length—in sexual isolation between *D. novamexicana* and *D. a. texana*. They also implicate a causal role

for the gene CG17821 in determining CHC phenotype via a global change in CHC elongation activity early

in the generation of these compounds. These data provide a strong example of how a recently derived

allele in a single gene with large phenotypic effects on a sexual signal could underpin complete

asymmetric sexual isolation between closely related allopatric species. Moreover, they suggest that

subsequent introgression of this allele into genomic backgrounds of other closely related populations

could lead to additional emerging yet complex barriers to reproduction.

Methods

Experimental Fly stocks

One stock from each of three species was obtained from the University of California San Diego

Drosophila Species Stock Center (DSSC): a *Drosophila novamexicana* stock from San Antonio, NM

(15010-1031.08); a *D. americana americana* stock from Chadron, NE (15010-0951.06); and, a *D.*

americana texana stock from Jamestown, SC, (15010-1041.29). All stocks were originally collected

between 1946 and 1950. *D. americana* is divided into subspecies according to presence of a

chromosomal fusion of the X- and 4-chromosomes in *D. a. americana* that shows a distinct latitudinal

cline (McAllister 2002). For simplicity we refer to them by their subspecies names. All fly stocks were

reared on standard cornmeal media prepared by the Bloomington Drosophila Stock Center (BDSC) at

Indiana University, and were kept at room temperature (~22 °C). Every assay in this study used virgins

isolated within 8 hours of eclosion and aged for 7 days prior to the start of experiments.

4x4 unperfumed mating assay

We performed trials in which four virgin males and four virgin females were paired and

observed for mating behavior, following the design used by Speith (1951) which allows for behavioral

interactions that might not otherwise be observed in single-pair no-choice assays. Within each trial, all

males are from a single subspecies, as are all females; crosses are varied by pairing males and females of

alternative lines. For each trial, 4 males and 4 females were transferred to a single vial without

anesthetization and observed for 3 hours. The number and duration of each copulation event was

recorded for each trial. This assay was repeated for a total of 5 replicates for each possible species

combination, in reciprocal (i.e., 9 cross types, each of N = 5). Each trial used 7-day old virgins and testing

was started within 30 minutes of lights on in the morning.

Courtship behavior assay

To quantify and evaluate differences in courtship behaviors between cross types, flies were

observed in single pair (no-choice) mating assays. We used a modified FlyPi setup—that combines a

Raspberry Pi (Raspberry Pi Foundation, Cambridge, UK), pi camera, and 3D printed parts (Chagas et al.

2017)—to record courtship behaviors. Assays were performed in a modified cell culture plate consisting

of six 3 cm-diameter culture wells, each with a small amount of cornmeal media in the bottom, that

allowed six total crosses to be recorded simultaneously. For each assay, individual virgin male and

female flies of a given cross were aspirated without anesthetization to a cell culture well; after 6 total

crosses were set up, the plate was videotaped for a contiguous 3-hour period. The six cross

combinations assessed in any particular video trial were randomized to account for variance that might

otherwise be explained by date. As in the 4x4 mating assay, we performed 5 replicates of all possible

species combinations in reciprocal.

Behavioral features were analyzed and scored manually by the same individual to avoid

subjective variation among researchers. Three courtship behaviors—male display events, male tapping

events, and male licking events—in addition to copulation were scored in each 1x1 trial using the

following criteria. A male "display" was counted when a male performed a combination of back and

forth movements and occasional wing flicks while maintaining sustained orientation in front of and

facing the female. "Tapping" events were defined when a male used his tarsus to touch the abdomen of

the female when oriented behind her. "Licking" events were defined when the male's mouthparts

contacted the female genital arch. For male display, tapping, and licking events, individual events were

scored separately only if a different behavior (including sitting still/walking away) was observed

between instances of the defined behavior. This criterion minimized overcounting of discrete behavioral

events, especially those with difficult to view aspects such as number of times a male extruded

mouthparts during a contiguous licking event. In addition to these behaviors, copulation success and

latency to copulation were also recorded for each trial. Note that females also engage in tapping and

other behaviors, but because these appear to be less consistent and are more difficult to observe and

score, they were not addressed in this study.

Extraction and quantification of cuticular hydrocarbons

Cuticular hydrocarbons were extracted from pooled samples by placing five 7-day old virgin flies

of a single sex and species identity in a 1.8 mL glass vial (Wheaton 224740 E-C Clear Glass Sample Vials)

with 120 µL of hexane (Sigma Aldrich, St Louis, MO, USA) spiked with 10 µg/mL of hexacosane (Sigma

Aldrich). After 20 minutes, 100 µL of the solution was removed to a sterilized 1.8 mL glass vial (Wheaton

224740 E-C Clear Glass Sample Vials) and allowed to evaporate overnight under a fume hood. Extracts

were stored at -20 °C until analysis. Five replicate samples consisting of 5 flies per sample (25 flies total)

were prepared for each sample type, and all replicates were extracted on the same day.

Gas chromatography mass spectrometry (GC/MS) analysis was performed on a 7820A GC

system equipped with a 5975 Mass Selective Detector (Agilent Technologies, Inc., Santa Clara, CA, USA)

and a HP-5ms column ((5%-Phenyl)-methylpolysiloxane, 30 m length, 250 µm ID, 0.25 µm film thickness;

Agilent Technologies, Inc.). Electron ionization (EI) energy was set at 70 eV. One microliter of the sample

was injected in splitless mode and analyzed with helium flow at 1 mL/ min. Two different temperature

gradients were used depending on the sample type. For CHC analysis of unmanipulated males and

females of each species, the following parameters were used: column was set at 50 °C for 0 min,

increased to 210 °C at a rate of 35 °C/min, then increased to 280 °C at a rate of 3 °C/min. The MS was set

to detect from m/z 33 to 500. For analysis of samples from perfuming trials, the parameters were

modified to increase resolution and sensitivity for less abundant compounds: the column was set at 40

°C and held for 3 min, increased to 200 °C at a rate of 35 °C/min, then increased to 280 °C at a rate of 3

°C/min and held for 15 minutes. Chromatograms and spectra were analyzed using MSD ChemStation

(Agilent Technologies, Inc.). CHCs were identified on the basis of retention time and electron ionization

fragmentation pattern. Compounds are identified in this study as: CXX:Y for alkenes or Me-CXX for

methyl-branched alkanes, where XX indicates the length of the carbon chain, and Y indicates number of

double bonds, e.g. C21:1 is a 21-carbon alkene with a single double bond.

The abundance of each compound was quantified by normalizing the area under each CHC peak

to the area of the hexacosane signal using homebuilt peak selection software (personal correspondence,

Dr. Scott Pletcher, Univ. of Michigan).

Perfuming manipulation and mating assay

'Perfuming' involves co-housing target flies in vials filled predominantly with flies of a desired

donor identity, so that the CHC profile of the target flies is altered via physical transfer of CHCs from

donor flies (Coyne et al. 1994, Dyer et al. 2014, Serrato-Capuchina et al. 2020). Perfuming was

performed by placing 2 'target' males with 15 'donor' males within a single vial. All males were 1-day old

virgins when perfuming vials were established, and the wings of donor males were removed (under

anesthetic) to distinguish them from target males. For the heterospecific perfuming treatments, target

males were co-housed with donor males of a different species; conspecific (control) treatments paired

donor and target males of the same species. Following 7 days of perfuming, 4 male flies from the same

perfuming conditions were transferred (without anesthetization) to vials containing 4 female *D. a.*

texana. Within each experiment (Nov-Tex, or Ame-Tex), we ran one trial of each perfuming condition (2

heterospecific-perfumed types, and 2 conspecific-perfumed types) in parallel on the same day (four 4x4

trials in total).

For CHC analysis, males were perfumed as described for the perfumed mating assay, for both

Nov-Tex and Ame-Tex experimental pairs, with the exception that 2-3 target males (rather than just 2)

were co-housed with 15-18 donor males, enabling 2 parallel perfuming vials to generate 5 target males

flies per identity.

Candidate gene selection

Our candidate list was generated by searching Flybase (Flybase.org) for annotated *Drosophila*

melanogaster genes with protein-coding domains (as identified by InterProt) that have known functions

in CHC synthesis (Pardy et al. 2018), resulting in an initial list of 34 genes. To this we added the

pigmentation genes *ebony* and *tan* as they have been shown to alter CHC variation among *D. americana*

group species (Lamb et al. 2020, Massey et al. 2019). With these 36 genes, we identified orthologs in

Drosophila virilis (as annotated in Flybase) and evaluated whether each of these loci had transcript

expression in our three species, using previously published whole-body transcriptome data generated

from the same three populations using RNA-seq (Davis and Moyle 2020). Our analyses used only gene

expression data from the control (ambient) conditions for both males and females from this study, and

excluded any desiccation stress treatment data. Of the 36 initial genes, 30 were found to have

unambiguous 1-to-1 orthologs in *D. virilis*, and of these, 23 had transcripts present within the Davis and

Moyle (2020) dataset. This final set of 23 genes was used to evaluate gene expression and sequence

variation among our three lines. For downstream analyses, we also identified alleles of these 23 loci in a

genome assembly of *D. a. americana* Anderson (15010-0951.00, also referred to as A01) generated by

Kim et al. (unpubl.), using the BLASTn function of BLAST+ version 2.6.0 (Camacho et al. 2009) and

retaining the top BLAST hits for each locus.

Statistical Analyses

All statistical analyses and figure construction in this study was performed with R version 3.4.3.

For the unperfumed 4x4 mating dataset, we used Kruskal-Wallis tests to compare copulation success

within 3 hours of observation with male or female species identity (one test for each sex). In addition,

we used Wilcoxon rank sum tests to perform post-hoc pairwise comparisons for each sex to evaluate

which species most differed from one another in female acceptance of males or in male courtship

success of females.

In 1x1 mating trials, mating rate was evaluated using a logistic regression with a chi-square test.

For copulation latency, the effects of male identity, female identity, and their interaction were evaluated

with an ANOVA, followed by a post-hoc Tukey HSD in which we assessed which male identities (species)

were specifically different for this trait. For each of the three courtship behaviors (displays, tappings and

lickings), we converted the count data for each trial to a rate per unit time within a trial, by dividing the

recorded counts for each trait by the latency to copulation (in minutes) or, if no copulation occurred, the

maximum amount of observed time (180 minutes). Describing male behaviors in terms of rates accounts

for differences among males in their courtship efficiency; for example, it allows us to differentiate males

that performed fewer courtship behaviors because they were rapidly accepted by a female, from males

that displayed lower courtship intensity across the total 3 hour monitored period. Differences in male

courtship behaviors were evaluated with ANOVA. Correlation of individual courtship behaviors with

copulation success was analyzed using a logistic regression (with z-test).

The perfumed 4x4 mating experiment was analyzed using planned contrasts that compared the

mating success of males that had the same target identity but different (con- versus hetero-specific)

donor perfumes, when paired with *D. a. texana* females. This enabled us to specifically assess the effects

of donor perfume variation on mating success of a given target male species. Each pairwise test was

performed using a non-parametric Mann-Whitney U test.

Our primary analyses of differences in CHC composition were done after summarizing major

axes of variation in CHC phenotypes using a set of Principal Component Analyses (PCA). Separate PCAs

were performed on each CHC experiment within this study: one for the initial (unperfumed) parental

populations (denoted U-PCA) that included both male and female samples, and one each for the

perfumed experimental pairs—Nov-Tex (N-PCA) and Ame-Tex (A-PCA)—with male samples only. Within

each dataset, a one-way ANOVA was used on each of the first 3 PCs to examine effects of sex and

species (for the unperfumed dataset) or male perfume identity (in each perfuming study) on CHC

composition. Additionally, for the perfumed datasets, T-tests were used to compare differences in PC

values between target males that were paired with different (con- and hetero-specific) donor males to

assess the effects of our perfuming manipulation on major axes of CHC variation. Factor loadings for N-

PC an A-PC analyses, as well as individual compound differences in perfuming pairs can be found in in

the supplement (Tables S7, S8, S9 and Figures S1, S2).

For all gene expression analyses, expression is quantified in transcripts per million (TPM), and

therefore is normalized within each sample. Here we analyzed datasets separated by sex, as Davis and

Moyle (2020) showed that the majority of genes have differential expression based on sex, whereas our

primary interest here is in differences between species that might be implicated in female mating

choices and male CHC profile variation. With the dataset for each sex, we ran one-way ANOVAs on every

expressed locus (11301 genes total)—including our 23 target candidate genes—in order to determine if

gene expression varied by species and to rank order each gene according to their resulting *P*-value. This

allowed us to interpret patterns of species variation specifically among candidate genes in the context of

species differences across all genes in the transcriptome.

Using the same dataset for our 23 candidate genes, we also evaluated patterns of molecular

evolution by calculating clade-wide d_N/d_S values for each gene. For each candidate gene, sequences

from each species were aligned to the *D. virilis* reference genome (Flybase version 1.7, www.flybase.org)

and aligned sequences analyzed in PAML v4.9 (Yang 2007) using the base model in CodeML. CodeML is a

maximum likelihood model for codon substitution that estimates the rates of nonsynonymous to

synonymous substitutions (d_N/d_S) to test for positive selection on a given gene between species. Gene

trees for each candidate gene were constructed in ClustalW (Thompson et al. 1994), as PAML uses a

tree-based model for computing d_N/d_S that is sensitive to use of the correct gene tree for a given gene.

Results of these analyses are reported in the supplement (Table S6).

We constructed additional gene trees for each candidate gene using orthologs from the

Anderson IN *D. a. americana* genome (from Kim et al. unpublished) in place of our (western) *D. a.*

americana sequences, using alignments of this sample with the other species generated from ClustalW

(Thompson et al. 1994). Additional gene trees were generated in the same manner for neighboring

genes in the region around CG17821/CG18609, to estimate size of a window of shared ancestry.

To assess evidence for genome-wide introgression, we used our transcriptome data in

conjunction with the *D. virilis* genome to generate a genome-wide set of SNPs for our three focal

populations and the *D. virilis* outgroup. To do so, we used bcftools (Li 2011) to generate a VCF (variant

calling format) file and call SNP sites between the 4 taxa after filtering for low depth and masking

heterozygous sites from individual taxa. With this, we calculated Patterson's *D*-statistic (Durand et al.

2011) using the Dtrios program from Dsuite (Malinsky et al. 2020) to calculate the 3 taxa *D*-statistic as

well as an overall *P*-value using standard errors generated from the default 20 jackknife blocks.

Acknowledgements

We would like to thank Matthew Gibson for help with implementing molecular evolution analyses, Zinan

Wang for advice about candidate gene selection, Bernard Kim for sequencing the *Drosophila americana*

genome and providing assemblies, Scott Pletcher for sharing hydrocarbon analysis software, and

Jonathan Massey for connecting JSD and LCM with JYY.

Funding

This work was supported by the IU Department of Biology (LCM, JSD), the Department of Defense

United States Army Research Office (Grant No. W911NF1610216) and the National Institutes of Health

(Grant No. 1P20GM125508) awarded to JYY. The GC/ MS analysis was performed in the UHM Microbial

Genetics and Analytical Laboratory (supported by NIH Grant No. 1P20GM125508).

References

Bartelt, R. J., Armold, M. T., Schaner, A. M., & Jackson, L. L. (1986). Comparative Analysis of Cuticular
Hydrocarbons in the Drosophila Virilis Species Group. *Comparative Biochemistry and Physiology*,
83B(4), 731–742.

Basolo, A. L. (1995). Phylogenetic evidence for the role of a pre-existing bias in sexual selection.
Proceedings. Biological Sciences, *259*(1356), 307–311.

Baxter, C., Mentlik, J., Shams, I., & Dukas, R. (2018). Mating success in fruit flies: courtship interference
versus female choice. *Animal Behaviour*, *138*, 101–108.

Billeter, J.C., Atallah, J., Krupp, J. J., Millar, J. G., & Levine, J. D. (2009). Specialized cells tag sexual and
species identity in Drosophila melanogaster. *Nature*, *461*(7266), 987–991.

Blomquist G. J., Bagnères A. G. (2010) Insect hydrocarbons: biology. biochemistry, and chemical ecology.
Cambridge University Press, New York

Bontonou, G., & Wicker-Thomas, C. (2014). Sexual Communication in the Drosophila Genus. *Insects*,
5(2), 439–458.

Butlin, R., Debelle, A., Kerth, C., Snook, R. R., Beukeboom, L. W., Castillo Cajas, R. F., Diao, W., Maan, M.
E., Paolucci, S., Weissing, F. J., van de Zande, L., Hoikkala, A., Geuverink, E., Jennings, J., Kankare,
M., Knott, K. E., Tyukmaeva, V. I., Zoumadakis, C., Ritchie, M. G., … Schilthuizen, M. (2012). What
do we need to know about speciation? *Trends in Ecology & Evolution*, *27*(1), 27–39.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009).
BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*, 1–9.

Chung, H., & Carroll, S. B. (2015). Wax, sex and the origin of species: Dual roles of insect cuticular
hydrocarbons in adaptation and mating. *BioEssays*, 822–830.

Chung, H., Loehlin, D., Dufour, H., Vaccarro, K., Millar, J. G., & Carroll, S. B. (2014). A Single Gene Affects
Both Ecological Divergence and Mate Choice in Drosophila. *Science*, *257082*.

Coyne, J. A., Kim, S. Y., Chang, A. S., Lachaise, D., & Elwyn, S. (2002). Sexual isolation between two sibling
species with overlapping ranges: Drosophila santomea and Drosophila yakuba. *Evolution*, *56*(12),
2424–2434.

Coyne, J. A., & Oyama, R. (1995). Localization of pheromonal sexual dimorphism in Drosophila
melanogaster and its effect on sexual isolation. *Proceedings of the National Academy of Sciences of*
the United States of America, *92*(21), 9505–9509.

Coyne, J. A., & Orr, H. A., 2004. Speciation. SinauerAssociates, Sunderland, MA

Coyne, J. A., Crittenden, A. P., & Maht, K. (1994). Genetics of a Pheromonal Difference Contributing to
Reproductive Isolation in Drosophila. *Science*, *265*(5177), 1461–1464.

Curtis, S., Sztepanacz, J. L., White, B. E., Dyer, K. a., Rundle, H. D., & Mayer, P. (2013). Epicuticular
Compounds of Drosophila subquinaria and D. recens: Identification, Quantification, and Their Role
in Female Mate Choice. *Journal of Chemical Ecology*, *39*(5), 579–590.

Davis, J. S., & Moyle, L. C. (2020). Constitutive and plastic gene expression variation associated with
desiccation resistance differences in the drosophila Americana species group. *Genes*, *11*(146).

Davis, J. S., & Moyle, L. C. (2019). Desiccation resistance and pigmentation variation reflects bioclimatic
differences in the Drosophila americana species complex. *BMC Evolutionary Biology*, *19*(1), 1–14.

Durand, E. Y., Patterson, N., Reich, D., & Slatkin, M. (2011). Testing for ancient admixture between
closely related populations. *Molecular Biology and Evolution*, *28*(8), 2239–2252.

Dyer, K. a, White, B. E., Sztepanacz, J. L., Bewick, E. R., & Rundle, H. D. (2014). Reproductive character
displacement of epicuticular compounds and their contribution to mate choice in Drosophila
subquinaria and Drosophila recens. *Evolution; International Journal of Organic Evolution*, *68*, 1163–
1175.

Franssen, S. U., Nolte, V., Tobler, R., & Schlotterer, C. (2015). Patterns of linkage disequilibrium and long
range hitchhiking in evolving experimental drosophila melanogaster populations. *Molecular
Biology and Evolution*, *32*(2), 495–509.

Frentiu, F. D., & Chenoweth, S. F. (2010). Clines in cuticular hydrocarbons in two *Drosophila* species with
independent population histories. *Evolution*, 64(6), 1784–1794.

Gaertner, B. E., Ruedi, E. A., McCoy, L. J., Moore, J. M., Wolfner, M. F., & Mackay, T. F. C. (2015).
Heritable variation in courtship patterns in drosophila melanogaster. *G3: Genes, Genomes,
Genetics*, *5*(4), 531–539.

Gaudet, P., Livstone, M. S., Lewis, S. E., & Thomas, P. D. (2011). Phylogenetic-based propagation of
functional annotations within the Gene Ontology consortium. *Briefings in Bioinformatics*, *12*(5),
449–462.

Gibbs, A. G., Fukuzato, F., & Matzkin, L. M. (2003). Evolution of water conservation mechanisms in
Drosophila. *The Journal of Experimental Biology*, 206, 1183–1192.

Havens, J. A., & Etges, W. J. (2013). Premating isolation is determined by larval rearing substrates in
cactophilic *Drosophila mojavensis*. IX. Host plant and population specific epicuticular hydrocarbon
expression influences mate choice and sexual selection. *Journal of Evolutionary Biology*, 26(3),
562–576.

Hoikkala, A., & Lumme, J. (1987). The Genetic Basis of Evolution of the Male Courtship Sounds in the
Drosophila virilis Group. *Evolution*, *41*(4), 827–845.

Hollocher, H., Ting, C. T., Wu, M. L., & Wu, C. I. (1997). Incipient speciation by sexual isolation in
Drosophila melanogaster: Extensive genetic divergence without reinforcement. *Genetics*, *147*(3),
1191–1201.

Jezovit JA, Levine JD, Schneider J. Phylogeny, environment and sexual communication across the
Drosophila genus. *J Exp Biol*. 2017;220(Pt 1):42-52.

Kaneshiro, K. Y. (1980). Sexual Isolation, Speciation and the Direction of Evolution. *Evolution*, *34*(3), 437–
444.

Kaneshiro, K. Y. (1976). Ethological Isolation and Phylogeny in the Planitibia Subgroup of Hawaiian
Drosophila. *Evolution*, *30*(4), 740.

Lamb, A. M., Wang, Z., Simmer, P., Chung, H., Patricia, J., States, U., Lansing, E., States, U., Biology, E.,
Program, B., Lansing, E., States, U., Biology, E., Program, B., Arbor, A., States, U., & Wittkopp, P. J.
(2020). ebony affects pigmentation divergence and cuticular hydrocarbons in *Drosophila*
americana and *D. novamexicana*. *Frontier in Ecology and Evolution*, *8*(June), 1–23.

Legendre, A., Miao, X. X., Da Lage, J. L., & Wicker-Thomas, C. (2008). Evolution of a desaturase involved
in female pheromonal cuticular hydrocarbon biosynthesis and courtship behavior in *Drosophila*.
Insect Biochemistry and Molecular Biology, *38*(2), 244–255.

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. *Bioinformatics*, *27*(21), 2987–
2993.

Malinsky, M., Matschiner, M., & Svardal, H. (2020). Dsuite - fast D-statistics and related admixture
evidence from VCF files. *BioRxiv*, 1–16.

Mas, F., & Jallon, J. M. (2005). Sexual isolation and cuticular hydrocarbon differences between
Drosophila santomea and *Drosophila yakuba*. *Journal of Chemical Ecology*, *31*(11), 2747–2752.

Massey, J. H., Rice, G. R., Firdaus, A. S., Chen, C. Y., Yeh, S. D., Stern, D. L., & Wittkopp, P. J. (2020). Co-
evolving wing spots and mating displays are genetically separable traits in *Drosophila*. *Evolution*,
74(6), 1098–1111.

Massey, J. H., Akiyama, N., Bien, T., Dreisewerd, K., Wittkopp, P. J., Yew, J. Y., & Takahashi, A. (2019).
Pleiotropic Effects of ebony and tan on Pigmentation and Cuticular Hydrocarbon Composition in
Drosophila melanogaster. *Frontiers in Physiology*, 10.

McAllister, B. F. (2002). Chromosomal and allelic variation in *Drosophila americana* : selective
maintenance of a chromosomal cline. *Genome*, *45*(1), 13–21.

Morales-Hojas, R., Reis, M., Vieira, C. P., & Vieira, J. (2011). Resolving the phylogenetic relationships and
evolutionary history of the *Drosophila virilis* group using multilocus data. *Molecular Phylogenetics*
and Evolution, *60*(2), 249–258.

Pardy, J. A., Rundle, H. D., Bernards, M. A., & Moehring, A. J. (2018). The genetic basis of female
pheromone differences between *Drosophila melanogaster* and *D. simulans*. *Heredity*, *122*(1), 93–
109.

Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T., Webster, T., &
Reich, D. (2012). Ancient admixture in human history. *Genetics*, *192*(3), 1065–1093.

Rajpurohit, S., Hanus, R., Vrkoslav, V., Behrman, E. L., Bergeland, A. O., Petrov, D. A., Cvacka, J., &
Schmidt, P. S. (2017). Adaptive dynamics of cuticular hydrocarbon in *Drosophila*. Journal of
Evolutionary Biology, 30(1), 66–80.

Ritchie, M. G. (2007). Sexual Selection and Speciation. *Annual Review of Ecology, Evolution, and*
Systematics, *38*(1), 79–102.

Roelofs, W. L., Liu, W., Hao, G., Jiao, H., Rooney, A. P., & Linn, C. E. (2002). Evolution of moth sex
pheromones via ancestral genes. *Proceedings of the National Academy of Sciences of the United*
States of America, 99(21), 13621–13626.

Ruedi, E. A., & Hughes, K. A. (2008). Natural genetic variation in complex mating behaviors of male
Drosophila melanogaster. *Behavior Genetics*, *38*(4), 424–436.

Schemske, D. W. (2000). UNDERSTANDING THE ORIGIN OF SPECIES. *Evolution*, *54*(3), 1069–1073.

Schluter, D. (2009). Evidence for Ecological Speciation and Its Alternative. *Science*, *323*(5915), 737–741.

Seddon, N., Botero, C. A., Tobias, J. A., Dunn, P. O., MacGregor, H. E. A., Rubenstein, D. R., Uy, J. A. C.,
Weir, J. T., Whittingham, L. A., & Safran, R. J. (2013). Sexual selection accelerates signal evolution
during speciation in birds. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1766).

Serrato-Capuchina, A., Schwochert, T. D., Zhang, S., & Roy, B. (2020). Pure species discriminate against
hybrids in the *Drosophila melanogaster* species subgroup. *bioRxiv*

Smadja, C., & Butlin, R. K. (2009). On the scent of speciation: The chemosensory system and its role in
premating isolation. *Heredity*, *102*(1), 77–97.

Spieth, H. T. (1951). Mating behavior and sexual isolation in the Drosophila virilis species group.
Behaviour, *3*(2), 105–145.

Sramkoski, L. L., Mclaughlin, W. N., Cooley, A. M., Yuan, D. C., & Wittkopp, P. J. (2020). Genetic
architecture of a body color cline in *Drosophila americana*. *BioRxiv*.

Szafer-Glusman, E., Giansanti, M. G., Nishihama, R., Bolival, B., Pringle, J., Gatti, M., & Fuller, M. T.
(2008). A Role for Very-Long-Chain Fatty Acids in Furrow Ingression during Cytokinesis in
Drosophila Spermatocytes. *Current Biology*, *18*(18), 1426–1431.

Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: Improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific gap
penalties and weight matrix choice. *Nucleic Acids Research*, *22*(22), 4673–4680.

Tomaru, M., Doi, M., Higuchi, H., & Oguma, Y. (2000). Courtship song recognition in the *Drosophila melanogaster* complex: Heterospecific songs make females receptive in *D. melanogaster*, but not in D. sechellia. *Evolution*, *54*(4), 1286–1294.

van Doorn, G. S., Edelaar, P., & Weissing, F. J. (2009). On the Origin of Species by Natural and Sexual Selection. *Science*, *581*(December), 1704–1708.

Wang, Z., Cui, Y., Song, L., & Fang, F. (2016). The hysteresis model of piezoelectric micro-positioning stage based on threshold optimization. *PLoS Biology*, *38*(3), 437–440.

Wilkins, M. R., Seddon, N., & Safran, R. J. (2013). Evolutionary divergence in acoustic signals: Causes and consequences. *Trends in Ecology and Evolution*, *28*(3), 156–166.

Wu, C. I., Hollocher, H., Begun, D. J., Aquadro, C. F., Xu, Y., & Wu, M. L. (1995). Sexual isolation in *Drosophila melanogaster*: A possible case of incipient speciation. *Proceedings of the National Academy of Sciences of the United States of America*, *92*(7), 2519–2523.

Yang, Z. (2007). PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution*, *24*(8), 1586–1591.

Yew, J. Y., & Chung, H. (2017). *Drosophila* as a holistic model for insect pheromone signaling and processing. *Current Opinion in Insect Science*, *24*, 15–20.

Yukilevich, R., & Peterson, E. K. (2019). The evolution of male and female mating preferences in *Drosophila* speciation. *Evolution*, 73(9), 1759–1773.

Yukilevich, R., & True, J. R. (2008). Incipient sexual isolation among cosmopolitan *Drosophila melanogaster* populations. *Evolution*, *62*(8), 2112–2121.