1　　A shift to shorter cuticular hydrocarbons accompanies sexual isolation among *Drosophila americana*
2　　　　　　　　　　　　　　　　　　　　　group populations

Jeremy S. Davis[1, 2]*, Matthew J. Pearcy[1], Joanne Y. Yew[3], Leonie C. Moyle[1]

[1]Dept. of Biology, Indiana University, Bloomington, IN, USA 47405

[2]Dept. of Biology, University of Kentucky, Lexington, KY, USA 40508

[3]Pacific Biosciences Research Center, University of Hawai'i at Mānoa, Honolulu, HI 96822

*Corresponding author: J.Davis@uky.edu

RRH: Sexual isolation and chemical signal evolution

Abstract

Because sensory signals often evolve rapidly, they could be instrumental in the emergence of

reproductive isolation between species. However, pinpointing their specific contribution to isolating

barriers, and the mechanisms underlying their divergence, remains challenging. Here we demonstrate

sexual isolation due to divergence in chemical signals between two populations of *Drosophila americana*

(SC and NE) and one population of *D. novamexicana,* and dissect its underlying phenotypic and genetic

mechanisms. Mating trials revealed strong sexual isolation between *Drosophila novamexicana* males

and SC *Drosophila americana* females*,* as well as more moderate bi-directional isolation between *D.*

americana populations. Mating behavior data indicates SC *D. americana* males have the highest

courtship efficiency and, unlike males of the other populations, are accepted by females of all species.

Quantification of cuticular hydrocarbon (CHC) profiles—chemosensory signals that are used for species

recognition and mate finding in *Drosophila*—shows that the SC *D. americana* population differs from the

other populations primarily on the basis of compound carbon chain-length. Moreover, manipulation of

male CHC composition via heterospecific perfuming—specifically perfuming *D. novamexicana* males

with SC *D. americana* males—abolishes their sexual isolation from these *D. americana* females. Of a set

of candidates, a single gene—elongase CG17821—had patterns of gene expression consistent with a

role in CHC differences between species. Sequence comparisons indicate *D. novamexicana* and our

Nebraska (NE) *D. americana* population share a derived CG17821 truncation mutation that could also

contribute to their shared "short" CHC phenotype. Together, these data suggest an evolutionary model

for the origin and spread of this allele and its consequences for CHC divergence and sexual isolation in

this group.

Introduction

Sensory signals can act as sexual cues that are critical for intraspecific mate evaluation and

reproductive success. Moreover, because they are frequently among the most rapidly evolving species

differences (Smadja and Butlin 2009, Seddon et al 2013, Wilkins et al 2013), divergence in sensory sexual

signals might often contribute to the earliest stages of reproductive isolation, in the form of prezygotic

barriers between lineages (Butlin et al 2012, Ritchie 2007). Among all such sexual signals, nowhere is the

diversity more evident than those acting as premating traits, including coloration and other visual

signals, acoustic signals, and complex pheromone blends. However, convincingly demonstrating the

connection between such sensory signal divergence and emerging reproductive isolation can be

challenging because it requires, first, identification and demonstration of the direct role of specific

signals mediating sexual isolation between species and, second, understanding the specific mechanistic

changes that have given rise to lineage differences in this signal.

Among insects, sexual signals draw on three primary sensory modalities—visual, auditory, and

chemosensory—and most species likely use a combination of all three modalities to identify potential

mates. Of these, many insect chemosensory signals primarily consist of cuticular hydrocarbons (CHCs)—

a broad group of carbon-chain compounds important for many essential functions including sex

pheromone signaling, but also environmental adaptation (Blomquist and Bagneres 2010, Chung and

Carrol 2015, Yew and Chung 2017). Unlike many auditory and visual cues, CHCs are produced and

received by both males and females, making them potentially important for both male and female mate

choice or species recognition. Indeed, studies on *Drosophila* chemical communication, particularly in the

melanogaster subgroup—have detected clear evidence for sexual isolation based on male choice of

female CHCs (Coyne et al. 1995, Billeter et al. 2009), while work in other species has demonstrated

female choice of species-specific male CHC profiles (Coyne et al. 2002, Mas and Jallon 2005, Curtis et al.

2013, Dyer et al. 2014). Elements of the biochemical pathways of CHC production and the underlying

genes are also relatively well understood in *D. melanogaster* (Pardy et al 2018). Accordingly, single

genes contributing to species-specific CHC differences have been implicated as causes of reproductive

isolation in several cases, including between *D. melanogaster* and *D. simulans* (desatF: Legendre et al.

2008) and *D. serrata* and *D. birchii* (mFAS: Chung et al. 2014). This framework provides an excellent

resource for identifying candidate genes for sensory sexual signal variation in other *Drosophila* systems.

Here we use the *Drosophila americana* group to investigate how species variation in CHC

profiles contributes to reproductive isolation, and to identify the underlying chemical and genetic

changes that may comprise this variation. This group includes two closely related (MRCA ~0.5 mya,

Morales-Hojas et al. 2008) species—*D. novamexicana* and *D. americana*—that occupy distinct

geographic and environmental habitats (Davis and Moyle 2019). *D. americana* is broadly distributed in

the United States from the east coast to the Rocky Mountains, and exhibits significant phenotypic and

genetic variation among populations throughout (e.g., Caletka and McAllister 2004, Davis and Moyle

2019), while *D. novamexicana* is localized to the arid southwestern US. Both species have been noted as

being associated with—and exclusively collected near—willow species of the genus *Salix*, though the

exact nature of this association is not known (Blight and Romano 1953, McAllister 2002). Classical

mating studies (Spieth 1951) have documented population-specific variation in reproductive isolation

between members of this group (see supplemental Table S1 and below). Prior analysis has also shown

qualitative differences in CHC composition between males (Bartelt at al. 1986), however the

contribution of hydrocarbons to premating isolation has not been assessed. With three populations

from this group, one *D. novamexicana* population and two *D. americana* populations—one southern

(South Carolina, hereafter SC) and one western (Nebraska, hereafter NE)—here we use a combination of

mating and behavioral studies, chemical analysis and manipulation, and gene expression and sequence

variation analyses to assess the role of CHCs in reproductive isolation. Together, our data suggest that

evolution of a male sexual signal—an overall shift in the relative abundance of longer versus shorter

cuticular hydrocarbons, due to novel mutation in an elongase gene—has produced complete premating

isolation between derived males and females from species that retain the ancestral trait and preference,

as proposed in classical models (Kaneshiro 1976, 1980) of the evolution of asymmetric sexual isolation.

Results

SC D. americana *females discriminate against heterospecific males in mating trials*

89   We found clear evidence of moderate to strong sexual isolation between SC *D. americana* and

the other two populations (Table 1). Mating rate (average proportion of females mated, in 4x4 mating

trials) ranged from very high in intrapopulation crosses and some interpopulation crosses, to <50% in

interpopulation combinations of female SC *D. americana* with each of the other species. Notably, *D.*

novamexicana males were never successful in mating with SC *D. americana* females, indicating strong

sexual isolation in this cross direction between these populations. In contrast, mating rate in the

reciprocal cross was 70%. The other pairing with mating rates at or below 50% involved both reciprocal

directions between *D. americana* populations. Across all cross combinations in the 4x4 mating trials, we

found that female identity (Kruskal-Wallis test: $\chi^2(2)$ = 8.99, *P* = 0.012), but not male identity (Kruskal-

Wallis test: $\chi^2(2)$ = 0.80, *P* = 0.67), significantly affected success. Post-hoc comparisons confirmed no

pairwise differences in male mating rate between any populations (Table S2). In contrast, the copulation

rate of SC *D. americana* females was significantly lower on average than both NE *D. americana* (*P-adj* =

0.025) and *D. novamexicana* (*P-adj* = 0.042), whereas NE *D. americana* and *D. novamexicana* females did

not differ (*P-adj* = 0.15). Together, these results indicate that SC *D. americana* females discriminate

against heteropopulation males more strongly than do females of the other two populations, with the

greatest discrimination against *D. novamexicana* males. This discrimination produces strongly

asymmetric sexual isolation between SC *D. americana* and *D. novamexicana*, while premating isolation

between the two *D. americana* populations is bi-directional and more moderate. Importantly, this

strong asymmetric isolation reiterates results documented by Spieth (1951, Table S1), using similar 4x4

mating experiments with 2 *D. novamexicana* and 9 *D. americana* lines—including the NE *D. americana*

and *D. novamexicana* populations used here. In this prior analysis, males from *D. novamexicana* showed

little to no mating success with southeastern *D. americana* populations (previously referred to as *D. a.*

texana) (0-12.5% mating frequency; Table S1), while *D. novamexicana* females were more receptive in

the reciprocal cross (mating frequency of 46.7-86.5%; Table S1). The consistency of these results 70

years apart suggests isolation observed here is not a bi-product of long-term laboratory culture, but

instead reflects differences present in the natural populations from which these lines were collected.

SC D. americana *males have greater courtship efficiency and mating success*

We video-recorded single-pair matings for each cross combination to evaluate population

differences in courtship strategies and whether these differed in interpopulation pairings. Three male

courtship behaviors were quantified from these 1x1 trials—display rate, tapping rate, and licking rate.

All three behaviors showed similar patterns of among-population variation, when each was evaluated

for the effect of male population, female population, or their interaction, using 2-way ANOVAs. Males

significantly differed with respect to display-rate ($F(2, 36) = 3.76$, $P = 0.03$) and tap-rate ($F(2, 36) = 4.76$,

$P = 0.017$), but only marginally for lick-rate ($F(2, 36) = 2.67$, $P = 0.083$). For all three behaviors, this

difference is due to SC *D. americana* males exhibiting higher rates (Tukey HSD post-hoc tests; Figure 1,

supplementary Table S3). In contrast, we detected no female identity effects on male behavioral rate

(display-rate: $F(2, 42) = 1.83$, $P = 0.18$; tap-rate $F(2, 36) = 0.58$, $P = 0. 56$; lick-rate: $F(2, 36) = 0.43$, $P =$

0.65), or any female x male interactions (display-rate: $F(2, 36) = 0.72$, $P = 0.58$; tap-rate $F(2, 36) = 0.45$, P

$= 0.77$; lick-rate: $F(2, 36) = 0.65$, $P = 0.63$). Because behavior rates are dependent on copulation latency,

and not simply the total number of behavioral events, these rates approximate the efficiency with which

each courtship behavior results in a mating. Taken together, our results indicate that SC *D. americana*

males have greater courtship efficiency (in terms of display and tap rates), and that males do not

significantly vary these behavioral rates depending on female population identity.

The single pair mating assays also broadly reiterated the patterns of moderate to strong sexual

isolation we observed for SC *D. americana* in 4x4 trials. As with our 4x4 mating assay, *D. novamexicana*

males never mated with SC *D. americana* females when paired individually, while mating rate between

the two *D. americana* populations was moderately reduced in both directions. A logistic regression

assessing how mating rate varied based on male identity, female identity, and their interaction showed

a significant interaction effect only (males: $\chi^2(2) = 2.97$, $P = 0.28$; females: $\chi^2(2) = 2.52$, $P = 0.23$;

males*females: $\chi^2(4) = 12.67$, $P = 0.013$)—a difference from 4x4-mating trials where female population

identity was the primary predictor of mating rate. This variation between the assays appears to be

driven by female NE *D. americana* accepting fewer *D. novamexicana* males in 1x1 trials (20%) compared

to 4x4 trials (100%, Table 1), suggesting that male density within mating trials might affect copulation

success in this particular species combination. Finally, copulation latency differed marginally based on

both male identity (ANOVA: $F(2) = 3.14$, $P = 0.055$) and the male by female interaction ($F(4) = 2.48$, $P =$

0.062), but did not differ based on female species identity ($F(2) = 0.68$, $P = 0.5147$). Post-hoc tests

suggest that the marginal male population effect is likely due to lower copulation latency in SC *D.*

americana males relative to *D. novamexicana* males (Table S4).

148
Populations and sexes differ in CHC composition

We found that both populations and sexes within populations showed different and distinctive

CHC profiles, with the largest differences detected between SC *D. americana* and the other two lines

(Figure 2). Across all samples (*n*=5 replicate samples for each identity) we detected 8 alkene compounds

(2 of which are sex-specific) and 4 methyl-branched alkane compounds present in at least one

population. A principal component analysis of profiles from unmanipulated flies (unmanipulated-PCA or

'U-PCA') summarizing the primary axes of CHC profile composition found that 95.3% of compound

variation across all samples was explained by the first three principal components (U-PC1 to 3). Of these,

CHC composition varied significantly for both population and sex for U-PC1 (pop: $F_{(2, 429.29)}$, $P <$

0.0001; sex: $F_{(1, 46.97)}$, $P < 0.0001$) and U-PC2 (pop: $F_{(2, 10.54)}$, $P = 0.00043$; sex: $F_{(1, 181.69)}$, $P <$

0.0001), but had no sex or population effect for U-PC3 (pop: $F_{(2, 0.31)}$, $P = 0.737$; sex: $F_{(1, 0.57)}$, $P =$

0.458)(Figure 2). Notably, SC *D. americana* of both sexes differed from the other two populations along

the U-PC1 axis (Tukey HSD; Nov-SC: *P-adjust* < 0.0001; NE-SC: *P-adjust* < 0.0001; Nov-NE: *P-adjust* =

0.28), which also explains most of the CHC variation between samples (69.9%). U-PC1 was positively

loaded for most of the shorter carbon chain length compounds in both alkene and methyl-branched

alkane classes of compounds, and negative values were strongly loaded for the compounds with longer

carbon chain length (Table S5); consequently, this axis can be interpreted as a composite of average

compound length across the CHC profile as a whole (Figure 2). Accordingly, both sexes of SC *D.*

americana had a higher abundance of longer-chain alkenes and methyl-branched alkanes than did either

NE *D. americana* or *D. novamexicana*.

In contrast to U-PC1, U-PC2 appeared to primarily differentiate sexes within populations, but

also NE *D. americana* males from males of the other two populations. This axis was most heavily loaded

for two compounds: C21:1 and Me-C28, followed by C27:1 and Me-C26, with much smaller loadings for

all other compounds (Table S5). C21:1 is a male-specific compound that is not detected in females; Me-

C28, Me-26, and C27:1 abundance was also consistently different between sexes, with males always

having more of these CHCs than females within each population (Figure 2). Therefore, negative U-PC2

values can be interpreted as primarily representing more male-specific profiles, while positive U-PC2

values correspond to more female-like profiles. With respect to the detected species difference in U-

PC2, post-hoc tests reveal that this was driven by NE *D. americana* (Tukey HSD; Nov-SC: *P-adjust* = 0.99;

NE-SC: *P-adjust* = 0.0062; Nov-NE: *P-adjust* = 0.0088). Furthermore, we find that abundance of the

individual male-specific compound C21:1 differs between NE *D. americana* and males of the other two

populations, but not between *D. novamexicana* and SC *D. americana* (Tukey HSD; Nov-SC: *P-adjust* =

0.98; NE-SC: *P-adjust* = 0.016; Nov-NE: *P-adjust* = 0.022). (Results from tests of individual compound

differences can be found in Table S6).

183
Interpopulation perfuming influences SC D. americana *female acceptance of intra- and inter-population*

males

We found that patterns of sexual isolation could be modified by specifically changing the CHC

profiles of males of different populations. To do this, we used perfuming assays to manipulate the

pheromone profile of males by co-housing them with either intra- or inter-population males, and then

evaluated how this manipulation influenced mating rates among populations. For these analyses, we

focused on two pairings in two separate, analogous experiments—*D. novamexicana* and SC *D.*

americana ("Nov-SC pair"), and the *D. americana* population pair ("NE-SC pair")— because SC *D.*

americana shows the strongest sexual isolation from the two other populations, and the largest

differences in CHC composition. Within each perfuming experiment, we generated four combinations of

target and donor male identities: each population perfumed by same-population males (control) and

each population perfumed by hetero-population males. Each class of perfumed males was evaluated for

mating rate specifically with SC *D. americana* females, because these females were the most

discriminating against heteropopulation males in our previous mating assays (Table 2). The observed

mating rate of males perfumed with same-population males (the control manipulation) was similar to

that observed in unperfumed mating trials: *D. novamexicana* and NE *D. americana* males perfumed with

their own population performed poorly with SC *D. americana* females, relative to SC *D. americana*

perfumed with their own males, which had a 100% mating rate (Table 2). In strong contrast, hetero-

population perfumed males showed altered mating rates relative to control perfumed males. For Nov-

SC pairings, *D. novamexicana* males perfumed with SC *D. americana* males successfully mated with SC *D.*

americana females 94% of the time, a significantly higher mating rate than *D. novamexicana* males

perfumed with their own males (25%; Mann-Whitney U-test: U = 0, *P* = 0.025). (Note that copulation

success between conspecific perfumed *D. novamexicana* males and SC *D. americana* remains low (25%)

but differs from the complete mating isolation observed in unperfumed trials, possibly because of

changes in behavior due to male-male co-housing during the perfuming manipulation.) The reciprocal

treatment also showed a significant effect of perfume source: male SC *D. americana* perfumed with *D.*

novamexicana males displayed copulation success of 19%, significantly lower success compared to

conspecific-perfumed SC *D. americana* (100%, Mann-Whitney U-test: U = 0, *P* = 0.018). For NE-SC

pairings, SC *D. americana* male mating rate was also significantly reduced when perfumed with NE *D.*

americana males (50%), compared to same-population perfumed SC *D. americana* males (100%, Mann-

Whitney U-test: U = 0, *P* = 0.018). In contrast, mating rate of NE *D. americana* males perfumed with SC

D. americana males increased slightly (56%), but did not significantly differ from the control (same-

population-perfumed) males (44%, Mann-Whitney U-test: U = 5.5, *P* = 0.54).

Overall, these results indicate that perfuming males with heteropopulation CHCs influences the

frequency of successful copulation with SC *D. americana* females. SC *D. americana* males perfumed with

profiles of either *D. novamexicana* or NE *D. americana* had significantly reduced mating rates with their

own females. Conversely, perfuming *D. novamexicana* males with SC *D. americana* males significantly

increased their mating rate with *D. americana* females, almost completely reversing the pattern of

sexual isolation observed for unmanipulated males. This large shift in mating rate for heteropopulation-

perfumed *D. novamexicana* indicates that differences in male CHC profiles play a critical role in sexual

isolation in the Nov-SC pair. In contrast with the three other classes of heteropopulation-perfumed

males, perfuming NE *D. americana* with SC *D. americana* males did not significantly change their mating

rate. It possible that this manipulation might have been less effective at altering the CHC profile of NE *D.*

americana males (see next section), and/or that mating isolation in the NE-SC pair might depend on

more complex factors than a simple shift in CHC composition alone (see Discussion).

Perfuming shifts CHC composition towards the donor male profile

Using CHCs extracted from an additional set of perfumed males (see Methods), we confirmed

that our heteropopulation perfuming manipulation produced quantitative changes in composite male

CHC profiles in both the Nov-SC and NE-SC pairs, by shifting male CHCs closer to the donor male profile.

PCAs were performed separately for each pairing (N-PCA for the Nov-SC pair, and A-PCA for the NE-SC

pair, hereafter), as were analyses of differences among classes of perfumed males. In both cases, of the

first three PCs of CHC composition, PC1 (i.e. N-PC1 or A-PC1) varied by both donor and target male

identity (Table S7), indicating that perfuming significantly shifted CHC profiles along the primary axis of

variation in both perfuming experiments. Nonetheless, the specific donor-target manipulation that was

most successful in this regard differed between the Nov-SC and NE-SC pairs. In the Nov-SC pair, *D.*

novamexicana males perfumed with SC *D. americana* males significantly differed in CHC composition

from control (*D. novamexicana*) perfumed samples (N-PC1) (t(5.9) = -2.8, *P* = 0.031); however,

heteropopulation-perfumed SC *D. americana* did not differ from same-population SC *D. americana*-

perfumed samples along the same axis (t(3.53) = 1.91, *P* = 0.14). In contrast, in the NE-SC pair, SC *D.*

americana perfumed with NE *D. americana* showed a significant shift in A-PC1 (t(4.85) = 4.23, *P* =

0.0088), but there was no significant shift for SC-perfumed NE *D. americana* compared to same-

population-perfumed controls (t(4.68) = 0.055, *P* = 0.96). The difference between these pairs might be

attributed to high variation seen among samples in some groups (in particular same-population

perfumed SC *D. americana*). Regardless, it is clear from these results that this perfuming assay can

produce significant, detectable differences in CHC profiles after heteropopulation perfuming, most

notably in *D. novamexicana* males.

Patterns of gene expression and sequence variation implicate an elongase gene that contributes to CHC

variation between species

From among a set of 23 candidate genes whose orthologs have functions related to cuticular

hydrocarbon variation, both transcriptome and sequence comparisons implicated one specific elongase

locus as potentially causal in CHC composition differences between our species. Using whole-

transcriptome RNA-seq data from the same three populations (previously generated in Davis and Moyle

2020), we found that 3 of our candidate genes in males and 2 candidates in females were in the upper

10[th] percentile of genes most differentially expressed between populations (i.e. they showed a stronger

species effect than 90% of all 11301 expressed genes in the transcriptome dataset) (Table 3). Of these

candidate genes, only CG17821 showed elevated gene expression specifically in SC *D. americana*

compared to the other two populations; this pattern was observed in both sexes but is more

pronounced in females (Figure 3).

Closer inspection of our CG17821 sequences revealed that alleles in NE *D. americana* and *D.*

novamexicana share a thymine insertion mutation that causes a premature stop codon 4 amino acids

upstream of the end of the gene, compared to the annotated gene model in outgroup *D. virilis* and the

allele in our SC *D. americana* stock (Figure 3); these four terminal amino acids are not present in RNA

transcripts for either NE *D. americana* and *D. novamexicana*. Among our three lines, this suggests an

association between the truncated protein product of CG17821 and the "short" CHC phenotype we

observe. This association is further supported by additional CHC and sequence data from another *D.*

americana population, originally collected in Anderson, IN (species stock center line 15010-0951.00,

henceforth IN *D. americana*). Lamb et al. 2020 characterized CHC divergence between females from this

IN *D. americana* stock and another *D. novamexicana* line (species stock center 15010-1031.04). The

profile for the *D. novamexicana* line had no long compounds (>C30), a low abundance of C29

compounds, and presence of multiple short alkenes (<C27), consistent with the "short" CHC profile we

observe for our *D. novamexicana* line. Likewise, the IN *D. americana* profile they observed is broadly

consistent with the "long" profile we observed in SC *D. americana*, as both show presence of

compounds longer than 30 carbons, a high abundance of C29 length alkenes, and absence of

compounds shorter than 27 carbons. In addition, using a genome assembly from the same IN *D.*

americana stock (Kim et al. 2020), we found that this population also has the non-truncated allele of

CG17821. These data are consistent with our inference that truncation of this allele could contribute to

the difference between "long" and "short" CHC phenotypes among populations in this group.

Evolutionary history of elongase CG17821

Several lines of evidence indicate that CG17821 has a complex evolutionary history within the *D.*

americana subgroup, that likely includes post-speciation introgression. The presence of the non-

truncated allele in outgroup *D. virilis* (Figure 3) indicates that the truncated allele is derived from a

thymine insertion mutation event that took place within the *D. americana* subgroup. Moreover, *D.*

novamexicana and our western *D. americana* (NE) line share an identical derived allele at CG17821,

indicating an evolutionary history at this locus that disagrees with expected phylogenetic relationships

among these three taxa (i.e., where the two *D. americana* populations are expected to be most closely

related). Gene trees for each of our 23 candidate genes confirmed that CG17821 has a topology that is

discordant with the expected species relationships, placing NE *D. americana* as sister to *D.*

novamexicana rather than grouping it with SC *D. americana* (Figure 3). Four other CHC candidate genes

also show this discordant topology, most notably CG18609 which is located immediately downstream

(within 1kB) of CG17821, according to the annotated genome of *D. virilis* (Figure 3). For 4 out of 5 of

these loci—including both CG17821 and CG18609—using the allele from the IN *D. americana* stock

instead produces a genealogy where the *D. americana* populations group together, as expected from

the species tree (Figure 3).

The observation of phylogenetically discordant sites can be due to several factors, notably

incomplete lineage sorting (ILS) or post-speciation introgression (see also Discussion). Two lines of

evidence strongly support introgression. First, evidence from SNP variants, across the whole

transcriptome and specifically at CG17821, is more consistent with introgression between *D.*

novamexicana and our western *D. americana* (NE) population—as determined by Patterson's D-statistic

(Durand et al. 2011). In particular, we found that genome-wide, of 34441 total SNPs detected in 11301

loci within the transcriptome dataset (Davis and Moyle 2020), 14902 supported *D. americana*

populations as sister (in accordance with the species tree), 13372 SNPs grouped *D. novamexicana* and

NE *D. americana*, and 6167 grouped *D. novamexicana* with SC *D. americana*. The significant excess of

shared variants between *D. novamexicana* and our NE *D. americana* line ($D = 0.369$, $P < 0.0001$), is

consistent with a history of introgression between the progenitors of these two populations. Second,

gene tree topologies at and around CG17821 also indicate this specific genomic region shares recent

ancestry between *D. novamexicana* and NE *D. americana* due to introgression. CG17821, the adjacent

downstream CHC candidate CG18609, and the next nearest gene (*List*, a neurotransmitter located ~14

kB downstream) all display shared derived SNPs between *D. novamexicana* and the NE *D. americana*

line, and discordant gene tree topologies that group these two lines as sister taxa. In contrast,

topologies for the next two nearest up- or down-stream genes in this region reflect the species tree,

indicating that recent shared ancestry between *D. novamexicana* and the NE *D. americana* extends

across a genomic window between 18 and 68 kB long around the specific region containing CG17821

(Figure 3C). Estimates of linkage disequilibrium (LD) from *D. melanogaster* autosomes indicate that most

LD decays within 200 base pairs, and r^2 (the correlation between SNPs) decreases to <0.1 within a 1 kB

window (Franssen et al. 2015); therefore, the size of the putatively-introgressed window observed here

is well outside the range expected under ILS, as discordant ancestry due to sorting from ancestral

variation is expected in small blocks (Hudson and Coyne 2002, Gao et al. 2015). Note that inversions

could also be responsible for capturing ancestrally-segregating variation in larger genomic regions than

expected from LD measures, and populations within this group are known to differ in the

presence/absence of inversions, including at the specific genomic region containing these genes (Reis et

al 2015). However, the inversion at this location is only present in southern *D. americana* (i.e., within the

"*D. a. texana*" chromosomal form) to the exclusion of NE *D. americana*, *D. novamexicana,* and *D. virilis*,

and therefore could not be responsible for the patterns of discordant variation we observe for these

genes here. Coupled with genetic and phenotypic evidence above, these results indicate introgression of

these genes between *D. novamexicana* and western *D. americana*—and not ILS—is most likely

responsible for their shared allele at CG17821 and therefore potentially for their similarity in "short"

CHC phenotypes.

We also evaluated whether there was evidence of elevated protein evolution at CG17821,

CG18609 or any other of our candidate genes, based on estimates of the ratio of nonsynonymous to

synonymous substitutions (d_N/d_S) at these loci, compared to a genome-wide average rate estimated

from our transcriptome-wide dataset (see Methods). For 4397 genes transcriptome-wide, the median

and mean d_N/d_S were 0.065 and 0.0991, respectively (± 0.133 s.d.) (Figure S3). Of our candidate genes,

only three had an estimated d_N/d_S greater than 1 standard deviation above this mean, including the two

candidates within the putatively introgressed region—CG17821 (d_N/d_S = 0.284) and CG18609 (d_N/d_S =

0.249)—as well as CG6660 (d_N/d_S = 0.38714)(full results in supplementary Table S8). Notably, these rates

of protein change fall within the top 8% of all genes analyzed (Figure S3, all gene data reported in

supplementary file). While these estimates are below the threshold for unambiguous evidence of

positive selection (d_N/d_S > 1), only 9 genes in the whole dataset met this criterion. Of the three candidate

loci that exhibit elevated protein evolution, CG6660 does not show patterns of genealogical, allelic, or

gene expression variation that match our observed CHC phenotypic variation (Figure 3A/B, and above).

For both CG17821 and CG18609, we also estimated d_N/d_S using just *D. virilis* and SC *D. americana*

sequence comparisons, to determine if elevated protein evolution in these genes is solely driven by the

branch leading to the *D. novamexicana*-NE *D. americana* allele, or if this pattern is more general across

the clade. We found our estimate of d_N/d_S was still modestly elevated between *D. virilis* and SC *D.*

americana (d_N/d_S = 0.284) even when this branch was removed—indicating CG17821 has experienced

consistently elevated protein evolution across the whole clade. In contrast, CG18609 protein evolution is

estimated to be lower (d_N/d_S = 0.147) when just considering divergence between *D. virilis* and SC *D.*

americana, suggesting that most of the accelerated protein evolution in this gene occurred after the

split of the *D. novamexicana*/NE *D. americana* lineage. The timing of elevated protein evolution in

CG18609 therefore also generally coincides with CHC profile differences observed here and, like

CG17821, CG18609 is also an elongase. However, CG18609 does not show patterns of gene expression

that match the "long" versus "short" CHC phenotypes we observe (instead, its expression is significantly

reduced in NE *D. americana*; Figure 3A) so while it's possible that this tandem elongase gene also plays a

role in CHC phenotype divergence, it is unlikely to be responsible for the primary axis of CHC variation

described here.

Finally, we note that our d_N/d_S analysis also revealed two odorant binding proteins (OBPs)—to

be among the fastest evolving loci in our transcriptome-wide dataset: the orthologs of *Obp99d* (second

fastest in our dataset; d_N/d_S = 1.589), and *Obp56g* (40[th] fastest; d_N/d_S = 0.689) (see supplementary file).

OBPs are thought to facilitate olfactory processing by chaperoning odorants to the olfactory receptor

neurons or terminating neuron activity by clearing odorants form the surrounding (Sun et al. 2018). OBP

genes have been previously shown to evolve quite rapidly in multiple insect groups (Foret and Maleszka

2006), although rates in these specific OBPs are lower across 6 species in the *Drosophila melanogaster*

group (d_N/d_S > 0.24; Vieira et al. 2007) than we detect here. Notably, both these OBPs are known to

expressed in chemosensory organs in *D. melanogaster*—including adult labellum (*Obp56g*; Galindo and

Smith 2001), antenna and maxillary palps (*Obp99d*; Hekmat-Scafe et al. 2002), and wing sensilla (both

loci; He at al. 2019)—consistent with roles in pheromone perception. Moreover, one of these loci—

Obp56g—is a known seminal fluid protein in the *D. melanogaster* group (Findlay et al 2008), that also

changes in expression within females specifically in response to mating (McGraw et al. 2004). Given

these roles, the elevated rates we observe here suggest both genes might contribute to variation in

behavioral responses to pheromone and other sexual stimuli among our species (see also discussion).

Discussion

Identifying genetic and evolutionary mechanisms involved in the earliest steps of reproductive

isolation between species is essential for understanding the factors that drive speciation. Evolutionary

divergence in sexual signals may be an especially potent contributor to this process (Schemske 2000,

Coyne and Orr 2004, Ritchie 2007, Schluter et al. 2009, van Doorn et al. 2009). However, demonstrating

the connection between sensory signal divergence and emerging reproductive isolation can be

challenging, as it requires identification and demonstration of the direct role of specific signals

mediating sexual isolation between species and knowledge of the specific mechanistic changes that

have given rise to lineage differences in this signal. Here we have demonstrated that sexual isolation

between laboratory populations in the *D. americana* group is based on female choice of male chemical

signals, and identified both the specific phenotypic shift between species in pheromone chemistry as

well as a genetic variant likely contributing to this phenotypic change and the mating isolation that

results from it. Together these data support a clear role for sensory signal divergence in the evolution of

premating isolating barriers between populations in this closely related group, and provide insight into

how relatively simple genetic and phenotypic mechanisms can cause strong isolation even at early

stages of evolutionary divergence.

Female choice of male CHC variation is responsible for isolation between D. novamexicana *and SC* D.

americana

Differences in copulation success between populations are the product of variation in male

choice (via differences in courtship intensity and copulation attempts, depending upon female identity),

female choice (via differences in acceptance rates of males, depending upon their identity), or a

combination of these factors. Disentangling these alternatives is critical for identifying the specific trait

and corresponding preference variation responsible for the emergence of prezygotic species barriers.

Here we showed that female choice of a male sensory signal results in strong sexual isolation among *D.*

americana group populations, specifically SC *D. americana* female preference for CHC profiles of their

own males and rejection of males with alternative profiles. The role of CHCs is exceptionally clear in the

case of premating isolation between male *D. novamexicana* and female SC *D. americana*: strong sexual

isolation in this cross can be almost entirely reversed by perfuming *D. novamexicana* males with SC *D.*

americana male CHCs. In contrast, we find little evidence for differential male preference of females,

even though female CHC profiles are also divergent between populations. Instead, our behavioral data

indicates that males did not alter their courtship behavior in response to female species identity.

Interestingly, these observations also suggest evidence for female choice of male courtship behaviors,

whereby SC *D. americana* male courtship consistently induced females to copulate sooner than

courtship behaviors of other population males. Moreover, our finding of strong asymmetric mating

isolation between male *D. novamexicana* and specific (southeastern) populations of *D. americana*

recapitulates patterns of isolation described in this group more than 70 years ago (Spieth 1951, Table

S1), indicating that these mating isolation patterns reflect natural variation among populations from

which these lines were collected.

These findings fit within a body of studies that have identified either male or female choice of

sensory signals as critical for sexual isolation among *Drosophila* species (patterns reviewed Yukilevich

and Patterson 2019). In *Drosophila melanogaster,* many studies have found that male choice of specific

female CHC compounds play a role in isolation between closely related heterospecifics (Coyne et al

1995, Billeter et al. 2009) as well as between intraspecific populations (Wu et al. 1995, Hollocher et al.

1997, Yukilevich and True 2008). Such patterns of CHC-mediated male mate-discrimination have also

been associated with allelic variation in CHC elongases. For example, *D. sechellia* reproductive isolation

from closely related *D. simulans* results from male mate-discrimination based on female CHC profiles

(Shahandeh et al. 2017) and the CHC elongase *eloF* has been demonstrated to inhibit interspecific

mating in the same species (Combs et al. 2018). Female choice of male CHC variation has received

comparably less mechanistic attention, however has been demonstrated as a prezygotic barrier in

several *Drosophila* groups, most notably between *D. santomea* and *D. yakuba* (Coyne et al. 2002, Mas

and Jallon 2005), between the mycophagous *D. subquinaria* and *D. recens* (Curtis et al. 2013, Dyer et al.

2013), and between *D. mojavensis* males reared on different cactus substrates (Havens and Etges 2013).

Our results therefore complement a growing body of evidence that shows female choice can be an

important factor dictating patterns of sexual isolation among closely related *Drosophila* populations and

species.

CHC divergence and the role of Elongases

Dissecting the finer details of phenotypic divergence in sensory signals can help pinpoint

underlying mechanisms and associated genes responsible, and more clearly demonstrate how these

signal changes contribute to isolation between populations. Here we found that CHC divergence

between our lines occurs primarily on the basis of compound length, with *D. novamexicana* and NE *D.*

americana profiles both having similar enrichment of shorter compounds ("short" phenotype) compared

to an enrichment of longer compounds for SC *D. americana* ("long" phenotype). These differences in

abundance of shorter- versus longer-chain compounds were observed across both sexes, and across

both alkenes and methyl-branched alkanes (Figure 2, Table S6). In contrast, we find no evidence for

variation in other features such as double bond or methyl branch location or number. These features

themselves suggest that the striking difference between 'shorter' and 'longer' profile phenotypes could

be due to variation in fatty-acid elongase activity, which globally influences the carbon chain length of

CHC precursors, and therefore can have consistent downstream effects on both alkenes and methyl-

branched alkanes, because both are modified after the elongation step (Pardy et al. 2018).

Strikingly, analyses of gene expression and sequence variation revealed that CG17821 and

CG18609—two putative fatty-acid elongases (Szafer-Glusman et al. 2008, Gaudet et al. 2011)—could

functionally contribute to our observed variation between longer and shorter CHC profiles. Sequences at

both loci are identical between NE *D. americana* and *D. novamexicana*, differ from SC and IN *D.*

americana lines, and show modestly elevated protein evolution on the branch leading to *D.*

novamexicana/NE *D. americana,* all of which broadly coincide with our observed differences between

"short" versus "long" CHC phenotypes. The variation we observed at CG17821 is particularly interesting,

because we observe a thymine insertion mutation that results in premature truncation of CG17821

specifically in *D. novamexicana* and NE *D. americana* (Figure 3) and because our gene expression data

also indicates reduced expression of the CG17821 allele specifically in these lines. In the latter case,

while we might expect gene expression differences to be tissue-specific (that is, observed specifically in

the oenocytes: the CHC-producing organs), our data indicate that the population-specific expression

signal at this locus is strong enough to detect from whole-body transcriptome data. Both observed

sequence and expression changes at CG17821 could individually produce a greater abundance of short

CHC products either by lowering elongase protein levels or reducing enzyme activity. Therefore,

although our current data cannot differentiate which change may have occurred first (or whether they

are pleiotropic), either could produce the pattern of phenotypic difference observed between

populations. Overall, these data suggest allelic variation in CG17821 primarily underlies the major axis of

CHC divergence observed between SC *D. americana* and the other populations, consistent with a

hypothesis of a simple underlying basis for chain length variation. Moreover, because this axis of CHC

divergence appears primarily responsible for sexual isolation between *D. novamexicana* and SC *D.*

americana, this points to a large role for simple allelic change at this genomic location in the emergence

of a strong isolating barrier between these two populations.

The evolutionary history of CHC divergence and consequences for past and future sexual isolation in this

group

Together, our data demonstrate sexual isolation between SC *D. americana* and *D. novamexicana*

is due to CHC divergence in compound length and suggest that variation between truncated and non-

truncated alleles of the elongase CG17821 contribute to this phenotypic variation. Moreover, both

genome-wide SNP variation, as well as localized variation specifically around this locus (Figure 3C),

indicate that this phenotypic variation involves a history of introgression. These data point to a model

hypothesis for the evolutionary history of transitions involved in the change in CHC profiles between

species and, potentially, in the emergence and expression of sexual isolation that depends upon this

phenotype.

First, the distribution of both "long" versus "short" CHC phenotypes (shown here and in Lamb et

al. 2020), and allelic variation CG17821, indicate that the "short" phenotype and the CG17821 truncated

allele are derived states that arose within the *D. americana* group. This shift most likely occurred in

western lineages that gave rise to contemporary *D. novamexicana*. The evolutionary forces responsible

for the persistence and spread of this phenotype are not yet known. The relationship between variation

in environmental factors, insect stress physiology, and features of CHC length and branching is known to

be complex. Prior work has associated aspects of CHC divergence with abiotic variation such as latitude

(Frentiu and Chenoweth 2010, Rajpurohit et al. 2017, but see Gibbs et al. 2003), and physiological traits

such as desiccation resistance (reviewed Chung and Carroll 2015), that suggest a role for natural

selection in shaping CHC composition, but the specific sources(s) of selection can be challenging to

pinpoint. In the *D. americana* group, species habitats are differentiated primarily on the basis of water

availability and these two species, as well as populations within them, differ in key physiological traits

such as desiccation resistance (Davis and Moyle 2019). However, sexual selection might also contribute

to shaping evolution at CHC loci—as evidenced by the importance of CHC variation for mating success

outlined here and elsewhere. We also find evidence that two odorant binding proteins (OBPs)—*Obp99d*

and *Obp56g*—are rapidly evolving across this group. OBPs are known to be important for mediating

olfactory behavioral responses during sexual interactions (Laughlin et al. 2008, Leal 2013, Sun et al.

2018) as well as host-plant preference (Matsuo et al. 2007, Comeault et al. 2017). Therefore these loci

could be evolving due to natural selection, sexual selection, or both, including in response to changes in

pheromone profiles described here, or to other factors such as this group's close but little investigated

habitat association with willow (*Salix* sp.) trees (Blight and Romano 1953, McAllister 2002, personal

observations). Interestingly, our analysis indicates that the elongase CG17821 has experienced

modestly elevated protein evolution across the whole *D. virilis* sub-clade; this suggested history of

sustained selection indicates this locus might have played important roles in long-term CHC-mediated

adaptive divergence across this group. In comparison, the downstream elongase CG18609 has evidence

of elevated protein change primarily on the branch leading to *D. novamexicana*/NE *D. americana*,

possibly suggesting that this acceleration occurred after impactful changes to CG17821.

Based on our data associating phenotypic divergence with sexual isolation, the appearance of

this new CHC phenotype would have reduced sexual compatibility between derived "short" males and

females with strong preferences for the "long" ancestral CHC profile. Persistence of this phenotype

would have required a broadening of female preference to accommodate males with the derived

("short" CHC) pheromone phenotype (e.g., as has been observed, for example, in male *Ostrinia* moths;

Roelofs et al. 2002). Our data support this expectation, as *D. novamexicana* females are more accepting

of both (putatively derived) *D. novamexicana* and (ancestral) SC *D. americana* male CHC phenotypes,

while SC *D. americana* females discriminate against derived "short" CHC phenotypes. Interestingly, this

model for the evolution of asymmetric sexual isolation is broadly consistent with Kaneshiro's (1976,

1980) model for peripatric speciation. Kaneshiro observed that females from derived populations

frequently have broad preferences for both derived and ancestral male phenotypes; he proposed that

this was due to relaxed selection on narrow female preferences in genetically bottlenecked island

populations, where founder effects have led to the loss of elements of male courtship. Although

Kaneshiro's model explicitly invokes genetic drift in male trait evolution, our observations indicate his

model for the origin of sexual isolation asymmetry could extend more generally to any case where

evolutionary change affects a trait important for male sexual signaling. In the case described here,

evolutionary change in male CHC profiles (possibly due to selection acting on a CHC elongase gene(s)),

accompanied by an apparent broadening of female preferences for these profiles in derived

populations, has resulted in the emergence of strong premating asymmetry specifically between

females with ancestral trait preferences and males with derived trait values—akin to the model outlined

by Kaneshiro.

Intriguingly, our data also support the subsequent movement of this CHC phenotype from *D.*

novamexicana into western *D. americana* lineages. One possible explanation for shared variation in the

CG17821-CG18609 locus and CHC phenotypes is that this arose from segregating ancestral variation

present in both *D. americana* and *D. novamexicana* (that is, is due to ILS). Prior evidence (Caletka and

McAllister 2004) as well as data here indicates that some observed site discordance between

populations of *D. americana* and *D. novamexicana* is consistent with ILS. However, our analysis strongly

supports the additional occurrence of introgression between *D. novamexicana* and western *D.*

americana, including specifically of this trait from *D. novamexicana* into western *D. americana*

populations. Both their significant excess of shared genome-wide variation (as indicated by the D-

statistic), and a genomic region of at least 18 kB of shared recent ancestry surrounding

CG17821/CG18609 that accompanies a similar shift to the "short" CHC phenotype in western *D.*

americana, support this inference. Note also that while *D. novamexicana* and *D. americana* are reported

to be allopatric, limited and sporadic field collections of *D. novamexicana* mean there is an incomplete

understanding of the density and extent of this species' historical range. As a result, this species could

have been in closer geographical contact with western *D. americana* populations during the period since

their initial split (~500KYA), which could help explain evidence for introgression after speciation inferred

here. Recent work (Sramkowski et al. 2020) describing rare *D. novamexicana*-like pigmentation alleles in

geographically disparate *D. americana* populations, similarly suggests evidence of more recent gene

exchange between these species.

Given the effect of the "short" CHC phenotype on sexual isolation between *D. novamexicana*

and the SC *D. americana* population, this introgression is expected to have consequences for

reproductive isolation among *D. americana* lineages. Interestingly, our data indicate that, even though

NE *D. americana* shows the general shift to shorter chain CHC length associated with the introgressed

region, its current patterns of sexual isolation differ from those seen in *D. novamexicana*. One

explanation might be that this introgression-mediated shift in CHCs occurred on a novel *D. americana*

genomic background, with potential consequences for the expression of introgressed CHC-affecting loci,

and therefore for patterns and strengths of CHC-mediated isolation. A concrete example of these

background effects can be seen for CG18609, where NE *D. americana* shares the *D. novamexicana* allele

but nonetheless exhibits significantly reduced expression of this locus compared to the other two

lineages (Figure 3A). A differential history of allelic exchange (with *D. novamexicana*) across the range of

D. americana, plus variable genomic background effects on the expression of CHC loci, could contribute

to the more complex mating relationships observed here, and elsewhere, among *D. americana*

populations. For example, Spieth's mating analyses (Table S1) identified up to 10-fold variation in mating

success among nine disparate populations of *D. americana*. The possibility that this variation is

influenced by differential gene exchange with *D. novamexicana* (including of loci with major effects on a

signaling phenotype) is testable with work examining mating success between geographically diverse *D.*

americana populations—particularly between eastern and western populations—and its covariation

with CHC phenotypic and genotypic variation.

Regardless of the collateral consequences for isolation among *D. americana* populations, our

data clearly support the role of divergent cuticular hydrocarbon profiles—specifically a general shift in

carbon chain length—in sexual isolation between our *D. novamexicana* and SC *D. americana*

populations. They also implicate a potentially causal role for the gene CG17821 in determining CHC

phenotype via a global change in CHC elongation activity early in the generation of these compounds.

These data provide a strong example of how a recently derived allele in a single gene with large

phenotypic effects on a sexual signal could underpin asymmetric sexual isolation between closely

related species. Moreover, they suggest multiple (behavioral, biochemical, and molecular) lines of

evidence that chemosensory processes are evolving rapidly and dynamically across this group.

Methods

Experimental Fly stocks

Three stocks were obtained from the University of California San Diego Drosophila Species Stock

Center (DSSC): a *Drosophila novamexicana* stock from San Antonio, NM (15010-1031.08); and two *D.*

americana stocks, one from Chadron, NE (15010-0951.06, NE *D. americana* throughout); and one from

Jamestown, SC, (15010-1041.29, SC *D. americana* throughout). All stocks were originally collected

between 1946 and 1953. *D. americana* has sometimes been divided into two subspecies according to

presence (*D. americana americana*) or absence (*D. a. texana*) of a chromosomal fusion of the X- and 4-

chromosomes that shows a distinct latitudinal cline (McAllister 2002), however because sub-specific

differences apart from this fusion have not been consistently supported, our two lines are treated as

populations from within a single heterogeneous species here. The *D. novamexicana* and NE *D.*

americana populations used here are the same as those collected and used by Spieth 1951. All fly stocks

were reared on standard cornmeal media prepared by the Bloomington Drosophila Stock Center (BDSC)

at Indiana University, and were kept at room temperature (~22 °C). Every assay in this study used virgins

isolated within 8 hours of eclosion and aged for 7 days prior to the start of experiments, similar to the 8

days used by Spieth (1951).

4x4 unperfumed mating assay

We performed trials in which four virgin males and four virgin females were paired and

observed for mating behavior, following the design used by Spieth (1951) which allows for behavioral

interactions that might not otherwise be observed in similar single-pair assays. Within each trial, all

males are from a single population, as are all females, so are no-choice with respect to the genotype of a

mating partner; crosses are varied by pairing males and females of alternative lines. For each trial, 4

males and 4 females were transferred to a single vial without anesthetization and observed for 3 hours.

The number and duration of each copulation event was recorded for each trial. This assay was repeated

for a total of 5 replicates for each possible population combination, in reciprocal (i.e., 9 cross types, each

of N = 5). Each trial used 7-day old virgins and testing was started within 30 minutes of lights on in the

morning.

Courtship behavior assay

To quantify and evaluate differences in courtship behaviors between cross types, flies were

observed in single pair (no-choice) mating assays. We used a modified FlyPi setup—that combines a

Raspberry Pi (Raspberry Pi Foundation, Cambridge, UK), pi camera, and 3D printed parts (Chagas et al.

2017)—to record courtship behaviors. Assays were performed in a modified cell culture plate consisting

of six 3 cm-diameter culture wells, each with a small amount of cornmeal media in the bottom, that

allowed six total crosses to be recorded simultaneously. For each assay, individual virgin male and

female flies of a given cross were aspirated without anesthetization to a cell culture well; after 6 total

crosses were set up, the plate was videotaped for a contiguous 3-hour period. The six cross

combinations assessed in any particular video trial were randomized to account for variance that might

otherwise be explained by date. As in the 4x4 mating assay, we performed 5 replicates of all possible

population combinations in reciprocal.

Behavioral features were analyzed and scored manually by the same individual to avoid

subjective variation among researchers. Three courtship behaviors—male display events, male tapping

events, and male licking events—in addition to copulation were scored in each 1x1 trial using the

following criteria. A male "display" was counted when a male performed a combination of back and

forth movements and occasional wing flicks while maintaining sustained orientation in front of and

facing the female. "Tapping" events were defined when a male used his tarsus to touch the abdomen of

the female when oriented behind her. "Licking" events were defined when the male's mouthparts

contacted the female genital arch. For male display, tapping, and licking events, individual events were

scored separately only if a different behavior (including sitting still/walking away) was observed

between instances of the defined behavior. This criterion minimized overcounting of discrete behavioral

events, especially those with difficult to view aspects such as number of times a male extruded

mouthparts during a contiguous licking event. In addition to these behaviors, copulation success and

latency to copulation were also recorded for each trial. Note that females also engage in tapping and

other behaviors, but because these appear to be less consistent and are more difficult to observe and

score, they were not addressed in this study.

Extraction and quantification of cuticular hydrocarbons

Cuticular hydrocarbons were extracted from pooled samples by placing five 7-day old virgin flies

of a single sex and species identity in a 1.8 mL glass vial (Wheaton 224740 E-C Clear Glass Sample Vials)

with 120 µL of hexane (Sigma Aldrich, St Louis, MO, USA) spiked with 10 µg/mL of hexacosane (Sigma

Aldrich). After 20 minutes, 100 µL of the solution was removed to a sterilized 1.8 mL glass vial (Wheaton

224740 E-C Clear Glass Sample Vials) and allowed to evaporate overnight under a fume hood. Extracts

were stored at -20 °C until analysis. Five replicate samples consisting of 5 flies per sample (25 flies total)

were prepared for each sample type, and all replicates were extracted on the same day.

Gas chromatography mass spectrometry (GC/MS) analysis was performed on a 7820A GC

system equipped with a 5975 Mass Selective Detector (Agilent Technologies, Inc., Santa Clara, CA, USA)

and a HP-5ms column ((5%-Phenyl)-methylpolysiloxane, 30 m length, 250 µm ID, 0.25 µm film thickness;

Agilent Technologies, Inc.). Electron ionization (EI) energy was set at 70 eV. One microliter of the sample

was injected in splitless mode and analyzed with helium flow at 1 mL/ min. Two different temperature

gradients were used depending on the sample type. For CHC analysis of unmanipulated males and

females of each species, the following parameters were used: column was set at 50 °C for 0 min,

increased to 210 °C at a rate of 35 °C/min, then increased to 280 °C at a rate of 3 °C/min. The MS was set

to detect from m/z 33 to 500. For analysis of samples from perfuming trials, the parameters were

modified to increase resolution and sensitivity for less abundant compounds: the column was set at 40

°C and held for 3 min, increased to 200 °C at a rate of 35 °C/min, then increased to 280 °C at a rate of 3

°C/min and held for 15 minutes. Chromatograms and spectra were analyzed using MSD ChemStation

(Agilent Technologies, Inc.). CHCs were identified on the basis of retention time and electron ionization

fragmentation pattern. Compounds are identified in this study as: CXX:Y for alkenes or Me-CXX for

methyl-branched alkanes, where XX indicates the length of the carbon chain, and Y indicates number of

double bonds, e.g. C21:1 is a 21-carbon alkene with a single double bond.

The abundance of each compound was quantified by normalizing the area under each CHC peak

to the area of the hexacosane signal using homebuilt peak selection software (personal correspondence,

Dr. Scott Pletcher, Univ. of Michigan).

Perfuming manipulation and mating assay

'Perfuming' involves co-housing target flies in vials filled predominantly with flies of a desired

donor identity, so that the CHC profile of the target flies is altered via physical transfer of CHCs from

donor flies (Coyne et al. 1994, Dyer et al. 2014, Serrato-Capuchina et al. 2020). Perfuming was

performed by placing 2 'target' males with 15 'donor' males within a single vial. All males were 1-day old

virgins when perfuming vials were established, and the wings of donor males were removed (under

anesthetic) to distinguish them from target males. For the hetero-population perfuming treatments,

target males were co-housed with donor males of a different population; same-population (control)

treatments paired donor and target males of the same identity. Following 7 days of perfuming, 4 male

flies from the same perfuming conditions were transferred (without anesthetization) to vials containing

4 female SC *D. americana*. Within each experiment (Nov-SC, or NE-SC), we ran one trial of each

perfuming condition (2 heterospecific-perfumed types, and 2 conspecific-perfumed types) in parallel on

the same day (four 4x4 trials in total).

For CHC analysis, males were perfumed as described for the perfumed mating assay, for both

Nov-SC and NE-SC experimental pairs, with the exception that 2-3 target males (rather than just 2) were

co-housed with 15-18 donor males, enabling 2 parallel perfuming vials to generate 5 target male flies

per identity, for each individual CHC extraction. This was then replicated 4 times for each identity to

reach N=4 biological replicates for this analysis.

Candidate gene selection

Our candidate list was generated by searching Flybase (Flybase.org) for annotated *Drosophila*

melanogaster genes with one of the protein-coding domains (as identified by InterProt) that have

known functions in CHC synthesis (Pardy et al. 2018): "fatty acid desaturase", "fatty acid desaturase

domain", "Cyp4g", "ELO family", or "fatty-acid-synthase"—resulting in an initial list of 34 genes. To this

we added the pigmentation genes *ebony* and *tan* as they have been shown to alter CHC variation among

both *D. melanogaster* (Massey et al. 2019) and *D. americana* group species (Lamb et al. 2020). With

these 36 genes, we identified orthologs in *Drosophila virilis* (via Flybase using OrthoDB v9.1, Zdobnov et

al. 2017) and then evaluated whether each of these loci had transcript expression in our three

populations. To do so, we used the BLASTn function of BLAST+ version 2.6.0 (Camacho et al. 2009) to

search for matches between the *D. virilis* orthologs and previously published whole-body transcriptome

data generated from the same three populations using RNA-seq (Davis and Moyle 2020). Our analyses

used only gene expression data from the control (ambient) conditions for both males and females from

this study, and excluded any desiccation stress treatment data. Of the 36 initial genes (listed in Table

S9), 30 were found to have unambiguous 1-to-1 orthologs in *D. virilis* and, of these, 23 had transcripts

present within the Davis and Moyle (2020) dataset. This final set of 23 genes was used to evaluate gene

expression and sequence variation among our three lines. For downstream analyses, we also identified

alleles of these 23 loci in a genome assembly of *D. americana* Anderson (15010-0951.00, also referred to

as A01) generated by Kim et al. 2020, using the BLASTn function of BLAST+ version 2.6.0 (Camacho et al.

2009) with 1/-1 match/mismatch scoring parameters, and retaining the top BLAST hits for each locus.

Statistical Analyses

All statistical analyses and figure construction in this study was performed with R version 3.4.3.

For the unperfumed 4x4 mating dataset, we used Kruskal-Wallis tests to compare copulation success

within 3 hours of observation with male or female species identity (one test for each sex). In addition,

we used Wilcoxon rank sum tests to perform post-hoc pairwise comparisons for each sex to evaluate

which species differed from one another in female acceptance of males or in male courtship success of

females. We also calculated the reproductive isolation index RI_1 for prezygotic barriers as defined by

Sobel and Chen (2014). This index is appropriate for comparisons between no-choice tests, and

described by the equation $RI_1 = 1 - $ (heterospecific success)/(conspecific success).

In 1x1 mating trials, mating rate was evaluated using a logistic regression with a chi-square test.

For copulation latency, a 2-way ANOVA was performed with latency in minutes as the dependent

variable, to test for the effects of male identity, female identity, and their interaction. This was followed

by a post-hoc Tukey HSD in which we assessed which male identities (populations) were specifically

different for copulation latency. For each of the three courtship behaviors (displays, tappings and

lickings), we converted the count data for each trial to a rate per unit time within a trial, by dividing the

recorded counts for each trait by the latency to copulation (in minutes) or, if no copulation occurred, the

maximum amount of observed time (180 minutes). Describing male behaviors in terms of rates accounts

for differences among males in their courtship efficiency; for example, it allows us to differentiate males

that performed fewer courtship behaviors because they were rapidly accepted by a female, from males

that displayed lower courtship intensity across the total 3 hour monitored period. For each of the three

behavior rates (display-rate, tap-rate, lick-rate) as dependent variables, we performed an ANOVA with

male identity, female identity, and their interaction as independent variables.

The perfumed 4x4 mating experiment was analyzed using planned contrasts that compared the

mating success of males that had the same target identity but different (con- versus hetero-specific)

donor perfumes, when paired with SC *D. americana* females. This enabled us to specifically assess the

effects of donor perfume variation on mating success of a given target male species. Each pairwise test

was performed using a non-parametric Mann-Whitney U test.

Our primary analyses of differences in CHC composition were done after summarizing major

axes of variation in CHC phenotypes using a set of Principal Component Analyses (PCA). Separate PCAs

were performed on each CHC experiment within this study: one PCA for the initial (unperfumed)

parental populations (denoted U-PCA), and one PCA for each of the perfumed experimental pairs—Nov-

SC (N-PCA) and NE-SC (A-PCA)(Figure S2). Within each dataset, a one-way ANOVA was used on each of

the first 3 PCs to examine effects of sex and species (for the unperfumed dataset) or male perfume

identity (in each perfuming study) on CHC composition. Additionally, for the perfumed datasets, T-tests

were used to compare differences in PC values between target males that were paired with different

(con- and hetero-specific) donor males to assess the effects of our perfuming manipulation on major

axes of CHC variation. Factor loadings for N-PC an A-PC analyses, as well as individual compound

differences in perfuming pairs can be found in in the supplement (Tables S10, S11, S12 and Figures S1,

S2).

For all gene expression analyses, expression is quantified in transcripts per million (TPM), and

therefore is normalized within each sample. Here we analyzed datasets separated by sex, as Davis and

Moyle (2020) showed that the majority of genes have differential expression based on sex, whereas our

primary interest here is in differences between species that might be implicated in female mating

choices and male CHC profile variation. With the dataset for each sex, we ran one-way ANOVAs on TPM

of every expressed locus (11301 genes total)—including our 23 target candidate genes—to determine

loci for which gene expression varied by population. Each gene was ranked according to their resulting

F-value (Table 3; un-corrected P-values are given in the supplement (Table S13)). This allowed us to

evaluate which candidate genes had more pronounced expression differences between populations,

compared to all other genes in the dataset. Candidate genes with greater differential expression

between populations than the majority of the transcriptome could suggest these genes impact observed

phenotypic differences, even if the number of tests performed make finding significance at alpha = 0.05

difficult.

We also estimated rates of nonsynonymous to synonymous substitutions (d_N/d_S) for 4397 genes

in this transcriptome-wide dataset, to quantify molecular evolution transcriptome-wide in this group

and to evaluate evidence for positive selection specifically in candidate loci. We used a pipeline for

obtaining genome-wide estimates of d_N/d_S modified from Wu et al. 2018. First, as in Davis and Moyle

2020, transcript sequences from each population were aligned to the $D.\ virilis$ reference genome

(Flybase version 1.7, www.flybase.org) to identify loci with expression in the populations studied here.

Then, for each gene with transcripts present in all populations and coding sequence (CDS) annotated in

the $D.\ virilis$ reference, a consensus fasta was generated for each line (population) for the longest splice

variant of a given gene. These consensus fasta sequences were then aligned to each other using PRANK

(Löytynoja & Goldman, 2005) with codons enforced and ten bootstrap replicates, allowing us to obtain

orthologous gene sequence alignments among the three populations used here and the $D.\ virilis$

outgroup. We then calculated d_N/d_S from these aligned sequences in PAML v4.9 (Yang 2007) using model

M0 in CodeML—a maximum likelihood model for codon substitution. As PAML uses a tree-based model

for computing d_N/d_S that is sensitive to use of the correct gene tree for a given gene (Mendes et al.,

2016) consensus gene trees were constructed individually for each gene using RAxML v8.3 with the

GTRGAMMA model with 100 bootstraps (Stamatakis, 2014). Lastly, for both CG17821 and CG18609—

genes with shared derived alleles for NE *D. americana* and *D. novamexicana*—we also computed the

d_N/d_S value using only sequences for *D. virilis* and SC *D. americana* to determine if observed estimates of

protein evolution in these genes is limited to the derived pair or if this pattern is consistent across the

clade. Full results of d_N/d_S for each gene are reported in the supplementary file, with annotation of any

known function of orthologs (according to Flybase) for the 100 loci with the highest estimated rates of

protein evolution.

To evaluate possible introgression of candidate gene alleles between populations in this group,

we constructed additional gene trees for each candidate gene using orthologs from the Anderson IN *D.*

americana genome (from Kim et al. 2020) in place of our western (NE) *D. americana* sequences using

RAxML v8.3 (Stamatakis, 2014), with alignments created using PRANK. Additional gene trees were

generated in the same manner for neighboring genes within 50 kbp around CG17821/CG18609, to

estimate size of a window of shared ancestry.

To assess evidence for genome-wide introgression, we used our transcriptome data in

conjunction with the *D. virilis* genome to generate a genome-wide set of SNPs for our three focal

populations and the *D. virilis* outgroup. To do so, we used bcftools (Li 2011) to generate a VCF (variant

calling format) file and call SNP sites between the 4 taxa after filtering for low depth and masking

heterozygous sites from individual taxa. With this, we calculated Patterson's *D*-statistic (Durand et al.

2011) using the Dtrios program from Dsuite (Malinsky et al. 2020) to calculate the 3 taxa *D*-statistic as

well as an overall *P*-value using standard errors generated from the default 20 jackknife blocks. We used

(((SC americana, NE americana), novamexicana), virilis)(figure 3B top tree) as the expected species tree

topology for this analysis.

Acknowledgements

We would like to thank Matthew Gibson for help with implementing molecular evolution analyses, Zinan

Wang for advice about candidate gene selection, Bernard Kim for sequencing the *Drosophila americana*

genome and providing assemblies, Scott Pletcher for sharing hydrocarbon analysis software, and

Jonathan Massey for connecting JSD and LCM with JYY.

This work was supported by the IU Department of Biology (LCM, JSD), the Department of Defense

United States Army Research Office (Grant No. W911NF1610216) and the National Institutes of Health

(Grant No. 1P20GM125508) awarded to JYY. The GC/ MS analysis was performed in the UHM Microbial

Genetics and Analytical Laboratory (supported by NIH Grant No. 1P20GM125508).

Table 1: Copulation rates and latency for each cross in unperfumed 4x4 and 1x1 mating trials. Isolation
index included is RI_1 as defined by Sobel and Chen (2014), and uses values from the 4x4 mating trials.

Female species	Male species	4x4 trials	1x1 trials		
		Mating rate[a] (mean ± SE)	Mating rate[a] (mean)	Cop. latency (mean ± SE)[b]	Isolation Index (RI_1)
D. novamexicana	*D. novamexicana*	0.90 ± 0.06	1.00	69.89 ± 22.73	
	NE *D. americana*	0.69 ± 0.12	0.60	147.48 ± 27.00	0.233
	SC *D. americana*	0.71 ± 0.18	0.60	83.63 ± 39.70	0.211
NE *D. americana*	*D. novamexicana*	1.00 ± 0	0.20	158.33 ± 21.67	0
	NE *D. americana*	1.00 ± 0	0.60	123.46 ± 29.01	
	SC *D. americana*	0.50 ± 0.18	0.60	99.42 ± 34.74	0.5
SC *D. americana*	*D. novamexicana*	0.00 ± 0	0.00	180 ± 0	1
	NE *D. americana*	0.40 ± 0.14	0.60	99.42 ± 34.74	0.52
	SC *D. americana*	0.85 ± 0.06	0.80	50.32 ± 33.07	

[a] n = 5, rate is a proportion out of maximum possible matings (4 or 1)
[b] n = 5, units are in minutes. Unmated trials are scored as a latency of 180 min—the maximum time observed.

806
807

808 **Table 2: Mean copulation success (number of matings out of 4 per trial) with SC *D. americana* females**
809 **for each perfuming identity in manipulated (perfumed) 4x4 mating assays.**

Target male	Donor male	Copulation success (Mean ± SE)	Pairwise Mann-Whitney U test	
			U-score	*P*-value
Nov-SC pair				
D. novamexicana	*D. novamexicana*	0.25 ± 0.15	} 0	0.0247
D. novamexicana	SC *D. americana*	0.94 ± 0.063		
SC *D. americana*	SC *D. americana*	1.00 ± 0	} 0	0.018
SC *D. americana*	*D. novamexicana*	0.19 ± 0.063		
NE-SC pair				
NE *D. americana*	NE *D. americana*	0.44 ± 0.12	} 5.5	0.54
NE *D. americana*	SC *D. americana*	0.56 ± 0.12		
SC *D. americana*	SC *D. americana*	1.00 ± 0	} 0	0.018
SC *D. americana*	NE *D. americana*	0.50 ± 0.063		

Note: *n* = 5 for all crosses

810
811

812 **Table 3: Molecular evolution (d_N/d_S) and species differences in quantitative gene expression for 23**
813 **candidate genes associated with CHC function, analyzed separately by sex, with *F*-value of ANOVA test**
814 **and rank compared to all genes in dataset. d_N/d_S of all genes reported in supplementary file.**

Candidate gene	*D. virilis* ortholog	InterProt domain	d_N/d_S[a]	Males		Females	
				F-value[b]	Rank %ile[b]	*F*-value[b]	Rank %ile[b]
Cyp4g1	GJ15981	Cyp4g	0.0469	0.143	95%	0.158	87%
Cyp4g15	GJ19152	Cyp4g	0.0862	2.07	33%	1.05	38%
Cyt-b5-r	Cyt-b5-r	fatty-acid desaturase	0.0001	2.32	29%	0.736	53%
ifc	GJ15437	fatty-acid desaturase	0.0266	0.122	96%	0.74	52%
CG17928	GJ17306	fatty-acid desaturase	0.133	1.09	56%	0.0233	96%
CG8630	GJ10413	fatty-acid desaturase	0.118	3.48	19%	1.46	27%
CG9743	GJ10408	fatty-acid desaturase	0.0712	2.83	24%	1.75	22%
desat2	desat2	fatty-acid desaturase	0.0780	0.216	93%	0.903	45%
Baldspot	GJ12451	elongase	0.0280	0.814	68%	0.916	45%
bond	GJ24166	elongase	0.0503	0.710	53%	0.949	79%
ELOVL	**GJ24115**	**elongase**	0.0291	**9.61**	**6%**	2.67	14%
CG5326	GJ24664	elongase	0.0081	4.14	16%	0.138	89%
CG17821	**GJ22070**	**elongase**	**0.284**	3.50	19%	**17.2**	**0.8%**
CG18609	**GJ22071**	**elongase**	**0.249**	**8.19**	**7%**	1.85	21%
CG30008	**GJ22296**	**elongase**	0.151	2.19	31%	**4.33**	**7%**
CG31522	GJ24118	elongase	0.0422	0.744	71%	0.380	73%
CG31523	**GJ24117**	**elongase**	0.0410	**6.57**	**9%**	0.0481	95%
CG33110	GJ24167	elongase	0.0178	0.466	83%	0.436	70%
CG6660	**GJ14350**	**elongase**	**0.387**	1.15	54%	1.20	33%
FASN1	GJ21736	fatty-acid-synthase	0.0274	1.42	46%	0.526	64%
FASN2	GJ21725	fatty-acid-synthase	0.0979	1.83	37%	1.02	39%
ebony	GJ14444	pigment	0.0280	0.673	33%	1.15	35%
tan	GJ14875	pigment	0.0269	1.49	45%	0.347	75%

[a]**Bold** genes in this column indicate d_N/d_S values over 1 standard deviation from mean d_N/d_S value of 4397 genes analyzed.
[b]**Bold** genes in these columns indicate genes in the top 10% of differentially expressed genes within sex.

816
817 **Figure 1: Male courtship behavior rates in 1x1 mating assays across all cross types. Points represent**
818 **individual trials, with boxes showing quartiles and the mean as a solid bar. Males (columns) show**
819 **significant differences in display rate and tap rate, and marginal differences in lick rate. Rates did not**
820 **differ based on female identity (rows) or male x female interaction (see results). Bar and whiskers**
821 **indicate mean and standard error (SE).**

823
Figure 2: Cuticular hydrocarbon composition of unmanipulated males and females of each species
stock. A) Relative log-scale abundance of compounds for males (upper) and females (lower), for each
major compound type (line = mean, box = central quartile, whiskers = S.E.). *n* **= 5 samples for each sex**
and line. Test statistics for species differences are in Table S3. B) First two principal components (U-
PC1 and U-PC2) of composite CHC variation among males (circle) and females (triangle), with percent
of total variance explained. Ellipses indicate 90% bivariate normal density for each species-sex group.
Species and sex significantly influenced both U-PC1 and U-PC2.

Figure 3: CHC candidate gene expression, genealogical relationships, and sequence variation among *D.*
americana **group lines. A) Gene expression variation showing mean and standard error between focal**

species lines for candidate genes with the largest differences in either males (upper, circles) or

females (lower, triangles). Gene expression is measured in transcripts per million (TPM, n = 3 for each

sample). B) Gene tree topologies for all 23 candidate genes using either western (NE) or eastern (IN)

populations for NE *D. americana*. C) Schematic of loci (green/blue boxes) in the genomic window

surrounding candidates CG17821 and CG18609. Blue boxes indicate a genealogy that groups the NE *D.*

americana population with *D. novamexicana*; gene topology of shaded blue region displayed below

line. Green boxes indicate loci with genealogies matching the expected species tree. D) 3' terminal

nucleotides and occurrence of inferred ancestral and derived (truncated) allele in CG17821 among

lineages with sequence data.

References

Bartelt, R. J., Armold, M. T., Schaner, A. M., & Jackson, L. L. (1986). Comparative Analysis of Cuticular
Hydrocarbons in the Drosophila Virilis Species Group. *Comparative Biochemistry and Physiology*,
83B(4), 731–742.

Basolo, A. L. (1995). Phylogenetic evidence for the role of a pre-existing bias in sexual selection.
Proceedings. Biological Sciences, *259*(1356), 307–311.

Baxter, C., Mentlik, J., Shams, I., & Dukas, R. (2018). Mating success in fruit flies: courtship interference
versus female choice. *Animal Behaviour*, *138*, 101–108.

Billeter, J.C., Atallah, J., Krupp, J. J., Millar, J. G., & Levine, J. D. (2009). Specialized cells tag sexual and
species identity in Drosophila melanogaster. *Nature*, *461*(7266), 987–991.

Blight, W. C., & Romano, A. (1953). Notes on a Breeding Site of *Drosophila americana* Near St . Louis ,
Missouri. *The American Naturalist*, 87(833), 111–112.

Blomquist G. J., Bagnères A. G. (2010) Insect hydrocarbons: biology. biochemistry, and chemical ecology.
Cambridge University Press, New York

Bontonou, G., & Wicker-Thomas, C. (2014). Sexual Communication in the Drosophila Genus. *Insects*,
5(2), 439–458.

Butlin, R., Debelle, A., Kerth, C., Snook, R. R., Beukeboom, L. W., Castillo Cajas, R. F., Diao, W., Maan, M.
E., Paolucci, S., Weissing, F. J., van de Zande, L., Hoikkala, A., Geuverink, E., Jennings, J., Kankare,
M., Knott, K. E., Tyukmaeva, V. I., Zoumadakis, C., Ritchie, M. G., … Schilthuizen, M. (2012). What
do we need to know about speciation? *Trends in Ecology & Evolution*, *27*(1), 27–39.

Caletka, B. C., & McAllister, B. F. (2004). A genealogical view of chromosomal evolution and species
delimitation in the Drosophila virilis species subgroup. *Molecular Phylogenetics and Evolution*,
33(3), 664–670.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009).
BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*, 1–9.

Chung, H., & Carroll, S. B. (2015). Wax, sex and the origin of species: Dual roles of insect cuticular
hydrocarbons in adaptation and mating. *BioEssays*, 822–830.

Chung, H., Loehlin, D., Dufour, H., Vaccarro, K., Millar, J. G., & Carroll, S. B. (2014). A Single Gene Affects
Both Ecological Divergence and Mate Choice in Drosophila. *Science*, *257082*.

Combs, P. A., Krupp, J. J., Khosla, N. M., Bua, D., Petrov, D. A., Levine, J. D., & Fraser, H. B. (2018). Tissue-
Specific cis-Regulatory Divergence Implicates eloF in Inhibiting Interspecies Mating in Drosophila.
Current Biology, *28*(24), 3969-3975.e3.

Coyne, J. A., Kim, S. Y., Chang, A. S., Lachaise, D., & Elwyn, S. (2002). Sexual isolation between two sibling
species with overlapping ranges: Drosophila santomea and Drosophila yakuba. *Evolution*, *56*(12),
2424–2434.

Coyne, J. A., & Oyama, R. (1995). Localization of pheromonal sexual dimorphism in Drosophila
melanogaster and its effect on sexual isolation. *Proceedings of the National Academy of Sciences of*
the United States of America, *92*(21), 9505–9509.

Coyne, J. A., & Orr, H. A., 2004. Speciation. SinauerAssociates, Sunderland, MA

Coyne, J. A., Crittenden, A. P., & Maht, K. (1994). Genetics of a Pheromonal Difference Contributing to
Reproductive Isolation in Drosophila. *Science*, *265*(5177), 1461–1464.

Curtis, S., Sztepanacz, J. L., White, B. E., Dyer, K. a., Rundle, H. D., & Mayer, P. (2013). Epicuticular
Compounds of Drosophila subquinaria and D. recens: Identification, Quantification, and Their Role
in Female Mate Choice. *Journal of Chemical Ecology*, *39*(5), 579–590.

Davis, J. S., & Moyle, L. C. (2020). Constitutive and plastic gene expression variation associated with
desiccation resistance differences in the Drosophila americana species group. *Genes*, *11*(146).

Davis, J. S., & Moyle, L. C. (2019). Desiccation resistance and pigmentation variation reflects bioclimatic
differences in the Drosophila americana species complex. *BMC Evolutionary Biology*, *19*(1), 1–14.

Durand, E. Y., Patterson, N., Reich, D., & Slatkin, M. (2011). Testing for ancient admixture between
closely related populations. *Molecular Biology and Evolution*, *28*(8), 2239–2252.

Dyer, K. a, White, B. E., Sztepanacz, J. L., Bewick, E. R., & Rundle, H. D. (2014). Reproductive character
displacement of epicuticular compounds and their contribution to mate choice in Drosophila
subquinaria and Drosophila recens. *Evolution; International Journal of Organic Evolution*, *68*, 1163–
1175.

Findlay, G. D., Yi, X., MacCoss, M. J., & Swanson, W. J. (2008). Proteomics reveals novel Drosophila
seminal fluid proteins transferred at mating. *PLoS Biology*, 6(7), 1417–1426.

Franssen, S. U., Nolte, V., Tobler, R., & Schlotterer, C. (2015). Patterns of linkage disequilibrium and long
range hitchhiking in evolving experimental drosophila melanogaster populations. *Molecular*
Biology and Evolution, *32*(2), 495–509.

Frentiu, F. D., & Chenoweth, S. F. (2010). Clines in cuticular hydrocarbons in two *Drosophila* species with
independent population histories. *Evolution*, 64(6), 1784–1794.

Gaertner, B. E., Ruedi, E. A., McCoy, L. J., Moore, J. M., Wolfner, M. F., & Mackay, T. F. C. (2015).
Heritable variation in courtship patterns in drosophila melanogaster. *G3: Genes, Genomes,*
Genetics, *5*(4), 531–539.

Gao, Z., Przeworski, M., & Sella, G. (2015). Footprints of ancient-balanced polymorphisms in genetic
variation data from closely related species. *Evolution*, *69*(2), 431–446.

Gaudet, P., Livstone, M. S., Lewis, S. E., & Thomas, P. D. (2011). Phylogenetic-based propagation of
functional annotations within the Gene Ontology consortium. *Briefings in Bioinformatics*, *12*(5),
449–462.

Gibbs, A. G., Fukuzato, F., & Matzkin, L. M. (2003). Evolution of water conservation mechanisms in
Drosophila. *The Journal of Experimental Biology*, 206, 1183–1192.

Havens, J. A., & Etges, W. J. (2013). Premating isolation is determined by larval rearing substrates in
cactophilic *Drosophila mojavensis*. IX. Host plant and population specific epicuticular hydrocarbon
expression influences mate choice and sexual selection. *Journal of Evolutionary Biology*, 26(3),
562–576.

He, Z., Luo, Y., Shang, X., Sun, J. S., & Carlson, J. R. (2019). Chemosensory sensilla of the Drosophila wing
express a candidate ionotropic pheromone receptor. *PLoS Biology*, *17*(5), 1–27.

Hekmat-Scafe, D. S., Scafe, C. R., McKinney, A. J., & Tanouye, M. A. (2002). Genome-Wide analysis of the
odorant-binding protein gene family in Drosophila melanogaster. *Genome Research*, *12*(9), 1357–
1369.

Hoikkala, A., & Lumme, J. (1987). The Genetic Basis of Evolution of the Male Courtship Sounds in the
Drosophila virilis Group. *Evolution*, *41*(4), 827–845.

Hollocher, H., Ting, C. T., Wu, M. L., & Wu, C. I. (1997). Incipient speciation by sexual isolation in
Drosophila melanogaster: Extensive genetic divergence without reinforcement. *Genetics*, *147*(3),
1191–1201.

Hudson, R. R., & Coyne, J. A. (2002). Mathematical Consequences of the Genealogical Species. *Evolution*,
56(8), 1557–1565.

Jezovit JA, Levine JD, Schneider J. Phylogeny, environment and sexual communication across the
Drosophila genus. *J Exp Biol*. 2017;220(Pt 1):42-52.

Kaneshiro, K. Y. (1980). Sexual Isolation, Speciation and the Direction of Evolution. *Evolution*, *34*(3), 437–
444.

Kaneshiro, K. Y. (1976). Ethological Isolation and Phylogeny in the Planitibia Subgroup of Hawaiian
Drosophila. *Evolution*, *30*(4), 740.

Kim, B. Y., Wang, J. R., Miller, D. E., Barmina, O., Delaney, E., Thompson, A., Comeault, A. A., Peede, D.,
Agostino, E. R. R. D., Aguilar, J. M., Haji, D., Matsunaga, T., Armstrong, E. E., Zych, M., et. Al. (2020).
Highly contiguous assemblies of 101 drosophilid genomes Bernard. *BioRxiv*.

Lamb, A. M., Wang, Z., Simmer, P., Chung, H., Patricia, J., States, U., Lansing, E., States, U., Biology, E.,
Program, B., Lansing, E., States, U., Biology, E., Program, B., Arbor, A., States, U., & Wittkopp, P. J.
(2020). ebony affects pigmentation divergence and cuticular hydrocarbons in *Drosophila*
americana and *D. novamexicana*. *Frontier in Ecology and Evolution*, *8*(June), 1–23.

Laughlin, J. D., Ha, T. S., Jones, D. N. M., & Smith, D. P. (2008). Activation of Pheromone-Sensitive
Neurons Is Mediated by Conformational Activation of Pheromone-Binding Protein. *Cell*, *133*(7),
1255–1265.

Leal, W. S. (2013). Odorant reception in insects: Roles of receptors, binding proteins, and degrading
enzymes. *Annual Review of Entomology*, *58*, 373–391.

Legendre, A., Miao, X. X., Da Lage, J. L., & Wicker-Thomas, C. (2008). Evolution of a desaturase involved
in female pheromonal cuticular hydrocarbon biosynthesis and courtship behavior in *Drosophila*.
Insect Biochemistry and Molecular Biology, *38*(2), 244–255.

Löytynoja, A., & Goldman, N. (2005). An algorithm for progressive multiple alignment of sequences with
insertions. *Proceedings of the National Academy of Sciences of the United States of America*,
102(30), 10557–10562.

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. *Bioinformatics*, *27*(21), 2987–
2993.

Malinsky, M., Matschiner, M., & Svardal, H. (2020). Dsuite - fast D-statistics and related admixture
evidence from VCF files. *BioRxiv*, 1–16.

Mas, F., & Jallon, J. M. (2005). Sexual isolation and cuticular hydrocarbon differences between
Drosophila santomea and *Drosophila yakuba*. *Journal of Chemical Ecology*, *31*(11), 2747–2752.

Massey, J. H., Rice, G. R., Firdaus, A. S., Chen, C. Y., Yeh, S. D., Stern, D. L., & Wittkopp, P. J. (2020). Co-
evolving wing spots and mating displays are genetically separable traits in *Drosophila*. *Evolution*,
74(6), 1098–1111.

Massey, J. H., Akiyama, N., Bien, T., Dreisewerd, K., Wittkopp, P. J., Yew, J. Y., & Takahashi, A. (2019).
Pleiotropic Effects of ebony and tan on Pigmentation and Cuticular Hydrocarbon Composition in
Drosophila melanogaster. *Frontiers in Physiology*, 10.

Matsuo, T., Sugaya, S., Yasukawa, J., Aigaki, T., & Fuyama, Y. (2007). Odorant-binding proteins OBP57d
and OBP57e affect taste perception and host-plant preference in Drosophila sechellia. *PLoS*
Biology, *5*(5), 0985–0996.

McAllister, B. F. (2002). Chromosomal and allelic variation in *Drosophila americana* : selective
maintenance of a chromosomal cline. *Genome*, *45*(1), 13–21.

McGraw, L. A., Gibson, G., Clark, A. G., & Wolfner, M. F. (2004). Genes Regulated by Mating, Sperm, or
Seminal Proteins in Mated Female Drosophila melanogaster Lisa. *Current Biology*, *14*, 1509–1514.

Morales-Hojas, R., Reis, M., Vieira, C. P., & Vieira, J. (2011). Resolving the phylogenetic relationships and
evolutionary history of the *Drosophila virilis* group using multilocus data. *Molecular Phylogenetics*
and Evolution, *60*(2), 249–258.

Pardy, J. A., Rundle, H. D., Bernards, M. A., & Moehring, A. J. (2018). The genetic basis of female
pheromone differences between *Drosophila melanogaster* and *D. simulans*. *Heredity*, *122*(1), 93–
109.

Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T., Webster, T., &
Reich, D. (2012). Ancient admixture in human history. *Genetics*, *192*(3), 1065–1093.

Rajpurohit, S., Hanus, R., Vrkoslav, V., Behrman, E. L., Bergeland, A. O., Petrov, D. A., Cvacka, J., &
Schmidt, P. S. (2017). Adaptive dynamics of cuticular hydrocarbon in *Drosophila*. Journal of
Evolutionary Biology, 30(1), 66–80.

Reis, M., Vieira, C. P., Lata, R., Posnien, N., & Vieira, J. (2018). Origin and consequences of chromosomal
inversions in the virilis group of Drosophila. *Genome Biology and Evolution*, *10*(12), 3152–3166.

Ritchie, M. G. (2007). Sexual Selection and Speciation. *Annual Review of Ecology, Evolution, and*
Systematics, *38*(1), 79–102.

Roelofs, W. L., Liu, W., Hao, G., Jiao, H., Rooney, A. P., & Linn, C. E. (2002). Evolution of moth sex
pheromones via ancestral genes. *Proceedings of the National Academy of Sciences of the United*
States of America, 99(21), 13621–13626.

Ruedi, E. A., & Hughes, K. A. (2008). Natural genetic variation in complex mating behaviors of male
Drosophila melanogaster. *Behavior Genetics*, *38*(4), 424–436.

Schemske, D. W. (2000). UNDERSTANDING THE ORIGIN OF SPECIES. *Evolution*, *54*(3), 1069–1073.

Schluter, D. (2009). Evidence for Ecological Speciation and Its Alternative. *Science*, *323*(5915), 737–741.

Seddon, N., Botero, C. A., Tobias, J. A., Dunn, P. O., MacGregor, H. E. A., Rubenstein, D. R., Uy, J. A. C.,
Weir, J. T., Whittingham, L. A., & Safran, R. J. (2013). Sexual selection accelerates signal evolution
during speciation in birds. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1766).

Serrato-Capuchina, A., Schwochert, T. D., Zhang, S., & Roy, B. (2020). Pure species discriminate against
hybrids in the *Drosophila melanogaster* species subgroup. *bioRxiv*

Shahandeh, M. P., Pischedda, A., & Turner, T. L. (2018). Male mate choice via cuticular hydrocarbon
pheromones drives reproductive isolation between Drosophila species. *Evolution*, *72*(1), 123–135.

Smadja, C., & Butlin, R. K. (2009). On the scent of speciation: The chemosensory system and its role in
premating isolation. *Heredity*, *102*(1), 77–97.

Sobel, J. M., & Chen, G. F. (2014). Unification of methods for estimating the strength of reproductive
isolation. *Evolution*, *68*(5), 1511–1522.

Spieth, H. T. (1951). Mating behavior and sexual isolation in the Drosophila virilis species group.
Behaviour, *3*(2), 105–145.

Sramkoski, L. L., Mclaughlin, W. N., Cooley, A. M., Yuan, D. C., & Wittkopp, P. J. (2020). Genetic
architecture of a body color cline in *Drosophila americana*. *BioRxiv*.

Stamatakis, A. (2014). RAxML version 8: A tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics*, *30*(9), 1312–1313.

Sun, J. S., Xiao, S., & Carlson, J. R. (2018). The diverse small proteins called odorant-binding proteins.
Open Biology, 8(12).

Szafer-Glusman, E., Giansanti, M. G., Nishihama, R., Bolival, B., Pringle, J., Gatti, M., & Fuller, M. T.
(2008). A Role for Very-Long-Chain Fatty Acids in Furrow Ingression during Cytokinesis in
Drosophila Spermatocytes. *Current Biology*, *18*(18), 1426–1431.

Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: Improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific gap
penalties and weight matrix choice. *Nucleic Acids Research*, *22*(22), 4673–4680.

Tomaru, M., Doi, M., Higuchi, H., & Oguma, Y. (2000). Courtship song recognition in the *Drosophila*
melanogaster complex: Heterospecific songs make females receptive in *D. melanogaster*, but not
in D. sechellia. *Evolution*, *54*(4), 1286–1294.

van Doorn, G. S., Edelaar, P., & Weissing, F. J. (2009). On the Origin of Species by Natural and Sexual
Selection. *Science*, *581*(December), 1704–1708.

Vieira, F. G., Sánchez-Gracia, A., & Rozas, J. (2007). Comparative genomic analysis of the odorant-binding
protein family in 12 Drosophila genomes: Purifying selection and birth-and-death evolution.
Genome Biology, *8*(11).

Wang, Z., Cui, Y., Song, L., & Fang, F. (2016). The hysteresis model of piezoelectric micro-positioning
stage based on threshold optimization. *PLoS Biology*, *38*(3), 437–440.

Wilkins, M. R., Seddon, N., & Safran, R. J. (2013). Evolutionary divergence in acoustic signals: Causes and
consequences. *Trends in Ecology and Evolution*, *28*(3), 156–166.

Wu, C. I., Hollocher, H., Begun, D. J., Aquadro, C. F., Xu, Y., & Wu, M. L. (1995). Sexual isolation in
Drosophila melanogaster: A possible case of incipient speciation. *Proceedings of the National*
Academy of Sciences of the United States of America, *92*(7), 2519–2523.

Wu, M., Kostyun, J. L., Hahn, M. W., & Moyle, L. C. (2018). Dissecting the basis of novel trait evolution in
a radiation with widespread phylogenetic discordance. *Molecular Ecology*,

Yang, Z. (2007). PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution*,
24(8), 1586–1591.

Yew, J. Y., & Chung, H. (2017). *Drosophila* as a holistic model for insect pheromone signaling and
processing. *Current Opinion in Insect Science*, *24*, 15–20.

Yukilevich, R., & Peterson, E. K. (2019). The evolution of male and female mating preferences in
Drosophila speciation. *Evolution*, 73(9), 1759–1773.

Yukilevich, R., & True, J. R. (2008). Incipient sexual isolation among cosmopolitan *Drosophila*
melanogaster populations. *Evolution*, *62*(8), 2112–2121.

Zdobnov, E. M., Tegenfeldt, F., Kuznetsov, D., Waterhouse, R. M., Simao, F. A., Ioannidis, P., Seppey, M.,
Loetscher, A., & Kriventseva, E. V. (2017). OrthoDB v9.1: Cataloging evolutionary and functional
annotations for animal, fungal, plant, archaeal, bacterial and viral orthologs. *Nucleic Acids*
Research, *45*(D1), D744–D749.

1051