Parallel evolution of direct development in frogs – Skin and thyroid gland development in

African Squeaker Frogs (Anura: Arthroleptidae: *Arthroleptis*)

Benjamin Naumann[1],*, Susan Schweiger[1], Jörg U. Hammel[2], Hendrik Müller[1,3,4],*

[1]Institut für Zoologie und Evolutionsforschung, Erbertstraße 1, 07743 Jena, Germany

[2]Helmholtz-Zentrum Geesthacht, Zentrum für Material- und Küstenforschung, Außenstelle am

DESY, Gebäude 66, Notkestraße 66, 22607 Hamburg, Germany

[3]Zentralmagazin Naturwissenschaftlicher Sammlungen, Martin-Luther-Universität Halle-

Wittenberg, Domplatz 4, 06108 Halle (Saale), Germany

[4]Department of Life Sciences, The Natural History Museum, Cromwell Road, London SW7

5BD, United Kingdom

*corresponding author: benjamin.naumann@uni-jena.de; hendrik.mueller@zns.uni-halle.de

Key words: *Cardioglossa*, tadpole, terrestrialization, heterochrony, developmental constraints

Abstract

Cases of parallel evolution offer the possibility to identify adaptive traits and to uncover

developmental constraints on the evolutionary trajectories of these traits. The independent

evolution of direct development, from the ancestral biphasic life history in frogs is such a case of

parallel evolution. In frogs, aquatic larvae (tadpoles) differ profoundly from their adult forms and

exhibit a stunning diversity regarding their habitats, morphology and feeding behaviors. The

transition from the tadpole to the adult is a climactic, thyroid hormone (TH)-dependent process of

profound and fast morphological rearrangement called metamorphosis. One of the organ systems

that experiences the most comprehensive metamorphic rearrangements is the skin. Direct-

developing frogs lack a free-swimming tadpole and hatch from terrestrial eggs as fully formed

froglets. In the few species examined, development is characterized by the condensed and

transient formation of some tadpole-specific features and the early formation of adult-specific

features during a "cryptic" metamorphosis. In this study we show that skin in direct-developing

African squeaker frogs (*Arthroleptis*) is also repatterned from a tadpole-like to an adult-like

histology during a cryptic metamorphosis. This repatterning correlates with an increase of thyroid

gland activity. A comparison with data from the Puerto Rican coqui (*Eleutherodactylus coqui*)

reveals that direct development might have evolved in parallel in these frogs by a comparable

heterochronic shift of thyroid gland activity. This suggests that the development of many adult-

features is still constrained by the ancestral dependency on thyroid hormone signaling.

Introduction

How do developmental constraints influence phenotypic evolution? A promising way to approach this question is studying cases of parallel evolution (Schluter, Clifford, Nemethy, & McKinnon, 2004). Parallel evolution can be defined as the independent origin of similar (derived) traits in two or more taxa sharing a common ancestry and bauplan (Futuyma & Kirkpatrick, 2017; Simpson, 1961). Investigating patterns of parallel evolution allows to identify adaptive traits and to uncover developmental constraints on the evolutionary trajectories of these traits (Schluter et al., 2004). A potential case of parallel evolution is the repeated, independent origin of direct development within frogs and other amphibians (San Mauro et al., 2014; Schweiger, Naumann, Larson, Möckel, & Müller, 2017; Thibaudeau & Altig, 1999; Wake & Hanken, 1996). Direct development is characterized by the loss of an aquatic larval phase (Hall & Olson, 2003) and may have evolved in response to environmental conditions that restricted the availability of suitable habitats for aquatic development (Goin & Goin, 1962; Liedtke et al., 2017; Müller et al., 2013; ten Brink, Onstein, & de Roos, 2020).

In frogs, direct development evolved several times independently (Figure 1A), making them a suitable model system to unravel developmental constraints that underlay patterns of parallel evolution (Goldberg, Candioti, & Akmentins, 2012; Heinicke et al., 2009). The ancestral biphasic life history of frogs includes a free-swimming larval stage called tadpole (McDiarmid & Altig, 1999). In contrast to other amphibian larvae, tadpoles differ extremely from their adult forms and exhibit a stunning diversity regarding their habitats, morphology and feeding behaviors (Altig & McDiarmid, 1999). In biphasic anurans, embryonic development from a fertilized egg to a tadpole happens more or less gradually, similar to other vertebrate groups. In contrast, the

transition from the tadpole to the adult is a climactic, thyroid hormone (TH)-dependent process of

profound and fast morphological rearrangement called metamorphosis (Shi, 1999) (Figure 1B). In

direct developing species, development from the embryo to the adult frog appears more gradual

without an obvious climactic phase (Callery, Fang, & Elinson, 2001) (Figure 1B). Tadpole-

specific traits such as the larval skeleton and its associated musculature, the cement glands, the

lateral line system and the coiled intestine are nearly or completely reduced (Callery et al., 2001;

Hanken, Klymkowsky, Alley, & Jennings, 1997). However, morphological studies on the

embryonic development of direct developing frogs are limited to only a few species (Goldberg et

al., 2012; Goldberg, Taucce, Quinzio, Haddad, & Candioti, 2020; Hanken, Klymkowsky,

Summers, Seufert, & Ingebrigtsen, 1992; Kerney, Meegaskumbura, Manamendra□Arachchi, &

Hanken, 2007; Schweiger et al., 2017). The only species in which direct development has been

investigated in greater detail is the Puerto Rican coqui, *Eleutherodactylus coqui* Thomas, 1966

(thyroid gland: Laslo, Denver, & Hanken, 2019; Callery & Elinson, 2000; Elinson & Fang, 1998;

Jennings & Hanken, 1998; Fang & Elinson, 1996; limb development: Hanken et al., 2001; cranial

development: Hanken et al., 1997; Kerney, Gross, & Hanken, 2010; Olsson, Moury, Carl,

Håstad, & Hanken, 2002; Schlosser, Kintner, & Northcutt, 1999; Schlosser & Roth, 1997; gross

embryonic development: Townsend & Stewart, 1985).

Although direct-developing frogs have departed profoundly from their ancestral ontogeny

(Hanken et al., 1997; Schlosser, 2008; Schweiger et al., 2017), studies in *E. coqui* have shown

that several developmental events are still under the control of TH (Callery & Elinson, 2000;

Hanken & Summers, 1988) and that it undergoes a cryptic metamorphosis (Callery & Elinson,

2000; Ziermann & Diogo, 2014). The ontogenetic repatterning seen in direct-developing frogs is

thought to be the result of changes in the expression of TH, which leads to changes in timing of

TH-dependent events during development ("heterochronic shift" hypothesis). At least some

developmental events are thought to have become decoupled from TH-regulation ("loss of

constraint" hypothesis), but so far data are only available for *E. coqui* and the general influence

of TH during the ontogeny of other direct-developing frogs remains unclear.

An interesting system to study putative developmental constraints on the evolution of direct

development is the skin. The frog skin hosts a complex microbiome and plays major roles in

immune response, pathogen defense, respiration, osmoregulation, camouflage and even

reproduction (Douglas, Hug, & Katzenback, 2020; Fernandes et al., 2011; Huang et al., 2016;

Katz, 1986; Varga, Bui-Marinos, & Katzenback, 2019). Among all tissues that remodel during

metamorphosis the skin exhibits the most extreme changes regarding histology and gene

expression (Yoshizato, 1992). It is exposed to completely different environments (aquatic vs.

terrestrial) depending on life history phases and often exhibits tadpole- and adult-specific

adaptations (Quinzio & Goldberg, 2019). The skin of pre-metamorphic tadpoles is a mostly two-

layered epithelium with unicellular glands and an outer mucus layer (Duellman & Trueb, 1994).

The underlying dermis consists of a stratum compactum and some melanophores in the

mesenchyme beneath (Figure 1C). During pro-metamorphosis, the cells of the epidermis

degenerate except for cells attached to the basal lamina. These basal cells start to proliferate

during metamorphic climax and build up the post-metamorphic five- to seven-layered adult frog

epidermis with an outer keratinized cell layer (stratum corneum) and different multicellular gland

types. The underlying dermis contains an outer (exterior to the stratum compactum) and an inner

(interior to the stratum compactum) melanophore layer (Figure 1C) (Fox, 1985; Kinoshita &

Sasaki, 1994; Tamakoshi, Oofusa, & Yoshizato, 1998). Metamorphic changes in biphasic species

are initiated by thyroid hormone (TH) production (Brown & Cai, 2007; Kulkarni, Singamsetty, &

Buchholz, 2010; Tata, 2006). Thyroid hormone is also involved in several developmental

processes in *E. coqui* indicating the presence of a cryptic metamorphosis in direct developing

species (Callery & Elinson, 2000; Hanken et al., 1992; Kulkarni et al., 2010; Schlosser, 2008).

In this study, we provide detailed data on skin development in direct developing African

Squeaker frogs (*Arthroleptis*) using histological and immuno-histochemical techniques. To

evaluate if potential changes in the skin and thyroid gland histology of *Arthroleptis* are due to

direct development or instead shared by biphasic Arthroleptidae, we investigated the same tissues

in a *Cardioglossa* tadpole, the sister genus of *Arthroleptis*. Additionally, we describe thyroid

gland development in *Arthroleptis* and infer its activity based on morphology and morphometric

measurements.

Results

Many aspects of the development of the embryonic pigmentation pattern, skin and thyroid glands

are similar in *A. wahlbergii* and *A. xenodactyloides*. Differences between the two species are

mentioned where present.

Development of the pigmentation pattern in *Arthroleptis* embryos

Embryos until mid TS5 lack any obvious, externally visible pigmentation and have a white-

yellowish color (Figure 2A; see also Schweiger et al., 2017). However, very few melanophores

were found in the epidermal layer in histological sections of an embryo at TS4 (see Figure 3A4).

At late TS5/early TS6 melanophores are recognizable extending dorso-ventrad until the ventral

border of the eye in the head region and until the dorsal-most quarter of the lateral body wall in

the trunk region (Figure 2B). Melanophores at this stage are spindle-shaped with long, thin

extensions (Figure 2B`). From late TS7 to TS8, melanophores continue to extend ventrad in the

head and the trunk regions (Figure 2C, D) until they completely cover the lateral body wall from

TS9 on (Figure 2E, F). At this stage, the body wall has completely enclosed the yolk (Figure

2E3). At TS11, the density of melanocytes of the head region starts to increase resulting in a

darkening of the skin (Figure 2G). Additionally, another type of melanophores becomes visible

all over the body. These melanophores are smaller, with a more spherical shape and with only a

few short or no extensions (Figure 2G`). At TS12, melanocyte density continues to increase

dorso-ventrad until the whole lateral body wall is heavily pigmented at TS15 (Figure 2H-K).

In summary, pigmentation development is characterized by a first dorso-ventrad wave of

melanophore development starting at late TS5 and a second wave made up by a morphologically

different type of melanophores starting at TS11.

Skin development in *Arthroleptis*

TS4 (Figure 3A) – The epidermis is single-layered with a sometimes indistinct border between

the underlying cells and tissues. In the dorsal body region, cells are large and cuboid with

spherical nuclei. Ventrally, cells are flattened and nuclei are ovoid to spindle-shaped. The cells

possess high amounts of intracellular yolk granules and some seem to have large vacuoles with

dark pigments. Ciliated cells are found in a scattered pattern all over the embryo. Very few

melanophores can be found intercalating between epidermal cells. A dense, undifferentiated

mesenchyme lies beneath the epidermis.

TS5 (Figure 3B) – The dorsal epidermis of the head as well as of the dorso-lateral trunk is two-

layered while the ventral epidermis of the trunk, covering the yolk sac, is single-layered. Most

cells are slightly flattened with ovoid nuclei. The amount of intracellular yolk granules has

decreased compared to TS4 but is still high. The number of ciliated cells has increased compared

to TS4. A sharp border between epidermal cells and the underlying mesenchyme is recognizable.

TS6 and TS7 (Figure 3C, D) – The number of epidermal layers has not changed compared to

TS5. Some unicellular mucus glands are recognizable. The amount of intracellular yolk granules

has further decreased compared to previous stages. A higher number of melanophores with long

cytoplasmic extensions are present in the mesenchyme directly beneath the epidermis.

TS8 (Figure 3E) – The number of epithelial layers is similar to previous stages. Epidermal cells

in both layers are flattened and show spindle-shaped nuclei. Some hypertrophied cells with a

granular content that might represent unicellular glands are present in the skin of the head.

Intracellular yolk granules are only detectable in the trunk epidermis. In the head, a distinct

stratum compactum is present beneath the epidermis with a dense layer of inner melanophores

interior to it. In the trunk, the stratum compactum is most distinct in the dorsal region and

becomes gradually less obvious in lateral and ventral areas. This pattern is mirrored by the

density of associated inner melanophores.

TS11 (Figure 3F) –The epidermis has become single-layered again in the head and dorsolateral

trunk region and epidermal cells exhibit extremely flattened nuclei. In some scattered areas of the

dorsal head, the epidermis is still two-layered. Intracellular yolk granules and ciliated cells are

completely absent. Multicellular gland primordia, consisting of several compact cells, can be

found all over the body. In the head, a few outer melanocytes are present between the stratum

compactum and basal epidermal cells. In the trunk, the stratum compactum appears denser

compared to the previous stage and the density of inner melanophores has increased. It is still

lowest in the ventral trunk region.

TS13 (Figure 3G) – The skin exhibits many features of an adult frog skin. The epidermis is two -

to three-layered. The majority of the basal epidermal cells are cuboid. The apical cell layer is

flattened and is stained dark-blue in Azan-stained section, indicating an increased keratinization

and the presence of a stratum corneum. The stratum corneum is more obvious in the head

compared to the trunk epidermis. Multicellular mucus and granular glands can be found in

various regions of the body. They are localized within a thin stratum spongiosum beneath the

basal epidermal cell layer. Mucus glands are outlined by a single cell layer with an inner lumen.

Granular glands are recognizable as large sacs filled with many granulated cells. The density of

outer melanophores within the now developed stratum spongiosum has increased compared to

previous stages.

TS15 (Figure 3H) – Hatching occurs at this stage and the skin of the froglet is similar to the skin

of an adult *Arthroleptis* (Supplementary material 2). The thickness of the epidermis varies from

two - to five cell layers but is three-layered in most areas. In the trunk region, the dark blue-

stained stratum corneum is now also recognizable as a distinct cellular layer. Multicellular glands

in the whole skin are more numerous and melanophore density has increased in the ventral trunk

region compared to TS13.

In summary, the embryonic ectoderm and underlying mesenchyme at TS4 differentiated into a

skin consisting of an epidermis and an underlying dermis both exhibiting pre-metamorphic

tadpole-typical features at TS8. At TS11 the apical epidermal layers seem to degenerate to a

certain degree while multicellular gland progenitors appear all over the body. The skin of

embryos at TS13 and TS15 exhibit many features typical for the post-metamorphic frog skin.

PCNA expression during epidermal development in *Arthroleptis*

Metamorphic skin remodeling is based on the degeneration of apical epidermal cells and the proliferation of remaining basal cells building the adult epidermis (Yoshizato, 1992). We therefore investigated the epidermal proliferation pattern in embryonic stages of *A. wahlbergii* using two different antibodies against proliferating cell nuclear antigen (PCNA). Signals from both antibodies were detected in the same tissues of two adjacent serial sections verifying antibody specificity. However, the signal from PCNA-1 antibody was always stronger than from the PCNA-2 antibody.

At TS8, scattered signals of the PCNA-1 antibody are detectable in nuclei of both, epidermal cell layers as well as some mesenchymal cells (Figure 4A, A`). Only a few PCNA-2 positive cells are detectable at this stage (Figure 4B, B`). At TS11, PCNA-1 and PCNA-2 signals are detectable in the majority of epidermal cells, indicating a strong increase in the proliferation rate of the epidermis (Figure 4C, D). A few PCNA-1 and PCNA-2 positive mesenchymal cells are also present. No antibody signals are detectable in the epidermis at TS13 and TS15, indicating a low proliferation rate (Figure 4E-G). In some multicellular glands however, PCNA-1 positive cells can be detected (Figure 4G).

In summary, PCNA reactivity (and therefore cell proliferation rate) of the skin is detectable in a scattered pattern at TS8, has increased tremendously at TS11 and is low or absent at TS 13 and TS15.

Thyroid gland development in *Arthroleptis* embryos

The thyroid glands are paired, ovoid structures located at the lateral edges of the hyoid plate (Figure 5A-C). Primordia are first detectable at TS8. They appear as spherical, condensed cell masses including scattered lumina which lack an epithelial lining. The primordia are located on the left and right side ventrally to the lateral edges of the developing hyoid plate (Figure 5A, D). First developing follicles are recognizable at TS10/11. The cells of the follicular epithelial are cuboid with little cytoplasm and large spherical nuclei (Figure 5E). At later TS11, thyroid glands have elongated, adopting a mature, adult-like morphology (Figure 5B). The follicles have increased in size and number and the follicular epithelium is now organized in a columnar-pattern. Nuclei of follicular epithelial cells are spherical to ovoid-shaped. Some colloid, recognizable as a bluish substance in Azan-stained sections, is present within follicular lumina (Figure 5F). At TS13 the follicles have further increased in size and number. The cells of the follicular epithelium appear slightly flattened compared to the previous stage. No colloid is detectable in the investigated specimen (Figure 5G). At TS15 the thyroid glands have continued to elongate and are almost spindle-shaped (Figure 5C). The size and number of follicles have further increased and the follicular lumina are filled with colloid in the investigated specimen. Erythrocytes are recognizable around and in between follicles indicating a continuous vascularization of the thyroid gland (Figure 5H).

We therefore measured the cell height of the follicular epithelium as well as the number and diameter of follicles (Figure 6B-D). All measurements were taken from sections from the central region of the thyroid gland (Cruz & Fabrezi, 2020). In *A. wahlbergi* we observed a huge increase in the follicle cell height from TS8 to TS10/11, followed by a gradual decrease in TS 13 and TS15. In the two examined stages of *A. xenodactyloides* cells were smaller compared to *A.*

wahlbergii. However, a similar but lower decrease in cell height from TS11 to TS15 is also

recognizable (Figure 6B). Follicle numbers and follicle number are more or less similar in *A.*

wahlbergii and *A. xenodactyloides* (Figure 6C, D). In *A. wahlbergi*, follicle number and diameter

decrease from TS8 to TS10/11 and then increases gradually from TS10/11 to TS15. In the two

specimens of *A. xenodactyloides*, an increase in follicle number and diameter can also be

observed.

In summary, our data indicate that thyroid glands differentiate histologically between TS8 and

TS10/11, become highly active at TS11 (follicle cell height, first detected colloid) and develop

and adult-like morphology between TS 13 and TS15 (increasing follicle diameter and number,

decreasing cell height).

Skin and thyroid gland histology in a *Cardioglossa* tadpole

The skin of *Cardioglossa* sp. at Gosner-stage 27 (Gosner, 1960) exhibits a pattern typical for the

majority of tadpoles (McDiarmid & Altig, 1999). The epidermis is two-layered. The basal layer

consists of cuboid cells with spherical nuclei while the apical layer is slightly flattened in the

head but not in the trunk region (Figure 7A1-A5). A dark-blue stained extracellular layer

resembling the keratinized apical cell layer present in late *Arthroleptis* embryos can be found.

Some unicellular gland cells are present in between basal and apical cells (Figure 7A1, A3). A

thick stratum compactum is present beneath the basal epidermal layer. Melanophores are

associated with the stratum compactum. Their density decreases dorso-ventrad in the head as well

as the trunk region (Figure 7A1-A5).

The thyroid glands exhibit a more elongated olive-shape and are localized in the same position as

described for *Arthroleptis* embryos (Figure 7B). At this developmental stage, well developed

follicle outlines by slightly flattened epithelial cells can be recognized. No colloid or

vascularization has been detected in the examined specimen. The flattened follicular epithelial

cells and the absence of colloid and vascularization indicate a low activity of TH-production.

Discussion

***Arthroleptis* exhibits accelerated body wall fusion and a phase of metamorphic skin**

pigmentation repatterning that correlates with increased thyroid gland activity

The skin of frogs harbors a variety of chromatophores (Duellman & Trueb, 1994). In this study

however we focus on the pattern of melanophores only. Different types of melanophores are

present in the adult frog skin (Yasutomi, 1987). This adult pigmentation pattern is established via

metamorphic changes in the dermal pigmentation, chromophore morphology, and biochemistry

of the tadpole skin (Yasutomi, 1987). This indicates a correlation of TH production and

metamorphic skin pigmentation repatterning in biphasic anurans (Smith-Gill & Carver, 1981).

In direct developing species, pigmentation patterning has only been described in *E. coqui*

(Townsend and Stewart, 1985). At TS7, first melanophores appear in the trunk region and

pigmentation increases slowly until it becomes heavy at TS10. Pigmentation of the head is

delayed becoming heavy at TS12. This general process of melanophore patterning is also seen in

Arthroleptis (Schweiger et al., 2017; this study) and we here provide the first detailed data on the

morphology of different melanophore types during skin development in a direct developing frog

species. In *Arthroleptis*, the first type of melanophore is spindle-shaped with long thin extensions.

This type is typical of embryonic melanophores in biphasic species (Smith-Gill & Carver, 1981;

Yasutomi, 1987). A second type of melanophore with a smaller, more circular-shape with a few

short or no extensions appears at around TS11. This melanophore type resembles melanophores

appearing during TH-mediated metamorphosis in biphasic species (Smith-Gill & Carver, 1981;

Yasutomi, 1987). It is generally accepted that follicular cell height, follicle diameter and number

(see Figure 6A) roughly reflect thyroid gland activity and TH production (Coleman, Evennett, &

Dodd, 1968; Grim et al., 2009). Therefore, the appearance of this second melanophore type

correlates with an increased thyroid activity. The timing of pigmentation repatterning and

melanophore type appearance in *Arthroleptis* is very similar to the TH-mediated metamorphic

transitions seen in the skin of biphasic species (Smith-Gill & Carver, 1981; Yasutomi, 1987).

Experimental studies on the pigmentation pattern (Elinson & Fang, 1998) and histological

investigation of thyroid gland activity (Jennings & Hanken, 1998) in *E. coqui* led to the

suggestion that TH is also involved in the closure of the pigmented body surface (Callery &

Elinson, 2000; Elinson & Fang, 1998). While the overall melanophore patterning and subsequent

darkening of the skin are similar in *Arthroleptis* and *E. coqui,* there is a difference in the timing

of the ventral fusion of the pigmented lateral body walls

This fusion appears much earlier in *Arthroleptis* (TS9, Figure 2) compared to *E. coqui* (TS12,

Elinson & Fang, 1998). In *E. coqui* this event correlates with a peak in thyroid gland activity

(Laslo et al., 2019) and has experimentally shown to be TH dependent (Callery & Elinson, 2000).

In *Arthroleptis*, histologically inferred thyroid activity peaks at TS10/11. Therefore fusion of the

lateral body walls seems slightly accelerated. This could be due to a higher TH-sensitivity of the

lateral body wall tissue or the earlier expression of TH that cannot be detected histologically.

Experimental and molecular/immuno-histochemical or mass spectroscopic data from *Arthroleptis*

are needed to clarify this.

Skin histology in *Arthroleptis* exhibits a phase of metamorphic repatterning that correlates

with increased thyroid gland activity

Skin and thyroid gland histology of the *Cardioglossa* tadpole are similar to many other tadpoles

at a comparable developmental stage (Cruz & Fabrezi, 2020). Alterations in skin development in

Arthroleptis therefore seem to be correlated with the evolution of direct development. However,

an apical extracellular layer resembling the stratum corneum in adult frog skin is present in the *C.*

manengouba tadpole. Tadpoles of various species of *Cardioglossa*, including *C. manengouba*,

have been frequently found buried in sediment and other substrate in streams (Blackburn, 2008;

Hirschfeld, Barej, Gonwouo, & Rödel, 2012) and this putative stratum corneum could be an

adaptation to their fossorial life style.

As reported for *E. coqui* (Fang & Elinson, 1996; Schlosser et al., 1999), major tadpole characters

such as cement glands, neuromasts and skein cells (Tamakoshi et al., 1998) were not observed in

Arthroleptis. Data for *I. henselii* are not available (Goldberg et al., 2020). In *Arthroleptis*, skin

development can be divided into four major phases (embryonic, "tadpole", metamorphic and

adult; Figure 8) that are also typical for many biphasic species (Fox, 1986; Robinson &

Heintzelman, 1987). At TS4 embryonic ectoderm is a single-cell layered epithelium not yet

differentiated into an epidermis. At around TS7 the ectodermal cells have differentiated into a

tadpole-like epidermis (two-layered, unicellular glands, inner melanophores). However, the

epidermis still exhibits some embryonic features (intracellular yolk granules, low melanophore

density, ciliated cells and the lack of a stratum compactum; Figure 8). At TS8, the skin exhibits

more mature tadpole-like features such as fewer intracellular yolk granules and ciliated cells, a

higher melanophore density, many unicellular glands and a stratum compactum. At TS11,

degeneration of the apical epidermal layer and unicellular glands together with the appearance of

multicellular gland progenitors and an additional outer melanophore layer (Figure 8) resembles

skin repatterning during the metamorphic climax of biphasic species (Fox, 1985; Gaupp, 1904;

Verma, 1965). In biphasic species, shortly before metamorphic climax, apical cells in the tadpole

epidermis undergo apoptosis except for the single layer of basal cells. The remaining basal cells

proliferate extensively during metamorphic climax building up the adult epidermis and gland

cells (Kinoshita & Sasaki, 1994; Schreiber & Brown, 2003; Yoshizato, 1992). In *Arthroleptis*,

metamorphic skin repatterning is seen at TS11, when the "tadpole" apical epidermal layer

degenerates and cells in the remaining basal layer start to proliferate (Figure 8; strong PCNA

signal). This patterning of the tadpole-like skin and increased PCNA activity correlate with an

increase of histologically inferred thyroid gland activity at TS10/11 (see Figure 6). From TS13

on, the skin of *Arthroleptis* exhibits major features of an adult frog skin, such as a multilayered

(three to five layers dorsally) epidermis with an apical stratum corneum, large multicellular

glands embedded into a stratum spongiosum and a layer of outer and inner melanophores

bordering the thick stratum compactum (Figure 8). This in term correlates with a decreased

thyroid gland activity and the absence of a PCNA signal in epidermal cells.

Regarding other direct developing species, detailed histological data are only available for two

developmental stages of *Ischnocnema henselii* (Goldberg et al., 2020). At TS6, the epidermis of

this species is a single-layered epithelium overlaying an undifferentiated mesenchymal dermis

(Goldberg et al., 2020). This is very similar to *Arthroleptis* embryos except that the epidermis is

two-layered in most body regions at this developmental stage. In *I. henselii* embryos at TS14,

epidermal cells start to proliferate forming a two-layered epidermis. Ciliated cells are still present

in many body areas, progenitors of multicellular glands are detectable and melanocyte density

has increased compared to TS6. The dermis has also differentiated and a stratum compactum is

present (Goldberg et al., 2020). This is different to *Arthroleptis,* where the features described for

I. henselii at TS14 are already present at TS11 (Figure 8). In contrast to *I. henselii*, the short

description of a stage likely corresponding to TS9 in *Eleutherodactylus* that exhibits a two-

layered epidermis (Adamson, Harrison, & Bayley, 1960), implies an earlier pattern of skin

differentiation more similar to *Arthroleptis*. This is very interesting since a differentiated thyroid

gland is already present at TS6 in *I. henselii*. This is much earlier compared to *Arthroleptis* (TS8;

this study) and *E. coqui* (TS10; Jennings & Hanken, 1998) and skin development does therefore

not seem to correlate with thyroid gland maturation in *I. henselii*.

This mismatch between very early TH gland maturation and very late skin remodeling in *I.*

henselli could be indicative of a loss of TH-dependency of skin remodeling and hence a loss of

constraint. Alternatively, skin maturation might still be dependent on TH-signaling, but TH-

production of the thyroid gland could be very low and a concentration necessary to induce skin

repatterning is reached very late. Another possibility is that TH-levels are normal but the

sensitivity of the skin tissue is decreased, requiring higher TH concentrations to induce

repatterning. Both of these TH-dependent scenarios would explain the late onset (i.e.

heterochronic shift) of skin remodeling.

Thyroid gland development and activity in direct developing frogs

As it has been shown that metamorphic changes in anurans, and amphibians in general, are under a complex hormonal control but TH seems to be a major regulator (Shi, 1999). There are two hypotheses how the ancestral thyroid axis has influenced the evolution of direct development (see Jennings & Hanken, 1998 and references therein). (1) Thyroid gland activity and TH production are activated early in development leading to a precocious formation of adult features. Direct development is constrained by the ancestral dependency on TH for adult development. Consequently, direct development evolved by a heterochronic shift (Gould, 1977) of ancestral developmental mechanisms rather than the loss of constraints from it ("heterochronic shift" hypothesis). (2) Metamorphosing tissues lost their ancestral TH-dependency. In this case, the loss of constraints would have been more important in the evolution of direct development than heterochronic shifts of ancestral mechanisms ("loss of constraint" hypothesis).

Experimental studies on *E. coqui* have shown that this species exhibits a mosaic pattern of heterochronic shifts and loss of constraints for different morphological features (Callery & Elinson, 2000; Hanken et al., 1997; Hughes, 1966; Hughes & Reier, 1972; Jennings & Hanken, 1998; Lynn, 1948; Lynn & Peadon, 1955; Townsend & Stewart, 1985).

There are many similarities between *E. coqui* and *Arthroleptis* regarding metamorphic repatterning processes that correlate with increased thyroid activity such as tail regression, remodeling of the "larval" into the adult hyobranchial apparatus and cranial muscle repatterning, (Hanken et al., 1997; Jennings & Hanken, 1998; Schweiger et al., 2017; Schweiger, Naumann, & Müller, in prep.; Townsend & Stewart, 1985). Most of these processes start slightly earlier in *Arthroleptis*, which correlates with an earlier appearance of the thyroid glands compared to *E. coqui* (Hanken & Jennings, 1998; this study). *Ischnocnema henselii* might be another example

where some features of skin development might have lost their dependency on TH, but the

interpretation of the data for *I. henselii* is hampered by the lack of a detailed investigation of

thyroid gland and skin development over several developmental stages.

However, our data indicate that many tadpole specific structures appear, differentiate and are

repatterned to the adult configuration during a short metamorphic phase although this cryptic

metamorphosis appears to be not as climactic as in biphasic frogs.

Developmental constraints, heterochrony and the parallel evolution of direct development

Direct development has evolved parallel in different groups of frogs. The comparison of

embryonic development of these direct developing frogs offers the possibility to identify how

developmental constraints may have influenced the observed pattern of parallel evolution. Many

developmental mechanisms are governed by major regulators such as TH. In case of a

heterochronic shift of this regulator (e.g. embryonic thyroid activity in direct developing frogs),

traits that are under the control of this regulator have to follow this shift. These traits are

constrained by this major regulator. Other traits that do not follow the heterochronic shift of this

regulator develop normally and are not constrained. This may lead to a decoupling of ancestrally

synchronous processes such as, e.g., growth and differentiation of the retinotectal system in *E.*

coqui (Schlosser, 2008).

Further investigations of the developmental timing and TH-dependency of anatomical features of

Eleutherodactylus, *Arthroleptis*, *Ischnocnema* and other direct developing frogs are needed.

These investigations will clarify if homologous features are generally under the control of TH

and therefore constrained by thyroid gland activity. Studies on direct development in frogs, or

amphibians in general, and various other cases such as the classic example of the evolution of

similar morphotypes in African cichlids (Elmer et al., 2014; Meier, Marques, Wagner, Excoffier,

& Seehausen, 2018) or the recently "re-discovered" dispersion/re-aggregation and diapause

phases in early killifish development (Furness, Reznick, Springer, & Meredith, 2015; Naumann

& Englert, 2018) will help to better understand how developmental processes constrain the

evolution of adaptive traits and to further clarify mechanisms and identifying major regulators

underlying parallel evolution.

Experimental Procedures

Specimens

Embryos of *Arthroleptis wahlbergii* Smith, 1849, collected in South Africa, and *A.*

xenodactyloides Hewitt, 1933, collected in Tanzania, were euthanized using 0.1 % tricaine

methanesulfonate (MS222, Fluka), fixed in 4% phosphate buffered formalin and stored in 70%

ethanol (Schweiger et al., 2017). Staging of embryos follows Townsend and Stewart (1985); TS

stages hereafter. One tadpole of *Cardioglossa manengouba* Blackburn, 2008 was available for

investigation. For a complete list of specimen see Supplementary material 1.

Histological sectioning and staining

For sectioning, embryos were dehydrated in an ethanol series (70%, 90%, 2 x 96%; 5 min each),

sectioned and subsequently rehydrated (2 x 96%, 90%, 70% ethanol, distilled water; 5 min each)

prior to staining. Embryos up to TS8 were embedded in Technovit 8100 (Kulzer, R0010022),

sectioned at 3 µm and stained with a mixture of basophilic Methylene blue and acidic Fuchsin

(Mulisch & Welsch, 2010). Embryos from TS10/11 on were decalcified for up to three days

(Osteomoll, Merck), embedded in Paraplast (Roth, X881.1), sectioned at 8 µm and stained with

Heidenhain's Azan (Mulisch & Welsch, 2010). All sections were made using a Microm HM 360

(Zeiss) and photographs were taken using an Olympus Dotslide BX51 microscope.

Fluorescent antibody staining

Paraffin sections were rehydrated as described before. For antigen retrieval, slides were

transferred to a plastic cuvette with citrate buffer (recipe see Schlosser, 2008) and heated in a

microwave (Bosch HMT 72M 420) for 10 minutes at 180W. Subsequently, slides were placed at

room temperature (RT, 21-25°C) to cool down for at least 5 minutes and then rinsed in distilled

water for 1 minute. Afterwards, slides were rinsed with PBS (3 x 5 minutes), placed into a wet

chamber and blocked with antibody diluent (DAKO) for 1 hour at RT. Primary antibodies against

proliferating cell nuclear antigen (PCNA-1, 1/200, sc-7907, Santa Cruz; PCNA-2, 1/200,

M087901, Dako) were applied overnight at 4°C. On the next day, slides were rinsed in PBS (3 x

5 minutes), blocked and incubated with secondary antibodies (Alexa-488 anti-mouse, #A28175,

Thermo Fisher Scientific and Alexa-568 anti-rabbit, #A-11011, Thermo Fisher Scientific) for 1

hour at RT. Afterwards, slides were rinsed with PBS (3 x 5 minutes), quickly washed with

distilled water and cover-slipped with Fluoroshield with DAPI (Sigma, F6057). Slides were

photographed using a Zeiss Axioplan Microscope equipped with a Spot camera and the Zeiss

Axioplan software.

Micro-CT scanning and 3D reconstruction

Specimens were contrasted using a solution of 1% polymolybdenic acid in 70% ethanol

(Metscher, 2009). *Arthroleptis* specimens were CT-scanned using a Nanotom S μCT scanner

(Phoenix X-Ray) and the *Cardioglossa* tadpole was studied using synchrotron radiation based x-

ray micro-CT. Imaging was performed at the Imaging Beamline P05 (IBL) (Greving et al., 2014;

Haibel et al., 2010; Wilde et al., 2016) operated by the Helmholtz-Zentrum-Geesthacht at the

storage ring PETRA III (Deutsches Elektronen Synchrotron – DESY, Hamburg, Germany) using

a photon energy of 24 keV. Projections were recorded using a CCD camera system (MicroLine

ML09000 - Finger Lake Instruments) with an effective pixel size of 2.42 μm. For each

tomographic scan 1801 projections at equal intervals between 0 and π have been recorded.

Tomographic reconstruction has been done by applying a filtered back projection algorithm

(FBP) implemented in a custom reconstruction pipeline (Moosmann et al., 2014) using Matlab

(Math-Works) and the Astra Toolbox (Palenstijn, Batenburg, & Sijbers, 2011; van Aarle et al.,

2016; van Aarle et al., 2015). For the tomographic reconstruction, raw projections were binned

two times resulting in an effective pixel size of the reconstructed volume of 4.83 μm. Three-

dimensional reconstructions and mixed surface-volume renderings of the thyroid gland were

prepared using AMIRA 5.4.2 (FEI Visualization, Science Group, Bordeaux, France).

Follicle cell height and follicle diameter measurements

Qualitative and quantitative descriptions of thyroid glands are based on sections from the middle

region of the paired thyroid lobes according to Cruz and Fabrezi (2020). Measurements of follicle

cell heights and follicle diameter were obtained from digitalized sections using ImageJ

(Schneider, Rasband, & Eliceiri, 2012). A box and whisker plot of follicle cell heights was

prepared using Microsoft Excel 2010. Numbers of follicles were counted and mean values of the

measured follicle diameters were calculated for each specimen investigated.

Photographs and Image processing

Photographs of whole embryos were taken using a Zeiss Discovery V12 stereomicroscope with

an attached Zeiss AxioCam digital camera. Brightness and Contrast of Images was adjusted using

either ImageJ or Adobe Photoshop CS6. Channel colors of fluorescent antibody staining were

assigned in ImageJ. In some images the "CLAHE" filter implemented in ImageJ was applied to

enhance local contrast.

Acknowledgements

We would like to thank Katja Felbel for valuable help in the laboratory. We thank Mark-Oliver

Rödel for critical reading of the manuscript and a photograph of an adult *C. manengouba* and

Mareike Hirschfeld for a photograph of a *C. manengouba* tadpole. We thank Lennart Olsson for

critical reading of the manuscript. We thank Paul Lukas for help with the Dot Slide Microscope.

We would also like to thank Christoph Englert, Birgit Perner and Dagmar Kruspe for the kind gift

of the two PCNA antibodies.

References

Adamson, L., Harrison, R., & Bayley, I. (1960). *The development of the whistling frog*
Eleutherodactylus martinicensis of Barbados. Paper presented at the Proceedings of the
Zoological Society of London.
Altig, R., & McDiarmid, R. (1999). Diversity: familial and generic characterizations.

Blackburn, D. C. (2008). A new species of *Cardioglossa* (Amphibia: Anura: Arthroleptidae) endemic to Mount Manengouba in the Republic of Cameroon, with an analysis of morphological diversity in the genus. *Zoological Journal of the Linnean Society, 154*(3), 611-630.

Brown, D. D., & Cai, L. (2007). Amphibian metamorphosis. *Developmental Biology, 306*(1), 20.

Callery, E. M., & Elinson, R. P. (2000). Thyroid hormone-dependent metamorphosis in a direct developing frog. *Proceedings of the National Academy of Sciences, 97*(6), 2615-2620.

Callery, E. M., Fang, H., & Elinson, R. P. (2001). Frogs without polliwogs: evolution of anuran direct development. *BioEssays, 23*(3), 233-241.

Coleman, R., Evennett, P., & Dodd, J. (1968). Ultrastructural observations on the thyroid gland of *Xenopus laevis* Daudin throughout metamorphosis. *General Comparative Endocrinology, 10*(1), 34-46.

Cruz, J. C., & Fabrezi, M. (2020). Histology and microscopic anatomy of the thyroid gland during the larval development of *Pseudis platensis* (Anura, Hylidae). *Journal of Morphology, 281*(1), 122-134.

Douglas, A. J., Hug, L. A., & Katzenback, B. A. (2020). Composition of the North American wood frog (*Rana sylvatica*) skin microbiome and seasonal variation in community structure. *bioRxiv*.

Duellman, W. E., & Trueb, L. (1994). *Biology of Amphibians*: JHU press.

Elinson, R., & Fang, H. (1998). Secondary coverage of the yolk by the body wall in the direct developing frog, *Eleutherodactylus coqui*: an unusual process for amphibian embryos. *Development Genes and Evolution, 208*(8), 457-466.

Elmer, K. R., Fan, S., Kusche, H., Spreitzer, M. L., Kautt, A. F., Franchini, P., & Meyer, A. (2014). Parallel evolution of Nicaraguan crater lake cichlid fishes via non-parallel routes. *Nature Communications, 5*(1), 1-8.

Fang, H., & Elinson, R. P. (1996). Patterns of Distal-less gene expression and inductive interactions in the head of the direct developing frog *Eleutherodactylus coqui*. *Developmental Biology, 179*(1), 160-172.

Feng, Y.-J., Blackburn, D. C., Liang, D., Hillis, D. M., Wake, D. B., Cannatella, D. C., & Zhang, P. (2017). Phylogenomics reveals rapid, simultaneous diversification of three major clades of Gondwanan frogs at the Cretaceous–Paleogene boundary. *Proceedings of the National Academy of Sciences, 114*(29), E5864-E5870.

Fernandes, T. L., Antoniazzi, M. M., Sasso-Cerri, E., Egami, M. I., Lima, C., Rodrigues, M. T.,

& Jared, C. (2011). Carrying progeny on the back: reproduction in the Brazilian aquatic

frog *Pipa carvalhoi*. *South American Journal of Herpetology, 6*(3), 161-176.

Fox, H. (1985). Changes in amphibian skin during larval development and metamorphosis. In M.

B. M Balls (Ed.), *Metamorphosis*: Clarendon Press, Oxford.

Fox, H. (1986). The skin of amphibia. *Biology of the Integument, Vertebrates, 2*, 78-148.

Furness, A. I., Reznick, D. N., Springer, M. S., & Meredith, R. W. (2015). Convergent evolution

of alternative developmental trajectories associated with diapause in African and South

American killifish. *Proceedings of the Royal Society B: Biological Sciences, 282*(1802),

20142189.

Futuyma, D., & Kirkpatrick, M. (2017). Evolution. Sinauer. *Sunderland, MA*.

Gaupp, E. (1904). *A. Ecker's und R. Wiedersheim's Anatomie des Frosches. Band 3. Lehre von*

den Eingeweiden, dem Integument und den Sinnesorganen: Vieweg.

Goin, O. B., & Goin, C. J. (1962). Amphibian eggs and the montane environment. *Evolution*,

364-371.

Goldberg, J., Candioti, F. V., & Akmentins, M. S. (2012). Direct-developing frogs: ontogeny of

Oreobates barituensis (Anura: Terrarana) and the development of a novel trait. *Amphibia-*

Reptilia, 33(2), 239-250.

Goldberg, J., Taucce, P. P., Quinzio, S. I., Haddad, C. F., & Candioti, F. V. (2020). Increasing

our knowledge on direct-developing frogs: The ontogeny of *Ischnocnema henselii*

(Anura: Brachycephalidae). *Zoologischer Anzeiger, 284*, 78-87.

Gosner, K. L. (1960). A simplified table for staging anuran embryos and larvae with notes on

identification. *Herpetologica, 16*(3), 183-190.

Gould, S. J. (1977). *Ontogeny and phylogeny*: Harvard University Press.

Greving, I., Wilde, F., Ogurreck, M., Herzen, J., Hammel, J. U., Hipp, A., . . . Beckmann, F.

(2014). *P05 imaging beamline at PETRA III: first results*. Paper presented at the

Proceedings of SPIE - Developments in X-Ray Tomography IX, San Diego, California,

United States.

Grim, K. C., Wolfe, M., Braunbeck, T., Iguchi, T., Ohta, Y., Tooi, O., . . . Tietge, J. (2009).

Thyroid histopathology assessments for the amphibian metamorphosis assay to detect

thyroid-active substances. *Toxicologic Pathology, 37*(4), 415-424.

Haibel, A., Ogurreck, M., Beckmann, F., Dose, T., Wilde, F., Herzen, J., . . . Mohr, J. (2010). *Micro- and nano-tomography at the GKSS Imaging Beamline at PETRA III.* Paper presented at the SPIE Optical Engineering + Applications.

Hall, B. K., & Olson, W. M. (2003). *Keywords and Concepts in Evolutionary Developmental Biology*: Harvard University Press.

Hanken, J., Carl, T. F., Richardson, M. K., Olsson, L., Schlosser, G., Osabutey, C. K., & Klymkowsky, M. W. (2001). Limb development in a "nonmodel" vertebrate, the direct□developing frog *Eleutherodactylus coqui. Journal of Experimental Zoology, 291*(4), 375-388.

Hanken, J., Klymkowsky, M. W., Alley, K. E., & Jennings, D. H. (1997). Jaw muscle development as evidence for embryonic repatterning in direct-developing frogs. *Proceedings of the Royal Society B: Biological Sciences*, 1349-1354.

Hanken, J., Klymkowsky, M. W., Summers, C. H., Seufert, D. W., & Ingebrigtsen, N. (1992). Cranial ontogeny in the direct□developing frog, *Eleutherodactylus coqui* (Anura: Leptodactylidae), analyzed using whole□mount immunohistochemistry. *Journal of Morphology, 211*(1), 95-118.

Hanken, J., & Summers, C. H. (1988). Skull development during anuran metamorphosis: III. Role of thyroid hormone in chondrogenesis. *Journal of Experimental Zoology, 246*(2), 156-170.

Heinicke, M. P., Duellman, W. E., Trueb, L., Means, D. B., MacCulloch, R. D., & Hedges, S. B. (2009). A new frog family (Anura: Terrarana) from South America and an expanded direct-developing clade revealed by molecular phylogeny. *Zootaxa, 2211*(1), 1-35.

Hewitt, J. (1933). Descriptions of some new reptiles and a frog from Rhodesia. *Occasional Papers of the National Museum of Southern Rhodesia, 2*, 45-50.

Hirschfeld, M., Barej, M. F., Gonwouo, N. L., & Rödel, M.-O. (2012). Tadpole descriptions of three *Cardioglossa* species from southwestern Cameroon (Amphibia: Anura: Arthroleptidae). *Salamandra, 48*(3), 147-156.

Huang, L., Li, J., Anboukaria, H., Luo, Z., Zhao, M., & Wu, H. (2016). Comparative transcriptome analyses of seven anurans reveal functions and adaptations of amphibian skin. *Scientific reports, 6*(1), 1-11.

Hughes, A. (1966). The thyroid and the development of the nervous system in *Eleutherodactylus martinicensis*: an experimental study. *Development, 16*(3), 401-430.

Hughes, A., & Reier, P. (1972). A preliminary study on the effects of bovine prolactin on embryos of *Eleutherodactylus ricordii*. *General and Comparative Endocrinology, 19*(2), 304-312.

Jennings, D. H., & Hanken, J. (1998). Mechanistic basis of life history evolution in anuran amphibians: thyroid gland development in the direct-developing frog, *Eleutherodactylus coqui*. *General Comparative Endocrinology, 111*(2), 225-232.

Katz, U. (1986). The role of amphibian epidermis in osmoregulation and its adaptive response to changing environment. In *Biology of the Integument* (pp. 472-498): Springer.

Kerney, R., Gross, J. B., & Hanken, J. (2010). Early cranial patterning in the direct￿developing frog *Eleutherodactylus coqui* revealed through gene expression. *Evolution & Development, 12*(4), 373-382.

Kerney, R., Meegaskumbura, M., Manamendra￿Arachchi, K., & Hanken, J. (2007). Cranial ontogeny in *Philautus silus* (Anura: Ranidae: Rhacophorinae) reveals few similarities with other direct￿developing anurans. *Journal of Morphology, 268*(8), 715-725.

Kinoshita, T., & Sasaki, F. (1994). Body-specific proliferation of adult precursor cells in *Xenopus* larval epidermis. *Histochemistry, 101*(6), 397-404.

Kulkarni, S. S., Singamsetty, S., & Buchholz, D. R. (2010). Corticotropin-releasing factor regulates the development in the direct developing frog, *Eleutherodactylus coqui*. *General and Comparative Endocrinology, 169*(3), 225-230.

Laslo, M., Denver, R. J., & Hanken, J. (2019). Evolutionary conservation of thyroid hormone receptor and deiodinase expression dynamics in ovo in a direct-developing frog, *Eleutherodactylus coqui*. *Frontiers in Endocrinology, 10*, 307.

Liedtke, H. C., Müller, H., Hafner, J., Penner, J., Gower, D. J., Mazuch, T., . . . Loader, S. P. (2017). Terrestrial reproduction as an adaptation to steep terrain in African toads. *Proceedings of the Royal Society B: Biological Sciences, 284*(1851), 20162598.

Lynn, W. G. (1948). The effects of thiourea and phenylthiourea upon the development of *Eleutherodactylus ricordii*. *The Biological Bulletin, 94*(1), 1-15.

Lynn, W. G., & Peadon, A. M. (1955). The role of the thyroid gland in direct development in the anuran, *Eleutherodactylus martinicensis*. *Growth, 19*(4), 263.

McDiarmid, R., & Altig, R. (1999). *Tadpoles: the Biology of Anuran Larvae*: University of Chicago Press.

Meier, J. I., Marques, D. A., Wagner, C. E., Excoffier, L., & Seehausen, O. (2018). Genomics of
parallel ecological speciation in Lake Victoria cichlids. *Molecular Biology and Evolution,*
35(6), 1489-1506.

Metscher, B. D. (2009). MicroCT for comparative morphology: simple staining methods allow
high-contrast 3D imaging of diverse non-mineralized animal tissues. *BMC Physiology,*
9(1), 11.

Moosmann, J., Ershov, A., Weinhardt, V., Baumbach, T., Prasad, M. S., LaBonne, C., . . .
Hofmann, R. (2014). Time-lapse X-ray phase-contrast microtomography for in vivo
imaging and analysis of morphogenesis. *Nature Protocols, 9*(2), 294-304.
doi:10.1038/nprot.2014.033

Mulisch, M., & Welsch, U. (2010). *Romeis Mikroskopische Technik*: Spektrum Akademischer
Verlag.

Müller, H., Liedtke, H. C., Menegon, M., Beck, J., Ballesteros-Mejia, L., Nagel, P., & Loader, S.
P. (2013). Forests as promoters of terrestrial life-history strategies in East African
amphibians. *Biology Letters, 9*(3), 20121146.

Naumann, B., & Englert, C. (2018). Dispersion/reaggregation in early development of annual
killifishes: Phylogenetic distribution and evolutionary significance of a unique feature.
Developmental Biology, 442(1), 69-79.

Olsson, L., Moury, D. J., Carl, T. F., Håstad, O., & Hanken, J. (2002). Cranial neural crest-cell
migration in the direct-developing frog, *Eleutherodactylus coqui*: molecular heterogeneity
within and among migratory streams. *Zoology, 105*(1), 3-13.

Palenstijn, W. J., Batenburg, K. J., & Sijbers, J. (2011). Performance improvements for iterative
electron tomography reconstruction using graphics processing units (GPUs). *Journal of*
Structural Biology, 176(2), 250-253. doi:10.1016/j.jsb.2011.07.017

Quinzio, S. I., & Goldberg, J. (2019). Transient integumentary structures in *Boana riojana*
(Anura, Hylidae) tadpoles. *Amphibia-Reptilia, 40*(4), 543-549.

Robinson, D. H., & Heintzelman, M. B. (1987). Morphology of ventral epidermis of *Rana*
catesbeiana during metamorphosis. *The Anatomical Record, 217*(3), 305-317.

San Mauro, D., Gower, D. J., Müller, H., Loader, S. P., Zardoya, R., Nussbaum, R. A., &
Wilkinson, M. (2014). Life-history evolution and mitogenomic phylogeny of caecilian
amphibians. *Molecular Phylogenetics and Evolution, 73*, 177-189.

Schlosser, G. (2008). Development of the retinotectal system in the direct-developing frog

Eleutherodactylus coqui in comparison with other anurans. *Frontiers in Zoology, 5*(1), 9.

Schlosser, G., Kintner, C., & Northcutt, R. G. (1999). Loss of ectodermal competence for lateral

line placode formation in the direct developing frog *Eleutherodactylus coqui*.

Developmental biology, 213(2), 354-369.

Schlosser, G., & Roth, G. (1997). Evolution of nerve development in frogs; pp. 94–112. *Brain,*

Behavior and Evolution, 50(2), 94-112.

Schluter, D., Clifford, E. A., Nemethy, M., & McKinnon, J. S. (2004). Parallel evolution and

inheritance of quantitative traits. *The American Naturalist, 163*(6), 809-822.

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of

image analysis. *Nature Methods, 9*(7), 671-675.

Schreiber, A. M., & Brown, D. D. (2003). Tadpole skin dies autonomously in response to thyroid

hormone at metamorphosis. *Proceedings of the National Academy of Sciences, 100*(4),

1769-1774.

Schweiger, S., Naumann, B., Larson, J. G., Möckel, L., & Müller, H. (2017). Direct development

in African squeaker frogs (Anura: Arthroleptidae: *Arthroleptis*) reveals a mosaic of

derived and plesiomorphic characters. *Organisms Diversity & Evolution, 17*(3), 693-707.

Schweiger, S., Naumann, B., & Müller, H. (in prep.). The ghost of the tadpole - Embryonic

development of the cranial musuloskeletal system in African squeaker frogs (*Arthroleptis*)

reveals heterochronic shifts and parallel evolution of differential metamorphosis in direct

developing frogs.

Shi, Y.-B. (1999). *Amphibian metamorphosis: from morphology to molecular biology*: Wiley-

Liss.

Simpson, G. G. (1961). Principles of animal taxonomy.

Smith-Gill, S. J., & Carver, V. (1981). Biochemical characterization of organ differentiation and

maturation. In *Metamorphosis* (pp. 491-544): Springer.

Smith, A. (1849). *Illustrations of the Zoology of South Africa: Consisting Chiefly of Figures and*

Descriptions of the Objects of Natural History Collected During an Expedition Into the

Interior of South Africa, in the Years 1834, 1835, and 1836; Fitted Out by" the Cape of

Good Hope Association for Exploring Central Africa" (Vol. 4).

Tamakoshi, T., Oofusa, K., & Yoshizato, K. (1998). Visualization of the initiation and sequential
expansion of the metamorphic conversion of anuran larval skin into the precursor of adult
type. *Development, Growth & Differentiation, 40*(1), 105-112.

Tata, J. R. (2006). Amphibian metamorphosis as a model for the developmental actions of
thyroid hormone. *Molecular and Cellular Endocrinology, 246*(1-2), 10-20.

ten Brink, H. t., Onstein, R. E., & de Roos, A. M. (2020). Habitat deterioration promotes the
evolution of direct development in metamorphosing species. *Evolution*.

Thibaudeau, G., & Altig, R. (1999). Endotrophic anurans: development and evolution. *Tadpoles:*
the Biology of Anuran Larvae, 170-188.

Thomas, R. (1966). New species of antillean *Eleutherodactylus*. *Quarterly Journal of the Florida*
Academy of Sciences, 28(4), 375-391.

Townsend, D. S., & Stewart, M. M. (1985). Direct development in *Eleutherodactylus coqui*
(Anura: Leptodactylidae): a staging table. *Copeia*, 423-436.

van Aarle, W., Palenstijn, W. J., Cant, J., Janssens, E., Bleichrodt, F., Dabravolski, A., . . .
Sijbers, J. (2016). Fast and flexible X-ray tomography using the ASTRA toolbox. *Optics*
Express, 24(22), 25129-25147. doi:Doi 10.1364/Oe.24.025129

van Aarle, W., Palenstijn, W. J., De Beenhouwer, J., Altantzis, T., Bals, S., Batenburg, K. J., &
Sijbers, J. (2015). The ASTRA Toolbox: A platform for advanced algorithm development
in electron tomography. *Ultramicroscopy, 157*, 35-47. doi:10.1016/j.ultramic.2015.05.002

Varga, J. F., Bui-Marinos, M. P., & Katzenback, B. A. (2019). Frog skin innate immune
defences: sensing and surviving pathogens. *Frontiers in Immunology, 9*, 3128.

Verma, K. (1965). Regional differences in skin gland differentiation in *Rana pipiens*. *Journal of*
Morphology, 117(1), 73-85.

Wake, D., & Hanken, J. (1996). Direct development in the lungless salamanders: what are the
consequences for developmental biology, evolution and phylogenesis? *International*
Journal of Developmental Biology, 40(4), 859-869.

Wilde, F., Ogurreck, M., Greving, I., Hammel, J. U., Beckmann, F., Hipp, A., . . . Schreyer, A.
(2016). Micro-CT at the imaging beamline P05 at PETRA III. *AIP Conference*
Proceedings, 1741(1), 030035. doi:10.1063/1.4952858

Yasutomi, M. (1987). Migration of epidermal melanophores to the dermis through the basement
membrane during metamorphosis in the frog, *Rana japonica. Pigment cell research, 1*(3),
181-187.

Yoshizato, K. (1992). Death and transformation of larval cells during metamorphosis of anura.

Development, growth and differentiation, 34(6), 607-612.

Ziermann, J. M., & Diogo, R. (2014). Cranial muscle development in frogs with different

developmental modes: direct development versus biphasic development. *Journal of*

morphology, 275(4), 398-413.

Figure legends

Figure 1. A, a family-level phylogeny of recent frogs based on Feng et al., 2017 (Feng et al., 2017). Blue boxes indicate the presence of an ancestral biphasic live history while orange boxes indicate direct development. B, a schematic graph showing the gradual morphological change during ontogeny of the direct developing *Arthroleptis* (*A. wahlbergii*; 1, embryo; 2, adult frog) and the more climactic metamorphic change in its biphasic sister genus *Cardioglossa* (*C. manengouba*; 1, tadpole; 2, adult frog). Photographs are not to scale. C, schematic diagrams of the organization of the tadpole and adult frog skin.

Figure 2. A-G, Photographs of the skin pigmentation of different developmental stages of *Arthroleptis wahlbergii*. The schematic embryo on the left illustrates the photographed regions (1, 2 and 3). Blue arrows indicate the position of the eye, red asterisks the position of the forelimb. The dotted grey lines and percentage number indicate the distance between the back (dorsal, 0%) and ventral midline (ventral, 100%). The black arrow in A2-D2 indicates the migration distance of the first melanophore type (shown in B`). The white arrow in H2-J2 indicates the migration distance of the second melanophore type (shown in G`). The black dotted line in C3-G3 indicates the ventral midline. Scale bar is 200 μm.

Figure 3. Histological cross sections of the skin of *Arthroleptis wahlbergii* at different developmental stages. The schematic embryo on top illustrates the photographed regions (1-5). Grey arrows indicate the epidermal cilia. Yellow arrows indicate intracellular yolk droplets.

Black arrows indicate epidermal nuclei. The yellow arrow in H3 indicates the glandular duct. The

number of epidermal cellular layers is indicated by a grey line and asterisks. Scale bar is 20 µm.

cc, ciliated cell; co, stratum corneum; ec, epithel cell; etm, epaxonic trunk muscles; g,

multicellular gland progenitor; g1, unicellular gland; g2, multicellular granular gland; g3,

multicellular mucus gland; htm, hypaxonic trunk muscles; m, mesenchyme; me, melanophore; sc,

stratum compactum; sp, stratum spongiosum; vc, vacuole.

Figure 4. Fluorescent PCNA antibody staining in cross sections through the skin of *Arthroleptis*

wahlbergii embryos at different developmental stages. Two different antibodies against PCNA

have been used. PCNA-1 is colored in yellow and shown in A, C, E and G. PCNA-2 is colored in

magenta and shown in B, D, F and H. Nuclei are stained with DAPI and colored in grey. A` and

B` Show a close up of the PCNA signal within the nucleus. Magenta arrows in B indicate the

scattered antibody signal. A-H, scale bar is 20 µm. A` and B`, scale bar is 10 µm. g2,

multicellular granular gland; g3, multicellular mucus gland.

Figure 5. A-C volume renderings of µCT scans of different developmental stages of *Arthroleptis*

xenodactyloides. The thyroid glands are colored in lilac. Renderings are not to scale. D-H,

histological cross-sections of the ventral head region of *A. wahlbergii* embryos at different

developmental stages. Blue arrows indicate absence of colloid within follicles. Blue asterisks

indicate the presence of colloid within follicles. Scale bar is 25 µm. hp, hyoid plate; th, thyroid

gland.

Figure 6. A, schematic illustration of thyroid gland follicels and the measured parameters. B, Box

plot of the follicle cell height in *Arthroleptis* at different developmental stages. C, diagram of the

follicle number in *Arthroleptis* at different developmental stages. D, diagram of the average

follicle diameter in *Arthroleptis* at different developmental stages. Aw, *A. wahlbergii*; Ax, *A.*

xenodactyloides.

Figure 7. A, histological cross sections of the skin of a *Cardioglossa manengouba* tadpole at

Gosner-stage 27. The schematic tadpole on top illustrates the photographed regions (1-5). Black

arrows indicate epidermal nuclei. The number of epidermal cellular layers is indicated by a grey

line and asterisks. B, volume renderings of a synchrotron scan of a *C. manengouba* tadpole at

Gosner-stage 27. The thyroid glands are colored in lilac. Renderings are not to scale. C,

histological cross sections of the ventral head region of the same tadpole as in A. Lilac arrows

indicate absence of colloid within follicles. Scale bar is 20 µm. bh, basihyal; ch, ceratohyal; co,

keratinized layer; ec, epithel cell; g1, unicellular gland; htm, hypaxonic trunk muscles; m,

mesenchyme; me, melanophore; mu, muscle; sc, stratum compactum; th, thyroid gland.

Figure 8. Schematic summary of the skin composition of *Arthroleptis* at different developmental

stages. Presumptive thyroid gland activity inferred from histological measurements is indicated in

lilac.

Supplementary material 1. Specimen list. ICH, immune-histochemistry; MB-aF, Methylene blue

acidic Fuchsin; PMC, polymolybdenic acid; µCT, micro-computed tomography.

Supplementary material 2. Adult skin histology. Histological cross sections of the skin of an adult (1.3 cm) *Arthroleptis wahlbergi*. A, dorsal head region. B, lateral head region. The orange arrow indicates the glandular duct. C, ventral head region. The number of epidermal cellular layers is indicated by a grey line and asterisks. Scale bar is 20 µm. co, stratum corneum; ec, epithel cell; g2, multicellular granular gland; g3, multicellular mucus gland; me, melanophore; sc, stratum compactum; sp, stratum spongiosum.

Figure 1

Figure 2

Figure 3

	head		trunk		
	dorsal	ventral	dorsal	lateral	ventral
TS 4	A1	A2	A3	A4	A5
TS 5	B1	B2	B3	B4	B5
TS 6	C1	C2	C3	C4	C5
TS 7	D1	D2	D3	D4	D5
TS 8	E1	E2	E3	E4	E5
TS 11	F1	F2	F3	F4	F5
TS 13	G1	G2	G3	G4	G5
TS 15	H1	H2	H3	H4	H5

Figure 4

Figure 5

TS8	TS11	TS15

A frontolateral   B frontolateral   C frontolateral

ventral   lateral

TS8	TS10/11	TS11	TS13	TS15

D   hp   th

E   hp   th

F   hp   th   *

G   th

H   hp   th *

Figure 6

Figure 7

Figure 8

Supplementary material 1

ID	TS stage	species	technique	staining
#01	3	*Arthroleptis wahlbergii*	photo	-
#02	4	*A. wahlbergii*	histology/photo	MB-aF/ -
#03	5	*A. wahlbergii*	histology/photo	MB-aF/ -
#04	5/6	*A. wahlbergii*	photo	-
#05	5/6	*A. wahlbergii*	photo	-
#06	6	*A. wahlbergii*	histology/photo	MB-aF/ -
#07	6/7	*A. wahlbergii*	photo	-
#08	7	*A. xenodactyloides*	histology	MB-aF
#09	7	*A. wahlbergii*	photo	-
#10	8	*A. xenodactyloides*	µCT	1% PMA
#11	8	*A. wahlbergii*	histology/ICH	MB-aF/anti-PCNA
#12	8	*A. wahlbergii*	photo	-
#13	8	*A. wahlbergii*	photo	-
#14	9	*A. xenodactyloides*	histology	MB-aF
#15	9	*A. wahlbergii*	photo	-
#16	10	*A. wahlbergii*	photo	-
#17	10	*A. wahlbergii*	photo	-
#18	11	*A. xenodactyloides*	µCT	1% PMA
#19	11	*A. xenodactyloides*	histology	Azan
#20	11	*A. wahlbergii*	histology/ICH	Azan/anti-PCNA
#21	11	*A. wahlbergii*	photo	-
#22	11	*A. wahlbergii*	photo	-
#23	11/12	*A. wahlbergii*	photo	-
#24	12	*A. wahlbergii*	photo	-
#25	12	*A. wahlbergii*	photo	-
#26	12/13	*A. wahlbergii*	photo	-
#27	13	*A. xenodactyloides*	histology	Azan
#28	13	*A. wahlbergii*	histology/ICH	Azan/anti-PCNA
#29	13	*A. wahlbergii*	photo	-
#30	13	*A. wahlbergii*	photo	-
#31	14	*A. wahlbergii*	photo	-
#32	14/15	*A. wahlbergii*	photo	-
#33	15	*A. xenodactyloides*	µCT	1% PMA
#34	15	*A. xenodactyloides*	histology	Azan
#35	15	*A. wahlbergii*	histology/ICH	Azan/anti-PCNA
#36	15	*A. wahlbergii*	photo	-
#37	15	*A. wahlbergii*	photo	-
#38	15	*A. wahlbergii*	photo	-
#39	adult, 1.3 cm	*A. wahlbergii*	histology	Azan
#40	adult, 2 cm	*A. wahlbergii*	histology	Azan
#41	tadpole	*Cardioglossa* sp.	histology	Azan

Supplementary material 2