Brain volumetric changes in the general population following the COVID-19 outbreak and lockdown

Tom Salomon[1], Adi Cohen[1,2], Gal Ben-Zvi[1,2], Rani Gera[1,2,3], Shiran Oren[1,2], Dana Roll[1], Gal Rozic[1], Anastasia Saliy[1], Niv Tik[2,4], Galia Tsarfati[5], Ido Tavor*[2,4,6], Tom Schonberg*[1,2,6], Yaniv Assaf*[1,2,6]

* corresponding authors equal contribution

1. School of Neurobiology, Biochemistry and Biophysics, Faculty of Life Science, Tel Aviv University, Tel Aviv, Israel

2. Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel

3. School of Psychological Sciences, Tel Aviv University, Tel Aviv, Israel

4. Department of Anatomy and Anthropology, Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

5. Division of Diagnostic Imaging, Sheba Medical Center, Tel-Hashomer, affiliated to the Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

6. The Strauss Center for Computational Neuroimaging, Tel Aviv University, Tel Aviv, Israel

Abstract

The COVID-19 outbreak introduced unprecedented health-risks, as well as pressure on the financial, social, and psychological well-being due to the response to the outbreak[1–4]. Here, we examined the manifestations of the COVID-19 outbreak on the brain structure in the healthy population, following the initial phase of the pandemic in Israel. We pre-registered our hypothesis that the intense experience of the outbreak potentially induced stress-related brain modifications[5–8]. Volumetric changes in $n = 50$ participants scanned before and after the COVID-19 outbreak and lockdown, were compared with $n = 50$ control participants that were scanned twice prior to the pandemic. The pandemic provided a rare opportunity to examine brain plasticity in a natural experiment. We found volumetric increases in bilateral amygdalae, putamen, and the anterior temporal cortices. Changes in the amygdalae diminished as time elapsed from lockdown relief, suggesting that the intense experience associated with the pandemic outbreak induced volumetric changes in brain regions commonly associated with stress and anxiety[9–11].

Main text

Since early 2020, the world has been coping with the outbreak of the coronavirus disease 2019 (COVID-19) pandemic that has infected millions with devastating numbers of deaths globally. As an initial response to the first wave of the outbreak, countries closed their borders and implemented a series of ad-hoc laws and orders to restrict the spread of the disease. Countries with major outbreaks such as China, Italy, and Spain enforced stringent restriction of movement for a limited period, referred to here as 'lockdown'. Although lockdowns contributed to restricting the health risks of the outbreak[12], they also had a negative impact on the social,

economic and psychological well-being of the general population, leading to one of the sharpest

declines in economic growth over the past decades[1,2]. Lockdowns also led to high rates of stress

and anxiety which were attributed in large to the social and financial consequences of responding

to the health crisis[4]. It is now evident that the indirect consequences of the pandemic affected a

much larger proportion of the population, having an impact of no lesser gravity than the actual

health risks that were meant to be prevented[3,13].

In Israel, a strict lockdown period was issued from mid-March until the end of April. During its

peak, most unessential businesses were closed and civilians' movement for non-essential

destinations was restricted for a radius of 100 meters from their homes. Prior to COVID-19, the

country had experienced a period of peak economic prosperity[14], which was interrupted by the

outbreak, leading to unprecedented unemployment rates (reaching nearly 30% of the work-force

in April 2020) and the collapse of several sectors such as aviation, tourism, and culture[15,16]. The

outbreak period was characterized with acute uncertainty and increase in anxiety, regarding both

the health and socioeconomic effects of the pandemic[17].

Over the past years several studies demonstrated brain plasticity detected using T1-weighted

magnetic resonance imaging (MRI)[18–20]. The current work was initiated as a reaction to the

outbreak of COVID-19 in Israel, aimed to study the structural brain plasticity in the general

population following a real-life event of global scale. For this purpose, we examined $n = 50$ test

group participants that were scanned with T1-weighted MRI prior to the outbreak and returned

for a follow-up scan after the lockdown period. The structural changes of the study group (before

versus after the outbreak) were compared to those of $n = 50$ control participants who were

scanned twice before the COVID-19 outbreak. The unique circumstances imposed due to the

COVID-19 lockdown created rare settings for a natural experiment to examine the effect of a

real-world intense event on brain plasticity.

All participants were healthy, without a history of neurological or psychiatric disorders, did not

show COVID-19 symptoms, and were not diagnosed carrying the virus (see the methods section

for further demographic information). The hypotheses and general design of the current study

were pre-registered prior to the completion of data collection and were based on a small

independent pilot sample with $N = 16$ participants ($n = 8$ participants in each group). The data

and analysis codes are openly shared online (project page: https://osf.io/wu37z/; preregistration:

https://osf.io/k6xhn/).

Prior to their follow-up MRI scan session, we asked participants of the post-lockdown test group

to fill in a short questionnaire regarding the lockdown period (see methods). Of the participants

who agreed to reply, 79.6% reported they did not leave their home for non-essential needs,

38.8% indicated an increased feeling of anxiety following the lockdown, 79.6% met no more

than 3 people, 34.7% anticipated that their future behavior will change after the lockdown,

44.9% did not meet with their parents at all, 42.9% indicated that their employment status was

reduced to part-time or unemployment, 46.8% reported they were concerned about their personal

future well-being. In an exploratory principal component analysis (PCA), we found that two

principal components best explain the variability in the data, explaining together 42.58% of the

variance. The first component was highly loaded with increased feelings of anxiety, and the

second was related to items describing increased social isolation (Figure 1). These analyses (not

included in the pre-registration) indicate that the pandemic outbreak had a significant impact on

the social and psychological well-being of most participants in our study.

Figure 1. Principal component analysis of COVID-19 questionnaire.
A principal component analysis (PCA) of the responses to the questionnaire revealed two main themes characterized the participants. **a.** An increased feeling of anxiety dominated the first principal component (x-axis), while the three items relating to social distancing (staying at home, meeting parents and meeting more than 3 people) contributed together to the second component (y-axis). **b.** Visualization of participants dispersion across the two principal components and their categorization into binary anxiety and isolation groups. High isolation was defined as directional response to all three social-isolation items described above. Points represent individual participants.

Based on our pilot study results and previous studies of stress-related morphological brain changes[5–8] we hypothesized that the focus of volumetric changes will be observed mainly in the amygdalae. The anatomical data were used as input for deformation and surface-based morphometry (SBM) analysis using the CAT12 toolbox (http://www.neuro.uni-jena.de/cat/, University of Jena) for SPM12 (http://www.fil.ion.ucl.ac.uk/spm/software/spm12/, Wellcome Trust Centre for Neuroimaging). The brain was segmented to 58 regions based on the cortical and subcortical nuclei classifications of the Hammers atlas (Hammers et al., 2003). Following surface reconstruction, each participant's individual gray matter volume was estimated for each of the 58 anatomically defined regions of interest (ROIs). This procedure accounted for the longitudinal nature of the data, performing the analysis on both scans simultaneously. To avoid voxel-based multiple comparisons, we performed a region-based analysis (following surface

projection to the Hammer atlas) and corrected for multiple comparisons using the Benjamini-Hochberg correction[21] to control for false discovery rate (FDR; at $p < 0.05$ following correction). Validation of this pipeline was performed using simulated data and by comparing the results with other software (see methods).

Using a linear mixed model, we examined volumetric changes, testing for regions with stronger effects for the test group, compared to the control group. Examining an interaction effect of session (baseline versus follow-up scans) and experimental group (test versus control) revealed ten anatomical brain regions (composed of bilateral five unique regions in both hemispheres) in which volumetric increases were observed uniquely for the test group (Table 1 and Figure 2). Most prominently, as we expected and pre-registered, we found a robust effect in the bilateral amygdalae. We also observed a significant effect bilaterally in the putamen, and in three anatomical regions within the ventral anterior temporal cortex adjacent to each other, namely in the medial part of the anterior temporal lobe, the fusiform gyrus, and the parahippocampal gyrus. To examine the spatial distribution within significant ROIs, we performed an additional post-hoc voxel-based analysis, which allowed us to visualize the changes within the significant ROIs (Figure 2a). Examining the post-hoc voxel-based results revealed that volumetric changes occurred throughout the entire surface of bilateral amygdalae, while in the putamen the effects occurred mainly in the dorsal area. In the ventral anterior temporal cortices, large connected clusters of volumetric change spanned throughout the three adjacent temporal ROIs, thus suggesting that the three ROIs shared a similar origin. In all ROIs, we ensured that the significant effect was apparent for the test group but not for the control group (see methods), suggesting that the reported interaction effects originated from volumetric changes in the test group following the COVID-19 outbreak and its related lockdown period (Figure 2b).

134   **Table 1. Surface based morphology analysis results**

Region	Hemi-sphere	Interaction estimate (95% CI)	Interaction p (FDR adj.)	Session estimate (95% CI) [a]	Session p (FDR adj.)
Amygdala	Left	0.09 [0.05, 0.13]	$2.4E^{-5}$ (0.001)	0.08 [0.05, 0.11]	$9.8E^{-6}$ ($2.1E^{-4}$)
	Right	0.08 [0.03, 0.13]	0.003 (0.030)	0.08 [0.05, 0.11]	$1.6E^{-5}$ ($2.3E^{-4}$)
Putamen	Left	0.19 [0.09, 0.29]	$4.1E^{-4}$ (0.006)	0.13 [0.06, 0.2]	$4.0E^{-4}$ (0.002)
	Right	0.17 [0.08, 0.26]	$2.4E^{-4}$ (0.005)	0.14 [0.08, 0.2]	$1.1E^{-5}$ ($2.1E^{-4}$)
Anterior temporal lobe (medial part)	Left	0.25 [0.12, 0.38]	$1.8E^{-4}$ (0.005)	0.15 [0.07, 0.23]	$4.7E^{-4}$ (0.003)
	Right	0.21 [0.07, 0.35]	0.004 (0.030)	0.15 [0.05, 0.25]	0.004 (0.023)
Parahippocampal gyrus	Left	0.09 [0.03, 0.15]	0.006 (0.035)	0.04 [0, 0.08]	0.029 (0.085)
	Right	0.11 [0.04, 0.18]	0.003 (0.030)	0.08 [0.03, 0.13]	0.002 (0.009)
Fusiform gyrus	Left	0.08 [0.03, 0.13]	0.007 (0.036)	0.06 [0.03, 0.09]	$3.8E^{-4}$ (0.003)
	Right	0.11 [0.04, 0.18]	0.002 (0.022)	0.05 [0, 0.1]	0.044 (0.111)

135   [a] Session estimate examined the effect of baseline versus follow-up scan in the post-lockdown
136   test group. This parameter was used to validate that the interaction effect observed between the
137   group stemmed from a robust effect in the test group (see methods).
138

Figure 2. Volumetric changes results.
An interaction effect for time (baseline versus follow-up scan) and group (test versus control) was evaluated on segmented surfaces in an SBM analysis. Significant interaction effects were observed bilaterally in the amygdala and putamen ROIs, as well as in three ventral temporal cortical ROIs. **a.** To examine spatial patterns within the identified ROIs, a post-hoc voxel-based analysis was conducted within each ROI mask. Light red contours represent segmentation borders of the ROIs. **b.** Individual distribution of the results in the control group (light colors) and test group (dark colors). Box-plot center, hinges, and whiskers represent the median, quartiles, and   from the hinges, respectively. A notch of   represent an estimated 95% confidence interval for comparing medians. Dots represent individual participants. Abbreviated ROI names: AntMedTeLo = anterior temporal lobe (medial part); FusGy = fusiform gyrus, ParHipGy = Parahippocampal gyrus.

To evaluate and control for the effect of time between scans and time from lockdown, we included in the model two additional covariates - the time between scans (TBS; which was generally longer for the test group) and time following lockdown (TFL; see supplementary methods for more details). The two covariates were not correlated with each other in our test

group sample (r = -0.106, $t(48)$ = -0.74, p = 0.463). Our reported regions demonstrated

significant volumetric change above and beyond these covariates. After FDR correction, no

region showed an effect of TBS. However, we did find a negative effect of TFL in the two

amygdalae ROIs and the left fusiform gyrus, suggesting that the volumetric changes in these

regions moderated as time following lockdown elapsed. Based on these results we estimated the

time to decay as the estimated number of days from lockdown until volumetric changes returned

to normal levels, similar to those of the control group (left amygdala: β_{TFL} = -0.41, $t(47)$ = -3.1, p

= 0.003, $p_{adj.}$ = 0.048, time to decay = 95 days; right amygdala: β_{TFL} = -0.54, $t(47)$ = -4.38, p =

6.7E^{-5}, $p_{adj.}$ = 0.002, time to decay = 83 days; left fusiform gyrus: β_{TFL} = -0.54, $t(47)$ = -4.44, p =

5.5E^{-5}, $p_{adj.}$ = 0.002, time to decay = 82 days; Figure 3).

Figure 3. Time following lockdown effect on volumetric changes.
The time from lockdown relief until the follow-up scan session (TFL) was introduced as addition
covariate to the model, revealing significant effect in the two amygdalae and left fusiform gyrus.
Points represent individual participants in the post-lockdown test, p-values were FDR adjusted
for multiple comparisons. Abbreviated ROIs: AntMedTeLo = anterior temporal lobe (medial
part); FusGy = fusiform gyrus, ParHipGy = Parahippocampal gyrus.

Additional exploratory analyses examined the association of volumetric changes and the reported

experience during lockdown. We found no strong association between the two, as reported in the

supplementary results.

In conclusion, our study demonstrates that volumetric change patterns in the brain occurred

following the COVID-19 initial outbreak period and restrictions. Previous studies demonstrated

brain plasticity using T1-weighted MRI following planned interventions[18–20]. The current work

uniquely demonstrates stark structural brain plasticity following a major real-life event.

Our findings show changes in gray matter in the amygdala, putamen and ventral anterior

temporal cortex. The changes in the amygdalae showed a temporal-dependent effect, related to

the time elapsed from lockdown but not the duration from the baseline scan. It should be noted

that although lockdown restrictions had initially reduced infection rates in Israel, just one month

after the lockdown was lifted, the number of infected cases started to rise again and reached

higher number of active infected cases by the end of data collection, compared with the peak

numbers during the actual lockdown period (approximately 2,000 daily new cases by the end of

July versus under 750 new daily cases during the peak of the lockdown period in April[22], see

supplementary Figure 1). This suggests that the effects observed in the current study are less

likely to be attributed to the concrete health risks of COVID-19, but rather to the first wave of

the outbreak, characterized with perceived uncertainty.

The current study was in many aspects unplanned; thus we are left with only partial answers as

to which specific components of the COVID-19 outbreak led to the neural changes observed in

the healthy participants that took part in our study. The involvement of the amygdala may

suggest that stress and anxiety could be the source of the observed phenomenon, due to its well-

recorded functional and structural associations[5–11]. Nevertheless, it is hard to draw clear

conclusions as many aspects of life have changed in this time period, and could have potentially

affected different regions in the brain – from limiting social interactions, increased financial

stress, changes in physical activity, work routine, and many more. The limited behavioral data

collected in the current study did not provide a strong connection to the imaging results, and thus

future work could try to better address the complex brain-behavioral associations in this real-life

experience. Nonetheless, our findings show healthy young adults, with no records of mental

health issues, were deeply affected by the outbreak of COVID-19. We suggest that policy makers

take into consideration the impact of their actions on the general well-being of the population

they seek to help, alongside the efficacy of disease prevention.

Acknowledgements

This work was supported by the Israeli Science Foundation granted to Yaniv Assaf (ISF

1314/15), Tom Schonberg (2004/15), and Ido Tavor (ISF 1603/18). Tom Salomon was supported

by the Nehemia Levtzion fellowship and the Fields-Rayant Minducate Learning Innovation

Research Center. We would also want to thank Dr. Daniel Barazany, the manager of the Strauss

MRI center, for his invaluable assistance throughout the study.

Author contributions

T.Sa. wrote the manuscript with Y.A, assisted with the study dsign and analyzed the data. Data

was collected by A.C., R.G, S.O, G.R., D.R, and A.S. Free-surfer analysis was made by G.B-Z.

N.T performed VBM validation analysis. G.T. provided support and medical supervision. I.T.,

T.Sc. and Y.A. conceived the study, wrote the manuscript and supervised the study. Y.A

performed the preprocessing and analysis of imaging data. All Author contributed intellectually

and reviewed the manuscript.

Competing interests

The authors declare no competing financial interests.

References

1. Zhang, D., Hu, M. & Ji, Q. Financial markets under the global pandemic of COVID-19.
Financ. Res. Lett. 101528 (2020). doi:10.1016/j.frl.2020.101528
2. Fernandes, N. Economic effects of coronavirus outbreak (COVID-19) on the world
economy Nuno Fernandes Full Professor of Finance IESE Business School Spain. *SSRN*
Electron. Journal, ISSN 1556-5068, Elsevier BV, 0–29 (2020).
3. Gruber, J. *et al.* Mental health and clinical psychological science in the time of COVID-
19: Challenges, opportunities, and a call to action. *Am. Psychol.* (2020).
doi:https://doi.org/10.1037/amp0000707
4. Taylor, S. *et al.* COVID stress syndrome: Concept, structure, and correlates. *Depress.*
Anxiety 1–9 (2020). doi:10.1002/da.23071
5. Ganzel, B. L., Kim, P., Glover, G. H. & Temple, E. Resilience after 9/11: Multimodal
neuroimaging evidence for stress-related change in the healthy adult brain. *Neuroimage*
40, 788–795 (2008).
6. Hölzel, B. K. *et al.* Stress reduction correlates with structural changes in the amygdala.
Soc. Cogn. Affect. Neurosci. **5**, 11–17 (2009).
7. Rogers, M. A. *et al.* Smaller amygdala volume and reduced anterior cingulate gray matter
density associated with history of post-traumatic stress disorder. *Psychiatry Res. -*
Neuroimaging **174**, 210–216 (2009).
8. Schienle, A., Ebner, F. & Schäfer, A. Localized gray matter volume abnormalities in
generalized anxiety disorder. *Eur. Arch. Psychiatry Clin. Neurosci.* **261**, 303–307 (2011).
9. Stevens, J. S. *et al.* Amygdala Reactivity and Anterior Cingulate Habituation Predict
Posttraumatic Stress Disorder Symptom Maintenance After Acute Civilian Trauma. *Biol.*

Psychiatry **81**, 1023–1029 (2017).

10.   Bryant, R. A. *et al.* Enhanced amygdala and medial prefrontal activation during

nonconscious processing of fear in posttraumatic stress disorder: An fMRI study. *Hum.*

Brain Mapp. **29**, 517–523 (2008).

11.   Mochcovitch, M. D., Da Rocha Freire, R. C., Garcia, R. F. & Nardi, A. E. A systematic

review of fMRI studies in generalized anxiety disorder: Evaluating its neural and

cognitive basis. *J. Affect. Disord.* **167**, 336–342 (2014).

12.   Vinceti, M. *et al.* Lockdown timing and efficacy in controlling COVID-19 using mobile

phone tracking. *EClinicalMedicine* (2020). doi:10.1016/j.eclinm.2020.100457

13.   Qiu, J. *et al.* A nationwide survey of psychological distress among Chinese people in the

COVID-19 epidemic: Implications and policy recommendations. *Gen. Psychiatry* **33**, 1–4

(2020).

14.   Bank of Israel Research Department. The Decline in Unemployment in Israel by

International Comparison. in *Bank of Israel Annual Report 2019* (2020).

15.   Bank of Israel Research Department. *The unemployment rate and its definition during the*

corona period. (2020).

16.   Bank of Israel Research Department. *Changes in credit card purchases as a result of the*

corona crisis. (2020).

17.   Tzur Bitan, D. *et al.* Fear of COVID-19 scale: Psychometric characteristics, reliability and

validity in the Israeli population. *Psychiatry Res.* **289**, 113100 (2020).

18.   Maguire, E. A. *et al.* Navigation-related structural change in the hippocampi of taxi

drivers. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 4398–4403 (2000).

19.   Jung, W. H. *et al.* Exploring the brains of Baduk (Go) experts: Gray matter morphometry,

resting-state functional connectivity, and graph theoretical analysis. *Front. Hum.*

Neurosci. **7**, 633 (2013).

20.   Draganski, B. *et al.* Changes in grey matter induced by training. *Nature* **427**, 311–312

(2004).

21.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate□: A Practical and

Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).

22.   Israel Ministry of Health. Coronavirus in israel - Overview (in Hebrew). (2020). Available

at: https://datadashboard.health.gov.il/. (Accessed: 31st August 2020)

23.   Champely, S. *et al.* Package 'pwr': Basic Functions for Power Analysis. *CRAN Repos.*
281   (2018).
282   24.   Manjón, J. V., Coupé, P., Martí-Bonmatí, L., Collins, D. L. & Robles, M. Adaptive non-
283   local means denoising of MR images with spatially varying noise levels. *J. Magn. Reson.*
284   *Imaging* **31**, 192–203 (2010).
285   25.   Dahnke, R., Yotter, R. A. & Gaser, C. Cortical thickness and central surface estimation.
286   *Neuroimage* **65**, 336–348 (2013).
287   26.   Kassambara, A. & Mundt, F. factoextra: Extract and Visualize the Results of Multivariate
288   Data Analyses. R package version 1.0.7 (2020). Available at: https://cran.r-
289   project.org/package=factoextra.
290

291   **Supplementary Methods**

Codes and Data Accessibility

Our sample size, hypotheses and analyses plan were pre-registered on the Open Science
Framework (OSF), soon after data collection began, but prior to completion of the data collection
and data analysis (project page: https://osf.io/wu37z/; preregistration: https://osf.io/k6xhn). All
behavioral processed imaging data along with the analysis codes are shared on the OSF project
page. Uncorrected and small-volume corrected statistical maps of the voxel-based results
described in the current work are available at https://neurovault.org/collections/8591/.

Participants

The study included two groups: A test group scanned before and after COVID-19 lockdown, and
a control group, scanned twice before COVID-19 lockdown. All participants had no background
of neurological disorders, did not show symptoms for COVID-19 and were not diagnosed as
carriers of the virus. The study was approved by the ethics committee of Tel Aviv University and
institutional review board (IRB) at the Sheba Tel-Hashomer medical center. Since the IRB
protocol allowed us to scan the participants several times over a long period of time, we were
able to collect the data from participants who were scanned prior to COVID-19 outbreak and
invite them back for a follow-up scan as part of the longitudinal study they have agreed to take
part in. Participants received monetary compensation for their time and gave their informed
consent to take part in a longitudinal experiment aimed to examine brain plasticity across several
sessions, which was initially not directly related to COVID-19 outbreak.

The test group comprised of $n = 50$ participants who were scanned before and after COVID-19

lockdown (Δ Time between scans: $M = 309.3$, $SD = 207.5$, range = 67 - 1460 days; Age: $M =$

30.1, $SD = 6.65$, range = 21 - 48; Females: n = 20, prop. = 40%). The lockdown period began on

March 25[th], and was gradually relieved throughout late April. We mark here May 1[st] as the

lockdown relief date, as on this day an issued 100-meters movement limit for non-essential needs

was lifted. The test group data collection started as soon as lockdown relief took place, for a

period of approximately 3 months, until the end of July, 2020 (Δ Time from lockdown relief: M

$= 57$, $SD = 24.62$, range = 9 - 89 days; see Supplementary figure 1 for the study timeline).

320
Supplementary Figure 1. Study timeline and outbreak data
On February 21[st], 2020, the first COVID-19 case in Israel was recorded. Daily new cases are
presented on green bars (right y-axis), along with the cumulative number of cases and recoveries
(left y-axis). Data was retrieved and modified based on the Israeli Ministry of Health reports[22]. A
lockdown was issued on March 25[th], which was gradually released until the removal of the 100-
meter restriction on May 1[st], marking lockdown onset and relief, respectively (shorter vertical
dashed line). MRI data of the test group was collected from May 10[th] to July 29[th] (longer vertical
dashed line). Red bars on top represent the number of participants scanned for the study each
day.
330

As a control group, we used the data of $n = 50$ participants who were scanned twice using a

similar protocol before COVID-19 lockdown (Δ Time between scans: $M = 126.7$, $SD = 190.4$,

range = 21 - 886 days; Age: M = 27.3, $SD = 5.63$, range = 19 - 42; Females: n = 23, prop. =

46%).

The minimal sample size was determined and pre-registered (https://osf.io/uktsn), based on a
80% power analysis conducted using R 'pwr' package[23], on a pilot study with $N = 16$
participants ($n = 8$ in each group). We decided to collect a minimum of $n = 37$ participants which
should provide 80% to detect the group and session interaction effect with $\alpha = .05$, within both
the left and right Amygdala. We originally committed to collect $n = 37$ participants in each
group, under the assumption it would be difficult to complete the sample due to COVID-19
limitations. Eventually, thanks to further relief in COVID-19 restrictions, we were able to extend
the sample size to $n = 50$ in each group.

The results remain generally consistent, even when the data included only the first $n = 37$
participant; demonstrating significant effects in the bilateral amygdalae, putamen,
parahippocampal gyrus and the left anterior temporal lobe. In this smaller sample, we did not
find significant effects in the right anterior temporal lobe, nor in the fusiform gyrus. We also
found volumetric increase effects that were not identified using the full sample in the left nucleus
accumbens, left cuneus, and left insula.

Imaging data

Acquisition protocol. Imaging data were acquired using a 3T Siemens Prisma scanner, with a
64-channel head coil. For the structural data, T1w high resolution (1-mm^3) whole brain images
were acquired with a magnetization prepared rapid gradient echo (MPRAGE) pulse sequence
with repetition time (TR) of 2.53s, echo time (TE) of 2.88ms, flip angle (FA) = 7°, field-of-view
(FOV) = 224 × 224 × 208 mm, resolution = 1 × 1 × 1. (see below).

Some participants were also scanned with diffusion-weighted echo-planar imaging (DW EPI)
sequence and some with functional gradient-echo EPI (GE EPI) in a resting state scan. The
analyses of these scans are beyond the scope of the current study.

Data processing and analysis. The T1w MPRAGE anatomical scans were used for a surface-
based morphometry (SBM) analysis. From the images we estimated the pial and inner surfaces
of the cortex and projected those into the Hammers atlas (Hammers et al., 2003). Data were
preprocessed in SPM12 (http://www.fil.ion.ucl.ac.uk/spm/software/spm12/, Wellcome Trust
Centre for Neuroimaging) and SPM based CAT12 (Computational Anatomy Toolbox 12;
http://www.neuro.uni-jena.de/cat/, University of Jena) extension. We deployed the CAT12
surface-processing pipeline, which includes skull striping, a denoising filter[24] projection-based
thickness estimation[25], partial volume correction, and spatial normalization to MNI space.

Surface-based volumetric data of cortical and subcortical regions were segmented based on the
Hammers Atlas, segmenting the volumetric data into 58 anatomically defined regions. Ten
additional ROIs of non-gray matter (ventricles, white matter, brain-stem and cerebellum ROIs)
were excluded from statistical analyses. To evaluate the effect of lockdown on volumetric
imaging data we ran a mixed linear model on the data within each one of the 58 anatomical
regions, examining the effect of session (baseline versus follow-up scan) and group interaction
(test versus control), controlling for time between scans (TBS) and time following lockdown
(TFL) covariates. Both covariates were mean centered before they were added to the model.

To identify our regions of interest we included only regions that showed both a significant
interaction effect (i.e. the volumetric difference between the two sessions was significantly
different for the test and control group), and a significant session effect within the test group (i.e.
a significant difference between the two sessions for the test group). Results were corrected for
multiple comparisons using the false discovery rate (FDR) correction, based on the number of
brain regions tested[21]. Following the analysis pipeline, we identified ten significant ROIs:
bilateral amygdalae, putamen, parahippocampal gyrus, the medial part of the anterior temporal
lobe, and the fusiform gyrus.

An additional ROI of the right inferior and middle temporal gyri showed a significant interaction
effect (interaction estimate = 0.17, 95% CI = [0.05, 0.29], $p = 6.1\text{E}^{-3}$, $p_{\text{adj.}}$ = 0.035), however
examining the test group separately, we could not identify a significant session effect (session
estimate = 0.04, 95% CI = [-0.02, 0.10], $p = 0.259$, $p_{\text{adj.}}$ = 0.424). Therefore, it is harder to
interpret that this interaction effect stemmed from the test group. Thus, we did not include this
ROI as one of our significant ROIs. A less robust session effect within the test group was also
observed for the left parahippocampal gyrus (session estimate = 0.04 [0.00, 0.08], $p = 0.029$, $p_{\text{adj.}}$
= 0.085) and right fusiform gyrus ROI (session estimate = 0.05 [0.00, 0.10], $p = 0.044$, $p_{\text{adj.}}$ =
0.111), however as these ROIs demonstrated strong interaction effects and their session effects
were significant before FDR correction, we decided to report them together with the other
significant regions. A similar procedure was used in the pre-registration (i.e. including only
regions that showed a significant interaction and an effect in the test group), however in the
analysis of the pilot for the pre-registration, we used uncorrected results due to the small sample
size.

To examine the spatial distribution of our effect within significant ROIs, that were identified
with the SBM analysis, we performed an additional post-hoc voxel-based analysis. We projected
the data on a voxel-based map, effectively examining which voxels demonstrated an interaction
effect. Then, we used anatomical masks of ROIs which were found to be significant in the SBM
analysis, to visualize the results within these regions, similarly to a small-volume correction
analysis (Figure 2a)

Pipeline validation

To validate the imaging processing protocol we used two approaches, before data collection was
completed (at the time of the pre-registration finalization). As many surface-based software
focus on analysis of cortical surfaces, rather than subcortical regions, we aimed to validate that
our SBM protocol using CAT12 could reliably identify subcortical morphological changes, such
as the ones we observed in the current study within the amygdala and putamen. To test the
detection capabilities of our protocol, we generated simulated data with volumetric changes in
the amygdala and ran the pipeline on the simulated data. A volumetric increase in the amygdala
was simulated using a hand-drawn polygon mask, surrounding the left amygdala on the original
T1w images. Within this polygon 3D mask, the signal intensity was artificially changed.
Following this procedure, both the original and modified T1w images underwent the same
CAT12 pipeline. The analysis was performed on 10 participants. We were able to identify the
simulated volumetric changes within the subcortical amygdala nuclei (see supplementary Figure
2).

416
417   **Supplementary figure 2.** Data simulation
418   Sub-cortical changes in the left amygdala were simulated in ten participants. The SBM pipeline
419   applied in CAT12 identified the simulated change effect. Data was projected from surfaces back
420   to a voxel-based map for visualization in the current figure.
421

In an additional validation procedure, we reanalyzed our results with an additional analysis
pipeline. Raw T1-weighted maps were preprocessed using FreeSurfer. Using this alternative
analysis pipeline, we found similar results, including robust effects on the temporal cortical
regions in addition to other cortical regions. It should be noted that the FreeSurfer pipeline
analyze only cortical surface, thus the analysis did not include subcortical regions of the
amygdala and putamen.
The results of both validations indicated our analysis pipeline could reliably identify regional
volumetric estimations. Using CAT12 provided an advantage by performing a longitudinal
analysis of subcortical regions, including the amygdala which was pre-hypothesized and of great
importance in the current work.

Behavioral data

Data collection. To evaluate participants' experience in the peak days of the COVID-19
outbreak, we asked them to fill out a 7-items questionnaire regarding their experience of the
COVID-19 lockdown (see supplementary Table 1 for a description of the items). The
questionnaires were filled out soon after the initiation of the study, when the lockdown's
stringent 100-meters limitation was lifted. Most participants filled out the questionnaire on the
day of the post-lockdown scan session, some filled it a few days before their second scanning
session. A total of $n = 77$ participants filled out the COVID-19 questionnaire and comprised the
potential pool of test group participants for the current study, out of which $n = 50$ were sampled
and scanned. One participant was scanned but did not complete the questionnaire, therefore this
participant's behavioral data were not used and analyses of the questionnaire were based on $n =$
49 valid participants.

444
445

446   **Supplementary Table 1.** COVID-19 lockdown questionnaire

Question	Possible Answers (prop.)	Binary outcome (prop.)
1. Did you stay home during the lockdown, except for essential needs / did not leave at all?	0 - no 1 - yes	0 - no (20.4%) 1 - yes (79.6%)
2. Did the lockdown increase your feeling of anxiety?	0 - no 1 - yes	0 - no (61.2%) 1 - yes (39.8%)
3. With how many people did you meet during the lockdown (including people you are living with at home)?	0 - none 1 - up to three people 2- up to five people 3 - up to ten people	0 - up to three (79.6%) 1 - more (20.4%)
4. Do you think your behavior will change following the lockdown?	0 - no 1 - yes	0 - no (65.3%) 1 - yes (34.7%)
5. How did your meeting with your parents' routine look like during the lockdown?	0 - same as before the lockdown 1 - with precaution measurements: distancing, mask, etc. 2 - did not visit at all	0 - as before or with precautions (44.9%) 1 - did not visit (55.1%)
6. What was your employment status during the lockdown?	0 - same as before lockdown 1 - full time working from home 2 - part time working from home 3 - Furlough / unemployed	0 - unemployed / part time (42.9%) 1 - same as before / full time from home (57.1%)
7. How concerned are you with the long-term effect of the lockdown, regarding yourself?	1 - 5 scale	0 - low, score 1-2 (53.1%) 1 - moderate-high, score 3-5 (46.9%)

Data analysis. Responses to the lockdown questionnaire were coded into binary responses,
based on the sample median, splitting the sample into relatively similar sized groups for each
item. To identify the main themes in the questionnaire which could be correlated with the
imaging data, we performed a PCA analysis on the binarized data, using the "factoextra" R
package[26]. We found two principal components, which explained 42.6% of the variance in the
sample data. These two components were extracted and correlated with the change in gray matter
volumetric data in our regions of interest.

455   **Supplementary results**

As an exploratory analysis, we examined whether the volumetric brain changes were associated
with the psychological constructs identified in our PCA analysis, based on participants' self-
reports. We used two linear models aimed to explain the variability in each of the principal
components, using the volumetric changes as our model features. Overall neither one of the two
PCs were well associated with the volumetric changes (Principal component 1 model: $R^2 =$
0.205, $F(10,38) = 0.98$, $p = 0.475$; Principal component 2 model: $R^2 = 0.33$, $F(10,38) = 1.89$, $p =$
0.077).
While we did find some sporadic ROIs showing significant contribution within the models
(measured as the significance of the ROI's β estimates), after FDR correction by the number of
features in the model, none of the ROIs demonstrated a significant association with the PCs ($p_{adj.}$
> 0.05).
Finally, we examined correlation patterns of the volumetric change for all brain regions aiming
to identify shared change patterns across multiple ROIs. Hierarchical clustering of the correlation
matrices revealed different patterns between the two groups (Supplementary figure 3). In the test
group, three principal groups of clusters could be identified - the first included the palladium,
hippocampus, amygdala, putamen, insula (all bilateraly), and right anterior cingulate cortex, the
second cluster included mostly occipito-temporal cortical and subcortical nuclei, and the third
included highly correlated regions of the frontal, parietal and occipital cortices. All regions that
passed the statistical threshold of the SBM analysis (Table 1), except for the left fusiform gyrus,
were grouped closely together within the first two clusters, and showed low to negative
correlations the ROIs of the third cluster. This analysis could suggest that the origin of the
volumetric change observed in the regions of the two clusters might be different. The regions
that appear in cluster 1 are often reported in the studies that explore brain changes following
stress, anxiety or traumatic events[5–8], while the regions of the second cluster are less associated
with specific phenomena.
A different pattern was observed in the control group, where the correlation pattern demonstrated
stronger volumetric synchrony, (Supplementary figure 3b). This could suggest that changes in
the control group were much more affected by within-participant effects, rather than an
exogenous effect (which could be the outbreak and lockdown in the test-group). It is important to
note in this context that in addition to the experience of the lockdown, the participants in the test
group also had longer time gaps between the two scanning sessions, which might provide an

alternative explanation for the stronger homogeneity in volumetric changes within the control

group.

490
Supplementary figure . Volumetric changes correlation matrices
Correlation coefficients were calculated between the volumetric change values for all ROI pairs,
analyzed separately for the test **(a)** and the control group **(b)**, clustered according to Euclidean
distances into dendrograms. In the test group, the first and second cluster contained the
subcortical nuclei (amygdala and putamen) and temporal ROI which were found significant in
the SBM interaction analysis, respectively (highlighted in italic bold font). In the control group, a
more homogeneous change pattern was observed with more robust correlation coefficients
between the ROIs. Pearson correlation coefficients are represented by the color scheme.