Brain volumetric changes in the general population following the COVID-19 outbreak and lockdown

Tom Salomon[1], Adi Cohen[1,2], Daniel Barazany[3], Gal Ben-Zvi[1,2], Rotem Botvinik-Nezer[1,2,4], Rani Gera[1,2,5], Shiran Oren[1,2], Dana Roll[1], Gal Rozic[1], Anastasia Saliy[1], Niv Tik[2,6], Galia Tsarfati[7], Ido Tavor*[2,3,6], Tom Schonberg*[1,2,3], Yaniv Assaf*[1,2,3]

* corresponding authors equal contribution

1. School of Neurobiology, Biochemistry and Biophysics, Faculty of Life Science, Tel Aviv University, Tel Aviv, Israel

2. Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel

3. The Strauss Center for Computational Neuroimaging, Tel Aviv University, Tel Aviv, Israel

4. Department of Psychological and Brain Sciences, Dartmouth College, Hanover, New Hampshire, USA

5. School of Psychological Sciences, Tel Aviv University, Tel Aviv, Israel

6. Department of Anatomy and Anthropology, Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

7. Division of Diagnostic Imaging, Sheba Medical Center, Tel-Hashomer, affiliated to the Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

Abstract

The COVID-19 outbreak introduced unprecedented health-risks, as well as pressure on the financial, social, and psychological well-being due to the response to the outbreak. Here, we examined the manifestations of the COVID-19 outbreak on the brain structure in the healthy population, following the initial phase of the pandemic in Israel. We pre-registered our hypothesis that the intense experience of the outbreak potentially induced stress-related brain modifications. Volumetric changes in $n = 50$ participants who were scanned before and after the COVID-19 outbreak and lockdown, were compared with $n = 50$ control participants who were scanned twice prior to the pandemic. The pandemic provided a rare opportunity to examine brain plasticity in a natural experiment. We found volumetric increases in bilateral amygdalae, putamen, and the anterior temporal cortices. Changes in the amygdalae diminished as time elapsed from lockdown relief, suggesting that the intense experience associated with the pandemic outbreak induced transient volumetric changes in brain regions commonly associated with stress and anxiety.

Main

Since early 2020, the world has been coping with the outbreak of the coronavirus disease 2019 (COVID-19) pandemic that infected millions with devastating numbers of deaths globally. As an initial response to the first wave of the outbreak, countries closed their borders and implemented a series of ad-hoc laws and orders to restrict the spread of the disease. Countries with major outbreaks such as China, Italy, and Spain enforced stringent restriction of movement for a limited period, referred to here as 'lockdown'. Although lockdowns contributed to restricting the health risks of the outbreak[1], they also had a negative impact on the social, financial and mental

well-being of the general population, leading to one of the sharpest declines in economic growth

over the past decades[2,3]. COVID-19 outbreak also led to high rates of stress and anxiety that

were largely attributed to implications beyond actual health risk, such as the difficulties of social

isolation and financial consequences of responding to the health crisis[4]. It is now evident that the

indirect consequences of the pandemic affected a much larger proportion of the population,

having an impact of no lesser gravity than the actual health risks that were meant to be

prevented[5,6].

In Israel, a strict lockdown period was issued from mid-March until the end of April. During its

peak, most unessential businesses were closed and civilians' movement for non-essential

destinations was restricted to a radius of 100 meters from their homes. Prior to COVID-19, the

country had experienced a period of peak economic prosperity[7], which was interrupted by the

outbreak, leading to unprecedented unemployment rates (reaching nearly 30% of the work-force

in April 2020) and the collapse of several sectors such as aviation, tourism, and culture[8,9]. The

outbreak period was characterized with acute uncertainty and increase in anxiety, regarding both

the health and socioeconomic effects of the pandemic[10].

Over the past years, several studies demonstrated the potential to detect brain plasticity using T1-

weighted magnetic resonance imaging (MRI)[11–13]. The current work was initiated as a reaction to

the outbreak of COVID-19 in Israel, aiming to study the structural brain plasticity in the general

population following a real-life event of global scale. For this purpose, we examined $n = 50$ test

group participants that were scanned with T1-weighted MRI prior to the outbreak and returned

for a follow-up scan after the lockdown period (see methods for a detailed timeline of post

COVID-19 follow-up scans). The structural changes of the study group (before versus after the

outbreak) were compared to those of $n = 50$ control participants who were scanned twice before

the COVID-19 outbreak. The unique circumstances imposed due to the COVID-19 lockdown

created rare settings for a natural experiment to examine the effect of a real-world intense event

on brain plasticity.

All participants were healthy, without a history of neurological or psychiatric disorders, did not

show COVID-19 symptoms, and were not diagnosed carrying the virus (see the methods section

for further demographic information). The hypotheses and general design of the current study

were pre-registered prior to the completion of data collection and were based on a small

independent pilot sample with $N = 16$ participants ($n = 8$ participants in each group). The data

and analysis codes are openly shared online (project page: https://osf.io/wu37z/; preregistration:

https://osf.io/k6xhn/).

Results

Prior to their follow-up MRI scan session, we asked participants of the post-lockdown test group

to fill in a short questionnaire regarding the lockdown period (see methods). Of the participants

who agreed to reply, 79.6% reported they did not leave their home for non-essential needs,

79.6% met no more than 3 people, 44.9% did not meet with their parents at all, 38.8% indicated

an increased feeling of anxiety following the lockdown, 34.7% anticipated that their future

behavior will change after the lockdown, 46.8% reported they were concerned about their

personal future well-being, 42.9% indicated that their employment status was reduced to part-

time or unemployment/furlough. In an exploratory factor analysis (EFA; using Varimax rotation,

see methods), we identified two main factors, explaining together 54.0% of the variance. The

first factor was highly loaded with questionnaire items that described increased social isolation,

and the second was mainly related to increased feelings of anxiety (Figure 1).

Figure 1. Exploratory factor analysis (EFA) of COVID-19 questionnaire.
Exploratory factor analysis of the responses to the questionnaire revealed two main themes characterized the participants. **a.** The first factor ('Social isolation') strongly related to the item indicating meeting no more than 3 people, as well as to other two related items of avoiding meeting parents and staying at home during lockdown. An increased feeling of anxiety dominated the second factor, along with changing future behavior and concerns regarding the long-term effects. **b.** Visualization of participants dispersion across the two factors and their categorization into binary anxiety and isolation groups. Points represent unique scores. Scores' frequency is represented by point size and participants' indices around their corresponding data points.

Based on our pilot study results and previous studies of stress-related morphological brain changes[14–17], we hypothesized that the focus of volumetric changes will be observed mainly in the amygdalae. T1-weighted anatomical MRI scans were used as input for deformation and surface-based morphometry (SBM) analysis using the CAT12 toolbox (http://www.neuro.uni-jena.de/cat/, University of Jena) for SPM12 (http://www.fil.ion.ucl.ac.uk/spm/software/spm12/, Wellcome Trust Centre for Neuroimaging). The brain was segmented to 58 regions based on the cortical and subcortical nuclei classifications of the Hammers atlas (Hammers et al., 2003). Following surface reconstruction, each participant's individual gray matter volume was estimated for each of the 58 anatomically defined regions of interest (ROIs). This procedure accounted for the longitudinal nature of the data, performing the analysis on both scans

simultaneously. To avoid voxel-based multiple comparisons, we performed a region-based

analysis (following surface projection to the Hammer atlas) and corrected for multiple

comparisons using the Benjamini-Hochberg correction[18] to control for false discovery rate (FDR;

using $p < 0.05$ following correction threshold). Validation of the pipeline was performed using

simulated data and by comparing the results with other software (see methods).

Using a linear mixed model, we examined volumetric changes, testing for regions with stronger

changes for the test group compared to the control group. Examining the interaction effect of

session (baseline versus follow-up scans) and experimental group (test versus control) revealed

ten anatomical brain regions (composed of bilateral five unique regions in both hemispheres) in

which volumetric increases were observed uniquely for the test group (Table 1 and Figure 2).

Most prominently, as we expected and pre-registered, we found a robust effect in the bilateral

amygdalae. We also observed a significant effect bilaterally in the putamen, and in three

anatomical regions within the ventral anterior temporal cortex adjacent to each other, namely in

the medial part of the anterior temporal lobe, the fusiform gyrus, and the parahippocampal gyrus.

To examine the spatial distribution within significant ROIs, we performed an additional post-hoc

voxel-based analysis, which allowed us to visualize the changes within the significant ROIs

(Figure 2a). Examining the post-hoc voxel-based results revealed that volumetric changes

occurred throughout the entire surface of bilateral amygdalae, while in the putamen the effects

occurred mainly in the dorsal area. In the ventral anterior temporal cortices, large connected

clusters of volumetric change spanned throughout the three adjacent temporal ROIs, thus

suggesting that the three ROIs shared a similar origin. To ensure that the reported effects

originated from volumetric changes in the test group following the COVID-19 outbreak and its

related lockdown period, we tested for ROIs where the significant interaction effect was

accompanied by a significant effect for the test group but not for the control group, and was

consistent beyond baseline scans effect or measurement protocol (see methods).

Table 1. Surface based morphology analysis results

Region	Hemi-sphere	Interaction estimate (95% CI)	Interaction p (FDR adj.)	Test group session estimate (95% CI) [a]	Test group session p (FDR adj.)	Change from baseline M (SE) [b]	
						Test	Control
Amygdala	Left	0.09 [0.05, 0.13]	$2.4E^{-5}$ (0.001)	0.08 [0.05, 0.11]	$9.8E^{-6}$ ($2.1E^{-4}$)	4.92% (1.06%)	-0.56% (0.6%)
	Right	0.08 [0.03, 0.13]	0.003 (0.030)	0.08 [0.05, 0.11]	$1.6E^{-5}$ ($2.3E^{-4}$)	4.47% (1.07%)	0.07% (0.86%)
Putamen	Left	0.19 [0.09, 0.29]	$4.1E^{-4}$ (0.006)	0.13 [0.06, 0.2]	$4.0E^{-4}$ (0.002)	3.31% (0.83%)	-0.5% (0.7%)
	Right	0.17 [0.08, 0.26]	$2.4E^{-4}$ (0.005)	0.14 [0.08, 0.2]	$1.1E^{-5}$ ($2.1E^{-4}$)	3.31% (0.67%)	-0.01% (0.68%)
Anterior temporal lobe (medial part)	Left	0.25 [0.12, 0.38]	$1.8E^{-4}$ (0.005)	0.15 [0.07, 0.23]	$4.7E^{-4}$ (0.003)	2.82% (0.75%)	-0.7% (0.76%)
	Right	0.21 [0.07, 0.35]	0.004 (0.030)	0.15 [0.05, 0.25]	0.004 (0.023)	2.93% (1.07%)	-0.37% (0.7%)
Parahippocampal gyrus	Left	0.09 [0.03, 0.15]	0.006 (0.035)	0.04 [0, 0.08]	0.029 (0.085)	1.22% (0.55%)	-0.5% (0.57%)
	Right	0.11 [0.04, 0.18]	0.003 (0.030)	0.08 [0.03, 0.13]	0.002 (0.009)	2.05% (0.61%)	-0.08% (0.54%)
Fusiform gyrus	Left	0.08 [0.03, 0.13]	0.007 (0.036)	0.06 [0.03, 0.09]	$3.8E^{-4}$ (0.003)	1.78% (0.53%)	-0.4% (0.56%)
	Right	0.11 [0.04, 0.18]	0.002 (0.022)	0.05 [0, 0.1]	0.044 (0.111)	1.36% (0.64%)	-0.72% (0.57%)

139 [a] Session estimate examined the effect of baseline versus follow-up scan in the post-lockdown
140 test group. This parameter was used to validate that the interaction effect observed between the
141 group stemmed from a robust effect in the test group (see methods).
142 [b] Volumetric change normalized to baseline scan (difference/ baseline).

144
Figure 2. Volumetric changes results.
An interaction effect for time (baseline versus follow-up scan) and group (test versus control)
was evaluated on segmented surfaces in an SBM analysis. Significant interaction effects were
observed bilaterally in the amygdala and putamen ROIs, as well as in three ventral temporal
cortical ROIs. **a.** To examine spatial patterns within the identified ROIs, a post-hoc voxel-based
analysis was conducted within each ROI mask. Light red contours represent segmentation
borders of the ROIs. **b.** Individual distribution of the results in the control group (light colors)
and test group (dark colors). For better visualization, units were normalized to baseline
(difference/baseline) and presented in percentage units. Box-plot center, hinges, and whiskers
represent the median, quartiles, and ‾   from the hinges, respectively. A notch of
‾ represent an estimated 95% confidence interval for comparing medians. Dots represent
individual participants. Abbreviated ROI names: AntMedTeLo = anterior temporal lobe (medial
part); FusGy = fusiform gyrus, ParHipGy = Parahippocampal gyrus.
158
To evaluate and control for the effect of time between scans and time from lockdown, we

included in the model two additional covariates - the time between scans (TBS; which was

generally longer for the test group) and time following lockdown (TFL; see methods for more

details). The two covariates were not correlated with each other in our test group sample ($r = -$

0.106, $t(48) = -0.74$, $p = 0.463$). Our reported regions demonstrated significant volumetric

change above and beyond these covariates. After FDR correction, no region showed an effect of

TBS. However, we did find a negative effect of TFL in the two amygdalae ROIs and the left

fusiform gyrus, suggesting that the volumetric changes in these regions moderated as time

following lockdown elapsed. Based on these results, we estimated the time to decay as the

estimated number of days from lockdown until volumetric changes returned to normal levels,

similar to those of the control group (left amygdala: $\beta_{TFL} = -0.41$, $t(47) = -3.1$, $p = 0.003$, $p_{adj.} =$

0.048, time to decay = 95 days; right amygdala: $\beta_{TFL} = -0.54$, $t(47) = -4.38$, $p = 6.7E^{-5}$, $p_{adj.} =$

0.002, time to decay = 83 days; left fusiform gyrus: $\beta_{TFL} = -0.54$, $t(47) = -4.44$, $p = 5.5E^{-5}$, $p_{adj.} =$

0.002, time to decay = 82 days; Figure 3).

Figure 3. Time following lockdown effect on volumetric changes.

The time from lockdown relief until the follow-up scan session (TFL) was added as an addition
covariate to the model, revealing significant effect in the two amygdalae and left fusiform gyrus.
Points represent individual participants in the post-lockdown test, *p*-values were FDR adjusted
for multiple comparisons. Abbreviated ROIs: AntMedTeLo = anterior temporal lobe (medial
part); FusGy = fusiform gyrus, ParHipGy = Parahippocampal gyrus.
180

In addition to the volumetric change effect reported here, we also performed an exploratory

analysis aimed to detect pattern of correlation change between all tested ROIs, and how do these

changes differ between the two groups (see supplementary results). We also performed

additional exploratory analyses to examine the association of volumetric changes and the

reported experience during lockdown. We found no strong association between the two, as

reported in the supplementary results.

Discussion

Our study demonstrates that volumetric change patterns in the brain occurred following the

COVID-19 initial outbreak period and restrictions in a sample of healthy participants, who were

not somatically affected by the pandemic. While previous studies demonstrated brain plasticity

using T1-weighted MRI following planned interventions[11–13], the current work outstands in its

unique demonstration of stark structural brain plasticity following a major real-life event.

Our findings show neural changes that were not caused directly due to COVID-19 infection, but

rather related to the societal effect. We show volumetric increase in gray matter in the

amygdalae, putamen and ventral anterior temporal cortices. The changes in the amygdalae

showed a temporal-dependent effect, related to the time elapsed from lockdown but not the

duration from the baseline scan. It should be noted that although lockdown restrictions had

initially reduced infection rates in Israel, just one month after the lockdown was lifted, the

number of infected cases started to rise again and reached higher number of active infected cases

by the end of data collection, compared with the peak numbers during the actual lockdown

period (approximately 2,000 daily new cases by the end of July versus under 750 new daily cases

during the peak of the lockdown period in April[19], see methods and Figure 4). This suggests that

the effects observed in the current study are less likely to be attributed to the concrete health

risks of COVID-19, but rather to the first wave of the outbreak, characterized with perceived

uncertainty and substantial unexpected changes in everyday life.

The current study was in many aspects unplanned; therefore, we are left with only partial

answers as to which specific components of the COVID-19 outbreak led to the neural changes

observed in the healthy participants that took part in our study. The involvement of the amygdala

may suggest that stress and anxiety could be the source of the observed phenomenon, due to its

well-recorded functional and structural associations[14–17,20–22]. Nevertheless, it is hard to draw

clear conclusions as many aspects of life have changed in this time period, and could have

potentially affected different regions in the brain – from limiting social interactions, increased

financial stress, changes in physical activity, work routine, and many more. The limited

behavioral data collected in the current study did not provide a strong connection to the imaging

results, and thus future work could try to better address the complex brain-behavioral

associations in this real-life experience. Nonetheless, our findings show that healthy young

adults, with no records of mental health issues, were deeply affected by the outbreak of COVID-

19. We suggest that policy makers take into consideration the impact of their actions on the

general well-being of the population they seek to help, alongside the efficacy of disease

prevention.

221   **Methods**

Data and codes availability

Our sample size, hypotheses and analyses plan were pre-registered on the Open Science

Framework (OSF), soon after data collection began, but prior to completion of the data collection

and data analysis (project page: https://osf.io/wu37z/; preregistration: https://osf.io/k6xhn). All

behavioral data, processed imaging data, and analysis codes are shared on the OSF project page.

Uncorrected and small-volume corrected statistical maps of the voxel-based results described in

the current work are available at https://neurovault.org/collections/8591/.

Participants

The study included two groups: A test group scanned before and after COVID-19 lockdown, and

a control group, scanned twice before COVID-19 lockdown. All participants had no background

of neurological disorders, did not show symptoms for COVID-19 and were not diagnosed as

carriers of the virus. The study was approved by the ethics committee of Tel Aviv University and

institutional review board (IRB) at the Sheba Tel-Hashomer medical center. Since the IRB

protocol allowed us to scan the participants several times over a long period of time, we were

able to collect the data from participants who were scanned prior to COVID-19 outbreak and

invite them back for a follow-up scan as part of the longitudinal study they have agreed to take

part in. Participants received monetary compensation for their time and gave their informed

consent to take part in a longitudinal experiment aimed to examine brain plasticity across several

sessions, which was initially not directly related to COVID-19 outbreak.

The test group comprised of $n = 50$ participants who were scanned before and after COVID-19

lockdown (Δ Time between scans: $M = 309.3$, $SD = 207.5$, range $= 67 - 1460$ days; Age: $M =$

30.1, $SD = 6.65$, range $= 21 - 48$; Females: n $= 20$, prop. $= 40\%$). The lockdown period began on

March 25[th] and was gradually relieved throughout late April. We mark here May 1[st] as the

lockdown relief date, as on this day an issued 100-meters movement limit for non-essential needs

was lifted. The test group data collection started as soon as lockdown relief took place, for a

period of approximately 3 months, until the end of July, 2020 (Δ Time from lockdown relief: M

$= 57$ days, $SD = 24.62$, range $= 9 - 89$ days; see Figure 4 for the study timeline).

250
Figure 4. Study timeline and outbreak data
On February 21[st], 2020, the first COVID-19 case in Israel was recorded. Daily new cases were
smoothed using 7-days moving average. Data were retrieved and modified based on the Israeli
Ministry of Health reports[19,23]. A lockdown was issued on March 25[th], which was gradually
released until the removal of the 100-meter restriction on May 1[st], marking lockdown onset and
relief, respectively (shorter vertical dashed line). MRI data of the test group were collected
between May 10[th] to July 29[th] (longer vertical dashed line). Red bars on top represent the number
of participants scanned for the study each day.

As a control group, we used the data of $n = 50$ participants who were scanned twice using a

similar protocol before COVID-19 lockdown (Δ Time between scans: $M = 126.7$, $SD = 190.4$,

range = 21 - 886 days; Age: M = 27.3, $SD = 5.63$, range = 19 - 42; Females: n = 23, prop. =

46%).

The minimal sample size was determined and pre-registered (https://osf.io/uktsn), based on a

80% power analysis conducted using R 'pwr' package[24], on a pilot study with $N = 16$

participants ($n = 8$ in each group). We decided to collect a minimum of $n = 37$ participants which

should provide 80% to detect the group and session interaction effect with α = .05, within both

the left and right amygdala, under the assumption that it would be difficult to complete the

sample due to COVID-19 limitations. Eventually, thanks to further relief in COVID-19

restrictions, we were able to extend the sample size to $n = 50$ in each group.

The results remain generally consistent, even when the data included only the first $n = 37$

participant; demonstrating significant effects in the bilateral amygdalae, putamen,

parahippocampal gyrus and the left anterior temporal lobe. In this smaller sample, we did not

find significant effects in the right anterior temporal lobe, nor in the fusiform gyrus. We also

found volumetric increase effects that were not identified using the full sample in the left nucleus

accumbens, left cuneus, and left insula.

Imaging data

Acquisition protocol. Imaging data were acquired using a 3T Siemens Prisma scanner, with a

64-channel head coil. For the structural data, T1w high resolution (1-mm^3) whole brain images

were acquired with a magnetization prepared rapid gradient echo (MPRAGE) pulse sequence

with repetition time (TR) of 2.53s, echo time (TE) of 2.88ms, flip angle (FA) = 7°, field-of-view

(FOV) = 224 × 224 × 208 mm, resolution = 1 × 1 × 1. Since for this project we combined data

from different labs, some of the participants had their scans aligned to the anterior commissure -

posterior commissure (AC-PC) line, while others were scanned with a 30° angle (50% of test

group participants and 80% of control group participants).

Some participants were also scanned with diffusion-weighted echo-planar imaging (DW EPI)

sequence and some with functional gradient-echo EPI (GE EPI) in a resting state scan. The

analyses of these scans are beyond the scope of the current study.

Data processing and analysis. The T1-weighted MPRAGE anatomical scans were used for a

surface-based morphometry (SBM) analysis. From the images we estimated the pial and inner

surfaces of the cortex and projected those into the Hammers atlas (Hammers et al., 2003). Data

were preprocessed in SPM12 (http://www.fil.ion.ucl.ac.uk/spm/software/spm12/, Wellcome

Trust Centre for Neuroimaging) and SPM based CAT12 (Computational Anatomy Toolbox 12;

http://www.neuro.uni-jena.de/cat/, University of Jena) extension. We deployed the CAT12

surface-processing pipeline, which includes skull striping, a denoising filter[25] projection-based

thickness estimation[26], partial volume correction, and spatial normalization to MNI space. Prior

to the preregistration of the protocol, we validated the reliability of our surface-based image

processing pipeline, by simulating artificial activity in the amygdala, and using FreeSurfer based

analysis of cortical regions (See supplementary materials for further details).

Surface-based volumetric data of cortical and subcortical regions were segmented based on the

Hammers Atlas, segmenting the volumetric data into 58 anatomically defined regions. Ten

additional ROIs of non-gray matter or non-cerebral structures (ventricles, white matter, brain-

stem and cerebellum ROIs) were excluded from statistical analyses. To evaluate the effect of

lockdown on volumetric imaging data we ran a mixed linear model on the data within each one

of the 58 anatomical regions, examining the effect of session (baseline versus follow-up scan)

and group interaction (test versus control), controlling for time between scans (TBS) and time
following lockdown (TFL) covariates. Both covariates were mean centered before they were
added to the model. To include TFL covariate in the same model with control group participants,
for which TFL was irrelevant, control groups' participants were assigned with the mean TFL;
thus, their data was not used to evaluate the effect of TFL.
To identify our regions of interest we only included regions that showed both a significant
interaction effect (i.e. the volumetric difference between the two sessions was significantly
different for the test and control group), and a significant session effect within the test group (i.e.
a significant difference between the two sessions for the test group). Results were corrected for
multiple comparisons using the false discovery rate (FDR) correction[18], based on the number of
brain regions tested. Following the analysis pipeline, we identified ten significant ROIs: bilateral
amygdalae, putamen, parahippocampal gyrus, the medial part of the anterior temporal lobe, and
the fusiform gyrus.
An additional ROI of the right inferior and middle temporal gyri showed a significant interaction
effect (interaction estimate = 0.17, 95% CI = [0.05, 0.29], $p = 6.1E^{-3}$, $p_{adj.} = 0.035$), however
examining the test group separately, we could not identify a significant session effect (session
estimate = 0.04, 95% CI = [-0.02, 0.10], $p = 0.259$, $p_{adj.} = 0.424$). Therefore, it is harder to
interpret that this interaction effect stemmed from the test group. Thus, we did not include this
ROI as one of our significant ROIs. A less robust session effect within the test group was also
observed for the left parahippocampal gyrus (session estimate = 0.04 [0.00, 0.08], $p = 0.029$, $p_{adj.}$
= 0.085) and right fusiform gyrus ROI (session estimate = 0.05 [0.00, 0.10], $p = 0.044$, $p_{adj.} =$
0.111), however as these ROIs demonstrated strong interaction effects and their session effects
were significant before FDR correction, we decided to report them together with the other
significant regions. A similar procedure was used in the pre-registration (i.e. including only
regions that showed a significant interaction and an effect in the test group); however, in the
analysis of the pilot for the pre-registration, we used uncorrected results due to the small sample
size.
To examine the spatial distribution of our effect within significant ROIs, that were identified
with the SBM analysis, we performed an additional post-hoc voxel-based analysis. We projected
the data on a voxel-based map, effectively examining which voxels demonstrated an interaction
effect. Then, we used anatomical masks of ROIs which were found to be significant in the SBM

analysis, to visualize the results within these regions, similarly to a small-volume correction
analysis (Figure 2a).

Behavioral data

Data collection. To evaluate participants' experience in the peak days of the COVID-19
outbreak, we asked them to think back on their experience during this time and fill out a 7-items
questionnaire regarding their experience of the COVID-19 lockdown (see Table 2 for a
description of the items). The questionnaires were filled out after the initiation of the study, when
the lockdown's stringent 100-meters limitation was lifted, thus the results represents the
participants' recalled experience of the lockdown. Most participants filled out the questionnaire
on the day of the post-lockdown scan session, some filled it a few days before their second
scanning session. A total of $n = 76$ participants filled out the COVID-19 questionnaire and
comprised the potential pool of test group participants for the current study, out of which $n = 50$
were sampled and scanned. One participant was scanned but did not complete the questionnaire,
therefore this participant's behavioral data were not used and analyses of the questionnaire were
based on $n = 49$ valid participants.

Table 2. COVID-19 lockdown questionnaire

Question	Possible Answers (prop.)	Binary outcome (prop.)
1. Did you stay home during the lockdown, except for essential needs / did not leave at all?	0 - no (20.4%) 1 - yes (79.6%)	0 - no (20.4%) 1 - yes (79.6%)
2. Did the lockdown increase your feeling of anxiety?	0 - no (61.2%) 1 - yes (38.8%)	0 - no (61.2%) 1 - yes (38.8%)
3. With how many people did you meet during the lockdown (including people you are living with at home)?	0 – none (0%) 1 - up to three people (57.1%) 2- up to five people (22.4%) 3 - up to ten people (20.4%)	0 - more than three (20.4%) 1 - up to three (79.6%)
4. Do you think your behavior will change following the lockdown?	0 - no (65.3%) 1 - yes (34.7%)	0 - no (65.3%) 1 - yes (34.7%)
5. How did your meeting with your parents' routine look like during the lockdown?	0 - same as before the lockdown (34.7%) 1 - with precaution measurements: distancing, mask, etc. (20.4%) 2 - did not meet at all (44.9%)	0 - as before or with precautions (55.1%) 1 - did not meet at all (44.9%)
6. What was your employment status during the lockdown?	0 - same as before lockdown (28.6%) 1 - full time working from home (28.6%) 2 - part time working from home (8.2%) 3 - Furlough / unemployed (34.7%)	0 - unemployed / part time (42.9%) 1 - same as before / full time from home (57.1%)
7. How concerned are you with the long-term effect of the lockdown, regarding yourself?	Ordinal scale 1 – not at all (28.6%) 2 (24.5%) 3 (30.6%) 4 (14.3%) 5 – very concerned (2%)	0 - low, score 1-2 (53.1%) 1 - moderate-high, score 3-5 (46.9%)

Exploratory factor analysis (EFA). Responses to the lockdown questionnaire were coded into binary responses, based on the sample median, splitting the sample into relatively similar sized groups for each item (Table 2). To identify the main themes in the questionnaire, which could be correlated with the imaging data, we performed an exploratory factor analysis (EFA) on the binarized data, using the "psych" R package[27].

Since EFA require large number of participants, we used the data from all available $n = 75$

participants who completed the questionnaire. Kaiser-Meyer-Olkin (KMO) factor adequacy test

revealed that the 'employment' item had a very low measure of sampling adequacy (MSA;

employment item's MSA = 0.26; which is far below the suggested minimal MSA of 0.5[28]) for

EFA. The 'employment' was also not loaded to any of the factors; therefore, we removed it from

the final EFA model. Overall, even after removing low KMO item, our data was found to be

weakly appropriate for factor analysis, with overall MSA = 0.48. Thus, considering the small

sample size and low fit of the data to EFA, the results of the analysis should be interpreted with

caution.

We performed polychoric correlations based EFA, which is suitable for binary variables, with

two factors and Varimax rotation, assuming orthogonality between the factors. Our selection of

number of factors was based on visual inspection of scree plot of eigenvalues, as well as by

comparing actual data to simulations of random data matrices. Using Oblimin rotation, which

allows for correlation between the factors, revealed very low correlation ($r = 0.08$), suggesting

that Varimax rotation was an appropriate choice for our model.

To examine the association of the behavioral data with the volumetric changes, while

maintaining relatively limited number of multiple comparisons, we used the two factors for these

analyses instead of each of the items. These two factors' scores for each participant were

extracted and correlated with the change in gray matter volumetric data in our regions of interest

(see supplementary materials for results of this exploratory analysis).

References

1. Vinceti, M. *et al.* Lockdown timing and efficacy in controlling COVID-19 using mobile phone tracking. *EClinicalMedicine* (2020). doi:10.1016/j.eclinm.2020.100457

2. Fernandes, N. Economic effects of coronavirus outbreak (COVID-19) on the world economy Nuno Fernandes Full Professor of Finance IESE Business School Spain. *SSRN Electron. Journal, ISSN 1556-5068, Elsevier BV,* 0–29 (2020).

3. Zhang, D., Hu, M. & Ji, Q. Financial markets under the global pandemic of COVID-19. *Financ. Res. Lett.* 101528 (2020). doi:10.1016/j.frl.2020.101528

4. Taylor, S. *et al.* COVID stress syndrome: Concept, structure, and correlates. *Depress. Anxiety* 1–9 (2020). doi:10.1002/da.23071

5. Gruber, J. *et al.* Mental health and clinical psychological science in the time of COVID-19: Challenges, opportunities, and a call to action. *Am. Psychol.* (2020). doi:https://doi.org/10.1037/amp0000707

6. Qiu, J. *et al.* A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: Implications and policy recommendations. *Gen. Psychiatry* **33**, 1–4 (2020).

7. Bank of Israel Research Department. The Decline in Unemployment in Israel by International Comparison. in *Bank of Israel Annual Report 2019* (2020).

8. Bank of Israel Research Department. *The unemployment rate and its definition during the corona period*. (2020).

9. Bank of Israel Research Department. *Changes in credit card purchases as a result of the corona crisis*. (2020).

10. Tzur Bitan, D. *et al.* Fear of COVID-19 scale: Psychometric characteristics, reliability and validity in the Israeli population. *Psychiatry Res.* **289**, 113100 (2020).

11. Maguire, E. A. *et al.* Navigation-related structural change in the hippocampi of taxi drivers. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 4398–4403 (2000).

12. Jung, W. H. *et al.* Exploring the brains of Baduk (Go) experts: Gray matter morphometry, resting-state functional connectivity, and graph theoretical analysis. *Front. Hum. Neurosci.* **7**, 633 (2013).

13. Draganski, B. *et al.* Changes in grey matter induced by training. *Nature* **427**, 311–312 (2004).

14.   Ganzel, B. L., Kim, P., Glover, G. H. & Temple, E. Resilience after 9/11: Multimodal
neuroimaging evidence for stress-related change in the healthy adult brain. *Neuroimage*
40, 788–795 (2008).

15.   Hölzel, B. K. *et al.* Stress reduction correlates with structural changes in the amygdala.
Soc. Cogn. Affect. Neurosci. **5**, 11–17 (2009).

16.   Rogers, M. A. *et al.* Smaller amygdala volume and reduced anterior cingulate gray matter
density associated with history of post-traumatic stress disorder. *Psychiatry Res. -*
Neuroimaging **174**, 210–216 (2009).

17.   Schienle, A., Ebner, F. & Schäfer, A. Localized gray matter volume abnormalities in
generalized anxiety disorder. *Eur. Arch. Psychiatry Clin. Neurosci.* **261**, 303–307 (2011).

18.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate : A Practical and
Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).

19.   Israel Ministry of Health. Coronavirus in israel - Overview (in Hebrew). (2020). Available
at: https://datadashboard.health.gov.il/. (Accessed: 31st August 2020)

20.   Stevens, J. S. *et al.* Amygdala Reactivity and Anterior Cingulate Habituation Predict
Posttraumatic Stress Disorder Symptom Maintenance After Acute Civilian Trauma. *Biol.*
Psychiatry **81**, 1023–1029 (2017).

21.   Bryant, R. A. *et al.* Enhanced amygdala and medial prefrontal activation during
nonconscious processing of fear in posttraumatic stress disorder: An fMRI study. *Hum.*
Brain Mapp. **29**, 517–523 (2008).

22.   Mochcovitch, M. D., Da Rocha Freire, R. C., Garcia, R. F. & Nardi, A. E. A systematic
review of fMRI studies in generalized anxiety disorder: Evaluating its neural and
cognitive basis. *J. Affect. Disord.* **167**, 336–342 (2014).

23.   Max, R., Ritchie, H., Ortiz-Ospina, E. & Joe, H. Coronavirus Pandemic (COVID-19). *Our*
World in Data (2020). Available at: https://ourworldindata.org/coronavirus. (Accessed:
1st September 2020)

24.   Champely, S. *et al.* Package 'pwr': Basic Functions for Power Analysis. *CRAN Repos.*
(2018).

25.   Manjón, J. V., Coupé, P., Martí-Bonmatí, L., Collins, D. L. & Robles, M. Adaptive non-
local means denoising of MR images with spatially varying noise levels. *J. Magn. Reson.*
Imaging **31**, 192–203 (2010).

26.   Dahnke, R., Yotter, R. A. & Gaser, C. Cortical thickness and central surface estimation.

Neuroimage **65**, 336–348 (2013).

27.   Revelle, W. Psych: Procedures for Psychological, Psychometric, and Personality

Researche. R package version 2.0.8 (2020).

28.   Kaiser, H. F. An index of factorial simplicity. *Psychometrika* **39**, 31–36 (1974).

Acknowledgements

This work was supported by the Israeli Science Foundation granted to Yaniv Assaf (ISF

1314/15), Tom Schonberg (2004/15), and Ido Tavor (ISF 1603/18). Tom Salomon was supported

by the Nehemia Levtzion fellowship and the Fields-Rayant Minducate Learning Innovation

Research Center.

Author contributions

T.Sa. wrote the manuscript with Y.A, assisted with the study design and analyzed the data. Data

was collected by A.C., R.B.-N., R.G, S.O, G.R., D.R, and A.S. Free-surfer analysis was made by

G.B-Z. N.T performed VBM validation analysis. D.B. designed the scanning protocols. G.T.

provided support and medical supervision. I.T., T.Sc. and Y.A. conceived the study, wrote the

manuscript and supervised the study. Y.A performed the preprocessing and analysis of imaging

data. All Author contributed intellectually and reviewed the manuscript.

Competing interests

The authors declare no competing financial interests.

Supplementary materials

Validation using different statistical modeling

In our work, we found volumetric change effect following COVID-19 outbreak and lockdown.

Our principle statistical analysis was based on the differences in volume values of the baseline

and follow-up scans of each participant. To easily compare the effect in each brain region, we

report and illustrate the change effect in units of percent change: $\frac{(V_{follow-up} - V_{baseline})}{V_{baseline}}$.

Supplementary Figure 1, provides a visualization of the results presented in Figures 2b and 3,

using the original units, without normalizing for baseline scan unites $(V_{follow-up} - V_{baseline})$.

476
Supplementary Figure 1. Volumetric change effects not normalized to baseline.
Visualization of the reported volumetric change effects, without normalizing the data to baseline
volume values. **a.** distribution of volumetric change values across the 10 signifiacant ROIs, as
presented in Figure 2b. **b.** The association of volumetric change effect and time from lockdown
release, as presented in Figure 3.

To further corroborate the validity of our results, we examined the volumetric change effect
through several additional confirmatory perspectives, which could help exclude potential
confounds. First, we examined whether the difference effect might be related to baseline scan

values. To make sure that the volumetric change effect that we observed in the test group is
robust above and beyond initial baseline scan, we examined a linear model with the follow-up
scan volume as our dependent variable and the baseline scan as an additional covariate. The
results showed consistent interaction effect. In all 10 significant ROIs, even when baseline scans
explained a significant portion of the variance in follow-up scans, the difference effect between
the two groups remained significant above and beyond the effect of baseline scan
(Supplementary figure 2).

Supplementary Figure 2. Volumetric change and baseline scan
Volumetric change (difference between follow-up and baseline scans, relative to baseline) of the
test group (dark grey) and control group (light grey), plotted against the volumetric values
measured at baseline. Small dots represent individual participants, large dots represent the groups
means. A significant interaction effect was observed above and beyond the variability explained
by baseline scan values, as test group participants overall showed more robust volumetric
changes than the control participants.

In another model, we also included the gender and the scan angle (AC-PC line versus 30° angle
from the AC-PC) as additional independent variables. Both gender and scan angle had no
significant contribution beyond other independent variables ($p_{adj.} > 0.05$ in all reported ROIs).
Adding gender had no effect on the reported results. However, the scan angle was correlated
with the group and the TFL variables, such that a larger proportion of control-group participants
were scanned with the 30°angle (80% versus 50% in the test group); and in the test group,

participants who were scanned with the 30° angle were mostly scanned in the later days of data
collection (TFL and scan-angle correlation: $r = 0.615$, $t(48) = 5.41$, $p = 2.0E^{-6}$). The addition of
the scan angle to the model therefore had an impact on the significance of other effects: (1) all
reported TFL effects became non-significant after FDR-correction; and (2) while no additional
regions became significant for the group effect, the FDR-corrected p-value dropped just slightly
below the statistical significance threshold for the right amygdala ($p_{adj.} = 0.058$), right anterior
medial temporal lobe ($p_{adj.} = 0.058$), right parahipocampal gyrus ($p_{adj.} = 0.058$), left
parahipocampal gyrus ($p_{adj.} = 0.074$), right fusiform gyrus ($p_{adj.} = 0.058$) and left fusiform gyrus
($p_{adj.} = 0.074$). The other regions remained significant (left amygdala $p_{adj.} = 0.003$, left anterior
medial temporal lobe $p_{adj.} = 0.007$, left putamen $p_{adj.} = 0.010$, and right putamen $p_{adj.} = 0.007$).
These results suggest that the scan angle had a slight impact on the significance of the effects,
probably due to poor balance of the condition across groups and time. Although some fail to
exceed the stringent statistical threshold after FDR by a small margin, overall the results remain
fairly consistent.
We further validated that our choice of measurement to analyze (difference of follow-up –
baseline scans) had no significant impact on our conclusions. In another analysis, we compared
the log-ratio between scans as our dependent variable, wherein log(follow-up/baseline) under the
null hypothesis is expected to be 0. Using the same analysis process with log-ratio scores as the
dependent variable resulted in the same 10 significant ROIs. Thus, our additional analysis
suggested that our reported results are robust to potential biases due to baseline scans and choice
of statistical analysis procedure.
Imaging pipeline validation
To validate the imaging processing protocol we used two approaches, before data collection was
completed (at the time of the pre-registration finalization). As many surface-based software
focus on analysis of cortical surfaces, rather than subcortical regions, we aimed to validate that
our SBM protocol using CAT12 could reliably identify subcortical morphological changes, such
as the ones we observed in the current study within the amygdala and putamen. To test the
detection capabilities of our protocol, we generated simulated data with volumetric changes in
the amygdala and ran the CAT12vpipeline on the simulated data. A volumetric increase in the
amygdala was simulated using a hand-drawn polygon mask, surrounding the left amygdala on
the original T1w images of 10 participants. Within this polygon 3D mask, the signal intensity

was artificially changed. Following this procedure, both the original and modified T1w images

underwent the same CAT12 pipeline. We were able to identify the simulated volumetric changes

within the subcortical amygdala nuclei (see supplementary Figure 3).

Supplementary Figure 3. Data simulation
Sub-cortical changes in the left amygdala were simulated in 10 participants. The SBM pipeline
applied in CAT12 identified the simulated change effect. Data were projected from surfaces back
to a voxel-based map for visualization in the current figure.

In an additional validation procedure, we reanalyzed our results with an additional analysis

pipeline. Raw T1-weighted maps were preprocessed using FreeSurfer. Using this alternative

analysis pipeline, we found similar results, including robust effects on the temporal cortical

regions in addition to other cortical regions. It should be noted that the FreeSurfer pipeline can

analyze only cortical surface, thus the analysis did not include subcortical regions of the

amygdala and putamen.

The results of both validations indicated our analysis pipeline could reliably identify regional

volumetric estimations. Using CAT12 provided an advantage by performing a longitudinal

analysis of subcortical regions, including the amygdala which was pre-hypothesized and of great

importance in the current work.

Exploratory factor analyses: association with volumetric changes

As an exploratory analysis, we examined whether the volumetric brain changes were associated

with the psychological constructs identified in our EFA, based on participants' self-reports. In a

previous version of this manuscript, we used principal component analysis (PCA) to identify our

factors of interest, however this procedure is less appropriate than EFA, and is therefore not

reported here. However, the results using PCA were fairly similar to the ones we found with

EFA   (previous   version   is   available   at:

https://www.biorxiv.org/content/10.1101/2020.09.08.285007v2).

We used two linear models to explain the variability in each of the factors, using the volumetric

changes as our model features.

Overall, neither one of the factors was well associated with the volumetric changes (Factor 1

model: $R^2 = 0.20$, $F(10,38) = 0.92$, $p = 0.522$; Factor 2 model: $R^2 = 0.22$, $F(10,38) = 1.10$, $p =$

0.383). Examining the contribution of individual ROIs within the models (measured as the

significance of the β estimates), did not reveal a significant association with the factors for any

one of the ROIs ($p_{adj.} > 0.05$; FDR correction by the number of features in the model). Adding

the two factors as covariates to the linear models examining the volumetric change effect, also

revealed no significant contribution of the factor to the models.

Thus, in our work we could not identify a clear association between the behavioral data and

volumetric changes in our detected ROIs.

Volumetric change correlation patterns

Finally, we examined correlation patterns of the volumetric change for all brain regions aiming

to identify shared change patterns across multiple ROIs. Hierarchical clustering of the correlation

matrices revealed different patterns between the two groups (Supplementary figure 4). In the test

group, three principal groups of clusters could be identified - the first included the palladium,

hippocampus, amygdala, putamen, insula (all bilaterally), and right anterior cingulate cortex, the

second cluster included mostly occipito-temporal cortical and subcortical nuclei, and the third

included highly correlated regions of the frontal, parietal and occipital cortices. All regions that

passed the statistical threshold of the SBM analysis (Table 1), except for the left fusiform gyrus,

were grouped closely together within the first two clusters, and showed low to negative

correlations the ROIs of the third cluster. This analysis could suggest that the origin of the

volumetric change observed in the regions of the two clusters might be different. The regions

that appear in cluster 1 are often reported in the studies that explore brain changes following

stress, anxiety or traumatic events[14–17], while the regions of the second cluster are less associated

with specific phenomena.

A different pattern was observed in the control group, where the correlation pattern demonstrated

stronger volumetric synchrony (Supplementary figure 4b). This could suggest that changes in the

control group were much more affected by within-participant effects, rather than an exogenous

effect (which could be the outbreak and lockdown in the test-group). It is important to note in

this context that in addition to the experience of the lockdown, the participants in the test group

also had longer time gaps between the two scanning sessions, which might provide an alternative
explanation for the stronger homogeneity in volumetric changes within the control group.

601
Supplementary figure 4. Volumetric changes correlation matrices
Correlation coefficients were calculated between the volumetric change values for all ROI pairs,
analyzed separately for the test **(a)** and the control group **(b)**, clustered according to Euclidean
distances into dendrograms. In the test group, the first and second cluster contained the
subcortical nuclei (amygdala and putamen) and temporal ROI which were found significant in
the SBM interaction analysis, respectively (highlighted in italic bold font). In the control group, a
more homogeneous change pattern was observed with more robust correlation coefficients
between the ROIs. Pearson correlation coefficients are represented by the color scheme.