Pseudomonas aeruginosa induced acute emphysema through
bacterial secretion in mice

Yajie Zhu[1], and Shuming Pan[1]*

[1]Department of Emergency Medicine, Xinhua Hospital, Shanghai Jiao Tong University
School of Medicine, Shanghai, 200092, China

*Correspondence：Shu-ming Pan, Tel: +8602125078999, Fax: +8602165030840,
E-mail: panshuming@xinhuamed.com.cn

Authors' Contributions

Yajie Zhu contributed to the original findings and the majority of the study conception,
data analysis, and manuscript preparation. Shuming Pan contributed partly to the study
conception, data analysis, and manuscript preparation. Shuming Pan was responsible
for the study supervision. Both authors critically reviewed the manuscript for important
intellectual content and read and approved the final manuscript.

Declaration of interests

The authors declare no conflicts of interest in this work.

Funding

Abstract

Pulmonary emphysema is the major pathological feature of chronic obstructive

pulmonary disease (COPD). It is a slowly progressive pathological condition and acute

lung infection is not considered to play a role in emphysema initiation. Recently, we

found that the secretion from *Pseudomonas aeruginosa* could cause severe lung

emphysema in mice rapidly. Since this bacterium is ubiquitous and secrets proteases,

we hypothesized that direct *P. aeruginosa* airway infection would have a similar effect.

To address this issue, we applied a unilateral lung injury model. *P. aeruginosa* secretion

was extracted and instilled intratracheally into the left lung of C57BL/6 and C3H/HeJ

mice, while the right lung was saved as self-control. Alveolar diameter and lung

compliance were measured. Later, we tested the effect of *P. aeruginosa* inoculation in

C57BL/6 mice, immunosuppressed C57BL/6 mice, and C3H/HeJ (TLR4 deficient)

mice. *P. aeruginosa* secretion extract caused acute panacinar emphysema and decreased

dynamic lung compliance. Different types of emphysema are transformable. However,

the *P. aeruginosa* infection could only elicit emphysema in C3H/HeJ mice and

immunosuppressed C57BL/6 mice, indicating normal immunity is essential to protect

the hosts from emphysema. *P. aeruginosa* could be an important pathogen in COPD

initiation. Our finding filled a major gap in COPD pathogenesis.

Key words: Emphysema, COPD, pathology, pathogenesis, unilateral lung injury,

Pseudomonas aeruginosa

Introduction

Chronic obstructive pulmonary disease (COPD) is the third leading cause of death globally, characterized by irreversible obstruction of the small airways (bronchiolitis) and destruction of lung parenchyma (emphysema), which lead to air trapping, dyspnea, and even respiratory failure[1]. The pathogenesis of COPD is largely attributed to cigarette smoking induced chronic lung inflammation; however, it could not explain the fact that many COPD patients did not smoke [2], and only 40% of heavy smokers developed emphysema[3]. Current animal models established by passive tobacco smoking or elastase instillation poorly reproduce the main features of human COPD[4]. Accumulating evidence suggest a better understanding of the disease mechanism is needed.

Pseudomonas aeruginosa is a gram-negative, aerobic, opportunistic bacterium with a wide distribution. In the lung, *P. aeruginosa* is known to prevalent among immunocompromised hosts, causing cystic fibrosis, bronchiectasis and nosocomial pneumonia [5]. *P. aeruginosa* would colonize after COPD formation and contribute to the acute exacerbation [6, 7]. It can be isolated from 4-15% of adults with COPD, a number probably greatly underestimated due to this organism's unique ability to form biofilms, making it difficult to detect[8]. *P. aeruginosa* expresses a variety of virulence factors including endotoxin, exotoxin, elastase, alkaline proteases and hemolysins [9]. The proteases can disrupt epithelial cell tight junctions, cleave collagen, IgG, IgA, complement, and degrade surfactant proteins, causing harm to the lung[10]. However, the manifestation of *P. aeruginosa* pneumonia reported is nonspecific compared with other bacteria[11].

Recently, we discovered accidentlly that distinguished from Klebsiella pneumoniae, Staphylococcus aureus, Escherichia coli and Proteus mirabilis, the secretion of *P. aeruginosa* could induce severe pulmonary emphysema in murine rapidly. Surprised by this finding, for bacterial factors were never reported to be an initiator in emphysema

development [4, 12], we hypothesized that this phenomenon was caused by the proteases secreted by *P. aeruginosa*. Thus, theoretically, *P. aeruginosa* infection shall lead to acute emphysema too. To address the issue, a unilateral lung instillation model of normal C57BL/6, immunosuppressed C57BL/6 and C3H/HeJ mice were used. We observed the dynamic pathology evolution and physiology changes in this study.

Materials and Methods

Ethical statement

All experiments referring to the use of animals in this study were approved by the Institutional Animal Care and Use Committee of Shanghai Xinhua Hospital affiliated to Shanghai Jiao Tong University School of Medicine (XHEC-F-2018-047).

Mice

C57BL/6 mice (10 weeks old) were purchased from Shanghai Sippr-BK laboratory animal Co. Ltd; C3H/HeJ mice (10weeks old) were purchased from Shanghai SLAC laboratory animal Co. Ltd. All mice were accommodated at the Model Animal Research Center of Xinhua Hospital in a specific pathogen-free animal facility under constant temperature and humidity, with sufficient qualified food and water for 1 to 2 weeks before use.

Isolation and identification of *P. aeruginosa*

P. aeruginosa strain was originally separated from sputum, confirmed through mass spectrometry and the strain kept at the bacteriology lab of Xinhua Hospital. To observe the morphology and culture characteristics, the purified strain was streaked on Sheep blood agar plates (Yi Hua, Shanghai) and nutrient agar plates (Comagal, Shanghai) and incubated at 37°C for 24-36 hours.

Bacterial secretion extraction and identification

1×10^6 CFU *P. aeruginosa* were streaked on sheep blood agar plate and incubated at

37°C for 24–48 hours. To harvest the *P. aeruginosa* secretion, all colonies on the blood

agar plate were scratched down, collected, and dissolved in 5ml PBS then centrifuged

at 3000 rcm for 10 minutes. The supernatant was collected, filtered through a 0.22μm

PES membrane filter unit (Millex, Merck), and stored at -20°C for later use. The

quantification of the supernatant protein concentration was carried through BCA

protein assay (Beyotime, China). Protein types in the secretion were identified by

protein mass spectrometry. Briefly, the secretion extract went through enzymatic

hydrolysis and was analyzed by a Q-Exactive mass spectrometer (Thermo, USA).

Proteome Discoverer (v.2.4) was used to search all of the raw data thoroughly against

the protein database (UniProt_ *Pseudomonas aeruginosa*). Names and abundances of

proteins were acquired as the result.

Live bacteria preparation

P. aeruginosa was streaked on sheep blood agar plate and incubated at 37°C for 8 hours.

A proper number of colonies were inoculated in sterile PBS and the turbidity was

adjusted to 2 McFarland standards (MCF), which equals 6×10^8 CFU/ml, by the

standard procedure for later use.

Immunosuppression model

The immunosuppressed model of C57BL/6 mice was established by intraperitoneal

administration of dexamethasone sodium phosphate (DEX) 25mg/kg per day for a

consecutive 5 days referring to the method described by Rasmussen KR and

colleges[13]. Control mice received the same dosage of normal saline intraperitoneally.

The immunosuppression status was confirmed on the sixth day by evaluating the

organ-to-body weight ratios of the spleen (relative spleen weight)[14]. C3H/HeJ mice

were used directly as immunodeficient mice for their lack of Toll-like receptor 4.

Unilateral lung instillation

In the unilateral lung injury mice model, the left lung was set as the instillation side

and the right lung was set as the control side. Mice were anesthetized through

injecting 1% pentobarbital sodium (50mg/Kg) intraperitoneally and put to a proper

supine position. The anterior cervical hair was removed and the skin disinfected with

75% ethanol. To expose the trachea, a small longitudinal incision was made and

endotracheal intubation performed between the second to fourth cricoid cartilage

using a 24G intravenous indwelling catheter (Introcan, BRAUN). The 19mm long

24G cannula was then inserted gently into the left main bronchus. A volume of 30ul *P.*

aeruginosa secretion extract was delivered into the left lung of each mouse using

microliter syringes (Gaoge, Shanghai). Different dosage of live bacteria was

inoculated through the same method. After the intratracheal instillation, mice were put

back to the mouse cage with a thermal pad underneath, resting in the left recumbent

position until full recovery from anesthesia.

Measurement of lung compliance

Dynamic lung compliances were measured 7 days after unilateral lung instillation of

bacterial secretion. The main surgery procedures were the same as above. A 22G

intravenous indwelling catheter (Introcan, BRAUN) was used for endotracheal

intubation. After exposure of the main trachea, 5mm of the catheter tip was inserted

into the main trachea and a ventilator for rodents (Vent Jr, Kent) was connected to the

catheter. Target tidal volume range between 0.2-0.5ml was set and the corresponding

peak pressure was recorded each (RR=100bpm). Lung dynamic compliance (C_{dyn}) was

calculated through linear regression by GraphPad Prism 8 software.

Lung histopathology

Mice were sacrificed at different time points set. The lung was inflated with 4%

paraformaldehyde fixative under constant pressure of 15 to 25 cm H_2O via the trachea

and the trachea then tied off. Lung and heart were harvested en bloc and submerged in

fixative for approximately 24 hours before the heart was removed and lung was

embedded in paraffin blocks for tissue sections (5μm). Hematoxylin and eosin staining

were performed subsequently. Slides were scanned (PANNORAMIC, 3DHISTECH),

reviewed and evaluated.

Alveolar mean linear intercept (Lm)

Air space enlargement was measured using the Short & Hennig method, 1967.

Briefly, HE stained sections were scanned and reviewed in Caseviewer software. 10

random fields of each right lung and left lung were randomly selected. Equally

crossed two lines (both 1000um) were drawn on the scope. The mean linear intercept

Lm was then calculated: $Lm=N*L/m$ [15]. In this case, $Lm=2*10^4/m$ (μm). To better

measure how much the left lung air space has enlarged compare to the right lung, Lm

Ratio was further calculated, Lm Ratio=Lm (left lung)/ Lm (right lung).

Statistics

The data were expressed as the mean ± standard error of mean. Statistical analysis

was performed in the GraphPad Prism 8 software. Pair-wise comparisons were made

using t-tests, whereas multiple comparisons were made using one-way analysis of

variance (ANOVA) test. The confidence interval was set at 95% for all tests. Groups

of three or four animals were used in each experiment.

Results

Identification of *P. aeruginosa*

P. aeruginosa was originally isolated and purified from a clinical sputum sample and

confirmed through mass spectrometry. The strain exhibited typical *P. aeruginosa*

characteristics including strong ginger odor, green metallic sheen, and beta-hemolytic

on blood agar with blue-green pigment producing on nutrient agar (**Figure 1**). The *P.*

aeruginosa secretion extract was translucid, light green colored with a strong ginger

odor. The total protein concentration was 0.35mg/ml.

***P. aeruginosa* secretion causes emphysema in different mice**

Two days after unilateral pulmonary instillation of *P. aeruginosa* secretion, prominent

panacinar emphysema with acute inflammation appeared in the left lung of both

C57BL/6 mice and C3H/HeJ mice while the right lung remained mostly unharmed

(**Figure 2**). The inflammation is characterized by neutrophil infiltration and erythrocyte

diapedesis. Lm Ratio indicates the alveolar of the left lung had enlarged more than three

times in C57BL/6 mice, and two times in C3H/HeJ mice.

Seven days after the instillation, inflammation faded and panacinar emphysema was

well developed and remained in C3H/HeJ mice without fibrosis. While interestingly in

C57BL/6 mice, the panacinar emphysema had changed to distal emphysema with most

of the left lung consolidated. Fibrosis is obvious, inflammation cells can be seen

excreting from alveolars and bronchus. To decide if the panacinar emphysema would

recur after inflammation resolution, we observed C57BL/6 mice models to three weeks

after instillation. Surprisingly, Large apical and subpleural bullae appeared.

Histopathology shows all four types of emphysema (centriacinar, panacinar, distal

acinar and irregular) coexisting in the left lung and compensatory emphysema

developed in the right lung after 21 days (**Figure 2**).

During the above modeling process, C57BL/6 mice underwent dyspnea, huddle, weight

loss and scaphoid abdomen while C3H/HeJ mice showed no obvious symptoms or

weight loss (**Figure 3**).

***P. aeruginosa* secrets heat-labile proteases**

After heated for 80°C, 30mins, the bacterial secretion lost the ability to cause

emphysema but still induced neutrophil infiltration (data not shown), indicating

different components are responsible for alveolar destruction and inflammation.

Protein mass spectrometry confirmed the protein types in the secretion include a high
abundance of metal-dependent hydrolase, urocanate hydratase and other enzymes
(supplementary material), suggesting the emphysema was caused by heat-labile
proteases.

Dynamic lung compliance decreased after instillation

197 Seven days after unilateral lung instillation of bacterial secretion, C_{dyn} of C57BL/6
198 mice decreased from 0.041 to 0.031 ml/cmH$_2$O while C_{dyn} of C3H/HeJ mice
199 decreased from 0.045 to 0.036 ml/cmH$_2$O. Peak pressure and targeted tidal volume
200 were not well linearly correlated as they used to (**Figure 3, D**).

P. aeruginosa infection induces emphysema in immunocompromised but not immunocompetent mice

Different bacteria dosages, infection time and other conditions in normal C57BL/6 mice
were tested, but none of the mice showed noteworthy emphysema (n≥10). A
representative case is the mouse was attacked with 15μl of 5 MCF _P. aeruginosa_, which
formed a prominent abscess full of _P. aeruginosa_ bacteria at the apex of the left lung
after 24 hours, but it only caused bronchiectasis-like pathology change in a few areas
(**Figure 4**). Whereas, all C3H/HeJ mice developed homogenous emphysema after live
bacteria instillation (n=7). 15μl of 0.5 MCF _P. aeruginosa_—1/10 of the dosage used in
C57BL/6 mice—was able to cause obvious centriacinar emphysema in C3H/HeJ. This
result reveals that emphysema induced by _P. aeruginosa_ is not dosage-dependent.

In order to decide if the emphysema induced is related to species or immunity
differences, groups of immunosuppressed C57BL/6 mice, C3H/HeJ mice, and normal
C57BL/6 mice were infected with 30μl of 2 MCF _P. aeruginosa_ for 9 hours (**Figure 5**).
Centriacinar pulmonary emphysema occurred in both immunosuppressed C57BL/6
mice and C3H/HeJ mice. A comparison of the Lm Ratios between different groups
reveals that _P. aeruginosa_ infection had caused airspace enlargement in two

immunocompromised mice groups but alveolar collapse and lung compression in
normal C57BL/6 mice group.

Discussion

In this study, we applied a unilateral lung injury model and found the secretion of *P.*
aeruginosa could cause acute emphysema in mice. Utilizing live *P. aeruginosa* bacteria,
we successfully induced emphysema in both immunosuppressed C57BL/6 and
C3H/HeJ mice, but not in any C57BL/6 mice with normal immunity. Our findings show
that the pathogenesis of COPD can be naturally acute and cigarette smoking irrelevant.
Emphysema induced by *P. aeruginosa* is not dependent on dosage but more related to
the immunity status of the hosts. Chronic inhalation of noxious particles could have
impaired immune function and contributed to COPD formation by creating a
circumstance that favors *P. aeruginosa* to survive and thrive [16, 17].

P. aeruginosa is a nonfastidious bacterium that can proliferate in distilled water. The
secretion of *P. aeruginosa* is a mixture of metal-dependent hydrolase, enzymes, and
bacterial endo/exotoxins which elicited strong inflammation in normal mice.
Interestingly, it seems the neutrophils played a positive role in preventing emphysema
formation. Immunosuppressed C57BL/6 as well as TLR4 deficient C3H/HeJ mice who
had lighter inflammatory response and neutrophils infiltration than C57BL/6 mice,
exhibited more severe airspace dilation.

Protease–antiprotease imbalance as a result of cigarette smoking induced chronic
inflammation is the potential mechanism of COPD, based on the observation that
patients with a genetic deficiency of α1-anti-trypsin have a predisposition to develop
emphysema[3]. Emphysema caused by lung instillation of porcine pancreatic elastase
as a widely used disease model supported this theory[18]. Many studies found elevated
matrix metalloproteinases (MMPs) concentration and increased elastin degradation in
COPD patients [19-21]. Neutrophil or macrophage elastases were held culprit[22]. But

it seems that the proteolytic enzymes secreted by various bacteria had not been taken

into account. Proteases secretion is a common feature in *P. aeruginosa* and other

bacteria[23], however, it didn't occur to us before that the excessive elastase may come

from a pathogen rather than our own neutrophils.

The difference of air space volume is hard to detect since lung tissue has a nature of

instability. In our study, even lungs were carefully inflated with fixative under constant

pressure, the many steps in the histopathology preparation process still impacted the

final pressure, which can be seen from the variation of alveolar diameters of control

lung between different mouse. We solved this difficulty by utilizing the unilateral

instillation model. Unilateral lung injury animals were readily used in a few studies[24,

25]. A mouse model of unilateral injury can provide valuable self-control, eliminate

systemic influence, and reduce the use of lab animals. By ensuring the pressure is equal

within the lung, minor changes could be detected and compared. The Lm Ratio of left

to right lung further revealed clearly how much the airspace has enlarged or compressed.

However, a unilateral lung injury model is not perfect. Sometimes contamination of the

control lung caused by the sneeze or cough of mice is not totally avoidable, especially

in bacterial infection models. Thus, the histology changes in the main bronchus of the

control lung should be interpreted carefully. We tried to do more quantitative

assessment of the lung structure according to standard recommendation[26], but found

it is impractical when a heavy inflammation occurs (Figure 2h). Due to this reason, we

had to sacrifice bacteria inoculated mice at hour 9, before the inflammatory response

makes it impossible to measure Lm.

There are multiple pieces of research studied the correlation between P. aeruginosa and

COPD[27, 28]. Pulmonary P. aeruginosa infection is not found of any differences than

other bacteria[29]. Whereas, most researches conducted used animals with normal

immunity. α1-anti-trypsin secreted by local macrophages and liver is a major inhibitor

of proteases normally presented in serum and tissue fluids. If the immunity is

unimpaired, proper inflammation response can increase α1-anti-trypsin level to
eliminate the destructive effects caused by bacteria proteases[12]. Our research
supported this concept by demonstrating that P. aeruginosa could only cause limited
bronchiectasis in normal mice.

Centriacinar and panacinar forms of emphysema are most commonly found in COPD
patients. The two types are distinct entities. Centriacinar emphysema is the type most
closely associated with tobacco smoking while panacinar emphysema is usually
associated with α1-anti-trypsin deficiency[30, 31]. However, we found the forms of
emphysema are related and transformable to each other. In C3H/HeJ mice, the acute
emphysema transited to stable irreversible panacinar emphysema without fibrosis while
in normal C57BL/6 mice, it went through dynamic change interweaved with
inflammation and developed to other types of emphysema as well as bullae.

On the contrary to the current paradigm taking lung infection as no participant in
emphysema initiation, we discovered for the first time that *P. aeruginosa* and its
secretion can lead to acute emphysema in immunodeficient animals, which is in comply
with its predilection to infect immunocompromised patients. Emphysema induced by P.
aeruginosa in mice recapitulates the main features of human emphysema and COPD.
The role of *P. aeruginosa* in COPD needs more exploration. In the future, strategies
aimed at developing a vaccine or specialized antibiotics targeting *P. aeruginosa* may be
useful for COPD prevention and treatment.

Conclusion

P. aeruginosa infection caused acute emphysema in immunocompromised mice by
bacterial secretion. Different types of emphysema are transformable. *P. aeruginosa*
could be an important pathogen in COPD initiation.

Acknowledgements

We would like to thank Dr. Feng Chen and Dr. Jingxian Liu (Bacteriology Lab, Xinhua

Hospital) as well as Professor Yong Zhang (Department of Immunology, SJTUSM) and

Professor Min Wang (Department of Histology and Embryology, SJTUSM) for their

skilled advice and assistance;

References

1. Rabe KF, Watz H. Chronic obstructive pulmonary disease. *Lancet (London, England)* 2017; 389: 1931-1940.

2. Hogg JC. Pathophysiology of airflow limitation in chronic obstructive pulmonary disease. *Lancet (London, England)* 2004; 364: 709-721.

3. MacNee W. Pathogenesis of chronic obstructive pulmonary disease. *Proceedings of the American Thoracic Society* 2005; 2: 258-266; discussion 290-251.

4. Liang GB, He ZH. Animal models of emphysema. *Chinese medical journal* 2019; 132: 2465-2475.

5. Restrepo MI, Babu BL, Reyes LF, Chalmers JD, Soni NJ, Sibila O, Faverio P, Cilloniz C, Rodriguez-Cintron W, Aliberti S. Burden and risk factors for Pseudomonas aeruginosa community-acquired pneumonia: a multinational point prevalence study of hospitalised patients. *Eur Respir J* 2018; 52.

6. Sethi S, Murphy TF. Infection in the pathogenesis and course of chronic obstructive pulmonary disease. *N Engl J Med* 2008; 359: 2355-2365.

7. Eklöf J, Sørensen R, Ingebrigtsen TS, Sivapalan P, Achir I, Boel JB, Bangsborg J, Ostergaard C, Dessau RB, Jensen US, Browatzki A, Lapperre TS, Janner J, Weinreich UM, Armbruster K, Wilcke T, Seersholm N, Jensen JUS. Pseudomonas aeruginosa and risk of death and exacerbations in patients with chronic obstructive pulmonary disease: an observational cohort study of 22 053 patients. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases* 2020; 26: 227-234.

8. Murphy TF, Brauer AL, Eschberger K, Lobbins P, Grove L, Cai X, Sethi S. Pseudomonas aeruginosa in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2008; 177: 853-860.

9. Li J, Ramezanpour M, Fong SA, Cooksley C, Murphy J, Suzuki M, Psaltis AJ, Wormald PJ, Vreugde S. Pseudomonas aeruginosa Exoprotein-Induced Barrier Disruption Correlates With Elastase Activity and Marks Chronic Rhinosinusitis Severity. *Frontiers in cellular and infection microbiology* 2019; 9: 38.

10. Sadikot RT, Blackwell TS, Christman JW, Prince AS. Pathogen-host interactions in Pseudomonas aeruginosa pneumonia. *Am J Respir Crit Care Med* 2005; 171: 1209-1223.

11. Fujitani S, Sun HY, Yu VL, Weingarten JA. Pneumonia due to Pseudomonas aeruginosa: part I: epidemiology, clinical diagnosis, and source. *Chest* 2011; 139: 909-919.

12. Aster VKAAJ. Robbins Basic Pathology. 2017.

13. Rasmussen KR, Healey MC. Experimental Cryptosporidium parvum infections in immunosuppressed adult mice. *Infect Immun* 1992; 60: 1648-1652.

14. Manente FA, Quinello C, Ferreira LS, de Andrade CR, Jellmayer JA, Portuondo DL, Batista-Duharte A, Carlos IZ. Experimental sporotrichosis in a cyclophosphamide-induced immunosuppressed mice model. *Medical mycology* 2018; 56: 711-722.

15. Dunnill MS. Quantitative methods in the study of pulmonary pathology. *Thorax* 1962; 17: 320-328.

16. Faure K, Sawa T, Ajayi T, Fujimoto J, Moriyama K, Shime N, Wiener-Kronish JP. TLR4 signaling

is essential for survival in acute lung injury induced by virulent Pseudomonas aeruginosa
secreting type III secretory toxins. *Respiratory research* 2004; 5: 1.

17. Drannik AG, Pouladi MA, Robbins CS, Goncharova SI, Kianpour S, Stämpfli MR. Impact of
cigarette smoke on clearance and inflammation after Pseudomonas aeruginosa infection.
Am J Respir Crit Care Med 2004; 170: 1164-1171.

18. Antunes MA, Rocco PR. Elastase-induced pulmonary emphysema: insights from experimental
models. *Anais da Academia Brasileira de Ciencias* 2011; 83: 1385-1396.

19. Chaudhuri R, McSharry C, Brady J, Donnelly I, Grierson C, McGuinness S, Jolly L, Weir CJ,
Messow CM, Spears M, Miele G, Nocka K, Crowther D, Thompson J, Brannigan M, Lafferty
J, Sproule M, Macnee W, Connell M, Murchison JT, Shepherd MC, Feuerstein G, Miller DK,
Thomson NC. Sputum matrix metalloproteinase-12 in patients with chronic obstructive
pulmonary disease and asthma: relationship to disease severity. *J Allergy Clin Immunol*
2012; 129: 655-663.e658.

20. Maclay JD, McAllister DA, Rabinovich R, Haq I, Maxwell S, Hartland S, Connell M, Murchison JT,
van Beek EJ, Gray RD, Mills NL, Macnee W. Systemic elastin degradation in chronic
obstructive pulmonary disease. *Thorax* 2012; 67: 606-612.

21. Wells JM, Parker MM, Oster RA, Bowler RP, Dransfield MT, Bhatt SP, Cho MH, Kim V, Curtis JL,
Martinez FJ, Paine R, 3rd, O'Neal W, Labaki WW, Kaner RJ, Barjaktarevic I, Han MK,
Silverman EK, Crapo JD, Barr RG, Woodruff P, Castaldi PJ, Gaggar A. Elevated circulating
MMP-9 is linked to increased COPD exacerbation risk in SPIROMICS and COPDGene. *JCI*
Insight 2018; 3.

22. Tanner L, Single AB. Animal Models Reflecting Chronic Obstructive Pulmonary Disease and
Related Respiratory Disorders: Translating Pre-Clinical Data into Clinical Relevance.
Journal of innate immunity 2020; 12: 203-225.

23. Galloway DR. Pseudomonas aeruginosa elastase and elastolysis revisited: recent developments.
Molecular microbiology 1991; 5: 2315-2321.

24. Dames C, Akyüz L, Reppe K, Tabeling C, Dietert K, Kershaw O, Gruber AD, Meisel C, Meisel A,
Witzenrath M, Engel O. Miniaturized bronchoscopy enables unilateral investigation,
application, and sampling in mice. *Am J Respir Cell Mol Biol* 2014; 51: 730-737.

25. Singer T, Brand V, Moehrlen U, Fehrenbach H, Purkabiri K, Ott SR, Stammberger U, Ochs M,
Hamacher J. Left-sided mouse intubation: description and evaluation. *Experimental lung*
research 2010; 36: 25-30.

26. Hsia CC, Hyde DM, Ochs M, Weibel ER. An official research policy statement of the American
Thoracic Society/European Respiratory Society: standards for quantitative assessment of
lung structure. *Am J Respir Crit Care Med* 2010; 181: 394-418.

27. Martínez-Solano L, Macia MD, Fajardo A, Oliver A, Martinez JL. Chronic Pseudomonas
aeruginosa infection in chronic obstructive pulmonary disease. *Clin Infect Dis* 2008; 47:
1526-1533.

28. Boutou AK, Raste Y, Reid J, Alshafi K, Polkey MI, Hopkinson NS. Does a single Pseudomonas
aeruginosa isolation predict COPD mortality? *Eur Respir J* 2014; 44: 794-797.

29. Curran CS, Bolig T, Torabi-Parizi P. Mechanisms and Targeted Therapies for Pseudomonas
aeruginosa Lung Infection. *Am J Respir Crit Care Med* 2018; 197: 708-727.

30. Finkelstein R, Ma HD, Ghezzo H, Whittaker K, Fraser RS, Cosio MG. Morphometry of small
airways in smokers and its relationship to emphysema type and hyperresponsiveness. *Am*
J Respir Crit Care Med 1995; 152: 267-276.
31. Kim WD, Eidelman DH, Izquierdo JL, Ghezzo H, Saetta MP, Cosio MG. Centrilobular and
panlobular emphysema in smokers. Two distinct morphologic and functional entities. *Am*
Rev Respir Dis 1991; 144: 1385-1390.

Figures

A

B

Sheep blood agar   Nutrient agar

391   **Figure 1 Culture characteristics of the *P. aeruginosa* strain.** The strain exhibit

392   typical green metallic sheen, beta-hemolysis and produce blue-green pigment.

Figure 2 Unilateral pulmonary instillation of *P. aeruginosa* secretion: Two days after instillation, acute panacinar emphysema with inflammation in the left lung of both C57BL/6 mice (b) and C3H/HeJ mice (e). Seven days after instillation, obvious fibrosis and distal emphysema formed in C57BL/6 mice (h) but not in C3H/HeJ (k). Three weeks after instillation, C57BL/6 mice developed large apical and subpleural bullae (n). Subtypes of emphysema coexisting in the left lung (o, p) and compensatory emphysema developed in the right lung (q). All right lungs served as self-control (c, f, i, m, q);

401   Hematoxylin and eosin (H&E) staining;

Figure 3: Lm Ratio and dynamic lung compliance: (A) Lm of the C57BL/6 mice (Figure 2-a) and the C3H/HeJ mice (Figure 2-d) 2 days after instillation. (B) Lm ratio indicates the alveolar of the left lung had enlarged more than three times in C57BL/6 group, and two times in C3H/HeJ group. (C) C57BL/6 mice endured obvious weight loss in comparison with C3H/HeJ mice. (D) Dynamic lung compliance of both mice decreased 7 days after modeling. The data represent the mean ± SEM of three animals. *P < .05; **P < .01; ***P< .001;

Figure 4 Unilateral live bacteria inoculation: 24 hours after infection, *P. aeruginosa* didn't induce emphysema in C57BL/6 mice. An abscess full of bacteria at the apex of the left lung formed, but only bronchiectasis-like pathological change (a, b). On the contrary, *P. aeruginosa* inoculation induced prominent centriacinar emphysema in C3H/HeJ mice (d, e). Control lungs (c, f); H&E staining;

Figure 5 Unilateral live bacteria inoculation in immunosuppressed C57BL/6: (A)

Histology of lung of normal C57BL/6 mice, C57BL/6 + DEX mice, and C3H/HeJ mice,

Centriacinar pulmonary emphysema developed in both C57BL/6+DEX mice and

C3H/HeJ mice, but not in normal C57BL/6 mice. All mice were inoculated with 30μl

of 2 MCF *P. aeruginosa* and sacrificed after 9 hours; H&E staining. (B) The decrease

of relative spleen weight of C57BL/6 mice treated with DEX shows successful

modeling of hypoimmunity. (C) Lm decreased in C57BL/6 mouse. (D) Lm ratio

between different groups shows *P. aeruginosa* infection caused significant airspace

enlargement in two immunocompromised mice group but airspace compression in the

normal C57BL/6 group. The data represent the mean ± SEM of four animals. *P < .05;

**P < .01; ***P< .001;