Research article

1   **Stochastic asymmetric repartition of lytic machinery in dividing CD8[+] T**

2   **cells generates heterogeneous killing behavior**

4   Fanny Lafouresse[1,#*], Romain Jugele[1,#], Sabina Müller[1], Marine Doineau[2], Valérie

5   Duplan-Eche[3], Eric Espinosa[1], Marie-Pierre Puissegur[1], Sébastien Gadat[2] and Salvatore

6   Valitutti[1,4]

8   [1]INSERM U1037, Centre de Recherche en Cancérologie de Toulouse (CRCT), Université

9   de Toulouse III-Paul Sabatier, 31057 Toulouse, France. Team: Molecular dynamics of

10   lymphocyte interaction, « Equipe labellisée Ligue Nationale contre le Cancer 2018 »

11   [2]Mathematics of decision making and statistics, Toulouse School of Economics, UMR

12   5604, Université Toulouse 1 Capitole, France

13   [3]INSERM, UMR1043, Centre de Physiopathologie de Toulouse Purpan, 31024 Toulouse,

14   France

15   [4]Department of Pathology, Institut Universitaire du Cancer-Oncopole de Toulouse, 31059

16   Toulouse, France.

17   [#] These authors contributed equally to this work

18   [*]Corresponding author: fanny.lafouresse@inserm.fr

Abstract

Cytotoxic immune cells are endowed with a high degree of heterogeneity in their lytic function, but how this heterogeneity is generated is still an open question. We therefore investigated if human CD8$^+$ T cells could segregate their lytic components during telophase, using imaging flow cytometry, confocal microscopy and live cell imaging. We show that CD107a$^+$-intracellular vesicles, perforin and granzyme B unevenly segregate in a constant fraction of telophasic cells during each division round. Mathematical modeling posits that unequal lytic molecule inheritance by daughter cells results from the random distribution of lytic granules on the two sides of the cleavage furrow. Finally, we establish that the level of lytic compartment in individual CTL dictates CTL killing capacity. Together, our results show the stochastic asymmetric distribution of effector molecules in dividing CD8$^+$ T cells. They propose uneven mitotic repartition of pre-packaged lytic components as a mechanism generating non-hereditary functional heterogeneity in CTL.

Introduction

Heterogeneity and plasticity of lymphocyte function are key components of successful adaptive immune responses. Accordingly, several studies put forth the notion that individual mouse and human lymphocytes exhibit high degrees of heterogeneity in both their phenotypic and functional characteristics (Beuneu et al., 2010; Buchholz et al., 2016, 2013; Ganesan et al., 2017; Kumar et al., 2018; Lemaitre et al., 2013; Newell et al., 2012). Functional heterogeneity is not limited to cell differentiation and acquisition of phenotypic and functional characteristics, but also involves late steps of immune cell responses such as $CD8^+$ cytotoxic T lymphocyte (CTL)- and natural killer (NK) cell- mediated cytotoxicity (Guldevall et al., 2016; Halle et al., 2016). Accordingly, we have previously shown that human CTL belonging to the same clonal population exhibit heterogeneity in their lytic function during sustained interaction with target cells (Vasconcelos et al., 2015). While, some CTL kill a limited number of target cells, others emerge as super-killer cells.

One proposed mechanism of functional heterogeneity generation in T lymphocytes is asymmetric cell division (ACD). ACD is a key mechanism to generate cell heterogeneity in biology. It plays a crucial role in embryogenesis by allowing the formation of two distinct cells from a single mother cell (Dewey et al., 2015; Knoblich, 2008). In immunology, ACD has been proposed as a process allowing mouse naive T lymphocytes to divide into short-lived effector T cells and memory T cells, after TCR-triggered division (Arsenio et al., 2015, 2014; Chang et al., 2011, 2007).

In the present work, we investigated the possibility that, in dividing human $CD8^+$ T cells, heterogeneous distribution of molecules relevant for cytotoxic function into nascent daughter cells might contribute to CTL killing heterogeneity.

To address this question, we employed imaging flow cytometry, 3D confocal laser

scanning microscopy, live-cell imaging and mathematical modeling to investigate whether

and how lytic components might differently segregate in telophase.

Our results show that both freshly isolated human peripheral blood $CD8^+$ T cells and clonal

CTL exhibit a heterogeneous repartition of lytic machinery in telophase during TCR-

triggered proliferation which is not part of a classical ACD process. Furthermore, we

demonstrate that heterogeneous lytic compartment repartition resets at each round of CTL

division and is consequently stationary but not hereditary. Finally, we show that the level

of lytic granule expression in individual CTL influences their killing ability.

Together, our results unveil a mechanism of stochastic uneven repartition of pre-packaged

lytic components within intracellular vesicles that generates functional plasticity during

division and contributes to lytic function heterogeneity of individual cells belonging to

clonal populations.

Results

Imaging flow cytometry reveals uneven repartition of lytic machinery in dividing human

CD8+ T cells

To investigate the mechanisms leading to the generation of CTL exhibiting heterogeneous killing ability, we first measured the distribution of lytic machinery components in dividing human CD8+ T cells. Telophase is the *bona fide* cell cycle phase where unambiguous measurement of molecular repartition in nascent daughter cells is performed (Chang et al., 2007; Filby et al., 2011). Lytic granule repartition during human CD8+ T cell division was evaluated using imaging flow cytometry, a technique that combines the advantages of both flow cytometry and microscopy (Basiji and O'Gorman, 2015; Doan et al., 2018; Hritzo et al., 2018). This approach allowed us to collect and analyze a substantial number of cells and to visualize and assess the repartition of molecules of interest within individual cells that were unambiguously identified as being in telophase. Cells in telophase were identified using a computer-assisted gating strategy, on the basis of nuclear and tubulin stainings (**Figure 1-figure supplement 1**). Nuclear staining with SYTOXorange© identified bi-nucleated cells with elongated shape corresponding to cells in the late steps of division (anaphase and telophase). The cells in telophase were identified (and discriminated from possible cellular doublets) on the basis of tubulin staining that allowed us to highlight their midbodies. **Figure 1-figure supplement 2A** shows how masks were applied to delimit the cells and measure the fluorescence intensity of markers of interest in the nascent daughter cells. Cells were also stained with Cell Trace Violet© (CTV), a probe that labels total cell proteins. As previously reported (Filby et al., 2011), we observed that total proteins distribute in nascent daughter cells within a range of 40-60% (**Figure 1-figure supplement**

2B). In our study, CTV staining served both as a marker of cell division (allowing us to

identify cells in the different division rounds (Quah and Parish, 2012)), and to define total

protein repartition in telophase (Filby et al., 2011). This procedure minimized the

possibility that, if some images were taken slightly on an angle, with one daughter cell

slightly more in focus than the other, the markers of interest would artificially appear as

asymmetric. Indeed, asymmetric distribution was defined as cells in telophase in which

repartition of the marker of interest in the nascent daughter cells was beyond the 40-60 %

limits observed for CTV repartition (**Figure 1-figure supplement 2B**). In addition, to

further exclude the possibility of measurement artifacts, we verified individual cells by

eyes and included in the analysis only cells in telophase that were on a even plane.

Specificity of staining for the various markers was validated (see Material and Methods

section).

In a first approach, $CD8^+$ T cells freshly isolated from healthy donor blood samples were

stimulated with immobilized anti-CD3/anti-CD28/ICAM-1 for 72 hours. Anti-CD3/anti-

CD28/ICAM-1 stimulation resulted in activation of human $CD8^+$ T cells as shown by cell

proliferation and CD137 up-regulation (**Figure 1-figure supplement 3**). Repartition of the

lysosomal marker CD107a was investigated in cells in telophase. As shown in **Figure 1A**,

while CTV distribution ranged between 40-60% in dividing T cells, 23 % of telophasic

$CD8^+$ T cells exhibited an uneven distribution of $CD107a^+$ vesicles overcoming the 40-

60% CTV range.

We next investigated the distribution in telophase of lytic components such as perforin and

granzyme B (GrzB), molecules known to be pre-stored in lytic granules. As shown in

Figure 1B and **C**, perforin and GrzB also unevenly segregated into the two nascent

daughter cells in telophase, indicating that daughter cells received a heterogeneous quantity

of lytic components.

The slope of the linear regression curve for the distribution of CD107a, perforin and GrzB

as compared to CTV was close to 0.1, indicating that these 3 molecules distributed

independently from total proteins.

To define whether uneven repartition of lytic components could be observed in fully

differentiated cells, such as memory cells, we investigated CD107a and perforin

distribution in telophase in purified human memory CD8$^+$ T cells. This analysis showed

that also memory CD8$^+$ T cells exhibited uneven repartition of CD107a and perforin in

telophase (**Figure 1-figure supplement 4**).

We next investigated whether lytic machinery asymmetric repartition could also be

observed in activated CD8$^+$ T cell populations composed of monoclonal cells such as

antigen-specific CTL clones. To address this question, we investigated CD107a repartition

in CTL undergoing cell division. For this study, we activated CTL clones using

immobilized anti-CD3/anti-CD28/ICAM-1 for 72 hours. We opted for this stimulation

condition since, in preparatory experiments, we observed that conjugation of CTL with

cognate target cells, results (during the 72 hours culture) in the creation of cellular clumps

and debris due to CTL killing activity, thus making it difficult and potentially misleading

to analyze cells by image flow cytometry and conventional microscopy. As shown in

Figure 1D, we observed that in clonal CTL undergoing cell division, 15% of the two

nascent daughter cells in telophase exhibited uneven distribution of CD107a, thus

confirming and extending observations obtained using CD8$^+$ peripheral blood T cells.

Taken together, the above results indicate that a lysosomal-associated membrane protein

known to be a marker of lytic granules and effector molecules involved in CTL lytic

function, unevenly segregate in 10-23 % of individual human CD8$^+$ T cells undergoing

division.

Confocal laser scanning microscopy confirms uneven repartition of lytic machinery in

dividing CD8$^+$ T cells

Image flow cytometry allows the unambiguously identification and capture of rare events

within a cell population, such as cells in telophase, albeit exhibiting a lower resolution

when compared to classical imaging methods. This notion prompted us to confirm results

obtained using imaging flow cytometry, with additional methods.

We therefore used 3D confocal laser scanning microscopy to measure CD107a content in

telophasic CD8$^+$ T cells following stimulation with immobilized anti-CD3/anti-

CD28/ICAM-1. Although this approach allowed us to collect a relatively small number of

cells in telophase (n=61 compared to n=908 obtained by image flow cytometry), it revealed

that 27% of the CD8$^+$ T cells in telophase exhibited uneven repartition of CD107a, above

a 1.5 threshold (corresponding to the 40-60% range used in imaging flow cytometry

experiments) (**Figure 2A**). **Figure 2B** depicts the maximum intensity projection (MIP) of

a z-stack of images on which measurements of fluorescence intensity were performed (left

panel) and a central z-section (right panel). The asymmetry of CD107a repartition in

nascent daughter cells is better appreciated by looking at the 3D reconstructions of the

dividing cell (**Video 1**).

Together, the above results indicate that confocal laser scanning microscopy provides

results that reinforce those we obtained using imaging flow cytometry and supports the

finding that lytic granules undergo uneven repartition in ~20% of dividing CD8$^+$ T cells.

Uneven repartition of lytic machinery is not accompanied by asymmetric segregation of

fate determining transcription factors and does not require a polarity cue

The observation that lytic components were unevenly inherited in daughter cells prompted

us to investigate whether this process was somehow related to mechanisms of cell fate

determining ACD, a process reported to play a role in mouse naive T lymphocytes

differentiation (Arsenio et al., 2015, 2014; Kaminski et al., 2016; Pham et al., 2014).

Indeed, it has been reported that ACD can result in the generation of one daughter cell

predisposed to become a short-lived effector cell (harboring a high level of the transcription

factors T-bet and c-myc, and of GrzB) and one daughter cell predisposed to become a

memory T cell (Widjaja et al., 2017). We investigated whether uneven repartition of fate

determining transcription factors T-bet and c-myc (Chang et al., 2011; Verbist et al., 2016),

might occur in telophase in freshly isolated peripheral blood CD8+ T cells stimulated with

anti-CD3/anti-CD28/ICAM-1 for 72 hours. As shown in **Figure 3A** and **B**, both T-bet and

c-myc did not unevenly segregate into the two nascent daughter cells during telophase.

Moreover, the slope of the linear regression curve for the distribution of T-bet and c-myc

as compared to CTV was close to 1, indicating that the repartition of these 2 molecules in

telophase followed that of total proteins.

To further define whether the observed uneven repartition of lytic components was or was

not related to ACD, we investigated whether uneven repartition of lytic components was

dependent on a polarity cue (e.g. localized TCR stimulation) as previously described for

ACD (Arsenio et al., 2015; Pham et al., 2014). **Figure 4A** and **B** shows that a polarity cue

was not required to induce uneven distribution of lytic molecules, since comparable

CD107a$^+$ vesicles segregation was observed in peripheral blood CD8$^+$ T cells stimulated

by either immobilized (anti-CD3/anti-CD28/ICAM-1) or soluble (PMA + ionomycin)

stimuli.

Overall, the above results demonstrate that uneven partitioning of lytic compartment in

telophase is not associated with asymmetric segregation of fate determining transcription

factors. Moreover, a polarity cue is not required. All in all, the above results show that, in

human CD8$^+$ T cells, lytic machinery uneven repartition is not related to described

mechanisms of fate determining ACD.

Asymmetric repartition of CD107a$^+$ vesicles reset at each division event and generates

heterogeneous daughter cells

We next investigated whether lytic machinery uneven repartition occurred during

subsequent divisions and whether this process could be involved in preserving lytic

machinery heterogeneity within CD8$^+$ T cell populations.

We considered the cells in the different rounds of division (identified by different peaks of

CTV dilution, **Figure 1-figure supplement 3)** and analyzed CD107a repartition in

telophasic cells. This analysis showed that, in all division rounds considered, a comparable

percentage of cells underwent heterogeneous repartition of CD107a (**Figure 5A and B**).

A complementary observation indicated that the heterogeneity process is stationary but not

hereditary: e.g. a daughter cell originating from a heterogeneous division has a constant

stationary probability to produce a new uneven division. We arrived to this conclusion by

generating CD107a fluorescence intensity (CD107a-FI) density curves of all telophasic

cells having undergone 0, 1 or 2 mitosis. Cells in telophase showing unequal CD107a-FI

repartition were then plotted on these curves (**Figure 5C**). The χ^2 statistical test showed

that these cells were randomly and independently distributed on the CD107a-FI density

curves, supporting the hypothesis that there is no inheritance in the decision to divide

unevenly (see Materials and Methods section).

We next asked whether this process might create a drift in lytic compartment content in

daughter cells leading to the emergence of cellular subsets expressing higher or lower

levels of CD107a. To address this question, we analyzed the total CD107a-FI in all G1

cells (either undivided or following each division round). As shown in **Figure 5D**, the total

CD107a-FI appeared to be broadly similar in the different rounds of division in the whole

populations, suggesting that uneven repartition of CD107a, in a relatively constant fraction

of cells at each division round, does not lead to the emergence of well-defined cellular

subsets expressing higher or lower levels of CD107a. We employed the Kolmogorov-

Smirnov goodness of fit test to determine whether the different curves followed the same

distribution or not. The test strongly rejected the hypothesis that the CD107a expression

curves follow the same distribution during the first two division rounds (see Materials and

Methods section), indicating that during these division events randomly heterogeneous

populations were generated. Nevertheless, our test also showed that the Kolmogorov

distance decreased when the number of divisions increased, indicating that CD107a-FI

density distribution seems to be convergent with a higher number of divisions. To define

where variability was located in the curves, we employed the χ^2 test. The test showed that

variability was distributed all over the curves (i.e. for all the CD107a-FI). Together,

Kolmogorov-Smirnov goodness of fit and χ^2 tests revealed a non-stationary variability in

the content of CD107a$^+$ vesicles in CD8$^+$ T cells during early division events.

Taken together, the above results indicate that asymmetric distribution of CD107a$^+$ vesicles

in telophase is not limited to the first division, but it is rather a stochastic process, inherent

to each division, that perpetuates variability in daughter cells.

Lytic granules randomly distribute on the two sides of the cleavage furrow

To gain direct information about the possibility that lytic components might stochastically

distribute in nascent daughter cells, we visualized lytic granule repartition during division

in individual CTL transfected with mCherry-tagged GrzB mRNA, by live cell microscopy.

mCherry-tagged GrzB showed no preferential localization within cell cytosol at the

different phases of the division and appeared to randomly partition into the two nascent

daughter cells. In some cases, nascent daughter cells exhibited approximately similar

repartition of lytic granules (**Figure 6A, Video 2**), in some other cases lytic granule

repartition appeared to be rather asymmetric (**Figure 6B, Video 3**). Furthermore, we

investigated cell division in 4D (3D plus time). Sorted CD8$^+$ T cells in G2/M phase were

loaded with LysoTracker Red (LTR) to stain their late endosomal lysosomal vesicles (of

which lytic granules are an important fraction (Faroudi et al., 2003)). Nascent daughter

cells were imaged to monitor distribution of LTR$^+$ vesicles and measure the integrated

fluorescence intensity. An example of one CD8$^+$ T cell distributing LTR$^+$ vesicles in a

symmetric fashion during division is shown in **Figure 6C** and **Video 4** (LTR distribution

ranged within 40-60% at all time points measured). One CD8$^+$ T cell that distributed in an

asymmetry fashion LTR$^+$ vesicles is shown in **Figure 6D** and **Video 5** (LTR distribution

ranged above or below 40-60% at all time points measured). Additional examples of cells

dividing in symmetric and asymmetric fashion are shown in **Figure 6-figure supplement**

1 and **Video 6.**

While lytic granules seemed to be overall randomly distributed between nascent daughter

cells, in some cases the videos showed that lytic granules did not behave completely

independently from each other and exhibited some clustering. We therefore used a

computational approach to establish whether the above-described process might be linked

to a random repartition of lytic components into the two nascent daughter cells. We first

calculated the probability to obtain an asymmetric distribution of lytic granules (*e.g.* a

repartition of the granules into the two daughter cells out of the 40-60% range) related to

the granule number per dividing cell. This computation is naturally handled with a binomial

modeling for the behavior of the population of n granules (see Materials and Methods

section). This analysis showed that for n <100 the probabilities that individual particles

distribute asymmetrically on the two sides of the cleavage furrow are relatively high

(**Figure 6E**). Using stimulated emission depletion (STED) on CTL stained for GrzB, we

estimated that 14 to 65 (mean = 37) lytic granules are contained within individual CTL.

Our estimation well-matched with numbers published in previous studies, ranging between

10 and 100 (Chiang et al., 2017; Clark et al., 2003; Kataoka et al., 1996; Peters et al., 1991).

These values are compatible with a significant probability of stochastic uneven repartition

of lytic granules in telophase, assuming that all lytic granules behave independently.

Since our videos indicate that lytic granules might form transitory aggregates within

confined intracellular spaces, we upgraded our mathematical simulation of lytic granule

repartition in telophase to include the possibility that lytic granules might not segregate

completely independently. We simulated particle correlation during cell division for 10 to

100 particles. To evaluate the correlation level between individual particles during cell

division (ranging from 0 = absence of correlation to 1 = 100% correlation) for a given

probability of asymmetric repartition (outside the interval [40%-60%]), we used a Monte-

Carlo approach (see Material and Methods section).

The analysis shows that for a probability of 20% asymmetric repartition of particles

(corresponding to 20% uneven repartition of lytic granules during cell division

experimentally measured by imaging flow cytometry and confocal imaging), particle

correlation has a relatively low value (4% for 37 particles, 0.04; CI95%, 0.035-0.045),

suggesting that lytic granules mainly segregate independently during cell division.

Taken together, cell imaging and computational results strongly suggest that the observed

stationary unequal distribution of lytic granules in telophase is the result of a stochastic

repartition of particulate cytosolic structures on the two sides of the cleavage furrow in

dividing cells.

The level of lytic granule content in individual CTL dictates CTL killing capacity

To assess the consequences of an uneven distribution of lytic compartment on CTL-

mediated cytotoxicity, we investigated cytotoxic efficacy in CTL expressing high and low

lytic granule content. Clonal CTL were loaded with LysoTraker blue, and cells containing

high (LysoTracker^{High}) and low (LysoTracker^{Low}) levels were FACS sorted. As shown in

Figure 7A, sorted LysoTracker^{High} and LysoTracker^{Low} CTL populations maintained their

difference in LysoTracker staining at least 24 hours after cell sorting. The cytotoxic

efficacy of sorted CTL populations was compared at different effector:target (E:T) ratios

by measuring the percentage of killed targets (7-AAD positive targets). For each ratio,

LysoTracker^{High} CTL were more efficient than LysoTracker^{Low} CTL in exerting

cytotoxicity (**Figure 7B-C**), although basal killing (in the absence of peptide stimulation)

was comparable between LysoTracker^{High} and LysoTracker^{Low} CTL (**Figure 7C**). The

above results indicated that lytic granule content is associated with killing efficacy. To

strengthen these findings, we performed additional experiments on sorted CTL for high

and low LysoTracker staining and measured CD107a surface exposure and CD8

internalization following 4 hour conjugation with target cells. Results show that

LysoTracker^{high} CTL exhibited a higher lytic granule secretion as detected by CD107a

exposure when compared to LysoTracker^{low} CTL (**Figure 7D**). However, productive TCR

engagement was comparable in both populations as detected by similar levels of CD8

internalization (Huang et al., 2019; Xiao et al., 2007) (**Figure 7E**).

Together, these results indicate that the lytic granule cargo of individual CTL and not their

activation properties directly impact killing behavior. They imply that stochastic uneven

distribution of lytic granules in dividing CTL determine heterogeneous killing behavior at

the single cell level.

Discussion

In the present study we found that, in both freshly isolated peripheral blood CD8[+] T cells and clonal CTL, ~ 20 percent of telophasic cells undergoes asymmetric distribution of the lytic compartment into the two daughter cells. Our results establish that CD8[+] killing capacity is associated to lytic compartment level and strongly suggest that uneven lytic machinery repartition produces CD8[+] T cell populations with heterogeneous killing capacities.

We used imaging flow cytometry, a technique that combines the advantages of flow cytometry and microscopy and allows the detection and analysis of rare cells within whole cell populations on the basis of their morphological and staining characteristics (Basiji and O'Gorman, 2015; Doan et al., 2018; Hritzo et al., 2018). We thus acquired and analyzed a significant number of relatively rare events of T cell divisions by precisely identifying cells in telophase. The use of CTV distribution as a parameter of global protein repartition in telophase together with the acquisition of an important number of cell divisions strengthens our analysis. In addition, we investigated lytic granule repartition in dividing CD8[+] T cells by 3D confocal laser scanning microscopy and 4D live cell imaging. These techniques allowed visualization of lytic granule repartition in telophase with a high time/space resolution and strengthened imaging flow cytometry data by providing unambiguous visualization of lytic granule partitioning.

Our results demonstrate that the uneven lytic machinery distribution is not related to ACD. In mouse T lymphocytes, ACD has been reported as a mechanism contributing to the generation of effector/memory daughter cells following division of an individual naive T cell in response to polarizing cues (Arsenio et al., 2015; Chang et al., 2007). Establishment

of asymmetry has been associated to the uneven inheritance by daughter cells of

transcription factors such as c-Myc and T-bet known for their role in the induction of

metabolic reprogramming and in the acquisition of T cell effector function respectively

(Chang et al., 2011; Verbist et al., 2016). Following the original observation of uneven

repartition of proteasomes in dividing mouse CD4$^+$ T cells leading to asymmetric

degradation of T-bet in daughter cells (Chang et al., 2011), additional cellular effectors

including metabolic and signaling pathways have been found to be implicated in fate

determining ACD in mouse naive T lymphocytes (Kaminski et al., 2016; Pollizzi et al.,

2016; Verbist et al., 2016). Our results, by showing that lytic granule repartition is not

accompanied by a detectable asymmetric segregation of T-bet and c-Myc and does not

require a polarity cue, suggest that the lytic machinery uneven distribution observed in

human CD8$^+$ T cells is not related to previously described ACD. Although we could not

detect an asymmetric repartition of classical lineage-determining transcription factor, in

our models, this observation does not exclude the possibility that ACD might play a role

in the differentiation of human naive T cells into effector and memory subsets during initial

antigen specific immune responses. It is therefore possible that the discrepancy between

our results and previous studies on ACD in mouse T lymphocytes arises from the different

nature of the cells involved in the study. It should also be noted that, beside ACD, other

mechanisms can contribute to the generation of different T lymphocyte populations from

naive lymphocytes and, more in general, can play a role in T lymphocyte heterogeneity.

Alternative models postulate that lymphocyte differentiation might be achieved via the

accumulation of progressive differences among daughter cells due to variation in the

quantity of the inherited proteins (Buchholz et al., 2016; Cobbold et al., 2018; Gerlach et

al., 2013; Girel and Crauste, 2019; Pham et al., 2014; Rohr et al., 2014; Schumacher et al.,

2010).

A puzzling question is how asymmetric distribution of lytic components in telophase is

generated. Our results provide a stepping-stone to answer this question. First, mathematical

analysis of our imaging flow cytometry data provides an interpretation of our results that

is compatible with a stochastic distribution of lytic components during cell division. On

one hand, mathematical analysis shows that the process of asymmetric distribution is

stationary in terms of the fraction of involved cells: e.g. occurs always on a similar

percentage of cells, at each division round, in different experiments and following different

stimuli. On the other hand, the heterogeneity process, although stationary is not hereditary:

e.g. a daughter cell originating from a heterogeneous division has a constant stationary

probability to produce a new uneven division. Second, live-cell imaging shows lytic

granule distribution during mitosis. We did not observe any specific pattern of lytic granule

repartition (polarization at the membrane or close to the cleavage furrow) before or during

cell division. Instead, lytic compartments appeared randomly distributed in cell cytosol.

Our observations are consistent with the mathematical modeling of intracellular vesicle

distribution showing the high probability of an uneven distribution of a relatively small

quantity of granules. In other words, pre-packaged molecular components within a few

relatively big vesicles might have higher probability to be asymmetrically partitioned in

telophase than molecular components dispersed throughout the cytosol.

Moreover, it should be noted, that our videos and results in Figure 6 suggest that, in a

limited number of division events, granules might not segregate completely independently

from each other, implying that some active mechanism of granule segregation might

contribute, to a minor extent, to lytic granule repartition in telophase.

Together, our results point out a mechanism of heterogeneity generation that is for the

most part stochastic and might be a general mechanism for generating heterogeneity in

dividing cells. The possibility that particulate material is unevenly distributed in telophase

into two nascent daughter cells has been proposed for other organelles and in other cellular

systems (Bergeland et al., 2001; Carlton et al., 2020; Sanghavi et al., 2018). Indeed, in

MDCK cells, microscopy analysis and mathematical modelling based on the laws of

probability suggested that endosomes/lysosomes partitioning between daughter cells is

stochastic (Bergeland et al, 2001). Others show that in telophasic cells, endosomal

compartments are clustered at the cleavage furrow, suggesting that microtubules are

involved in this process. However, no mechanism ensuring endosomal compartment

anchorage to either spindle has been revealed, suggesting that this repartition is stochastic.

Similarly, in Dictyostelium cells, it has been demonstrated that dynein and kinesin motors

drive phagosomes segregation independently of each other and stochastically (Shanghavi

et al 2018). To our knowledge, our present study is the first to relate a mechanism of a

random segregation of organelles to functional heterogeneity of immune cells.

What could be the functional role of asymmetric molecular segregation during mitosis in

human CD8[+] T cells? We propose that a mechanism of asymmetric distribution in

telophase (that is stationary at each division, but not inherited by daughter cells) can be

instrumental to randomly generate short-lived CTL cohorts harboring functional

heterogeneity while ensuring globally reproducible antigen specific CD8[+] T cell responses.

This process might confer robustness to CTL responses through population averaging

(Buchholz et al., 2016; Hodgkin et al., 2014).

It is interesting to note that our results present analogies with previously published data in

which asymmetric segregation of internalized exogenous antigen was found to occur

during B cell division (Thaunat et al., 2012). Together with this previous study, our results

reveal an intriguing capacity of both T and B cells to stochastically distribute in telophase

their acidic compartments: MHC Class II compartments for B cells and lytic granules for

$CD8^+$ T cells. Thus, stochastic distribution in telophase appears to be a major mechanism

ensuring a high variability of both humoral and cellular adaptive immune responses during

lymphocyte clonal expansion.

454   **Material and Methods**

456   **Key resources table**

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
cell line (*Homo-sapiens*)	HLA-A2 restricted CD8+ T cell clone (VLAELVKQI)	(Khazen et al., 2016)		
cell line (*Homo-sapiens*)	HLA-A2 restricted CD8+ T cell clone (NLVPMVATV)	(Khazen et al., 2016)		
cell line (*Homo-sapiens*)	HLA-A2 restricted CD8+ T cell clone (VLAELVKQI)	(Khazen et al., 2016)		
cell line (*Homo-sapiens*)	JY (EBV-transformed B cells)	(Khazen et al., 2016; Vasconcelos et al., 2015)		
biological sample (*Homo-sapiens*)	Buffy coats of Healthy donors	EFS, Toulouse, France		with consent and approval AC-2014-2384
antibody	Anti-human CD3 (Human monoclonal, TR66)	Enzo	cat# ALX-804-822 RRID:AB_2051037	(1 µg/ml)
antibody	Anti-human CD28 (Mouse monoclonal, CD28.2)	eBioscience	cat# 16-0289-81 RRID:AB_468926	(1 µg/ml)
recombinant protein	Recombinant human ICAM-1-Fc fusion protein	R&D Systems	cat# 720-IC	(0.5µg/ml)
antibody	Anti-human CD107a (Mouse monoclonal, H4A3)	BD Pharmingen	cat# 555798 RRID:AB_396132	(10µg/ml)

antibody	Anti-human CD107a AlexaFluor® 647 (Mouse monoclonal, H4A3)	BD Pharmingen	cat# 562622 RRID:AB_2737684	(diluted at 1/100)
antibody	Anti-human Granzyme B (Mouse monoclonal, GB11)	Thermo Scientific	cat# MA1-80734 RRID:AB_931084	(10µg/ml)
antibody	Anti-human Granzyme B AlexaFluor® 647 (Mouse monoclonal, GB11)	BD Pharmingen	cat# 561999 RRID:AB_10897997	(10µg/ml)
antibody	Anti-human T-bet (Rabbit polyclonal, Tbx21)	Abcam	cat# ab181400	(10µg/ml)
antibody	Anti-human C-myc (Mouse monoclonal, 9E10)	Thermo Scientific	cat# MA1-980 RRID:AB_558470	(10µg/ml)
antibody	anti-human α-tubulin (Rabbit polyclonal)	Abcam	cat# ab15246 RRID:AB_301787	(diluted at 1/100)
antibody	anti-mouse IgG1 Alexa Fluor 647 (Goat polyclonal)	Invitrogen	cat# A21240 RRID:AB_2535809	(10µg/ml)
antibody	anti-mouse IgG1 Alexa Fluor 488 (Goat polyclonal)	Invitrogen	Cat # A-21121 RRID:AB_2535764	(10µg/ml)
antibody	anti-rabbit (H+L) AlexaFluor®488 (Goat polyclonal)	Invitrogen	cat# A11034 RRID:AB_2576217	(10µg/ml)
antibody	anti-rabbit (H+L) AlexaFluor®647 (Donkey polyclonal)	Invitrogen	cat# A31573 RRID:AB_2536183	(10µg/ml)

antibody	anti-rabbit AlexaFluor®555 (Goat polyclonal)	Invitrogen	cat# A21428 RRID:AB_2 535849	(10µg/ml)
antibody	anti-mouse IgG Abberior Star 580 (Goat polyclonal)	Abberior Instruments	cat# 52403	(10µg/ml)
antibody	Anti-human CD107a-PEcy7 (mouse monoclonal, H4A3)	BD Pharmingen	cat# 561348 RRID:AB_1 0644018	(diluted at 1/50)
antibody	Anti-human CD8-FITC (mouse monoclonal, HIT8A)	BD Pharmingen	cat# 555634 RRID:AB_3 95996	(diluted at 1/50)
recombinant DNA reagent	MGC Human GZMB Sequence verified cDNA (Clone Id: 5223876)	GE Healthcare BIO Sciences	cat# MHS6278-202801737	
recombinant DNA reagent	mCherry-SEpHluorin	(Koivusalo et al., 2010)	Addgene cat# 32001	
recombinant DNA reagent	pT7-GZMB-mCherry-SEpHluorin	This paper		
sequence-based reagent	Primer : XhoI-T7-GzB Forward caaCTCGAGTAATAC GACTC ACTATAGGGAGACC CGGTA CCatgcaaccaatcctgcttct gcc	This paper		
sequence-based reagent	Primer: EcoRI-GzB-noSTOP-R caaGAATTCcggcgtg gcgtttcatggttttctttatcc ag	This paper		
peptide	CMV peptide p65 (NV-9)	GeneCust	cat# 181329	

recombinant protein	Human rIL-2	Miltenyi Biotec	cat# 130-097-748	(150 IU/ml)
recombinant protein	Human rIL-15	Miltenyi Biotec	cat# 130-095-766	(50ng/mL)
commercial assay or kit	EasySep Negative human CD8+ T cell isolation kit	StemCell Technologies	cat# 17953	
commercial assay or kit	EasySep human Memory CD8+ T cell enrichment kit	StemCell Technologies	cat# 19159	
software, algorithm	IDEAS SpotCount Threshold (M03,nucleus,60)	Amnis, Luminex		
software, algorithm	IDEAS Area Range Threshold (M02,tubulin,75), 50-5000, 0-0.5	Amnis, Luminex		
software, algorithm	Fiji	(Schindelin et al., 2012)		
software, algorithm	Imaris Software	Oxford Instruments		
software, algorithm	ZEN ZEISS Efficient Navigation			
software, algorithm	Huygens Professional version 18.10 using CMLE algorithm with SNR:7	Scientific Volume Imaging, USA		STED images were deconvolved
software, algorithm	Python software version 3.5			χ^2 of independence test, χ^2 of homogeneity

				test and Kolmogorov-Smirnov goodness of fit test
software, algorithm	GraphPad Prism software version 5 for windows			Paired Student's t-test
software, algorithm	FlowJo software	TreeStar		
other	SYTOX^{TM} Orange Dead Cell Stain	Thermo Fisher Scientific	cat# S11368	Manufacturer recommended dilution
other	DAPI	Molecular Probes, Invitrogen^{TM}	cat# D1306 RRID:AB_26 29482	
other	Hoechst 33342	ThermoFisher Scientific	cat# 1399	(200 ng/ml)
other	CellTrace^{TM} Violet Cell Proliferation kit	ThermoFisher Scientific	cat# C34557	(5 µM)
other	LysoTraker Blue (DND22) Dye	Molecular probes	cat# L7525	(200 nM)
other	LysoTraker Red (DND99) Dye	Molecular probes	cat#L7528	(200 nM)
other	7-Aminoactinomycin D (7-AAD)	BD Pharmingen^{TM}	cat# 559925	(0.25µg)
other	Ibidi µ-slide chambered coverslips Angiogenesis	Ibidi, Biovalley	cat# 81506	
other	Ibidi µ-slide chambered coverslips 8 well	Ibidi, Biovalley	cat# 80821	
other	Nunc Lab-Tek chamber slides 8 wells	Nunc, ThermoFisher	cat#1 54526	

	Micromesh array (100μm)	Microsurface, Tebu- Bio	cat# MMA-0500-100-08-01	
other				

Experimental model and subject details

Peripheral blood mononuclear cells were isolated from buffy coats of healthy donors

obtained through the Etablissement Français du Sang (EFS, Toulouse, France). Blood

samples were collected and processed following standard ethical procedures (Helsinki

protocol), after obtaining written informed consent from each donor and approval by the

French Ministry of the Research (transfer agreement AC-2014-2384). Approbation by the

ethical department of the French Ministry of the Research for the preparation and

conservation of cell lines and clones starting from healthy donor human blood samples has

been obtained (authorization No DC-2018-3223).

Cell culture and stimulating conditions

Total human CD8$^+$ T cells were purified from healthy donor blood samples using the

EasySep Negative human CD8$^+$ T cell isolation kit (StemCell Technologies). CD8$^+$ T cells

were routinely ~90% pure. Memory human CD8$^+$ T cells were purified from healthy donor

blood samples using the EasySep Human Memory CD8$^+$ T cell enrichment kit (StemCell

Technologies), cells were routinely ~90% CD8$^+$RO$^+$RA$^-$.

HLA-A2 restricted CD8+ T cell clones, specific for the NLVPMVATV peptide or the

VLAELVKQI peptide of the CMV protein pp65 were cultured in complete RPMI/HS

medium (RPMI 1640 medium supplemented with 5% human AB serum; Inst.

Biotechnologies J. Boy, Reims), minimum essential amino acids, HEPES, sodium pyruvate

(Invitrogen), 2-mercaptoethanol (5 µM, Gibco) and 150 IU/ml human rIL-2 and 50ng/ml

rIL-15). Clones were re-stimulated every 2-3 weeks in 24-wells plate with 1×10^6 irradiated

(35 Gy) allogeneic peripheral blood mononuclear cells (isolated on Ficoll Paque Gradient

from fresh heparinized blood samples of healthy donors, obtained from EFS) and 1×10^5

irradiated EBV-transformed B cells. Complete RPMI/HS-Medium was supplemented with

1 µg/ml PHA.

EBV-transformed B cells (JY) HLA-A2+ were used as target cells and cultured in RPMI

1640 GlutaMAX supplemented with 10% FCS and 50 µM 2-mercaptoethanol, 10 mM

HEPES, 1X MEM non-essential amino acids, 1X sodium pyruvate, 10 µg/mL

ciprofloxacine. Profiling of JY cells has been done using STR.

All cell lines are routinely screened for mycoplasma contamination using MycoAlert

mycoplasma detection kit (Lonza, Basel, SW).

For imaging flow cytometry (ImageStream®X, Merk) and confocal laser-scanning

microscopy human CD8$^+$ T cells or CD8$^+$ T cell clones were stimulated for 72h with

immobilized anti-CD3 (1 µg/ml, TR66, (Valitutti, 1995)), anti-CD28 (1 µg/ml, clone

CD28.2, eBioscience) and immobilized recombinant ICAM1-Fc fusion protein (0.5µg/ml,

R&D Systems) in complete RPMI/HS medium. For confocal laser-scanning, cells were

plated on anti-CD3/CD28/ICAM1 coated Nunc Lab-Tek Chamber Slide™ system 8 wells

at 500 000 cells / well. For image stream analysis, cells were plated on anti-

CD3/CD28/ICAM1 coated 24 well plates at 1.5×10^6 cells / well.

498
Image Stream analysis

<u>*Staining and acquisition strategy*</u>. Cells were first stained with CellTrace ^{TM} Violet Cell

Proliferation Kit (CTV) in PBS (5 µM, 20 min, 37°C). After 72 hours of stimulation (cf:

Cell culture and stimulating condition), cells were fixed in 1% PFA (10 min, 37°C) and

permeabilized in permeabilization buffer (PBS 3% BSA, 0.1% saponin, Sigma) for 5 min.

Cells were incubated for 45 min with the indicated antibodies: AlexaFluor® 647 anti-human

CD107a antibody (diluted at 1/100, clone H4A3; BD Pharmingen ™), anti-human Perforin

(10µg/ml, clone δG9; BD Pharmingen ™), AlexaFluor® 647 anti-human Granzyme B

antibody (10µg/ml, clone GB11, BD Pharmingen ™), anti-human T-bet (Tbx21) (10µg/ml,

clone 4B10; Abcam), anti-human C-myc (10µg/ml, clone 9E10; Thermo scientific), anti-

human α-tubulin (diluted at 1/100, rabbit polyclonal; Abcam). The following secondary

antibodies were used: AlexaFluor®488 or 647 goat anti-mouse IgG1 (10µg/ml; Invitrogen),

AlexaFluor®488 or 647 anti-rabbit (H+L) (10µg/ml; Invitrogen). For image acquisition,

cells were adjusted to 10.10^6 - 20.10^6 /mL in FACS buffer (PBS, 1% FCS, 5% Hepes, 0.1%

Azide) containing SYTOX™ Orange Dead Cell Stain (recommended dilution, Thermo

Fisher Scientific) for nuclear staining. Cells were acquired using ImageStream®X (IsX;

Amnis, Luminex) technology.

Telophase discrimination strategy. Amnis IDEAS software was used to analyze IsX data

and identify cells in telophase. As in classical cytometry data analysis, cells in G2/M phase

were first selected according to their DNA content (fluorescence of SYTOX orange). A

mask based on nuclear staining was employed (SpotCount Threshold (M03, nucleus, 60))

to visualize the nuclei of cells/events in the G2/M fraction at the single cell level. A second

mask (Area Range (Threshold (M02, tubulin, 75), 50-5000, 0-0.5)) based on the α-tubulin

staining (to clearly identify the narrow intracellular bridge of highly condensed α-tubulin

that participates to midbody formation) was employed to distinguish telophases from

anaphases or cell-doublets. Finally, the results from both masks were used to manually

verify that selected cells were cells unambiguously in telophase.

Analysis of cell protein distribution during telophase using Fiji. Staining intensities of α-

tubulin, CTV and of the different markers of interest were analyzed on Fiji to determine

the percentage of proteins inherited by the two nascent daughter cells in telophase.

Watershed function of Fiji software was used on the α-tubulin staining intensity to

determine the specific areas corresponding to the two nascent daughter cells in telophase.

The obtained areas were converted to masks that were next applied to measure CTV and

the fluorescence of the different markers of interest. This procedure allowed us to

determine the intensity of fluorescence in the two nascent daughter cells in telophase

respectively. The percentage of staining in each nascent daughter cell was determined as:

Fluorescence Intensity of daughter cell 1 / (Fluorescence Intensity of daughter cell 1 +

Fluorescence Intensity of daughter cell 2) x 100. To test the specificity of the staining with

the different antibodies used to study molecular repartition in telophase, we measured the

fluorescent intensity of secondary antibodies or isotype controls as compared to specific

antibodies. This analysis gave the following values of MFI: CD107a 70.527 isotype control

13.621; perforin 716.312, secondary mouse antibody 56.383; GrzB 677.445 isotype control

13.621; T-Bet 356.228 secondary mouse antibody 56.383; c-Myc 1.434.537 secondary

rabbit antibody 14.231.

3D laser scanning microscopy on fixed cells

After 72 hours of stimulation, cells were fixed in 1% PFA (10 min, 37°C). Permeabilization

and staining with antibodies were performed in PBS 3% BSA, 0.1% saponin (Sigma) for

5 min and 45 min respectively. The following antibodies were used: anti-human CD107a

(10µg/ml, cloneH4A3, BD Pharmingen ™) followed by AlexaFluor®488 goat anti-mouse

IgG1 (10µg/ml; Invitrogen), anti-human α-tubulin (diluted at 1/100, rabbit polyclonal;

Abcam) followed by AlexaFluor®555 goat anti-rabbit (10µg/ml; Invitrogen). Nuclei were

labeled with DAPI (1µg/ml, 5 min). The samples were mounted in 90% glycerol-PBS

containing 2.5% DABCO (Sigma) and examined using a LSM710 (Zeiss) confocal

microscope with a ×63 plan-Apochromat objective (1.4 oil) with an electronic zoom of 4.

Cells in telophase were identified on the basis of nuclear and tubulin marker staining. 3D

images (using the z-stack function) were acquired for every cell identified as being in

telophase. CD107a fluorescence intensity in the two nascent daughter cells was measured

on 2-D image projections obtained applying the Sum function of Fiji Software to z-stack

series. Since the background noise made the watershed function unsuitable to use, a region

of interest (ROI) corresponding to the nascent daughter cell was manually drawn on the

basis of brightfield and tubulin staining. We determined the percentage of CD107a staining

in each nascent daughter cell as: CD107a intensity of daughter cell 1 / (CD107a intensity

of daughter cell 1 + CD107a intensity of daughter cell 1) × 100.

Stimulated Emission Depletion Microscopy

CTL were seeded on poly-L-lysin coated high performance coverslips and fixed in 3%

PFA (10 min, 37°C). Permeabilization and staining were performed in PBS 3% BSA, 0.1%

saponin (Sigma) for 5 min and 60 min respectively. Cells were stained with an anti-human

Granzyme B antibody (10µg/ml, clone GB11, Thermo Scientific) followed by a goat anti-

mouse IgG Abberior Star 580 (Abberior Instruments). Coverslips (high performance

D=0.17mm +/-0.005, ZEISS, Germany) were mounted on microscopy slides using

Mowiol-DABCO.

STED images were acquired with a Leica SP8 STED 3X microscope (Leica Microsystems,

Germany) using a HC PL APO CS2 100X/1.4 oil immersion objective. To optimize

resolution without bleaching in 3-D, the 775 nm STED lasers line was applied at the lowest

power that can provide sufficient improvement in resolution compared to confocal. Z-stack

series were acquired sequentially with the pulsed 532 nm laser. For image acquisition, we

used the following parameters: 3 time average/line, 400 Hz scan speed. STED images were

subsequently deconvoluted with Huygens Professional (SVI, USA) using the CMLE

algorithm, with a signal to noise ratio (SNR) of 7. 3-D image visualization was performed

using the Fiji software.

Live cell imaging

For 3D live cell imaging, the T7 GZMB sequence was obtained by PCR amplification as a

XhoI-EcoRI fragment from pCMV-SPORT6-GZMB by using XhoI-T7-GZB forward

primer and EcoRI-GRZB noSTOP reverse primer (Employed primers: Name: XhoI-T7-

GzB F

caaCTCGAGTAATACGACTCACTATAGGGAGACCCGGTACCatgcaaccaatcctgcttctgcc

Name: EcoRI-GzB-noSTOP-R   caaGAATTCcggcgtggcgtttcatggttttctttatccag).

XhoI-EcoRI fragment was cloned as a mCherry-SEpHlurin fusion construct in the

pmCherry-SEpHlurin vector to produce the vector pGZMB-mCherry-SEpHluorin

available to *in vitro T7 transcription. The plasmid* pCMV-SPORT6-GZMB and pmCherry-

SEpHlurin were purchased from Addgene.

For efficient transfection of human CTL with tagged molecules allowing to monitor lytic

granule repartition during mitosis, we synthetized capped and tailed poly(A) mCherry-

tagged Granzyme B mRNA by *in vitro* transcription from the plasmid pGZMB-mCherry-

SEpHluorin. One microgramme of pGZMB-mCherry-SEpHluorin was first linearized by

NotI digestion to be used as templates for *in vitro* transcription by the T7 RNA polymerase

using mMESSAGE mMACHINE T7 Ultra kit as per manufacturer's protocol.

Human CTL were transfected using a GenePulser Xcell electroporation system (BioRad).

$1x10^6$ CTL (5days after restimulation therefore in expansion phase) were washed and

resuspended in 100μl Opti-MEM medium (Gibco) at RT with 2μg mCherry-tagged

Granzyme B mRNA (*square wave* electrical pulse at 300V, 2ms, 1 pulse). 18 hours after

transfection the transfection efficacy was verified by FACS analysis (typically 50-80%).

Transfected CTL were seeded into poly-D-lysine-coated eight-well chambered slides

(Ibidi, Munich, Germany) before imaging. Chambered slides were mounted on a heated

stage within a temperature-controlled chamber maintained at 37°C and constant CO_2

concentrations (5%) and inspected by time-lapse laser scanning confocal microscopy

(LSM880, Zeiss, Germany with 1 image /30 seconds) for additional 5-6 hours using a Tile

Scan mode to enlarge the acquisition fields and capture the rare cells undergoing

spontaneous division during the time of acquisition.

For 4D live cell imaging, 72 hours after stimulation, $CD8^+$ T cells were stained with

Hoechst (200 ng/ml, ThermoFisher Scientific) to sort cells in G2/M phase by flow-

cytometry (BD FACSAria SORP, BD Biosciences). Sorted cells were stained with

LysoTracker Red (200 nM ThermoFisher) for 30 min at 37°C and washed. 20 000 cells in

5% HS/IL2/IL15 complete RPMI medium supplemented with 10 mM HEPES were seeded

into poly-D-lysine-coated eight-well chambered slides (Ibidi, Munich, Germany) pre-

coated with PDMS micromesh arrays (Microsurfaces, Melburn, Australia) containing 100-

μm-diameter wells. Cells were 4D imaged (time and z-stack) on a heated stage within a

temperature-controlled chamber maintained at 37°C and constant CO_2 concentrations (5%)

and inspected over night by time-lapse laser scanning confocal microscopy with a Plan-

Apochromat 40x/1.3 Oil DIC M27 using an LSM780 or LSM880, Zeiss, Germany) or by

spinning disk time-lapse microscopy using a spinning-disk microscope (Nikon) running on

Metamorph software. A camera emCCD Evolve (Photometrics) was used for acquisitions.

Image analysis was performed using Fiji software and 4-D videos and snapshots were

generated with Imaris software.

Cytotoxicity assay

CTL were incubated with 200nM LysoTracker Blue® a probe staining the acidic lytic

compartment of these cells (Faroudi et al., 2003) for 30 minutes at 37°C/5% CO_2 in 5%

FCS/RPMI/HEPES. After washing, cells expressing the highest and lowest 5-10 %

LysoTracker Blue staining were sorted using a FACSARIA-SORP (BD Biosciences). CTL

were used for standard over-night killing assays on the day of cell. Target cells were left

unpulsed or pulsed with 10μM antigenic peptide during 2 hours at 37°C/5% CO2, washed

three times and subsequently transferred to a 96 well U-bottom plate at 10×10^3 cells/100μl

RPMI, 5% FCS/HEPES. CTL were added to the target cells at the indicated effector (CTL):

target (JY) ratio, in 100μl RPMI, 5% FCS/HEPES. Cells were pelleted for 1 min at 455 g

and incubated at 37°C/5% CO2 overnight. Before FACS analysis, 0.25μg 7-

Aminoactinomycin D (7-AAD; BD Biosciences) and FITC conjugated anti-CD8 antibody

were added to each sample in order to measure the percentage of dead target cells. For the

CD107a exposure and CD8 internalization assay, sorted CTL were incubated with target

cells at 0.5:1 E/T ratio for 4 hours. Cells were stained with PE-cy7 conjugated anti-CD107a

antibody and FITC conjugated anti-CD8 antibody for 30 min in FACS-buffer (1% human

serum, 1% Foetal Calf Serum in PBS), washed, acquired on a Fortessa flow cytometer (BD

Biosciences) and analysed by using FlowJo software (TreeStar).

Statistical methods

Paired Student's t-test, was performed to determine the statistical significance of

differences between the groups (GraphPad Prism software version 5).

Chi-square of independence test was performed to determine the independence between

the level of expression of a given marker and the capacity of a cell in telophase to

asymmetrically distribute this marker (Python software version 3.5).

Kolmogorov-Smirnov goodness of fit test was performed to compare law between

probability distribution of a marker of interest in cells in G1 (Python software version 3.5).

Chi-square of homogeneity test was performed (in addition Kolmogorov-Smirnov

goodness of fit test) to determine where the probability distribution of a marker of interest

varies (Python software version 3.5).

Statistical procedures

Independence Chi2 test between heterogeneous cells and all cells	Test statistic (χ^2)	$\chi^2_{1-\alpha,\mathrm{dl}}$	p-value (p)	Degree of freedom (dl)
CD107a, Experiment 1, 0 division	4,060439	11.07	0.540748	5
CD107a, Experiment 1, 1 division	3.565087	11.07	0.613563	5
CD107a, Experiment 1, 2 divisions	1.614763	7.815	0.656047	3

	$D_{n,m}$	$\chi^2_{1-\alpha,dl}$	p-value	dl
CD107a, Experiment 2, 0 division				
CD107a, Experiment 2, 1 division	0.278928	7.815	0.963942	3
CD107a, Experiment 2, 2 divisions	0.413804	7.815	0.937376	3
CD107a, Experiment 3, 0 division	2.36867	15.51	0.967574	8
CD107a, Experiment 3, 1 division	2.092976	9.488	0.718663	4
CD107a, Experiment 3, 2 divisions	0.655225	9.488	0.956734	4

661   Table 1: Results of independence Chi-square test in telophase

662   In the independence Chi-square test, we compare the theoretical effective ($e_{i,j}$) to the

663   observed effective ($n_{i,j}$). The test statistic is defined by:

$$\chi^2 = \sum_{i,j} \frac{(n_{i,j} - e_{i,j})^2}{e_{i,j}}$$

We compare it to $\chi^2_{1-\alpha,dl}$, the quantile of the χ^2 distribution associated to the $1-\alpha$

quantile. The quantile with $1-\alpha = 95\%$ is the value such that $P(X < \chi^2_{0.95,dl}) = 95\%$

where P stands for the probability distribution of the Chi-square statistics with the

associated degree of freedom dl.

We reject the hypothesis of independence between division of heterogeneous cells and

division of all cells in one experiment when $\chi^2 \geq \chi^2_{1-\alpha,dl}$ or when the p-value p satisfies

$p < \alpha = 5\%$.

The red boxes represent the situations where we do not reject the hypothesis of

independence of division between heterogeneous cells and all cells in one experiment. We

shall observe that we never reject the hypothesis of independence.

Kolmogorov-Smirnov test Experiment 1	0 division		1 division		2 divisions	
	$D_{n,m}$	p-value	$D_{n,m}$	p-value	$D_{n,m}$	p-value
0 division						
1 division	0.13148	0				
2 divisions	0.220034	0	0.116283	0		

	0 division	1 division	2 divisions

Kolmogorov-Smirnov test Experiment 1	$D_{n,m}$	p-value	$D_{n,m}$	p-value	$D_{n,m}$	p-value
0 division	██████	██████				
1 division	0.087873	0	██████	██████		
2 divisions	0.0891924	0	0.04634	0.03582	██████	██████
3 divisions	0.054621	0.0159	0.067702	0.001185	0.047275	0.116534

Kolmogorov-Smirnov test Experiment 1	0 division		1 division		2 divisions	
	$D_{n,m}$	p-value	$D_{n,m}$	p-value	$D_{n,m}$	p-value
0 division	██████	██████				
1 division	0.14714	0.002607	██████	██████		
2 divisions	0.209553	0	0.143594	0	██████	██████
3 divisions	0.190642	0	0.121757	0	0.038549	0.3545

Table 2: Results of Kolmogorov-Smirnov test on G1

The Kolmogorov-Smirnov test consists in analyzing if two independent samples follow the same law comparing their cumulative distribution function. We denote the two samples $X_1, X_2, \dots X_n$ and $Y_1, Y_2, \dots Y_m$. If we denote by F_n and F_m their cumulative distribution respectively, the test statistic is defined by:

$$D_{n,m} = \quad x \in R |F_n(x) - F_m(x)|$$

We compare it to $d_{n,m,1-\alpha}$, the quantile of the associated Kolmogorov-Smirnov distribution.

We then reject the hypothesis of adequation between cells of one division and cells of one other division in one experiment when $D_{n,m} \geq d_{n,m,1-\alpha}$ or when the p-value p satisfies $p < \alpha = 5\%$.

The red boxes represent the situation where we do not reject the hypothesis of adequation between cells in one division and cells in another division. The white box represents the situation where we reject this hypothesis.

Probability of an asymmetric repartition of lytic granules

695   To obtain a tractable formula for the computation of the probability of an asymmetric

696   repartition of lytic granules, we use a binomial model that translates that each granule

697   possesses a probability of 0.5 to attain each of the two daughter cells. The binomial model

698   also assumes that all the granules behave independently of each other.

699   In that case, the probability of an asymmetric division for n granules is then equal to

$$p_n = 2^{-n} \sum_{k<0.4n} \frac{n!}{k!\,(n-k)!} + 2^{-n} \sum_{k>0.6n} \frac{n!}{k!\,(n-k)!}$$

To evaluate the correlation level between particles (between 0 and 1) for a given probability

of asymmetric repartition (outside the interval [40%-60%]), we use a Monte-Carlo

approach where we sampled a sequence of correlated random variables distributed

according to a Bernoulli distribution of parameter 0.5 since to the best of our knowledge

there is no explicit formula to calculate a such probability of asymmetric repartition. Even

with a Monte-Carlo approach, the simulation is a little bit involved: if r is the correlation

level and if X_i is the value of the random variable at step i, then X_{i+1} is obtained by :

$$X_{i+1} = X_i Y_i + Z_i(1 - Y_i)\,(Formula A)$$

where Z_i is a Bernoulli distribution of parameter 0.5 and Y_i a Bernoulli distribution of

parameter r. We shall verify that when $X_1, X_2, ... X_n$ are sampled according to Formula A,

they are Bernoulli distributed and pairwise correlated with a correlation r. Hence, we then

mimic the correlated division with this model and then estimate the probability of

asymmetric repartition with 5000 Monte-Carlo simulations for each value of r and a size

of n=90 cells. We then evaluate the desired probability for r varying in a regularly spaced

grid from 0 to 1 with a space equal to 0.02.

Acknowledgements

We thank Dr. Stephane Manenti for discussion and critical reading of the manuscript ; Dr.

Hellen Robey and Dr. Pauline Gonnord for discussion. We thank Dr. Liza Filali for advice

in image analysis and Dr. Juliet Foote for critical reading of the manuscript. We thank the

flow cytometry and imaging core facilities of the INSERM UMR 1043, CPTP and of the

INSERM UMR 1037, CRCT, Toulouse, France.

This work was supported by grants from the Laboratoire d'Excellence Toulouse Cancer

(TOUCAN) (contract ANR11-LABEX), Region Occitanie (contracts RCLE R14007BB,

671 34 No 12052802, and RBIO R15070BB, No 14054342), Fondation Toulouse Cancer

Santé (contract 2014CS044) from the Ligue Nationale contre le Cancer (Equipe labellisée

2018) and from Bristol-Myers Squibb (No CA184-575). R.J. was supported by the Ligue

Nationale contre le Cancer.

The funders had no role in study design, data collection and analysis, decision to publish,

or preparation of the manuscript.

Competing Interests statement

The authors declare no competing financial interests.

References

Arsenio J, Kakaradov B, Metz PJ, Kim SH, Yeo GW, Chang JT. 2014. Early specification of
CD8+ T lymphocyte fates during adaptive immunity revealed by single-cell gene-
expression analyses. *Nat Immunol* **15**:365–372. doi:10.1038/ni.2842
Arsenio J, Metz PJ, Chang JT. 2015. Asymmetric Cell Division in T Lymphocyte Fate
Diversification. *Trends Immunol* **36**:670–683. doi:10.1016/j.it.2015.09.004

742 Basiji D, O'Gorman MR. 2015. Imaging flow cytometry. *J Immunol Methods* **423**:1–2.
743   doi:10.1016/j.jim.2015.07.002
744 Bergeland T, Widerberg J, Bakke O, Nordeng TW. 2001. Mitotic partitioning of endosomes and
745   lysosomes. *Current Biology* **11**:644–651. doi:10.1016/S0960-9822(01)00177-4
746 Beuneu H, Lemaitre F, Deguine J, Moreau HD, Bouvier I, Garcia Z, Albert ML, Bousso P. 2010.
747   Visualizing the functional diversification of CD8+ T cell responses in lymph nodes.
748   *Immunity* **33**:412–23. doi:10.1016/j.immuni.2010.08.016
749 Buchholz VR, Flossdorf M, Hensel I, Kretschmer L, Weissbrich B, Graf P, Verschoor A,
750   Schiemann M, Hofer T, Busch DH. 2013. Disparate individual fates compose robust
751   CD8+ T cell immunity. *Science* **340**:630–5. doi:10.1126/science.1235454
752 Buchholz VR, Schumacher TN, Busch DH. 2016. T Cell Fate at the Single-Cell Level. *Annu Rev*
753   *Immunol* **34**:65–92. doi:10.1146/annurev-immunol-032414-112014
754 Carlton JG, Jones H, Eggert US. 2020. Membrane and organelle dynamics during cell division.
755   *Nat Rev Mol Cell Biol* **21**:151–166. doi:10.1038/s41580-019-0208-1
756 Chang JT, Ciocca ML, Kinjyo I, Palanivel VR, McClurkin CE, Dejong CS, Mooney EC, Kim JS,
757   Steinel NC, Oliaro J, Yin CC, Florea BI, Overkleeft HS, Berg LJ, Russell SM, Koretzky
758   GA, Jordan MS, Reiner SL. 2011. Asymmetric proteasome segregation as a mechanism
759   for unequal partitioning of the transcription factor T-bet during T lymphocyte division.
760   *Immunity* **34**:492–504. doi:10.1016/j.immuni.2011.03.017
761 Chang JT, Palanivel VR, Kinjyo I, Schambach F, Intlekofer AM, Banerjee A, Longworth SA,
762   Vinup KE, Mrass P, Oliaro J, Killeen N, Orange JS, Russell SM, Weninger W, Reiner
763   SL. 2007. Asymmetric T lymphocyte division in the initiation of adaptive immune
764   responses. *Science* **315**:1687–91. doi:10.1126/science.1139393
765 Chiang SCC, Wood SM, Tesi B, Akar HH, Al-Herz W, Ammann S, Belen FB, Caliskan U, Kaya
766   Z, Lehmberg K, Patiroglu T, Tokgoz H, Ünüvar A, Introne WJ, Henter J-I, Nordenskjöld
767   M, Ljunggren H-G, Meeths M, Ehl S, Krzewski K, Bryceson YT. 2017. Differences in
768   Granule Morphology yet Equally Impaired Exocytosis among Cytotoxic T Cells and NK
769   Cells from Chediak–Higashi Syndrome Patients. *Front Immunol* **8**.
770   doi:10.3389/fimmu.2017.00426
771 Clark RH, Stinchcombe JC, Day A, Blott E, Booth S, Bossi G, Hamblin T, Davies EG, Griffiths
772   GM. 2003. Adaptor protein 3–dependent microtubule-mediated movement of lytic
773   granules to the immunological synapse. *Nat Immunol* **4**:1111–1120. doi:10.1038/ni1000
774 Cobbold SP, Adams E, Howie D, Waldmann H. 2018. CD4+ T Cell Fate Decisions Are
775   Stochastic, Precede Cell Division, Depend on GITR Co-Stimulation, and Are Associated
776   With Uropodium Development. *Front Immunol* **9**:1381. doi:10.3389/fimmu.2018.01381
777 Dewey EB, Taylor DT, Johnston CA. 2015. Cell Fate Decision Making through Oriented Cell
778   Division. *J Dev Biol* **3**:129–157. doi:10.3390/jdb3040129
779 Doan M, Vorobjev I, Rees P, Filby A, Wolkenhauer O, Goldfeld AE, Lieberman J, Barteneva N,
780   Carpenter AE, Hennig H. 2018. Diagnostic Potential of Imaging Flow Cytometry. *Trends*
781   *in Biotechnology* **36**:649–652. doi:10.1016/j.tibtech.2017.12.008
782 Faroudi M, Utzny C, Salio M, Cerundolo V, Guiraud M, Muller S, Valitutti S. 2003. Lytic versus
783   stimulatory synapse in cytotoxic T lymphocyte/target cell interaction: Manifestation of a
784   dual activation threshold. *Proceedings of the National Academy of Sciences of the United*
785   *States of America* **100**:14145–14150. doi:10.1073/pnas.2334336100
786 Filby A, Perucha E, Summers H, Rees P, Chana P, Heck S, Lord GM, Davies D. 2011. An
787   imaging flow cytometric method for measuring cell division history and molecular
788   symmetry during mitosis. *Cytometry Part A* **79A**:496–506. doi:10.1002/cyto.a.21091
789 Ganesan AP, Clarke J, Wood O, Garrido-Martin EM, Chee SJ, Mellows T, Samaniego-Castruita
790   D, Singh D, Seumois G, Alzetani A, Woo E, Friedmann PS, King EV, Thomas GJ,
791   Sanchez-Elsner T, Vijayanand P, Ottensmeier CH. 2017. Tissue-resident memory

features are linked to the magnitude of cytotoxic T cell responses in human lung cancer.
Nature Immunology **18**:940–950. doi:10.1038/ni.3775

Gerlach C, Rohr JC, Perie L, van Rooij N, van Heijst JW, Velds A, Urbanus J, Naik SH, Jacobs
H, Beltman JB, de Boer RJ, Schumacher TN. 2013. Heterogeneous differentiation
patterns of individual CD8+ T cells. *Science* **340**:635–9. doi:10.1126/science.1235487

Girel S, Crauste F. 2019. Model-Based Assessment of the Role of Uneven Partitioning of
Molecular Content on Heterogeneity and Regulation of Differentiation in CD8 T-Cell
Immune Responses. *Frontiers in Immunology* **10**:16.

Guldevall K, Brandt L, Forslund E, Olofsson K, Frisk TW, Olofsson PE, Gustafsson K,
Manneberg O, Vanherberghen B, Brismar H, Karre K, Uhlin M, Onfelt B. 2016.
Microchip Screening Platform for Single Cell Assessment of NK Cell Cytotoxicity. *Front*
Immunol **7**:119. doi:10.3389/fimmu.2016.00119

Halle S, Keyser KA, Stahl FR, Busche A, Marquardt A, Zheng X, Galla M, Heissmeyer V, Heller
K, Boelter J, Wagner K, Bischoff Y, Martens R, Braun A, Werth K, Uvarovskii A,
Kempf H, Meyer-Hermann M, Arens R, Kremer M, Sutter G, Messerle M, Forster R.
2016. In Vivo Killing Capacity of Cytotoxic T Cells Is Limited and Involves Dynamic
Interactions and T Cell Cooperativity. *Immunity* **44**:233–45.
doi:10.1016/j.immuni.2016.01.010

Hodgkin PD, Dowling MR, Duffy KR. 2014. Why the immune system takes its chances with
randomness. *Nat Rev Immunol* **14**:711. doi:10.1038/nri3734-c1

Hritzo MK, Courneya J-P, Golding A. 2018. Imaging flow cytometry: A method for examining
dynamic native FOXO1 localization in human lymphocytes. *Journal of Immunological*
Methods **454**:59–70. doi:10.1016/j.jim.2018.01.001

Huang M, Zhang J, Chen W. 2019. FACS isolation of low percentage human antigen-specific
CD8+ T cells based on activation-induced CD3 and CD8 downregulation. *Journal of*
Immunological Methods **472**:35–43. doi:10.1016/j.jim.2019.06.013

Kaminski MM, Liedmann S, Milasta S, Green DR. 2016. Polarization and asymmetry in T cell
metabolism. *Semin Immunol* **28**:525–534. doi:10.1016/j.smim.2016.10.002

Kataoka T, Sato M, Kondo S, Nagai K. 1996. Estimation of pH and the Number of Lytic
Granules in a $CD8^+$ CTL Clone Treated with an Inhibitor of Vacuolar Type H^+-
ATPase, Concanamycin A. *Bioscience, Biotechnology, and Biochemistry* **60**:1729–1731.
doi:10.1271/bbb.60.1729

Khazen R, Muller S, Gaudenzio N, Espinosa E, Puissegur MP, Valitutti S. 2016. Melanoma cell
lysosome secretory burst neutralizes the CTL-mediated cytotoxicity at the lytic synapse.
Nat Commun **7**:10823. doi:10.1038/ncomms10823

Knoblich JA. 2008. Mechanisms of asymmetric stem cell division. *Cell* **132**:583–97.
doi:10.1016/j.cell.2008.02.007

Koivusalo M, Welch C, Hayashi H, Scott CC, Kim M, Alexander T, Touret N, Hahn KM,
Grinstein S. 2010. Amiloride inhibits macropinocytosis by lowering submembranous pH
and preventing Rac1 and Cdc42 signaling. *J Cell Biol* **188**:547–563.
doi:10.1083/jcb.200908086

Kumar BV, Kratchmarov R, Miron M, Carpenter DJ, Senda T, Lerner H, Friedman A, Reiner SL,
Farber DL. 2018. Functional heterogeneity of human tissue-resident memory T cells
based on dye efflux capacities. *JCI Insight* **3**:e123568. doi:10.1172/jci.insight.123568

Lemaitre F, Moreau HD, Vedele L, Bousso P. 2013. Phenotypic CD8+ T cell diversification
occurs before, during, and after the first T cell division. *J Immunol* **191**:1578–85.
doi:10.4049/jimmunol.1300424

Newell EW, Sigal N, Bendall SC, Nolan GP, Davis MM. 2012. Cytometry by time-of-flight
shows combinatorial cytokine expression and virus-specific cell niches within a
continuum of CD8+ T cell phenotypes. *Immunity* **36**:142–52.
doi:10.1016/j.immuni.2012.01.002

Peters PJ, Borst J, Oorschot V, Fukuda M, Krähenbühl O, Tschopp J, Slot JW, Geuze HJ. 1991.
Cytotoxic T lymphocyte granules are secretory lysosomes, containing both perforin and
granzymes. *The Journal of Experimental Medicine* **173**:1099–1109.
doi:10.1084/jem.173.5.1099
Pham K, Sacirbegovic F, Russell SM. 2014. Polarized cells, polarized views: asymmetric cell
division in hematopoietic cells. *Front Immunol* **5**:26. doi:10.3389/fimmu.2014.00026
Pollizzi KN, Sun I-H, Patel CH, Lo Y-C, Oh M-H, Waickman AT, Tam AJ, Blosser RL, Wen J,
Delgoffe GM, Powell JD. 2016. Asymmetric inheritance of mTORC1 kinase activity
during division dictates CD8+ T cell differentiation. *Nat Immunol* **17**:704–711.
doi:10.1038/ni.3438
Quah BJ, Parish CR. 2012. New and improved methods for measuring lymphocyte proliferation
in vitro and in vivo using CFSE-like fluorescent dyes. *J Immunol Methods* **379**:1–14.
doi:10.1016/j.jim.2012.02.012
Rohr JC, Gerlach C, Kok L, Schumacher TN. 2014. Single cell behavior in T cell differentiation.
Trends Immunol **35**:170–7. doi:10.1016/j.it.2014.02.006
Sanghavi P, D'Souza A, Rai Ashim, Rai Arpan, Padinhatheeri R, Mallik R. 2018. Coin Tossing
Explains the Activity of Opposing Microtubule Motors on Phagosomes. *Current Biology*
28:1460-1466.e4. doi:10.1016/j.cub.2018.03.041
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P,
Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**:676–682. doi:10.1038/nmeth.2019
Schumacher TN, Gerlach C, van Heijst JW. 2010. Mapping the life histories of T cells. *Nat Rev*
Immunol **10**:621–31. doi:10.1038/nri2822
Thaunat O, Granja AG, Barral P, Filby A, Montaner B, Collinson L, Martinez-Martin N,
Harwood NE, Bruckbauer A, Batista FD. 2012. Asymmetric segregation of polarized
antigen on B cell division shapes presentation capacity. *Science* **335**:475–9.
doi:10.1126/science.1214100
Valitutti S. 1995. Sustained signaling leading to T cell activation results from prolonged T cell
receptor occupancy. Role of T cell actin cytoskeleton. *Journal of Experimental Medicine*
181:577–584. doi:10.1084/jem.181.2.577
Vasconcelos Z, Muller S, Guipouy D, Yu W, Christophe C, Gadat S, Valitutti S, Dupre L. 2015.
Individual Human Cytotoxic T Lymphocytes Exhibit Intraclonal Heterogeneity during
Sustained Killing. *Cell Rep* **11**:1474–85. doi:10.1016/j.celrep.2015.05.002
Verbist KC, Guy CS, Milasta S, Liedmann S, Kaminski MM, Wang R, Green DR. 2016.
Metabolic maintenance of cell asymmetry following division in activated T lymphocytes.
Nature **532**:389–93. doi:10.1038/nature17442
Xiao Z, Mescher MF, Jameson SC. 2007. Detuning CD8 T cells: down-regulation of CD8
expression, tetramer binding, and response during CTL activation. *Journal of*
Experimental Medicine **204**:2667–2677. doi:10.1084/jem.20062376
883

 Figure Legends

Figure 1: Lytic components are asymmetrically distributed in dividing CD8$^+$ T cells.

(**A-C**) Freshly isolated polyclonal CD8$^+$ T cells or (**D**) CTL clones were stimulated by

immobilized anti-CD8/anti-CD28/ICAM-1 during 72h and stained with antibodies directed

against the indicated markers. Cells in telophase were identified using Imaging Flow

Cytometry **(A)** Left panel: Each dot represents one nascent daughter cell. Only one of the

two nascent daughter cells in telophase is plotted. The percentage of staining for CD107a

in the presented cell (*x axis*) is plotted against the percentage of staining for total cell

proteins (CTV, *y axis*). Asymmetric cells were defined as cells in telophase in which

repartition of CD107a in the nascent daughter cells was beyond the 40-60% observed for

CTV repartition (n=908 from 3 independent experiments). Right panel: example of

asymmetric and symmetric cell distribution of CD107a, as detected by Imaging Flow

Cytometry. **(B)** Left panel: The percentage of staining for perforin in the presented nascent

daughter cell is plotted as in panel A. Asymmetric cells were defined as indicated in panel

A (n=191 from 3 independent experiments). Right panel: example of asymmetric and

symmetric cell distribution of perforin. **(C)** Left panel: The percentage of staining for GrzB

in the presented nascent daughter cell is plotted as in panel A. Asymmetric cells were

defined as indicated in panel A (n=728 from 2 independent experiments). Right panel:

example of asymmetric and symmetric cell distribution of GrzB. (**D**) Left panel: The

percentage of staining for CD107a is plotted as in panel A. Asymmetric cells were defined

as indicated in panel A (n=352 from 3 independent experiments). Right panel: example of

asymmetric and symmetric cell distribution of CD107a.

Numbers highlighted in blue in the plots indicate the % of cells exhibiting asymmetric

repartition of the marker of interest. Red lines indicate the global distribution of the data.

Red numbers indicate the slope of the linear regression curve for marker distribution. See

Figure S1, S2, S3 and S4.

Figure 2: CD107a$^+$ vesicles uneven segregation in telophase is confirmed by confocal

laser scanning microscopy. Freshly isolated polyclonal CD8$^+$ T cells were stimulated by

immobilized anti-CD8/anti-CD28/ICAM-1 during 72h and stained with antibodies directed

against CD107a. Cells in telophase were identified using confocal laser scanning

microscopy. **(A)** Analysis of CD107a repartition in dividing cells. The fold increase of

CD107a staining in the brighter nascent daughter cell as compared to the other nascent

daughter cell is shown. The dotted red line indicates the limit between symmetric and

asymmetric cells (1,5 fold increase, corresponding to a 60-40% variation) (n=61 from 2

independent experiments). Each dot represents one CD8$^+$ T cell in telophase. **(B)** Example

of an asymmetric cell in division. Green CD107a, cyan DAPI, red Tubulin. A maximum

intensity projection (MIP) of a z-stack of images (left panel) and one z-section (right panel)

are shown. See Video 1.

Figure 3: Fate determining transcription factors do not undergo uneven distribution

in telophase. Freshly isolated polyclonal CD8$^+$ T cells were stimulated by immobilized

anti-CD8/anti-CD28/ICAM-1 during 72h and stained with antibodies directed against T-

bet (**A**) or c-Myc (**B**). (**A**): T-bet analysis (n=926 from 3 independent experiments). (**B**): c-

Myc analysis (n=703 from 3 independent experiments).

Numbers highlighted in blue in the plots indicate the % of cells exhibiting asymmetric

repartition of the marker of interest. Red lines indicate the global distribution of the data.

Red numbers indicate the slope of the linear regression curve for marker distribution.

Figure 4: A polarity cue is not necessary for asymmetric repartition of lytic

machinery. (A) Freshly isolated polyclonal CD8[+] T cells were stimulated using

immobilized anti-CD8/anti-CD28/ICAM-1 (left) or with PMA/ionomycin (right) during

72 hours and stained with antibodies directed against CD107a. Each dot represents one

nascent daughter cell. Only one of the two nascent daughter cells in telophase that were

identified by Imaging Flow Cytometry is plotted. The percentage of staining for CD107a

in the presented nascent daughter cell (*x axis*) is plotted against the percentage of staining

for total cell proteins (CTV, *y axis*). Asymmetric cells were defined as in Figure 1. Left:

CD107a analysis when cells were stimulated with immobilized stimuli (n=1185 from 3

independent experiments). Right: CD107a analysis when cells were stimulated with

PMA/ionomycin (n=644 from 3 independent experiments). Numbers highlighted in blue

in the plots indicate the % of cells exhibiting asymmetric repartition of the marker of

interest. Red lines indicate the global distribution of the data. Red numbers indicate the

slope of the linear regression curve for CD107a distribution. **(B)** Histograms represent the

mean and standard deviation of the percentage of asymmetric cells in the 3 independent

experiments. No statistical difference was revealed by paired t-test.

Figure 5: Asymmetric repartition of CD107a$^+$ vesicles reset at each division event.

(A, B) Freshly isolated polyclonal CD8$^+$ T cells were stimulated using immobilized anti-

CD8/anti-CD28/ICAM-1 during 72h and stained with antibodies directed against CD107a.

Cells in telophase were identified by Imaging Flow Cytometry. The number of divisions

accomplished and the cell cycle phase were determined on the basis of CTV and SYTOX

nuclear staining. **(A)** Each dot represents one nascent daughter cell. Only one of the two

nascent daughter cells in telophase that were identified by Imaging Flow Cytometry is

plotted. The percentage of staining for CD107a in the presented nascent daughter cell (*x*

axis) is plotted against the percentage of staining for total cell proteins (CTV, *y axis*).

Asymmetric cells were defined as in Figure 1. Numbers highlighted in blue in the plots

indicate the % of cells exhibiting asymmetric repartition of the marker of interest. Red lines

indicate the global distribution of the data. Red numbers indicate the slope of the linear

regression curve for CD107a distribution. See Figure S3. **(B)** Histograms represent the

mean and standard deviation of the percentage of asymmetric cells in 3 independent

experiments. No statistical difference was revealed by paired t-test.

(C, D) Statistical analysis of cells in telophase and in G1. **(C)** Cells in telophase are plotted

against their CD107a FI. The different curves represent cells having undergone 0, 1 or 2

mitoses. Each dot indicates one cell undergoing asymmetric CD107a repartition as

compared to its CD107a FI. The χ^2 statistical test showed that cells undergoing uneven

repartition of lytic machinery in telophase were randomly distributed all over the CD107a

expression curves (See Materials and Methods section). **(D)** Plots show cells in G1 from

three different experiments. Curves represent the distribution of CD107a florescence

intensity for all cells in G1. Individual plots, marked with different colors, show cells in

G1 at different rounds of division. The Kolmogorov-Smirnov goodness of fit test rejected

the hypothesis that the CD107a expression curves follow the same distribution at the

different division round (See Supplementary Results). The χ^2 test showed that variability

was distributed all over the curves. See Figure S3.

Figure 6: Lytic granules randomly distribute on the two sides of the cleavage furrow

(A and B) Snapshots depict typical cells in division undergoing even (**A**) or uneven (**B**)

repartition of lytic granules (mCherry-tagged GrzB, red) in telophase as detected by live

cell imaging. Images are from Video 2 and 3 respectively. Results are from 3 independent

experiments. **(C and D)** Snapshots depict Imaris software reconstructions of typical cells

undergoing even (**C**) or uneven (**D**) repartition of LTR$^+$ (red) lytic granules in division as

detected by 4D live cell imaging. Images are from Video 4 and 5 respectively. Results are

from 4 independent experiments. See Videos 4-6. (**E**) Binomial modeling for the behavior

of the population of n granules. The curve shows the probability of lytic granule

asymmetric repartition in telophase as a function of lytic granule number. (**F**) Monte-Carlo

simulation of particle correlation as a function of lytic granule number and probability of

lytic granule asymmetric repartition.

Figure 7: CTL expressing high level of lytic granules have better killing capability.

Clonal CTL were FACS-sorted on the basis of their LysoTracker Blue staining. (**A**)

Representative FACS histograms showing LysoTracker Blue staining levels on

LysoTracker^{high} and LysoTracker^{low} sorted-CTL at the indicated day (D) after cell sorting.

Numbers indicate mean fluorescence intensity. Results are representative of 3 independent

998 experiments (**B-C**) LysoTracker^{High} and LysoTracker^{Low} CTL-mediated cytotoxicity was

999 evaluated by FACS analysis by measuring 7-AAD uptake in target cells either pulsed or

1000 not with antigenic peptide following overnight incubation with CTL at the indicated E/T

1001 ratio. (**B**) Cytotoxicity is expressed as the % of 7-AAD$^+$-pulsed target cells minus % of 7-

1002 AAD$^+$-unpulsed target cells (basal). Results are from 3 independent experiments. Each dot

1003 represents results from one experiment performed in triplicate. Means +/- SEM are shown.

1004 Paired t-tests were performed and P-values are indicated. (**C**) Histograms shown are from

1005 one representative experiment. Numbers indicate the percentage of 7-AAD positive target

1006 cells. (**D**) LysoTracker^{High} and LysoTracker^{Low} CTL CD107a exposure after a 4 hours

1007 incubation with target cells pulsed or not with antigenic peptide (E/T ratio 0.5:1) was

1008 evaluated by FACS analysis. Each dot represents results from 4 independent experiments

1009 performed either in duplicate or triplicate. Means +/- SEM are shown. Paired t-tests were

1010 performed and P-values are indicated. (**E**) CD8 expression in LysoTracker^{High} and

1011 LysoTracker^{Low} CTL after a 4 hours incubation with target cells pulsed with antigenic

1012 peptide (E/T ratio : 0.5:1) was evaluated by FACS analysis. Results are normalized on CD8

1013 MFI level of LysoTracker^{High} and LysoTracker^{Low} CTL cultured in the absence of target

1014 cells. Each dot represents results from 4 independent experiments performed either in

1015 duplicate or triplicate. Means +/- SEM are shown.

 Supplementary figure legends

 Figure 1-supplement 1 : Gating strategy for Imaging Flow Cytometry (IsX)

 acquisition. Based on the brightfield illumination, all events were plotted for their aspect

 ratio (length/width, equal 1 for perfectly round cells) and their area. Cells in telophase were

 defined as those exhibiting a low aspect ratio and a big area. The region of interest (gray)

 included cell doublets and cells in anaphase and telophase. Based on the intensity of DNA

 staining (represented in linear axis) cells in G2/M were selected. We then applied a mask

 on the IsX image gallery (as described in material and methods section) to define the limits

 of the nuclei. This strategy was used to determine the number of nuclei present in each

 gated cell. To unambiguously identify cells in telophase we applied a mask on α-tubulin

 staining allowing to detect condensed microtubules in an elongated shape (as described in

 material and methods section). This procedure allowed us to detect the midbody (a

 structure characteristic of telophase formed by highly condensed α-tubulin that bridges the

 2 nascent daughter cells). Cells included the described gates were finally visually

 inspected. All the cells recognized as in telophases on the basis of nuclear and tubulin

 staining were included in the analysis of the markers of interest.

 Figure 1-supplement 2 : Analysis and representation of the repartition of markers of

 interest in dividing cells (A) Analysis of individual cells in telophase. All IsX generated

 TIFF files were analyzed using the Fiji software. For each telophase cell we used 3 TIFF

 image corresponding to: i) CTV staining; ii) α-tubulin staining iii) and marker of interest.

 To standardize analysis, we used macro programming on Fiji (described in supplementary

 results section). To determine a rupture zone between the 2 nascent daughter cells we

1039   applied watershed function on tubulin mask. The watershed masks were used to determine

1040   the 2 nascent daughter cells in which the fluorescence intensities of CTV and of the markers

1041   of interest were measured (yellow lines). **(B)** Example of a cell exhibiting asymmetric

1042   distribution in telophase of a marker of interest. The yellow lines highlight the nascent

1043   daughter cell exhibiting a higher content of the marker of interest.

1044

Figure 1-supplement 3 : CD8$^+$ T cells are efficiently stimulated on coated anti-

CD3/anti-CD28/ICAM1. Freshly isolated polyclonal CD8$^+$ T cells previously stained

with CTV were stimulated 72 hours using immobilized anti-CD3/anti-CD28/ICAM1. (**A**)

Imaging Flow Cytometry shows that stimulated cell undergo several rounds of division as

shown by CTV staining dilution. (**B**) Flow cytometry shows upregulation of CD137

expression in stimulated cells.

1051

Figure 1-supplement 4 : Uneven lytic granule segregation in telophase in CD8$^+$

memory T cells. The panels show staining for CD107a and perforin in human CD8$^+$

memory T cells stimulated and analyzed as **in Figure 1A-B**. CD107a n=978 from 3

independent experiments; perforin n=1127 from 3 independent experiments. Numbers

highlighted in blue in the plots indicate the % of cells exhibiting asymmetric repartition of

the marker of interest. Red lines indicate the global distribution of the data. Red numbers

indicate the slope of the linear regression curve for marker distribution.

1059

1060   **Figure 6-supplement 1 : Lysotracker randomly distribute on the two sides of the**

1061   **cleavage furrow**

1062 **(A and B)** Snapshots depict Imaris software reconstructions of typical cells undergoing

1063 uneven **(A)** and even **(B)** repartition of LTR$^+$ (red) lytic granules in division as detected by

1064 4D live cell imaging. Images are from Video 6.

1065

1066 **Video legends**

1067 **Video 1: 3D visualization of CD107a repartition in a telaphasic CD8$^+$ T cell** The video

1068 shows 3D reconstruction of a cell in telophase. CD107a (green), α-tubulin (red) and DAPI

1069 (cyan). The images presented in **Figure 2B** has been extracted from this video.

1070

Video 2 and 3: Visualization by time-lapse confocal laser scanning microscopy of cell

division. Human CTL were transfected during their expansion phase with mcherryGrzB.

18 hours after transfection, cells were inspected by time-lapse laser scanning confocal

microscopy for additional 5-6 hours using a Tile Scan mode to enlarge the acquisition filed

and to capture rare cells undergoing spontaneous division during the time of acquisition.

Video 2 shows a typical cell undergoing even repartition of GrzB$^+$ granules. Video 3 shows

a typical cell undergoing uneven repartition of GrzB$^+$ granules. Snapshots of Video 2 and

3 are shown in **Figure 6 A** and **B.**

1079

Video 4-6: Visualization by 4D time-lapse microscopy of cell division. G2/M sorted

CTL were loaded with Hoechst (blue) and LysoTracker Red (LTR, red) and inspected by

time-lapse laser scanning confocal microscopy (video 4-5) or spinning-disk microscopy

(video 6) for 12h-16h. **Video 4** shows 4D reconstruction (using Imaris software) of a

typical cell undergoing even repartition of LTR$^+$ granules. **Video 5** shows 4D

reconstruction (using Imaris software) of a typical cell undergoing uneven repartition of

LTR$^+$ granules. Snapshots of Video 4 and 5 are shown in **Figure 6 C** and **D. Video 6** shows

4D reconstruction (using Imaris software) of one typical cell undergoing uneven repartition

of LTR$^+$ granules and one typical cell undergoing even repartition of LTR$^+$ granules.

Snapshots of Video 6 are shown in **Figure 6-figure supplement 1.** Results are from 4

independent experiments.

Figure 1

Figure 2

A

fold increase

3.0
2.5
2.0
1.5
1.0

27%

telophase

B

MIP

Z-section

5μm

DAPI   Tubulin   CD107a

Figure 3

A

B

Figure 4

A

polarized stimulation

unpolarized stimulation

B

Figure 5

Figure 6

A Symmetric
0:00:00 0:43:50 0:52:48

B Asymmetric
0:00:00 0:38:11 0:43:50

C Symmetric
1:01:07 1:11:54 1:13:42

D Asymmetric
1:28:05 1:33:28 1:37:04

E

Probability of asymmetric repartition

Total number of granules in each cell

F

Probability of asymmetric repartition

Particle correlation during division

n particles

Figure 7

A

B

C

D

E

Figure 1- figure supplement 1

Figure 1- figure supplement 2

A

B

Figure 1- figure supplement 3

A

B

Time : 0 hour

Time : 72 hours

Figure 1- figure supplement 4

Figure 6- figure supplement 1

A

B