Distinguishing between interaction and dispersion effects in the analysis of gene-environment interaction

Benjamin W. Domingue[1,†], Klint Kanopka[2], Travis T. Mallard[3], Sam Trejo[4], and
Elliot M. Tucker-Drob[5,†]

[1]Graduate School of Education, Stanford University & Center for Population Health Sciences, Stanford
Medicine; bdomingu@stanford.edu
[2]Graduate School of Education, Stanford University; kkanopka@stanford.edu
[3]Department of Psychology, University of Texas at Austin; travis.mallard@utexas.edu
[4]La Follette School of Public Affairs & Department of Sociology, University of Wisconsin–Madison;
sam.trejo@wisc.edu
[5]Department of Psychology & Population Research Center, University of Texas at Austin;
tuckerdrob@utexas.edu
[†]Correspondence about the paper should be sent to these authors.

Abstract

Genotype-by-environment interaction (GxE) occurs when the size of a genetic effect varies
systematically across levels of the environment and when the size of an environmental effect
varies systematically across levels of the genotype. However, total variance in the phenotype
may shift as a function of the moderator irrespective of its etiology such that the *proportional*
effect of the predictor is constant. We expand the traditional GxE regression model to directly
account for heteroscedasticity associated with both the genotype and the measured environ-
ment. We then derive a test statistic, ξ, for inferring whether GxE can be attributed to an
effect of the moderator on the dispersion of the phenotype. We apply this method to identify
genotype-by-birth year interactions for Body Mass Index (BMI) that are distinguishable from
general secular increases in the variance of BMI or associations of the genetic predictors (both
PGS and individual loci) with BMI variance. We provide software for analyzing such models.

Key Words: gene-by-environment interaction, gene $\times$ environment interaction, GxE, G$\times$E,
vQTL, heteroscedasticity.

1   Introduction

Genotype-by-Environment interaction (GxE) studies test whether the genotype-phenotype associ-ation varies in magnitude across the range of a measured environmental variable and whether the environment-phenotype association varies in magnitude across the range of the measured genotype. Investigations of GxE have been of particular interest in the study of complex traits, leading to a variety of methods for estimating polygenic GxE [1, 2]. Development of these analytic approaches coincides with a large increase in the availability of genomic data, particularly in the context of population-based studies that also contain a broad range of phenotypes and environmental mea-sures. Here, we consider circumstances where either the genetic or the environmental measure selected for GxE is associated with the magnitude of variance in the target phenotype. Such het-eroscedasticity introduces a key interpretational problem to results from standard GxE approaches: any observed GxE may arise as an artifact of the fact that the total variance in the phenotype shifts across the range of the genetic or environmental moderator, irrespective of etiology. We resolve this problem by expanding the standard regression-based model for testing GxE in both single locus and polygenic score settings by introducing a simple yet flexible approach to directly model heteroscedastic residuals, and we develop a test statistic for inferring whether GxE can be plausibly attributed to a more general effect of the moderator on the dispersion of the phenotype.

All else equal, when the dispersion of a phenotype is directly controlled by the level of the moderator, the standard regression-based model will identify a significant interaction between that moderator and *any* correlate of the phenotype, whether genetic or non-genetic. For example, when the effectiveness, intensity, or dosage of a phenotype-altering intervention varies proportionally to baseline levels of the phenotype (e.g. when hours of individualized education are assigned based on previous year's test scores; or when intensity of a weight-loss intervention is determined from baseline body mass index), then the variance of the phenotype is expected to increase in response to the intervention, and the unstandardized effect sizes for all correlates of baseline levels of the phenotype are expected to increase. Similarly, when a genetic variant confers greater plasticity in a phenotype, then the variance of the phenotype is expected to differ across alleles (such that the variant is identified as a variance quantitative trait locus, vQTL). We would then expect the variant to moderate the effect sizes for all correlates of that phenotype proportionally to one another. Only when a moderator acts preferentially on specific mechanisms of variation in the phenotype are interactions with various predictors expected to depart from expectations that follow from differences in total variance across the range of the moderator.

Figure 1 illustrates how heteroscedasticity across the range of a measured environmental moderator (E; left panel) or genetic moderator (G; right panel) can produce the impression of GxE. In the left panel, we illustrate a scenario in which a PGS is associated with a constant proportion of variance in the phenotype across the range of the measured environment. That the proportion of variance explained by the PGS at any given location on the x-axis is constant indicates that the expected percentile location within the distribution of the phenotype will also be constant across the range of E (assuming that the shape, but not dispersion, of the distribution remains constant across the range of E). Nevertheless, a clear fan-spreading effect is observed, incorrectly implying that the PGS is increasingly penetrant at higher levels of E. The right panel represents an analogous scenario with respect to a single genetic variant that moderates the variance of the phenotype (i.e a vQTL [3–7]). In this scenario, the environment is associated with a constant proportion of variance in the phenotype across the range of the genotype, but appears to become increasingly predictive of the phenotype as the effect allele count increases as a result of heteroscedasticity. By directly modeling this heteroscedasticity, we are able to distinguish between scenarios such as these—in which GxE is an epiphenomenon of variance modulation—from those representing more meaningful patterns of GxE.

Figure 1: Hypothetical examples of an Environmental moderator (left) or a genetic moderator (right) of total variance in a phenotype producing the impression of GxE. Left: Residuals of the regression of the Phenotype on the Measured Environment are heteroscedastic. high, average, and low polygenic scores (PGSs) are associated with constant percentiles of the phenotype at any given location along the x axis. However, because variance of the phenotype expands across the range of the Measured Environment, the PGS accounts for increasing unstandardized variance across this range. Right: A variance quantitative trait locus (vQTL) in which the effect allele is associated with greater variance in the phenotype. Unstandardized scores on the phenotype that are associated with high, average, and low levels of the measured environment (E) become more distinct with increasing number of effect alleles. However, because the total variance in the phenotype expands across the x axis, the *percentile* locations of these scores within each genotype (0, 1, or 2) is constant across all levels of the genotype.

We develop and validate a flexible heteroscedastic regression model for testing GxE. We use this model to elucidate the well-studied interaction between birth cohort and genetics linked to body mass index (BMI). We test whether both PGS-by-Birth Year and SNP-by-Birth Year interactions on BMI can be attributed to more general increases in the total dispersion of BMI over historical time. That is, we ask whether birth year is associated specifically with amplification of genetic risk for BMI or more generally with amplification of all variation in BMI.

2 Methods

In settings wherein focus is on a measured genotype, as indexed by a single-locus allele count or a polygenic score (PGS), the standard regression model for GxE is

$$Y_i = \beta_0 + \beta_1 G_i + \beta_2 E_i + \beta_3 G_i \cdot E_i + e_i, \tag{1}$$

where Y_i is the phenotype for person i, G_i is the measured genotype, E_i is the measured environment, and e_i is an error term. Under this model, the effect of G_i on Y_i is allowed to vary as a linear function of E_i, as indexed by the regression coefficient β_3. However, a crucial assumption is that the residuals, e_i, are homoscedastic across all levels of E_i (e.g., $e_i \sim \mathrm{N}(0, \sigma_\epsilon^2)$).

The standard regression model given by Eqn 1 can be expanded so as to relax the assumption of homoscedasticity of residuals across E_i by specifying the following *environmental heteroscedasticity model*,

$$Y_i = \tau_0 + \tau_1 E_i + \pi_0 G_i + \pi_1 G_i \cdot E_i + \lambda_0 \epsilon_i + \lambda_1 E_i \cdot \epsilon_i. \tag{2}$$

Here, τ_1 is the main effect of the measured environment, π_0 is the main effect of G_i, π_1 is the GxE analogue to β_3 in Eqn 1, λ_0 is the main effect of the error ϵ_i (which we assume, without loss of generality given λ_0, to have unit variance), and λ_1 is the coefficient used to index heteroscedasticity. Rather than simply adjusting standard errors and p values of the standard regression model (Eqn 1) using a robust estimation method, this model explicitly estimates heteroscedasticity as a model parameter. As we articulate next, directly modelling heteroscedasticity allows us to overcome specification bias that would result from estimating a standard homoscedastic GxE model, even with a robust estimator. We are purposeful in choosing a function form for the heteroscedasticity term, $E_i \cdot \epsilon_i$, that directly parallels that specified for the GxE term, $G_i \cdot E_i$.

The heteroscedasticity model is a flexible model allowing for both conventional GxE and heteroscedasticity. In order to test whether a more general pattern of heteroscedasticity as a function

of E_i is sufficient to account for the observed GxE, we propose a restricted form version of this model that does not directly model GxE but instead allows for the total variance of the phenotype to vary as a function of E_i. This *scaling model* represents a scenario wherein GxE—in the sense of a significant estimate of β_3 in Eqn 1—arises as a function of differences in the total variance of Y_i as a function of E_i. We choose this nomenclature given that the model emphasizes the importance of the *scale* of the outcome. Under this model, the proportional contribution of G_i is constant over the range of E_i, but the scale of Y_i systematically varies across the range of E_i; i.e., E_i acts as a "dimmer" [8].

The scaling model for moderation of the variance of Y_i as a function of E_i takes the form

$$Y_i = a_0 + a_1 E_i + (b_0 + b_1 E_i) Y_i^\star. \tag{3}$$

Here, $Y_i^\star$ is an unobserved factor representing unexplained variation in Y_i incremental of E_i. The $b_0 + b_1 E_i$ coefficient on the $Y^\star$ term produces heteroscedasticity in Y_i as a function of E_i. We let

$$Y_i^\star = h G_i + e \epsilon_i \tag{4}$$

where G is a measured genotype standardized to have unit variance and ϵ_i is an unobserved error term (we assume it is also scaled to have unit variance). Finally, we identify the units of $Y^\star$ by specifying $e = \sqrt{1 - h^2}$. The penetrance of the measured genotype is thus controlled via the relative magnitudes of h and e. Note, in particular, that in terms of $Y_i^\star$, the penetrance of the measured genotype is constant. The test that we introduce relies on this property; i.e., when properly scaled, the penetrance of the measured genotype to Y_i is constant. Suppose that $b_1 = 0$. In that case, Y_i is affected solely by environmental and genetic main effects. If $b_1 \neq 0$, then the role of $Y_i^\star$ with respect to Y_i varies as a function of E, and as we elucidate next, so do the raw (but not relative) contributions of G_i and ϵ_i to Y.

In Section A of the supplemental information (SI), we show that the environmental heteroscedasticity model reduces to the environmental scaling model when

$$\frac{\pi_0}{\lambda_0} = \frac{\pi_1}{\lambda_1}. \tag{5}$$

We use this observation to derive the test statistic ξ_E and associated hypothesis test of whether

an empirical estimate of GxE is distinguishable from the scaling model as

$$\xi_E \equiv \widehat{\pi_0}\widehat{\lambda_1} - \widehat{\pi_1}\widehat{\lambda_0}, \tag{6}$$

$$H_0 : \xi_E = 0. \tag{7}$$

We provide parallel heteroscedasticity and scaling models for moderation of the variance of Y_i as a function of G_i (leading to an analogous test statistic, ξ_G) in Section A.3 of SI.

We conduct a variety of simulation studies (see Section C of SI). In summary, results of these simulation studies suggest that ξ_E is a reliable indicator of whether the scaling model is the basis for GxE. When a test of the statistic fails to reject $H_0 : \xi_E = 0$ and there is significant GxE (π_1) observed, we cannot rule out that the scaling model is driving observed GxE. When the test suggests rejection of $H_0 : \xi_E = 0$, alternative forms of GxE are implicated. Software to estimate the models considered here is described in Section D of SI.

3 Heteroscedasticity in context of BMI-linked genetics and birth year

We apply our heteroscedastic GxE model to the well-studied interaction between birth year and genetics linked to body mass index (BMI) [9–12], a literal textbook example (see Section 11.3 of [13]). We test whether previous reports of increasing penetrance of PGS for BMI for later-born birth cohorts can be plausibly attributed to more general increases in the total dispersion of BMI, including variation unique of its genetic etiology, across birth cohorts [14, 15]. We thus ask whether birth year is associated specifically with amplification of genetic risk for BMI or more more generally with amplification of all variation in BMI. We then go on to apply our heteroscedastic GxE model to examine individual locus-by-year interactions for BMI. To illustrate how our approach can be used to conduct SNP-level analyses, we apply our technique using the top hits from a recent BMI GWAS [16] to investigate whether they are associated with heteroscedastic GxE.

3.1 Polygenic Score Analysis

We consider GxE in the context of a long-running cohort study—the Health and Retirement Study (HRS) [17]—of US adults over 50 and their spouses (N=11,586). Respondents are interviewed every two years and data are collected on their early-life, anthropometrics, socioeconomic status, and

mental and physical well-being. Previous examinations of increases in the penetrance of the BMI polygenic score across birth year have used this data [9–11]. We use a polygenic score for BMI [16] as constructed by the HRS [18]; further description of the data are in E of SI.

Results are presented in Table 2. We observe considerable evidence for stronger prediction of BMI by the PGS for more recently born individuals (i.e., $\pi_1 > 0$); this is consistent with earlier findings indicating secular increases in the penetrance of a BMI PGS [9–11]. However, we also observe a birth year-linked increase in non-PGS variance in BMI (i.e., $\lambda_1 > 0$), raising the possibility that the obtained GxE may be attributable to a more general pattern of increasing variance in BMI with birth year. In sex-pooled analyses, probabilities associated with tests of $H_0 : \xi = 0$ are approximately 0.02. However, findings vary slightly by gender; especially for males, we are unable to reject the null of the scaling model. To illustrate the differences in implications in our sex-stratified analyses, we also conduct a posterior predictive check analysis [19], see Figure 2; there is less change in the penetrance of BMI PGS across birth years for males as compared to females. Using Eqn 43, we can decompose variation in π_1 estimates. For females and males, 41% (95% CI: 10-57%) and 25% (95% CI: 0-54%) of the GxE (π_1) effect cannot be attributed to a simple scaling term. In an analysis of all respondents, this proportion is 37% (95% CI: 9-52%). Collectively these results indicate substantial evidence for heteroscedasticity as a function of birth year; the simple homoscedastic GxE model does not represent the data adequately. Further discussion of this analysis and findings related to genetic heteroscedasticity are discussed in the SI.

Table 1: Estimates from parameters of the full heteroscedasticity model (Eqn 49) in analysis of GxE for Box-Cox transformed BMI as a function of birthyear in the UK Biobank for those SNPs with significant π_1 estimates following Bonferonni adjustment. The ξ_E and ξ_G estimates reported are obtained from the environmental and genetic heteroscedasticity models, respectively. We show probabilities for parameters when the maximal probability in a column is larger than $1e-6$.

SNP	π_0	π_1	$\Pr_{\pi_1}$	λ_0	λ_1	λ_2	$\Pr_{\lambda_2}$	$\Pr_{\xi_E}$	$\Pr_{\xi_G}$
rs543874_A	-0.0271	-0.0085	1.90e-07	0.9873	0.0369	-0.0036	1.48e-03	4.39e-06	1.04e-07
rs13021737_G	0.0279	0.0062	1.33e-04	0.9873	0.0369	0.0036	1.53e-03	1.46e-03	1.75e-04
rs1558902_T	-0.0503	-0.0077	2.12e-06	0.9864	0.0368	-0.0096	1.76e-17	2.96e-04	5.14e-07
rs2075650_A	0.0113	-0.0060	2.43e-04	0.9875	0.0371	-0.0002	8.53e-01	2.01e-04	5.78e-04

3.2 SNP Analyses

We conducted heteroscedastic regression analyses for 96 marker SNPs for the genome-wide significant loci identified in [16] using independent data from N=380,605 participants from UK Biobank

Table 2: Estimates from parameters of the full heteroscedasticity model (Eqn 49) in analysis of GxE for BMI as a function of birth year in the HRS. The ξ_E and ξ_G estimates reported are obtained from the environmental and genetic heteroscedasticity models, respectively. We show probabilities for parameters when the maximal probability in a column is larger than $1e-6$.

Gender	BMI[a]	N	τ_1	π_0	π_1	$\Pr_{\pi_1}$	λ_0	λ_1	λ_2	$\Pr_{\lambda_2}$	$\Pr_{\xi_E}$	$\Pr_{\xi_G}$
All	Std	11586	0.19	0.25	0.06	1.78e-11	0.93	0.12	0.09	7.44e-50	1.76e-02	7.89e-04
All	Trans	11586	0.17	0.26	0.04	1.79e-05	0.95	0.06	0.02	1.14e-03	2.68e-02	5.15e-04
M	Std	5022	0.19	0.26	0.04	4.69e-04	0.93	0.12	0.09	1.63e-22	3.91e-01	2.75e-01
M	Trans	5022	0.17	0.26	0.03	2.11e-02	0.95	0.07	0.03	1.33e-03	3.83e-01	1.65e-01
F	Std	6564	0.20	0.25	0.06	6.65e-09	0.93	0.12	0.09	3.52e-31	2.24e-02	1.21e-03
F	Trans	6564	0.18	0.26	0.04	2.08e-04	0.94	0.06	0.02	1.11e-02	3.82e-02	1.75e-03

[a] Std denotes the standardized mean BMI shown in the middle panel of Figure E.1. Trans denotes the Box-Cox transformation.

(UKB; not used in the BMI GWAS [16]). Replication results for the main effects can be found in Table F.2. Note that the UKB is highly non-representative [20] potentially affecting the generalizability of our results.

The environmental heteroscedasticity parameter (λ_1) was positive and significant in all models, suggesting greater variance in BMI among those born later in the 20th century. With the full heteroscedasticity model, we identified four SNPs that exhibited significant gene-by-birth year effects, as indicated by the π_1 parameter at the Bonferroni-adjusted alpha level. Parameter estimates from the full heteroscedasticity model for these SNPs are reported in Table 1, along with ξ_E and ξ_G estimates obtained from the environmental and genetic heteroscedasticity models, respectively. Note that rs1558902 is a variant in the FTO gene locus. This finding replicates earlier results [21] which reported increasing penetrance of a variant with FTO over historical time using a different data set. However, ξ_G was highly significant for each of these four SNPs, allowing us to reject the null that observed GxE is driven entirely by scaling.

With the full heteroscedasticity model, we identified six SNPs that exhibited significant effects on the dispersion of BMI, as indicated by the λ_2 parameter at the Bonferroni adjusted alpha level. Parameter estimates for these SNPs are reported in the supplement (Table F.1), along with ξ_E and ξ_G estimates obtained from the environmental and genetic heteroscedasticity models, respectively. Of the 4 Bonferonni significant variants for SNP-by-Year interaction described above, only the FTO variant (rs1558902) appears among the 6 Bonferonni significant variants for vQTL. This finding replicates earlier results [3] reporting that the FTO gene locus is associated with greater variability in BMI. The p value of ξ_G for this variant is 5.1e-07, indicating the the above-reported SNP-by-birth-year interaction on BMI for FTO is not simply attributable to a general modulation of variance by rs1558902. Note that although the ξ_G statistic is not significant for

Figure 2: Gray lines represent genetic penetrance as a function of birth year simulated based on either the scaling model (Eqn 13, on left) or the standard homoscedastic GxE model (Eqn 1, on right). Red lines represent genetic penetrance as a function of birth year estimated from the real data using the heteroscedastic regression model (Eqn 8). Analyses based on standardized BMI data.

rs7903146, rs9925964, and rs2287019, these SNPs do not exhibit GxE at Bonferonni-corrected levels, rendering the ξ_G moot.

To examine the total evidence of signal for GxE and genetic heteroscedasticity in the 96 SNPs, we provide a QQ plot of -1og10(p) for parameters indexing SNP-controlled heteroscedasticity of BMI vQTL and gene-by-birth year GxE interaction for Box-Cox transformed BMI in the left panel of Figure 3. We observe considerable departure of the observed -1og10(p) values relative to the expectation under the null, indicating strong enrichment of signal for both vQTL and GxE. In the right panel of Figure 3, we provide the scatter plots of parameter estimates for SNP main effects ($\pi0$), GxE (π_1), for vQTL (λ_2) for Box-Cox transformed BMI. We observe moderate-to-strong associations between the main effect of a SNP and both its GxE and vQTL effects, indicating that, among these variants selected on the basis of marking genome-wide significant loci for BMI in an external meta-analysis, those with larger main effects on BMI exhibit larger interactions with

¹⁹³ birth year and more pronounced moderation of BMI variance. Results of [6] suggest that similar

¹⁹⁴ patterns may apply genome-wide.

Figure 3: Left: QQ plot of -1og10(p) for parameters indexing SNP-by-birth year interaction (GxE, as indicated by the π_1 parameter) and SNP-controlled heteroscedasticity of BMI (vQTL, as indicated by the λ_2 parameter) for Box-Cox transformed BMI. The considerable departure of the observed -1og10(p) values relative to the expectation under the null indicates strong enrichment of signal for both vQTL and GxE in these 96 SNPs previously associated with mean BMI in independent data [16]. Right: Multipanel Scatterplot of the parameters for the SNP main effect (π_0), SNP-by-Birth Year interaction (π_1), and vQTL (λ_2) for Box-Cox transformed BMI. We analyzed 96 marker SNPs for the genome-wide significant loci for BMI identified in [16] in independent data.

¹⁹⁵ Our results indicate that SNP moderation of birth year effects on BMI and SNP moderation

¹⁹⁶ of BMI dispersion were in the same direction as the SNP effects on BMI means. We observe

¹⁹⁷ considerable sign congruence between SNP main effects (π_0) and SNP-by-Birth-Year interaction

¹⁹⁸ (π_0) for 3 out of 4 SNPs exhibiting Bonferonni significant π_1 estimates and for 71 out of 96 of all

¹⁹⁹ SNPs tested. We also observe considerable sign congruence between SNP main effects (π_0) and

²⁰⁰ vQTL effects (λ_2) for 6 out of 6 SNPs exhibiting Bonferonni significant λ_2 estimates, and for 75

²⁰¹ out of 96 of all SNPs tested.

²⁰² Finally, we note that the main effect of birth year (τ_1) was approximately -0.058 in all models

²⁰³ (all associated probabilities less than 1e-250), indicating that later-born UK Biobank participants

²⁰⁴ tend to have lower BMI. Given substantial epidemiological evidence from representative samples

²⁰⁵ indicating increasing BMI with birth year, we speculate that this negative association may be

²⁰⁶ driven by selection bias in the UKB sample [20]. However, note that, despite decreasing mean

²⁰⁷ BMI with birth year in this sample, we observe increasing variance in BMI with birth year (as

²⁰⁸ indicated by the positive λ_1 parameter), and increasing penetrance of SNPs with birth year (as

indicated by strong sign congruence between π_0 and π_1 parameters). This further suggests that variance moderation and SNP-by-Birth Year interactions are unlikely to have resulted from positive associations between BMI mean and variance.

4   Discussion

When GxE is detected using the conventional regression-based model, it is common to interpret the interaction in terms of the specific genetic and environmental measures included in the interaction model. However, GxE may arise more generally when the genetic or environmental predictor moderates the total amount of variation in the phenotype. When the total amount of variation in a phenotype varies across the range of an environmental measure, traditional homoscedastic GxE models will detect interactions between that measure and all other correlates of the phenotype, both genetic and non-genetic. Here, we have delineated an expanded *heteroscedastic* GxE regression model that explicitly allows for heteroscedasticity in the phenotype as a function of the genetic and environmental measures. We use this model to derive a test statistic, ξ, which compares this heteroscedasticity model to a simpler *scaling model* in which GxE arises from more general differences in phenotypic variance, irrespective of etiology. When the scaling model holds, differences in phenotypic variance across the range of the genetic or environmental moderator induce a form of GxE that is only apparent in unstandardized units; the proportional contribution of the remaining predictors to phenotypic variance remains constant across the range of the moderator.

We provide an application of the heteroscedastic regression model to empirical data, replicating previous observations that that genetic predictors of BMI have become increasingly penetrant in recent years. However, we additionally observed that the total variance in BMI, including variation not associated with the genetic predictors, increased with participant birthyear. We found that, amongst males within the sample, the interaction between PGS and birth year was indistinguishable from a scaling model in which the total variance in BMI increases over birth year. For males, although the unstandardized variance in BMI accounted for by the PGS increased with birth year, the variance in BMI unaccounted for by the PGS also increased, such that the proportion of variance accounted for in BMI did not significantly vary with birth year. While we reject the null that the scaling model produced data for females, we still estimate that a large proportion of the observed GxE is due to scaling. Other work has considered the potential moderation of genetic risk for BMI as a function of diet [22, 23], other lifestyle characteristics [24], and educational attainment [25]. The issues discussed here may be a relevant feature of such empirical work; i.e.,

environmentally-linked heteroscedasticity may play a role in driving some previously observed GxE findings.

In instances in which the units and scaling of the phenotype are intrinsically-meaningful, biologically-meaningful, or policy-relevant, GxE that is observed on the unstandardized scale may be of high substantive and practical relevance. However, even when GxE arising from heteroscedasticity is deemed meaningful, it may still be important to discern whether interactions are specific to the variables selected for GxE or representative of a more general pattern that would be obtained for many correlates of the phenotype. It will also stil be important to understand whether the moderator is associated with differences in the percentile location of expected values along the distribution of the phenotype, or simply with differences in raw values (e.g. has a policy specifically increased the income of a particular subgroup, or increased the income of all subgroups, such that relative income of the subgroup of interest remains constant?). The heterosedastic regression models introduced here can be used to answer these questions.

Issues surrounding the scaling of the phenotype not fully explored here have the potential to have substantive effects on GxE results. For instance, skew in the distribution of the phenotype may produce both GxE and heteroscedasticity by inducing a dependency between the mean and the variance of the distribution of the phenotype. Such forms of GxE and may altogether dissappear under the appropriate tranformation, an issue that geneticists have be aware of since the days of Fischer [26]. Importantly, heteroscedasticity is not always simply an epiphenomenon of non-normality. For instance, in our empirical analysis of BMI, we report results of GxE analyses after a normalizing (Box-Cox) transform of BMI, and continue to detect both GxE and heteroscedasticity.

GxE research faces a number of conceptual and statistical challenges [8]. Thoughtfully and rigorously engaging with these challenges is particularly salient given the substantial recent increases in the availability of both genetic resources [27] and computational tools [28] for such research. Our proposed test is designed to help clarify the nature of potential GxE findings and to shed additional light on the processes contributing to variation in the phenotype; in particular, we can assess the extent to which the genetic etiology is relatively stable across the range of the environmental measure. Using the model and test statistic presented here, researchers can test for whether variance in the phenotype that is not explained by the genetic predictor shifts across the range of the environment and whether a scaling model may account for the obtained pattern of GxE. When the scaling model holds, we would suggest that the environment largely acts as a "dimming" mechanism on phenotypic variation [8] without altering the proportional contribution of genetic variation to the phenotype.

References

1. Purcell, S. Variance components models for gene–environment interaction in twin analysis. *Twin Research and Human Genetics* **5,** 554–571 (2002).

2. Ni, G. *et al.* Genotype–covariate correlation and interaction disentangled by a whole-genome multivariate reaction norm model. *Nature communications* **10,** 1–15 (2019).

3. Yang, J. *et al.* FTO genotype is associated with phenotypic variability of body mass index. *Nature* **490,** 267–272 (2012).

4. Wang, H. *et al.* Genotype-by-environment interactions inferred from genetic effects on phenotypic variability in the UK Biobank. *Science advances* **5,** eaaw3538 (2019).

5. Conley, D. *et al.* A sibling method for identifying vQTLs. *PloS one* **13,** e0194541 (2018).

6. Young, A. I., Wauthier, F. L. & Donnelly, P. Identifying loci affecting trait variability and detecting interactions in genome-wide association studies. *Nature genetics* **50,** 1608–1614 (2018).

7. Marderstein, A. R. *et al.* Leveraging phenotypic variability to identify genetic interactions in human phenotypes. *bioRxiv* (2020).

8. Domingue, B., Trejo, S., Armstrong-Carter, E. & Tucker-Drob, E. M. Interactions between polygenic scores and environments: Methodological and conceptual challenges. *Sociological Science* (In press).

9. Conley, D., Laidley, T. M., Boardman, J. D. & Domingue, B. W. Changing Polygenic Penetrance on Phenotypes in the 20 th Century among Adults in the US Population. *Scientific reports* **6,** 30348 (2016).

10. Liu, H. & Guo, G. Lifetime socioeconomic status, historical context, and genetic inheritance in shaping body mass in middle and late adulthood. *American sociological review* **80,** 705–737 (2015).

11. Walter, S., Mejıa-Guevara, I., Estrada, K., Liu, S. Y. & Glymour, M. M. Association of a genetic risk score with body mass index across different birth cohorts. *Jama* **316,** 63–69 (2016).

12. Demerath, E. W. *et al.* The positive association of obesity variants with adulthood adiposity strengthens over an 80-year period: a gene-by-birth year interaction. *Human heredity* **75,** 175–185 (2013).

13. Mills, M. C., Barban, N. & Tropf, F. C. *An Introduction to Statistical Genetic Data Analysis* (MIT Press, 2020).

14. Flegal, K. M., Kruszon-Moran, D., Carroll, M. D., Fryar, C. D. & Ogden, C. L. Trends in obesity among adults in the United States, 2005 to 2014. *Jama* **315,** 2284–2291 (2016).

15. Ogden, C. L. *et al.* Trends in Obesity Prevalence by Race and Hispanic Origin—1999-2000 to 2017-2018. *JAMA.*

16. Locke, A. E. *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature* **518,** 197–206 (2015).

17. Juster, F. T. & Suzman, R. An overview of the Health and Retirement Study. *Journal of Human Resources,* S7–S56 (1995).

18. Ware, E., Schmitz, L., Gard, A. & Faul, J. HRS Polygenic Scores—Release 3: 2006–2012 Genetic Data. *Ann Arbor: Survey Research Center, University of Michigan* (2018).

19. Gelman, A., Meng, X.-L. & Stern, H. Posterior predictive assessment of model fitness via realized discrepancies. *Statistica sinica,* 733–760 (1996).

20. Fry, A. *et al.* Comparison of sociodemographic and health-related characteristics of UK Biobank participants with those of the general population. *American journal of epidemiology* **186,** 1026–1034 (2017).

21. Rosenquist, J. N. *et al.* Cohort of birth modifies the association between FTO genotype and BMI. *Proceedings of the National Academy of Sciences* **112,** 354–359 (2015).

22. Qi, Q. *et al.* Sugar-sweetened beverages and genetic risk of obesity. *New England Journal of Medicine* **367,** 1387–1396 (2012).

23. Qi, Q. *et al.* Fried food consumption, genetic risk, and body mass index: gene-diet interaction analysis in three US cohort studies. *Bmj* **348,** g1610 (2014).

24. Fang, J. *et al.* Polygenic risk, adherence to a healthy lifestyle, and childhood obesity. *Pediatric Obesity* **14** (2019).

25. Komulainen, K. *et al.* Education as a moderator of genetic risk for higher body mass index: prospective cohort study from childhood to adulthood. *International Journal of Obesity* **42,** 866–871 (2018).

26. Tabery, J. Biometric and developmental gene–environment interactions: Looking back, moving forward. *Development and Psychopathology* **19,** 961–976 (2007).

27. Lambert, S. A. *et al.* The Polygenic Score Catalog: an open database for reproducibility and systematic evaluation. *medRxiv* (2020). 333 334

28. Shin, J. & Lee, S. H. GxEsum: genotype-by-environment interaction model based on summary statistics. *BioRxiv* (2020). 335 336

Acknowlededements 337

The authors would like to thank Dan Benjamin, Dalton Conley, Michel Nivard, Paul Rathouz, 338
Mijke Rhemtulla, and Patrick Turley for helpful comments on an early version of this manuscript. 339
This research was conducted using the UK Biobank Resource (Application No. 36046). This work 340
was supported in part by the National Science Foundation Graduate Research Fellowship Program 341
under Grant No. DGE-1656518 (ST), by the Institute of Education Sciences under Grant No. 342
R305B140009 (ST), by NIH grants R01MH120219, R01AG054628, and R01HD083613 (EMTD), 343
and by the Jacobs Foundation (EMTD). Any opinions expressed are those of the authors alone 344
and should not be construed as representing the opinions of funding agencies. The authors declare 345
no competing interests. 346

Distinguishing between interaction and dispersion effects in

the analysis of gene-environment interaction

Supplemental Information (SI)

A Methods

A.1 Description of the Heteroscedastic Regression Model and Deriva-

tion of ξ

353 The *environmental heteroscedasticity model* is specified as

$$Y_i = \tau_0 + \tau_1 E_i + \pi_0 G_i + \pi_1 G_i \cdot E_i + \lambda_0 \epsilon_i + \lambda_1 E_i \cdot \epsilon_i. \tag{8}$$

Here, τ_1 is the main effect of the measured environment, π_0 is the main effect of G_i, π_1 is the
GxE directly analogous to β_3 in Eqn 1, λ_0 is the main effect of the error ϵ_i (which we assume,
without loss of generality given λ_0, to have unit variance), and λ_1 is the coefficient used to index
heteroscedasticity.
This model will produce very similar unstandardized estimates of GxE as the standard GxE
model given by Eqn 1. In other words, π_1 in Eqn 8 is directly comparable to β_3 in Eqn 1.
However, the heteroscedasticity model further models whether variance in the phenotype that is
not accounted for by the genetic predictor (G_i) also varies as a function of the environment (E_i)
in the form of an $E_i \epsilon_i$ interaction, as captured by the λ_1 parameter. Importantly, when G_i only
index portions of total genetic etiology of a trait [1–3], the $E_i \epsilon_i$ interaction may result from GxE
with respect to the genetic factors not indexed by Gi.
In order to test whether a more general pattern of heteroscedasticity as a function of E_i is
sufficient to account for the observed GxE, we propose a restricted form version of this model that
does not directly model GxE but instead allows for the total variance of the phenotype to vary as a
function of E_i. This *scaling model* represents a scenario wherein GxE, in the sense of a significant
estimate of β_3 in Eqn 1, arises as a function of differences in the total variance of Y as a function
of E_i. The scaling model for moderation of the variance of Y_i as a function of E_i takes the form

$$Y_i = a_0 + a_1 E_i + (b_0 + b_1 E_i) Y_i^{\star}. \tag{9}$$

371   Here, E_i is a measured environment and $Y_i^{\star}$ is an unobserved factor representing unexplained
372 variation in Y_i incremental of E_i. We introduce a scaling transformation that produces het-
373 eroscedasticity in Y_i as a function of E_i via the $b_0 + b_1 E_i$ coefficient on the $Y^{\star}$ term. We let

$$Y_i^{\star} = hG_i + e\epsilon_i \tag{10}$$

where G is a measured genotype standardized to have unit variance and ϵ_i is an unobserved
error term (we assume it is also scaled to have unit variance). Finally, we set the scale of $Y^{\star}$
by specifying $e = \sqrt{1 - h^2}$. The penetrance of the measured gentoype is thus controlled via the
relative magnitudes of h and e. Note, in particular, that in terms of $Y_i^{\star}$, the penetrance of the
measured genotype is constant. The test that we introduce relies on this property; i.e. that when
properly scaled, the penetrance of the measured genotype to Y_i is constant in this model. Suppose
that $b_1 = 0$. In that case, Y_i is affected solely by environmental and genetic main effects. If $b_1 \neq 0$,
then the role of $Y_i^{\star}$ with respect to Y_i varies as a function of E, and as we elucidate next, so do
the raw (but not relative) contributions of G_i and ϵ_i to Y.

Substituting Eqn 10 into Eqn 9, and setting $e = \sqrt{1-h^2}$, yields

$$Y_i = a_0 + a_1 E_i + (b_0 + b_1 E_i)(hG_i + e\epsilon_i) \tag{11}$$

$$= a_0 + a_1 E_i + b_0 h G_i + b_1 h E_i G_i + b_0 e \epsilon_i + b_1 e E_i \epsilon_i \tag{12}$$

$$= a_0 + a_1 E_i + b_0 h G_i + b_1 h E_i G_i + b_0 (\sqrt{1-h^2}) \epsilon_i + b_1 (\sqrt{1-h^2}) E_i \epsilon_i \tag{13}$$

It can be seen that the environmental scaling model (Eqn 13) constitutes a constrained form of the environmental heteroscedasticity model (Eqn 8). Both equations take the same form in terms of E and G. However, whereas the coefficients in Eqn 8 are unconstrained, these coefficients are linked to one another in Eqn 13 via the the ratio h/e which, as noted above, is fixed under the scaling model. Thus, Eqn 8 is equivalent to Eqn 13 when

$$\frac{\pi_0}{\lambda_0} = \frac{\pi_1}{\lambda_1}. \tag{14}$$

because, under Eqn 13

$$\frac{b_0 h}{b_0 e} = \frac{h}{e} = \frac{b_1 h}{b_1 e}. \tag{15}$$

i.e.

$$\frac{b_0 h}{b_0 \sqrt{1-h^2}} = \frac{h}{\sqrt{1-h^2}} = \frac{b_1 h}{b_1 \sqrt{1-h^2}}. \tag{16}$$

We use Eqn 14 as the basis for a hypothesis test. In particular, we consider the test statistic

$$\xi_E \equiv \widehat{\pi_0} \widehat{\lambda_1} - \widehat{\pi_1} \widehat{\lambda_0}, \tag{17}$$

$$H_0 : \xi_E = 0. \tag{18}$$

When $\xi_E = 0$, the environmental heteroscedasticity model is indistinguishable from the simpler scaling model, whereas when ξ_E is differs from 0, the environmental scaling model is rejected.

A.2   Estimation

We estimate Eqn 8 using maximum likelihood by assuming $\epsilon_i \sim \text{Normal}(0,1)$. We assume that Y is mean-centered and that both G_i and E_i are non-random. Consider $\mathbb{E}(Y_i)$ and $\mathbb{V}(Y_i)$. We first have

$$\mathbb{E}(Y_i) = \tau_1 E_i + \pi_0 G_i + \pi_1 G_i \cdot E_i \tag{19}$$

since $\mathbb{E}(\epsilon) = 0$. We then have

$$\mathbb{V}(Y_i) = \mathbb{V}((\lambda_0 + \lambda_1 E_i)\epsilon) = (\lambda_0 + \lambda_1 E_i)^2 \tag{20}$$

since we assume $\mathbb{V}(\epsilon) = 1$. If we write

$$\theta = \{\tau_1, \pi_0, \pi_1, \lambda_0, \lambda_1\}, \tag{21}$$

note that both $\mathbb{E}(Y_i)$ and $\mathbb{V}(Y_i)$ are functions of θ.

Estimation of Eqn 8 is based on the likelihood,

$$L = \prod \phi\left(Y_i; \mathbb{E}(Y_i), \mathbb{V}(Y_i)\right) \tag{22}$$

where ϕ is the normal density and we use the above expressions for $\mathbb{E}(Y_i)$ and $\mathbb{V}(Y_i)$. We can thus solve for $\theta = \{\tau_1, \pi_0, \pi_1, \lambda_0, \lambda_1\}$ using ML based on

$$\log L = \sum \log \phi\left(Y_i; \mathbb{E}(Y_i), \mathbb{V}(Y_i)\right). \tag{23}$$

In particular, we obtain estimators via

$$\widehat{\theta_{\mathrm{ML}}} = \mathrm{argmax}_\theta \sum \log \phi\left(Y_i; \mathbb{E}(Y_i), \mathbb{V}(Y_i)\right). \tag{24}$$

To obtain ML estimates, it is convenient to also have the gradient. Here, we have

$$\frac{\partial \log L}{\tau_1} = \sum \frac{\partial \log L}{\partial \mu_i} E_i \tag{25}$$

$$\frac{\partial \log L}{\pi_0} = \sum \frac{\partial \log L}{\partial \mu_i} G_i \tag{26}$$

$$\frac{\partial \log L}{\pi_1} = \sum \frac{\partial \log L}{\partial \mu_i} G_i E_i \tag{27}$$

$$\frac{\partial \log L}{\lambda_0} = \sum \frac{\partial \log L}{\partial \sigma_i} \tag{28}$$

$$\frac{\partial \log L}{\lambda_1} = \sum \frac{\partial \log L}{\partial \sigma_i} E_i \tag{29}$$

where $\mu_i = \mathbb{E}(Y_i)$, $\sigma_i = \sqrt{\mathbb{V}(Y_i)}$, and

$$\frac{\partial \log L}{\partial \mu_i} = \sum \frac{Y_i - \mu_i}{\sigma_i^2} \tag{30}$$

$$\frac{\partial \log L}{\partial \sigma_i} = (Y_i - \mu_i)^2 \sigma_i^{-3} - \sigma_i^{-1}. \tag{31}$$

Using these estimates, we can compute a test statistic. Replacing the values in Eqn 15 with estimates from Eqn 8, we propose to test

$$\frac{\widehat{\pi_0}}{\widehat{\lambda_0}} = \frac{\widehat{\pi_1}}{\widehat{\lambda_1}} \rightarrow \tag{32}$$

$$\xi_E \equiv \widehat{\pi_0}\widehat{\lambda_1} - \widehat{\pi_1}\widehat{\lambda_0}, \tag{33}$$

$$H_0 : \xi_E = 0. \tag{34}$$

If H_0 holds, the underlying form of GxE cannot be distinguished from the scaling phenomenon in which the total variance in Y_i varies as a function of E. Alternatively put, failure to reject H_0 constitutes a failure to reject the hypothesis that the *proportional* contributions of G_i and ϵ_i to Y_i do not vary with E.

H_0 can be tested via a Wald test for nonlinear restrictions. To test this nonlinear restriction, we consider (see Ref [4], Eqn 6-29)

$$W = \widehat{\xi_E} \cdot (\mathbb{V}(\xi_E))^{-1} \cdot \widehat{\xi_E}. \tag{35}$$

Given that we are testing a single restriction, W has a χ^2 distribution with one degree of freedom in large samples. To estimate $\mathbb{V}(\xi_E)$, we use the Hessian of Eqn 23 (see B) and the delta method. Computationally, we do this via [5]. Note also that conventional tests of π_1 and λ_1 can be used to offer insight into the existence of GxE and potential heteroscedasticity; we make use of such tests below.

A.3   Extensions

A.3.1   Decomposition of π_1

We can decompose the π_1 estimate in Eqn 8 in a way that helps elucidate the contribution of heteroscedasticity to observed GxE. Consider an expanded version of the environmental scaling

model (Eqn 9) wherein we add an explicit GxE term:

$$Y_i = a_0 + a_1 E_i + a_2 G_i E_i + (b_0 + b_1 E_i) Y_i^\star. \tag{36}$$

This expanded scaling model (in conjunction with Eqn 10) is equivalent (same number of free parameters and same model fit) to the environmental heteroscedasticy model, but decomposes the GxE effect into a component that is accounted for by scaling and a component unaccounted for by scaling (note that all environmental heteroscedasticity is assumed to operate through scaling). We can therefore equate Eqn 36 and Eqn 8 and recover the following system of equations:

$$a_1 = \tau_1 \tag{37}$$
$$b_0 h = \pi_0 \tag{38}$$
$$a_2 + b_1 h = \pi_1 \tag{39}$$
$$b_0 e = \lambda_0 \tag{40}$$
$$b_1 e = \lambda_1 \tag{41}$$
$$h^2 + e^2 = 1. \tag{42}$$

Using estimates from Eqn 8 for the quantities on the right, we can solve for the parameters on the left. In particular, we can decompose π_1 into a_2 and $b_1 h$. The term $a2$ represents the degree to which observed data depart from the scaling model. Moreover, we can calculate the ratio

$$\frac{a_2}{a_2 + b_1 h} \tag{43}$$

to describe the the magnitude of this departure as a proportion of the total GxE effect.

A.3.2 Genetically-linked heteroscedasticity

A version of the heteroscedasticity model presented in Eqn 8 above can be constructed so as to relax the assumption of homoscedasticity of residuals across G_i as follows

$$Y_i = \tau_0 + \tau_1 E_i + \pi_0 G_i + \pi_1 G_i \cdot E_i + \lambda_0 \epsilon_i + \lambda_2 G_i \cdot \epsilon_i. \tag{44}$$

Here λ_2 is the coefficient used to index heteroscedasticity as a function of G_i. We assume homoscedasticity of residuals as a function of E_i by omitting λ_1. We refer to Eqn 44 as a *genetic heteroscedasticity model*. When a single variant is entered for G, this model consitutes a vQTL model that includes a term for GxE. Note that this model is equivalent to the model presented in Eqn 8, but with G_i entered for E_i and E_i entered for G_i.

The scaling model for moderation of the variance of Y_i as a function of G_i takes the form

$$Y_i = d_0 + d_1 G_i + (f_0 + f_1 G_i) Y_i^\star. \tag{45}$$

and

$$Y_i^\star = s E_i + \sqrt{1 - s^2} \epsilon_i \tag{46}$$

As in the scaling model for moderation the variance of Y_i as a function of E_i, ϵ_i is assumed to have unit variance. In terms of $Y_i^\star$, the effect of E_i constant across G_i. However, in terms of Y_i, the effect of E_i changes across G_i as result of heteroscedasticity across G_i.

It follows that the hypothesis test comparing the GxE model allowing for heterocedasticity as a function of G_i (given by Eqn 44) to its reduced scaling model (given by Eqns 45 and 46, is

$$\xi_G \equiv \widehat{\tau_1} \widehat{\lambda_2} - \widehat{\pi_1} \widehat{\lambda_0}, \tag{47}$$
$$H_0 : \xi_G = 0. \tag{48}$$

Finally, we can fit a *full heteroscedasticity model* that allows for moderation of residual variance

in Y_i as a function of both G_i and E_i as follows

$$Y_i = \tau_0 + \tau_1 E_i + \pi_0 G_i + \pi_1 G_i \cdot E_i + \lambda_0 \epsilon_i + \lambda_1 E_i \cdot \epsilon_i + \lambda_2 G_i \cdot \epsilon_i. \tag{49}$$

This model may be particularly useful when there is nonzero gene-environment correlation between the two moderators, i.e. $\mid r(G_i, E_i) \mid > 0$, such that G_i and E_i can be mutually controlled when modelling heteroscedasticity. However, to avoid added complexity, we recommend that comparisons with the scaling models rely on the ξ_E and ξ_G tests from the separate environmental heteroscedasticity and genetic heteroscedasticity models, respectively.

B The Hessian

We use the Hessian $(\frac{\partial^2 \log L}{\partial \theta^2})$ for construction of the covariance matrix of the estimation errors for the model parameters under the null model. The constituent parts of the Hessian are as follows.

$$\frac{\partial^2 \log L}{\partial \tau_1^2} = \sum_i -E_i^2 \sigma_i^{-2} \tag{50}$$

$$\frac{\partial^2 \log L}{\partial \tau_1 \partial \pi_0} = \sum_i -E_i G_i \sigma_i^{-2} \tag{51}$$

$$\frac{\partial^2 \log L}{\partial \tau_1 \partial \pi_1} = \sum_i -E_i^2 G_i \sigma_i^{-2} \tag{52}$$

$$\frac{\partial^2 \log L}{\partial \tau_1 \partial \lambda_0} = \sum_i -2E_i(Y_i - \mu_i)\sigma_i^{-3} \tag{53}$$

$$\frac{\partial^2 \log L}{\partial \tau_1 \partial \lambda_1} = \sum_i -2E_i^2(Y_i - \mu_i)\sigma_i^{-3} \tag{54}$$

$$\frac{\partial^2 \log L}{\partial \pi_0^2} = \sum_i -G_i^2 \sigma_i^{-2} \tag{55}$$

$$\frac{\partial^2 \log L}{\partial \pi_0 \partial \pi_1} = \sum_i -E_i G_i^2 \sigma_i^{-2} \tag{56}$$

$$\frac{\partial^2 \log L}{\partial \pi_0 \partial \lambda_0} = \sum_i -2G_i(Y_i - \mu_i)\sigma_i^{-3} \tag{57}$$

$$\frac{\partial^2 \log L}{\partial \pi_0 \partial \lambda_1} = \sum_i -2E_i G_i(Y_i - \mu_i)\sigma_i^{-3} \tag{58}$$

$$\frac{\partial^2 \log L}{\partial \pi_1^2} = \sum_i -E_i^2 G_i^2 \sigma_i^{-2} \tag{59}$$

$$\frac{\partial^2 \log L}{\partial \pi_1 \partial \lambda_0} = \sum_i -2E_i G_i(Y_i - \mu_i)\sigma_i^{-3} \tag{60}$$

$$\frac{\partial^2 \log L}{\partial \pi_1 \partial \lambda_1} = \sum_i -2E_i^2 G_i(Y_i - \mu_i)\sigma_i^{-3} \tag{61}$$

$$\frac{\partial^2 \log L}{\partial \lambda_0^2} = \sum_i \left[\sigma_i^{-2} - 3(Y_i - \mu_i)^2 \sigma_i^{-4} \right] \tag{62}$$

$$\frac{\partial^2 \log L}{\partial \lambda_0 \partial \lambda_1} = \sum_i \left[E_i \sigma_i^{-2} - 3E_i(Y_i - \mu_i)^2 \sigma_i^{-4} \right] \tag{63}$$

$$\frac{\partial^2 \log L}{\partial \lambda_1^2} = \sum_i \left[E_i^2 \sigma_i^{-2} - 3E_i^2(Y_i - \mu_i)^2 \sigma_i^{-4} \right] \tag{64}$$

$$\tag{65}$$

C   Simulation Study

We now consider both the null distribution of the test statistic under different configurations of
the relevant parameters from the scaling model (Eqn 13) and the performance of our test statistic
in the case of generating models that do not conform to the scaling model (i.e. when Eqn 1 is
used to generate a model with GxE and homoscedastic residuals, or Eqn 8 is used to general a
model with heteroscedastic residuals but no GxE). We primarily describe the simulation models
in the context of the heteroscadasticity and scaling models in which the measured environment,
E_i moderates the residual variance of the phenotype, Y_i, as given by Eqns 8 and 13. However,
as described in A.3, these models are equivalent to those in which the measured phenotype, G_i
moderates the residual variance of Y_i (after G_i and E_i are substituted for one another). Thus
simulation results apply to both scenarios.

C.1   Distribution of the test statistic under H_0

We first consider the behavior of the test statistic when the null hypothesis is true. We generate
data according to the *scaling model* (Eqn 13), estimate the *heteroscedasticity model* (Eqn 8), and
then calculate the ξ_E test statistic and associated Wald-based probability. We start by illustrating
that the appropriate transformation of ξ_E has a χ^2 distribution. To do this, we generate 10,000
datasets from the scaling model (Eqn 13) with $b_0 = 1, b_1 = 0.15, h = 0.5, N = 5000$. We compute
ξ_E and then transform it to W via Eqn 35. Results are in Figure C.1. The null distribution of
W neatly aligns with the density of $\chi^2(1)$. Below, we report on ξ_E given that it is a relatively
intuitive function of the estimated parameters but also exploit the fact that ξ_E can be mapped to
W and inference can be done via reference to $\chi^2(1)$.

Figure C.1: Comparison of distribution of W under the null hypothesis to $\chi^2(1)$.

We next considered the behavior of ξ_E under the null, for configurations of the relevant param-
eters in the scaling model (Eqn 13) meant to capture the relevant features of an analysis in which
G is a PGS. We considered variation in four parameters. We used three different sample sizes
(1000, 5000, 10000). We used two values respectively for b_0 (0.8, 1) and b_1 (0.05, 0.1). Finally,
we used two values for h (0.1 and 0.5). The values of h suggest weakly (0.1) to strongly (0.5)
penetrant genetic predictors; they were chosen to represent the range of potential PGS predictors
given existing technology [1]. Recall that the value of e is fixed as a function of h. Note that we
considered two different distributions for E; we sampled E from both the standard normal and
the uniform distributions (where the uniform distribution was scaled to have the same standard
deviation as the standard normal). As polygenic scores for complex traits are expected to be
normally distributed [6], G was sampled from the standard normal distribution. To help develop
intuition, suppose $b_0 = 0.8$, $b_1 = 0.05$, and $h = 0.5$. In that case, $\mathbb{V}(y|E = 2) = 0.736$ compared to
$\mathbb{V}(y|E = -2) = 0.422$, a ratio of 1.74 (i.e., there is heteroscedasticity). Finally, it is important to
note that the scale of ξ_E is dependent on several features (e.g. scale of G and E); patterns shown

here are informative about the behavior of ξ_E but the values themselves do not necessarily generalize to other cases (e.g., if E or G does not have unit variance). In contrast, W and associated tests are invariant to scaling of G and E, but sensitive to sample size.

Results are shown in Table C.1. Note that the estimates of ξ_E are unbiased (i.e., the means are nearly zero). Further, as we would anticipate, the false positive rate (FPR) of the test is insensitive to the parameters; note also that the mean probability associated with the test of ξ_E via Eqn 35 is near 0.05, as we would expect when the null is true.[1]

In contrast, the variance of the test statistic (and, thus, statistical power) depends on the sample size and simulation parameters. In particular, note the decline in the SD of the test statistics as a function of the sample size. Statistical power will also depend upon h; this test will have less power with less penetrant genetic predictors.

Table C.1: Distribution of ξ_E under the null (mean and SD) and associated test (Eqn 35) for different values of parameters in Eqn 13. For each set of parameters, 1000 datasets are simulated. Normal and Uniform refer to the distributions used to generate the environmental variable.

Simulation Parameters				Results (Normal E_i)				Results (Uniform E_i)			
N	b_0	b_1	h	Mean(ξ_E)	SD(ξ_E)	Mean($\Pr_{\xi_E}$)	FPR(ξ_E)	Mean(ξ_E)	SD(ξ_E)	Mean($\Pr_{\xi_E}$)	FPR(ξ_E)
1000	0.80	0.05	0.10	1.58e-04	0.02	0.50	0.06	-9.05e-04	0.01	0.51	0.04
1000	1.00	0.05	0.10	3.71e-04	0.03	0.50	0.06	1.11e-03	0.02	0.50	0.05
1000	0.80	0.10	0.10	-5.53e-05	0.02	0.49	0.06	3.63e-04	0.01	0.50	0.05
1000	1.00	0.10	0.10	3.66e-04	0.03	0.49	0.05	-7.39e-04	0.02	0.50	0.05
1000	0.80	0.05	0.50	3.46e-04	0.02	0.49	0.05	-6.60e-05	0.01	0.49	0.05
1000	1.00	0.05	0.50	1.30e-03	0.02	0.51	0.04	-1.98e-04	0.02	0.50	0.04
1000	0.80	0.10	0.50	3.35e-04	0.02	0.50	0.05	7.74e-04	0.01	0.50	0.04
1000	1.00	0.10	0.50	-6.83e-04	0.03	0.51	0.05	-6.09e-05	0.02	0.50	0.05
5000	0.80	0.05	0.10	1.31e-04	0.01	0.51	0.05	-2.31e-04	0.01	0.52	0.04
5000	1.00	0.05	0.10	1.78e-04	0.01	0.52	0.04	-1.95e-04	0.01	0.49	0.05
5000	0.80	0.10	0.10	-2.32e-04	0.01	0.49	0.05	4.81e-06	0.01	0.51	0.04
5000	1.00	0.10	0.10	1.83e-04	0.01	0.49	0.05	5.02e-04	0.01	0.51	0.04
5000	0.80	0.05	0.50	1.59e-04	0.01	0.51	0.06	-1.48e-04	0.01	0.50	0.05
5000	1.00	0.05	0.50	-8.10e-05	0.01	0.47	0.05	2.47e-04	0.01	0.50	0.05
5000	0.80	0.10	0.50	1.95e-04	0.01	0.49	0.06	-8.94e-05	0.00	0.50	0.05
5000	1.00	0.10	0.50	4.67e-04	0.01	0.49	0.05	3.49e-05	0.01	0.51	0.05
10000	0.80	0.05	0.10	1.29e-04	0.01	0.50	0.04	-2.16e-05	0.00	0.49	0.07
10000	1.00	0.05	0.10	-9.81e-05	0.01	0.50	0.04	6.69e-05	0.01	0.50	0.05
10000	0.80	0.10	0.10	1.89e-04	0.01	0.49	0.04	1.85e-04	0.00	0.51	0.05
10000	1.00	0.10	0.10	-9.32e-05	0.01	0.50	0.05	4.98e-05	0.01	0.52	0.04
10000	0.80	0.05	0.50	2.62e-04	0.01	0.50	0.05	-4.27e-05	0.00	0.51	0.05
10000	1.00	0.05	0.50	2.56e-05	0.01	0.49	0.05	-1.42e-04	0.01	0.50	0.04
10000	0.80	0.10	0.50	-2.30e-04	0.00	0.49	0.06	-1.89e-04	0.00	0.51	0.04
10000	1.00	0.10	0.50	-3.71e-04	0.01	0.49	0.06	-6.03e-05	0.01	0.49	0.05

We next considered the behavior of of ξ_E under the null for configurations of parameters meant to capture the relevant features of an analysis in which G is an allele count for a single locus (e.g. a SNP). Results are in Table C.2. The table is similar to Table C.1, but where we now consider allele counts based on variants simulated under a binomial distribution with different minor allele frequencies (denoted p). Based on Extended Data Figure 2 in [7], we chose effect sizes of $h^2 \in \{.05\%, .1\%\}$. Note that after generating G under a binomial distribution, it was standardized, such that h can be interpreted as a standardized effect size. We examine behavior using a sample size of $N = 10000$, which might be used for confirmatory tests of a select number of variants previously identified in a larger independent GWAS, and a sample size of $N = 1000000$, which might be used for genome-wide discovery. Simulation results indicate that the heteroscedasticity model and the ξ_E statistic behave as expected across the range of conditions examined.

C.2   Performance under alternatives

We now turn to study the behavior of the heteroscedasticity model and associated ξ_E test statistic when the null hypothesis is false (i.e., when Eqn 13 is not the data generating model). We consider

---

[1]We use $\Pr_\beta$ to refer to probability associated with a test of the null hypothesis that some parameter β equals zero.

Table C.2: Distribution of ξ_E under the null (mean and SD) and associated test (via Eqn 35) for different values of parameters in Eqn 13 based on a standardized single-locus allele count. For each set of parameters, 100 datasets are simulated. The environmental variable is sampled from the standard normal distribution.

N	b_0	b_1	h	MAF	$\widehat{\pi_1}$	$\widehat{\lambda_1}$	Mean(ξ_E)	SD(ξ_E)	Mean(Pr_{ξ_E})	$\frac{\mathbb{E}(SE(\xi_E))}{SD(\xi_E)}$
10000	0.80	0.05	0.02	0.01	6.91e-04	0.05	3.25e-04	6.50e-03	0.51	0.97
10000	1.00	0.05	0.02	0.01	6.52e-04	0.05	4.67e-04	9.76e-03	0.49	1.02
10000	0.80	0.10	0.02	0.01	3.25e-03	0.10	-7.18e-04	6.35e-03	0.47	0.95
10000	1.00	0.10	0.02	0.01	2.53e-03	0.10	-3.72e-04	9.29e-03	0.52	1.05
10000	0.80	0.05	0.03	0.01	1.17e-03	0.05	3.92e-04	6.18e-03	0.52	1.02
10000	1.00	0.05	0.03	0.01	1.81e-03	0.05	-1.64e-04	1.13e-02	0.47	0.88
10000	0.80	0.10	0.03	0.01	2.29e-03	0.10	8.04e-04	6.13e-03	0.48	0.98
10000	1.00	0.10	0.03	0.01	2.43e-03	0.10	7.48e-04	9.56e-03	0.49	1.01
100000	0.80	0.05	0.02	0.01	1.27e-03	0.05	-1.12e-04	1.96e-03	0.52	1.02
100000	1.00	0.05	0.02	0.01	6.32e-04	0.05	4.75e-04	3.10e-03	0.47	1.01
100000	0.80	0.10	0.02	0.01	2.00e-03	0.10	1.59e-04	1.84e-03	0.52	1.03
100000	1.00	0.10	0.02	0.01	2.49e-03	0.10	-2.23e-04	2.97e-03	0.50	1.02
100000	0.80	0.05	0.03	0.01	1.21e-03	0.05	2.92e-04	2.11e-03	0.47	0.94
100000	1.00	0.05	0.03	0.01	1.40e-03	0.05	1.70e-04	2.88e-03	0.52	1.09
100000	0.80	0.10	0.03	0.01	3.04e-03	0.10	7.05e-05	2.16e-03	0.46	0.88
100000	1.00	0.10	0.03	0.01	2.53e-03	0.10	5.93e-04	3.04e-03	0.53	1.00
10000	0.80	0.05	0.02	0.05	3.28e-03	0.05	-1.73e-03	5.97e-03	0.52	1.06
10000	1.00	0.05	0.02	0.05	1.03e-03	0.05	9.44e-05	1.15e-02	0.41	0.86
10000	0.80	0.10	0.02	0.05	2.20e-03	0.10	1.36e-04	5.53e-03	0.53	1.08
10000	1.00	0.10	0.02	0.05	2.61e-03	0.10	-8.78e-06	1.04e-02	0.47	0.93
10000	0.80	0.05	0.03	0.05	7.91e-04	0.05	6.14e-04	6.44e-03	0.47	0.98
10000	1.00	0.05	0.03	0.05	1.51e-03	0.05	-4.78e-06	1.13e-02	0.47	0.88
10000	0.80	0.10	0.03	0.05	4.31e-03	0.10	-7.83e-04	6.14e-03	0.50	0.98
10000	1.00	0.10	0.03	0.05	4.16e-03	0.10	-9.08e-04	9.64e-03	0.48	0.99
100000	0.80	0.05	0.02	0.05	1.16e-03	0.05	-5.90e-05	1.98e-03	0.54	1.00
100000	1.00	0.05	0.02	0.05	7.56e-04	0.05	3.44e-04	3.06e-03	0.49	1.02
100000	0.80	0.10	0.02	0.05	1.74e-03	0.10	3.91e-04	2.03e-03	0.48	0.93
100000	1.00	0.10	0.02	0.05	2.20e-03	0.10	3.24e-05	3.20e-03	0.47	0.95
100000	0.80	0.05	0.03	0.05	1.72e-03	0.05	-9.93e-05	1.83e-03	0.54	1.09
100000	1.00	0.05	0.03	0.05	1.31e-03	0.05	2.68e-04	3.31e-03	0.46	0.95
100000	0.80	0.10	0.03	0.05	3.69e-03	0.10	-3.60e-04	2.15e-03	0.44	0.88
100000	1.00	0.10	0.03	0.05	3.09e-03	0.10	5.50e-05	3.35e-03	0.45	0.91
10000	0.80	0.05	0.02	0.50	-4.27e-04	0.05	1.22e-03	6.81e-03	0.48	0.92
10000	1.00	0.05	0.02	0.50	1.53e-03	0.05	-4.68e-04	8.87e-03	0.53	1.11
10000	0.80	0.10	0.02	0.50	1.60e-03	0.10	5.75e-04	6.84e-03	0.45	0.87
10000	1.00	0.10	0.02	0.50	3.90e-03	0.10	-1.55e-03	9.61e-03	0.53	0.99
10000	0.80	0.05	0.03	0.50	1.35e-03	0.05	1.87e-04	6.44e-03	0.51	0.98
10000	1.00	0.05	0.03	0.50	3.36e-03	0.05	-1.68e-03	9.08e-03	0.56	1.08
10000	0.80	0.10	0.03	0.50	4.00e-03	0.10	-6.84e-04	5.63e-03	0.52	1.06
10000	1.00	0.10	0.03	0.50	3.78e-03	0.10	-5.68e-04	9.75e-03	0.50	0.99
100000	0.80	0.05	0.02	0.50	1.34e-03	0.05	-1.60e-04	2.00e-03	0.48	1.00
100000	1.00	0.05	0.02	0.50	7.07e-04	0.05	4.14e-04	2.91e-03	0.51	1.08
100000	0.80	0.10	0.02	0.50	2.21e-03	0.10	3.19e-05	1.92e-03	0.51	0.99
100000	1.00	0.10	0.02	0.50	2.00e-03	0.10	2.02e-04	2.73e-03	0.55	1.11
100000	0.80	0.05	0.03	0.50	1.89e-03	0.05	-2.42e-04	2.24e-03	0.44	0.89
100000	1.00	0.05	0.03	0.50	1.99e-03	0.05	-4.30e-04	3.26e-03	0.52	0.96
100000	0.80	0.10	0.03	0.50	3.00e-03	0.10	1.23e-04	2.04e-03	0.44	0.92
100000	1.00	0.10	0.03	0.50	3.51e-03	0.10	-3.25e-04	2.96e-03	0.50	1.02

two different data generating models: (i) a model with GxE and homoscedastic residuals, generated according to Eqn 1 (ii) a model with no GxE but with heteroscedastic residuals, generated according to Eqn 8. In these analyses, we aim to probe whether the test statistic allows us to reject H_0 under different configurations of the relevant data-generating parameters.

We first study a circumstance in which the generating model is GxE with homoscedastic residuals. We use Eqn 1 for our data-generating model and suppose that e_i is white-noise (i.e., $e_i \sim \mathrm{N}(0, \sigma^2)$) as opposed to heteroscedastic as a function of E_i. In particular, we set $\sigma^2 = 1$. We focus on two tests: a test of whether $\pi_1 = 0$ and a test of whether $\xi_E = 0$. Data are analyzed with the environmental heteroscadasticity model (Eqn 8, which subsumes Eqn 1).

Results are shown in Table C.3. When both N and b_3 are relatively small, we are poorly powered to detect GxE and to reject the scaling hypothesis (i.e., both Pr_{π_1} and Pr_{ξ_E} are relatively large). Importantly, when GxE is not detected, the ξ_E test is moot. In contrast, power to detect GxE and to reject scaling are both excellent when N or b_3 is relatively large. Throughout we reject $\lambda_1 = 0$ for $\alpha = .05$ at the appropriate FPR.

Table C.3: Behavior of ξ_E when true model is homoscedastic GxE (Eqn 1). Behavior is shown for 1000 simulated datasets based on each set of parameters.

Simulation Parameters				Results				
N	β_1	β_2	β_3	Mean(Pr_{π_1})	Power(π_1)	Mean(Pr_{ξ_E})	Power(ξ_E)	FPR(λ_1)
1000	0.10	0.00	0.05	2.45e-01	3.48e-01	2.48e-01	3.48e-01	0.05
1000	0.20	0.00	0.05	2.26e-01	3.33e-01	2.31e-01	3.35e-01	0.05
1000	0.10	0.20	0.05	2.36e-01	3.56e-01	2.38e-01	3.57e-01	0.07
1000	0.20	0.20	0.05	2.16e-01	3.87e-01	2.21e-01	3.80e-01	0.05
1000	0.10	0.50	0.05	2.40e-01	3.34e-01	2.41e-01	3.32e-01	0.05
1000	0.20	0.50	0.05	2.23e-01	3.57e-01	2.26e-01	3.41e-01	0.06
1000	0.10	0.00	0.15	7.68e-04	9.98e-01	8.40e-04	9.97e-01	0.05
1000	0.20	0.00	0.15	9.08e-04	9.97e-01	1.01e-03	9.95e-01	0.04
1000	0.10	0.20	0.15	8.63e-04	9.98e-01	8.68e-04	9.97e-01	0.06
1000	0.20	0.20	0.15	6.13e-04	9.97e-01	6.72e-04	9.98e-01	0.05
1000	0.10	0.50	0.15	8.49e-04	9.96e-01	9.04e-04	9.97e-01	0.06
1000	0.20	0.50	0.15	9.02e-04	9.97e-01	1.14e-03	9.95e-01	0.06
5000	0.10	0.00	0.05	1.30e-02	9.37e-01	1.30e-02	9.32e-01	0.04
5000	0.20	0.00	0.05	9.69e-03	9.53e-01	1.03e-02	9.47e-01	0.06
5000	0.10	0.20	0.05	1.39e-02	9.37e-01	1.39e-02	9.32e-01	0.05
5000	0.20	0.20	0.05	1.26e-02	9.34e-01	1.34e-02	9.30e-01	0.05
5000	0.10	0.50	0.05	1.29e-02	9.43e-01	1.29e-02	9.43e-01	0.04
5000	0.20	0.50	0.05	1.52e-02	9.36e-01	1.60e-02	9.36e-01	0.05
5000	0.10	0.00	0.15	3.05e-13	1.00e+00	1.36e-12	1.00e+00	0.06
5000	0.20	0.00	0.15	1.62e-17	1.00e+00	9.32e-17	1.00e+00	0.04
5000	0.10	0.20	0.15	1.58e-15	1.00e+00	1.16e-15	1.00e+00	0.05
5000	0.20	0.20	0.15	2.74e-17	1.00e+00	5.83e-17	1.00e+00	0.04
5000	0.10	0.50	0.15	7.99e-18	1.00e+00	2.30e-17	1.00e+00	0.05
5000	0.20	0.50	0.15	1.48e-14	1.00e+00	2.87e-14	1.00e+00	0.05

We next consider the circumstance in which there is no GxE, but there are G and E main effects, and there is heteroscedasticity as a function of E so as to parallel Eqn 20. In particular, we generate data via

$$Y_i = b_1 G_i + b_2 E_i + \epsilon_i \tag{66}$$

with $\epsilon_i \sim \mathrm{N}(0, (\lambda_0 + \lambda_1 E_i)^2)$. We use $\lambda_0 = 2$ and vary λ_1 between 0.15 and 0.3 (if $\lambda_1 = 0$, there is no heteroscedasticity). We fit the environmental heteroscedasticity model (eqn 8) and focus on tests of $\pi_1 = 0$ (which is true under the generating model), $\lambda_1 = 0$ (which is false under the generating model), and $\xi_E = 0$ (which is false under the generating model).

Results are shown in Table C.4. First, note that the appropriate FPR of .05 is obtained for $\pi_1 = 0$ at alpha=.05, irrespective of λ_0 and λ_1. This indicates that heteroscedasticity has not induced the impression of GxE under these conditions. We can also uniformly detect that $\lambda_1 \neq 0$ suggesting that errors are in fact heteroscedastic. However, tests of ξ_E are relatively weak here (e.g., power is 0.2 at best). Importantly, as the tests of π_1 appropriate indicate no GxE, the question of whether there is scaling GxE is moot.

Table C.4: Behavior of ξ_E when true model does not have GxE but does have heteroscedastic error (Eqn 66). Behavior is shown for 1000 simulated datasets based on each set of parameters.

Simulation Parameters					Results					
N	π_0	τ_1	λ_0	λ_1	$\text{Mean}(\text{Pr}_{\pi_1})$	$\text{FPR}(\pi_1)$	$\text{Mean}(\text{Pr}_{\lambda_1})$	$\text{Power}(\lambda_1)$	$\text{Mean}(\text{Pr}_{\xi_E})$	$\text{Power}(\xi_E)$
1000.00	0.10	0.00	2.00	0.15	0.50	0.06	2.03e-02	0.90	0.49	0.05
1000.00	0.20	0.00	2.00	0.15	0.50	0.05	1.55e-02	0.93	0.49	0.06
1000.00	0.10	0.50	2.00	0.15	0.49	0.07	1.74e-02	0.92	0.48	0.07
1000.00	0.20	0.50	2.00	0.15	0.50	0.05	1.70e-02	0.93	0.48	0.06
1000.00	0.10	0.00	2.00	0.30	0.50	0.05	1.62e-06	1.00	0.48	0.05
1000.00	0.20	0.00	2.00	0.30	0.50	0.05	8.32e-07	1.00	0.45	0.09
1000.00	0.10	0.50	2.00	0.30	0.50	0.06	5.46e-06	1.00	0.49	0.07
1000.00	0.20	0.50	2.00	0.30	0.49	0.05	8.40e-07	1.00	0.46	0.07
5000.00	0.10	0.00	2.00	0.15	0.51	0.06	1.25e-08	1.00	0.51	0.06
5000.00	0.20	0.00	2.00	0.15	0.49	0.05	9.85e-08	1.00	0.46	0.09
5000.00	0.10	0.50	2.00	0.15	0.49	0.06	6.29e-09	1.00	0.48	0.06
5000.00	0.20	0.50	2.00	0.15	0.50	0.05	5.07e-08	1.00	0.46	0.08
5000.00	0.10	0.00	2.00	0.30	0.49	0.06	6.79e-36	1.00	0.45	0.10
5000.00	0.20	0.00	2.00	0.30	0.50	0.06	1.95e-37	1.00	0.33	0.22
5000.00	0.10	0.50	2.00	0.30	0.49	0.05	7.32e-38	1.00	0.44	0.09
5000.00	0.20	0.50	2.00	0.30	0.49	0.06	1.48e-38	1.00	0.32	0.23

D   Software

Code to estimate the heteroscestacity model and to estimate ξ_E is available in an R package.[2] In particular, the documentation illustrates functionality via simple simulations similar to those found in Section C.

---

[2]https://github.com/ben-domingue/scalingGxE

E Data

E.1 HRS Data

The first wave of HRS data collection was in 1992, the most recent in 2018 (wave 13). We focus on N=11,586 respondents of European ancestry. Descriptive information for the environment and outcome are shown in Figure E.1. We use data from the RAND HRS files on BMI, see [8] for additional documentation. In particular, we consider the mean BMI over all waves. The mean BMI was skewed, so we also considered a transformed version (see Figure E.1). HRS respondents were born across a wide span of birthyears centered around 1940. To ensure that results aren't driven by small numbers of observations with extreme birth years, we analyzed those born between 1920–1965. All data (polygenic score, birthyear, and BMI) are standardized to have mean zero and unit variance for analysis.

Figure E.1: Top: Density of BMI and birthyear in HRS data as well as QQ plot of the Box-Cox transformed version (meant to reduce skew). We standardize respondent's average BMI taken over all waves; this mean BMI is skewed (skew equals 1.14). This is a common finding with respect to BMI (see, for example, the Figure in [9]). Thus, we also consider a version of this variable in which it has been transformed, via the Box-Cox transformation, to closely approximate a normal distribution. All variables (birthyear, polygenic score, and outcome) are standardized (to have zero mean and unit variance) prior to analysis. Bottom: Illustration of increasing variance in BMI distribution as a function of birth year for analytic sample.

E.2 UKB data

We used data from UK Biobank to conduct a series of SNP-level gene-by-environment association analyses. Genotyping and imputation procedures were performed by the original UK Biobank investigators, as described in a previous publication [10].[3] In the present study, we applied additional thresholds to minimize potential confounding, removing participants with (i) low call rate, (ii) extreme heterozygosity, (iii) chromosomal aneuploidy, (iv) discordant self-reported and chromosomal sex, or (v) missing data for the relevant outcome and covariates. We further focused our

---

[3]See also https://www.ukbiobank.ac.uk/scientists-3/genetic-data/.

analyses on unrelated participants of non-Hispanic European ancestry to minimize the threat of population stratification.

Rather than perform a genome-wide scan of gene-by-environment interactions, we focused on our analyses on loci that have been previously associated with BMI. Specifically, we selected the 97 loci identified in a large genome-wide association study of body mass index that did not include UK Biobank [11]. We successfully identified and extracted the lead SNP for 96/97 (99%) of these loci. For imputed SNPs, dosages were converted to hard calls. Any converted hard calls with a certainty less than .90 were coded as missing.

As some UK Biobank participants had multiple observations for BMI, we computed the mean BMI for each individual. We applied a Box-Cox transformation to reduce skew. Box-Cox transformed BMI was then adjusted for sex, batch, and the first 40 principal components of ancestry, which we estimated ourselves [12]. All variables (SNP, birth year, and body mass index) were standardized to have mean zero and unit variance prior to analysis. The final analytic samples size was N=380,605.

 # F Supplemental Resultse

 ## F.1 Polygenic score analysis

To better illustrate the differences in the sex-stratified analyses, we consider a graphical analysis
inspired by the notion of posterior predictive checks [13]. Using estimates from the empirical data,
we construct simulated data sets using Eqn 13 and Eqn 1 and the HRS data. We then consider the
predicted percentage of phenotypic variance explained by the genetic predictor across birthyears.
Based on Eqn 8, we compute this percentage as

$$\frac{Vg}{Vp} = \frac{(\pi_0 + \pi_1 E)^2}{(\pi_0 + \pi_1 E)^2 + (\lambda_0 + \lambda_1 E)^2}. \tag{67}$$

Results of this graphical analysis are in Figure 2. For males (top row), the empirical data (in
red) produce a pattern that falls well within the distribution of results from data simulated from
the scaling model, but departs more appreciably from results from data simulated under the
conventional homoscedastic GxE model. The proportional contribution of the PGS to BMI does
not significantly increase with birth year for males. For females, there is more change in penetrance
than would be anticipated under the scaling model but less change than would be anticipated under
the conventional homoscedastic GxE model. As compared to males,the proportional contribution
of the PGS to BMI significantly increases with birth year for females. Thus, for females, the full
heteroscedasticity model, in which the contributions of PGS and non-PGS factors independently
shift with birth year, may be the most appropriate analytic tool.

  We also examined the potential for genetic heteroscedasticity using the genetic heteroscedas-
ticity model (Eqn 44). We observe evidence for moderation of residual variance in BMI by the
BMI PGS, as indicated by the λ_2 parameter. Interestingly, this result holds even for the Box-Cox
transformed version of BMI, suggesting that the heteroscedasticity is not entirely an artifact of
mean-variance associations induced by the skewed distribution of BMI. One interpretation of these
results is that the polygenic score for BMI also captures some degree of "plasticity" (in the sense
of [14]). Results of the ξ_G tests (see final column of Table 2) are similar to those of the ξ_E in that
we cannot reject the null of the scaling model for the male respondents.

 ## F.2 SNP analysis

Tables F.1 and F.2 contain results discussed in main text.

Table F.1: vQTL discoveries for BMI. From the 96 marker SNPs for genome-wide significant
loci for BMI in [11], those reaching significance at $p < .05/96$ for moderation of variance, as
indicated by the λ_2 coefficient from the full heterscedasticity model. Parameters are reported
for the full heteroscedasticity model, with ξ_E and ξ_G parameters from the environmental and
genetic heteroscedasticity models, respectively

SNP	π_0	π_1	Pr_{π_1}	λ_0	λ_1	λ_2	Pr_{λ_2}	Pr_{ξ_E}	Pr_{ξ_G}
rs2207139_A	-0.02	-0.00	3.76e-03	0.99	0.04	-0.00	8.93e-05	1.39e-02	3.84e-03
rs7903146_C	0.01	-0.00	6.48e-01	0.99	0.04	0.01	3.22e-06	8.09e-01	9.02e-01
rs9925964_A	0.02	-0.00	9.07e-01	0.99	0.04	0.00	1.10e-04	6.11e-01	7.50e-01
rs1558902_T	-0.05	-0.01	2.12e-06	0.99	0.04	-0.01	1.76e-17	2.96e-04	5.14e-07
rs6567160_T	-0.03	-0.00	1.65e-02	0.99	0.04	-0.01	2.84e-08	8.97e-02	1.40e-02
rs2287019_C	0.02	-0.00	8.71e-01	0.99	0.04	0.01	8.15e-06	8.27e-01	6.66e-01

Table F.2: Replication of main effects at $p < .05/96$ for the 96 marker SNPs for the genome-wide significant loci for BMI in [11]. Main effects (π_0) come from the full heteroscedasity model.

SNP	π_0	SE(π_0)	$\Pr_{\pi_0}$	Replication?
rs977747_G	-0.0119	0.0016	1.3380e-13	TRUE
rs657452_G	-0.0116	0.0016	6.3858e-13	TRUE
rs11583200_T	-0.0103	0.0016	1.8037e-10	TRUE
rs3101336_C	0.0152	0.0016	1.8616e-21	TRUE
rs12566985_A	-0.0133	0.0016	1.3812e-16	TRUE
rs12401738_G	-0.0103	0.0016	1.4297e-10	TRUE
rs11165643_T	0.0125	0.0016	6.5757e-15	TRUE
rs17024393_T	-0.0146	0.0016	1.8806e-19	TRUE
rs543874_A	-0.0271	0.0016	2.2391e-63	TRUE
rs2820292_C	0.0131	0.0016	2.6984e-16	TRUE
rs13021737_G	0.0279	0.0016	1.6738e-68	TRUE
rs10182181_A	-0.0243	0.0016	2.0290e-51	TRUE
rs11126666_G	-0.0021	0.0016	1.9697e-01	FALSE
rs1016287_C	-0.0128	0.0016	1.4586e-15	TRUE
rs11688816_G	0.0083	0.0016	3.0389e-07	TRUE
rs2121279_C	-0.0050	0.0016	2.0019e-03	FALSE
rs1460676_T	-0.0083	0.0016	2.7992e-07	TRUE
rs1528435_T	0.0101	0.0016	2.8226e-10	TRUE
rs17203016_A	-0.0071	0.0016	1.0494e-05	TRUE
rs7599312_G	0.0116	0.0016	9.6039e-13	TRUE
rs492400_T	-0.0072	0.0016	7.8542e-06	TRUE
rs2176040_G	-0.0014	0.0016	3.7490e-01	FALSE
rs6804842_G	0.0088	0.0016	5.6154e-08	TRUE
rs2365389_C	0.0092	0.0016	1.0286e-08	TRUE
rs3849570_C	-0.0075	0.0016	2.5934e-06	TRUE
rs13078960_T	-0.0109	0.0016	1.3418e-11	TRUE
rs16851483_G	-0.0124	0.0016	1.3755e-14	TRUE
rs1516725_C	0.0162	0.0016	4.0089e-24	TRUE
rs10938397_A	-0.0208	0.0016	1.5818e-38	TRUE
rs17001654_C	-0.0069	0.0016	2.3970e-05	TRUE
rs13107325_C	-0.0187	0.0016	2.1694e-31	TRUE
rs11727676_T	0.0014	0.0016	3.9716e-01	FALSE
rs2112347_T	0.0189	0.0016	5.4850e-32	TRUE
rs7715256_T	-0.0106	0.0016	3.8649e-11	TRUE
rs205262_A	-0.0185	0.0016	8.4528e-31	TRUE
rs2033529_A	-0.0132	0.0016	2.5074e-16	TRUE
rs2207139_A	-0.0209	0.0016	3.0425e-38	TRUE
rs9400239_C	0.0098	0.0016	1.0575e-09	TRUE
rs9374842_T	0.0059	0.0016	2.1120e-04	TRUE
rs13201877_A	-0.0048	0.0016	2.8052e-03	FALSE
rs13191362_A	0.0086	0.0016	8.0189e-08	TRUE
rs1167827_G	0.0155	0.0016	4.3736e-22	TRUE
rs2245368_T	-0.0133	0.0016	2.0238e-16	TRUE
rs9641123_G	-0.0052	0.0016	1.1563e-03	FALSE
rs6465468_G	-0.0032	0.0016	4.6521e-02	FALSE

SNP	π_0	$SE(\pi_0)$	Pr_{π_0}	Replication?
rs17405819_T	0.0137	0.0016	1.2564e-17	TRUE
rs16907751_C	0.0074	0.0016	4.3644e-06	TRUE
rs2033732_C	0.0065	0.0016	4.8454e-05	TRUE
rs4740619_T	0.0140	0.0016	2.4629e-18	TRUE
rs10968576_A	-0.0149	0.0016	1.5486e-20	TRUE
rs6477694_T	-0.0088	0.0016	4.7682e-08	TRUE
rs1928295_T	0.0083	0.0016	1.9969e-07	TRUE
rs10733682_G	-0.0101	0.0017	1.0969e-09	TRUE
rs7899106_A	-0.0086	0.0016	1.0639e-07	TRUE
rs17094222_T	-0.0070	0.0016	1.2704e-05	TRUE
rs11191560_T	-0.0085	0.0016	1.2063e-07	TRUE
rs7903146_C	0.0094	0.0016	4.1177e-09	TRUE
rs4256980_G	0.0113	0.0016	1.8886e-12	TRUE
rs11030104_A	0.0219	0.0016	1.6990e-42	TRUE
rs2176598_C	-0.0136	0.0016	1.9700e-17	TRUE
rs3817334_C	-0.0179	0.0016	6.0633e-29	TRUE
rs12286929_G	0.0113	0.0016	2.4293e-12	TRUE
rs7138803_G	-0.0179	0.0016	1.0766e-28	TRUE
rs11057405_G	0.0122	0.0016	2.3471e-14	TRUE
rs12429545_G	-0.0125	0.0016	1.3652e-14	TRUE
rs9540493_G	-0.0085	0.0016	1.2686e-07	TRUE
rs1441264_A	0.0126	0.0017	4.0919e-14	TRUE
rs10132280_C	0.0143	0.0016	1.3063e-18	TRUE
rs12885454_C	0.0110	0.0016	5.7772e-12	TRUE
rs11847697_C	-0.0072	0.0016	7.7682e-06	TRUE
rs7141420_T	0.0145	0.0016	5.3645e-19	TRUE
rs3736485_G	-0.0082	0.0016	3.5057e-07	TRUE
rs16951275_T	0.0159	0.0016	3.9682e-23	TRUE
rs7164727_T	0.0104	0.0016	9.0124e-11	TRUE
rs758747_C	-0.0088	0.0016	7.5808e-08	TRUE
rs12446632_G	0.0149	0.0016	2.1385e-20	TRUE
rs2650492_G	-0.0124	0.0016	1.7199e-14	TRUE
rs3888190_C	-0.0191	0.0016	1.1586e-32	TRUE
rs4787491_G	0.0126	0.0016	4.9604e-15	TRUE
rs9925964_A	0.0165	0.0016	1.0272e-24	TRUE
rs2080454_A	-0.0057	0.0016	4.8633e-04	TRUE
rs1558902_T	-0.0503	0.0016	1.0423e-214	TRUE
rs9914578_C	-0.0064	0.0016	6.7263e-05	TRUE
rs1000940_A	-0.0105	0.0016	5.9037e-11	TRUE
rs12940622_G	0.0129	0.0016	1.1756e-15	TRUE
rs1808579_C	0.0131	0.0016	3.3014e-16	TRUE
rs7239883_A	-0.0073	0.0016	6.8097e-06	TRUE
rs7243357_T	0.0092	0.0016	9.8026e-09	TRUE
rs6567160_T	-0.0311	0.0016	5.0802e-83	TRUE
rs17724992_A	0.0111	0.0016	5.1966e-12	TRUE
rs29941_G	0.0093	0.0016	7.1063e-09	TRUE
rs2075650_A	0.0113	0.0016	1.7586e-12	TRUE
rs2287019_C	0.0183	0.0016	4.9748e-30	TRUE
rs3810291_A	0.0191	0.0016	8.5285e-33	TRUE
rs6091540_C	0.0134	0.0016	1.1496e-16	TRUE
rs2836754_C	0.0076	0.0016	1.8681e-06	TRUE

References

1. Wainschtein, P. *et al.* Recovery of trait heritability from whole genome sequence data. *BioRxiv,* 588020 (2019).

2. Tucker-Drob, E. M. Measurement Error Correction of Genome-Wide Polygenic Scores in Prediction Samples. *bioRxiv,* 165472 (2017).

3. Tropf, F. C. *et al.* Hidden heritability due to heterogeneity across seven populations. *Nature human behaviour* **1,** 757–765 (2017).

4. Greene, W. H. *Econometric Analysis* Fifth (McGraw-Hill. Inc. New York, 2002).

5. Komashko, O. nl WaldTest: Wald Test of Nonlinear restrictions and Nonlinear CI. *R package version* **1** (2016).

6. Fisher, R. A. The correlation between relatives on the supposition of Mendelian inheritance. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* **52,** 399–433 (1919).

7. Okbay, A. *et al.* Genome-wide association study identifies 74 loci associated with educational attainment. *Nature* **533,** 539–542 (2016).

8. Bugliari, D. *et al.* RAND HRS data documentation, version P. *RAND Center for the Study of Aging* (2016).

9. Flegal, K. M., Kruszon-Moran, D., Carroll, M. D., Fryar, C. D. & Ogden, C. L. Trends in obesity among adults in the United States, 2005 to 2014. *Jama* **315,** 2284–2291 (2016).

10. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562,** 203–209 (2018).

11. Locke, A. E. *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature* **518,** 197–206 (2015).

12. Mallard, T. T. *et al.* Multivariate GWAS of psychiatric disorders and their cardinal symptoms reveal two dimensions of cross-cutting genetic liabilities. *bioRxiv,* 603134 (2020).

13. Gelman, A., Meng, X.-L. & Stern, H. Posterior predictive assessment of model fitness via realized discrepancies. *Statistica sinica,* 733–760 (1996).

14. Johnson, R., Sotoudeh, R. & Conley, D. Polygenic Scores for Plasticity: A New Tool for Studying Gene-Environment Interplay. *bioRxiv.*