Phenology and flowering overlap drive specialization in pollinator networks

Paul Glaum┼* 1,2 & Thomas J. Wood┼* 3, Jonathan R. Morris 4 & Fernanda S. Valdovinos 1,2

1. Dept. of Environmental Science & Policy, University of California-Davis, Davis, CA 95616
2. Dept. of Ecology & Evolutionary Biology, University of Michigan, Ann Arbor, MI 48109
3. Laboratory of Zoology, University of Mons, Mons, 7000, Belgium
4. School for Environment and Sustainability, University of Michigan, Ann Arbor, MI 48109

* These authors contributed equally to this work and are listed alphabetically.

┼Materials and Correspondence to:
Paul Glaum – prglaum@ucdavis.edu
Thomas J. Wood – thomasjames.wood@umons.ac.be

Classification: Article
Keywords: diet breadth, specialization, phenology, temporal resource density, pollination networks, pollen, seasonality, adaptive foraging

Variation in diet breadth and specialization stems from fundamental interactions species

have with their environment[1-3]. Consequently, understanding the drivers of this variation is key to

understanding ecological and evolutionary processes, and will facilitate the development of

predictive tools as ecological networks respond to environmental change[4,5]. Diet breadth in wild

bees has been an area of focus due to both their close mutualistic dependence on plants, and

because both groups are under threat from global biodiversity loss[6]. Though many of the principles

governing specialization for pollinators have been identified[7,8], they remain largely unvalidated.

Using mechanistic models of adaptive foraging in pollinators[9,10], we show that while temporal

resource overlap has little impact on specialization in pollinators with extended flight periods,

reduced overlap increases specialization as pollinator flight periods decrease. These results are

corroborated empirically using pollen load data taken from bees with shorter (genus *Andrena*) and

longer (genus *Lasioglossum*) flight periods across environments with both high and low temporal

resource overlap. This approach reveals how interacting phenologies structure plant-pollinator

networks and drive pollinator diet breadth via the temporal overlap of floral resources.

The dietary specialization of an organism sits at the nexus of numerous fundamental processes in

ecology and evolutionary biology. Diet breadth helps define organismal niches[1], mediates migration and

dispersal potential[2,11], shapes inter- and intraspecific competition[3,12], influences species persistence in the

face of environmental disturbance[4], and can affect rates of speciation and extinction[5]. Even the simple

assembly of the collection of dietary ranges of organisms in a community defines one of the major

concepts in ecology, the food web. Therefore, identifying the drivers influencing diet breadth and

specialization provides fundamental steps towards understanding a multitude of essential biological

questions.

Much of the progress in understanding specialization in consumers comes from studies of

herbivory[1], particularly on insects[13]. Using wild bees and plant-pollinator networks as our study system,

we extend the scope of those studies to include mutualistic interactions. A mutualism is a ubiquitous

ecological interaction in which participant species benefit each other[14], and pollination therefore differs

from other insect herbivory because of the reproductive benefit offered to both consumer and consumed

species. Bees depend on removing pollen from flowers to rear their offspring, providing a pollination

service at the same time[15]. Pollination is also of particular interest given its role in supporting terrestrial

biodiversity[6] and agricultural output[16], a role threatened by widespread declines in both pollinators and

insect pollinated plants[6,17].

Research on insect herbivores has largely focused on how adaptation to plant traits and defenses

drives consumer concentration on single plant families[18,19]. However, other drivers of diet breadth have

been identified (ref **7,8**, see Table S1). Here, we expand on the fundamental effect of resource density on

a consumer's diet breadth and specialization[7], not in a spatial context, but temporally. This resource

density is important in its interaction with consumer phenology. For long-lived organisms (e.g., predatory

vertebrates), opportunities to access a single resource type principally depend on the resource's spatial

distribution. If a single resource type is physically dense, then a predator can specialize on it without

needing to turn to alternatives, whereas if it is patchily distributed then a strategy of specialization is less

optimal. However, for a comparatively short-lived organism like a bee whose adult flight period can be

measured in days, the distribution of a resource type in time is the principal determinant of accessibility.

A plant species that blooms for 30 days is *de facto* dense, continuously available resource for a bee

species whose flight period lasts for only 25 days, but sparse and patchily distributed for a species which

flies for 90 days. The temporal pattern of flowering can therefore be considered the functional equivalent

of resource density for short-lived organisms. These ideas have proven influential but require concrete

theoretical and empirical validation to solidify their conclusions.

Expanding upon this idea of temporal density, we hypothesize that the degree of temporal overlap

in the availability of different resources (flowering overlap among co-occurring plant species) in

combination with the fundamental life history phenologies of bees (adult flight period) determines

realized diet breadth. Applied to plant-pollinator systems, we hypothesize that reduced flowering overlap

will reduce the diet breadth of short-lived but not of long-lived pollinators (Fig. 1). Shorter-lived

pollinators will experience markedly different resource availability based on the degree of temporal

overlap of co-flowering species. High flowering overlap will allow shorter-lived pollinators a greater

number of options while low overlap will restrict options. Longer-lived pollinators will be less constricted

by the temporal overlap of co-occurring resources, often having equal number of options regardless of the

degree of overlap (Fig. 1). Importantly, our hypothesis does not indicate that flight period will necessarily

correspond with being more or less specialized. Instead, it describes how temporal resource overlap will

affect diet breadth as a function of a pollinator's flight period.

We employed a multi-faceted approach to test our hypothesis. First, we leverage advances in

modeling adaptive foraging in ecological networks[9,10] to directly address the phenological mechanism

driving diet breadth across different pollinator flight periods. Second, we utilize our intercontinental

pollen load dataset of shorter lived *Andrena* bees and longer lived *Lasioglossum* bees from both a highly

seasonal and a less seasonal environment which function as low and high temporal resource overlap

treatments, respectively, to empirically validate our model outputs.

Results

Overview

Plant-pollinator dynamics are modeled using a dynamic consumer-resource approach that

incorporates adaptive foraging of pollinators to mechanistically model pollinators' consumption of floral

rewards and reproductive services to plant species[9,10] (see Supplementary Methods Table S2). The

model's adaptive foraging mechanism serves as a useful tool to implement and measure changing

pollinator diet breadth (see Supplementary Methods). Phenology is integrated into the base model through

modified sinusoidal wave functions which produce unique phenologies for every plant i (T_{p_i}) and animal

pollinator j (T_{a_j}). These functions T_{p_i} and T_{a_i} regulate floral reward production, adaptive foraging rates,

pollinator visitation rates, and consequently, plant and pollinator reproduction across time (Tables S3-S4).

The malleability of our phenology functions facilitates direct control over the availability of specific

resources and the activity patterns of the animal pollinators over time in simulations (see Supplementary

Methods, Fig S1-S5). The quantitative degree of overlap in co-occurring floral resources can be measured

as the Total Resource Overlap (TRO) of the entire plant community, and the Averaged Resource Overlap

(ARO) per plant species. Activity patterns of animal pollinators are set up to produce a range of different

flight period lengths separated by differing lengths in between flight periods (Fig S6). Simulations use

three fully connected bipartite network frameworks across 2072 combinations of plant and pollinator

phenologies to produce 62160 unique networks used to model over 22 million plant-pollinator

interactions (see Supplementary Methods).

Our pollen load data (see Supplementary Methods) comprises a large sample set of two bee

genera, *Andrena* (Andrenidae) and *Lasioglossum* (Halictidae), collected both in the state of Michigan,

USA (Nearctic) and the United Kingdom (Palearctic). Restricting our geographic scale to these two

specific Holarctic locations aids in our ability to compare across both bee and plant communities due to

their shared evolutionary history and overall similarity (see Supplementary Methods). The highly

seasonal continental climate of Michigan[20] produces lower degrees of overlap and highly seasonal

flowering communities, while the more mild oceanic climate of the British Isles[20] and its consistent

temperatures produces longer, overlapping flowering times among plant species[21].

Diet breadth or the degree of specialization in pollinators is measured both in-model with output

on pollinator foraging effort and empirically with relative abundance of different pollen in bee pollen

loads. We used two metrics that were applicable to both theoretical and empirical results for better direct

comparisons, because simply counting unique taxonomic groups in the pollen data will not be applicable

to theoretical results. We developed the first metric and labeled it Deviation from Generalism (DFG) (see

Supplementary Methods, Table S5). Briefly, DFG is the normalized summation of all pairwise differences

of pollinator foraging effort (or relative pollen load) on potential floral resources (see Supplementary

Methods, Table S5). It ranges from 0 (perfect generalist) to 1 (perfect specialist). The second metric was

the Coefficient of Variation (CV) of a pollinator's foraging effort (or pollen load). Higher CV values

occur with outlying values indicating specialization. With model results, both DFG and CV were applied

to final foraging levels at the end of simulations as well as the average foraging levels across the last 1000

time steps (see Supplementary Methods, Fig S7).

Model Results

Our simulation results (Figs. 2, 3) support our hypothesis (Fig. 1) demonstrating strong effects of

temporal resource overlap on the diet breadth of short- but not long-lived pollinators. Fig. 2 shows the

results of one network as an illustration of how resource overlap explains large portions of the variation in

diet breadth of pollinators with shorter flight periods (Fig. 2a, c), but little for the diet breadth of

pollinators with longer flight periods (Fig. 2b, d). The transition from high explanatory power to low

explanatory power is apparent as pollinator flight period increases regardless if specialization was

measured at the end of simulations or averaged across the last 1000 model time steps (Fig 3). This result

is also consistent across varying proportional lengths of time between pollinator flight periods using

either the DFG or CV metric across all networks tested (Fig S8-Fig S10). As predicted, low levels of

resource overlap drive pollinators with short flight periods to specialize because potential resource

options are limited at any given moment in time. As temporal resource overlap increases, potential

options increase resulting in a more generalized diet breadth (Fig. 2a, c). Long-lived bees, on the other

hand, do not experience the same limitation of potential diet options due to temporal resource overlap,

because they are active most of the entire flowering season and can potentially access most or all of the

flowering plants. Results were consistent regardless of whether resource overlap was measured as total

resource overlap (TRO) or average resource overlap (ARO).

Pollen data results

Our empirical results support our hypothesis, showing more specialized diets of short-lived bees

in more seasonal environments (Michigan) with lower flowering overlap, than in less seasonal

environments (UK) with higher flowering overlap. Our pollen data from Michigan presents more single

family specialists than the less seasonal UK dataset (Fig S11). Dietary specialization between bee genera

and regions varied significantly at both the level of botanical family (χ^2=27.1, p<0.001, Fig 4a) and genus

(χ^2=25.3, p<0.001, Fig 4b). In accordance with our hypothesis, there was an effect of region at the

botanical family level, but only for the short-lived *Andrena* where species in Michigan were significantly

more specialized than their British counterparts. In contrast, there were no differences for long-lived

Lasioglossum. At the botanical genus level, the same trend was more strongly expressed, with *Andrena* in

the UK showing a more generalized diet than their Michigan counterparts, but with no differences from

Lasioglossum in either Michigan or the UK. Results were consistent when analyzed using the DFG metric

(Fig S12).

Discussion

188   Our model results mechanistically demonstrate how differing phenologies interact with varying

189   degrees of temporal resource overlap to alter pollinator diet breadth. These results are corroborated

190   through our regional pollen load data which also presents the first empirical validation of benchmark

191   theories on adaptive foraging by consumers in general[7] and pollinators in particular[8]. Furthermore, our

192   results suggest broader implications regarding the drivers of bee diet breadth and diversification.

193   Global patterns of herbivory, particularly in insects, show strong trends of specialization at the

botanical family level[13]. These patterns have been linked to phytochemical diversity[19,22,23] which

consequently drives an increase of both dietary specialization and species diversification in many

herbivorous insect groups across a latitudinal gradient towards the tropics, most clearly seen and well-

studied in Lepidoptera[22,24]. However, bees break sharply from this pattern, with a relatively low diversity

in the tropics when compared to Mediterranean and xeric environments[8,25,26]. Tropical environments are

dominated by highly generalized, often social species that visit a huge variety of botanical families[27,28].

Tropical areas have the highest levels of global phytochemical diversity, but their flowering patterns have

wider phenological variation than temperate areas at both the inter- and intraspecific levels. This results in

tropical floral communities that *de facto* show higher resource overlap in any particular year[29,30] than

temperate communities. Our theoretical and empirical results suggest that bees' strong departure from the

typical global pattern in herbivory[13] may be due to a relatively stronger effect of phenology on their diet

breadth than the phytochemical restrictions suggested for other insect herbivores. In fact, tropical bees

have not evolved anything approaching the same high degree of dietary specialization as seen in the

tropical Lepidoptera despite existing in the same extremely phytochemically diverse landscapes. For

further discussion of this point, see Supplementary Discussion.

Our modeling developments present a flexible operational basis going forward. Integration of

time dependent functional components into species behavior and traits show that even high dimensional

network models can be further expanded in tractable ways to include the dynamics of organismal

phenology. Additionally, our metrics for diet breadth complement each other (see Supplementary

Methods) and reflect past methods (Fig S10). These metrics also integrated well with empirical pollen

load data sets that can be used to vet theoretical predictions. By expanding pollen datasets globally and

incorporating empirically vetted plant-pollinator phenology, future research can address how foraging

competition occurs across overlapping phenologies and what effects changing climatic conditions can

have on network stability.

218
219   **Conflict of interest statement:** We declare no conflict of interest.
220
221   **Acknowledgements:** Part of this project was supported by the National Science Foundation grant DEB-
222   1834497 awarded to F.S.V.
223
224   **Author Contributions:** F.S.V. conceived the project. T.J.W., P.G., F.S.V., and J.R.M. developed the
225   conceptual basis for the project. T.J.W. identified suitable collection sites and bee genera for the empirical
226   study design, collected bee and pollen data, and identified specimens. P.G. and F.S.V. developed dynamic
227   model, simulation design, code, and diet breadth metrics. P.G. implemented simulations and analyses.
228   T.J.W. and P.G. wrote the first draft, and all authors edited and revised the manuscript.
229
230   **Supplementary Information:** Supplementary Methods, Supplementary Discussion, Supplementary
231   Figures and Tables.
232
233   **Data Availability:** Pollen load data is available as Supplementary Table S9.
234
235   **Code Availability:** Simulation code and simulation data are available at the repository:
236   https://github.com/fsvaldovinos/Phenology. Phenology parameters used in the simulation portion of our
237   study are available in Table S10.
238

239   **References**

240   1. Kartzinel, T.R. et al. DNA metabarcoding illuminates dietary niche partitioning by African large
241   herbivores. *PNAS* **112(26)**, 8019-8024 (2015).
242   2. Bommarco, R., Biesmeijer, J.C., Meyer, B., Potts, S.G., Pöyry, J., Roberts, S.P., Steffan-
243   Dewenter, I. & Öckinger, E.Dispersal capacity and diet breadth modify the response of wild bees
244   to habitat loss. *Proc. Roy. Soc. B* **277(1690)**, 2075-2082 (2010).

3. Svanbäck, R. & Bolnik, D.I. Intraspecific competition drives increased resource use diversity within a natural population. *Proc. Roy. Soc. B* **274(1161)**, 839-844 (2007).
4. Wood, T.J. et al. Narrow pollen diets are associated with declining Midwestern bumble bee species. *Ecology* **100(6)** e02697 (2019).
5. Thompson, J. Rapid evolution as an ecological process. *TREE* **13(8)**, 329-332 (1998).
6. Biesmeijer, J.C., Roberts, S.P., Reemer, M., Ohlemüller, R., Edwards, M., Peeters, T., Schaffers, A.P., Potts, S.G., Kleukers, R., Thomas, C.D. & Settele, J. Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science* **313(5785)**, 351-354 (2006).
7. MacArthur, R.H. & Pianka, E.R. On optimal use of a patchy environment. *The American Naturalist* **100(916)**, 603-609 (1966).
8. Waser, N.M., Chittka, L., Price, M.V., Williams, N.M. & Ollerton, J. Generalization in pollination systems, and why it matters. *Ecology* **77(4)**, 1043-1060 (1996).
9. Valdovinos, F.S., Moisset de Espanés, P., Flores, J.D. & Ramos-Jiliberto, R. Adaptive foraging allows the maintenance of biodiversity of pollination networks. *Oikos* **122(6)**, 907-917 (2013).
10. Valdovinos, F.S., Brosi, B.J., Briggs, H.M., Moisset de Espanés, P., Ramos-Jiliberto, R. & Martinez, N.D. Niche partitioning due to adaptive foraging reverses effects of nestedness and connectance on pollination network stability. *Ecology Letters* **19**, 1277-1286 (2016).
11. Boyle, W.A., Conway, C.J. & Bronstein, J.L Why do some, but not all, tropical birds migrate? A comparative study of diet breadth and fruit preference. *Evolutionary Ecology* **25(1)**, 219-236 (2011).
12. deVries, M.S. The role of feeding morphology and competition in governing the diet breadth of sympatric stomatopod crustaceans. *Biology Letters* **13(4)**, 20170055 (2017).
13. Forister, M.L., Novotny, V., Panorska, A.K., Baje, L., Basset, Y., Butterill, P.T., Cizek, L., Coley, P.D., Dem, F., Diniz, I.R. and Drozd, P. The global distribution of diet breadth in insect herbivores *PNAS* **112(2)**, 442-447 (2015).
14. Bronstein, J.L. Our current understanding of mutualism. *The Quarterly Review of Biology* **69(1)**, 31-51 (1994).
15. Westerkamp, C. Pollen in bee–flower relations. Some considerations on melittophily. *Bot. Acta.* **109**, 325–332 (1996).
16. Garibaldi, L.A., Steffan-Dewenter, I., Winfree, R., Aizen, M.A., Bommarco, R., Cunningham, S.A., Kremen, C., Carvalheiro, L.G., Harder, L.D., Afik, O. & Bartomeus, I. Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science* **339**, 1608-1611 (2013).
17. Burkle, L. A., Marlin, J. C., & Knight, T. M. Plant-pollinator interactions over 120 years: loss of species, co-occurrence, and function. *Science* **339** (6127), 1611-1615 (2013).
18. Ali, J.G. & Agrawal, A.A. Specialist versus generalist insect herbivores and plant defense. *Trends in Plant Science* **17(5)**, 293-302 (2012).
19. Hunter, M. The Phytochemical Landscape: Linking Trophic Interactions and Nutrient Dynamics. Princeton University Press, 376 pp (2016).
20. Beck, H., Zimmermann, N., McVicar, T. et al. Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Sci Data* **5**, 180214 https://doi.org/10.1038/sdata.2018.214 (2018).
21. Baude, M., Kunin, W.E., Boatman, N.D., Conyers, S., Davies, N., Gillespie, M.A., Morton, R.D., Smart, S.M. & Memmott, J. Historical nectar assessment reveals the fall and rise of floral resources in Britain. *Nature* **530**, 85-88 (2016).
22. Becerra, J. On the factors that promote the diversity of herbivorous insects and plants in tropical forests. *PNAS* **112(19)**, 6098-6103 (2015).
23. Richards, L.A., Dyer, L.A., Forister, M.L., Smilanich, A.M., Dodson, C.D., Leonard, M.D. & Jeffrey, C.S. Phytochemical diversity drives plant–insect community diversity. *PNAS* **112(35)**, 10973-10978 (2015).

24. Dyer, L.A., Singer, M.S., Lill, J.T., Stireman, J.O., Gentry, G.L., Marquis, R.J., Ricklefs, R.E.,
Greeney, H.F., Wagner, D.L., Morais, H.C. & Diniz, I.R. Host specificity of Lepidoptera in
tropical and temperate forests. *Nature* **448(7154)**, 696-699 (2007).
25. Heithaus, E.R. Community structure of neotropical flower visiting bees and wasps: diversity and
phenology. *Ecology* **60(1)**, 190-202 (1979).
26. Moldenke, A.R. The role of host-plant selection in bee speciation processes. *Phytologia*, **43**, 433-
460 (1979).
27. Eltz, T., Brühl, C.A., van der Kaars, S., Chey, V.K. & Linsenmair, K.E. Pollen foraging and
resource partitioning of stingless bees in relation to flowering dynamics in a Southeast Asian
tropical rainforest. *Insectes sociaux* **48**, 273-279 (2001).
28. Galgani-Barraza, P., Moreno, J.E., Lobo, S., Tribaldos, W., Roubik, D.W. & Wcislo, W.T.
Flower use by late nineteenth-century orchid bees (Eufriesea surinamensis, Hymenoptera,
Apidae) nesting in the Catedral Basílica Santa María la Antigua de Panamá. *Journal of
Hymenoptera Research* **74**, 65-81 (2019).
29. Newstrom, L.E., Frankie, G.W. & Baker, H.G. A new classification for plant phenology based on
flowering patterns in lowland tropical rain forest trees at La Selva, Costa Rica. *Biotropica* **26**,
141-159 (1994).
30. Stevenson, P.R., Castellanos, M.C., Cortés, A.I. & Link, A. Flowering patterns in a seasonal
tropical lowland forest in western Amazonia. *Biotropica* **40**, 559-567 (2008).

345 **Figures & Figure Legends**
346

347
348
Figure 1: Graphical diagram of our hypothesis. Our hypothesis stems from the temporal overlap of
flowering resources available to pollinators. Highly seasonal delimitations between different flowering
populations create low levels of temporal overlap amongst different flowering resources. On the other
hand, fewer seasonal delimitations between different flowering populations create higher levels of
temporal overlap amongst different flowering resources. a) In long flight-period pollinators (e.g.
Lasioglossum), differing levels of temporal resource overlap have limited to no effect on the degree of
diet options available to the pollinator. b) This is predicted to lead to a non-significant relationship
between overlap and specialization in long flight period pollinators. c) In short flight-period pollinators
(e.g. *Andrena*), low and high temporal resource overlap can limit or expand pollinator diet options
respectively. d) This is predicted to lead to a significant relationship between overlap and specialization in
short flight-period pollinators.
360

361
362
Figure 2: Temporal resource overlap drives specialization more strongly in short-lived than in long-
lived pollinators. Model output on pollinator specialization taken across average plant overlap (ARO) for
short flight period (flight period = 20 time steps) and long flight period pollinators (flight period = 300
time steps). Each dot represents the diet breadth metric of a single pollinator population based on their
foraging effort per plant species. Diet breadth was measured at the end of simulations using **a)** DFG in a
short flight period bee, F-statistic: 1.05e+05 on 1 and 18498 DF, p-value: $< 2.2e\text{-}16$, **b)** DFG in a long
flight period bee, F-statistic: 1089 on 1 and 18498 DF, p-value: $< 2.2e\text{-}16$, **c)** CV in a short flight period
bee, Residual deviance: 671.94 on 18498 degrees of freedom, **d)** CV in a long flight period bee, Residual
deviance: 8244.4 on 18498 degrees of freedom. Example data taken from a network framework with 30
plant and 50 pollinator species. Higher values of both DFG and CV metrics indicate higher levels of
specialization (narrower diet breadth) while lower values indicate a more generalized feeding range
(wider diet breadth).
375

376
377
Figure 3: Interactive effect of temporal resource overlap and pollinator flight period on pollinator diet breadth. Representative sample of generalized linear models (Gamma distribution with log link functions) showing change in effect of Total Resource Overlap (TRO) on pollinator diet breadth in model simulations. Flight period labels indicate the length of the flight period and the number of phenology cycles between subsequent flight periods (see Supplementary Methods). Diet breadth shown as the degree of specialization measured by the CV of pollinator foraging effort per plant species taken **a)** at the end of simulations and **b)** averaged across the last 1000 time steps. Higher CV values correspond with a higher level of specialization (narrower diet breadth) while lower CV values indicate a more generalized feeding range (wider diet breadth). Example data taken from a network framework with 30 plant and 50 pollinator species.

389
390
Figure 4: Empirical measures of dietary specialization. Short-lived *Andrena* bees show higher levels
of specialization in highly seasonal (Michigan) compared to the weakly seasonal (UK), with no
differences for long-lived *Lasioglossum* bees using the CV metric. Moreover, only in Michigan is there a
difference in specialization between the two bee genera, showing a minimal impact of flight period in
weakly seasonal environments. Effects are consistent at both the (a) botanical family and (b) botanical
genus level. Bars headed by different letters are significantly different (p<0.05). Error bars show standard
error.
398