Title

Regulation of *Wolbachia* proliferation by the amplification and deletion of an
addictive genomic island

Authors

Elves H Duarte[1,2*], Ana Carvalho[1], Sergio López-Madrigal[1], Luís Teixeira[1,3*]

(1) Instituto Gulbenkian de Ciência, Oeiras, Portugal.

(2) Faculdade de Ciências e Tecnologia, Universidade de Cabo Verde, Palmarejo, Cabo
Verde.

(3) Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal.

* corresponding authors:

Elves H Duarte: ehelegam@gmail.com

Luís Teixeira: lteixeira@igc.gulbenkian.pt

Abstract

Wolbachia is one of the most prevalent bacterial endosymbionts, infecting approximately
40% of terrestrial arthropod species. *Wolbachia* is often a reproductive parasite but can
also provide fitness benefits to its host, as for example protection against viral pathogens.
This protective effect is currently being applied to fight arboviruses transmission by
releasing *Wolbachia*-transinfected mosquitoes. Titre regulation is a crucial aspect of
Wolbachia biology. Higher titres can lead to stronger phenotypes and fidelity of
transmission but can have a cost to the host. Since *Wolbachia* is maternally transmitted,
its fitness depends on host fitness, and, therefore, its cost to the host needs to be
controlled. Understanding this and other aspects of *Wolbachia* biology has been hampered
by the lack of genetic tools. Here we developed a new forward genetic screen to identify
Wolbachia over-proliferative mutant variants. We characterized in detail two of these new
mutants, *w*MelPop2 and *w*MelOctoless, and show that the amplification or loss of the
Octomom genomic region causes their over-proliferation. These results confirm previous

data and expand on the complex role of this genomic region in the control of *Wolbachia* proliferation. Both new mutants shorten the host lifespan and increase antiviral protection. Moreover, we show that *Wolbachia* proliferation rate in *Drosophila melanogaster* depends on the interaction between Octomom copy number, the host developmental stage, and temperature. Our analysis also suggests that the life shortening phenotype and antiviral protection of *Wolbachia* are dependent on related, but different, properties of the endosymbiont; the rate of proliferation and the titres at time of infection, respectively. Altogether, we demonstrate the feasibility of a novel and unbiased experimental approach to study *Wolbachia* biology, which can be further adapted to characterize other genetically intractable bacterial endosymbionts.

Introduction

Intracellular maternally-transmitted bacterial symbionts (bacterial endosymbionts) are widespread in insects [1]. These bacteria can be mutualistic by complementing the diets of their hosts, or by increasing fitness of their hosts in other particular ecological contexts, and may expand the range of ecological niches of their insect hosts [1]. Alternatively, bacterial endosymbionts can be parasitic, many times manipulating the reproduction of their hosts and providing a benefit to infected females over uninfected counterparts [1]. Understanding the interaction of endosymbionts with their host is crucial to understand much of insect biology. A key aspect of this interaction is the regulation of endosymbiont titres which influence the strength of induced phenotypes and cost to the hosts [2,3].

Wolbachia is one of the most prevalent bacterial endosymbionts in arthropods, being found in approximately 40% of terrestrial arthropod species [4]. *Wolbachia* is broadly known as a host reproduction manipulator [5]. However it can also be mutualistic, by, for example, providing a vitamin to its host [6] or protection against viral pathogens [7,8].

Since insects are vectors of many viruses, the discovery of *Wolbachia* antiviral phenotype, in *Drosophila melanogaster*, boosted its use as a vector control agent [9]. *Aedes aegypti* mosquitoes trans-infected with *Wolbachia* have increased resistance to viruses, including dengue, chikungunya, Zika, and yellow fever viruses, and, therefore, reduced vectorial capacity [10–13]. Release of *Wolbachia*-infected mosquitoes in dengue-endemic areas is likely to reduce dengue burden [14,15]. Despite the preliminary successful results of this strategy we still lack knowledge on several fundamental aspects of *Wolbachia* biology and interaction with viral pathogens, which hinders predicting the long-term outcome of *Wolbachia*-based interventions in vector control.

Wolbachia titres are a critical factor regulating its biology and interaction with the host [3]. Titres correlate positively with transmission fidelity and the strength of induced phenotypes, including the *Wolbachia* pathogen blocking phenotype [3,16–20]. In contrast, higher titres are associated with a reduction in host lifespan [16,17,21,22]. This may have a cost to *Wolbachia* because, since it is vertically transmitted, its fitness depends on the host fitness. Thus, *Wolbachia* titres regulation by the symbiont or the host may be under selection. The interaction between titres, cost, and strength of antiviral resistance is also relevant for the deployment of *Wolbachia*-carrying mosquitoes to block arboviruses transmission. Although several host and environmental factors (e.g. temperature) have been shown to affect *Wolbachia* titres, less is known about *Wolbachia* genetic factors [3].

Comparison of *Wolbachia* variants in controlled conditions (e.g. same host genetic background) shows the role of *Wolbachia* genetic factors in titre regulation [16,16,23]. So far, a single *Wolbachia* genetic factor, the Octomom region, has been shown to influence proliferation [16,17]. This region, comprised of eight predicted genes, is amplified in the pathogenic *w*MelPop. The degree of the Octomom region amplification determines the proliferation rate of *w*MelPop and the extent of its life shortening phenotype [17].

The genetic intractability of *Wolbachia*, which remains unculturable so far, hampers the identification of more genetic modifications altering *Wolbachia* proliferation. Hence, unbiased approaches such as genetic screen could contribute to our understanding of the genetic bases of *Wolbachia*-host interactions. Here, we developed a screening strategy in *Wolbachia* to isolate novel over-proliferating variants. The strategy was based on random mutagenesis, which has been applied before to other unculturable bacteria [24]. We fed the mutagen ethyl methanesulfonate (EMS) to *D. melanogaster* females carrying *Wolbachia* and screened for over-proliferative *Wolbachia* in their progeny. This approach allowed us to isolate new mutant over-proliferating *Wolbachia* variants. We identified the causative genetic changes in *Wolbachia* for this over-proliferation and made a detailed phenotypical characterization in terms of growth and antiviral protection. We identified a new mutation leading to *Wolbachia* over-proliferation and revealed a complex role for the Octomom region in the regulation of *Wolbachia* growth. Moreover, we demonstrated the feasibility of a novel and unbiased experimental approach to study *Wolbachia* biology.

Results

Isolation of over-proliferative *Wolbachia* in an unbiased forward genetic screen

We implemented a classical forward genetic screen in order to isolate new over-proliferative *Wolbachia* variants. We attempted to mutagenize *Wolbachia* by feeding the mutagen EMS to *Wolbachia*-carrying *D. melanogaster* females. EMS is extensively used in *D. melanogaster* [25] and has been previously used to mutagenize intracellular bacteria in cell culture [24]. We then tested *Wolbachia* titres, by real-time quantitative PCR (qPCR), in the progeny of treated females, since this bacterium is maternally transmitted. We used flies with the variant *w*MelCS_b as our starting variant because of its genetic proximity to the over-proliferative and virulent *w*MelPop variant [16,22,26] and its potential to easily become over-proliferative [17].

Putative mutagenized *Wolbachia* cells within the host would be in a mixed population, which would make it harder to assess their specific phenotype. However, we hypothesized that over-proliferating *Wolbachia* cells would overtake the population and that the resulting higher titres could be detectable. Moreover, we decided to pre-treat some of the EMS exposed females with tetracycline to reduce the *Wolbachia* population in these females and their progeny. This *Wolbachia* titre reduction should decrease competition for any new mutated *Wolbachia*, increase drift, and maybe facilitate fixation of new variants. To set up the conditions for tetracycline treatment we tested different doses of this antibiotic on females, without EMS. The titre of *Wolbachia* in the progeny of treated females are reduced to approximately 0 to 90% of the titres in controls (S1 Fig, $p < 0.001$ for all doses compared with control, at generation 1). We then followed the subsequent progeny of these flies to test how many fly generations it takes to recover normal *Wolbachia* titres. Except for higher tetracycline doses which lead to infection loss, *Wolbachia* titres recovered to normal within four fly generations (S1 Fig; linear mixed model [lmm], $p > 0.05$ for all doses compared with control at generation 4).

To optimize the screen, we also tested for the effect of different EMS doses on the fecundity of *Drosophila* females and *Wolbachia* titres. We observed that increasing doses of EMS reduce female fecundity (S2A-B Fig, linear model [lm], $p < 0.001$ for both egg number and adult progeny per female). Moreover, we found that EMS feeding strongly reduces *Wolbachia* titres in the next generation, in a dose-dependent manner (S2C-D Fig, non linear model [nls] fit, $p < 0.001$). Titres were reduced by up to 90% when 8,000 mM EMS was supplied, leading to the loss of *Wolbachia* in the next generation in some lines (S2C-D Fig). This titre reduction due to the EMS treatment could hamper the identification

of over-proliferating *Wolbachia* in a screen in the first generation after treatment. Given these results and the recovery time after tetracycline treatment detailed above, we quantified *Wolbachia* titres at the first generation (F1), the immediate progeny of EMS-treated females, and at the fourth generation after treatment (F4) were we would expect *Wolbachia* titres to recover after the severe reduction due to EMS treatment.

We screened approximately 1000 F1 progeny of EMS-treated females, in a range of experimental conditions, and at least one F4 female descendent. We varied EMS dose from 10 mM to 8,000 mM, and tetracycline dose from 0 µg/ml to 12.5 µg/ml, in different combinations (S1 Table). The relative *Wolbachia* titre was determined when females were ten days old after they laid eggs so that any putatively interesting progeny could be followed up.

In three independent batches of EMS-treated flies, we detected females with 3 to 14-fold more *Wolbachia* than controls, suggesting the presence of over-proliferative variants (Fig 1 and S3 Fig). In two batches, over-proliferating *Wolbachia* was identified in the F1 and in the other batch in the F4. We assessed Wolbachia titres in the next generation and found that the over-proliferative phenotype was inherited. Subsequent selections allowed us to establish *Drosophila* lines carrying new potentially over-proliferative *Wolbachia* variants.

Fig 1. Isolation of over-proliferative *Wolbachia* variants by a forward genetic screen.
(A and B) Relative *Wolbachia* titres in a control (*w*MelCS_b) and EMS-treated flies (Lines 1A and 2A). 5–10 virgin females were randomly collected each generation for egg-laying and *Wolbachia* titre measured using qPCR. Bacterial titres are normalized to that of control flies. The progeny of the female with the higher *Wolbachia* titre was used to set up the next generation. The selection of the other putative over-proliferating Wolbachia line in panel B in shown in S3A Fig. (C) Relative titres of over-proliferating *Wolbachia* variants in an isogenic genetic background. Both lines kept the over-proliferative phenotype (p < 0.001). Each dot represents Wolbachia titre of a single female.

We designed the screen to find new mutants of *Wolbachia,* and not host mutants, that lead
to the endosymbiont proliferation. However, EMS will most likely also induce mutations in
the host germline, in the nuclear or mitochondrial genomes, that can be transmitted. To
minimize the influence of new host nuclear mutations on our screen, we backcrossed the
EMS-treated females and their progeny, at every generation, with males from the control
isogenic line. To verify that new mutations in the host were not the cause of *Wolbachia*
over-proliferation, we replaced the first, second and third chromosomes of *Drosophila*
females carrying the over-proliferating *Wolbachia* variants, in line 1A and line 2A, with the
chromosomes of the control line, through the use of balancer chromosomes (S4 Fig.). We
repeated *Wolbachia* titres quantification and found that the over-proliferative phenotypes
were maintained (Fig 1C; lmm, p-value< 0.001 for both).

Since mitochondria are maternally transmitted, the experiments described above cannot
exclude the possibility that *Wolbachia* over-proliferation is mitochondria-determined
(directly or indirectly). Thus, the mitogenome of the lines showing higher *Wolbachia* titres
was Illumina sequenced and aligned to the mitochondrial reference genome release 6
(GenBank: KJ947872.2:1–14,000). A summary of the alignment features is given in S2
Table. We did not find SNPs or indels unique to the mitochondria of EMS-fed flies, which
shows that flies with over-proliferative *Wolbachia* did not inherit mutated mitochondria (S3
Table). Therefore, we concluded that the observed *Wolbachia* over-proliferative
phenotypes did not result from mutations in the host genome, neither nuclear nor
mitochondrial.

Identification of genetic basis of the new over proliferative variants

To identify the mutations associated with over-proliferation, we sequenced and assembled
the genomes of these over-proliferative *Wolbachia*. A summary of the sequencing results
is given in S2 table. We performed a hybrid assemble with short (Illumina) and long-reads
(Nanopore) and obtained single and circular *Wolbachia* chromosomes differing in length
(S4 Table).

To test our assembly pipeline we assembled a previously characterized Cluster III *w*Mel
Wolbachia variant, named *w*Mel [16], which derives from the line used for the original
*w*Mel reference genome (GenBank: AE017196.1) [27]. The new *w*Mel genome (GenBank:
CP046925.1) was also circular and comparable in size, structure and number of ORFs
with previously published *w*Mel genomes [27,28], including the *w*Mel reference genome
(S4 Table). We found, however, nine differences (2 SNPs and seven indels) relative to

*w*Mel reference genome, which we confirmed using Sanger sequencing (S5 Table). These
results validate our sequencing pipeline.

The only difference between the genome of the over-proliferative *Wolbachia* variant in Line
1A and *w*MelCS_b was an amplification of the Octomom region (Fig 2A and S1 Text).
There were three more copies of this region, giving a genome size difference of 62,814bp.
The Octomom region amplification, and lack of other differences, was also confirmed by
the mapping of the Illumina sequencing reads of Line 1A on the genome of wMelCS_b
(GenBank: CP046924.1) (Fig 2B). These results show that Octomom amplification is the
cause of over-proliferation, consistently with previous findings with the variant *w*MelPop
[16,17,29]. Given the nature of the genomic change inducing the over-proliferation we
named this variant *w*MelPop2.

As shown before for *w*MelPop [17], we observed variation in the Octomom copy number in
*w*MelPop2-carrying flies. For further analyses of the phenotype of these variant we
established, through selection (as in [17]), *Drosophila* lines carrying *w*MelPop2 with
specific low (2-3) and high Octomom (8-9) copy number (S5 Fig).

Interestingly, the genome of the over-proliferative *Wolbachia* variant in line 2A only differs
from *w*MelCS_b by a deletion of a 20,938bp genomic fragment that includes the full-length
Octomom region and one of its flaking direct repeats (Fig 2A and S1 Text). Mapping the
Illumina sequencing reads of this variant on the genome of *w*MelCS_b (GenBank:
CP046924.1) also confirmed this deletion has the only difference between the two (Fig
2B). The absence of all Octomom genes in this line is also confirmed by qPCR (S6 Fig).
These results identify Octomom loss as the cause of its over-proliferative phenotype.
Thus, we named this variant *w*MelOctoless.

Fig 2. Both amplification and deletion of Octomom lead to *Wolbachia* over-proliferation.
(A) Gene synteny across Octomom region and its flanking region. Assembled genomes of *w*MelCS_b,
*w*MelOctoless and *w*MelPop2 were aligned using Mauve v2.4.0 to identify mutations causing over-
proliferation. The three-fold amplification of Octomom in Line 1A (*w*MelPop2) and its deletion in Line 2A
(*w*MelOctoless) were the only mutations identified when compared to *w*MelCS_b. For illustrative purposes,
we used MultiGeneBlast v1.1.13 (http://multigeneblast.sourceforge.net/) to find, in Line 1A, 2A and a newly
assembled *w*Mel genome, homologous proteins to those of *w*Mel locus WD0501–WD0519 (GenBank:
AE017196.1). Genomes were annotated using the NCBI Prokaryotic Genome Annotation Pipeline v4.10. (B)
The relative coverage in a genomic region, including the Octomom region. Illumina paired-end reads were
mapped to *w*MelCS_b genome (GenBank: CP046924.1). We confirmed the amplification or deletion of
Octomom as the only difference between over-proliferating *Wolbachia* and *w*MelCS_b.

A second line isolated together with *w*MelOctoless, line 2B, also lost the Octomom region

(S7 Fig). And this was the only observed difference with *w*MelCS_b when mapping the

Illumina reads on *w*MelCS_b (S7 Fig). No differences were observed when the Illumina

reads were mapped to the *w*MelOctoless genome (GenBank: CP046923.1). Since

*w*MelOctoless and line 2B were identified in the same batch they may be not independent.

However, and importantly, we obtained the same results with another independent over-

proliferative line, isolated in a different batch of treatment, line 3A (S3 and S7 Fig). Mapping the Illumina sequence reads of this line identifies the loss of Octomom as the only difference with *w*MelCS_b, and identifies no difference to *w*MelOctoless. We named this line *w*MelOctoless2. These results further confirm that loss of the Octomom region leads to an over-proliferative phenotype in *Wolbachia*.

We also sequenced by short and longs reads and made a hybrid assembly of *w*MelPop genome and compared it to wMelPop2. We only detected and confirmed the two SNPs previously identified between *w*MelCS_b and *w*MelPop (position 920,191: T in *w*MelPop and C in *w*MelPop2; and position 1,005,339: A in *w*MelPop and G in *w*MelPop2 [16]). We also compared the mitogenome of flies carrying *w*MelPop and *w*MelPop2 and found one single substitution (position 10,793: G in wMelPop and A in wMelPop2) (Table S3). We confirmed this SNP using Sanger sequencing (S2 Text).

In summary, we were able to identify the genomic changes associated with the new over-proliferative variants and all map to loss or amplification of the Octomom region.

Deletion and amplification of the Octomom region differently impact titres and growth of *Wolbachia*

In order to characterize better the phenotypes of the new *Wolbachia* variants *w*MelOctoless and *w*MelPop2 we analysed their growth, together with *w*MelCS_b and *w*MelPop, in adult males kept at 18ºC, 25ºC, and 29ºC (Fig 3 and S8 Fig). These flies were all reared at 25ºC and placed at the different temperatures when 0-1 days old. At this initial point, at adult eclosion, there are differences in titres between lines carrying different copy numbers of Octomom genomic region (S9 Fig, $p < 0.028$ for all comparisons). Flies with *Wolbachia* with one copy of Octomom, *w*MelCS_b, have the lowest relative titre of *Wolbachia*. Flies with variants with low amplification of the Octomom region have approximately twice the titres of *Wolbachia*, while flies with variants with high copy number of this region have three times more *Wolbachia* than *w*MelCS_b. Flies with *w*MelOctoless have the highest titres, approximately 4-fold higher than flies carrying *w*MelCS_b. Therefore, the deletion or amplification of the Octomom region impact *Wolbachia* titres in adults at eclosion.

A

B

C

Fig 3. The amplification or deletion of Octomom increase *Wolbachia* proliferation rate.
Wolbachia proliferation dynamics at 18ºC (A), 25ºC (B) and 29ºC (C). *D. melanogaster* males used in these
experiments developed at 25ºC, were collected on the day of eclosion and aged to specific time-point at a
given temperature (18ºC, 25ºC or 29ºC). Ten males were collected at each time-point for *Wolbachia* titre
measurement using qPCR. *Wolbachia* titres were normalized to that of 0-1 days-old *w*MelCS_b-infected
males. A replicate of the experiment is given in S8 Fig. Exponential models were used to estimate Wolbachia
doubling time, and a summary of the results is given in Table 1. Both replicates were used for statistical
analysis. Each dot represents relative *Wolbachia* titre of a single male.

To analyse growth during adult life we fitted an exponential model to these data, from

which we also estimated doubling time of the *Wolbachia* variants, at the different

temperatures (Table 1). There is a wide range of doubling times for different *Wolbachia*

variants at different temperatures, from approximately one day to seventeen days. A model with all the data shows a complex interaction between growth rate, *Wolbachia* variant and temperature (lmm , $p < 0.001$).

Wolbachia variants	Adults			Larvae
	18ºC	25ºC	29ºC	25ºC
	Doubling time (days) (±SE)	Doubling time (days) (±SE)	Doubling time (days) (±SE)	Doubling time (days) (±SE)
*w*MelCS_b	16.79	13.90	4.07	0.68
	(14.80-19.39)	(11.49-17.59)	(3.67-4.56)	(0.60-0.79)
*w*MelOctoless	9.41	10.46	3.37	0.48
	(8.75-10.18)	(9.03-12.43)	(3.09-3.71)	(0.44-0.54)
*w*MelPop2 (3 copies)	17.68	3.50	1.28	n.d.
	(15.49-20.59)	(3.23-3.80)	(1.20-1.38)	
*w*MelPop2 (8-9 copies)	11.86	2.26	1.07	0.57
	(10.83-13.10)	(1.96-2.66)	(0.98-1.17)	(0.52-0.65)
*w*MelPop (2-3 copies)	15.44	5.30	1.38	n.d.
	(13.74-17.61)	(4.86-5.83)	(1.29-1.49)	
*w*MelPop (8-9 copies)	13.82	2.79	0.88	n.d.
	(12.43-15.56)	(2.35-3.44)	(0.82-0.95)	

Table 1 – Doubling time, in days, of *Wolbachia* variants at different temperatures or developmental stages. N.d. - not determined.

A direct comparison between *w*MelOctoless with *w*MelCS_b shows that this new variant replicates faster than *w*MelCS_b (lmm, p < 0.001), although it is a relatively small difference at all temperatures (in the full model with all variants, however, the growth of *w*MelOctoless and *w*MelCS_b is only statistically different at 18ºC, Table 1). Both strains interact equally with temperature. Their growth rate does not significantly change between 18ºC and 25ºC ($p = 0.94$), but increases at 29ºC ($p < 0.001$).

We also compared the effect of Octomom copy number in a model taking genotype (wMelCS_b, for the control and new variants generated here, or wMelPop) and Octomom copy number (zero, one, low and high) as factors. At 18ºC the amplification of the Octomom copy number does not lead to a statistically significant increase of growth rate, compared with wMelCS_b (lmm, $p > 0.125$ for both low and high Octomom copy number compared with control). At this temperature only the deletion of Octomom leads to a significantly higher growth, as seen above. However, variants with Octomom amplification

grow more at 25ºC than at 18ºC, and grow more at 29ºC than at 25ºC ($p < 0.001$ for all comparisons). At both 25ºC and 29ºC, variants with low Octomom copy numbers have a higher growth rate than the variants with one copy or deletion of Octomom region ($p < 0.001$ for all comparisons). At these temperatures the variants with high Octomom copy numbers grow faster than all the other variants ($p < 0.003$ for all comparisons). These results confirm that the degree of amplification of the Octomom region controls the intensity of the over-proliferation of these variants, as shown before [17]. The data also demonstrate a strong interaction between temperature and the increased proliferation of variants with amplification of the Octomom region. The effect of the amplification is not significant at 18ºC and becomes increasingly stronger with higher temperature. On the other hand, loss of Octomom leads to a smaller effect in growth, but similar at all temperatures. Therefore, although both genomic mutations lead to an increase in *Wolbachia* titres they have different impacts in the growth rates and their interaction with temperature.

The statistical model taking into account the *w*MelCs_b or *w*MelPop genotype, which differ only in two SNPs (see above), indicated a significant difference in growth between them, at 25ºC ($p < 0.001$). This could indicate that these two SNPs also influence growth of *Wolbachia*. However, this could also be due to the fact that the copy number of the Octomom region was not equally controlled in *w*MelPop and *w*MelPop2 lines during these experiments. *w*MelPop2 low copy number line had 2-3 copies of Octomom, while the wMelPop line had 3 copies. To test if *w*MelPop and *w*MelPop2 indeed vary in proliferation rate, we repeated this experiment with a more tightly controlled Octomom copy number in these two lines, at 25ºC (S10 Fig A-D). Both *w*MelPop and *w*MelPop2 carrying 3 copies of Octomom grow faster than wMelCS_b (lmm, $p < 0.001$ for both) and there is no difference in growth between them ($p = 0.32$). This indicates that the genetic differences between these lines do not affect their growth and that they are equally influenced by Octomom copy number.

Rapid proliferation of *Wolbachia* during the host development

We also analysed the growth of *w*MelCS_b, *w*MelOctoless and *w*MelPop2 (8-9 Octomom copies) during host development. *D. melanogaster* develops very fast from egg to adult, in only 10 days at 25ºC. We predicted that *Wolbachia* would grow much faster during this period than during adult life, in order to grow from the small population, present in the egg, to the population spread throughout the adult tissues. We, therefore, estimated absolute

numbers of *Wolbachia* genome copies in individuals at the different stages of development, using qPCR and a calibration curve. Assuming one genome per *Wolbachia* cell [30], these number correspond to *Wolbachia* cells. Embryos with 0-2h had between 4,000 and 4,700 *Wolbachia* genome copies on average, with no statistically significant difference between *Wolbachia* variants, (lm, *p* = 0.89 for the effect of *Wolbachia* variant, Fig 4, Table 2). Interestingly, at the end of the experiment, in newly eclosed adults, and as also observed above, there are significant differences between the three variants (lm, *p* < 0.009 for all comparisons, Table 2, Fig 4). Also, males carry significantly less *Wolbachia* than females (*p* = 0.029). Adults carried from approximately 600,000 to 4,500,000 *Wolbachia* genome copies.

Fig 4. *Wolbachia* proliferates rapidly during larval development.
Wolbachia genome copies per individual throughout *Drosophila* development. Dots represent either data from a pool of 10 individuals (eggs and larvae) or from a single individual (pupae and adults). *Wolbachia* proliferation in the first 120 hours were analysed using a exponential model. A summary is given in Table 1. Vertical dashed lines separate developmental stages (i.e. eggs, larvae, pupae, and adults).

| | Genome copy number (± SE) | | |
| | *Wolbachia* variants | | |
	*w*MelCS_b	*w*MelOctoless	*w*MelPop2
Embryos	4,100 (±1,200)	4,000 (±1,400)	4,700 (±1,500)
Adults			
Males	619,000 (±179,000)	3,334,000 (±975,000)	918,000 (±591,000)
Females	841,000 (±245,000)	4,528,000 (±928,000)	2,439,000 (±323,000)

Table 2. *Wolbachia* genome copies in embryos and in newly eclosed adult flies.

Mean (and standard error) of *Wolbachia* genome copies were estimated using qPCR.

Wolbachia growth seems to be restricted to the period between egg and white prepupae (120h), since there is no significant growth from this stage to adults (lm, *p* = 0.44). From eggs to white prepupae there is rapid exponential growth of all variants (Fig 5, Table 1). *w*MelCS_b has an estimated doubling time of approximately 16h, *w*MelPop2 of 14h, and *w*MelOctoless of 12h. These different doubling times probably explain how *Wolbachia* variants reach different amounts per individual host in adults, starting from the same estimated amount in embryos. However, in this analysis the difference between growth rates is not statistically significant (*p* = 0.10 for interaction between *Wolbachia* variants and growth). The growth rates of these variants are, therefore, very similar during this stage, and much faster than in adults. At the same temperature, we estimated doubling times in adults of *w*MelCS_b, *w*MelOctoless, and *w*MelPop2 (high-copy) to be, approximately, 13.9, 10.5, and 2.3 days, respectively. Therefore, *Wolbachia* growth at different stages of *D. melanogaster* can vary dramatically, and the different variants respond differently to host development stages.

We also asked if *Wolbachia* Octomom copy number changed in *w*MelCS_b and *w*MelPop2, throughout development, as *Wolbachia* is proliferating so fast, and found no evidence of so (lm, *p* = 0.88, S11 Fig). However, as shown before [31], during adult life there was a small increase of Octomom copy number with age, in *w*MelPop and *w*MelPop2 (an increase of 0.032 per day, lmer, *p* = 0.009, S11 Fig).

Wolbachia variants with a deletion or amplification of the Octomom region differently induce a life-shortening phenotype

The over-proliferation of *w*MelPop has been associated with a shortening of the host lifespan [16,22]. We, therefore, tested if these new over-proliferative variants also shorten

the lifespan of their host, at different temperatures, in males (Fig 5, S12 Fig). We also
performed this assay in females at 25ºC, with similar results to males at 25ºC (S12 Fig).
There was a significant interaction between survival, *Wolbachia* variant, and temperature
(Cox proportional hazard model with mixed effects (CHR), $p < 0.001$). All lines, including
the *Wolbachia*-free line have a shorter lifespan at 25ºC than 18ºC, and even shorter at
29ºC ($p < 0.001$ for all these comparisons). *w*MelCS_b did not affect the host lifespan at
any temperature ($p > 0.16$ for all comparisons with the line without *Wolbachia*).

*w*MelOctoless strongly reduces host lifespan at all tested temperatures ($p < 0.001$, each
comparison with *w*MelCS_b) (Fig 5, S12 Fig). This deleterious effect is stronger at 18ºC,
where wMelOctoless is the tested variant with the highest impact on lifespan, although
very similar and not statistically different from *w*MelPop2 with high Octomom copy number
($p < 0.001$, for all comparisons with other lines, $p = 1$ when compared with *w*MelPop2 with
high Octomom copy number). At 25ºC the effect of wMelOctoless on host lifespan is
weaker ($p = 0.001$), and similar at 25ºC and 29ºC ($p = 0.95$). These results demonstrate
that the new over-proliferative *w*MelOctoless also induces a cost to the host in terms of
lifespan and the effect interacts with temperature, being stronger at lower temperature.

Fig 5. *w*MelOctoless and *w*MelPop2 are pathogenic.
Lifespan of males with different *Wolbachia* variants at 18°C (A), 25°C (B), and 29°C (C). For survival analyses, fifty males were collected on the day of eclosion and kept in groups of 10 per vial until all flies had died. Flies were transferred to new vials every five days. A replicate of these experiments is given in S12A-C Fig. (D) Coefficients of a Cox mixed model, which represent the effect of *Wolbachia* on the lifespan of flies relative to the lifespan of *Wolbachia*-free flies. Both replicate experiments performed at each temperature were pooled for statistical comparisons. Bars represent the standard error of the coefficient, and letters statistical significant groups after *p*-value correction. A significant interaction between *Wolbachia* variant and temperature determined the lifespan (*p* < 0.001). (E-G) The correlation between the strength of life-shortening phenotype and *Wolbachia* doubling time at 18°C (E), 25°C (F), and 29°C (G). The y-axis represents the strength of *Wolbachia* life-shortening phenotype (estimated using Cox mixed models) and the x-axis *Wolbachia* doubling time (in days). The Pearson correlation coefficient (*r*) and its significance (*p*) are given in each panel.

*w*MelPop2, as *w*MelPop, also shortens host lifespan (Fig 5, S12 Fig). The variants containing high copy number of Octomom shorten lifespan at all temperatures (*p* < 0.001, for each comparison with *w*MelCS_b). This effect is much stronger at 25°C than at 18°C (*p* < 0.001 for contrasts between both lines *w*MelCS_b), and similar at 25°C and 29°C (*p* > 0.21 for these contrasts). At these two higher temperatures the lines carrying the variants

with high copy number of Octomom have the shortest lifespan of all tested lines ($p < 0.001$ for all comparisons). *w*MelPop2 and *w*MelPop with low copy number of Octomom always have a weaker effect than high copy number variants ($p < 0.001$ for all these comparisons), demonstrating the effect of the degree of amplification in this phenotype. As observed with the high copy number variants their effect increases with temperature and is stronger at 25ºC than 18ºC, and stronger at 29ºC than 25ºC ($p < 0.05$ for these comparisons). In fact, they are not pathogenic at 18ºC and an effect on host lifespan is only observed at 25ºC and 29ºC. These data confirm the association of Octomom region amplification with host lifespan shortening, an increase in the severity of the phenotype with increase in the Octomom copy number, and the increase in the severity of the phenotype with temperature.

*w*MelPop2 and *w*MelPop, high and low copy number variants, had significantly different effects in several comparisons at the three tested temperatures (Fig 5). This indicated that there could be differences in this phenotype between these two lines. However, the Octomom copy number was not exactly the same between these lines. Therefore, and as done above in the analysis of proliferation, we compared the lifespan phenotype in *w*MelPop2 and *w*MelPop lines with tightly controlled Octomom copy number (S10 Fig). At 25ºC lines with both *w*MelPop and *w*MelPop2 with 3 copies of the Octomom region had a shorter lifespan than the line with *w*MelCS_b ($p < 0.001$), and no difference between them ($p = 0.29$). These results show that *w*MelPop2 and *w*MelPop have the same phenotype.

To further demonstrate that the life shortening phenotypes were due to the new *Wolbachia* variants, and not an EMS-induced mutation in the host nuclear genome, we performed reciprocal crosses between flies with *w*MelCS_b and flies carrying either *w*MelOctoless or *w*MelPop2 (carrying 3 or 8-9 copies of Octomom) and followed the survival of their female progeny at 29ºC. Females progeny from reciprocal crosses should be identical in the nuclear genome but differ in the *Wolbachia* variant that is maternally transmitted. We observed maternal segregation of the life-shortening phenotype, thus demonstrating that the *Wolbachia* variant carried by the lines is the cause of the phenotype (S13 Fig). The relative strength of the life-shortening phenotype of these female progenies matches what was observed above for the maternal lines. Moreover, all the tested lines that inherited wMelCS_b had a similar lifespan ($p > 0.78$ for all comparisons), indicating no contribution of the host genotype in this set of experiments.

The life shortening phenotype of *w*MelPop has been associated with its over proliferation since its discovery [22]. We tested if these phenotypes were correlated by taking

advantage of the data on proliferation and shortening of lifespan, at different temperatures,
we collected from these set of variants. At all temperatures, we found a negative
correlation between the strength of the life-shortening phenotype and *Wolbachia* doubling
time (Fig 5, $|r| > 0.86$, $p < 0.03$, for all correlations). We found a much weaker or no
significant correlation between the strength of the life-shortening phenotype and
Wolbachia titres in 0-1 day-old adults (S14 Fig). These results show that over-proliferative
variants shorten the host lifespan and the strength of this phenotype correlates with their
proliferation rates.

Wolbachia variants with deletion or amplification of the Octomom region provide
stronger protection against DCV

Previous studies established a link between *Wolbachia* titres and the strength of anti-viral
protection [16–20]. To test if wMelOctoless and wMelPop also provide a stronger
protection against viruses, we infected flies with *Drosophila* C virus (DCV), by pricking, and
followed their survival for 40 days at 18ºC. All *Wolbachia* variants tested provided
protection against DCV (CHR, $p < 0.001$ for all comparisons with the *Wolbachia* free line*,*
Fig 6A and S15 Fig). Importantly, the survival rate of flies pricked with a buffer solution was
not different from each other (Fig 6B, 6D and S15B Fig, $p > 0.05$ for all comparisons).
*w*MelCS_b was the least protective variant, while *w*MelOctoless the one that provided the
highest protection. In general the over-proliferative *Wolbachia* variants confer stronger
protection to DCV than *w*MelCS_b, although this difference is not always significant (Fig
6C, $p < 0.001$ for all comparisons, except for *w*MelPop (2-3 copies), $p = 0.10$).

Fig 6. *w*MelOctoless and *w*MelPop2 provide strong protection against DCV.
Survival of males carrying different *Wolbachia* variants after a challenge with DCV (A) or a buffer solution
(B). 50 3-5 days-old *Drosophila* males were pricked with DCV (10^9 TCID$_{50}$/ml), or a buffer solution, and
survival curves were determined at 18°C for 40 days. A replicate of the experiment is given in S15 Fig. Both
survival curves of DCV-infected (C) or control-pricked flies (D) were analysed using Cox mixed models. The
resulting coefficients represent the effect of *Wolbachia* infection on survival relative to the *Wolbachia*-free
flies. Bars represent the standard error of the estimate, and the letters the statistical significant groups after
p-value correction. *Wolbachia* infection improved the survival of DCV-infected flies (p < 0.001) but did not
change the survival of buffer-pricked flies in this timeframe (*p* > 0.5). (E) Correlation between the strength of
anti-viral protection (represented as the coefficients of a Cox mixed model in the y-axis) and *Wolbachia* titres
at the day of eclosion, as a proxy for *Wolbachia* titer in the day of infection. The direction and strength of the
correlation were estimated using Person correlation coefficient.

We also tested the correlation between the antiviral protection and *Wolbachia* proliferation. We found no correlation at either 18ºC (p = 0.21), the temperature where the flies were kept after infection, or 25ºC (p = 0.35), the temperature in which flies developed and were kept until being infected with DCV (Fig S14). We also tested the correlation between the antiviral strength of *Wolbachia*-induced anti-viral protection with *Wolbachia* titres in 0-1 day-old flies, as a proxy for *Wolbachia* titre at the day of infection, which is 3-5 day-old flies. We found a significant correlation between these parameters (Fig 6F, p = 0.024, |r| = 0.87). Overall, the new over-proliferative variants give more protection to viruses than *w*MelCS_b, and the strength of this protection is correlated with *Wolbachia* levels at the time of infection.

Discussion

We developed a new forward genetic screen and identified new *Wolbachia* over-proliferative variants. We characterized in detail two of these new mutants, *w*MelPop2 and *w*MelOctoless, and identified the genetic bases of their over-proliferation. *w*MelPop2 had an amplification of the Octomom region, which has been previously shown to lead to over-proliferation in the wMelPop variant [16,17]. *w*MelOctoless, on the other hand, had a deletion of this same Octomom region. These results further confirm and develop the complex role of this genomic region in the control of *Wolbachia* proliferation. An extensive phenotypic characterization of two of these lines showed both *Wolbachia* variants to shorten the host lifespan, as well as to increase antiviral protection. Moreover, we show that *Wolbachia* proliferation rate in *D. melanogaster* depends on the interaction between Octomom copy number, the host developmental stage, and temperature. Our analysis also suggests that the life shortening and antiviral protection phenotypes of *Wolbachia* are dependent on the related, but different, properties of the endosymbiont, rate of proliferation and titres at time of infection, respectively.

An unbiased approach for genetically intractable symbionts

A difficulty in studying obligate intracellular bacteria is that they are often genetically intractable, mostly because of their dependency on the host cells to survive and difficulty in isolating and growing them clonally. We decided to try to directly mutagenize and screen *Wolbachia* in the host. The main difficulty of this approach is how to identify a new mutant and link it with a phenotype in the *Wolbachia* population present within a host. We

calculated here that newly emerged female adults, with *w*MelCS_b, carry approximately

840,000 *Wolbachia* genome copies, probably corresponding to the same number of

Wolbachia cells [30]. Since EMS induces random mutations, we expected mosaicism in

the *Wolbachia* population at the individual fly level. Each new mutant, when generated,

would be a unique cell within these 840,000 other cells. Since the *Wolbachia* phenotypes

are normally measured at the individual host levels (e.g. *Wolbachia* titres, antiviral

protection), the properties of individual or small numbers of mutant *Wolbachia* could be

diluted and unmeasurable.

We hypothesized, however, that over-proliferating *Wolbachia* cells could overtake the

population and that the resulting higher titres could be detectable. Indeed, fast proliferative

Wolbachia can be selected at the level of a single host [31]. To increase the probability of

isolating rare over-proliferating *Wolbachia* variants, we also relied on the bottleneck

imposed in the vertical transmission of *Wolbachia*. We calculated here that single embryos

carry approximately 4,000 *Wolbachia* genomes, which is consistent with previous

estimates [32]. Moreover, we treated flies with tetracycline to further reduce this

population, and increase drift. By screening at the immediate progeny (F1) of EMS-treated

females or three generations later (F4) we were able to select new over-proliferative

Wolbachia mutants.

Genetic bases of *Wolbachia* over-proliferation

After discarding the possibility that mutations in the host nuclear or mitochondrial genomes

were the cause of *Wolbachia* over-proliferation, we performed *de novo* assembly of the

ancestral, *w*MelCS_b, and the new mutant variants, *w*MelPop2 and *w*MelOctoless. The

assemblies generated complete full chromosomes of these *Wolbachia*. *w*Mel and

*w*MelPop were also assembled. This allowed us to identify single nucleotide differences

and structural differences between these genomes [33]. To validate our pipeline we

compared our *w*Mel genome assembly to the reference *w*Mel genome. We identified only

seven indels and two SNPs, which we confirmed to be present in our line, by Sanger

sequencing. Our assembly results also allowed us to confirm two previously identified

SNPs between *w*MelCS_b (but also *w*MelOctoless and *w*MelPop2) and *w*MelPop.

Additionally, our assembly provides a significant improvement over the previous *w*MelPop

genome [26].

The only differences between the new over-proliferative variants and *w*MelCS_b were

structural differences in the Octomom region. *w*MelPop2 has an amplification of this

region. The assembly confirms that the Octomom region is amplified in tandem [17], and that all copies are located in the *Wolbachia* genome. We had shown before that amplification of the Octomom region and the degree of this amplification determined *w*MelPop over-proliferative phenotype [16,17]. If the Octomom region copy number in *w*MelPop reverted to one, the variant became phenotypically identical to *w*MelCS_b [17]. We now show that if we start from *w*MelCS_b with one copy of Octomom, an increase in the copy number of this genomic region leads to an over-proliferative phenotype. Moreover, *w*MelPop and *w*MelPop2 variants, carrying the exact same copy number of Octomom, have identical phenotypes. These results further confirm the role of amplification of Octomom region in the over-proliferation of *Wolbachia*.

*w*MelOctoless has a deletion of the entire Octomom region. The deletion leaves behind one of the direct repeats flanking the Octomom region [16], suggesting that excision might be mediated by recombination. The causal link between deletion of this region and over-proliferation is further supported by an independent over-proliferative variant isolated in the screen, *w*MelOctoless2. This *Wolbachia* also has a deletion of Octomom as the only difference with *w*MelCS_b. These data show that deletion of the Octomom region leads to an over-proliferative phenotype in *Wolbachia*. Thus, we identified the second known *Wolbachia* mutation with a clear link between genotype and phenotype.

The mutations identified in the new variants are deletions and amplifications. We did not detect any new SNPs in these variants. However, we expected EMS to mainly induce single nucleotide mutations. It is possible that the deletion or amplification of the Octomom region in these over-proliferative variants were independent of the EMS treatment. For instance, loss of the Octomom region has been twice reported, in cell culture [26,34], suggesting that it can occur spontaneously. However, chemical mutagens such as EMS can activate DNA damage response and transposable elements [35], and some genes of the Octomom region and its flaking genes are predicted to be potentially involved in transposition and DNA repair [16]. Therefore, we cannot rule out that the EMS treatment induced the mutations in this genomic region.

Opposing mutations lead to a similar *Wolbachia* over-proliferative phenotype

Both the deletion or the amplification of the Octomom region causing an over-proliferative phenotype seems to be a paradox. The resolution of this paradox and the mechanisms leading to these phenotypes will rely on the functional characterization of genes in the

Octomom region. These genes may be involved in interaction with the host, transcriptional
regulation or DNA repair [16].

One possibility is that the amplification of the Octomom region and over-expression of
particular genes in this region lead, mechanistically, to the same result as the absence of
the genes. There are many examples of over-expression of a gene leading to a dominant
negative phenotype. For instance, both over-expression or loss of a protein forming a
gradient, abolish the gradient [36]. Also, over-expression of a member of protein complex
may lead to loss of stoichiometry and therefore loss of functional complexes [37]. Another
possibility is that the phenotypes are mediated by loss and amplification of different genes
in the Octomom region. The second hypothesis is supported by the fact that the two
mutant variants have similar but not identical phenotypes, and interact differently with
temperature. Furthermore, *w*MelPop and *w*MelPop2 have a higher rate of proliferation
than *w*MelOctoless at 25ºC and 29ºC. Therefore, their phenotype is stronger than the
phenotype associated with the Octomom region loss-of-function.

Although complex, these results help to explain some of the data from an over-proliferative
variant trans-infected in *Aedes aegypti*. The *D. melanogaster* variant *w*MelPop was
transinfected into *Aedes albopictus* cells, and then transinfected into *A. aegypti*. In the
process of cell culture adaptation the Octomom region, which was amplified in *w*MelPop,
was deleted [26]. The *w*MelPop-PGYP variant in *A. aegypti* lacks, therefore, the Octomom
region. Nonetheless, this variant over-proliferates and is highly pathogenic in *A. aegypti*
[38]. If one assumed the same genetic basis for the pathogenicity of *w*MelPop in *D.*
melanogaster and *w*MelPop-PGYP in *A. aegypti*, one might conclude that the Octomom
region was not related with these phenotypes [26]. However, similarly to
wMelPop/wMelPop-PGYP variants our results show that both amplification and deletion of
the Octomom region lead to increased *Wolbachia* pathogenicity. Thus, deletion of the
Octomom region in *w*MelPop-PGYP is probably why it is also pathogenic.

The Octomom region has the properties of a genomic island and it is not part of the
Wolbachia core genome, since many *Wolbachia* strains lack this region [16,39]. Although
the *w*Mel strain can lose this region and be viable in several scenarios (here and [26,34],
natural variants of *w*Mel without Octomom are not known [40]. The over-proliferative
phenotype, and the associated shortening of host lifespan, may lead to a fitness
disadvantage to a host harbouring such mutant. This in turn may lead to loss of these
variants from the host population. Therefore, this genomic region, absent in many other

strains of *Wolbachia*, and through its integration in the regulation of proliferation, became
addictive to the *w*Mel strain.

***Wolbachia* proliferation, pathogenicity, and antiviral protection**

We found a complex interaction between temperature and proliferation of the different
Wolbachia variants in adults. *w*MelOctoless proliferates faster than *w*MelCS_b, to a similar
extent, at all temperatures. *w*MelPop and *w*MelPop2, however, strongly interact with
temperature and proliferate much faster at higher temperatures. We also confirmed here
that the Octomom copy number in these variants modulates the proliferation rate [17], and
interacts with temperature.

Throughout the range of tested temperatures, the different variants have very different
proliferation rates. The titres of *w*MelPop2 at 25ºC, for instance, double every 2 days, while
wMelCS_b titres only double every 14 days. Interestingly the proliferation rates of
*w*MelCS_b, *w*MelOctoless, and *w*MelPop2 during larval development were much faster
and very similar between them. During development, at 25ºC, their doubling time varies
between 12 to 16 hours. Thus, rapid exponential proliferation of *Wolbachia* occurs
simultaneously with the fast growth of the host in the larval stages. These results show
that although *w*MelCS_b has a relatively low proliferation rate in adult flies, it is capable of
a very fast proliferation. Moreover, these data show that *Wolbachia*, somehow, coordinates
its proliferation with host developmental stage. The Octomom region influences this
coordination. While in *w*MelCS_b the proliferation rate is severely reduced in the adult
stages, in *w*MelOctoless is less reduced, and in wMelPop and wMelPop2 is still kept very
high. The over-proliferative phenotypes of these variants, in particular of *w*MelPop and
*w*MelPop2, can be interpreted as an inability of the *Wolbachia* variants to properly respond
to the host developmental stage or its own population density within the host.

The new over-proliferative variants shorten the lifespan of *D. melanogaster*, as *w*MelPop
does. Furthermore, we showed this phenotype to result from the interaction of *Wolbachia*
genotype and temperature. *w*MelOctoless had a similar life shortening phenotype at all
temperatures, although it was stronger at 18ºC. *w*MelPop and *w*MelPop2 responded
strongly to temperature, being much more costly at higher temperatures, as shown before
for *w*MelPop [17,22,41]. Octomom copy number also influenced this interaction. While low
copy number variants had no phenotype at 18ºC, the high copy number *w*MelPop and
*w*MelPop2 are pathogenic at this temperature. Therefore these variants can also be
pathogenic at 18ºC, contrary to previous data [29,41,42]. Interestingly, we find at all

temperatures a significant correlation between the proliferation rate of the *Wolbachia* variants and the life shortening phenotype. The faster the variants proliferate the shorter the host lifespan.

These over-proliferative variants also increased antiviral resistance, with *w*MelOctoless conferring the strongest protection. This phenotype correlated poorly with proliferation rates at 25ºC or 18ºC, the temperature before and after infection with DCV, respectively. However, the strength of the antiviral resistance correlated with the titres of *Wolbachia* near the day of infection. Thus, the cost of harbouring *Wolbachia* in terms of lifespan, and the benefit of the antiviral protection, correlate with related but different parameters. Therefore, it may be possible to select for highly protective *Wolbachia* variants without necessarily having a high cost to the host. These would be *Wolbachia* variants with high titres but low proliferation in adults. Such variants would be particularly useful in the use of *Wolbachia*-transinfected mosquito to prevent arboviruses transmission.

In summary, our results show the feasibility of forward genetic screens to study *Wolbachia* biology. Similar strategies may be used in the future to study other aspects of *Wolbachia*-host interactions or the biology of other genetically intractable endosymbionts. The new over-proliferative variant *w*MelPop2 confirms the causal link between amplification of the Octomom region and increase proliferation. Whereas the new loss-of-function mutant *w*MelOctoless reveals that this region is also required to control *Wolbachia* proliferation.

Materials and Methods

Fly genotypes, infection status, and maintenance

Flies were reared on fly food, supplemented with live yeast, at 25ºC, 70% humidity. Fly food was composed of molasses (37.5g/L), sugar (62.5g/L), cornflour (58.3g/L), yeast extract (16.7g/L), and agar (8.3g/L) in distilled water. The mixture was sterilized by autoclaving and cooled to 45ºC. For each litre of food we added 29.2 mL of a solution with 100g of methylparaben and 0.2g of Carbendazim for 1L absolute ethanol.

All fly stocks used had the Drosdel w^{1118} isogenic background [16,43].

The bacterial community associated with the fly stocks was homogenized as in Pais *et al.* [44], with minor modifications. Briefly, we collected eggs for 6 hours in fresh agar plates supplemented with live yeast and sterilized the eggs surface by consecutive washes on

2.1% sodium hypochlorite (NaOCl) solution (10 minutes), 70% ethanol (5 minutes) and

sterile water (5 minutes). Next, we transferred axenic eggs to sterile fly food supplemented

with 40µL of 1:1 overnight culture of *Acectobater* OTU 2753 and *Lactobacillus* OTU 1865

[44]. We confirmed the presence of these bacterial species by squashing five females

aged 3–6 days in sterile 1x PBS, plating 30µL of the lysate in mannitol plates, incubate

them at 25ºC for 72h, and identify bacteria by colony morphology.

Selection of *D. melanogaster* lines carrying *Wolbachia* with specific Octomom copy number

To select for flies carrying *w*MelPop and *w*MelPop2 with a desired Octomom copy number,

we proceeded as in Chrostek and Teixeira [17], with minor modifications. Briefly, we

allowed 5–20 virgin females to cross with 2–3 *Wolbachia*-uninfected males of the Drosdel

w^{1118} isogenic background in individual vials, and lay eggs for 3-4 days. Females were

then collected in individual tubes for DNA extraction and Octomom copy number

determination by qPCR.

Determination of time for *Wolbachia* titres recovery

Flies with *w*MelCS_b developed in fly food supplemented with tetracycline at the

concentrations 1.5625µg/ml, 3.125µg/ml, 6.25µg/ml, 12.5µg/ml, 25µg/ml, and 50µg/ml.

Three isofemale lines were established from each dose. In the F1, we randomly selected

four virgin females for egg-laying and *Wolbachia* titre measurement using qPCR. We set

up the next four generations using the progeny of a female with the median *Wolbachia*

titres.

Forward genetic screen

We attempted to mutagenize *Wolbachia in vivo* by feeding its host with the mutagen EMS.

DrosDel w^{1118} isogenic flies carrying *w*MelCS_b were first raised in standard fly food or fly

food supplemented with tetracycline (from 1.5625 to 12.5µg/ml). Virgin females were

collected, starved for 6h, and then fed on a 1% sucrose solution with EMS concentrations

ranging from 10 to 8,000mM. Control flies fed on sucrose solution only. A dye was added

to the feeding solution to confirm intake and feeding proceeded for 13h (overnight).

EMS-fed females (G0), and control females, were mated individually with 2–3 *Wolbachia*-
free Drosdel w^{1118} isogenic males, egg-laying was allowed for 3–4 days, and parents
discarded. From the F1 progeny we collected virgin females, mated them individually with
2–3 *Wolbachia*-free Drosdel w^{1118} isogenic males, and allowed egg laying for 3-4 days.
These females were collected when 10 days old, and *Wolbachia* titres determined by
qPCR. We followed the progeny of F1 females showing 50% or higher increase in
Wolbachia titres relative to control flies in the same conditions. We have also transferred
the progeny of these F1 for three more generations, without selection, and repeated the
determination of *Wolbachia* titres in F4 females. In the same batch of experiments we may
have tested more than one F1 or F4 progeny from each G0 female. Hence, over-
proliferative *Wolbachia* variants isolated in the same batch of treated females may be a
result of a single event in the G0 female.

Real time quantitative PCR

DNA extraction for qPCR was performed as described before [17].

The qPCR reactions were performed in the QuantStudioTM 7 Flex (Applied Biosystems).
The reaction mix and PCR program used were described before [16]. The specificity of the
amplicons were confirmed by analysing the melting curve profile.

Relative levels of the target genes was determined using the Pfaffl method [45]. To
quantify relative *Wolbachia* titres were performed using *Drosophila RpL32* gene as
calibrator, and *Wolbachia wsp* as the target gene. To determine the copy number of the
Octomom region, *Wolbachia wsp* gene was used as the calibrator and *WD0513* used as
the target gene. For determination of copy number of other Octomom region genes, or
control genes, *wsp* was also used as a calibrator.

For absolute quantification of *Wolbachia* genome copies the full-length *Wolbachia wsp*
gene was cloned into a pMT/V5 *Drosophila* expression vector (Invitrogen). The plasmid
was amplified in *Escherichia coli* strain DH5-α, purified using midiprep (QIAGEN) and its
concentration determined using Qubit® 2.0 (Thermo Fisher Scientific). Standard curves of
1:10 serial dilutions were run to calibrate the assay each time.

Determination of lifespans

For each replicate a total of 50 males or 50 females were collected on the day of eclosion. Flies (10 per vials) were then incubated at 18ºC, 25ºC or 29ºC, being transferred to new fresh vials every four (females) or five (males) days. The number of dead flies was recorded daily until all the flies had died. Censored observations (i.e. flies lost or trapped in the vial plug) were recorded and taken into account during data analysis.

Protection against *Drosophila* C Virus

Viral protection assays were blinded (i.e. virus and buffer pricking occurred without previous knowledge of their *Wolbachia* status). We produced and titrated the virus solution as described in Teixeira *et al.* [8]. We infected 50 3-5 days old males by dipping insect needles (Austerlitz Insect Pins) into a virus solution (DCV at 10^9 $TCID_{50}$/ml in 50mM Tris-HCl, pH 7.5) and pricking flies anaesthetized under CO2 in the thorax. An equal number of males were pricked with a buffer solution (50mM Tris-HCl, pH 7.5) and served as controls. After pricking, flies were incubated in groups of 10 individuals per vial, and kept at 18ºC. Survival was followed as above.

Wolbachia proliferation during development and in adults

To determine *Wolbachia* growth during development flies carrying *w*MelCS_b, *w*MelOctoless, and wMelPop2 laid eggs for 2 hours in apple juice agar plates supplemented with live yeast. Eggs were transferred to fly food-containing bottles and allowed to develop at 25ºC. For *Wolbachia* titre assessment we sampled eggs (2 hours), L1 larvae (24 hours post-egg-laying), newly moulted L2 larvae (48 hours), L3 larvae (72 and 84 hours), white prepupae pupae (120 hours), P8 staged pupae (168 hours), and newly eclosed adult males and females (240 hours). Ten samples per time point were analysed. Samples included ten individuals each for eggs and larvae and one individual each for pupae and adults. Except for adults who were collected within 24 hours post-emergence, all samples were collected within two hours interval.

For assessment of titres dynamics in adult flies, newly eclosed males, raised at 25ºC, were incubated at 18ºC, 25ºC, and 29ºC. Flies were collected every three (29ºC), seven (25ºC) or ten days (18ºC) for *Wolbachia* titre measurement. Ten individuals were processed for each time point, and the experiment was performed twice. Each sample consisted of a single fly.

Wolbachia genomes sequencing and quality control

For *Wolbachia* genomic sequencing (Illumina and Oxford Nanopore) we enriched the
sample for *Wolbachia* cells before DNA purification. To this end, approximately 500 10-
days old flies were squashed for 5 minutes in 10ml Schneider's Insect Medium (Thermo
Fisher Scientific) using glass beads. Next, we pelleted host debris by centrifugation at
1,000g for 5 min and filtered the supernatant solution thought a 5μm pore. Bacterial cells
were pelleted by centrifugation at 13,000rpm for 15 minutes, and DNA was extracted. All
centrifugations were carried out at 4ºC. DNA was extracted with a phenol-chloroform
isolation protocol and resuspended in 10mM Tris-HCl (pH 8).

Sequencing was performed at the Genomics Facility at the Instituto Gulbenkian de
Ciência, (Portugal). Both Illumina and Oxford Nanopore sequencing was done on genomic
DNA extracted from the same biological material. Illumina 300bp paired-end libraries were
prepared using the Pico Nextera kit according to the manufacturer's instructions and
sequenced by the MiSeq sequencing. Data quality was assessed via FastQC v.0.11.5 [46]
and reads were trimmed using Trimmomatic v.0.36 [47]. Genomic samples for Oxford
Nanopore sequencing were processed with minimal shearing to maximize the size of the
libraries. After ligation (kit SQK-LSK108), libraries were sequenced in MinION Mk1b
portable sequencing device using SpotON flow cell (R9.4.1). The status of the sequencing
pores was monitored using MinKNOW (v2.0.1). Sequencing lasted for up to 48 hours.
Albacore (v2.3.1) and Porechop (v0.2.2) were used for base-calling and read trimming,
respectively.

Genome assembly

Quality reads matching *Drosophila* genome (BioProject: PRJNA13812) were removed
from the datasets through a custom Python script before proceeding with *Wolbachia*
genome assembly. Illumina and Oxford Nanopore reads were mapped to the fly genome
using BWA mem v0.7.12-r1039 [48] and minimap2 v2.17-r941 [49] respectively. We used
Unicycler v0.4.8-beta [50] assembly pipeline on the remaining reads in order to assemble
the *Wolbachia* genomes. Briefly, Unicycler uses Illumina data to produce a repeats-limited
image graph using Spades v3.9.0 [51], which was further refined through Bandage v0.8.1
[52]. Both small short nodes and nodes with no homology with *w*Mel genome
(AE017196.1) upon blastn v2.8.1+ [53] search were removed. Next, repeats were resolved

by bridging Spades' assemblies with Oxford Nanopore long reads. The resulting draft
assemblies were polished using Racon v1.3.1 [54] and Pilon v1.23 [55] and rotated so that
genomes begin at the *dna*A gene.

We further refined our genome assemblies using a two-stepped pipeline. First, we mapped
Illumina reads to the corresponded draft genomes to identify mismatches, which were later
corrected via Sanger sequencing. Next, we improved genome assemblies by aligning
them using Mauve v2.4.0 [56], and the predicted differences were also confirmed by
Sanger sequencing. Only two predicted SNPs which are unique to *w*MelPop were real.

Read mapping and mutation calling

We identified mutations relative to a genome following a previously published pipeline [57].
It consisted of mapping quality checked reads to a reference genome using BWA mem
v0.7.12-r1039 algorithm [48] and saving the output as Sequence Alignment/Map file format
(SAM). After conversion to the Binary Alignment/Map format (BAM), the file was sorted,
duplicates removed and indexed using SAMtools v0.1.19 [58]. Next, we generate mpileup
files, also using SAMtools (option '-d 1,000,000'), after which was converted to Variant Call
Format (VCF) files using BCFtools v1.9-209. We visually confirmed all inferred mutations
in IGV v2.4.2 [59]. We did not consider mutations associated with homopolymer regions or
in regions with low coverage (<10X). The set of mutations were compared between
Wolbachia variants using custom Python and R scripts.

To compare the set of mutations in the mitochondria of flies infected with different
Wolbachia variant, we mapped Illumina reads to the *D. melanogaster* Release 6 genome
sequence (KJ947872.2:1–14,000) and proceed as previously. Mutations following the
criteria previously described were also compared by using custom scripts.

Statistical analysis

All the statistical comparisons were performed in R v4.0.0 [60].

To compare *Wolbachia* titres across multiple groups, we used log linear models (LMM) or
generalized linear mixed models assuming Gamma errors (GLMM). The effect of EMS on
Wolbachia titre was tested using non-linear regression. We estimated the doubling time of
Wolbachia variants using the equation $\log(2)/\beta$, with β being the coefficients of a log linear
model.

The lifespan datasets and survival curves after challenge with DCV were analysed with mixed effect Cox models [61].

The significance of correlations were tested using Pearson's correlation coefficient.

In multiple comparisons were necessary, the p-values were adjusted as proposed by Holm [62]. When multivariate techniques were applied, all the relevant covariates were included in the model, and the final model was selected as proposed by Burnham & Anderson [63].

Acknowledgments

We are thankful to João Costa and the Bioinformatics Unit and the Genomics Facility at the Instituto Gulbenkian de Ciência for help with sequencing and assembling the *Wolbachia* genomes. We are thankful to the Fly Facility at IGC for support to the *Drosophila* work.

Funding

This work was funded by the FCT grant IF/00839/2015, and the ERC grant 773260.

E.D. was funded with the fellowship SFRH/BD/113757/2015 from Fundação para a Ciência e Tecnologia, in the context of the Graduate Program Science for the Development.

The fly work at the Fly Facility of Instituto Gulbenkian de Ciencia (Oeiras, Portugal), was partially supported by the research infrastructure Congento, co-financed by Lisboa Regional Operational Programme (Lisboa2020), under the PORTUGAL 2020 Partnership Agreement, through the European Regional Development Fund (ERDF) and Fundação para a Ciência e Tecnologia (Portugal) under the project LISBOA-01-0145-FEDER-022170.

The NGS analysis at the Genomics Unit of Instituto Gulbenkian de Ciencia (Oeiras, Portugal), was partially supported by ONEIDA project (LISBOA-01-0145-FEDER-016417) co-funded by FEEI - "Fundos Europeus Estruturais e de Investimento" from "Programa Operacional Regional Lisboa 2020" and by national funds from FCT - "Fundação para a Ciência e a Tecnologia."

References

1. Moran NA, McCutcheon JP, Nakabachi A. Genomics and Evolution of Heritable Bacterial Symbionts. Annual review of genetics. 2008;42: 165–190. doi:10.1146/annurev.genet.41.110306.130119

2. Jaenike J. Coupled population dynamics of endosymbionts within and between hosts. Oikos. 2009;118: 353–362.

3. López-Madrigal S, Duarte EH. Titer regulation in arthropod-Wolbachia symbioses. FEMS microbiology letters. 2019;366: 118–9. doi:10.1093/femsle/fnz232

4. Zug R, Hammerstein P. Still a host of hosts for wolbachia: analysis of recent data suggests that 40% of terrestrial arthropod species are infected. PLoS ONE. 2012;7: e38544. doi:10.1371/journal.pone.0038544

5. Werren JH, Baldo L, Clark ME. Wolbachia: master manipulators of invertebrate biology. Nature reviews Microbiology. 2008;6: 741–751. doi:10.1038/nrmicro1969

6. Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. Wolbachia as a bacteriocyte-associated nutritional mutualist. Proceedings of the National Academy of Sciences of the United States of America. 2010;107: 769–774. doi:10.1073/pnas.0911476107

7. Hedges LM, Brownlie JC, O'Neill SL, Johnson KN. Wolbachia and virus protection in insects - Supp. Science (New York, NY). 2008;322: 702. doi:10.1126/science.1162418

8. Teixeira L, Ferreira A, Ashburner M. The bacterial symbiont Wolbachia induces resistance to RNA viral infections in Drosophila melanogaster. PLoS biology. 2008;6: e1000002. doi:10.1371/journal.pbio.1000002

9. Ross PA, Turelli M, Hoffmann AA. Evolutionary Ecology of Wolbachia Releases for Disease Control. Annual review of genetics. 2019;53: 93–116. doi:10.1146/annurev-genet-112618-043609

10. Aliota MT, Walker EC, Yepes AU, Velez ID, Christensen BM, Osorio JE. The w Mel Strain of Wolbachia Reduces Transmission of Chikungunya Virus in Aedes aegypti. PLoS Neglected Tropical Diseases. 2016;10: e0004677. doi:10.1371/journal.pntd.0004677

11. Dutra HLC, Rocha MN, Dias FBS, Mansur SB, Caragata EP, Moreira LA. Wolbachia Blocks Currently Circulating Zika Virus Isolates in Brazilian Aedes aegypti Mosquitoes. Cell host & microbe. 2016. doi:10.1016/j.chom.2016.04.021

12. Moreira LA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AT, Hedges LM, et al. A Wolbachia symbiont in Aedes aegypti limits infection with dengue, Chikungunya, and Plasmodium. Cell. 2009;139: 1268–1278. doi:10.1016/j.cell.2009.11.042

13. van den Hurk AF, Hall-Mendelin S, Pyke AT, Frentiu FD, McElroy K, Day A, et al. Impact of Wolbachia on Infection with Chikungunya and Yellow Fever Viruses in the Mosquito Vector Aedes aegypti. PLoS Neglected Tropical Diseases. 2012;6: e1892. doi:10.1371/journal.pntd.0001892

14. Nazni WA, Hoffmann AA, NoorAfizah A, Cheong YL, Mancini MV, Golding N, et al.
Establishment of Wolbachia Strain wAlbB in Malaysian Populations of Aedes aegypti
for Dengue Control. CURBIO. 2019;0. doi:10.1016/j.cub.2019.11.007

15. O'Neill SL, Ryan PA, Turley AP, Wilson G, Retzki K, Iturbe-Ormaetxe I, et al. Scaled
deployment of *Wolbachia* to protect the community from *Aedes* transmitted
arboviruses. Gates Open Research. 2018;2: 36. doi:10.12688/gatesopenres.12844.1

16. Chrostek E, Marialva MSP, Esteves SS, Weinert LA, Martinez J, Jiggins FM, et al.
Wolbachia Variants Induce Differential Protection to Viruses in Drosophila
melanogaster: A Phenotypic and Phylogenomic Analysis. PLoS genetics. 2013;9:
e1003896. doi:10.1371/journal.pgen.1003896

17. Chrostek E, Teixeira L. Mutualism breakdown by amplification of Wolbachia genes.
PLOS Biol. 2015;13: e1002065. doi:10.1371/journal.pbio.1002065

18. Lu P, Bian G, Pan X, Xi Z. Wolbachia Induces Density-Dependent Inhibition to
Dengue Virus in Mosquito Cells. PLoS Neglected Tropical Diseases. 2012;6: e1754.
doi:10.1371/journal.pntd.0001754

19. Osborne SE, Leong YS, O'Neill SL, Johnson KN. Variation in antiviral protection
mediated by different Wolbachia strains in Drosophila simulans. PLoS pathogens.
2009;5: e1000656. doi:10.1371/journal.ppat.1000656

20. Osborne SE, Iturbe-Ormaetxe I, Brownlie JC, O'Neill SL, Johnson KN. Antiviral
protection and the importance of Wolbachia density and tissue tropism in Drosophila
simulans. Applied and environmental microbiology. 2012;78: 6922–6929.
doi:10.1128/AEM.01727-12

21. McMeniman CJ, Lane RV, Cass BN, Fong AWC, Sidhu M, Wang Y-F, et al. Stable
Introduction of a Life-Shortening Wolbachia Infection into the Mosquito Aedes aegypti
- Supp Material. Science (New York, NY). 2009;323: 141–144.
doi:10.1126/science.1165326

22. Min KT, Benzer S. Wolbachia, normally a symbiont of Drosophila, can be virulent,
causing degeneration and early death. Proceedings of the National Academy of
Sciences of the United States of America. 1997;94: 10792–10796.

23. Ant TH, Sinkins SP. A Wolbachia triple-strain infection generates self-incompatibility in
Aedes albopictus and transmission instability in Aedes aegypti. Parasites & Vectors.
2018;11: 1–7.

24. McClure EE, Chávez ASO, Shaw DK, Carlyon JA, Ganta RR, Noh SM, et al.
Engineering of obligate intracellular bacteria: progress, challenges and paradigms.
Nature reviews Microbiology. 2017;984: 249. doi:10.1038/nrmicro.2017.59

25. Ashburner M, Golic KG, Hawley RS. Drosophila : a laboratory handbook. Cold Spring
Harbor, N.Y. : Cold Spring Harbor Laboratory Press; 2005. Available:
http://lccn.loc.gov/2004019474

26. Woolfit M, Iturbe-Ormaetxe I, Brownlie JC, Walker T, Riegler M, Seleznev A, et al.
Genomic evolution of the pathogenic Wolbachia strain, wMelPop. Genome biology
and evolution. 2013;5: 2189–2204. doi:10.1093/gbe/evt169

27. Wu M, Sun LV, Vamathevan J, Riegler M, Deboy R, Brownlie JC, et al.
Phylogenomics of the reproductive parasite Wolbachia pipientis wMel: a streamlined
genome overrun by mobile genetic elements. PLoS biology. 2004;2: E69.
doi:10.1371/journal.pbio.0020069

28. Basting PJ, Bergman CM. Complete Genome Assemblies for Three Variants of the
Wolbachia Endosymbiont of Drosophila melanogaster. Microbiol Res Announc.
2019;8: 741.

29. Chrostek E, Teixeira L. Comment on Rohrscheib et al. 2016 "Intensity of mutualism
breakdown is determined by temperature not amplification of Wolbachia genes." PLoS
pathogens. 2017;13: e1006540. doi:10.1371/journal.ppat.1006540

30. McGarry HF, Egerton GL, Taylor MJ. Population dynamics of Wolbachia bacterial
endosymbionts in Brugia malayi. Molecular and Biochemical Parasitology. 2004;135:
57–67. doi:10.1016/j.molbiopara.2004.01.006

31. Chrostek E, Teixeira L. Within host selection for faster replicating bacterial symbionts.
PLoS ONE. 2018;13: e0191530. doi:10.1371/journal.pone.0191530

32. Christensen S, Camacho M, Sharmin Z, Momtaz AJMZ, Perez L, Navarro G, et al.
Quantitative methods for assessing local and bodywide contributions to Wolbachia
titer in maternal germline cells of Drosophila. BMC Microbiol. 2019;19: 1–17.
doi:10.1186/s12866-019-1579-3

33. Alkan C, Coe BP, Eichler EE. Genome structural variation discovery and genotyping.
Nature reviews Genetics. 2011;12: 363–376.

34. Grobler Y, Yun CY, Kahler DJ, Bergman CM, Lee H, Oliver B, et al. Whole genome
screen reveals a novel relationship between Wolbachia levels and Drosophila host
translation. PLoS pathogens. 2018;14: e1007445. doi:10.1371/journal.ppat.1007445

35. Burr B, Burr FA. Activation of Silent Transposable Elements. Plant Transposable
Elements. Boston, MA: Springer US; 1988. pp. 317–323.

36. Duchek P, Rørth P. Guidance of Cell Migration by EGF Receptor Signaling During
Drosophila Oogenesis. Science (New York, NY). 2001;291: 131–133.
doi:10.1126/science.291.5501.131

37. Prelich G. Gene overexpression: uses, mechanisms, and interpretation. Genetics.
2012;190: 841–854. doi:10.1534/genetics.111.136911

38. McMeniman CJ, Lane RV, Cass BN, Fong AWC, Sidhu M, Wang Y-F, et al. Stable
Introduction of a Life-Shortening Wolbachia Infection into the Mosquito Aedes aegypti.
Science (New York, NY). 2009;323: 141–144. doi:10.1126/science.1165326

39. Iturbe-Ormaetxe I, Burke GR, Riegler M, O'Neill SL. Distribution, expression, and
motif variability of ankyrin domain genes in Wolbachia pipientis. Journal of
bacteriology. 2005;187: 5136–5145. doi:10.1128/JB.187.15.5136-5145.2005

40. Richardson MF, Weinert LA, Welch JJ, Linheiro RS, Magwire MM, Jiggins FM, et al.
Population Genomics of the Wolbachia Endosymbiont in Drosophila melanogaster.
PLoS genetics. 2012;8: e1003129. doi:10.1371/journal.pgen.1003129

41. Reynolds KT, Thomson LJ, Hoffmann AA. The effects of host age, host nuclear
background and temperature on phenotypic effects of the virulent Wolbachia strain
popcorn in Drosophila melanogaster. Genetics. 2003;164: 1027–1034.

42. Rohrscheib CE, Frentiu FD, Horn E, Ritchie FK, van Swinderen B, Weible MW, et al.
Intensity of Mutualism Breakdown Is Determined by Temperature Not Amplification of
Wolbachia Genes. PLoS pathogens. 2016;12: e1005888.
doi:10.1371/journal.ppat.1005888

43. Ryder E, Blows F, Ashburner M, Bautista-Llacer R, Coulson D, Drummond J, et al.
The DrosDel collection: a set of P-element insertions for generating custom
chromosomal aberrations in Drosophila melanogaster. Genetics. 2004;167: 797–813.
doi:10.1534/genetics.104.026658

44. Pais IS, Valente RS, Sporniak M, Teixeira L. Drosophila melanogaster establishes a
species-specific mutualistic interaction with stable gut-colonizing bacteria. PLOS Biol.
2018;16: e2005710. doi:10.1371/journal.pbio.2005710

45. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR.
Nucleic acids research. 2001;29: e45.

46. Andrews, Simon. FastQC: A quality control tool for high throughput sequence data.
2010. Available: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

47. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics (Oxford, England). 2014;30: 2114–2120.
doi:10.1093/bioinformatics/btu170

48. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics (Oxford, England). 2009;25: 1754–1760.
doi:10.1093/bioinformatics/btp324

49. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics (Oxford,
England). 2018;34: 3094–3100.

50. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome
assemblies from short and long sequencing reads. bioRxiv. 2016; 096412.
doi:10.1101/096412

51. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes:
A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing.
Journal of Computational Biology. 2012;19: 455–477. doi:10.1089/cmb.2012.0021

52. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de
novogenome assemblies: Bioinformatics (Oxford, England). 2015;31: 3350–3352.

53. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
tool. Journal of molecular biology. 1990;215: 403–410.

54. Vaser R, Sovic I, Nagarajan N, Sikic M. Fast and accurate de novo genome assembly
from long uncorrected reads. Genome research. 2017;27: 737–746.

55. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: An
Integrated Tool for Comprehensive Microbial Variant Detection and Genome
Assembly Improvement. PLoS ONE. 2014;9: e112963.

56. Darling ACE, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of conserved
genomic sequence with rearrangements. Genome research. 2004;14: 1394–1403.

57. Li H. A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data. Bioinformatics
(Oxford, England). 2011;27: 2987–2993.

58. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics (Oxford, England). 2009;25:
2078–2079. doi:10.1093/bioinformatics/btp352

59. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
Integrative genomics viewer. Nature biotechnology. 2011;29: 24–26.

60. Team RC. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria; 2012. Available: URL http://www.R-project.org/

61. Therneau TM. coxme: Mixed Effects Cox Models. 2020. Available: https://CRAN.R-
project.org/package=coxme

62. Holm S. A simple sequentially rejective multiple test procedure. Scandinavian Journal
of Statistics. 1979;6: 65–70.

63. Burnham KP, Anderson DR. Model selection and multimodel inference : a practical
information-theoretic approach. New York : Springer; 2002.

1053

1054

Supporting information

S1 Fig. *Wolbachia* recovers from severe titre reduction within 3-4 fly generations.

Relative *Wolbachia* titres of the progeny of tetracycline-treated flies. *w*MelCS_b-carrying females laid eggs in food containing varying doses of tetracycline. The progeny of three females were used to set up the experiment. At the first generation, four females were randomly selected for egg-laying in antibiotic-free fly food and *Wolbachia* titre was measured using qPCR. Titres of untreated females were used to normalize the qPCR results. The progeny of a female with the median titre was used to set up the next generation. *Wolbachia* titre in the F1 was significantly determined by the concentration of the antibiotic ($p < 0.001$ for all doses compared with control at generation 1), but recovered to normal within four fly generations ($p > 0.05$ for all doses compared with control at generation 4).

S2 Fig. EMS decreases female fecundity and *Wolbachia* titre in the next generation in a dose-dependent manner.

The total number of eggs (A) and adults (B) of females treated with varying doses of EMS. The reproductive
output of 10 females was determined in the first ten days after EMS treatment by daily transferring females to
new vials for egg laying. Females fed on a sucrose solution served as controls. Each dot represents the total
number of eggs (A) or adults (B) laid by individual females during ten days. The effect of EMS on the
reproductive output of females was estimated using a non-linear model and was highly significant ($p < 0.001$
for both numbers of eggs and adults per female). (C-D) *Wolbachia* titres in the F1 progeny of flies treated
with varying EMS doses. *Wolbachia* titre was quantified on individual females (n=5–13), after laying eggs for
three days. *Wolbachia* titres were normalized against the titres of untreated females. Dashed red lines
represent the mean value predicted using non-linear models. The effect of EMS on *Wolbachia* titres in the
next generation was highly significant ($p < 0.001$ for both panels).

1078

1079 **S3 Fig. Isolation of over-proliferative *Wolbachia* variants.**

1080 (A-D) Relative *Wolbachia* titres in a control (*w*MelCS_b) and EMS-treated *Drosophila* lines. Lines to set up
1081 the next generation was selected as described for Fig 1. Line 2B was isolated in the same batch as for Line
1082 2A (wMelOctoless) and they may be not independent. Likewise, Lines 3A (*w*MelOctoless2), 3B, and 3C were
1083 also isolated in the same batch.

1084

S4 Fig. Generation of isogenic *Drosophila* lines with *w*MelPop2 and *w*MelOctoless.

The first, second and third chromosomes of flies carrying *w*MelPop2 and *w*MelOctoless were replaced
through the use of balancer chromosomes. *Wolbachia* infection (and also mitochondria) was kept in the
stock by crossing females with *Wolbachia* with selected males. The mitochondria are only shown in females
because of its strictly maternal transmission. All males were free of *Wolbachia* infection. Dashed lines
indicate the genotype selected from the previous cross. Virgin female in the first cross were considered
mutant in all chromosomes (*), for illustrative purposes. Question marks (?) represent recombined
chromosomes.

1093

S5 Fig. Identification the genetic bases of *Wolbachia* over-proliferating variants.

The relative coverage in a selected genomic region. As detailed for Fig 2B, Illumina paired-end reads were
mapped to *w*MelCS_b (GenBank: CP046924.1) genome, and the number of reads mapping to each position
were normalized by dividing to the median coverage across the genome. Coverage information for
*w*MelCS_b, *w*MelPop2 and *w*MelOctoless is also given in Fig 2B. We identified the deletion of Octomom as
the cause of over-proliferation in lines 2B and line 3A (*w*MelOctoless2), as no other difference was found
when compared to *w*MelCS_b.

1101

1102 **S6 Fig. Confirmation of deletion of Octomom genes in *w*MelOctoless by qPCR.**

1103 The deletion of individual Octomom genes was confirmed using qPCR. The copy number of three genes
1104 outside the Octomom region were also determined. Five females carrying *w*MelCS_b or *w*MelOctoless were
1105 used in the analysis. The copy number of *w*MelOctoless genes is relative to that of *w*MelCS_b.

1106

S7 Fig. Selection for stocks with a desired Octomom copy number.

The relative copy number of genomic WD0513 in *Wolbachia*-carrying stocks throughout 30 fly generations. Each generation, 5-20 females were randomly collected for egg-laying for 3-4 days and used to determine the relative copy number of WD0513 as a proxy for the Octomom copy number. The progeny of a single female was used to set up the next generation. qPCR results were normalized to that of *w*MelCS_b, which has a single copy of Octomom per genome.

1113

S8 Fig. Proliferation dynamics of over-proliferative *Wolbachia* variants at different temperatures.

Wolbachia proliferation dynamics at 18ºC (A), 25ºC (B) and 29ºC (C). *Wolbachia* titres were determined and analysed as described for Fig 3.

1117

S9 Fig. Octomom copy number determines *Wolbachia* titres on the day of eclosion.

Relative *Wolbachia* titres on the day of adults eclosion. Males developed at 25ºC were collected within 24 hours after eclosion for *Wolbachia* titre measurement using qPCR. Data used in this figure were given in Fig 3 and S8 Fig (time point 0). Letters represent significant groups after *p*-value correction.

1122

S10 Fig. *w*MelPop2 and *w*MelPop are phenotypically indistinguishable.

(A) WD0513 copy number of *w*MelPop2 and *w*MelPop in two experimental replicates. Using WD0513 as a proxy, the Octomom copy number of *w*MelPop2 and *w*MelPop was tightly controlled prior to phenotypic comparison. (B) *Wolbachia* relative titres at 25ºC. The progeny of *w*MelPop2- and *w*MelPop-infected females carrying three copies of Octomom was used to set up the experiments. Males that developed at 25ºC were collected upon eclosion, aged to specific time-points and used to determine *Wolbachia* titers using qPCR. *Wolbachia* titers were normalized to that of *w*MelCS_b-carrying flies collected on the day of eclosion. Proliferation rates of *w*MelPop2 and *w*MelPop were not different (p = 0.32). (C) Lifespan of males (solid lines) and females (dashed lines) flies at 25ºC. The experiment was performed twice, as described for Fig 3. Males were transferred to new vials every five days, while females every four days. (D). Coefficients of a Cox mixed model, representing the effect of *w*MelPop2 and *w*MelPop on the lifespan relative to *w*MelCS_b-carrying flies. *w*MelPop2 and *w*MelPop was equally pathogenic (p = 0.29).

1135

S11 Fig. Octomom copy number dynamics throughout *Drosophila* development and during the adult life.

(A) Relative copies of WD0513 throughout the fly development. Samples were collected and processed as described in Fig 4. Relative copies of WD0513 was determined using qPCR and are normalized to *w*MelCS_b. There is no significant change of Octomom copy number in wMelPop2 during fly development (p = 0.88). (B) The relative copies of WD0513 during the adult life at 18ºC, 25ºC, and 29ºC. Relative copies

1142 of WD0513 was determined and normalized as previously described. There was a small increase of
1143 Octomom copy number with adult age (p = 0.009).

1144

S12 Fig. New over-proliferative Wolbachia variants are pathogenic.

Lifespan of males with different *Wolbachia* variants at 18ºC (A), 25ºC (B), and 29ºC (C). An experimental
replicate of these experiments is given in Fig 5A-C, respectively. (D) Lifespan of females at 25ºC in two
experimental replicates. Experiments consisted of collecting 50 females upon adult eclosion and following
their survival at 25ºC until all flies had died. Females were kept on groups of 10 per vial and transferred to
new vials every four days. (E) Coefficients of a Cox mixed model, representing the effect of *Wolbachia*
variants on the survival of *Drosophila* females. Bars represent the standard error of the estimates, and letters
statistically significant groups.

1153

S13 Fig. *Wolbachia* variants and not the differences in the host genetic background are the cause of
the life shortening phenotype.

(A-B) Lifespan of females at 29ºC. Virgin females carrying *w*MelCS_b were crossed with males with
*w*MelOctoless or *w*MelPop2. Reciprocal crosses were also performed. The survival of resulting female
progeny, which have the same genetic background but differ in *Wolbachia* infection, were assayed at 29ºC.
For both experimental replicates, we determined the lifespan of 50 female progeny from each cross.
Females were maintained in groups of 10 per vial and transferred to new vial every four days. (C)
Coefficients of a Cox mixed model, representing the effect of *Wolbachia* variant on the survival of females.
Significance of each group was accessed after *p*-value correction for multiple comparisons, and are
represented by letters.

1164

S14 Fig. Correlation between the strength of life-shortening phenotype or anti-viral protection with
***Wolbachia* titre and proliferation.**

(A-C) Correlation between the strength of life-shortening phenotype at 18ºC (A), 25ºC (B), and 29ºC (C) and
Wolbachia titres at the day of eclosion. (D-E) Correlation between the strength of *Wolbachia* anti-viral
protection and *Wolbachia* doubling times at 18ºC (D) and 25ºC (E). The direction and significant of all
correlations was estimated using Person correlation coefficient and are given in each panel.

1171

1172 **S15 Fig. Over-proliferative *Wolbachia* variants confer strong protection against DCV.**

1173 Survival of males males after a DCV (A) or buffer (B) challenge. An experimental replicate is given in Fig 6.
1174 The experiment was set up and analysed as described in Fig 6.

1175

1176 **S1 Table. Number of progeny of EMS-treated females screened for over-proliferating *Wolbachia***
1177 **variants.**

1178 Progenies of EMS-treated females were collected as virgin, mated with 2-3 *Wolbachia*-free Drosdel *w*[1118]
1179 isogenic males, allowed to egg-laying for 3–4 days and used in qPCR reations to determine *Wolbachia* titres
1180 relative to the progeny of control females.

1181

1182 **S2 Table. Coverage statistics of the sequencing project.**

1183 Coverage statistics (median, the first and third quantiles) of Illumina reads across different genomes mapped
1184 to either *Wolbachia* or mitochondria of *D. melanogaster* Release 6 genome sequence (KJ947872.2:1–
1185 14,000) using BWA mem v0.7.12-r1039. *Wolbachia* reference genome used for mapping the different reads
1186 are indicated. ND – not determined.

1187

S3 Table. Flies with *w*MelCS_b, *w*MelOctoless and *w*MelPop2 carry an identical mitochondrial
genome.

Illumina reads of fly stocks varying on *Wolbachia* variants were mapped to the *D. melanogaster* Release 6
genome sequence (KJ947872.2:1–14,000) using BWA mem v0.7.12-r1039. The mitogenome of flies infected
with *w*MelCS_b, *w*MelOctoless and *w*MelPop2 was identical. We found a SNP unique to flies infected with
*w*MelCS-like variants (G→A on position 10,793) but absent in flies infected with *w*MelPop or *w*Mel. We
confirmed the substitution using Sanger sequencing (S2 Text).

1195

1196 **S4 Table. Assembly and annotation statistics.**

1197 *Wolbachia* genomes were assembled using the Unicycler v0.4.8-beta pipeline and annotated using NCBI
1198 Prokaryotic Genome Annotation Pipeline v4.10. We found variant in *Wolbachia* genome size but the resulting
1199 genomes encoded for the same number of transfer, ribosomal and non-coding RNAs.

1200

1201 **S5 Table. SNPs and indels differences between different *w*Mel genomes.**

1202 The genome of a newly assembled Cluster III *Wolbachia* variant, called *w*Mel, was compared to *w*Mel
1203 reference genome. Nine differences were identified and validated using Sanger sequencing.

1204

1205 **S1 Text. Alignment of a genomic region, including Octomom, in different Wolbachia variants.**

1206 An alignment summary of a genomic region, including Octomom, in *w*MelCS, *w*MelPop2 and *w*MelOctoless.

1207

1208 **S2 Text. Confirmation of a SNP in the mitochondrial genome.**

1209 Sanger sequencing of a PCR amplicon containing a SNP present in *w*MelCS_b, *w*MelPop2 and
1210 *w*MelOctoless, but absent in *w*MelPop and *w*Mel.

1211

1212

Relative *Wolbachia* titer

☿ $\dfrac{X_{iso}*}{X_{iso}*}$; $\dfrac{2_{iso}*}{2_{iso}*}$; $\dfrac{3_{iso}*}{3_{iso}*}$ X ♂ $\dfrac{FM7}{Y}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{3_{iso}}{3_{iso}}$

☿ $\dfrac{X_{iso}*}{FM7}$; $\dfrac{2_{iso}*}{2_{iso}}$; $\dfrac{3_{iso}*}{3_{iso}}$ X ♂ $\dfrac{X_{iso}}{Y}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{3_{iso}}{3_{iso}}$

☿ $\dfrac{X_{iso}}{FM7}$; $\dfrac{?}{2_{iso}}$; $\dfrac{?}{3_{iso}}$ X ♂ $\dfrac{X_{iso}}{Y}$; $\dfrac{Gla}{Cyo}$; $\dfrac{3_{iso}}{3_{iso}}$

☿ $\dfrac{X_{iso}}{X_{iso}}$; $\dfrac{?}{Cyo}$; $\dfrac{?}{3_{iso}}$ X ♂ $\dfrac{X_{iso}}{Y}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{3_{iso}}{3_{iso}}$

☿ $\dfrac{X_{iso}}{X_{iso}}$; $\dfrac{2_{iso}}{Cyo}$; $\dfrac{?}{3_{iso}}$ X ♂ $\dfrac{X_{iso}}{Y}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{TM2}{TM6}$

☿ $\dfrac{X_{iso}}{X_{iso}}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{?}{TM6}$ X ♂ $\dfrac{X_{iso}}{Y}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{3_{iso}}{3_{iso}}$

☿ $\dfrac{X_{iso}}{X_{iso}}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{3_{iso}}{TM6}$ X ♂ $\dfrac{X_{iso}}{Y}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{3_{iso}}{3_{iso}}$

☿ $\dfrac{X_{iso}}{X_{iso}}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{3_{iso}}{3_{iso}}$ X ♂ $\dfrac{X_{iso}}{Y}$; $\dfrac{2_{iso}}{2_{iso}}$; $\dfrac{3_{iso}}{3_{iso}}$

■ *Wolbachia* ● Mitochondria

Relative coverage

Copy number of individual genes

Relative copies of *WD0513*

A

Relative *Wolbachia* titer at 18°C

B

Relative *Wolbachia* titer at 25°C

C

Relative *Wolbachia* titer at 29°C

Relative *Wolbachia* titer at the day of eclosion

A Percentage of males alive at 18°C

B Percentage of males alive at 25°C

C Percentage of males alive at 29°C

Wolbachia-free
*w*MelCS_b (1 copy)
*w*MelOctoless (0 copies)
*w*MelPop2 (3 copies)
*w*MelPop2 (8-9 copies)
*w*MelPop (2-3 copies)
*w*MelPop (8-9 copies)

D Percentage of females alive at 25°C
Replicate 1

Percentage of females alive at 25°C
Replicate 2

Days post eclosion

E Coefficients of Cox mixed model

Wolbachia-free	a
*w*MelCS_b (1 copy)	a
*w*MelOctoless (0 copies)	b
*w*MelPop2 (3 copies)	d
*w*MelPop2 (8-9 copies)	f
*w*MelPop (2-3 copies)	c
*w*MelPop (8-9 copies)	e

Coefficients ± SE

A

Percentage of females alive at 29°C (Replicate 1)

B

Percentage of females alive at 29°C (Replicate 2)

Days post eclosion

C

Coefficients of Cox mixed model

A

Percentage of flies alive after DCV infection

B

Percentage of flies alive after buffer pricking

Days post infection

- *Wolbachia*-free
- *w*MelCS_b (1 copy)
- *w*MelOctoless (0 copies)
- *w*MelPop2 (3 copies)
- *w*MelPop2 (8-9 copies)
- *w*MelPop (2-3 copies)
- *w*MelPop (8-9 copies)