Individual differences in honey bee (*Apis mellifera*) behavior enabled by plasticity in brain gene regulatory networks

Beryl M. Jones[1,*], Vikyath D. Rao[2,3], Tim Gernat[2,4], Tobias Jagla[4], Amy Cash-Ahmed[2], Benjamin E. R. Rubin[5], Troy J. Comi[5], Shounak Bhogale[6], Syed S. Husain[7], Charles Blatti[2], Martin Middendorf[4], Saurabh Sinha[2,6], Sriram Chandrasekaran[7,8,‡], Gene E. Robinson[2,9,10,‡]

[1]Program in Ecology, Evolution, and Conservation Biology, University of Illinois at Urbana–Champaign, Urbana, IL, USA
[2]Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana–Champaign, Urbana, IL, USA
[3]Department of Physics, University of Illinois at Urbana–Champaign, Urbana, IL, USA
[4]Swarm Intelligence and Complex Systems Group, Department of Computer Science, Leipzig University, Leipzig, Germany
[5]Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ, USA
[6]Center for Biophysics and Quantitative Biology, University of Illinois at Urbana–Champaign, Urbana, IL, USA
[7]Department of Biomedical Engineering, University of Michigan, Ann Arbor, MI, USA
[8]Center for Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, USA
[9]Neuroscience Program, University of Illinois at Urbana–Champaign, Urbana, IL, USA
[10]Department of Entomology, University of Illinois at Urbana–Champaign, Urbana, IL, USA

*Current address: Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA
‡Joint corresponding authors, csriram@umich.edu and generobi@illinois.edu

Abstract

Understanding the regulatory architecture of phenotypic variation is a fundamental goal in biology, but connections between gene regulatory network (GRN) activity and individual differences in behavior are poorly understood. We characterized the molecular basis of behavioral plasticity in queenless honey bee colonies, where individuals engage in both reproductive and non-reproductive behaviors. Using high-throughput behavioral tracking, we discovered these colonies contain a continuum of phenotypes, with some individuals specialized for either egg-laying or foraging and "generalists" that perform both. Brain gene expression and chromatin accessibility profiles were correlated with individual behavioral variation, with generalists intermediate in behavior and molecular profiles. Models of brain GRNs constructed for individuals revealed that transcription factor (TF) activity was highly predictive of behavior,

and behavior-associated regulatory regions had more TF motifs. These results provide new

insights into the important role played by brain GRN plasticity in the regulation of behavior, with

implications for social evolution.

Introduction

Understanding the genomic regulatory architecture of phenotypic plasticity is necessary

to achieve comprehensive knowledge of the mechanisms and evolution of complex traits. While

a growing body of knowledge exists on specific regulatory mechanisms involved in

developmental plasticity, less is known about the regulation of behavioral plasticity. Behavioral

plasticity is of special interest and presents unique challenges, as behavioral traits derive from

the integrated actions of genetic, transcriptomic, and neuronal networks (Sinha et al., 2020).

Over the past 20 years, a close relationship between behavioral variation and brain gene

expression has been documented across a range of organisms and behaviors (e.g., Zayed and

Robinson, 2012). Still, the regulatory architecture underlying connections between the genome,

brain, environment, and behavior are not well resolved, in part because behavior is itself a

complex phenotype with substantial variation between individuals. To fully understand how

genomic and transcriptomic variation is transduced into behavioral plasticity, we need both high-

dimensional behavioral data at the individual level as well as information on regulatory

genomics for those same individuals.

Modification of gene regulatory networks (GRNs) has emerged as an important driver of

plasticity during the development and evolution of morphological phenotypes. For example,

gains and losses of *cis*-regulatory elements (e.g., binding sites for transcription factors (TFs))

influence species-specific wing melanization patterns in *Heliconius* butterflies and *Drosophila*

flies (Prud'homme et al., 2006; Reed et al., 2011; Werner et al., 2010). Pelvic loss in stickleback

fish convergently evolved through deletion of a tissue-specific enhancer of the TF *Pitx1* in

multiple natural populations (Chan et al., 2010). In other cases, similar morphological novelties

arose independently through modification of distinct developmental programs, as observed for

beak size variation across clades of finches (Mallarino et al., 2012). Recruitment of genes

involved in developmental plasticity in the evolution of novel phenotypes is thought to be

facilitated by the fact that TFs and other regulatory genes often have great temporal flexibility,

with extensive variation in expression across developmental time (Dufour et al., 2020).

Similar to its role in morphological variation, plasticity in GRNs is theorized to influence behavioral variation, over both organismal and evolutionary time scales (Sinha et al., 2020). Brain gene expression is often responsive to environmental stimuli (Chandrasekaran et al., 2011; Cummings et al., 2008; Mukherjee et al., 2018; Rittschof et al., 2014; Whitfield et al., 2003) and the regulatory activity of many TFs is context-specific with respect to behavioral state (Chandrasekaran et al., 2011; Hamilton et al., 2019). These results demonstrate that GRNs are plastic not only across developmental timescales but also influence real-time behavioral variation. Still, the link between changes in GRNs and behavioral plasticity is weaker than for developmental plasticity (Sinha et al., 2020), and to our knowledge, no empirical studies have linked brain GRN plasticity to individual differences in behavior.

Eusocial insects are ideal for studying how GRN activity influences both developmental and behavioral plasticity at the individual scale. Eusociality is characterized by a reproductive division of labor between queen and worker castes, representing a developmentally plastic polyphenism well-studied in many species (e.g., Holldobler and Wilson, 1990; Michener, 1974; O'Donnell, 1998; Wheeler, 1986). Queens are specialized for reproductive functions, including mating and egg-laying, and in species with complex eusociality, have levels of fecundity orders of magnitude greater than their solitary ancestors. Workers, on the other hand, typically do not perform reproductive behaviors and in many cases are sterile or unable to mate, instead performing many different non-reproductive behaviors in a colony that are essential for colony growth and development. Species with complex eusociality also often show additional within-caste behavioral plasticity, with individuals specializing on specific subsets of tasks based on differences in worker age, morphology, or genetic predisposition. The extensive behavioral plasticity observed in colonies of eusocial species may be linked to ancestral developmental plasticity (Kapheim et al., 2020), highlighting the interconnectedness of gene regulation in both developmental and behavioral phenotypes relevant for social behavior (Sinha et al., 2020).

We studied the relationship between brain GRN activity and behavioral plasticity at the individual scale. We focused on a recently discovered, surprising form of behavioral plasticity among worker honey bees. Honey bee workers do not mate, but they possess functional ovaries and can produce viable haploid eggs. Laying workers (LW) are rare in queenright colonies (Ratnieks, 1993; Visscher, 1996) but frequent in situations of permanent queenlessness, when colonies lose their queen and then fail to rear a replacement queen. In these cases, up to 50% of

workers may activate their ovaries (Sakagami, 1954) and some of these workers lay eggs,

producing thousands of drones prior to colony death (Page Jr and Erickson Jr, 1988). Recently, it

was discovered that some LWs engage in both reproductive and non-reproductive behaviors

(Naeger et al., 2013), a level of behavioral plasticity not previously described in honey bee

workers. Studying honey bee workers in LW colonies enables investigation of the molecular

architecture of behavioral variation typically only seen when comparing queens and workers,

without the confounds of caste-specific developmental and physiological differences.

Recent advances in machine learning and automatic behavioral tracking have enabled the

study of individual behavior for thousands of members within social insect colonies (Crall et al.,

2015; Gernat et al., 2018; Greenwald et al., 2015; Mersch et al., 2013; Wario et al., 2015). We

used automatic behavioral tracking, genomics, and the extensive behavioral plasticity present in

honey bee colonies with LW to test the hypothesis that individual differences in behavior are

associated with changes in the activity of brain GRNs, i.e., changes in the expression of TFs and

their target genes. Our results provide key insights into the mechanisms underlying the regulation

of individual differences in behavior by brain GRNs.

Results

Extensive variation in behavior across laying workers

To define the behavior of individual bees, we deployed a high-resolution, automatic

behavior monitoring system on six LW colonies in which each bee (n=800 per colony) was

individually barcoded, similar to Gernat et al. (2018). Our extension of this system identifies the

location and heading direction of each individual once per second, and uses convolutional neural

networks and machine learning to detect behaviors. For each individual across seven days of

tracking (when bees were 15-21 days old), egg-laying events and foraging trips were detected

from images of the hive interior and entrance (Fig. 1A). A total of 115,281 egg-laying events and

96,086 foraging trips were predicted for the six colonies (File S1).

Colonies exhibited considerable variation in the proportion of bees engaged in egg-laying

and/or foraging. With the exception of colony F, more workers were identified as layers than

foragers (Fig. 1B). Across all colonies, a high proportion of bees were observed laying eggs

(54% with at least two egg-laying events on at least one day) or foraging (28% with at least two

foraging trips on at least one day) during the recording period, while 10.8% of bees performed

both egg-laying and foraging on the same day at least once during the seven days of tracking. A small number of these "generalist" bees (1.3%; 45 individuals) were exceptional in their consistent high performance of both measured behaviors, with a minimum of two egg-laying events and two foraging trips on the same day, across at least three days. Three-day ethograms of an egg-layer, generalist, and forager are shown in Fig. 1C. Ovary dissection of a subset of individuals revealed that 100% of specialized egg-layers and generalists had active ovaries (ovary scores of 3-5; Hess 1942), compared with only 54% of the specialized foragers (Fig. 1D, File S2). Of the foragers with activated ovaries, 13/14 had 5 or fewer predicted egg-laying events, compared with generalists and layers, which laid an average of 206 eggs (range: 64-774).

The daily and lifetime behavior of each bee was summarized using two behavioral scores: the "specialist" score, which describes how specialized an individual was on either egg-laying or foraging, and the "generalist" score, which describes how much an individual performed both egg-laying and foraging. Scores were derived from daily normalized ranks within colonies to allow comparison across days and colonies with differing overall activity levels; bees that performed neither egg-laying nor foraging across the experiment have both specialist and generalist scores of 0. Scores were mapped onto a two-dimensional color space for visualization of behavior over time (Fig. 2, File S1).

Figure 1. (A) Automatic behavior monitoring was performed inside the hive and at the hive entrance to predict egg-laying and foraging events in six colonies (N=800 bees per colony at the start of each trial). Hive images were captured 1/s for 24 h/day, and entrance images 2/s for 12 h/day beginning when adult bees were 15 days old. **(B)** Proportion of bees alive each day categorized as layers (purple), foragers (green), generalists (orange), or others (gray). For colonies A-C, individuals were from single source colonies headed by a naturally mated queen. For colonies D-F, individuals from two source colonies headed by queens each inseminated by semen from a single different drone (single drone inseminated, SDI) were mixed. Different source colonies are indicated by pattern and hue. **(C)** Ethograms for three individuals selected for sequencing (bCodes shown below group labels) across three days of tracking. **(D)** Distribution of ovary scores for individuals selected for sequencing. Insets are images from bees with ovary scores of 1, 3, and 5. L: layer, G: generalist, F: forager.

Influence of worker source colony on behavior

To study the influence of source colony (including genetics and development) on behavior, experimental colonies were assembled with workers from different source colonies headed by unrelated queens. Bees in colonies A-C were derived from colonies with naturally mated queens. Queens of *Apis mellifera* mate multiply with up to ~20 males and produce workers with a mix of paternal genotypes (Adams et al., 1977; Estoup et al., 1994; Lobo and Kerr, 1993); workers derived from these colonies were therefore assumed to be a mix of many patrilines. In contrast, experimental colonies D-F were assembled of workers obtained from two different source colonies, each of which was headed by a queen artificially inseminated with the semen of a single different drone (SDI). Workers within each SDI source colony are highly genetically related compared with workers from a naturally mated queen colony (average relatedness = 0.75 due to haplodiploidy).

Using SDI colonies allowed us to more easily explore whether the genetic and environmental differences between source colonies would lead to segregation of reproductive and non-reproductive behavior when mixed into the same (queenless) environment. In both colonies D and E, which were replicates of the same two SDI queens' offspring, the behavior of workers differed considerably by source: one source colony (SDI 1) comprised the majority of foragers, while the other (SDI 2) contained the majority of egg-layers (Fig. 2). In colony F the two SDI source colony progeny contributed more equally to foraging, while the most specialized egg-laying bees were predominantly from just one source colony (Fig. 2, SDIs 3 and 4). However, even in colonies where SDI source was clearly influential, specialized foragers and layers were identified from both sources, indicating that colony genetics and development are not the only contributors to individual variation in the likelihood of performing these behaviors. Similar patterns of specialization were observed in colonies A-C and D-F (Fig. 2), indicating that they were not an artifact of decreased intracolonial genetic diversity.

Figure 2. Daily behaviors of individual bees (rows) across time in each colony. Colored rectangles indicate specialist and generalist scores represented in 2D color space as shown in legend. Individuals are sorted by median lifetime specialist score. Single-drone inseminated (SDI) queen source is shown to the right of each row for colonies D-F, where workers were known offspring of two SDI queens per colony. In colonies A-C workers were offspring of naturally mated queens.

Specialized behavioral groups are highly transcriptionally and epigenetically distinct

A subset of highly specialized foragers, egg-layers, and generalist individuals were

selected from two experimental colonies (from only one source SDI each) for brain gene

expression and chromatin accessibility profiling (Fig. 3A). Principal component analysis (PCA)

on behavioral data for these individuals shows these three groups are behaviorally distinct, with

generalists intermediate and more variable than forager and layer groups (Fig. 3B).

Consistent with strong behavioral differentiation, foragers and layers exhibited

widespread differences in brain gene expression, with differential expression of nearly half

(46%) of all genes expressed in the brain (Fig. 3C, File S3). Differences in brain gene expression

were much stronger between foragers and layers (4506 differentially expressed genes, DEGs)

than for generalists relative to the two specialist groups (648 generalist vs. layer and 374

generalist vs. forager DEGs). Generalists shared transcriptional profiles of both foragers and

layers, with nearly all genes differentially expressed between generalists and either specialized

group also present on the forager vs. layer DEG list (Fig. 3C).

Forager vs. layer DEGs were enriched for cytoplasmic translation and transport gene

ontology (GO) biological processes, along with many metabolic and biosynthetic processes (File

S3). All enriched GO terms but one (114 of 115) were for genes more highly expressed in

foragers relative to layers (forager-biased genes). The only GO term enriched in layer-biased

genes relative to foragers, cytoplasmic translation, was also the only enriched GO term for genes

overexpressed in generalists relative to foragers. Similarly, GO terms enriched in generalist-

biased genes (relative to layers) included many of the transport terms enriched among forager-

biased genes (File S3).

In addition to differences in brain gene expression, layers and foragers showed

differences in accessible chromatin in the brain based on the Assay for Transposase-Accessible

Chromatin using sequencing (ATAC-seq; Buenrostro et al., 2013). 1794 differentially accessible

peaks (DAPs) were identified between foragers and layers, proximal to 1207 genes (Fig. 3D, File

S4). Forager-biased DEGs and genes proximal to forager-biased DAPs overlapped significantly,

1.2x more than expected by chance (p=0.01 for hypergeometric test of overlap). Genes proximal

to peaks of accessible chromatin (regardless of differential status) were on average more highly

expressed than genes without proximal peaks (p<0.0001, Kolmogorov-Smirnov test), supporting

a signal of transcriptional activation near ATAC-seq peaks (Fig. S1). No DAPs were identified

Figure 3. (A) Daily rank-normalized behavior of individuals (rows) selected for brain RNAseq and ATACseq analysis converted to 2D colorspace from specialist and generalist scores. **(B)** Principal Component Analysis (PCA) of behavioral variation for individuals chosen for brain RNAseq and ATACseq analysis. Metrics included number of eggs laid, number of foraging events, proportion of foraging trips with evidence of nectar collection, proportion of trips with evidence of pollen collection, and proportion of trips with evidence of both nectar and pollen collection. **(C)** Euler diagram for overlaps of pairwise differentially expressed genes (DEGs) between behavioral groups. Note that one gene was overlapping between F vs. G and G vs. L (but not F vs. L) and is not represented in the diagram due to graphical constraints. **(D)** Euler diagram for overlaps of genes proximal to pairwise differentially accessible chromatin peaks (DAPs) between behavioral groups. **(E)** PCs from PCA of brain transcriptomic profiles regressed against specialist score (PC1: R^2=0.947, p<0.0001; PC2: R^2=0.838, p<0.001). **(F)** PCs from PCA of brain chromatin accessibility regressed against specialist score (PC2: R^2=0.584, p<0.001; PC3: R^2=0.543, p<0.0001; PC4: R^2=0.187, p<0.0045; PC1: p>0.05). L: layer, G: generalist, F: forager.

for foragers relative to generalists, and there were only 16 DAPs (assigned to 13 genes) for

generalists relative to layers (Fig. 3D). These 13 genes also had DAPs for foragers relative to

layers (Fig. 3D). DAPs between foragers and layers were enriched for 148 GO terms, including

developmental processes, morphogenesis, and metabolism (File S4). Similar to differentially

expressed genes, GO enrichment signal came from those DAPs with a bias in foragers (i.e., more

accessible in foragers relative to layers); no significantly enriched GO terms were identified from

layer-biased peaks, despite 44% of differential peaks being more accessible in layers compared

to foragers.

Brain gene expression and chromatin accessibility are correlated with behavioral variation

Our high-resolution behavioral data allowed us to test whether molecular and behavioral

variation were connected not only at the group level, but for individuals as well. Using PCA, we

found that degree of individual behavioral specialization was significantly correlated with

measures of both brain gene expression and chromatin accessibility (Fig. 3E-F). Among PCs for

gene expression, PCs 1 and 2, which explained 31.1% and 11.9% of the total variance in gene

expression, respectively, were significantly correlated with individual behavioral specialist score

(Fig. 3E). Generalists showed intermediate values of these PCs, consistent with an intermediate

brain transcriptomic profile. Genes with extreme PC loading values (upper and lower 5% of

loadings) for PC1 were enriched for transmembrane and ion transport, functions related to

aerobic and cellular respiration, and energy transport (File S5). PC2 extreme loading genes were

enriched for processes relating to detection of light, phototransduction, and sensory perception

(File S5). Extreme loadings for both PC1 and PC2 overlapped significantly with DEGs in the

pairwise comparison of layers and foragers (PC1: RF=1.2, p=1.39e-06, PC2: RF=1.7, p=8.39e-

91).

Similarly, PCA of chromatin accessibility data revealed PCs that were correlated with

behavioral variation. Accessibility PCs 2, 3, and 4 were all significantly correlated with the

individual behavioral specialist score (Fig. 3F). Genes with extreme PC loading values for each

correlated PC showed enrichment for multiple GO terms, including biological processes related

to cell-cell adhesion, locomotion, axon guidance, neuron projection guidance, and synapse

organization (File S6). Many GO terms (11 of 37) were enriched for extreme loading genes of

both PCs 2 and 3, while synapse organization was the only enriched term for loadings of PC4.

GRN activity links molecular and behavioral phenotypes

To test the role of transcription factor (TF) and gene regulatory plasticity in the regulation

of LW behavioral phenotypes, we conducted TF motif analyses and brain GRN reconstruction,

individualized for each bee.

TF motifs were strongly enriched in peaks of differential accessibility or in the regulatory

regions of DEGs between specialized layers and foragers. 77 out of 223 motifs (functionally

validated in *Drosophila melanogaster*, Zhu et al., 2011) were enriched in layer vs. forager DAPs

(q-val<0.01, File S7), and 14 motifs were specifically enriched in the regulatory regions of

forager-upregulated DEGs (q-val<0.2, File S7). Nine motifs were common to both sets (Fig. 4),

including binding sites for TFs involved in regulating nervous system development (hairy, side,

sr, and klu), transcription (max/mnt), juvenile hormone (JH) signaling (tai and tai/met),

chromatin modification (trl), and circadian rhythms (cwo). These motifs were centrally enriched

within DAPs (Fig. 4A), and showed two peaks of elevated binding probability in the promoter

regions of forager-biased DEGs, one ~3kb upstream of transcriptional start sites (TSSs) and a

second overlapping TSSs (Fig. 4B). By contrast, only two TF motifs were significantly enriched

in the regulatory regions of specialist vs. generalist DEGs (mad and ken, File S7), and no motifs

were enriched within DAPs between generalists and either specialist group (File S7), likely due

in part to the small number of DAPs distinguishing generalists and specialists (Fig. 3D). Many of

the motifs enriched within DAPs or DEG promoters are binding sites for TFs that were

themselves differentially expressed, including cwo, tai/met, side, h, and sr (File S3).

Figure 4. Shared motifs enriched within DAPs and DEG promoters for specialist foragers vs. layers. **(A)** Motifs enriched within DAPs show maximum binding probabilities near peak summits. **(B)** Motifs enriched in promoter regions of forager>layer DEGs show elevated binding probabilities ~3kb upstream of and overlapping TSSs. Motif names and sequences are from FlyFactor (Zhu et al. 2011) for *Drosophila melanogaster*.

The relationship between TF activity and behavior was so strong that it was possible to

predict individual behavior based solely upon the expression of TFs in the brain using a

machine-learning algorithm and leave-one-out cross validation (Fig. 5A). TF expression

correctly predicted 100% of foragers and 94% of layers. By contrast, it was not possible to

predict generalists based on brain TF expression (only 1 of 8 correctly classified).

Brain GRN reconstruction (Chandrasekaran et al., 2011), performed on an individualized

basis, further highlights the strong influence of gene regulatory plasticity on behavior. GRN

activity was largely consistent between individuals within each behavioral group (Fig. 5B).

Further, at the individual level, the activity of many TF modules (TFs and their predicted targets)

showed significant correlations with individual variation in several behavioral metrics, including

numbers of eggs laid (50 TF modules), specialist behavioral score (77 TF modules), and number

of foraging trips (74 TF modules) (File S8). At the individual level, 23 TF modules were

correlated with all 9 behavioral and physiological metrics (Fig. S3, File S8). These behaviorally

correlated TF modules include TFs involved in JH signaling (usp, Kr-h1, and Blimp-1), histone

acetylation (trx), neuronal remodeling (Kr-h1, Hr51, trx), and circadian rhythms (opa and Hr51).

Figure 5. (A) Class prediction analysis based on brain TF expression correctly classified all but one specialist (L: layer, F: forager) but only one generalist (G). Normalized expression (logCPM) of 4 of the top 20 informative TFs for class prediction analysis are shown (others in Fig. S2). Median of points is represented by bold horizontal line within shaded 95% confidence interval, with length of shape and smoothed curve showing range and density of data, respectively. **(B)** TF modules (rows) with significant up/downregulation in at least 10 individuals, sorted by hierarchical clustering. Individuals (columns) are ordered by specialist score, with darkly colored blocks indicating correctly classified individuals based on TF expression prediction analysis and lightly colored blocks indicating incorrect classification. TF modules showed patterns of differentiation between L and F, while G were more variable in module activity. Labeled modules are those with TFs shown in panel (A) or discussed in text.

Comparative analyses of LW colony behavioral phenotypes and other social insect phenotypes

The performance of both egg laying and foraging by individuals in LW colonies,

previously reported in Naeger et al. (2013), is unusual for honey bees; these behaviors are

otherwise confined to separate castes (queens and workers). This raises the question of whether

the mechanisms underlying LW behavior reflect caste-related molecular differences. We

compared our gene expression results to previous studies of queens, workers, and worker

subcastes in various species of social insects to ask whether the molecular architecture of LW

phenotypes may be useful in the context of understanding additional social phenotypes.

In comparison with honey bee subcastes, forager-biased genes in LW colonies showed

significant overlap with forager-biased genes in two studies of queenright colonies (when

compared with nurses) (RF=1.7 p=1.707e-09; Alaux et al., 2009; RF=1.9 p=1.740e-07; Whitfield

et al., 2003; File S9). Layer-biased genes in this study overlapped with genes upregulated in

nurses relative to foragers in these queenright colonies (RF=1.7, p=3.656e-10; Alaux et al., 2009;

RF=2.0, p=1.116e-13; Whitfield et al., 2003; File S9).

In addition, differences in brain gene expression between egg-layers and foragers

mirrored caste-related differences across species. Genes differentially expressed between

foragers and egg-layers in this study were enriched for previously identified queen vs. worker

brain DEGs in *Megalopta genalis* bees, which facultatively engage in both reproductive and non-

reproductive behaviors (RF:1.3, p=0.009; Jones et al., 2017; File S9). Overlap was in the

expected direction, with queen-biased genes in *M. genalis* overlapping layer-biased genes

(RF:2.5, p=0.003) and worker-biased genes overlapping forager-biased genes (RF:1.6, p=0.01).

Further, worker-upregulated DEGs in the primitively eusocial wasp, *Polistes metricus*

overlapped significantly with forager-upregulated genes in this study (RF:2.6, p<0.0001; Toth et

al., 2010). In comparison with honey bee unmated queens and workers, overlap was significant

but in an unexpected direction: queen- and worker-biased genes overlapped with forager- and

layer- upregulated genes, respectively (RF:1.4, p=5.495e-08 and RF:1.2, p=0.008; Grozinger et

al., 2007; File S9).

Additionally, forager vs. layer DEGs in this study were enriched for genes identified as

under selection in two studies of social evolution. Forager vs. layer DEGs overlapped

significantly with genes undergoing positive selection in honey bees (RF=1.1, p=0.015; Harpur

et al., 2014; File S9) and across highly eusocial species relative to solitary or primitively eusocial

species (Woodard et al., 2011; File S9). Genes under selection in highly eusocial lineages were

enriched specifically for genes identified here as upregulated in foragers relative to layers

(RF=1.4, p=0.009), but not for layer-biased DEGs (p=0.106) (File S9). Forager vs. layer DEGs

were not significantly enriched for genes that were identified in a third study as under selection

in social lineages of bees (p=0.262; Kapheim et al., 2015; File S9). Many of the forager vs. layer

DEGs also found to be undergoing positive selection were related to metabolism (Fig. S4).

TFs involved in LW plasticity previously implicated in social evolution

Given that differences in brain gene expression between egg-layers and foragers reflect

caste-related differences, we also tested whether there is overlap between TFs involved in LW

plasticity and those previously implicated in social evolution. Indeed, many of the TFs we

identified above as related to behavioral plasticity based on motif enrichment, group predictive

analysis, or brain GRN activity were previously known to be associated with social behavior on

an evolutionary timescale. A comparative analysis of the genomes of ten bee species (Kapheim

et al., 2015) identified 13 TF motifs with associations between binding strength and social

complexity. Nine of those 13 motifs were also detected above as enriched within specialist DAPs

or DEG regulatory regions (p=0.015, hypergeometric test of overlap), and 6 of those 9 are

binding sites for TFs included in the above individualized GRNs. Along with TF module

correlation and behaviorally-predictive TF expression, these results highlight a set of 15 TFs as

compelling candidates in social plasticity and evolution, with significant associations in at least 3

of the 5 analyses (Fig. 6). The 15 TFs have functions related to known mechanisms associated

with social behavior, including brain development (Hamilton et al., 2016), JH signaling

(Woodard et al., 2011), and chromatin changes via histone acetylation (Simola et al., 2015).

Figure 6. Candidate TFs regulating specialist behavioral phenotypes based on evidence across all analyses (descriptions of categories in Methods). Names given are for *Drosophila melanogaster* motifs (Zhu et al. 2011), with homology to honey bee genes as in Kapheim et al. (2015). Color of bar in first two columns indicates whether there was stronger enrichment among forager-biased (green) or layer-biased (purple) peaks or genes.

Discussion

Uncovering the regulatory mechanisms involved in behavioral plasticity is important to

fully understand how behavioral phenotypes develop and evolve. We used automatic behavioral

tracking and genomics to uncover the role of brain GRN activity in the extensive behavioral

variation observed in colonies of laying worker honey bees. We discovered that continuous

phenotypic variation is associated with continuous variation in both brain gene expression and

brain chromatin accessibility, and that TF activity is predictive of behavioral phenotype at the

individual level. These results provide new mechanistic insights into the important role played by

brain GRNs in the regulation of behavioral plasticity, with implications for understanding the

mechanisms and evolution of complex traits.

Our high-dimensional behavioral data revealed a near continuous distribution of

phenotypes along an axis of egg-laying and foraging, two behaviors that are typically expressed

separately in the queen and worker castes of honey bee colonies. Consistent with previous

reports of ovary activation in queenless colonies (Page Jr and Erickson Jr, 1988; Sakagami,

1954), over half of workers tracked laid eggs. Some of these workers also engaged in foraging,

consistent with the observations of Naeger et al. (2013), which supports the suggestion that some

laying workers are not "selfish" reproducers but engage in activities that may benefit the colony

as a whole. We also showed a striking decoupling between ovary status and behavior, unlike in many other social insect species (Barth et al., 1975; Michener, 1974; Wilson, 1971). Two-thirds (14/21) of the foragers had activated ovaries, but most were infrequently predicted to lay eggs, demonstrating that ovary activation alone is not a strong predictor of exactly which individuals will lay eggs. This decoupling of reproductive physiology from reproductive behavior is consistent with the evolutionary co-option of reproductive signaling pathways for non-reproductive behaviors, a phenomenon well documented in honey bees (Amdam et al., 2008; Graham et al., 2011; Page et al., 2012).

Like the task specialization observed in typical, queenright colonies of honey bees and many other social insects (Oster and Wilson, 1979), the majority of individuals in LW colonies showed consistency in performance of either egg-laying or foraging, but not both. Task specialization can contribute to increased efficiency in social insects, either through learning or reduction of task switching costs (Holldobler and Wilson, 1990; Jeanson et al., 2008; Trumbo and Robinson, 1997; c.f. Dornhaus, 2008). In queenless colonies of honey bees, specialization along a reproductive/non-reproductive axis may lead to increased production of haploid males prior to the death of workers, with specialized foragers collecting food for these developing drones while specialized egg-layers work to produce thousands of drones synchronously in these terminal colonies (Page Jr and Erickson Jr, 1988). These findings suggest that LW honey bees may display a form of colony organization that is adaptive, as opposed to one of chaos and competition, which has long been thought to characterize LW colonies (Morse, 1990; Ratnieks et al., 2006; Ratnieks and Wenseleers, 2008; Dadant & Sons, 1975; Wenseleers and Ratnieks, 2006). Worker derived drones have viable sperm (Gençer and Kahya, 2011) and therefore may provide a permanently queenless honey bee colony with a final fitness opportunity if the males can successfully mate with queens. It is difficult to evaluate this hypothesis because the incidence of permanently queenless colonies is not known in natural populations of honey bees. However, production of drones by workers in LW colonies is similar to that observed in bumble bees, where worker competition over male production is a normal part of the colony cycle after queen death (Cnaani et al., 2002; Free, 1955), or even prior to queen death in some species (Velthuis and Duchateau, 2011).

Consistent with many other studies of behavior and brain gene expression across animal species (e.g., Bukhari et al., 2019; Mello et al., 1992; Whitfield et al., 2003), we identified robust

brain transcriptional signatures associated with specific behavioral states. Beyond these group

level differences, we also discovered that large components of this molecular variation were

correlated with behavior at the individual level, and both behavior and brain gene regulatory

activity were continuous across bees. Our finding that both brain gene expression and chromatin

accessibility vary continuously with behavioral phenotype suggests that behavioral plasticity is

subserved by continuously varying molecular programs, rather than threshold-based or quantized

changes.

At the individual bee level, changes in the expression of TFs, accessibility of TF motifs

in enhancers and promoters, and activity of TF module target genes were all strongly associated

with behavioral state. This is highlighted by the results of our predictive analysis, where 97% of

specialists were accurately predicted to phenotype based on TF expression alone, despite the

small number of TFs relative to all differentially expressed genes. Spatial and temporal

integration of discrete events such as TF binding, aggregated at the whole brain level and across

TFs and genes, may lead to the continuous variation we observed in gene expression and

chromatin accessibility (e.g., Araya et al., 2014).

In addition to predicting the collective behavioral phenotypes of individual bees, our

analysis of GRNs allowed us to probe the influence of TF module activity on single behaviors.

We identified a set of 23 TF modules that were associated with all aspects of behavior and

physiology we measured. These TFs appear to coordinate sets of behaviors that are not overtly

linked (e.g., proportion of nectar foraging trips and number of eggs laid) but may be influenced

by the same regulatory machinery. Three of these modules are activated by TFs downstream of

JH, a hormone with numerous well-studied roles in social insect behavior, including the

regulation of oogenesis in queens and age-related division of labor in workers (Amdam et al.,

2008; Hamilton et al., 2017; Page et al., 2012). Our results are consistent with a role of JH

signaling in queenless colonies of worker honey bees, regulating a behavioral division of labor

between specialized egg-layers and foragers while also likely influencing ovary development and

oogenesis in the subset of individuals that lay eggs. These findings match previous work

describing differences in JH titers between egg-laying workers and foragers in queenless

colonies (Robinson et al., 1992), and suggest that mechanisms underlying variation in egg-laying

behavior may be similar to nurse/forager differences in queenright colonies. Overlap in brain

gene expression profiles between nurses and egg-layers further supports this conclusion.

While behavioral specialization appears to be common among members of queenless
honey bee colonies, the finding of even a small number of generalist bees who perform both egg-
laying and foraging has intriguing implications. The presence of these generalists suggests that
despite the long divergence from a solitary ancestor (~85 my, Branstetter et al., 2017), honey
bees retain great flexibility for performance of multiple behaviors that are typically confined to
either the queen or worker caste. Latent plasticity in social insects that is inducible under extreme
conditions is also seen in morphologically and temporally defined worker subcastes (Robinson,
1992; Simola et al., 2015; Wilson, 1980). Generalists showed high variation in behavior, and
similarly were difficult to predict phenotypically based on TF activity, unlike specialists. Further,
brain GRN activity in these individuals was less defined, with fewer TF modules showing
significant up- or down-regulation in generalist individuals compared with specialists. Combined
with PCA on brain gene expression and chromatin accessibility, these findings suggest that
generalists are molecularly intermediate between specialized groups.

Our discovery of intermediate generalist phenotypes in laying worker colonies provides
molecular support for one of the leading theories of eusocial evolution, the Ovarian Ground Plan
Hypothesis (OGPH). The OGPH posits that the emergence of queen and worker castes from
solitary ancestors involved the genetic decoupling of reproductive and non-reproductive
behavioral programs through changes in gene regulation acting on ancestral plasticity (Gadagkar,
1997; Turillazzi and West-Eberhard, 1996; West-Eberhard, 1987). The phenotypic continuum
we observed in laying worker colonies, with both reproductive and non-reproductive specialists
as well as generalists, suggests that this decoupling process is at least partially reversible and/or
incomplete in honey bees, unlike in eusocial species where workers lack reproductive anatomy
and corresponding behaviors entirely (e.g., ants and higher termites). Additionally, molecular
characterization of this behavioral variation, especially our TF analyses, supports the hypothesis
that incremental changes in gene regulatory network activity led to the decoupling of solitary
behavioral programs into distinct queen and worker castes. This hypothesis is consistent with
previous research linking changes in TF activity with social evolution (Kapheim et al., 2020,
2015). If correct, this hypothesis provides a framework for understanding the evolution of
eusociality at the molecular level.

Methods

Bees and colony setup

<u>Source colonies</u>

Honey bee colonies were maintained according to standard beekeeping practices at the University of Illinois Bee Research Facility in Urbana, Illinois. One-day-old adult worker bees were obtained by removing sealed honeycomb frames of late-stage pupae from source colonies in the field and housing them in an incubator inside emergence cages at 34 °C and 50% relative humidity. Bees were removed from frames daily to collect adults less than 24 hours old. Prior to establishing the colonies of barcoded bees, 16 source colonies were screened for worker egg-laying ("laying worker", LW) potential by stocking Plexiglas cages with 50-100 one-day-old workers and holding them in queenless, broodless conditions. Cages contained small pieces of 3D-printed honeycomb (similar to Fine et al., 2018) to provide a standardized location for workers to lay eggs, as well as 50% sucrose solution and pollen paste (45:45:10 ratio by weight of pollen, honey, and water) provided ad libitum and refreshed daily. Cages were monitored daily to count eggs. We found, as in other studies, variation in the timing and extent of LW development among different source colonies (Fig. S5), reflecting the effect of genotypic and/or environmental differences on laying worker potential (Miller III and Ratnieks, 2001; Page and Robinson, 1994; Robinson et al., 1990; Velthuis, 1970). When possible, source colonies were chosen from among those screened that displayed high levels of worker egg-laying in cages within 14 days.

To reduce genetic variation among bees used for sequencing, experimental colonies D-F were established from a mix of two source colonies each headed by a queen of either *Apis mellifera ligustica* or *Apis mellifera carnica* origin who had been artificially inseminated with semen from a single drone (SDI) (queen rearing and inseminations performed by Sue Cobey, Honey Bee Insemination Service; Washington State University; US stocks of bees are primarily, but not completely *ligustica* or *carnica*). Experimental colonies A-C were established from naturally mated, *Apis mellifera ligustica* source colonies. Honeycomb frames of late-stage pupae were removed from source colonies and maintained in an indoor incubator. Worker bees were collected from these frames each day to obtain 0-24 hr old individuals for barcoding. A total of 800 bees were used for each experimental colony, collected and barcoded over 1-2 days upon eclosion (File S10).

 <u>Barcoding bees</u>

Bees were tagged with "bCode" barcodes as in Gernat et al. (2018). Unique sets of

bCodes were used to differentiate bees barcoded on different days, as well as to differentiate bees

from different source colonies in colonies D-F. To attach bCodes to bees, workers were

anesthetized on ice and then positioned using soft forceps (BioQuip, Compton, CA). A small

drop of Loctite Super Glue Gel Control (Henkel, Düsseldorf, Germany) was applied to the center

of the thorax of each bee, followed by a bCode positioned with its left and right edge parallel to

the anteroposterior axis of the bee. Bees were carefully placed in plastic dishes until they

recovered from cold anesthetization, at which point the glue was dry. After waking, all bees were

placed in a large container with Fluon®-coated walls (Insect-a-Slip, BioQuip) where honey was

provided *ad libitum* until placement into a custom observation hive, described below. At the end

of each barcoding day, bees were carefully transferred into the observation hive.

Behavioral tracking

<u>Hive monitoring</u>

Barcoded bees were housed in a glass-walled observation hive with a one-sided plastic

honeycomb frame, as in Gernat et al. (2018). Bees were unable to access the back side of the

honeycomb, and could exit the hive through a plastic tube to the outside. Colonies were

maintained in a dark room with a heater and humidifier that kept the room at approximately 32°C

and 50% relative humidity.

Infrared light (not visible to bees) was used to illuminate the hive from both the front and

back while capturing hive images. Images were acquired at one-second resolution with a

monochrome Prosilica GX6600 machine vision camera (Allied Vision, Stadtroda, Germany) fitted

with a Nikkor AF 135 mm f/2 D DC prime lens (Nikon, Minato City, Japan). Additional details

about image acquisition can be found in Gernat et al. (2018). Images were saved to a redundant

array of independent disks, then copied onto a computing cluster (Biocluster, UIUC) for analysis

after the end of each experimental recording period.

<u>Entrance monitoring</u>

Colonies of barcoded bees were given access to the outside via a tube connected through

an exterior wall of the Bee Research Facility to an entrance equipped with an automated flight

activity monitor as in Geffre et al. (2020). This monitor included a maze to slow down incoming

and outgoing bees, and a Raspberry Pi camera (5 megapixel v1.3, Adafruit, New York, NY) that

imaged the maze twice per second from 07:00 until 19:00 daily. The camera was controlled by a

Raspberry Pi 2B computer running the Raspian 8 operating system, using a custom shell script and

the raspistill program to record images.

Barcode detection

Barcodes were detected in hive images as in Gernat et al. (2018) and filtered to facilitate

subsequent behavioral analyses. Filtering involved removal of potential tracking errors, including

removal of barcodes that did not pass read error correction. In addition, records for barcodes that

were read twice in the same image were removed, as were hive image records of the same barcode

identified more than 5 cm/second between successive detections, which are likely to be

misidentifications. An average of 94.51% of detections remained after these filtering steps (range

across colonies: 91.94-97.11%). Finally, the time of death of each bee was estimated using the last

time she was observed for at least 4 minutes during a 5-minute window above the third row of

honeycomb cells from the bottom of the hive; dead bees tend to accumulate below this level prior

to being removed by other bees (Gernat et al., 2018). Records for bees following their time of

death were filtered out so behavioral scores (below) were calculated only over times in which bees

were alive.

In entrance monitor images, barcodes were similarly detected as in hive images, but with

parameters adjusted for images produced by the Raspberry Pi camera. Fast-moving bees were not

filtered out in entrance images, because bees do move quickly through the entrance monitor and

due to the relatively small number of bees that fit into the maze, spurious fast movement due to

bCode decoding errors is unlikely.

Egg-laying detector

<u>Annotated image library</u>

Hive images from three experimental colonies and across 12 different days were used for

manual annotation of egg-laying events. The software Fiji (Schindelin et al., 2012) was used to

mark the bCode positions of all workers laying eggs in an initial set of 1500 hive images, followed

by an additional set of 782 images, each annotated by three independent observers. After the initial

identification of egg-laying bees in these images, the two seconds before and after each egg-laying

event were also annotated for those bees. Bees not marked as laying eggs with visible bCodes were

considered non-egg-laying for training of the CNN, below.

<u>CNN training and performance estimation</u>

Two convolutional neural networks (CNNs) were trained on the annotated egg-laying

images, using TensorFlow^{TM} (Abadi et al., 2016). The first CNN used images cropped to include

just a small rectangular region behind the barcode of each bee. For egg-laying bees, these images

show the honeycomb, because their abdomen is backed into the comb and thus not visible. For

non-layers, these images show the abdomen. The CNN was trained to differentiate between these

two cases. The second CNN was applied to images of bees that were identified as potential egg-

layers by the first CNN. It used slightly larger images that showed the entire bee and was trained

to use information about the bee's posture and her immediate surroundings to identify false

positives, which were subsequently filtered out.

Application of a CNN to an image results in a score between 0 and 1 that reflects the

likelihood of that image showing the event of interest. Deciding whether a score is sufficiently

high for assuming that the event took place involves thresholding that score. To choose thresholds

for each CNN score and a minimum egg-laying duration, a calibration set of images, which were

not used for training the CNNs, was used to estimate the performance of the egg-laying detector

for different threshold combinations. Thresholds were chosen from this calibration set to maximize

the detector's positive predictive value, then were applied to an independent test set of images that

had also never been seen by the detector to obtain unbiased performance values. Based on the

performance estimation on the test set of images, the egg-laying detector had the following

performance: 99.71% accuracy, 35.39% sensitivity, 100% specificity, 100% positive predictive

value, and 99.71% negative predictive value. Minimizing false positives came at a cost to

sensitivity, but bees who lay eggs will likely do so more than once over the course of the

experiment and can thus still be identified as egg-layers (honey bees possess multiple ovarioles,

each of which can develop eggs simultaneously (Hess, 1942)). Egg-laying detections were further

aggregated into events: subsequent detections that occurred within 10 seconds and 11.2 mm (the

width of two honeycomb cells) of one another were assumed to belong to the same egg-laying

event and were merged.

Filtering and annotation of entrance data

Raw detections of bees in the entrance were filtered as in Geffre et al. (2020). Briefly, a bee must traverse at least one-third the distance of the entrance monitor to be counted, and traversals that occurred within 10 seconds of each other were merged into a single event. These traversal events were then determined to be incoming or outgoing based on the positional coordinates of the bee at the start and end times of each event. Numbers of foraging trips (File S1) was inferred from series of outgoing and incoming events.

Incoming foraging trips were additionally annotated with trophallaxis data to determine whether a forager likely returned with nectar. CNNs trained to identify pairs of bees engaged in trophallaxis as well as the direction of trophallaxis (i.e., which bee was donor and which was recipient; Gernat et al., unpublished) were used to annotate incoming trips for all bees. Parameters used for the detector resulted in the following performance metrics based on test images: 88.7% sensitivity, 99.6% specificity, 90.4% positive predictive value, 99.6% negative predictive value, and 88.9% accuracy in determining trophallactic role (donor or receiver) of each bee. If a bee was a trophallaxis donor within 5 minutes after returning from a trip (Seeley, 2009), with no trophallaxis reception prior to the donation, that foraging trip was annotated as a nectar trip. Additionally, incoming trips were manually annotated for pollen on the hindlegs of returning bees for colonies D-F. Combining these nectar and pollen data for each trip, the proportion of foraging trips with nectar ("p.nectar"), pollen ("p.pollen") or both ("p.both") were calculated per bee in these colonies.

Specialist and generalist scores

In order to characterize the activity of egg-laying and foraging for each bee, two behavioral scores were created. The "specialist" score describes how specialized an individual was on either egg-laying (scores near -1) or foraging (scores near +1) relative to other bees in the colony; bees that consistently performed both egg-laying and foraging, or that performed neither behavior, have specialist scores near 0. The generalist score ranges from 0 to 1 and describes the degree to which an individual performed both egg-laying and foraging behaviors, differentiating bees with specialist scores near 0 based on the performance (or not) of egg-laying and foraging. Scores were created by first counting the number of egg-laying and foraging events per day. Bees were then ranked for each behavior relative to other bees in the colony on the same day, with tying ranks being assigned the minimal rank (e.g., if three bees were tied between the 4th and 8th ranked bees,

they all received a rank of 5). Ranks were then normalized by dividing by the maximum rank, so

that all ranks were in the range [0,1]. The normalized rank space for each bee (i.e., normalized

egg-laying rank and normalized foraging rank) was then mapped to behavioral scores (and

corresponding color space) using the following formulae in polar coordinates (ρ,Θ) on the two-

dimensional rank space: generalist score = $(1/2)\rho^2\sin^4 2\Theta$, specialist score= $\sin(\Theta-\pi/4)\rho^4\cos^4 2\Theta$.

Note that the numerical value of the scores has no biological meaning, but is simply a mapping

from rank space to the space of colors as shown in Fig. S6.

Selection of bees for sequencing

The median of specialist and generalist scores was weighted to emphasize the latter part of

the experiment; days 15-21 received a weight of 1-7, respectively, and each day's score was

multiplied by this weight. These scores were used to characterize the overall behavior of each bee

in the colony. The rank approach allowed for normalization across days with different overall

levels of activity in the colony, and the median score across days provides an overall assessment

of the lifetime behavior of each bee. These weighted median scores were used to rank all bees, and

the top ranking specialists and generalists from two colonies were selected for brain RNA

sequencing (RNAseq) and Assay for Transposase-Accessible Chromatin using sequencing

(ATACseq). Scores for each sequenced bee (n=45, 25 from colony E, 20 from colony F), as well

as total numbers of detected egg-laying and foraging events per bee, are provided in File S2.

To examine variation in behavior within and among groups, principal component analysis

(PCA) was performed on the following set of behavioral traits (see also File S2): number of eggs

laid, number of foraging events, proportion of trips with evidence of nectar collection, proportion

of trips with evidence of pollen collection, and proportion of trips with evidence of both nectar and

pollen. PCA was performed in R using the prcomp function and plotted using the ggplot2 package.

Tissue dissection and homogenization

At the end of behavioral tracking, bees were collected from each colony and stored at -

80°C. For bees selected for sequencing, abdomens of each bee were carefully removed on dry ice

and incubated for 16 hours at -20°C in RNA-later ICE (Life Technologies, Carlsbad, CA). Ovaries

were imaged and assessed for ovary development using a 1-5 scale adapted from (Hess, 1942) to

assign an ovary score; a score of 3-5 indicates ovary activation. These dissections confirmed that

egg-layers and generalists had activated ovaries, while many foragers did not. Ovary scores, as
well as number of ovarioles as determined from dissections, are given in File S2.

The head of each bee was freeze-dried at 300 milliTorr for 55 minutes, and whole brains
were removed from the head capsule in a dry ice ethanol bath (Schulz and Robinson, 1999).
Dissected brains were stored individually in 1.5 mL microcentrifuge tubes at -80°C until
extractions.

Brains were individually homogenized in 150 µL phosphate buffered saline (1X PBS,
Corning, Corning, NY, cat. #21-040-CV) with protein inhibitor complex (PIC, Complete Tablets,
EDTA-free Protease Inhibitor Cocktail from Roche, Basel, Switzerland, cat. #04693132001) using
a motorized pestle for 20 seconds. 50 µL of this homogenate was then pipetted into 450 µL cold
PBS+PIC and placed on ice for ATAC-seq library preparation (see below). The remaining 100 µL
homogenate was mixed with 500 µL RLT buffer (Qiagen, Hilden, Germany) with 1% β-
mercaptoethanol for use in the Qiagen RNeasy Mini Kit RNA extraction protocol (see below).

RNAseq library preparation and sequencing

Whole brain RNA was extracted from the 600 µL homogenate in RLT buffer after an
additional 30 second homogenization following the Qiagen RNeasy Mini Kit protocol, including
a DNase (Qiagen) treatment to remove genomic DNA. RNA quantities were determined for each
sample using a Qubit RNA HS Assay Kit (Invitrogen, Carlsbad, CA). High RNA integrity for all
samples was confirmed with Bioanalyzer 2100 RNA Pico chips (Agilent, Santa Clara, CA) prior
to library preparation.

RNAseq libraries were constructed and sequenced by the W.M. Keck Center for
Comparative and Functional Genomics at the Roy J. Carver Biotechnology Center (University of
Illinois at Urbana-Champaign). Libraries were constructed from 500 ng RNA per sample using the
TruSeq Stranded mRNA HT kit (Illumina, San Diego, CA) on an ePMotion 5075 robot
(Eppendorf, Hamburg, Germany). Libraries were uniquely barcoded, quantified, and pooled for
sequencing across 6 lanes with 100 nt single-end sequencing on the Illumina HiSeq 4000.

ATACseq library preparation and sequencing

The 500 µL tissue homogenate was additionally homogenized by aspirating through a 20
gauge needle followed by a 23 gauge needle 5 times each. Samples were centrifuged at 500g for
5 minutes at 4°C. Supernatant was removed, and cells were resuspended in 50 µL cold PBS+PIC.

15 µL of this cell suspension (approximately 1/10th of the total brain, ~100k cells) was placed into

a new microcentrifuge tube, and this was centrifuged at 500g for 5 minutes at 4°C as an additional

cell washing step. Supernatant was removed, and cells were gently resuspended in 50 µL cold lysis

buffer prepared as in Buenrostro et al. (2015). The remainder of the ATACseq library protocol

followed Buenrostro et al. (2015), with the exception of the final purification step, where a 0.8:1

ratio of Ampure XP beads (Beckman Coulter, Brea, CA) to sample was used to purify each library.

In addition to sample libraries, input libraries were constructed from thoracic genomic DNA from

a random bee from each colony per sequencing batch using 50 ng of genomic DNA (extracted

using the Gentra Puregene Tissue Kit from Qiagen, cat. #158667, following manufacturer's

protocol for DNA purification from 25 mg tissue but with 6 µL proteinase K and 4 µL RNase A

at the appropriate steps). Genomic DNA was transposed with Nextera Tn5 Transposase (Nextera

Kit, Illumina) following the ATACseq protocol immediately following the cell lysis step

(Buenrostro et al., 2015), again using an 0.8:1 Ampure XP bead clean-up at the end of the protocol.

A Qubit dsDNA HS Assay Kit (Invitrogen) was used to quantify each library, and library size and

quality was assessed using a Bioanalyzer High-Sensitivity DNA Analysis kit (Agilent).

ATACseq libraries, including input libraries, were pooled at equal nM concentrations and

a bead clean-up (0.8:1 ratio of Ampure XP beads to sample) was performed on the pool prior to

submission for sequencing. QC on the final pool was performed using qPCR and an AATI

Fragment Analyzer by the Keck Center. Libraries were sequenced across three lanes with 100 nt

paired-end sequencing on the Illumina HiSeq 4000 by the Keck Center.

Data processing and analysis

RNAseq

Sequencing of RNAseq libraries (n=45, 25 from colony E, 20 from colony F) produced

1,487,641,973 reads which survived quality and adapter trimming using Trimmomatic (version

0.36, parameters used: ILLUMINACLIP: 2:35:30 LEADING:20 TRAILING:20 MINLEN:30).

Trimmed reads were aligned to the *Apis mellifera* HAv3.1 genome (NCBI accession

GCA_003254395.2) using STAR (version 2.5.3) and default parameters, resulting in an average

of 96.7% reads mapping uniquely. The program featureCounts from the Subread package (version

1.5.2) was used to assign mapped reads to gene features from the GFF file from NCBI associated

with the *A. mellifera* HAv3.1 genome. On average, 84.8% of uniquely mapped reads were assigned

to gene features using featureCounts.

Gene counts were imported into R for differential expression analysis using edgeR. Genes

with less than 1 CPM in at least 2 samples were removed, and remaining count values were

normalized using the TMM method. Gene-wise variances were calculated by estimating tagwise

dispersions in edgeR on filtered gene count matrices for each group separately and plotted using

ggplot2. Tagwise dispersion estimates were followed by quasi-likelihood F-tests for each pairwise

comparison of groups. Differentially expressed genes (DEGs) for each pairwise comparison are

given in File S3.

ATACseq

ATACseq libraries (n=48, 25 from colony E, 20 from colony F, 3 input libraries) produced

1,110,401,018 paired-end reads which survived quality and adapter trimming using Trimmomatic

(version 0.38, parameters used: ILLUMINACLIP: 2:15:10 HEADCROP:10 LEADING:20

TRAILING 20 SLIDINGWINDOW:4:15 MINLEN:30). An average of 98.1% of reads mapped to

the *Apis mellifera* HAv3.1 genome using bwa mem (version 0.7.17, default parameters).

Duplicates were marked and removed prior to further processing using picard (version 2.10.1,

average duplication level 30.2%).

Peaks were called from deduplicated BAM files using MACS2 (version 2.1.1, command:

callpeak, with parameters: --nomodel -g 2.5e8 --nolambda --keep-dup all --slocal 10000) using the

appropriate colony and sequencing batch input as control. Peaks were called on each colony and

behavioral group separately, then merged and sorted using BEDTools (version 2.26.0, sort and

merge commands). This resulted in a total of 11,614 merged peaks with an average width of 721

bp. Mapped reads were counted to each peak per individual using featureCounts from the Subread

package (version 1.5.2). An average of 51.0% of reads were mapped to called peaks.

Peak counts were imported into R for differential accessibility analysis using edgeR. Peaks

with less than 1 CPM in at least 2 samples were removed, and remaining count values were

normalized using the TMM method. Gene-wise variances were calculated by estimating tagwise

dispersions in edgeR on filtered gene count matrices for each group separately and plotted using

ggplot2. Tagwise dispersion estimates were followed by quasi-likelihood F-tests for each pairwise

comparison of groups. Differentially accessible peak (DAP) results for each pairwise comparison

are given in File S4.

Functional annotation of differential expression and chromatin accessibility

Differential expression

Differentially expressed gene (DEG) lists were functionally annotated using Gene Ontology (GO) by first mapping putative orthologs between *Apis mellifera* and *Drosophila melanogaster* using reciprocal best BLASTP hits (e-value cutoff = 1e-5). Only DEGs with putative *D. melanogaster* orthologs were included for GO enrichment, and the background list used was all tested genes (those which passed the minimum expression threshold) with putative *D. melanogaster* orthologs. Enrichment tests for biological processes were conducted using GOrilla (Eden et al., 2009) with all significant DEGs (FDR-corrected p-value<0.05) against the background list. GO enrichment results for all DEG lists are given in File S3.

Differential accessibility

To functionally annotate DAPs, the midpoint coordinate of the 11,614 peaks identified with MACS2 were assigned to genes based on proximity to honey bee gene features (*Apis mellifera* HAv3.1 genome). The following features were considered per gene: promoters (1 kb upstream), introns, exons, 5' UTR, 3' UTR, upstream (10 kb upstream), and downstream (10 kb). Peaks not associated with any gene feature were classified as intergenic. When peaks were associated with multiple genes (e.g., the intron of one gene and the promoter of another), they were assigned to individual genes based on the following priority: promoter (highest priority), exon, 5' UTR, 3' UTR, intron, upstream, downstream (lowest priority). If a peak was present in the same highest priority class for multiple genes, it was randomly assigned to one gene. In this way, each peak was assigned to either a single gene or considered intergenic. Of the 11,614 peaks, 1822 were assigned to the promoter region of a gene, 776 to exons, 1326 to 5' UTRs, 273 to 3' UTRs, 4666 to introns, 1155 to upstream regions, 773 to downstream regions, and 823 peaks were located in intergenic regions.

As before with GO enrichment for DEGs, Differentially accessible peaks (DAPs) were functionally annotated by mapping peak-associated genes to putative orthologs in *D. melanogaster* using BLASTP. The background list for enrichment analyses was the list of peaks which met the minimum accessibility count threshold for analysis and which had putative orthologs in *D. melanogaster*. GOrilla (Eden et al., 2009) was used for enrichment tests. GO enrichment results for all DAP lists are given in File S4.

Motif enrichment of DAPs and DEG regulatory regions

Transcription factor (TF) motif enrichment analysis in this study was performed similarly to the methods described in Whitney et al. (2014). The overall approach is as follows, with details below. For each TF motif, 1) genomic windows were scored for the presence of the motif, 2) window scores were combined into scores for genomic segments of interest, representing either gene regulatory regions or accessibility peaks, 3) a set of motif targets was created using a fixed cutoff on the segment scores, and 4) a statistical test for enrichment was performed between segments that were motif targets and those that were significant in differential analysis.

Motif scores for genomic windows

First, we divided the honey bee genome (version HAv3.1, NCBI accession GCA_003254395.2) into 500 bp windows with 250 bp shifts. We gathered a collection of 223 representative TFs (Kapheim et al., 2015) and downloaded their DNA binding specificities (motifs) characterized as position weight matrices (PWMs) from FlyFactorSurvey (Zhu et al., 2011). Separately for each TF motif, we ran the Stubb algorithm (Sinha et al., 2003) on all genomic windows to score them for the presence of that TF's binding sites. Tandem repeats in the windows were masked using the Tandem Repeat Finder (Benson, 1999) before calculating the Stubb scores to avoid scoring the repeats as weak binding sites. Since the honey bee genome has significant local G/C heterogeneity (Sinha et al., 2006), we converted the raw Stubb scores for each window into G/C content-normalized empirical p-values. This was done by determining the rank of each window among all genomic windows of similar G/C content (when grouped into 20 G/C bins).

Scores for genomic segments

We defined two different collections of genomic segments (accessibility peaks and gene regulatory regions) to analyze with motif enrichment in this study. Since the genomic segments may overlap with a variable number of our genomic windows, we defined a length-adjusted motif score for each segment. This score was calculated using the score of the best scoring window in that segment for the given motif and the number of windows overlapping the segment, as follows:

$$sc_{seg} = 1 - (1 - pval_{best})^N$$

where,

sc_{seg} = *length-adjusted motif score for the segment*

N = number of windows that overlap with the scoring window

pval$_{best}$ = best G/C normalized empirical p-value among the N overlapping windows.

<u>Statistical test for TF enrichment</u>

TF enrichment was analyzed for two sets of regions: DAPs (Differentially Accessible

Peaks) and DEGs (Differentially Expressed Genes) (File S7).

For analysis of DAPs, the collection of genomic segments was defined as the combination

of all DAPs and randomly selected non-accessible parts of genome that had the same distribution

of lengths as those DAPs. The number of randomly selected genomic segments was set to 10 times

the number of DAP segments. For each motif, the top 200 scoring segments from the collection

were defined as the TF motif target set. Hypergeometric p-values were calculated for each motif-

DAP set pair (File S7) to quantify the significance of the overlap between the corresponding TF

motif target set and DAP set.

For DEGs, the collection of genomic segments was the regulatory regions of all genes in

the honey bee annotation (Amel_OGSv3.21). Each regulatory region was defined as 5kb upstream

to   2kb   downstream   of   the   transcriptional   start   site   of   its   gene

(http://veda.cs.uiuc.edu/beeMotifScores/). The top 500 scoring segments from the gene universe

were selected as the TF motif target set for each motif. Finally, the significance of the overlap for

each motif-DEG set pair (File S7) was calculated with the Hypergeometric p-value.

All p-values were then converted to q-values using the "qvalue" function in the R software

package qvalue (Storey et al., 2019) to control the false discovery rate from multiple hypothesis

testing.

For motifs enriched both within DAPs and DEG upstream regions, CentriMo (Bailey and

Machanick, 2012) from MEME Suite was used to calculate and plot the probability of motif

binding across 2 kb windows centered on the peak summit for DAPs and 7 kb windows (5kb

upstream and 2kb downstream of the transcriptional start site (TSS)) for DEGs. These probabilities

are shown in Fig. 4.

Individualized Gene Regulatory Network (GRN) analysis

To understand how TFs orchestrate transcriptional changes in the brain, we reconstructed

a gene regulatory network (GRN) model using the ASTRIX approach (Chandrasekaran, 2014;

Chandrasekaran et al., 2011). ASTRIX uses gene expression data to identify interactions between

TFs and their target genes. The ASTRIX algorithm has been previously used to infer brain GRN

models for various organisms including the honey bee (Bukhari et al., 2017; Saul et al., 2017;

Shpigler et al., 2017). These models showed significantly high accuracy in predicting gene

expression changes in the brain and identified TFs that regulate social behaviors.

Here we applied ASTRIX using the gene expression data of the 45 individual bees along

with a list of honey bee TFs as input to identify regulatory interactions. We normalized the

transcriptomics data prior to GRN construction using the ComBat algorithm (Johnson et al., 2007)

to minimize batch and colony effects in the data. The effectiveness of the normalization was

checked using PCA. Any TF predicted to interact with a given target gene by ASTRIX had to pass

through two criteria: 1) share a significant degree of mutual information with the target gene (p-

value $< 10^{-6}$), and 2) explain at least 10% of the variance of the target gene, quantified by Least

angle regression algorithm. Similarly, each target gene included in the GRN must be predicted

with a correlation of at least 0.8 by the ASTRIX model using expression levels of TFs.

The GRN model built by ASTRIX predicted 2,190 genes with a Pearson's correlation of

0.8 or higher using expression levels of TFs. Overall, the GRN inferred by ASTRIX contains 4,500

interactions between 190 TFs and the 2,190 target genes. The full GRN is in File S8.

To determine TFs correlated with specific behaviors, we first identified genes that were

strongly correlated with specific behavior scores across all individuals (FDR p-value of correlation

< 0.001). TFs whose targets were over-represented among the behavior-correlated genes were then

determined. Significance of the overlap between the list of behavior-correlated genes with targets

of each TF ("TF module") was estimated using the hypergeometric test.

Finally, to identify TF modules associated with expression changes in each individual

("Individualized TF modules"), genes that were upregulated or downregulated in each individual

were identified using z-transformation. Genes in each individual with z-scores above 2 (i.e., 2

standard deviations above mean) or below -2 were considered to be differentially expressed in an

individual. This list of genes was then overlapped with TF modules to identify modules

significantly associated with each individual using the hypergeometric test of overlap.

We used a Random Forests classification algorithm for predicting individual behavioral

group from TF expression levels. A leave-one-out cross validation analysis was performed wherein

the algorithm was trained using data from the remaining 44 individuals and then used to predict

the behavior of the 45[th] individual using its TF levels. The model achieved an accuracy of 82% in

predicting behavior. Performance of the model was evaluated by comparison with random

shuffling of the behavior labels. We made predictions 100 times with a different set of shuffled

labels and compared the accuracy of predictions (i.e., total individuals for which behavioral group

was correctly predicted) between the random model and the Random Forest algorithm using a t-

test ($p=1 \times 10^{-8}$). This suggests that TF expression levels can accurately forecast the behavior of

the individual, especially for specialists. The relative importance of each TF in predicting behavior

was determined using Out-of-bag predictor importance estimation, wherein each predictor's value

is permuted and the corresponding impact on model accuracy is determined (importance scores

given in File S8). The random forest classification algorithm was implemented in MATLAB with

default parameters for the number of predictors sampled (square root of the number of predictors,

in this case 258 TFs) and default values for the tree depth (n - 1, where n is the training data size).

Selection of candidate TFs involved in specialized phenotypes

Candidate TFs displayed in Fig. 6 were drawn from multiple analyses presented in this

paper and in Kapheim et al. (2015). "Enriched within DAPs" indicates enrichment of the TF motif

within forager vs. layer DAPs from the analysis of ATACseq data within this manuscript (see

Motif enrichment of DAPs and DEG promoters and File S7*)*. Similarly, "Enriched near DEGs"

indicates enrichment of the TF motif among putative regulatory regions of forager vs. layer DEGs

(see *Motif enrichment of DAPs and DEG promoters* and File S7*)*. "Module correlated with

behavior" indicates that TF module expression is significantly correlated with at least one

behavioral metric across individuals (see *Individualized Gene Regulatory Network (GRN) analysis*

and File S8). "Group Predictive TF" indicates the TF is among the 20 most informative for

predicting individual group membership based on TF expression (see *Individualized Gene*

Regulatory Network (GRN) analysis and File S8). "Implicated in eusocial evolution" indicates that

the TF motif was previously found to be associated with social evolution in Kapheim et al. (2015).

Acknowledgements

We thank A. Sankey, A. Ray, S. Bransley, J. Cullum, K. Wilk and J. Falk for assistance

in the field, A. Hernandez and the staff at the Carver Biotechnology Center for sequencing

services, administrators of Biocluster (UIUC) for computational support, and M.B. Sokolowski,

M. Hudson, A.M. Bell, members of the Robinson lab, and three anonymous reviewers for

comments that improved this manuscript. Funding: This research was supported by Grant

R01GM117467 from the National Institute of General Medical Sciences (GER and N.

Goldenfeld, PIs), the Christopher Foundation (GER), and the Illinois Sociogenomics Initiative

(GER). Author contributions: B.M.J. and G.E.R. conceptualized the study and designed

experiments; B.M.J. performed behavioral experiments; V.D.R., T.G., and T.J. developed

methodology and software to collect and analyze data; B.M.J. and A.C-A. performed molecular

work; B.M.J., B.E.R.R., T.J.C., S.B., S.S.H., and C.B. performed or aided in data analysis;

M.M., S.S., S.C., and G.E.R. provided mentorship and project oversight throughout data

collection, analysis, and interpretation. B.M.J. and G.E.R. drafted the initial manuscript; all

authors contributed to editing and final writing of manuscript. Competing interests: The authors

declare no competing interests. Data and materials availability: The raw sequence data reported

in this paper have been deposited at the National Center for Biotechnology Information (NCBI)

Sequence Read Archive, Accession PRJNA593999. Requests for materials should be addressed

to B.M.J.

Competing interests

No competing interests declared.

References

Abadi M, Barham P, Chen J, Chen Z, Davis A, Dean J, Devin M, Ghemawat S, Irving G, Isard

M, Kudlur M, Levenberg J, Monga R, Moore S, Murray DG, Steiner B, Tucker P,

Vasudevan V, Warden P, Wicke M, Yu Y, Zheng X, Brain G, Osdi I, Barham P, Chen J,

Chen Z, Davis A, Dean J, Devin M, Ghemawat S, Irving G, Isard M, Kudlur M, Levenberg

J, Monga R, Moore S, Murray DG, Steiner B, Tucker P, Vasudevan V, Warden P, Wicke

M, Yu Y, Zheng X. 2016. TensorFlow : A system for large-scale machine learning. *OSDI*

16:265–283. doi:10.1126/science.aab4113.4

Adams J, Rothman ED, Kerr WE, Paulino ZL. 1977. Estimation of the number of sex alleles and

queen matings from diploid male frequencies in a population of Apis mellifera. *Genetics*

86:583–596.

Alaux C, Le Conte Y, Adams HA, Rodriguez-Zas S, Grozinger CM, Sinha S, Robinson GE.

2009. Regulation of brain gene expression in honey bees by brood pheromone. *Genes,*

Brain Behav **8**:309–319. doi:10.1111/j.1601-183X.2009.00480.x

Amdam G V, Page Jr RE, Tsuruda JM. 2008. Sensory response system of social behavior tied to

female reproductive traits. *PLoS One* **3**:e3397.

Araya CL, Kawli T, Kundaje A, Jiang L, Wu B, Vafeados D, Terrell R, Weissdepp P,

Gevirtzman L, Mace D, Niu W, Boyle AP, Xie D, Ma L, Murray JI, Reinke V, Waterston

RH, Snyder M. 2014. Regulatory analysis of the *C. elegans* genome with spatiotemporal

resolution. *Nature* **512**:400–405. doi:10.1038/nature16075

Bailey TL, Machanick P. 2012. Inferring direct DNA binding from ChIP-seq. *Nucleic Acids Res*

40:e128.

Barth RH, Lester LJ, Sroka P, Kessler T, Hearn R. 1975. Juvenile hormone promotes dominance

behavior and ovarian development in social wasps (*Polistes annularis*). *Experientia*

31:691–692. doi:10.1007/BF01944632

Benson G. 1999. Tandem repeats finder: A program to analyze DNA sequences. *Nucleic Acids*

Res. doi:10.1093/nar/27.2.573

Branstetter MG, Danforth BN, Pitts JP, Faircloth BC, Ward PS, Buffington ML, Gates MW,

Kula RR, Brady SG. 2017. Phylogenomic insights into the evolution of stinging wasps and

the origins of ants and bees. *Curr Biol* **27**:1019–1025. doi:10.1016/j.cub.2017.03.027

Buenrostro J, Giresi P, Zaba L, Chang H, Greenleaf W. 2013. Transposition of native chromatin

for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and

nucleosome position. *Nat Methods* **10**:1213–1218.

Buenrostro JD, Wu B, Chang HY, Greenleaf WJ. 2015. ATAC-seq: A Method for Assaying

Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* **109**:21.29.1-21.29.9.

Bukhari SA, Saul MC, James N, Bensky MK, Stein LR, Trapp R, Bell AM. 2019. Neurogenomic

insights into paternal care and its relation to territorial aggression. *Nat Commun* **10**:4437.

doi:10.1038/s41467-019-12212-7

Bukhari SA, Saul MC, Seward CH, Zhang H, Bensky M, James N, Zhao SD, Chandrasekaran S,

Stubbs L, Bell AM. 2017. Temporal dynamics of neurogenomic plasticity in response to

social interactions in male threespined sticklebacks. *PLoS Genet* **13**:e1006840.

doi:10.1371/journal.pgen.1006840

Chan YF, Marks ME, Jones FC, Villarreal G, Shapiro MD, Brady SD, Southwick AM, Absher

DM, Grimwood J, Schmutz J, Myers RM, Petrov D, Jónsson B, Schluter D, Bell MA,

Kingsley DM. 2010. Adaptive evolution of pelvic reduction in sticklebacks by recurrent
deletion of a pitxl enhancer. *Science* **327**:302–5. doi:10.1126/science.1182213

Chandrasekaran S. 2014. Predicting phenotype from genotype through reconstruction and
integrative modeling of metabolic and regulatory networksA Systems Theoretic Approach
to Systems and Synthetic Biology I: Models and System Characterizations. pp. 307–325.
doi:10.1007/978-94-017-9041-3_12

Chandrasekaran S, Ament SA, Eddy JA, Rodriguez-Zas SL, Schatz BR, Price ND, Robinson GE.
2011. Behavior-specific changes in transcriptional modules lead to distinct and predictable
neurogenomic states. *Proc Natl Acad Sci* **108**:18020.

Cnaani J, Schmid-Hempel R, Schmidt JO. 2002. Colony development, larval development and
worker reproduction in *Bombus impatiens* Cresson. *Insectes Soc* **49**:164–170.
doi:10.1007/s00040-002-8297-8

Crall JD, Gravish N, Mountcastle AM, Combes SA. 2015. BEEtag: A low-cost, image-based
tracking system for the study of animal behavior and locomotion. *PLoS One* **10**:e0136487.
doi:10.1371/journal.pone.0136487

Cummings ME, Larkins-Ford J, Reilly CRL, Wong RY, Ramsey M, Hofmann HA. 2008. Sexual
and social stimuli elicit rapid and contrasting genomic responses. *Proc R Soc B Biol Sci*
275:393–402. doi:10.1098/rspb.2007.1454

Dadant & Sons. 1975. Activities of queenless bees In: Dadant & Sons, editor. The Hive and the
Honey Bee. Carthage, Illinois: Journal Printing Company. pp. 245–247.

Dornhaus A. 2008. Specialization does not predict individual efficiency in an ant. *PLoS Biol*
6:2368–2375. doi:10.1371/journal.pbio.0060285

Dufour HD, Koshikawa S, Finet C. 2020. Temporal flexibility of gene regulatory network
underlies a novel wing pattern in flies. *Proc Natl Acad Sci U S A* **117**:11589–11596.
doi:10.1073/pnas.2002092117

Eden E, Navon R, Steinfeld I, Lipson D, Yakhini Z. 2009. GOrilla: A tool for discovery and
visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* **10**:48.
doi:10.1186/1471-2105-10-48

Estoup A, Solignac M, Cornuet JM. 1994. Precise assessment of the number of patrilines and of
genetic relatedness in honeybee colonies. *Proc R Soc B Biol Sci* **258**:1–7.
doi:10.1098/rspb.1994.0133

Fine JD, Shpigler HY, Ray AM, Beach NJ, Sankey AL, Cash-Ahmed A, Huang ZY,
Astrauskaite I, Chao R, Zhao H, Robinson GE. 2018. Quantifying the effects of pollen
nutrition on honey bee queen egg laying with a new laboratory system. *PLoS One*
13:e0203444. doi:10.1371/journal.pone.0203444

Free JB. 1955. The behaviour of egg-laying workers of bumblebee colonies. *Br J Anim Behav*
3:147–153. doi:10.1016/S0950-5601(55)80053-6

Gadagkar R. 1997. The evolution of caste polymorphism in social insects genetic release
followed by diversifying evolution. *J Genet* **76**:167–179. doi:10.1007/BF02932215

Geffre AC, Gernat T, Harwood GP, Jones BM, Gysi DM, Hamilton AR, Bonning BC, Toth AL,
Robinson GE, Dolezal AG. 2020. Honey bee virus causes context-dependent changes in
host social behavior. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.2002268117

Gençer HV, Kahya Y. 2011. Are sperm traits of drones (*Apis mellifera* L.) from laying worker
colonies noteworthy? *J Apic Res* **50**:130–137. doi:10.3896/IBRA.1.50.2.04

Gernat T, Rao VD, Middendorf M, Dankowicz H, Goldenfeld N, Robinson GE. 2018.
Automated monitoring of behavior reveals bursty interaction patterns and rapid spreading
dynamics in honeybee social networks. *Proc Natl Acad Sci* **115**:1433–1438.
doi:10.1073/pnas.1713568115

Graham AM, Munday MD, Kaftanoglu O, Page RE, Amdam G V., Rueppell O. 2011. Support
for the reproductive ground plan hypothesis of social evolution and major QTL for ovary
traits of Africanized worker honey bees (*Apis mellifera* L.). *BMC Evol Biol* **11**:95.
doi:10.1186/1471-2148-11-95

Greenwald E, Segre E, Feinerman O. 2015. Ant trophallactic networks: simultaneous
measurement of interaction patterns and food dissemination. *Sci Rep* **5**:12496.

Grozinger CM, Fan Y, Hoover SER, Winston ML. 2007. Genome-wide analysis reveals
differences in brain gene expression patterns associated with caste and reproductive status
in honey bees (*Apis mellifera*). *Mol Ecol* **16**:4837–4848.

Hamilton AR, Shpigler H, Bloch G, Wheeler DE, Robinson GE. 2016. Endocrine Influences on
Insect Societies.Hormones, Brain and Behavior: Third Edition. doi:10.1016/B978-0-12-
803592-4.00037-7

Hamilton AR, Traniello IM, Ray AM, Caldwell AS, Wickline SA, Robinson GE. 2019. Division
of labor in honey bees is associated with transcriptional regulatory plasticity in the brain. *J*

Exp Biol **222**:jeb200196. doi:10.1242/jeb.200196

Harpur BA, Owayss AA, Lebon JMD, Zayed A, Alqarni AS, Molodtsova D, Kent CF. 2014. Population genomics of the honey bee reveals strong signatures of positive selection on worker traits. *Proc Natl Acad Sci* **111**:2614–2619.

Hess G. 1942. Über den einfluss der weisellosigkeit und des fruchtbarkeitsvitamins e auf die ovarien der bienenarbeiterin. *Schweiz Bienen Ztg* **2**:33–110.

Holldobler B, Wilson EO. 1990. The Ants. Cambridge, MA: Belknap/Harvard Univ. Press.

Jeanson R, Clark RM, Holbrook CT, Bertram SM, Fewell JH, Kukuk PF. 2008. Division of labour and socially induced changes in response thresholds in associations of solitary halictine bees. *Anim Behav* **76**:593–602. doi:10.1016/j.anbehav.2008.04.007

Johnson WE, Li C, Rabinovic A. 2007. Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics* **8**:118–127. doi:10.1093/biostatistics/kxj037

Jones BM, Kingwell CJ, Wcislo WT, Robinson GE. 2017. Caste-biased gene expression in a facultatively eusocial bee suggests a role for genetic accommodation in the evolution of eusociality. *Proc R Soc B Biol Sci* **284**:20162228.

Kapheim KM, Jones BM, Pan H, Li C, Harpur BA, Kent CF, Zayed A, Ioannidis P, Waterhouse RM, Kingwell C, Stolle E, Avalos A, Zhang G, McMillan WO, Wcislo WT. 2020. Developmental plasticity shapes social traits and selection in a facultatively eusocial bee. *Proc Natl Acad Sci U S A* **117**:13615–13625. doi:10.1073/pnas.2000344117

Kapheim KM, Pan H, Li C, Salzberg SL, Puiu D, Magoc T, Robertson HM, Hudson ME, Venkat A, Fischman BJ, Hernandez A, Yandell M, Ence D, Holt C, Yocum GD, Kemp WP, Bosch J, Waterhouse RM, Zdobnov EM, Stolle E, Kraus FB, Helbing S, Moritz RFA, Glastad KM, Hunt BG, Goodisman MAD, Hauser F, Grimmelikhuijzen CJP, Pinheiro DG, Nunes FMF, Soares MPM, Tanaka ÉD, Simões ZLP, Hartfelder K, Evans JD, Barribeau SM, Johnson RM, Massey JH, Southey BR, Hasselmann M, Hamacher D, Biewer M, Kent CF, Zayed A, Blatti C, Sinha S, Johnston JS, Hanrahan SJ, Kocher SD, Wang J, Robinson GE, Zhang G. 2015. Genomic signatures of evolutionary transitions from solitary to group living. *Science* **348**:1139–1143.

Lobo J, Kerr W. 1993. Estimation of the number of matings in *Apis mellifera*: extensions of the model and comparison of different estimates. *Ethol Ecol Evol* **5**:337–345.

Mallarino R, Campàs O, Fritz JA, Burns KJ, Weeks OG, Brenner MP, Abzhanov A. 2012.

Closely related bird species demonstrate flexibility between beak morphology and underlying developmental programs. *Proc Natl Acad Sci U S A* **109**:16222–16227. doi:10.1073/pnas.1206205109

Mello C V, Vicario DS, Clayton DF. 1992. Song presentation induces gene expression in the songbird forebrain. *Proc Natl Acad Sci U S A* **89**:6818–6822. doi:10.1073/pnas.89.15.6818

Mersch DP, Crespi A, Keller L. 2013. Tracking individuals shows spatial fidelity is a key regulator of ant social organization. *Science* **340**:1090. doi:10.1126/science.1234316

Michener CD. 1974. The Social Behavior of the Bees: A Comparative Study. Cambridge, MA: Harvard University Press.

Miller III DG, Ratnieks FLW. 2001. The timing of worker reproduction and breakdown of policing behaviour in queenless honey bee (*Apis mellifera* L.) societies. *Insectes Soc* **48**:178.

Morse RA. 1990. Laying workers In: Morse RA, Flottum K, editors. The ABC & XYZ of Bee Culture. The A. I. Root Company. pp. 291–293.

Mukherjee D, Ignatowska-Jankowska BM, Itskovits E, Gonzales BJ, Turm H, Izakson L, Haritan D, Bleistein N, Cohen C, Amit I, Shay T, Grueter B, Zaslaver A, Citri A. 2018. Salient experiences are represented by unique transcriptional signatures in the mouse brain. *Elife* **7**:e31220. doi:10.7554/eLife.31220

Naeger NL, Peso M, Even N, Barron AB, Robinson GE. 2013. Altruistic behavior by egg-laying worker honeybees. *Curr Biol* **23**:1574–1578. doi:10.1016/j.cub.2013.06.045

O'Donnell S. 1998. Reproductive caste determination in eusocial wasps (Hymenoptera: Vespidae). *Annu Rev Entomol* **43**:323–346. doi:10.1146/annurev.ento.43.1.323

Oster GF, Wilson EO. 1979. Caste and Ecology in the Social Insects. Princeton: Princeton University Press. doi:10.1007/s00265-010-1009-x

Page Jr RE, Erickson Jr EH. 1988. Reproduction by worker honey bees (Apis mellifera). *Behav Ecol Sociobiol* **23**:117–126. doi:10.1007/BF00299895

Page Jr RE, Rueppell O, Amdam G V. 2012. Genetics of reproduction and regulation of honey bee (*Apis mellifera* L.) social behavior. *Annu Rev Genet* **46**:19–97.

Page RE, Robinson GE. 1994. Reproductive competition in queenless honey bee colonies (*Apis mellifera* L.). *Behav Ecol Sociobiol* **35**:99–107. doi:10.1007/BF00171499

Prud'homme B, Gompel N, Rokas A, Kassner VA, Williams TM, Yeh SD, True JR, Carroll SB.

2006. Repeated morphological evolution through *cis*-regulatory changes in a pleiotropic gene. *Nature* **440**:1050–3. doi:10.1038/nature04597

Ratnieks FLW, Foster KR, Wenseleers T. 2006. Conflict resolution in insect societies. *Annu Rev Entomol* **51**:581–608. doi:10.1146/annurev.ento.51.110104.151003

Ratnieks FLW, Wenseleers T. 2008. Altruism in insect societies and beyond: voluntary or enforced? *Trends Ecol Evol* **23**:45–52. doi:10.1016/j.tree.2007.09.013

Reed RD, Papa R, Martin A, Hines HM, Counterman BA, Pardo-Diaz C, Jiggins CD, Chamberlain NL, Kronforst MR, Chen R, Halder G, Nijhout HF, McMillan WO. 2011. optix drives the repeated convergent evolution of butterfly wing pattern mimicry. *Science* **333**:1137.

Rittschof CC, Bukhari SA, Sloofman LG, Troy JM, Caetano-Anollés D, Cash-Ahmed A, Kent M, Lu X, Sanogo YO, Weisner PA, Zhang H, Bell AM, Ma J, Sinha S, Robinson GE, Stubbs L. 2014. Neuromolecular responses to social challenge: Common mechanisms across mouse, stickleback fish, and honey bee. *Proc Natl Acad Sci* **111**:17929–17934.

Robinson GE. 1992. Regulation of division of labor in insect societies. *Annu Rev Entomol* **37**:637–665.

Robinson GE, Page RE, Fondrk MK. 1990. Intracolonial behavioral variation in worker oviposition, oophagy, and larval care in queenless honey bee colonies. *Behav Ecol Sociobiol* **26**:315–323. doi:10.1007/BF00171096

Robinson GE, Strambi C, Strambi A, Huang ZY. 1992. Reproduction in worker honey bees is associated with low juvenile hormone titers and rates of biosynthesis. *Gen Comp Endocrinol* **87**:471–480. doi:10.1016/0016-6480(92)90055-O

Sakagami SF. 1954. Occurrence of an aggressive behaviour in queenless hives, with considerations on the social organization of honeybee. *Insectes Soc* **1**:331–343. doi:10.1007/BF02329618

Saul MC, Seward CH, Troy JM, Zhang H, Sloofman LG, Lu X, Weisner PA, Caetano-Anolles D, Sun H, Zhao SD, Chandrasekaran S, Sinha S, Stubbs L. 2017. Transcriptional regulatory dynamics drive coordinated metabolic and neural response to social challenge in mice. *Genome Res* **27**:959–972. doi:10.1101/gr.214221.116

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A. 2012. Fiji: An open-source platform for biological-image analysis. *Nat Methods* **9**:676. doi:10.1038/nmeth.2019

Schulz DJ, Robinson GE. 1999. Biogenic amines and division of labor in honey bee colonies: Behaviorally related changes in the antennal lobes and age-related changes in the mushroom bodies. *J Comp Physiol A* **184**:481–488. doi:10.1007/s003590050348

Seeley TD. 2009. The Wisdom of the Hive: the social physiology of honey bee colonies, Proceedings of the Royal Society B: Biological Sciences. doi:10.1098/rspb.2010.0491

Shpigler HY, Saul MC, Murdoch EE, Cash-Ahmed AC, Seward CH, Sloofman L, Chandrasekaran S, Sinha S, Stubbs LJ, Robinson GE. 2017. Behavioral, transcriptomic and epigenetic responses to social challenge in honey bees. *Genes, Brain Behav* **16**:579–591. doi:10.1111/gbb.12379

Simola DF, Graham RJ, Brady CM, Enzmann BL, Desplan C, Ray A, Zwiebel LJ, Bonasio R, Reinberg D, Liebig J, Berger SL. 2015. Epigenetic (re)programming of caste-specific behavior in the ant *Camponotus floridanus*. *Science* **351**:aac6633.

Sinha S, Jones BM, Traniello IM, Bukhari SA, Halfon MS, Hofmann HA, Huang S, Katz PS, Keagy J, Lynch VJ, Sokolowski MB, Stubbs LJ, Tabe-Bordbar S, Wolfner MF, Robinson GE. 2020. Behavior-related gene regulatory networks: A new level of organization in the brain. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.1921625117

Sinha S, Ling X, Whitfield CW, Zhai C, Robinson GE. 2006. Genome scan for *cis*-regulatory DNA motifs associated with social behavior in honey bees. *Proc Natl Acad Sci U S A* **103**:16352–16357. doi:10.1073/pnas.0607448103

Sinha S, van Nimwegen E, Siggia ED. 2003. A probabilistic method to detect regulatory modules. *Bioinformatics* **19 Suppl 1**:i292-301. doi:10.1093/bioinformatics/btg1040

Storey JD, Bass AJ, Dabney A, Robinson D. 2019. qvalue: Q-value estimation for false discovery rate control. R package version 2.14.1.

Toth AL, Várala K, Henshaw MT, Rodriguez-Zas SL, Hudson ME, Robinson GE. 2010. Brain transcriptomic analysis in paper wasps identifies genes associated with behaviour across social insect lineages. *Proc R Soc B Biol Sci* **277**:2139–2148. doi:10.1098/rspb.2010.0090

Trumbo ST, Robinson GE. 1997. Learning and task interference by corpse-removal specialists in honey bee colonies. *Ethology* **103**:966–975. doi:10.1111/j.1439-0310.1997.tb00138.x

Turillazzi S, West-Eberhard M. 1996. Natural History and Evolution of Paper-Wasps. Oxford:

Oxford University Press.

Velthuis HHW. 1970. Ovarian development in *Apis mellifera* worker bees. *Entomol Exp Appl* **13**:377.

Velthuis HHW, Duchateau MJ. 2011. Ovarian development and egg laying in workers of *Bombus terrestris*. *Entomol Exp Appl* **51**:199–213.

Wario F, Wild B, Couvillon MJ, Rojas R, Landgraf T. 2015. Automatic methods for long-term tracking and the detection and decoding of communication dances in honeybees. *Front Ecol Evol* **3**:103. doi:10.3389/fevo.2015.00103

Wenseleers T, Ratnieks FLW. 2006. Enforced altruism in insect societies. *Nature* **444**:50. doi:10.1038/444050a

Werner T, Koshikawa S, Williams TM, Carroll SB. 2010. Generation of a novel wing colour pattern by the Wingless morphogen. *Nature* **464**:1143–1148. doi:10.1038/nature08896

West-Eberhard MJ. 1987. Flexible strategy and social evolution, Animal societies: theories and facts. Tokyo: Japan Scientific Societies Press.

Wheeler DE. 1986. Developmental and physiological determinants of caste in social Hymenoptera: Evolutionary implications. *Am Nat* **128**:13–34.

Whitfield CW, Cziko AM, Robinson GE. 2003. Gene expression profiles in the brain predict behavior in individual honey bees. *Science* **302**:296–299. doi:10.1126/science.1086807

Whitney O, Pfenning AR, Howard JT, Blatti CA, Liu F, Ward JM, Wang R, Audet JN, Kellis M, Mukherjee S, Sinha S, Hartemink AJ, West AE, Jarvis ED. 2014. Core and region-enriched networks of behaviorally regulated genes and the singing genome. *Science* **346**:1256780. doi:10.1126/science.1256780

Wilson EO. 1980. Caste and division of labor in leaf-cutter ants (Hymenoptera: Formicidae: *Atta*). *Behav Ecol Sociobiol* **7**:157–165. doi:10.1007/BF00299521

Wilson EO. 1971. The insect societies. Cambridge: Harvard University Press.

Woodard SH, Fischman BJ, Venkat A, Hudson ME, Varala K, Cameron SA, Clark AG, Robinson GE. 2011. Genes involved in convergent evolution of eusociality in bees. *Proc Natl Acad Sci* **108**:7472–7477. doi:10.1073/pnas.1103457108

Zayed A, Robinson GE. 2012. Understanding the relationship between brain gene expression and social behavior: Lessons from the honey bee. *Annu Rev Genet* **46**:591–615.

Zhu LJ, Christensen RG, Kazemian M, Hull CJ, Enuameh MS, Basciotta MD, Brasefield JA,

1143   Zhu C, Asriyan Y, Lapointe DS, Sinha S, Wolfe SA, Brodsky MH. 2011. FlyFactorSurvey:

1144   A database of *Drosophila* transcription factor binding specificities determined using the

1145   bacterial one-hybrid system. *Nucleic Acids Res*. doi:10.1093/nar/gkq858

1146

1147 **Supplemental Figures**

1148

1149 **Fig. S1.** Histogram (bars) and density (lines) of normalized (logCPM) gene expression for genes
1150 with (dark gray) and without (light gray) nearby peaks of chromatin accessibility. Distributions
1151 are significantly different (p<0.0001, Kolmogorov-Smirnov test).

1152

Fig. S2. Normalized expression (logCPM, scaled to a maximum of 1 to allow for comparison across TFs) of the top 20 most informative TFs for class prediction analysis plotted against individual specialist score. Points are colored by behavioral group (Layer: purple, Generalist: orange, Forager: green). Although some of the top predictors individually show weak correlation with specialist score, the random forest machine learning algorithm combined multiple weak predictors together in a single model to accurately classify the three behavioral groups, suggesting the TFs act combinatorially to influence behavior.

Direct PPI   First-order PPI   Second-order PPI

Fig. S3. Network of 23 TFs with module expression significantly correlated with 9 behavioral and physiological metrics (see File S2) measured across individuals. Edges indicate known interactions based on MIST database for Drosophila melanogaster. First-order PPI indicates one intermediate protein between linked nodes, while second-order PPI indicates two intermediate proteins. PPI = protein-protein interaction.

1166

Fig. S4. Annotated honey bee genes in the glycolysis/gluconeogenesis KEGG pathway (Kyoto
Encyclopedia of Genes and Genomes, http://www.genome.jp/kegg/). Genes are colored to
indicate whether they are differentially expressed (DEG) between foragers (F) and layers (L),
and hashed if they have a nearby differentially accessible peak (DAP). Thick borders indicate
genes which were found to be under selection in at least one of three studies of selection in bees
(Harpur et al., 2014; Kapheim et al., 2015; Woodard et al., 2011). "Not expressed" genes refer to
those not expressed above the minimum threshold in the F vs. L dataset, while non-DEG genes
were expressed but not differentially expressed in F vs. L at an FDR-corrected p<0.05.

Fig. S5. Smoothed average egg counts for laying workers in laboratory cages. Solid lines indicate bees from source colonies headed by a naturally-mated queen, while dashed lines indicate bees from source colonies headed by queens instrumentally inseminated with semen from a single drone (SDI).

$$\text{specialist score} = \sin\left(\theta - \frac{\pi}{4}\right)\rho^4 \cos^4 2\theta$$

$$\text{generalist score} = \frac{1}{2}\rho^2 \sin^4 2\theta$$

Fig. S6. Formulae and color-space mapping for specialist (left) and generalist (right) behavioral scores.

Supplemental Files

File S1: Daily egg-laying counts, foraging counts, specialist scores, and generalist scores for individual bees

File S2: Detailed behavioral and physiological information for sequenced bees

File S3: Differentially expressed genes (DEGs) and Gene Ontology (GO) enrichment of DEGs for each pairwise comparison of specialists and generalists

File S4: Differentially accessible peaks (DAPs) and Gene Ontology (GO) enrichment of DAPs for each pairwise comparison of specialists and generalists

File S5: Lists of genes with upper and lower 5% of PC loadings for gene expression and Gene Ontology (GO) enrichment results

File S6: Lists of genes with upper and lower 5% of PC loadings for chromatin accessibility and Gene Ontology (GO) enrichment results

File S7: TF motif enrichment of DAPs and DEGs

File S8: Predicted gene regulatory network (GRN), GRN module correlations with behavior and physiological measurements, and importance scores of TFs from class prediction analysis

File S9: Overlaps and statistics for comparative gene expression datasets

File S10: Details of experimental setup for recorded colonies