Pervasive selection pressure in wild and domestic pigs

J. Leno-Colorado[1], S. Guirao-Rico[12], M. Pérez-Enciso[134], S. E. Ramos-Onsins[1*]

[1] Centre for Research in Agricultural Genomics (CRAG) Consortium CSIC-IRTA-UAB-UB. 08193

Bellaterra, Spain.

[2] Institute of Evolutionary Biology, CSIC-Universitat Pompeu Fabra, Barcelona, Spain.

[3] Universitat Autònoma de Barcelona, Department of Animal Science, Bellaterra, Spain.

[4] Institut Català de Recerca I Estudis Avançats (ICREA), Carrer de Lluís Companys 23, Barcelona, Spain.

*Corresponding author: sebastian.ramos@cragenomica.es

11   **ABSTRACT**

Animal domestication typically affected numerous polygenic quantitative traits, such as behavior,

development and reproduction. However, uncovering the genetic basis of quantitative trait variation is

challenging, since they are caused by small allele-frequency changes. To date, only a few causative

mutations related to domestication processes have been reported, strengthening the hypothesis that small

effect variants have a prominent role. So far, approaches on domestication have been limited to the detection

of the global effect of domestication on deleterious mutations and on strong beneficial variants, ignoring

the importance of variants with small selective effects. To overcome these difficulties, here we propose to

estimate the proportion of beneficial variants based on the asymptotic MacDonald Kreitman (MK) method,

according to estimates of variability based on frequency spectrum. We applied this approach to the pig

species, analyzing 46 complete genome sequences from 20 European wild boars, 6 Iberian and 20 Large

White pigs at different molecular scales: gene, metabolic pathway and whole-genome.

Descriptive variability analyses on pig populations indicate that domestic and wild pig populations do not

differ in nonsynonymous fixed mutations. Instead, most variants are shared among them, despite that the

phenotypes of wild and domestic individuals are clearly divergent. Additionally, asymptotic MK plots

based on summary statistics show that small effects variants may affect the final calculation of α, the

proportion of beneficial mutations. The distribution of fitness effects inferred with Approximate Bayesian

Computation analysis indicates that both wild and domestic pigs display an important quantity of

deleterious mutations at low frequency (~83% of total mutations) and a high number of nearly-neutral

mutations (~17%) that may have a significant effect on the evolution of domestic and wild populations.

Exclusive mutations show that recent demographic changes have severely affected the fitness of

populations, especially of the local Iberian breed. Finally, the median proportion of the strong favorable

mutations are very scarce in all cases ($\leq$ 0.2%). The median estimated alpha values (weak and strong

favorable) are 0.9% for wild and domestic pigs.

Keywords: Domestication, Distribution of fitness effects, Proportion of beneficial mutations, Approximate

Bayesian calculation, Polygenic selection, Population genomics

INTRODUCTION

Domestic animal histories are evolutionary experiments that have often lasted for millennia, with the result of dramatic phenotypic changes to suit human needs. In addition, domestic species can be structured into subpopulations (breeds) that are partly or completely genetically isolated and can display a wide catalog of specific phenotypes. Therefore, they offer a material of utmost interest to study the interplay of demography and accelerated adaptation. However, as their demographic history can be quite complex, many events remain unknown or poorly documented nowadays.

The pig (*Sus scrofa*) is a particularly interesting species because of its domestication history and its relatively well-annotated genome. *S. scrofa* originated in Southeast Asia ~ 4 MYA and spread throughout Eurasia ~1.2 MYA, colonizing all climates except the driest (Frantz et al. 2013). Subsequently, the pig was domesticated from local wild boars independently in both Asia and Europe ~9,000 years ago. To complicate the story, modern European domestic pig breeds were crossed with Asian domestic pigs during the late 17th century and onwards. In breeds such as Large White (LW), approximately 30% of the genome is estimated to be of Asian origin (Bosse, Megens, Madsen, et al. 2014). Nevertheless, some local European breeds, such as the Iberian breed (IB), were spared genetic contact with Asian pigs and no evidence of genetic introgression has been found in this breed (Alves et al. 2003, Esteve-Codina et al. 2013). Moreover, domestic breeds have different recent demographic histories. For instance, the IB breed suffered a dramatic reduction of its effective population size during the last century (Alves et al. 2006), whereas commercial breeds such as Duroc or LW have been introgressed with Asian pigs (Bosse, Megens, Frantz, et al. 2014).

Differences in the effective population size, demographic histories and artificial selective pressures between pig populations could result in differences among their evolutionary rates. In addition to possible differences in the rate of evolution between populations, there may be differences in the rate of evolution between genes within genomes. For instance, it is known that the strength of the selection is affected by the position of the genes in the networks in which they participate. Genes that are more central in a network and are more connected with other genes are more evolutionarily constrained, while peripheral genes are more prone to be under adaptive selection (Fraser et al. 2002; Hahn and Kern 2005; Montanucci et al. 2011; Alvarez-Ponce and Fares 2012). Furthermore,

it has been observed that the evolutionary rate, within a metabolic pathway, increases as we move

downstream, possibly because upstream genes are more pleiotropic, since they are involved in

more functions and hence these genes are probably more conserved (Rausher, Miller, and Tiffin

1999; Riley, Jin, and Gibson 2003; Livingstone and Anderson 2009; Ramsay, Rieseberg, and

Ritland 2009).

So far, the nature of the underlying genetic changes caused by domestication and ensuing artificial

breeding is still under debate. While the most prevalent view is that regulatory changes have been

targeted (Anderson 2013), several other studies underline the influence of protein coding changes

(Rubin et al. 2012). Other authors have reported an increase in the rate of deleterious mutations in

domestic pigs compared to their wild counterparts (Cruz, Vilà, and Webster 2008; Renaut and

Rieseberg 2015; Pérez-Enciso et al. 2017; Leno-Colorado et al. 2017). Others, as in Makino et al.

(2018) detected a general pattern of reduction of variability in domestic populations in relation to

their wild counterpart, and a higher nonsynonymous/synonymous ratio across the frequency

spectrum. These patterns were compatible with the effect of strong bottlenecks in domestic

populations and the higher accumulation of deleterious mutations. Interestingly, the same authors

observed that the opposite trend has been observed in pigs. Moreover, most of the previous studies

have focused on genes of major effect with clear signals of selective sweeps. In those studies, the

hallmarks of positive selection were detected by a valley of reduced variation and/or population

differentiation that spans a relatively large region (e.g., Amaral et al. 2011, Rubin et al. 2012,

Frantz et al. 2013, Wilkinsonet al. 2013) but also by haplotype structure and homozygosity blocks

(e.g., Fang et al. 2011, Bosse et al. 2012, Li et al. 2013). Some of these studies have detected recent

breed specific signals of selection attributed to the domestication process (Li et al. 2014, Kim et

al. 2015). Nevertheless, the signals were too scarce to explain the domestication process. Other

studies have tried to elucidate the effect that domestication has at a whole-genome scale and on

the fitness of individuals of domestic populations (e.g., Cruz et al. 2008, MacEachern et al. 2009,

Kono et al. 2016, Perez-Enciso et al. 2016, Makino et al. 2018, Chen et al. 2018, Orlando and

Librado 2019). For instance, an excess of deleterious variants has been observed in a number of

domestic animal and plants (e.g., contrasting nonsynonymous versus synonymous polymorphism

ratios, Chen et al. 2018, using the MacDonald framework, MacEachern et al. 2009, contrasting

ancestors with ancient DNA, Orlando and Librado 2019, combining the frequency of

polymorphisms with functional effects and divergence, Kono et al. 2016, Makino et al. 2018).

Kono et al. (2016) and Perez-Enciso et al. (2016) found an excess of detrimental variants affecting

phenotypes of interest, suggesting, as we previously mention above, that protein sequence may

have a stronger influence than regulatory changes in the domestication process. Kono et al. (2016)

also showed that null alleles are uncommon in domestic animal species (also reviewed by

Anderson 2013), suggesting that phenotypic changes involved in domestication are produced by

the accumulation of consecutive mutations that modify these functions under selection. Finally,

the possible presence of beneficial mutations during the domestication process has also been

reported (Perez-Enciso 2016).

Here, we are interested in determining the proportion and the selective effects of protein-coding

variants in wild and domestic pig genomes to understand their role in the domestication process.

Particularly, we aimed to test the role of both new and extant mutations in the domestication

process and whether the phenotypes associated with domestic breeds are the product of a large

number of variants with weak selective effects, as suggested by previous results. To achieve this,

we have investigated the differential effects of selection on coding sequences at the different

molecular scales (gene, metabolic pathway and whole-genome) in two domestic and one wild pig

population using the McDonald-Kreitman framework (McDonald and Kreitman 1991, Eyre-

Walker 2006, Fay 2011) and have inferred the distribution of fitness effects (DFE) while taking

into account the effect of different demographic scenarios. Interestingly, the analysis was

performed using variability estimators that allow including positions with missing data (Ferretti,

Raineri, and Ramos-Onsins 2012).

Our results support the hypothesis that changes in allele frequencies in coding variants with weak

positive selective effect have been relevant for pig domestication, as evidenced by a relatively high

number of nonsynonymous variants segregating at medium and high frequencies and by the

obtained estimates of the Distribution of Fitness Effects in domestic pig populations.

MATERIALS AND METHODS

Biological samples

We analyzed a sample of 46 pig (*Sus scrofa*) genomes (Table S1). These pigs correspond to European wild boars (WB, n = 20) and domestic pigs, which are represented by the Iberian Guadyerbas (IB, n = 6) and LargeWhite (LW, n = 20) breeds. These two domestic breeds were selected because they have very different interesting features: IB is a local breed that has been under weak artificial selection intensity and with no documented evidence of Asian introgression. LW, in contrast, is a commercial breed undergoing strong artificial selection with a deliberate admixture with Asian pigs (Bosse, Megens, Madsen, et al. 2014; Groenen 2016). To analyze the divergence between the different breeds, we used the consensus ancestral reference sequence obtained from combining the information from several *Sus* species (*S. barbatus*, *S. cebifrons*, *S. verrucosus*, *S. celebensi*, approximately 4.2 MYA of divergence) and the African warthog (*Phacochoerus africanus*, around ~10 MYA of divergence) as an outgroup, as detailed in Bianco et al. (2015). The sequences are available in public databases (Rubin et al. 2012; Ramírez *et al.* 2014; Bianco *et al.* 2015; Frantz et al. 2015; Moon *et al.* 2015, Esteve-Codina *et al.* 2013, Leno *et al.* 2017) and were downloaded from the short read archive (SRA, http://www.ncbi.nlm.nih.gov/sra).

Mapping and genotyping analysis

Raw reads for each pig genome were mapped against the reference genome assembly (Sscrofa10.2, Groenen et al. 2012) using *BWA mem* option (H. Li and Durbin 2009). PCR duplicates were removed using *SAMtools rmdup* v 0.1.19 (H. Li et al. 2009) and mapped reads were realigned around indels with the *GATK IndelRealigner* tool (McKenna et al. 2010). Genotype calling was performed with *SAMtools mpileup* and *bcftools call* v 1.3.0 (H. Li et al. 2009) for each individual separately. We set a minimum (5x) and a maximum depth (twice the average sample's depth plus one) to call a SNP. Base quality was set to 20 (*P*-value=1e-2). Homozygous blocks (regions of contiguous positions with the same nucleotide as the reference genome) were also called, following the same criteria (i.e., minimum and maximum coverage and base quality) as with the SNPs and using *samtools depth* utility, *BEDtools* (Quinlan 2014) and custom scripts (available at https://github.com/miguelperezenciso/NGSpipeline, Pérez-Enciso *et al*., 2017). This resulted in a

gVCF file per individual with the combined information about variant calls and non-varying
positions. Next, each *gVCF* file was converted into a fasta file and all fasta files were subsequently
merged to obtain a multindividual *gVCF* file (Pérez-Enciso et al. 2017), which comprised the
whole set of the SNPs of the 46 pigs.

Analysis of the population structure of the samples

A principal component analysis (PCA) was performed using the total number of SNPs to analyze
the population structure. First, genotypes were converted to alternative allele frequency, being 0
for the homozygous reference genotype (0/0), 0.5 for the heterozygous genotype (0/1) and 1 for
the homozygous alternative genotype (1/1). For cases of missing genotype (./.), these were
replaced by the average SNP frequency across all individuals. We used the function *tcrossprod()*
from R version 3.3.1 (2016) to obtain the matrix of covariates from the frequencies matrix. Finally,
we obtained the principal components from the Eigen-value decomposition with the R function
eigen().

Estimation of levels and patterns of variability

Genetic diversity and divergence per pig population were estimated using *mstatspop* software
(Nevado, Ramos-Onsins, and Perez-Enciso 2014; Bianco et al. 2015; Guirao-Rico et al. 2018,
available from the authors, https://github.com/cragenomica/mstatspop). The *multi-VCF* file was
converted into a *tfasta* (transposed *fasta*) file and *mstatspop* was run on either the whole genome,
using 5 Mb windows and at each functional coding region. We used four different estimators of
nucleotide variability that takes into account missing data (Ferretti, Raineri, and Ramos-Onsins
2012): Watterson (Watterson 1975), Tajima (Tajima 1983), Fu&Li (Fu and Li 1993) and
Fay&Wu's estimators (Fay and Wu 2000). Specifically, variability was estimated using the
Ferretti, Raineri, and Ramos-Onsins (2012) expression:

$$\hat{\theta} = \frac{1}{L} \sum_{x=1}^{L} \sum_{i=1}^{n_x-1} i\omega_{i,n_x} \xi_i(x), \qquad \frac{1}{L} \sum_{x=1}^{L} \sum_{i=1}^{n_x-1} \omega_{i,n_x} = 1$$

184   *(Equation 1)*

185   where (ω_i) is the weight for the different variability estimators such as $\omega_i = n/(i(n-i)(1+\partial_{i,n-i}))$ for

186   the Watterson estimator, $\omega_i = n/(1+\partial_{i,n-i})$ for the Tajima estimator (both for folded spectrum), $\omega_i =$

187   i for the Fay&Wu estimator and $\omega_i = 1$, $\omega_{i>1} = 0$ for the Fu&Li estimator (Achaz 2009).

Filtering for artifactual effects

A preliminary analysis of the variability showed a moderate negative correlation (~ 0.3) between

the levels of variability and divergence and the proportion of missing data for each gene. To

eliminate this artifactual correlation, we plotted the estimators of variability and divergence versus

the ratio of missing data and eliminated those genes that showed a ratio of missing data greater

than 0.3. Since this filtering was not enough to completely remove the bias, we also removed genes

with extreme values of variability and divergence (higher than 99% quantile of the total genes).

The remaining $\sim$13,500 genes (70% of the total annotated genes) showed a low or null correlation

with missing data and were used in the present analysis (Table S2).

Estimation of the proportion of adaptive of variants

Under the neutral model, the majority of polymorphisms segregating in a population are neutral

and only a small number of positively selected variants segregates for a short time on their way to

loss or fixation. Hence, most of the positive selected variants are only observed as fixed variants.

In addition, functional positions (nonsynonymous positions) are constrained compared to

nonfunctional positions (synonymous positions), and hence their evolutionary ratios are smaller.

In the neutral scenario, polymorphism and divergence (excluding the adaptive fixed variants) are

proportional to the mutation rate and to the constriction factor in the case of nonsynonymous

positions (McDonald and Kreitman 1991, Eyre-Walker 2006, Fay 2011). That is:

$$\frac{\theta_n}{\theta_s} = \frac{(1-\alpha)K_n}{K_s},$$

209   *(Equation 2)*

210 where θ_n the nonsynonymous variability, θ_s is the synonymous variability, K_n the nonsynonymous

211 divergence, K_s the synonymous divergence and α is the proportion of adaptive variants that have

212 been fixed. To estimate the proportion of nonsynonymous substitutions that are adaptive (α) we

213 reorder the previous expression:

214
$$\alpha = 1 - \frac{K_s}{K_n}\frac{\theta_n}{\theta_s}$$

(Equation 3)

A higher ratio of nonsynonymous to synonymous divergence versus polymorphisms suggests that

positive selection has fixed adaptive variants ($\alpha > 0$) and the opposite case ($\alpha < 0$) suggests the

presence of deleterious mutations segregating in the population.

If we consider that weak deleterious mutations are segregating in the population, we expect that

their relative proportion will be higher at lower frequency variants and low or null for fixed

deleterious mutations. Following the same notation as in equation 3:

222
$$\frac{\theta_{in}(1 - \beta i)}{\theta_{is}} = \frac{(1 - \alpha - \beta d)K_n}{K_s},$$

223 *(Equation 4)*

224 where i refers to the frequency at which the calculation of variability is estimated, βi is the

225 proportion of weakly deleterious polymorphic mutations at frequency i, βd is the proportion of

226 weakly deleterious fixed mutations. $\beta d < \beta i$ was assumed at any frequency. Then, solving for the

227 proportion of fixed adaptive variants (α):

228
$$\alpha = 1 - \beta d - (1 - \beta i)\frac{K_s}{K_n}\frac{\theta_{in}}{\theta_{is}}$$

(Equation 5)

We see that in case calculating α without considering the effects of deleterious mutations, it would

be underestimated depending on the frequency at which the estimates of variability are calculated.

If we assume that the detrimental variants would never be fixed, a good estimator of α using

equation 3 would be the one that estimates variability based on high frequencies, as it would hardly

contain detrimental mutations. This is in agreement with the arguments used in Messer and Petrov

2013.

Similarly, if we also consider that weak positively selected variants are segregating in the
population, we expect that their relative proportion, compared to neutral ones, is higher at higher
frequencies:

$$\frac{\theta_{in}(1 - \beta i - \gamma i)}{\theta_{is}} = \frac{(1 - \alpha - \beta \, \mathrm{d} - \gamma d)K_n}{K_s},$$

(Equation 6)

242 where γi is the proportion of weakly advantageous polymorphic mutations at frequency *i*, and
243 γd is the proportion of weakly advantageous fixed mutations. Again, solving for the proportion of
244 fixed adaptive variants (α):

$$\alpha + \gamma d = 1 - \beta d - (1 - \beta i - \gamma i)\frac{K_s}{K_n}\frac{\theta_{in}}{\theta_{is}}$$

(Equation 7)

In this case, the presence of adaptive variants segregating in the population would affect the
estimates of variability based on high frequency variants when using equation 3, which would
result in an underestimation of the proportion of fixed adaptive variants (α). Note that adaptive
variants stabilized at intermediate frequencies are not considered in this approach, which can be
an important source of adaptation considering the infinitesimal model.

If we focus on the effects on the polymorphisms, equation 6 suggests that the ratio of
nonsynonymous to synonymous polymorphisms would increase due to mutations having both
positive and negative effects. It is expected that the number of mutations with negative selection
coefficients would rapidly decrease as we move to intermediate and high frequencies, while the
opposite trend is expected for mutations with positive selection coefficients. Hence, higher ratios
of nonsynonymous to synonymous polymorphisms at higher frequencies may be explained by the
presence of advantageous mutations segregating in the population.

Furthermore, in cases where two populations are from the same species and no fixed mutations
between them we can estimate the possible differential effect of the selection (positive and
negative) at any frequency among populations from the ratios of synonymous to nonsynonymous
polymorphisms of the two populations ($R\_\beta\gamma i$):

265
$$\frac{\theta_{in1}(1 - \beta i1 - \gamma i1)}{\theta_{is1}} = \frac{\theta_{in2}(1 - \beta i2 - \gamma i2)}{\theta_{is2}}$$

266 and

267
$$\frac{(1 - \beta i1 - \gamma i1)}{(1 - \beta i2 - \gamma i2)} = \frac{\theta_{is1}\theta_{in2}}{\theta_{in1}\theta_{is2}} = R\_\beta\gamma i$$

(Equation 8)

In addition, a comparison of the $R\_\beta\gamma i$ values calculated using different variability estimators
(hereafter $R\_\beta\gamma i$ pattern) can be used to inform about the effects of the different types of selection.
Importantly, different demographic effects (e.g., bottlenecks) together with the presence of
mutations with small selective effects may also disturb the ratios of variability and hence must be
taken into account when interpreting the results.

The effect of linkage disequilibrium between selective (detrimental or adaptive) and neutral
variants should not affect the expected estimate of the proportion of adaptive variants, as it would
affect both synonymous and nonsynonymous positions in the same proportion. On the contrary,
the interaction of variants with opposite selective effects would possibly reduce the effect of
selection and would have a significant consequence on the estimation of adaptive fixed variants
(Hill and Robertson 1966; Booker and Keightley 2018).

Bootstrap analysis

Nonparametric bootstrap analysis was performed to estimate the null distribution of the α statistic
for each variability estimator and pig population. In each case, coding positions for synonymous
and nonsynonymous (separately) were randomly chosen with replacement and the α statistic was
calculated as in equation 2. This process was repeated 100 times.

Simulations

We carried out forward simulations using the software *SLiM* (Haller and Messer 2017) in order to

assess the interaction of the different selective effects and demographic factors affecting the

evolution of pig populations during domestication. We explored the expected values of nucleotide

diversity, divergence, α and $R\_\beta\gamma$ under 63 different scenarios. For each scenario, we simulated

three populations corresponding to wild, domestic and an outgroup species. We first simulated

nine different scenarios that were classified into three main groups: i) standard neutral model

(SNM); ii) a model with negative selection (NS) and iii) a model with positive selection (PS). For

the models with selection, we let that selection operate from the ancestral species to the present

time. Each group of scenarios (SNM, NS and PS) was simulated with a constant effective

population size for the three populations or with a reduction or an expansion of the effective

population size in the branch leading to domestic pigs. A second group of simulations was

performed under more complex scenarios. In those simulations, we incorporated the combined

effect of negative and positive selective effects (using gamma and exponential distributions for the

selective coefficients, respectively) plus demographic effects such as expansion and reduction of

the effective population size in the domestic simulated populations and with or without migration

from the wild into the domestic populations (in total 54 complex simulated scenarios). Figure S1

shows a general scheme for the simulated populations and Table S3A-B shows the parameter

values used in these simulations. The obtained results were analyzed using the *mstatspop* software.

307   Approximate Bayesian computation (ABC) analysis

We used the ratio of the estimates of nucleotide variability ($\theta n/\theta s$) per nucleotide for

nonsynonymous versus synonymous positions (Fu&Li, Watterson, Tajima and Fay&Wu) and of

divergence (*Kn/Ks*) to infer the distribution of fitness effects (DFE) in coding regions. We

compared three evolutionary models that differ in the shape of the DFE using the algorithm

proposed by Tataru et al. (2017), which are the following: (i) model A: a model with a deleterious

gamma DFE with the mean and the shape of the gamma distribution as model parameters, (ii)

model C: a model with a gamma distribution of deleterious variants with two parameters (shape

and mean) and an exponential distribution of beneficial variants with one parameter (mean), and

the additional parameter of the proportion of beneficial versus deleterious variants, and (iii) model

D: a model with a discrete distribution of a priori values of possible selective coefficients (positive

and negative) and considering as parameters the proportion of each (positively and negatively
selected mutations). Some of the additional parameters, such as demographic or linkage effects,
were considered as nuisances, while others, such as errors in the polarity of unfolded mutations,
were fixed. Table S4 shows the parameters and the prior distributions used in the analysis. We
used *polyDFEv2* (Tataru et al 2019) to obtain the expected unfolded site-frequency spectrum
(SFS). With the aim of performing the ABC using summary statistics, the code of *polyDFEv2* was
slightly modified in order to print the SFS and the parameters for a large number of conditions.
For each model, one million iterations were run using different parameter conditions and the
resulting SFS for each condition were kept to later calculate the ratios of variability, divergence
and the α statistic. ABC analysis was performed using the R library *abc* (Csillery et al. 2012). We
performed a cross validation analysis to evaluate the ability of the approach to distinguish between
models using the *cv4postpr()* function, as suggested in the *abc* library documentation. The
confusion matrix indicated that these three models were quite distinguishable with a probability of
true classification from model A versus C/D of 0.70, from model C versus A/D of 0.60 and from
model D versus A/C of 0.67, with a tolerance of 0.05 (Table S5). Posterior probabilities of each
model given the observed data were obtained using the *postpr()* function and considering a
multinomial logistic and a rejection approach. Additionally, a goodness of fit analysis, which
compares the median of the distance between the accepted summary statistics and the observed
ones, was also performed to select the best model. Once the best model was chosen, the ability to
infer the parameters of the model was assessed using the *cv4abc()* function. Prediction errors for
the parameter inference of each model are shown in Table S6 and Figure S2. The parameters of
the best model were inferred with the *abc()* function using a local linear regression and a rejection
approach. Posterior predictive simulations were performed with the α statistic (instead of with the
ratios of variability and of divergence, to avoid circularity in the analysis) to determine whether
the simulated data generated from the estimated parameter of our best model resembled the
observed data (1000 replicates). Finally, the α values can be simply estimated using equation 10
from Tataru et al. (2017), which is a simple proportion of positive selective coefficients (*s*) values
in the case of the discrete distribution.

Pathway analysis

We downloaded the complete list of pathways and genes of *Sus scrofa* from KEGG v.20170213

(http://www.genome.jp/kegg/, Kanehisa et al. 2008). The list contained 471 pathways and 5480

genes. The median and mean number of genes per pathway was 26 and 43, respectively, and ranged

from 1 to 949. We filtered the pathways according to their size, removing pathways with less than

10 and more than 150 genes in order to discard pathways that were not informative or too generic

and complex. The final list contained 171 pathways and 3449 genes.

To analyze the selection pressure of each gene according to its position in the pathway, we obtained

different topological parameters. For that, we first downloaded the XML file of each pathway from

KEGG v.20170213. These files were analyzed with the *iGraph* R package (Csardi G. and Nepusz

T. 2006) to obtain the topological descriptors of each gene in each pathway. For each gene, three

different measures were computed: *betweenness* (number of shortest paths going through a vertex),

in-degree (number of in-going edges) and *out-degree* (number of out-going edges). These

parameters are measures of the importance of a gene within a pathway: *betweenness* is a centrality

feature, *in-degree* suggests the facility of a protein to be regulated and *out-degree* reflects the

regulatory role of a protein. We tested whether negatively and positively selected genes differed

in any of these statistics using a nonparametric Wilcoxon rank test, due to the extreme leptokurtic

distributions involved.

RESULTS

Predominance of shared variants and similar selective effects on pig populations

We found a total of 6,684,142 SNPs in autosomes, with 149,440 SNPs of these located in coding regions. We found that 12.5% of the SNPs in the coding regions are shared among the three populations, 32.2% are shared between at least two populations, 31.2% are exclusive to LW, 2.2% are exclusive to IB and 34.4% are exclusive to WB (Table 1). Based on the PCA analysis and using the total number of SNPs, we found that the individuals of each breed cluster together and are well separated from other breeds (Figure S3). The proportion of private SNPs in each population is in accordance with its specific demographic history (Esteve-Codina et al. 2013, Bosse, Megens, Madsen, et al. 2014; Bosse, Megens, Frantz, et al. 2014).

For each breed, each coding position was classified as polymorphic, fixed (i.e., different alleles from the outgroup) or ancestral allele (i.e., same allele as in the outgroup), with the aim of identifying those variants that appeared previously or posteriorly to the domestication process (Table 2). Surprisingly, we found very few fixed mutations between populations, indicating that the phenotypic traits of each population are not associated with fixed coding variants. Similarly, we found very few fixed coding variants in domestic (IB or LW) versus wild (WB). There are few variants fixed in the domestic that are polymorphic in the wild population, suggesting that these variants were previously present in wild breeds or, alternatively, were transferred into WB by gene flow from introgressed domestic breeds. Most of the variants that are exclusive of a single breed are polymorphic, which is in agreement with the recent origin of these variants. We found a large number of fixed variants in the IB that are polymorphic in LW and WB, likely due to a reduction of the effective population size of the IB breed. The ratio of nonsynonymous to synonymous polymorphism was always lower than one and with similar values in the three populations regardless of the variability estimator used. This result suggests that apparently, there are no differential effects of selection between domestic and wild populations.

Limited influence of genomic context and the network topology on selective patterns

The heterogeneity in the recombination rate, the gene density, the %GC and the distribution of
CpG islands across the genome can affect the local levels of variability. A previous study on the
IB breed detected a strong correlation between recombination and variability, although no
correlation was observed between variability and gene density or GC content (Esteve-Codina et
al. 2013). However, the effect of these factors on the estimation of the proportion of adaptive
nonsynonymous mutations (α) has not been previously studied. We observed no correlation
between α and recombination, gene density, missing rate, %GC and CpG in any of the three breeds
(P-values > 0.01).

Next, we investigated the effect of gene network topology on the selective patterns. It has been
claimed that topology limits the 'evolvability' of genes and that highly connected genes are more
constrained and, consequently, less likely to be targets of positive selection. We compared the
network topology features (*betweenness*, *out-degree* and *in-degree*) of genes within pathways
regarding the value of α, grouping genes with positive versus negative α values. We found that
genes with negative α values show significant large values of the *betweenness* statistic in the three
pig breeds compared to genes with positive α values (*P*-value < 0.01; Figure S4). LW and WB
(but not IB, possibly because the low sample size) show significant values (*P*-values < 0.01) of the
in-degree statistic for genes with negative α values compared to genes with positive *a* values.
However, we did not observe significant differences in the *out-degree* values between genes with
negative and positive *a* values in any of the three breeds (Figure S4). These results suggest that, in
the three breeds, genes that are more central in a pathway are more evolutionary constrained
compared to peripheral genes. In addition, in LW and WB, the genes that are more constrained
tended to have a higher number of upstream genes that regulated them, which is also in agreement
with the central position of these genes in the pathway. We did not observe significant differences
in *in-degree* statistic in the IB breed between genes with negative and positive *a* values, perhaps
because of a relaxation of functional constraints as a consequence of the reduction of its effective
population size.

Levels of nucleotide variation at protein coding regions is compatible with the history of the
surveyed pig populations and with positive selection

To assess the selective effect of domestication, we first studied the pattern of variation at

synonymous and nonsynonymous positions using four estimators of variability that differentially

weight the SNP frequencies (See Material and Methods). Estimates of the levels of variability per

nucleotide at the genome level using different estimators are shown in Figure 1 and detailed in

Table S7. We expect that, under the Standard Neutral Model (SNM), all estimates of variability

should be similar while differences among them may indicate demographic and/or selective

effects. We observed that the levels of variability are different for each estimator within breeds

and also for the same estimator for different breeds. However, we observed a similar ratio of

nonsynonymous to synonymous polymorphisms for all estimators of variability, suggesting that

demographic effects are responsible for the differences in the levels of variability (Figure 1). The

less variable population is the IB breed, which shows far fewer singletons compared to WB and

LW, probably as a consequence of the reduction of its population size. Note than in all three

populations, high-frequency variants are proportionally more abundant than those at intermediate

frequencies, which would be compatible with the accepted demographic history of the surveyed

populations (i.e., introgression in the LW) but also with the presence of pervasive positive selection

in all three populations.

α's and $R\_\beta\gamma$ ratios might show a differential effect of selection due to domestication

The differential effect of selection in the domestic and wild populations can be studied by

comparing the α values. Figure 2 and Table S8 show the genome-wide α values calculated using

the four variability estimators for each population. As expected, the α values are negative when α

is calculated using the estimate of variability based on low-frequency variants ($\alpha_{Fu\&Li}$), probably

reflecting the relatively high proportion of deleterious versus neutral mutations that are segregating

at low frequencies. We observed a similar value of $\alpha_{Fu\&Li}$ in all populations, suggesting a similar

proportion of segregating deleterious mutations, irrespective of the domestication process or other

demographic events (Figure 2A). Moreover, we observed less negative values of α, or even

positive (for LW in Figure 2A), when α is calculated based on variants at high frequencies,

according to expectations that point to a progressive elimination of deleterious mutations at higher

frequencies. Nevertheless, the pattern of α (i.e., the comparative a values calculated using the four

different variability estimators within each population) is very different in each population. WB

and IB show very low negative values of α for most of the estimators of variability, except for

$\alpha_{Fay\&Wu}$ in WB, which is zero. LW is the only breed that shows a linear increase of the α negative

values across the SFS, being even positive when calculated based on high frequencies ($\alpha_{Fay\&Wu}$).

We did not observe similar patterns of α between domestic breeds compared to WB (Figure 2A).

The differences in the ratio of synonymous to nonsynonymous variability between the two

different breeds is summarized by the $R\_\beta\gamma$ ratio (Figure 3A). We observed deviations from $R\_\beta\gamma$

= 1 when the ratio was calculated based on high-frequency variants ($\alpha_{Fay\&Wu}$). Although the ratio

of the two populations is difficult to interpret because of their different underlying demographics,

some trends can be observed. WB shows an excess of nonsynonymous variants segregating at

intermediate frequencies (WB-IB, WB-LW), which might be explained by a past bottleneck that

increased deleterious mutations at intermediate frequencies. In addition, the $R\_\beta\gamma$ ratio in LW-IB

shows an incremental pattern from low to high frequencies, which is compatible with an increase

of nonsynonymous beneficial variants on their way to fixation in LW.

α's and $R\_\beta\gamma$ ratios based on from exclusive and shared polymorphisms might reflect

changes in selection patterns before and after domestication

We observed a high ratio of nonsynonymous to synonymous exclusive singletons ($\alpha_{Fu\&Li}$, Figure

2B), suggesting that they have deleterious effects in all populations. Nevertheless, the values of α

calculated based on intermediate frequency variants (α_{Tajima}) in the WB and IB populations are

lower than to those based on low-frequency variants, which point to the action of positive selection

maintaining nonsynonymous variants at higher frequencies in the case of WB and to an attenuated

effect of deleterious mutations in IB due to a population size decline. In the case of LW, the

observed pattern of α is similar to that calculated with all SNPs (Figure 2B). Likewise, the $R\_\beta\gamma$

statistic shows the same pattern as that calculated using all SNPs but with a higher magnitude of

its value (Figure 3B).

On the other hand, the α values based on shared variants are in general more moderate (closer to

zero) than those based on exclusive or all SNPs (Figure 2C), likely because shared nonsynonymous

polymorphisms are older and hence, expected to be more functionally constrained than exclusive

ones. Additionally, the values of α based on singletons ($\alpha_{Fu\&Li}$) are less negative than those based

on intermediate-frequency variants. Again, this pattern might indicate that selection is involve in

the increase of the ratio nonsynonymous to synonymous polymorphisms up to intermediate

frequencies. The $R\_\beta\gamma$ statistic shows similar patterns than those observed for all variants but with

values much closer to 1, indicating a small or moderate selective effect on the shared variants

compared to all variants (Figure 3C).

When we calculated the α values from shared variants only between the two domestic breeds, we

found an inverse pattern regarding that calculated from all SNPs in each population, with high

positive values of α based on low frequencies and very negative values when α is calculated based

on high-frequency variants (Figure 2D). Some possible explanations might be the active

elimination of new nonsynonymous variants to preserve differences among breeds ($\alpha_{Fu\&Li}$) and

either the effect of the ancestral population structure (wild versus domestic), or the presence of

nonsynonymous variants targeted by the process of domestication that shifts them toward high

frequencies ($\alpha_{Fay\&wu}$).

Absolute values of α are dependent of the molecular scale but patterns remain similar

In addition to the genome-wide analysis, α was estimated using three additional molecular scale

levels: i) gene level, ii) genes within windows of 5 Mb, and iii) genes within the same pathway.

Figure 4 shows the median of the distributions of the α values for each scale level. In general, the

distribution of α values estimated at the genome-wide level are concordant with those estimated at

the gene level, genes within windows and pathway level in each breed. However, differences in

the value of α within each breed are notorious depending on the scale level examined. The median

estimates of α are generally lower at the gene scale level and most of them are very negative, while

at the genome-wide scale, the α values are closer to zero. However, the distribution of α values

can have a large variance at the gene scale since few variants are used for its estimation. We

identified the regions and pathways that showed extreme α values (Table S9 and S10). We found

a large number of genes having $\alpha = 1$ (highest value) because the number of polymorphic

nonsynonymous variants per gene was zero. We also found a moderately high correlation of α

values between breeds (rho ~ 0.7, Pearson correlation using pathways) suggesting that in general,

these breeds are under similar selective effects. For shared and exclusive variants, we generally

observed the same pattern, from genes to whole-genome, that is, larger α values at the gene level

and closer to zero α values at the larger scale. Only shared variants of the IB breed exhibited similar

α values from genes to the whole-genome scale. The differences in absolute α values could be

explained because of the nature of this ratio statistic, in which the mean can be more displaced to

more negative values due to a reduced number of functional variants (*i.e.,* few or null segregating

nonsynonymous variants).

522   Simulated data under scenarios that include the joint effect of demography and selective

523   events were more similar to the observed data

We used computer simulations to study how the different demographic and selective events

occurred during domestication affected nucleotide variation. We simulated populations mimicking

the process of domestication using *SLiM* software (Haller and Messer 2017) coupled with several

demographic events, including changes of the population size and/or migration. We analyzed the

genome-wide patterns of α and the $R\_\beta\gamma$ statistic produced by 63 simulated scenarios that included

different demographic events and selective forces acting separately (simple scenarios) or jointly

(complex scenarios). The results of the simulation study are summarized in Figures S5-S46. The

observed patterns of α calculated from all variants in the surveyed populations are not compatible

with simple scenarios that only consider demographic or positive selection forces (Figure S5).

Rather, simulated α trends (irrespective of the magnitude of α) fit a scenario with a predominant

effect of negative selection (Figure S5). However, the $R\_\beta\gamma$ statistic do not fit any of the simulated

simple scenarios (Figure S6). When more complex scenarios were considered, the general trends

of the patterns of α generated by scenarios with both negative and positive selection resembled

those observed in WB and LW (Figures S7-S13). Notice that this is true only for those scenarios

that include some demographic or migration events (Figures S10-S12). On the other hand, the IB

population would fit a scenario without positive selection and with a recent population size

reduction (Figure S11). The trends in the $R\_\beta\gamma$ statistic are, in broad strokes, concordant with the

conclusions extracted from the comparison between the observed and simulated patterns of α

(Figures S13-S18).

The observed patterns of α values from exclusive variants are different from those based on the total variants in WB and IB populations (Figure S19). These patterns cannot be fully explained by any of the complex simulated scenarios proposed when considering all variants, except for the IB population, which might be compatible with a scenario with negative selection and population size reduction (Figures S22-S26). The observed $R\_\beta\gamma$ are also compatible with the scenarios combining both types of selection and population size reduction (WB-IB) or with scenarios that include migration (WB-LW; Figures S27-S32). Finally, the observed patterns of α and $R\_\beta\gamma$ statistics calculated from shared variants are also compatible with scenarios having deleterious plus beneficial mutations (Figures S33-S46).

A model that assumes a discrete distribution of beneficial and deleterious mutations would fit better the observed data in the ABC analysis

We used an approximate Bayesian computation (ABC) analysis to infer the DFE separately for each population using the ratios of nonsynonymous to synonymous variants obtained from the whole-genome analysis (see Materials and Methods). Three different models implemented in *polyDFE2* software (Tataru et al. 2019) were tested. Model A, which assumes a gamma distribution of deleterious mutations; model C, which assumes a gamma distribution for deleterious mutations and an exponential distribution for beneficial mutations; and model D, that assumes a discrete distribution of beneficial and deleterious mutations. In these models, we included demographic and linkage effects as nuisance parameters (Tataru et al. 2017). Goodness of fit (GoF) analysis, which is a measure of the adjustment of the prior chosen models to the data, revealed that the simulated data under the different models do not always fit well to the observed data (Table S11A), suggesting that the real data fit only to a very restrictive parameter range of each model. This is especially pronounced for exclusive variants, where the simulations under the different models fit only marginally to the observed data. Indeed, the model with a wider parameter range versus the real data is usually the discrete model D. However, if we also take into account the values of the posterior probabilities, which is the probability assigned to each model relative to the other models of the analysis, we found that the best fit model differed among populations (model C to WB and LW and model D to the IB breed; Table S11A). Finally, posterior predictive

analysis indicated that the observed α values (Figure 5) and variability

nonsynonymous/synonymous ratios (Figure S47) cannot be obtained using the estimated

parameters of any model, although they were closer to model D. The parameters of the DFE

inferred for each population are shown in Table S12. The obtained results indicate that the DFE is

quite similar among all three populations, which is not entirely surprising because they share a

long-term history. Despite there is a lot of uncertainty in the inferred estimates, the obtained results

show that the DFE contains a large fraction of very deleterious variants, with approximately 83%

of the variants being strongly deleterious (S ≤ -200; considering the posterior distribution and using

the rejection method), and with approximately 17% of the variants being neutral, weak beneficial

and weak deleterious mutations ($-2 \leq S \leq +2$). The median proportion of adaptive mutations (α)

estimated from the discrete distribution (by summing weak and strong beneficial proportion of

mutations in Model C) is approximately 0.9% (Table S12A).

586   The inferred DFE is different when based on exclusive and shared variants

Although the inference of the DFE is going to be distorted by choosing only a subsection of the

variants (e.g., exclusive variants are mostly very recent), we considered that it can give some clues

about past events that could be more related to the domestication processes. Regarding to exclusive

variants, the simulations under the three different models show a low fit to the observed data.

Indeed, in some cases, the GoF is less than 1% (Table S11B). Although the posterior probability

is higher for model C, the posterior predictive simulations show that none of the models can

reproduce well the observed data. In general, the posterior predictive simulations under model D

yield more similar values to the observed data. However, only the posterior predictive simulations

under this model are reasonably similar to the observed data for the LW breed (Figure 6, Figure

S48). The results obtained for shared variants are very similar to those considering the total

variants, supporting the hypothesis that shared mutations have a predominant effect (compared to

exclusive variants) at the whole genome. Finally, the median proportion of adaptive variants (α)

estimated from the discrete distribution (Model C) gave similar results to the total mutations, but

surprisingly estimated a slightly higher proportion for the Iberian breed (Table S12C; Figure S49-

S50).

603
DISCUSSION

The polygenic nature of the domestication traits in animals often precludes identifying their
underlying genes since the domestic phenotypes are caused by subtle allele-frequency changes of
variants distributed throughout the genome that are very difficult to detect. In addition, the study
of the effects of selection using genome sequences that contain a nonnegligible fraction of missing
data is challenging and needs the use of appropriate methods to account for these positions.
Statistics that exploit the frequency of the variants while accounting for missing data are
particularly appropriate for such analyses.

Here, we provide a novel approach that combines the use of different estimators of variability that
account for missing data with the asymptotic approach proposed by Messer and Petrov (2013) and
Uricchio et al. 2019). This approach allowed us to study and to interpret the effects of the
domestication process on genomic variation as an alternative to the use of the full SFS. Although
it can be less precise (we used only four statistics to capture the entire trend of α across the SFS),
it helps to reduce the variance and facilitates the visual interpretation. The analyses performed here
include an exhaustive comparative study of the observed patterns of functional versus neutral
diversity in domestic pigs and wild boars, a forward simulation study for several diverse
evolutionary scenarios and the inference of DFE parameters given different selective models.

Note that the DFE was inferred using Bayesian calculations (ABC) instead of exact Bayesian or
Likelihood methods. Despite ABC requires additional steps and validation analysis and is in
general less precise, it allows contrasting models and inferring parameters from complex datasets
or data containing missing information (Beaumont et al. 2002).

Selection pressure on pigs and the process of domestication

A number of analyses with the aim to explain the process of domestication have been performed
already, using the MacDonald and Kreitman extension methods, or using other estimates such as
variability or divergence at functional or synonymous positions (MacEachern et al. 2009, Kono et
al. 2016, Makino et al. 2018). For instance, Kono et al. (2016) analyzed derived frequencies in

domesticated barley populations at different functional classes (deleterious, tolerated) and

observed a higher quantity of deleterious variants at low frequencies compared to tolerated or to

synonymous mutations. Makino et al. (2018) investigated the ratio of functional to neutral variants

at different frequencies in domestic and in wild populations of several animal and plant species,

including Asian and European pigs. They observed generally lower levels of synonymous and

nonsynonymous variation and a higher ratio of functional to neutral variants in the domestic

species compared with their wild counterparts. This ratio was negatively correlated with the

frequency of the variants, consistent with a higher number of detrimental variants in domestic

populations. The authors claimed that these patterns were compatible with the expected effect of

a bottleneck as a consequence of domestication, and with an increase in nonsynonymous variants

produced by the lesser efficacy of purifying selection at smaller population sizes, although the

presence of positive selection (hitchhiking) was not discarded. However, the opposite pattern was

observed in European wild boars and domestic pigs (a higher variability and a lower ratio of

nonsynonymous to synonymous in domestic pig populations compared to wild boars). The authors

argue that this can be explained by the highly variable patterns produced by bottlenecks, the strong

population contraction of European wild boars during the last glaciation and the presence of gene

flow between wild and domestic populations.

As in Makino et al. (2018), we do not observe an increase in functional diversity in domestic versus

wild populations. This result may also be explained by several recent events occurring in these

populations: (i) differences in the recent history of local and commercial domesticated populations,

that is, high inbreeding in Iberian local pigs but recent gene flow of the commercial pigs with

Asian pigs; (ii) demographic effects in the wild boar population that may have reduced their

diversity (Groenen et al 2016) or increase the variance of the patterns (see simulations, Figures

S5-S46); (iii) differential adaptive forces in local versus commercial pigs, with a recent high

selective pressure in this last population.

The two domestic breeds analyzed here have very different recent histories: the IB is a local

Spanish breed (Guadyerbas) that suffered a strong bottleneck during the 1970s (Esteve-Codina et

al. 2013) and with no evidence of introgression whereas the LW breed was admixed with pigs of

Asian origin (Bosse, Megens, Madsen, et al. 2014). Currently, approximately 20–35% of the LW

genome has been estimated to be of Asian origin (Groenen et al. 2012, Bosse et al. 2012, Frantz et

al. 2015, Bianco et al. 2015, Ai et al. 2015). Accordingly, the IB breed shows the lowest levels of

synonymous and nonsynonymous variation among the breeds studied, probably because of its

small effective population size and because the individuals from the IB sample come from a very

closed population of pigs. However, we expected to find a higher variability in LW compared to

WB due to the process of Asian introgression that this breed has undergone. Surprisingly, we

detected very similar levels of variability between them. However, the high levels of variability

were observed for variants that belonged to different frequency ranges in these two populations:

singletons in WB and in high-frequency derived alleles in LW. This difference may be due mainly

to the effects of gene flow in LW but also in some extend to the selective programs applied to the

commercial breed.

Domestication hallmarks at pig coding regions

The paucity of fixed variants at coding positions in the three breeds indicates that the observed

heritable phenotypic differences among the breeds are either due to: i) few selective sweeps, ii)

positive selection at noncoding functional regions which were not analyzed in this work, iii)

changes in the frequencies of nonsynonymous variants without being fixed. In the first hypothesis,

we expect that domestication process should fix the adaptive variants for those genes underlaying

the phenotypes of interest. However, we found no fixed variants between domestic breeds and

wild pigs, although they show the right breed phenotype. We checked the α values for those genes

that were previously reported to show signals of positive selection using other approaches

(Groenen 2016). We found that these genes show little or no nonsynonymous polymorphisms or

fixed variants (Table S13). This absence of variability is typical from regions under selective

sweeps, although not necessarily implicating that these genes are the targets of domestication since

there are no variants fixed or close to fixation at their coding regions. We only found significant

values of α over zero at gene KIT in the IB breed, genes IGF2R and JMJD1C in the LW breed and

gene LRRTM3 in the WB population. The second hypothesis implies that the functional regions

implicated in domestication would be out of coding regions (promotors, enhancers, etc. (e.g., Li et

al 2018, Rubin et al 2012, Anderson 2012). However, although being a promising hypothesis, we

did not analyze those regions because it requires a very accurate analysis of homology and their

associated functionality, which is very complicated at the genome level, especially for non-model

species. The third hypothesis suggests that the domesticated phenotype is caused by a moderate

change in the frequency of a relatively large number of variants with small selective effects. In this

case, depending on the size of the selective effect, the number of fixed shared variants may be

significant at the genome level, or alternatively, there would be only changes in the frequencies of

the variants without reaching fixation. In the last case, the functional variants involved in

domestication should be segregating in the analyzed populations. These positively selected

variants segregating at high frequencies, together with the presence of deleterious mutations also

segregating at low frequencies, would be reflected as an excess of non-neutral compared to

divergence. Hence, negative α values calculated using statistics based on high frequencies variants

should be observed in cases where there is a significant proportion of positive selection variants

that have not yet being fixed. We have observed this pattern in WB and the IB breed, but not in

LW. Furthermore, the estimation of DFE from ABC analysis showed that the global proportion of

beneficial mutations (weak and strong) is relatively small (~0.9%) and similar in all wild and

domestic populations. Nevertheless, this proportion of mutations is substantial in absolute

numbers. Although speculative, these mutations may change the fate of these populations that are

affected by natural or artificial selection.

We expected that shared variants between populations would be enriched by selective pressures

that predate domestication. Although they may reflect biological constraints at the species level,

these shared polymorphisms can also be the source of phenotypic variation in a polygenic selective

scenario such that a change in their frequencies (in an infinitesimal scenario) would result in the

observed phenotypic differences among the breeds. Furthermore, private variants (those

segregating only in one breed) may reflect recent and breed-specific selective hallmarks and hence,

would be responsible for the observed differences between domesticated and wild breeds. In both

cases, shared and exclusive polymorphisms are contributing to the differences in the SFS between

functional and nonfunctional positions. We expect that adaptive changes would increase the ratio

of nonsynonymous to synonymous polymorphisms and that this should be reflected as an increase

in the negative value of the α statistic. Our simulated domestication process indicates that the effect

of positive selection irrespective of being either strong and affecting a small percentage of variants

or weak and affecting a large percentage of variants is not reflected as marked changes in the
estimated patterns of α. This could be due to the short time since the change in the DFE occurred
but also by the interaction of positive and negative selection and demographic processes. In fact,
the observed α patterns are compatible with demographic effects (population size reduction in WB
and IB and gene flow in LW) but also with the effect of positive selection in LW. Finally, the
obtained results in the ABC analysis based on total variants show a clear genome-wide effect of
the action of purifying selection. We also observed a minor effect of purifying selection in IB and
WB when the analysis was performed based on exclusive variants, which suggest a reduction of
the population size of these two populations. Nevertheless, we had some difficulties in adjusting
the observed data to pertinent DFE models, especially when the analysis was performed based on
exclusive variants. Although the models used here are very simple and contain few parameters,
the real observations contained high heterogeneity that could not be fitted to these models. The
reasons may be technical, conceptual (undetected correlations that distort model assumptions) or
biological (too simplistic models to explain the real data). In any case, a model that assumes a
discrete distribution of beneficial and deleterious mutations (model D) seems to generally explain
the observed data better. The change of the DFE when the analysis was performed based on shared
variants is undistinguishable from that based on all variants, indicating that exclusive variants
should be more useful to detect the effects of the change of selective effects.

Final remarks

The observed patterns of variability are compatible with the presence of deleterious mutations
segregating in all three breeds and with weak signals of positive selection. Nevertheless, when the
variants are split into shared and exclusive, we observed patterns that are in line with the simulated
data under different demographic scenarios joint with the action of positive and negative selection.
We found a clear effect of deleterious mutations at low-frequency variants and a mild effect of
positive selection at high frequencies. Additional analyses contrasting evolutionary models that
consider the effects of standing variation whose effect change under domestication may shed more
light and will help to understand the patterns of variation due to the domestication process.

ACKNOWLEDGMENTS

We acknowledge L. Silió and M. C. Rodríguez for helpful comments on the manuscript. This work was

supported by Ministerio de Economia y Competitividad grants AGL2013-41834-R (MEC, Spain),

AGL2016-78709-R (MEC, Spain) and by the CERCA Programme/Generalitat de Catalunya. We

acknowledge financial support from the Spanish Ministry of Economy and Competitiveness,

through the "Severo Ochoa Programme for Centres of Excellence in R&D" 2016-2019 (SEV-

2015-0533). S.G-R. was supported by a Beautriu de Pinós postdoctoral fellowship (AGAUR; 2014

BP-B 00027).

LITERATURE CITED

Achaz G. 2009. "Frequency Spectrum Neutrality Tests: One for All and All for One." *Genetics*
183: 249–58. https://doi.org/10.1534/genetics.109.104042.

Ai H, Fang X, Yang B, Huang Z, Chen H, Mao L, et al. 215 "Adaptation and possible ancient
interspecies introgression in pigs identified by whole- genome sequencing." *Nat Genet.*
2015;47:217–25.

Alvarez-Ponce, David, and Mario A. Fares. 2012. "Evolutionary Rate and Duplicability in the
Arabidopsis Thaliana Protein-Protein Interaction Network." *Genome Biology and Evolution*
4 (12): 1263–74. https://doi.org/10.1093/gbe/evs101.

Alves, E,, C Ovilo, C Rodríguez, and L Silió. 2003. "Mitochondrial DNA sequence variation and
phylogenetic relationships among Iberian pigs and other domestic and wild pig
populations." *Animal Genetics* 34:319-324.

Alves, Estefania, A I Fernández, Carmen Barragán, C Ovilo, C Rodríguez, and L Silió. 2006.
"Inference of Hidden Population Substructure of the Iberian Pig Breed Using Multilocus
Microsatellite Data." *Spanish Journal of Agricultural Research* 4 (1): 37–46.
http://www.inia.es/gcontrec/pub/alves-fernandez-barragan-..._1141287881015.pdf.

Amaral AJ, Ferretti L, Megens HJ, Crooijmans RPMA, Ni H, Ramos-Onsins SE, Perez-Enciso
M, Schook LB, Groenen MAM. 2011 "Genome-wide footprints of pig domestication and
selection revealed through massive parallel sequencing of pooled DNA." *PLoS One.*
2011;6(4):e14782. doi:10.1371/journal.pone.0014782

Andersson L. 2012 "How selective sweeps in domestic animals provide new insight into
biological mechanisms." J Intern Med. 2012;271:1–14.

Andersson L. 2013 "Molecular consequences of animal breeding." *Curr Opin Genet Dev*.
23(3):295-301. doi: 10.1016/j.gde.2013.02.014.

Beaumont MA, Zhang W, Balding DJ. Approximate Bayesian computation in population
genetics. Genetics. 2002;162(4):2025-2035.

Bianco, Erica, Bruno Nevado, Sebastian E. Ramos-Onsins, and Miguel Pérez-Enciso. 2015. "A
Deep Catalog of Autosomal Single Nucleotide Variation in the Pig." *PLoS ONE* 10 (3): 1–
21. https://doi.org/10.1371/journal.pone.0118867.

Booker, Tom R, and Peter D Keightley. 2018. "Understanding the Factors That Shape Patterns of
Nucleotide Diversity in the House Mouse Genome." *Molecular Biology and Evolution* 35
(12): 2971–88. https://doi.org/10.1093/molbev/msy188.

Bosse, Mirte, Hendrik-Jan Megens, Laurent A. F. Frantz, Ole Madsen, Greger Larson, Yogesh
Paudel, Naomi Duijvesteijn, et al. 2014. "Genomic Analysis Reveals Selection for Asian
Genes in European Pigs Following Human-Mediated Introgression." *Nature
Communications* 5: 4392. https://doi.org/10.1038/ncomms5392.

Bosse, Mirte, Hendrik-Jan Megens, Ole Madsen, Laurent A. F. Frantz, Yogesh Paudel, Richard
P. M. A. Crooijmans, and Martien A. M. Groenen. 2014. "Untangling the Hybrid Nature of
Modern Pig Genomes: A Mosaic Derived from Biogeographically Distinct and Highly
Divergent Sus Scrofa Populations." *Molecular Ecology* 23 (16): 4089–4102.
https://doi.org/10.1111/mec.12807.

Bosse, Mirte, Hendrik Jan Megens, Ole Madsen, Yogesh Paudel, Laurent A. F. Frantz, Lawrence
B. Schook, Richard P.M.A. Crooijmans, and Martien A.M. Groenen. 2012. "Regions of
Homozygosity in the Porcine Genome: Consequence of Demography and the
Recombination Landscape." *PLoS Genetics* 8 (11).
https://doi.org/10.1371/journal.pgen.1003100.

Chen J, Ni P, Li X, Han J, Jakovlić I, Zhang C, Zhao S. 2018 "Population size may shape the
accumulation of functional mutations following domestication." *BMC Evolutionary Biology*
18:4 DOI 10.1186/s12862-018-1120-6

Cruz F, Vilà C, and Webster MT. 2008. "The Legacy of Domestication: Accumulation of
Deleterious Mutations in the Dog Genome." *Molecular Biology and Evolution* 25 (11):
2331–36. https://doi.org/10.1093/molbev/msn177.

Csardi G., and Nepusz T. 2006. "The Igraph Software Package for Complex Network Research."
InterJournal, Complex Systems 1695. http://igraph.org.

Csilléry K, Olivier François O. and Blum MGB. 2012 " abc: an R package for approximate
Bayesian computation (ABC) " Methods in Ecology and Evolution 2012, 3, 475–479.

Esteve-Codina, Anna, Yogesh Paudel, Luca Ferretti, Emanuele Raineri, Hendrik-Jan Megens,
Luis Silió, María C Rodríguez, Martien A. M. Groenen, Sebastian E. Ramos-Onsins, and
Miguel Pérez-Enciso. 2013. "Dissecting Structural and Nucleotide Genome- Wide
Variation in Inbred Iberian Pigs." *BMC Genomics* 14 (148): 1. https://doi.org/10.1186/1471-
2164-14-148.

Eyre-Walker, Adam. 2002. "Changing Effective Population Size and the McDonald-Kreitman
Test." *Genetics* 162: 2017–24.
http://www.genetics.org/content/genetics/162/4/2017.full.pdf.

Eyre-Walker A. 2006 "The genomic rate of adaptive evolution." *Trends Ecol Evol* 21(10):

569–575.

Fang M, Larson G, Ribeiro HS, Li N, Andersson L. 2011 "Contrasting mode of evolution at a
coat color locus in wild and domestic pigs." *PLoS Genet.* 5:e1000341.

Fay, Justin C, and Chung-I Wu. 2000. "Hitchhiking Under Positive Darwinian Selection."
Genetics 155: 1405–13.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1461156/pdf/10880498.pdf.

Fay JC, Wyckoff GJ, Wu CI. 2001 "Positive and negative selection on the human ge-

nome." *Genetics* 158(3):1227–1234.

Ferretti, Luca, Emanuele Raineri, and Sebastian E. Ramos-Onsins. 2012. "Neutrality Tests for
Sequences with Missing Data." *Genetics* 191 (4): 1397–1401.
https://doi.org/10.1534/genetics.112.139949.

Fisher, R. A. 1919. "XV.—The Correlation between Relatives on the Supposition of Mendelian

Inheritance." *Transactions of the Royal Society of Edinburgh* 52 (02): 399–433.

https://doi.org/10.1017/S0080456800012163.

Frantz, Laurent A. F., Joshua G. Schraiber, Ole Madsen, Hendrik-Jan Megens, Mirte Bosse,

Yogesh Paudel, Gono Semiadi, et al. 2013. "Genome Sequencing Reveals Fine Scale

Diversification and Reticulation History during Speciation in Sus." *Genome Biology* 14 (9):

R107. https://doi.org/10.1186/gb-2013-14-9-r107.

Frantz, Laurent A. F., Joshua G. Schraiber, Ole Madsen, Hendrik-Jan Megens, Alex Cagan,

Mirte Bosse, Yogesh Paudel, Richard P. M. A. Crooijmans, Greger Larson, and Martien A.

M. Groenen. 2015. "Evidence of Long-Term Gene Flow and Selection during

Domestication from Analyses of Eurasian Wild and Domestic Pig Genomes." *Nature*

Genetics 47 (10): 1141–48. https://doi.org/10.1038/ng.3394.

Fraser, Hunter B, Aaron E Hirsh, Lars M Steinmetz, Curt Scharfe, and Marcus W Feldman.

2002. "Evolutionary Rate in the Protein Interaction Network." *Science* 296 (5568): 750–52.

https://doi.org/10.1126/science.1068696.

Fu, Yun-Xin, and Wen-Hsiung Li. 1993. "Maximum Likelihood Estimation of Population

Parameters." *Genetics* 134: 1261–70.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1205593/pdf/ge13441261.pdf.

Groenen, Martien A. M. 2016. "A Decade of Pig Genome Sequencing: A Window on Pig

Domestication and Evolution." *Genetics, Selection, Evolution : GSE Sel Evol* 48 (23): 1–9.

https://doi.org/10.1186/s12711-016-0204-2.

Groenen, Martien A. M., Alan L. Archibald, Hirohide Uenishi, Cristopher K. Tuggle, Yasuhiro

Takeuchi, Max F. Rothschild, Claire Rogel-Gaillard, et al. 2012. "Analyses of Pig Genomes

Provide Insight into Porcine Demography and Evolution." *Nature* 491 (7424): 393–98.

https://doi.org/10.1038/nature11622.

Guirao-Rico, Sara, Oscar Ramirez, Ana Ojeda, Marcel Amills, and Sebastian E. Ramos-Onsins.

2018. "Porcine Y-Chromosome Variation Is Consistent with the Occurrence of Paternal

Gene Flow from Non-Asian to Asian Populations." *Heredity* 120 (1): 63–76.

https://doi.org/10.1038/s41437-017-0002-9.

Hahn, Matthew W., and Andrew D. Kern. 2005. "Comparative Genomics of Centrality and

Essentiality in Three Eukaryotic Protein-Interaction Networks." *Molecular Biology and*

Evolution 22 (4): 803–6. https://doi.org/10.1093/molbev/msi072.

Haller, Benjamin C., and Philipp W. Messer. 2017. "SLiM 2: Flexible, Interactive Forward

Genetic Simulations." *Molecular Biology and Evolution* 34 (1): 230–40.

https://doi.org/10.1093/molbev/msw211.

Hill, W G, and A Robertson. 1966. "The Effect of Linkage on Limits to Artificial Selection."

Genetical Research 8 (3): 269–94. http://www.ncbi.nlm.nih.gov/pubmed/5980116.

Kanehisa, Minoru, Michihiro Araki, Susumu Goto, Masahiro Hattori, Mika Hirakawa, Masumi

Itoh, Toshiaki Katayama, et al. 2008. "KEGG for Linking Genomes to Life and the

Environment." *Nucleic Acids Research* 36 (SUPPL. 1): 480–84.

https://doi.org/10.1093/nar/gkm882.

Kim H, Song KD, Kim HJ, Park W, Kim J, Lee T, et al. 2015 "Exploring the genetic signature of

body size in Yucatan miniature pig." *PLoS One*. 10:e0121732.

Kono TJY, Fu F, Mohammadi M, Hoffman PJ, Liu C, Stupar RM, Smith KP, Tiffin P, Fay JC,

Morrell PL. 2016 "The Role of Deleterious Substitutions in Crop Genomes." *Mol. Biol.*

Evol. 33(9):2307–2317. https://doi:10.1093/molbev/msw102

Leno-Colorado, Jordi, Nick J Hudson, Antonio Reverter, and Miguel Pérez-Enciso. 2017. "A

Pathway-Centered Analysis of Pig Domestication and Breeding in Eurasia." *G3* 7 (7):

2171–84. https://doi.org/10.1534/g3.117.042671.

Li C, Zou C, Cui Y, Fu Y, Fang C, Li Y, Li J, Wang W, Xiang H, Li C. 2018 "Genome-wide

epigenetic landscape of pig lincRNAs and their evolution during porcine domestication."

Epigenomics. 2018 Dec;10(12):1603-1618. doi: 10.2217/epi-2017-0117.

Li, Heng, and Richard Durbin. 2009. "Fast and Accurate Short Read Alignment with Burrows-

Wheeler Transform." Journal Article. *Bioinformatics* 25 (14): 1754–60.

https://doi.org/10.1093/bioinformatics/btp324.

Li, Heng, Bob Handsaker, Alec Wysoker, Tim Fennell, Jue Ruan, Nils Homer, Gabor Marth,

Gonçalo R. Abecasis, Richard Durbin, and Subgroup Genome Project Data Processing.

2009. "The Sequence Alignment/Map Format and SAMtools." Journal Article.

Bioinformatics 25 (16): 2078–79. https://doi.org/10.1093/bioinformatics/btp352.

Li, Mingzhou, Shilin Tian, Long Jin, Guangyu Zhou, Ying Li, Yuan Zhang, Tao Wang, et al.
2013. "Genomic Analyses Identify Distinct Patterns of Selection in Domesticated Pigs and
Tibetan Wild Boars." *Nature Genetics* 45 (12): 1431–38. https://doi.org/10.1038/ng.2811.

Li M, Tian S, Yeung CK, Meng X, Tang Q, Niu L, Wang X, Jin L, Ma J, Long K, Zhou C, Cao
Y, Zhu L, Bai L, Tang G, Gu Y, Jiang A, Li X, Li R. 2014 "Whole-genome sequencing of
Berkshire (European native pig) provides insights into its origin and domestication." *Sci
Rep.* 4:4678. doi: 10.1038/srep04678.

Livingstone, Kevin, and Stephanie Anderson. 2009. "Patterns of Variation in the Evolution of
Carotenoid Biosynthetic Pathway Enzymes of Higher Plants." *Journal of Heredity* 100 (6):
754–61. https://doi.org/10.1093/jhered/esp026.

MacEachern S, McEwan J, McCulloch A, Mather A, Savin K, Goddard M. 2009 "Molecular
evolution of the Bovini tribe (Bovidae, Bovinae): is there evidenceof rapid evolution or
reduced selective constraint in Domestic cattle?" *BMC Genomics* (10) *179;* https:// doi:
10.1186/1471-2164-10-179.

Makino T, Rubin CJ, Carneiro M, Axelsson E, Andersson L, Webster MT. 2018 "Elevated
Proportions of Deleterious Genetic Variation in Domestic Animals and Plants." *Genome
Biol Evol.* 10(1):276-290. https://doi: 10.1093/gbe/evy004.

McDonald, J H, and M Kreitman. 1991. "Accelerated Protein Evolution at the Adh Locus in
Drosophila." *Nature* 351: 652–54.

McKenna, Aaron, Matthew Hanna, Eric Banks, Andrey Sivachenko, Kristian Cibulskis, Andrew
Kernytsky, Kiran Garimella, et al. 2010. "The Genome Analysis Toolkit: A MapReduce
Framework for Analyzing next-Generation DNA Sequencing Data." *Genome Research* 20
(9): 1297–1303. https://doi.org/10.1101/gr.107524.110.

Messer, Philipp W., and Dmitri A. Petrov. 2013. "Frequent Adaptation and the McDonald-
Kreitman Test." *Proceedings of the National Academy of Sciences of the United States of
America* 110 (21): 8615–20. https://doi.org/10.1073/pnas.1220835110.

Montanucci, Ludovica, Hafid Laayouni, Giovanni Marco Dall'Olio, and Jaume Bertranpetit.
2011. "Molecular Evolution and Network-Level Analysis of the N-Glycosylation Metabolic
Pathway across Primates." *Molecular Biology and Evolution* 28 (1): 813–23.

https://doi.org/10.1093/molbev/msq259.

Moon, Sunjin, Tae-Hun Kim, Kyung-Tai Lee, Woori Kwak, Taeheon Lee, Si-Woo Lee, Myung-
927   Jick Kim, et al. 2015. "A Genome-Wide Scan for Signatures of Directional Selection in
928   Domesticated Pigs." *BMC Genomics* 16 (1): 1–12. https://doi.org/10.1186/s12864-015-
929   1330-x.

930   Nevado, Bruno, Sebastian E. Ramos-Onsins, and Miguel Perez-Enciso. 2014. "Resequencing
931   Studies of Nonmodel Organisms Using Closely Related Reference Genomes: Optimal
932   Experimental Designs and Bioinformatics Approaches for Population Genomics."
933   *Molecular Ecology* 23 (7): 1764–79. https://doi.org/10.1111/mec.12693.

934   Orlando L, Librado P. 2019 "Origin and Evolution of Deleterious Mutations in Horses." *Genes*
935   10, 649; doi:10.3390/genes10090649

936   Pérez-Enciso, M., G. de los Campos, N. Hudson, J. Kijas, and A. Reverter. 2017. "The
937   'Heritability' of Domestication and Its Functional Partitioning in the Pig." *Heredity* 118:
938   160–68. https://doi.org/10.1038/hdy.2016.78.

939   Quinlan, Aaron R. 2014. "BEDTools: The Swiss-Army Tool for Genome Feature Analysis."
940   *Current Protocols in Bioinformatics / Editoral Board, Andreas D. Baxevanis ... [et Al.]* 47
941   (January): 11.12.1-11.12.34. https://doi.org/10.1002/0471250953.bi1112s47.

942   Ramírez, Oscar, William Burgos-Paz, Encarna Casas, Maria Ballester, Erica Bianco, Iñigo
943   Olalde, Gabriel Santpere, et al. 2014. "Genome Data from a Sixteenth Century Pig
944   Illuminate Modern Breed Relationships." *Heredity* 114 (2): 175–84.
945   https://doi.org/10.1038/hdy.2014.81.

946   Ramsay, Heather, Loren H Rieseberg, and Kermit Ritland. 2009. "The Correlation of
947   Evolutionary Rate with Pathway Position in Plant Terpenoid Biosynthesis." *Molecular*
948   *Biology and Evolution* 26 (5): 1045–53. https://doi.org/10.1093/molbev/msp021.

949   Rausher, Mark D, Richard E Miller, and Peter Tiffin. 1999. "Patterns of Evolutionary Rate
950   Variation Among Genes of the Anthocyanin Biosynthetic Pathway." *Molecular Biology and*
951   *Evolution* 16 (2): 266–74.
952   https://watermark.silverchair.com/mbev_16_02_0266.pdf?token=AQECAHi208BE49Ooan
953   9kkhW_Ercy7Dm3ZL_9Cf3qfKAc485ysgAAAdwwggHYBgkqhkiG9w0BBwagggHJMIIB

xQIBADCCAb4GCSqGSIb3DQEHATAeBglghkgBZQMEAS4wEQQMB_IVBQyX1n-

F9I4pAgEQgIIBj8r0BkU_bdeIWEuY_7bUxMFToA8Yo-26snF_yPY.

Renaut, Sebastien, and Loren H. Rieseberg. 2015. "The Accumulation of Deleterious Mutations

as a Consequence of Domestication and Improvement in Sunflowers and Other Compositae

Crops." *Molecular Biology and Evolution* 32 (9): 2273–83.

https://doi.org/10.1093/molbev/msv106.

Riley, Rebecca M., WeI Jin, and Greg Gibson. 2003. "Contrasting Selection Pressures on

Components of the Ras-Mediated Signal Transduction Pathway in Drosophila." *Molecular*

Ecology 12 (5): 1315–23. https://doi.org/10.1046/j.1365-294X.2003.01741.x.

Rubin, Carl-Johan, Hendrik-Jan Megens, Alvaro Martinez Barrio, Khurram Maqbool, Shumaila

Sayyab, Doreen Schwochow, Chao Wang, et al. 2012. "Strong Signatures of Selection in

the Domestic Pig Genome." *Proceedings of the National Academy of Sciences of the United*

States of America 109 (48): 19529–36. https://doi.org/10.1073/pnas.1217149109.

Tajima, Fumio. 1983. "Evolutionary Relationship of DNA Sequences in Finite Populations."

Genetics 105: 437–60.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1202167/pdf/437.pdf.

Uricchio LH, Petrov DA, Enard D. 2019 "Exploiting selection at linked sites to infer the rate and

strength of adaptation." *Nat Ecol Evol.* 3(6):977-984. doi: 10.1038/s41559-019-0890-6.

Watterson, G.A. 1975. "On the Number of Segregating Sites in Genetical Models without

Recombination." *Theoretical Population Biology* 7 (2): 256–76.

https://doi.org/10.1016/0040-5809(75)90020-9.

Wilkinson S, Lu ZH, Megens HJ, Archibald AL, Haley C, Jackson IJ, Groenen MA, Crooijmans

RP, Ogden R, Wiener P. 2013 "Signatures of diversifying selection in European pig

breeds." PLoS Genet. 2013 Apr;9(4):e1003453. doi: 10.1371/journal.pgen.1003453.

Zeder, Melinda A. 2012. "The Domestication of Animals." *Journal of Anthropological Research*

68 (2): 161–90. https://doi.org/10.3998/jar.0521004.0068.201.

TABLES

Table 1. Number of SNPs (in whole genome, in genes and in coding regions), classified by its
presence in each population.

Table 2. Combinations of SNPs from coding regions according to its allelic status in each
population (A: Ancestral allele, F: Fixed allele, P: Polymorphic allele). SNPs that are missing in
any of the populations are not considered in this table.

FIGURES

Figure 1. Levels of variation at synonymous (A) and nonsynonymous (B) sites for each pig
population and variability estimates and for shared and exclusive variants. WB; wild boar
population; IB, Iberian breed; LW, Large White breed.

Figure 2. Estimates of α for each pig population. Total variants (A), exclusive variants (B), shared
variants (C) and shared variants between IB and LW (D). Bootstrap intervals at 95% are indicated
by a line at each bar. WB; wild boar population; IB, Iberian breed; LW, Large White breed.

Figure 3. Estimates of R_$\beta\gamma$ for each pig population and for all, exclusive and shared variants.
WB; wild boar population; IB, Iberian breed; LW, Large White breed.

Figure 4. Estimates of the median values of α for each pig population, different molecular scales
and for all, exclusive and shared variants. For each population, the order of different α's is: Fu&Li,
Watterson, Tajima and Fay&Wu. WB; wild boar population; IB, Iberian breed; LW, Large White
breed.

Figure 5. Posterior distribution of the α values for total variants. Four different estimators of alpha
(Fu&Li, Watterson, Tajima and Fay&Wu, see Materials and Methods) are used. Box plots indicate
simulated distributions of α values. Red line indicates observed α values.

Figure 6. Posterior distribution of the α values for exclusive variants. Four different estimators of
alpha (Fu&Li, Watterson, Tajima and Fay&Wu, see Materials and Methods) are used. Box plots
indicate the simulated distributions of α values. Red line indicates observed α values.

1007

1008　**SUPPLEMENTARY MATERIAL**

1009　See Supplementary Material file added to see the Tables (S1-S13) and Figures (S1-S50).

Table 1. Number of SNPs present in the whole genome, genes and coding regions and classified according to their presence in pig populations.

	Number of SNPs	Shared between WB, IB and LW	Shared between WB and IB	Shared between WB and LW	Shared between IB and LW	Exclusive of WB	Exclusive of IB	Exclusive of LW
Whole-genome	24,869,699	7,293,787	666,927	4,017,107	138,378	4,239,052	385,504	8,128,944
Genes	6,684,142	1,964,562	98,433	1,152,555	48,351	1,138,370	100,550	2,181,321
Coding regions	149,440	18,611	3,252	25,044	1,177	51,432	3,356	46,568

Table 2. Number of synonymous and nonsynonymous SNPs for different combinations of the allelic status. SNPs that are missing in any of the populations are not considered in this table.

IB	LW	WB	Synonymous	Non-synonymous
F	F	F	20297	9342
P	P	P	11712	7597
A	A	F	0	0
A	F	A	0	0
F	A	A	3	5
A	A	P	30314	20988
A	P	A	26027	15035
P	A	A	1833	1588
A	F	F	0	0
F	A	F	1	0
F	F	A	1	0
A	P	P	10128	7930
P	A	P	1676	1254
P	P	A	700	363
A	F	P	11	1
A	P	F	0	2
F	A	P	30	30
P	A	F	0	0
F	P	A	8	4
P	F	A	1	1
F	P	P	4924	2378
P	F	P	242	139
P	P	F	81	52
F	F	P	1140	489
F	P	F	4911	2073
P	F	F	38	22
			114078	69293

F: position with a fixed derived variant. **P**: Polymorphic position. **A**: position with the ancestral varian

Figure 1

A

B

Figure 2

Figure 4

Figure 5

Figure 6