Title:

The *Azolla* fern symbiosis sexual reproduction requires far-red light and involves responsive CMADS1 homologue, miR319-controlled GAMYB, miR529 in the fern, and transporters in the symbiont

Authors:

Laura W. Dijkhuizen[1], Badraldin E. S. Tabatabaei[2], Paul Brouwer[1], Niels Rijken[1], Rushi S. Mehta[1], Erbil Güngör[1], Henriette Schluepmann[1]

Addresses:

1, Department of Biology, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands

2, Department of Biotechnology, College of Agriculture, Isfahan University of Technology, Isfahan, 84156-83111, Iran

Author for correspondence:

H. Schluepmann, tel +31 30 2533289, e-mail h.schlupmann@uu.nl

Total word count: 6404

word counts for each section:

Introduction:	1041
M&M:	1291
Results:	2498
Discussion:	1574

Number of Figures: 7, all of which are in color.

Tables : 2

Supporting Information: Supplemental methods
Table S1-5
Fig. S1-8

Summary

- Regulation of the transition from vegetative to spore-forming phases in ferns is largely
unknown yet a pre-requisite for their domestication. External and endogenous cues regulating
the formation of sporocarps were researched in *Azolla* ferns, that are heterosporous and
constitute a symbiosis with the cyanobacteria *Nostoc azollae*.

- Once external cues were identified, small RNA and mRNA from eukaryotic and prokaryotic
partners in the symbiosis were analyzed for differential accumulation upon transition to
sexual reproduction.

- *A. filiculoides* fern sporocarp formation required far-red light (FR) and was suppressed by
medium nitrogen; FR-induced sporocarps were viable in crosses that verified species
attribution of Dutch strains but not those from Iran's Anzali lagoon. FR-responsive transcripts
encoded the *Azolla* MIKC^C homologue of CMADS1, miRNA319-controlled GAMYB in the fern,
ycf2 in chloroplasts, and transporters in *N. azollae*. Loci of conserved microRNA in the fern
lineage included miR172, but FR only induced mirR529 and miR535, and reduced miR159 and
miR319.

- The clade of the FR-responsive MIKC^C radiated separately in ferns from the seed plant clade
containing *AtSOC1* and floral homeotic genes A, C, D and E, yet the data supports that regulons
controlling seed plant flowering originate from the diploid to haploid phase transition in the
common ancestor of seed plants and ferns.

Keywords: *Azolla*, Dual RNA-sequencing, fern, GAMYB/MIKC^C transcription factors, *Nostoc azollae*,
miRNA, Sexual reproduction, Symbiosis.

INTRODUCTION (1041)

Ferns from the genus *Azolla* thrive floating on freshwater. They require no nitrogen whilst growing prolifically due to an unusual symbiosis with the cyanobacterium *Nostoc azollae* which fixates dinitrogen with light energy (Brouwer *et al.*, 2017). Previously used as biofertilizer, they now are considered a candidate crop for sustainable production of plant protein in subsiding delta regions but have never been domesticated (Brouwer *et al.*, 2014, 2018). What controls their sexual reproduction is completely unknown yet central to containment, propagation, breeding schemes or the establishment of biodiversity and germplasm banks.

Azolla ferns are heterosporous with diploid sporophytes generating two types of sori mostly called sporocarps (Nagalingum *et al.*, 2006; Coulter, 1910). Unlike homosporous ferns, the haploid phase of *Azolla* ferns, the gametophytes, are not free-living but contained inside the sporocarps. The sexual reproduction is therefore controlled by the sporophyte as in the case of seed plants. Unlike seed plants, gametophytes do not develop inside the sporocarps when these are still on the sporophytes and therefore fertilization is not guided by structures on the sporophyte such as flowers in angiosperms. *Azolla* sporocarps typically detach when mature and sink to the sediment of ditches; they constitute resting stages with nutrient reserves that survive drying or freezing and, thus, are key for dissemination practices (Brouwer *et al.*, 2014).

The developmental transition to reproductive growth of the sporophyte (RD) occurs when shoot apices of the sporophytes will form initials for sporocarps in addition to branches, leaves and roots. RD is not well studied in ferns. It is analogous, or possibly homologous, to the transition in angiosperms of shoot apical meristems (SAM) to inflorescence meristems: the meristem is switching to making branches with leaves and sporocarps whereby the making of sporocarps is ancestral to the making of leaves (Vasco *et al.*, 2016). Gene expression in lycophytes and ferns of class III HD-Zip transcription factors (C3HDZ) was observed in both leaves and sporangia even though ferns and lycophyte fossil records have shown that the lineages evolved leaves independently. Vasco *et al*. (2016), therefore, suggested that the common C3HDZ role in sporangium development was co-opted for leaf development when leaf forms, the micro and megaphylls, evolved in lycophytes and ferns, respectively.

In angiosperm SAM transition to RD, the LEAFY transcription factor (TF) induces floral homeotic genes and a shoot meristem identity gene; LEAFY functions solely when bound to proteins in a complex (Sayou *et al.,* 2016). LEAFY is conserved in plant lineages, but LEAFY does not promote formation of reproductive

structures in seed-free plants including ferns where, nonetheless, the protein was reported to maintain

apical stem cell activity (Plackett *et al.*, 2018). The pattern emerging, therefore, is that the LEAFY complex

is not among those that were already in place to link exogenous cues to the transition of SAM to RD in the

common ancestor of seed plants and ferns.

In angiosperms, MIKC^C-type TF are defined by the conserved MADS box, Intervening, Keratin and C-

terminal domains and form characteristic complexes that integrate endogenous and exogenous cues

turning SAM into RD (Theißen *et al.*, 2018). In *Arabidopsis thaliana (Arabidopsis)*, for example, gibberellic

acid (GA)-dependent and photoperiod/temperature signaling is integrated by MIKC^C from the FLC and SVP

clades that, unsurprisingly, are floral repressors. Consistent with separate transitions for inflorescence

and flower meristems in Arabidopsis, the MIKC^C floral integrators act on another, SOC1, that in turn

induces the expression of MIKC^C floral homeotic genes. Homeotic genes of particular interest specify

ovules, and stamens. These modules could be more ancient than those specifying carpels and sepals but

specification of ovules/stamens in angiosperms may not be homologous to that of mega- and

microsporocarps in *Azolla* ferns because heterospory evolved independently in ferns and angiosperms

(Sessa & Der, 2016).

In addition to MIKC^C TF, micro RNA (miR) are known to control phase transitions in angiosperms, including

RD. The miR156/172 age pathway controls flowering: miRNA156 decreases with the age of plants, and

thus also its repression of the targets *SQUAMOSA PROMOTER BINDING-LIKES* (*SPL*). In the annual

Arabidopsis, SPL promote expression of mir172 that targets another type of floral repressor, the AP2/TOE

TF (Hyun *et al.*, 2019). In the perennial *Arabis alpina*, the MIKC^C FLC homologue and miR156 regulate

expression of *SPL15* thus integrating temperature and age-dependent RD. Moreover, AP2/TOE and NIN

TF are known to mediate repression of *SOC1,* and thus RD, in Arabidopsis on high nitrate (Gras *et al.*, 2018;

Olas *et al.*, 2019). SOC1 is further known to control the steady state of miR319 that inhibits TCP protein

functions necessary for the transition to RD (Lucero *et al.*, 2017). Inextricably, therefore, transition to RD

is linked to conserved miRNA modules in angiosperms but we know little of miRNA loci in ferns (Berruezo

et al., 2017; You *et al.*, 2017).

Azolla ferns and Arabidopsis were shown to share responses to nitrogen starvation, suggesting that the

ancestor of both had already evolved some of the responses to exogenous cues we know from

angiosperms (Brouwer *et al.*, 2017). GA applications to *Azolla* ferns, however, were reported to not be

effective to induce RD in *Azolla*, neither was stress treatment (Kar *et al.*, 2002). Systematic research of the

symbiosis responses to day-length and light quality is lacking. The coordinate development of the

cyanobacterial symbiont, *N. azollae,* and fern during the induction of sporocarps is documented:

hormogonia generated in the shoot apical colony are attracted to the developing sporocarps in the shoot

apex; attraction to either the leaf cavities or the sporocarps is likely mediated by secretions from

specialized trichomes (Perkins & Peters, 1993). How the development of symbionts and host are

coordinated upon transition to RD is unknown. Sequences of both genomes are now available, setting the

stage for gene-expression profiling of both the pro- and eukaryote RNA by dual RNA sequencing.

Here we identify external cues that induce or inhibit transition to RD in *A. filiculoides* in laboratory settings.

We test the viability of reproductive structures induced in different fern strains in crosses, herby defining

species attributions, and support the results obtained with phylogenetic analyses of the strains. To reveal

endogenous mechanisms mediating transition to RD, we profile transcript accumulation by dual RNA-

sequencing comparing sporophytes induced for RD by far-red light, at a time point when the reproductive

structures are not yet visible. We assign induced MIKC^C and MYB TF to known clades using phylogenetic

analyses. In addition, we profile small RNA from the symbiosis, then identify miRNA and their targets in

the fern lineage. Finally, we discover regulatory modules of conserved miRNA and their targets responsive

to induction of RD in ferns.

MATERIALS AND METHODS (1353)

Plant Materials, growth conditions and crosses

Ferns were collected from different locations in the Netherlands, grown in liquid medium at a constant

temperature of 22^{o}C and a 16 h light period as described earlier (Brouwer *et al.*, 2017). Ferns from Spain

and Iran were processed to DNA on site. GPS coordinates from the collection sites of the strains used in

phylogenetic analyses are in Supporting Information (SI) Table S**1**. For sporocarp induction, ferns were

first raised under 16 h tube light (TL) periods, then transferred for induction under 16 h TL with far-red

(FR) Light Emitting Diodes - LED (Phillips GrowLED peaking at 735 nm, SI Fig. **S1a,b**). Photosynthetic Photon

Flux Density varied in the range 100-120 µmol m^{-2} s^{-1}. Sporophytes were generally kept at densities that

covered the surface. When testing sporocarp induction sporophytes were kept at set densities (with fresh

weight resets once per week, SI Fig. S**1c,d,e**) and set red to FR light ratios (SI Table S**2**). Phenotypic change

in mature sporophytes were photographed using a Nikon D300s with an AF Micro-Nikkor 60mm f/2.8D

objective or bellows on a rail to generate image reconstructions with Helicon focus software**.** Mega- and

microsporocarps were collected manually using tweezers. To initiate crosses, microsporocarps were

Dounce-homogenized to release massulae and then added to megasporocarps in distilled water (3-6 ml,

in 3 cm diameter petri dish), gently shaking to obtain clumps. The crosses were kept at room temperature

(about 18°C) under 16 h light conditions under TL at about 100 µmol m^{-2} s^{-1}. As sporelings emerged they

were transferred to liquid medium (Brouwer *et al.*, 2017).

RNA extraction and quantitative RT-PCR were as in (Brouwer *et al.*, 2017) and are detailed in Methods

S1.

Phylogenetic analyses are described in details in Methods **S1**. Chloroplastic marker regions were as in SI

Table S**3**. The MIKC^C tree was subsampled from individual genome browsers with 1 KP as a guide (Leebens-

Mack *et al.*, 2019); the R2R3 MYB tree used data from respective genome repositories, the *Azolla* genome

(Li *et al.,* 2018) and Jiang & Rao, 2020.

Reference genomes used to align Dual RNA sequences onto in the *A. filiculoides* Galgenwaard strain are

detailed in Methods **S1**.

Dual-RNA sequencing and data processing

A. filiculoides sporophytes were from the sequenced accession Galgenwaard (Li *et al.*, 2018). The ferns

were maintained on liquid medium without nitrogen in long-day TL, then transferred to TL with and

without FR for 7 days, on fresh liquid medium without nitrogen; harvest was 2 h into the light period by

snap freezing. All samples were grown separately and harvested as triplicate biological replicates. Total

RNA was extracted using the Spectrum Plant Total RNA (Sigma-Aldrich) using protocol B, then DNAse

treated as described in (Brouwer *et al.*, 2017), and cleaned using the RNeasy MinElute Cleanup Kit

(Qiagen). rRNA depletions were carried out as per protocol using Ribo-Zero rRNA Removal Kit (Plant Leaf,

Illumina), and the RNA was then cleaned again using the RNeasy MinElute Cleanup Kit. Stranded libraries

were then synthesized using the Ovation Complete Prokaryotic RNA-Seq Library Systems kit (Nugen). The

libraries thus obtained were characterized and quantified before sequencing using a single lane Hiseq4000

and 50 cycles PE reading (Illumina).

Reads obtained were de-multiplexed then trimmed and quality filtered with trimmomatic (Bolger *et al.,*

2014), yielding 13-60 million quality PE reads per sample. Alignment of the reads to the genome assembly

and GFF vs1 of *A. filiculoides* (Azfi_vs1; Li *et al.*, 2018) was with STAR-aligner default settings (Dobin *et al.*,

2013) and was combined with the STAR read count feature (PE read counts corresponding to the HTseq

"Union" settings, (Anders *et al.*, 2015)). Read count tables were fed into DESeq2 (Love *et al.*, 2013) for

analysis of differentially accumulating transcripts.

STAR was also used to align reads to the *N. azollae* and the chloroplast genomes, and read counts then

extracted using STAR default parameters or VERSE (Zhu *et al.*, 2016) with HTSeq setting "Union"

comparing single and paired end read counts on the polycistronic RNA. The three approaches yielded

similarly ranked genes but the Padj statistics of differential expression in count tables from the forced

single read counts yielded more dispersion in the data: only 20 *N. azollae* genes had P- adjusted values

smaller than 0.1 compared to up to 67 with the PE- approaches, we therefore present the PE-approach

using STAR read counts only.

STAR alignment to concatenated genomes of the *Azolla* symbiosis using default settings yielded a large

proportion of multi-mappers which interfered with PE read counting leading to losses of as much as 30%

of the reads. The strategy would reduce force mapping but is not necessary to extract differentially

accumulating transcripts. We, however, used alignments allowing no mismatches to concatenated

genomes when mapping the sRNA reads below, then recovering reads aligning to each genome using

Samtools fasta from the indexed .bam files.

Small RNA data sequencing and data processing

Ferns were maintained on liquid medium without nitrogen with 16 h TL, then transferred to TL with (F)

and without (T) FR for 7 days as described under Dual-RNA sequencing. In addition, sporophytes in F were

also grown on medium with 2 mM NH_4NO_3 (FN); ferns without cyanobacteria (C) were grown as in FN. All

sporophytes were harvested 7 d into the treatment, 2 h into the light period snap frozen. Samples were

harvested from triplicate growth experiments and thus represent triplicate biological replicates. Small

RNA was extracted as per protocol using the mirPremier microRNA Isolation Kit (Sigma-Aldrich), then

DNAse treated (Brouwer *et al.*, 2017), then cleaned using the RNeasy MinElute Cleanup Kit (Qiagen)

protocol for small RNA. Libraries of the sRNA were then generated using the SeqMatic TailorMix miRNA

Sample Preparation 12-reaction Kit (Version 2) as per protocol. Libraries were characterized using an

Agilent Technologies 2100 Bioanalyzer and quantified before sequencing using NextSeq500 and 1 x 75 b

read chemistry (Illumina).

Reads thus obtained were de-multiplexed, trimmed and quality filtered with trimmomatic, then aligned

with STAR (Dobin *et al.*, 2013) (set at maximal stringency and to output Bam files) to the concatenated

fasta files of the genomes Azfi_vs1, its chloroplast, and *N. azollae* 0708. The resulting bam files were then

split for each genome, and Samtools fasta (Li *et al.*, 2009) used to extract reads aligning to each genome.

fastx collapser (http://hannonlab.cshl.edu/fastx_toolkit) was then used to extract read counts for each

identical sequence, with the sequence defining the name in the lists obtained. A data matrix was

generated in bash, then exported to Excel or Rstudio for further analyses.

To determine differential expression, the lists obtained were restricted to sRNA with at least 10 reads per

sample joined in bash, and the tables thus obtained, first restricted to containing only reads of 20-22 nt

sRNA (to minimize gel excision bias during sequencing library preparation), then exported to R (Booth *et*

al., 2018). DESeq2 was then used to display variation and dispersion in the data and identify differentially

accumulating sRNA using Padj values (Love *et al.*, 2013).

miRNA discovery

The miRNA loci in *A. filiculoides* were analyzed as in SI Fig. S**2**. Pre-miRNA were predicted with miRDEEP-

P2 (Kuang *et al.*, 2019); multi-mappers min. 100 and max. loop length 500) then extracted and subjected

to secondary structure prediction using RandFold default parameters (Bonnet *et al.*, 2004; Gruber *et al.*,

2008). miRNA candidates containing less than four mismatches within the hairpin structure were viewed

in IGV along with RNA expression data aligned to Azfi_vs1. Predicted miRNAs that were expressed in *A.*

filiculoides were then compared in miRbase (Kozomara *et al.*, 2019). Prediction of miRNA targets in

Azfi_vs1 was using the intersect of both psRNATarget (Dai *et al.*, 2018) and TargetFinder (Srivastava *et al.*,

2014) with a cutoff of 5.0. Predicted targets were linked to the closest homolog using the Mercator

annotation (Lohse *et al.*, 2014) of Azfi_vs1 predicted proteins.

Data sharing

The data that support the findings of this study are openly available in [repository name] at [URL],

reference number [reference number]. Data will be uploaded upon acceptance of the manuscript and will

include 1) read files for the dual RNA seq as well as small RNA seq on ENA 2) phylogeny files on Gitup.

RESULTS (2498 words)

Far-red light induces, while presence of nitrogen in the medium represses, sexual reproduction in *A.*

filiculoides

A young fern growing at low density typically spreads over the water surface (Fig. **1a**) in contrast to the

three dimensional web of a fern that underwent transition to reproductive development in dense

canopies (Fig. **1b**). Unlike in the glasshouse, *A. filiculoides* ferns never sporulated under tube light (TL, SI

Fig. S**1a**), TL did not emit any photons in the far-red range (FR, SI Fig. S**1a**). We, therefore, added far-red

LED (FR, SI Fig. S**1b**) during the entire 16 h light period. This initially resulted in the elongation of the stem

internodes of the fern (Fig. **1c**) and, after 2-3 weeks, in the formation of the sporocarps (Fig. **1d**).

Sporocarps were positioned as pairs on the branch side of branch points and were predominantly

arranged as a mega- and microsporocarp pair (Fig. **1e**).

To test the optimum light for sporocarp induction, decreasing ratios of red to far-red LED were applied to

ferns that had been raised in TL light and were then maintained at three differing densities (SI Table S**2**,

SI Fig. S**1c,d,e**). Far-red light was required for sporocarp formation regardless of the density of the fern

culture (Fig. **1f**). Increasing far-red whilst keeping a constant photosynthetic active radiation (PAR) yielded

more sporocarps per fern mass. The sporocarps per fern mass increased with time of exposure to the far-

red LED up until five weeks and then decreased, presumably because ripe sporocarps detached. The

sporocarps per fern mass, furthermore, increased with the density of the fern culture. Reducing the PAR

at highest far-red intensity revealed that PAR limits the production sporocarps at high far-red light (Fig.

1f_FR4).

Adding 2 mM ammonium nitrate to the medium drastically reduced the spore count when inducing

sporocarp formation with far-red LED (Fig. **1g**). Both, ammonium and nitrate added individually inhibited

sporocarp formation (data not shown), and inhibition explained why sporophytes without *N. azollae* kept

on various nitrogen sources never sporulated.

The ratio of mega to microsporocarp observed was approximately equal over nine weeks of induction (SI

Fig. S**1f**); and the marker *AzfiSOC1* (Brouwer *et al.*, 2014) was maximally induced when sporocarps became

visible four weeks after induction started.

FR-induced sporocarps are viable which permits reciprocal crosses between strains to test phylogeny-

derived species attributions

We next wondered whether the sporocarps induced with far-red LED were viable. We randomly tested

reciprocal crosses of the sequenced strain *A. filiculoides* Galgenwaard (Li *et al.*, 2018), six further strains

from the Netherlands, one from Gran Canaria (Spain) and another from the Anzali lagoon (Iran) (SI Table

S**1**). All strains could be induced to sporulate under far-red supplemented TL except Anzali. All sporulating

strains tested in crosses could be crossed suggesting that the ferns from the Netherlands and Spain were

A. filiculoides (Table **1**).

To research the taxonomy of the strains, intergenic regions of the choloroplast rRNA, used for phylogenetic studies previously (Madeira *et al.,* 2016), were amplified and sequenced to compute phylogenetic relations. The regions trnL-trnF (Fig. **2a**), and trnG-trnR (Fig. **2b**) yielded similar trees confirming the close relationship of the *A. filiculoides* strains from the Netherlands and Spain. The trees further revealed that the Anzali strain clusters along with two stains from South America (Southern Brazil and Uruguay, SI Table S**1**). Use of the nuclear ITS region for phylogeny reconstruction proved futile when amplifying the region on the *A. filiculoides* genome *in silico*: the ITS sequence is variable within the genome (Fig. **2c**) and variations are very large when comparing to other species of *Azolla* (SI Fig. S**3a**).

The dorsal-side of the upper leaf lobes of the Anzali strain had two-celled, distinctly pointed, trichomes unlike *A. filiculoides* that mostly, but bot exclusively, have single celled and rounded trichomes (SI Fig. S**3b**). Consistent with its differing response to FR light, and its phylogenetic position based on the chloroplast sequences, the Anzali strain, we conclude, is not *A. filiculoides*.

Dual RNA sequencing by rRNA depletion profiles RNA from organelle, prokaryotic symbiont and fern nucleus

To research molecular processes altered during transition to reproductive development, ferns maintained on TL were transferred to TL with and without FR for a week in triplicate cultures, then collected 2 h into the 16 h light period snap frozen. Total RNA was then extracted, its plant rRNA depleted, and stranded sequencing libraries generated which were depleted for rRNA from gram-negative bacteria. Libraries were sequenced to obtain 12 to 60 million paired end (PE) 50 b reads, quality filtered and trimmed. To align the reads obtained, nuclear and chloroplast genomes with annotations for the *A.filiculoides* Galgenwaard strain were available (Li *et al.,* 2018) but not for its cyanobacterial symbiont.

Assembly of the *N. azollae* genome from metagenome short reads of the Galgenwaard strain yielded a bin with 385 contigs that proved to be highly complete (95.8%) based on single copy marker genes. Contigs aligning to the *N. azollae* 0708 reference genome (Ran *et al.*, 2010) were well over 99.5% identical (Fig. **3a**). This degree of nucleotide identity was well above the maximum mismatch of STAR-aligner's default setting; PE reads were thus aligned to the Galgenwaard strain nuclear and chloroplast genomes, and the *N. azollae* 0708 genome, and read counts derived from their GFF annotation files. Alignments revealed a high proportion of uniquely mapped reads to the fern nuclear and *N. azollae* genomes suggesting efficient removal of rRNA (Fig. **3b**). The larger proportion of PE reads mapping to multiple loci in the chloroplast indicated that fern chloroplast rRNA was not as efficiently removed (Fig. **3b**), yet read counts per gene

were still high because few genes are encoded in the chloroplast genome (84 without rRNA and tRNA

genes). Read counts in genome features further revealed that sense alignments dominated and thus that

the libraries were stranded (Fig. **3c**).

Dispersion of sense read count per gene was low which permitted identification of differentially

accumulating transcripts (defined as DEseq2 Padj <0.1; Love *et al.,* 2014) comparing ferns on TL and TL

with FR: 318 from the fern nucleus, 67 from *N. azollae* and 5 from the fern chloroplast (SI Table S**4**). We

next wondered whether the changes in transcript abundance reflected adaptations to light quality and

the transition to RD.

FR triggers small transcriptional changes reflecting light-harvesting adaptations in chloroplasts and *N.*

azolla,* yet large changes in transporters and transposases of *N.azollae

After a week exposure to FR, increased chloroplast transcripts encoded ycf2 (log2 of 2.9), energizing

transport of proteins into the chloroplast, and RNA polymerase subunit rpoA (log 2 of 1.3) (SI Table S**4**, *A.*

filiculoides chloroplast). Decreased chloroplast transcripts included psbD, previously reported as

responsive to light quality in plant lineages (Shimmura *et al.*, 2017).

Consistent with the high levels of N_2 fixation measured in *Azolla* ferns, the highest *N. azollae* read counts

were in transcripts of the *nif* operon, genes encoding photosystem I and II proteins and ATP synthase

subunits; in addition, read counts for *ntcA* and metabolic enzymes supporting high N_2 fixation were high

(SI Table S**5**). Read counts from the *N. azollae* transcripts thus reflected the activity of the symbiont and

we thus further analyzed the data for differential expression.

In *N. azollae*, FR led to accumulation of the PsbA transcript and a reduction of photosystem I reaction

center subunit XII, geranylgeranyl hydrogenase ChlP, and phycobilisome rod-core linker polypeptide

CpcG2 transcripts (SI Table S**4**, *N. azollae*). But these changes in light harvesting-related genes were small

compared to changes in hypothetical proteins of unknown function, transposases and highly expressed

transporters (TrKA, MFS and ABC transporters). We conclude that one week induction of reproductive

structures with FR triggered few transcriptional changes reflecting light-harvesting adaptations in *N.*

azollae. The larger differential accumulation of transporter transcripts reflect changes in metabolite

trafficking and communication with the host fern.

FR alters transcripts from MIKC[C] and R2R3MYB implicated in the reproductive transition of seed plants

As in the case of *N. azollae* symbiont, the ferns largest changes in transcript accumulation under TL with

FR were in genes with unknown function, then in secondary- and lipid-metabolism (SI Table S**4**, *A.*

filiculoides nucleus). We noted the large induction of the SWEET-like transporter (Azfi_s0221.g058704

log2 7.5 fold), of transcripts related to cysteine-rich peptide signaling (Azfi_s0046.g030195 log2 7.1 fold;

Azfi_s0745.g084813 log2 4.0 fold) and of transcripts from jasmonate metabolism. Transcription factors

(TF) characterize the RD in seed plants and were, therefore, analyzed in more depth. Transcripts of MIKC^C

TF and of R2R3MYB TF were strikingly induced in sporophytes under TL with far-red LED (**Table 2**).

Prediction of the most induced TF transcript encoding a SOC1-like MIKC^C (Azfi_s0028.g024032, log2 7.2

fold) missed the K-domain because this TF was not expressed in vegetative sporophytes. Manual

annotation allowed computation of a phylogenetic tree that placed both induced *Azolla* MIKC^C in the clade

of fern sequences radiating separately from the angiosperm and gymnosperm MIKC^C clade that gave rise

to AtSOC1, FLC, and to the A,C,D and E-function floral homeotic genes (SI Fig. S**4**). Induction of the

AzfiSOC1-paralogue was confirmed by RT-PCR (SI Fig. S**2g**). In Arabidopsis, AtSOC1 is known to control the

steady state of miR319 that inhibits TCP TF functions necessary for the transition to an inflorescence

meristem at the shoot apex (Lucero *et al.*, 2017); the link may also operate in *Azolla* ferns since a TCP TF

(Azfi_s0168.g054602) was induced in ferns on TL with far-red LED (SI Table S**4**, *A. filiculoides* nucleus).

Transcripts of R2R3MYB TF could be assigned to families using data from Jiang & Rao, 2020 (**Table 2**). The

highly induced class VIII-E TF (log2 6.1 fold) likely controls changes in secondary metabolism together with

the similarly changed bHLH TF (Güngör *et al.*, 2020), and therefore may not be linked to RD-directly. In

contrast, induced class V (log 2 2.1 fold) and VII (log 2 1.9 fold) TF are known to affect RD, with the GA

pathway associated class VII GAMYB from seed plants known to be regulated by miR319.

Prominence of transcription factors in Table **2** suspected to controlling RD as in seed plants begged the

question as to whether any RD pathways with conserved miRNA were conserved in the common ancestor

of seed plants and ferns. This required to first characterize miRNA loci in the *A.filiculoides* symbiosis.

Small RNA profiles are characteristic for fern nucleus, chloroplast or *N. azollae*

Short RNA reads were obtained from sporophytes after a week growth under TL without and with FR from

the same cultures as those used for dual RNA sequencing. In addition, reads were obtained from

sporophytes grown under TL with FR on medium with 2 mM NH_4NO_3, and from sporophytes without *N.*

azollae on medium with 2 mM NH_4NO_3. All samples were collected two hours into the 16 h light period.

The sRNA sequencing library preparation chemistry did not distinguish pro- from eukaryote sRNA and

therefore was inherently a dual profiling method. Reads obtained were aligned to concatenated genomes

which yielded small RNA profiles characteristic for the fern nucleus, chloroplast and *N. azollae* (Fig. **4**; SI

Fig. S**5**).

The most abundant sRNA encoded in the fern nucleus were 21, 20 and 24 nt long, but the most diverse

were 24 nt long, consistent with abundant 21 nt miRNA discretely cut from a hair-pin precursor by dicer,

and the 24 nt siRNA generated by random processes (Fig. **4a**). The chloroplast most abundant sRNA was

also the most diverse at 28 nt, but the chloroplast have an unusual peak of sRNA of low diversity at 34 nt

(Fig. **4b**). The sRNA encoded in *N. azollae* are generally shorter than those of the chloroplast, yet, they

have a conspicuous peak of low diversity at 42 nt length, consistent in length with CRISPR RNA (Fig. **4c**).

Conserved miRNA in *A. filiculoides* include miR172 and 396, but only miR529 and 535 increased in

response to FR

Fern genome-encoded reads of 20-25 nt length were used for miRNA predictions allowing a hairpin of

maximally 500 bp. Loci predicted were then verified against miRNA criteria (Axtell & Meyers, 2018) as, for

example, in the case of the miRNA 172 locus Azfi_s0021g15800 (Fig. **5**): the locus was expressed as a pre-

miRNA (Fig. **5a**, RNAseq all conditions) and both the miRNA and miRNA* are found in replicate samples

and cut precisely with two base overhangs (Fig. **5b,c**, sRNAseq). The miR172 reads were derived from the

fern since they were also found in sporophytes without *N. azollae* (Fig. **5,** RNAseq -*N. azollae*). The locus

encoded a stable long hairpin which when folded released -65.76 kcal mol^{-1} at 37°C. MiR172 targets in

seed plants include the AP2/TOE transcription factors, and alignment of the target sites to all annotated

AP2/TOE transcripts from *A. filiculoides* revealed target sites for the AzfimiR172 (SI Fig. S**6**). The miR156a

locus was similarly verified (SI Fig. S**7**).

Mature miRNA were compared in miRbase (Kozomara *et al.*, 2019) which identified 11 of the conserved

miRNA families of 21 nt in the *A. filiculoides* genome and thus in the fern lineage (Fig. **6a**). The list included

miR396, as well as miR172, not yet confirmed in other seed-free plant lineages. MiRNA predictions further

uncovered a number of novel miRNA (miRNF).

Differential expression analysis of 20-22 nt reads exhibited more dispersion than the dualRNAseq of long

RNA, yet, it reliably identified sRNA with low Padj values when comparing ferns under TL with and without

FR (SI Fig. S**5d,e**). Reads from miR529c&d and miR535 increased (Padj < 0.1) whilst those of miR159,

miR319 and miRNF4 decreased (Padj <0.16) in sporophytes on TL with FR compared to TL light only (Fig.

6b). We conclude that the miRNA156/miR172 known to reciprocally control flowering are not altered in

the ferns undergoing transition to RD under TL with FR.

miR319/GAMYB is responsive in sporophytes induced by FR

miRNA targets were predicted combining psRNATarget (Dai *et al.*, 2018) and TargetFinder (Srivastava *et*

al., 2014) then annotated with Mercator (Lohse *et al.*, 2014; Schwacke *et al.*, 2019). Predictions were

consistent with analyses in other land plant lineages (SI Table S**6**): miR172 targeted AP2/TOE-like

transcription factors and miR156, 529 and miR319 SPL-like proteins.

Transcripts of neither SPL nor AP2/TOE-like targets, however, were significantly altered in sporophytes

comparing TL with and without FR; this could be due to poor Azfi_vs1 annotation and incomplete

prediction of miR targets (SI Table S**6**, F vs T Padj). Nevertheless, consistent with an increased miR529c&d

the SPL from the Azfi_s0173.g055767 transcript is reduced (log2 0.74 fold) but with low significance (Padj

of 0.356). In contrast, transcripts of two GAMYB (Azfi_s0004.g008455 and Azfi_s0021.g015882) increased

up to 4 fold in sporophytes with decreased miR319 on TL with FR. The location of mir319 binding was

identified for the GAMYB encoded in Azfi_s0004.g008455 (SI Fig. S**8**).

Phylogenetic analyses revealed that the GAMYB targeted by miR319 were on the same leaf as the

GAMYB33 from *Arabidopsis thaliana* and were placed next to each other probably because they arose

from gene duplication (Fig. **7**). This is consistent with the whole genome duplication observed at the base

of *Azolla* evolution (Li *et al.*, 2018). We conclude, therefore, that miR319/GAMYB is responsive in

sporophytes undergoing transition to RD caused by FR.

DISCUSSION (1574)

Far-red light induced stem elongation and reproductive development in ferns, an analogy to

angiosperms?

The responses resemble a shade-avoidance syndrome but signal transduction pathways that mediate it in

ferns remain largely uncharacterized (Inoue *et al.*, 2017). Moreover, pathways causing shade response

components differ as they use alternative phytochromes and interacting factors (Possart *et al.*, 2014; Xie

et al., 2020). Unlike *A. filiculoides*, Arabidopsis will transition to RD in the TL used for indoors cultivation

over the past decades. Most studies on the transition to RD, therefore, were done with red-light gated

plants, i.e. conditions leading to artefactual peaking of CONSTANS protein accumulation and florigen (FT)

expression (Song *et al.*, 2018). Arabidopsis as well as many other seed plants flower early when in ratios

of red to far-red light approaching one, the ratio encountered in day light, compared to TL. Far-red light-

dependent flowering appeared mediated by PHYB and D, eventually leading to SOC1 expression in the

meristem (Halliday *et al.*, 1994; Aukerman *et al.*, 1997; Lazaro *et al.*, 2015). ft mutant Arabidopsis,

however, flowered earlier under incandescent light (Martinez-Zapater & Somerville, 1990) or in TL with

far-red LED compared to in TL (Schluepmann *et al.,* unpublished) suggesting that FT is not solely involved.

Signaling was also reported via PHYB-regulated PIF4 protein complexes that bind the promoters of

miRNA156/172 directly with interference from PHYA- regulated TF (Sánchez-Retuerta *et al.*, 2018; Sun *et*

al., 2018; Xie *et al.*, 2020). In spite of the apparent commonalities and the fern lineage representing the

sister lineage of seed plants, RD pathways from Arabidopsis may not operate in *Azolla*: the highly induced

MIKC[C] turned out a paralogue of SOC1, and miR156/172 steady states were unaltered by the far-red

triggered transition to RD in *A. filiculoides* (Fig. **6b**; SI Table S**6**).

PHYB in Arabidopsis has been shown to sense temperature by thermal reversion of the Pfr to fr state,

explaining why many responses to light quality can be tweaked by temperature regimes, in seed plants

(Legris *et al.*, 2017; Klose *et al.*, 2019). Phytochromes in ferns radiated separately from those of seed

plants, but have a similar domain structure (Li *et al.*, 2015). Temperature signaling by phytochromes has

not been researched in ferns. Initiation of fern, liverwort and bryophyte sporangia, however, is dependent

on temperature and photoperiod and differed with each species analyzed (Labouriau, 1958, Benson-

Evans, 1961; Nishihama *et al.*, 2015). Moreover, thermal reversion of phytochromes from some

cyanobacteria was described i*n vitro* as early as 1997 (Yeh *et al.*, 1997), but any effect on signal

transduction *in vivo* was not researched. Even though far-red LED did not induce sporocarps in the *Azolla*

strain Anzali, differing temperature regimes, therefore, may still work. Morphological/physiological

differences and phylogenetic analyses placed the Anzali strain closely to strains of a different species

collected in Uruguay/Southern Brazil which did not cluster with the representative *A. caroliniana* species.

Position of the Anzali strain was consistent with observations of distinct *Azolla* ferns in the Anzali lagoon

(Farahpour-Haghani *et al.*, 2017), and suggests that not only *A. filiculoides* may have been released as

nitrogen biofertilizer.

Whether analogous or homologous, repression of reproductive transition under dominating red light may

reflect an ecological strategy: when no neighbors are close, there is sufficient space to prolong vegetative

development and no need for dispersal by RD. Consistently, density of the *A. filiculoides* canopy had an

additional impact on the number of sporocarps observed per gram plant (Fig. **1f**) that may stem from

alternative cues such as volatile organic compounds released by neighbors (Vicherová *et al.*, 2020). Taken

together, results signify that for the first time RD of the *Azolla* fern symbiosis is amenable to experimental

enquiry.

410 Far-red light responsive accumulation of transcripts and small RNA implicate miR529, key MIKC^C TF and

411 the miR319/GAMYB module in fern RD

The GAMYB clade arose before land-plants evolved and before GA signaling (Aya *et al.*, 2011; Bowman *et*

al., 2018, Jiang & Rao, 2020, Fig. **7**). *Azolla* GAMYB induced under FR, therefore, likely function in RD by

mediating cues perceived by the sporophyte into an ancestral gametophyte regulon. GAMYB from the

moss *Physcomitrella patens* were required for the formation of antheridiophores and spores but we do

not know whether the *Pp*GAMYB regulon plays a role in RD of the moss sporophytes. GAMYB in mosses

and lycophytes regulate expression of enzymes assigned to sporopollenin biosynthesis which are also

upregulated in *Azolla* sporocarps (Aya *et al.*, 2011; Brouwer *et al.,* 2014). *At*GAMYB are known to be

regulated by miR159 in Arabidopsis and by the related and more ancestral miR319 in seed-free plant

lineages (Achard *et al.*, 2004; Palatnik *et al.*, 2007). The *At*GAMYB/mir159 is part of the GA pathway

promoting flowering in Arabidopsis under short days (Millar *et al.*, 2019). Specifically, AtMYB33 was shown

to directly act on the promotors of both miR156 and AtSPL9 (Guo *et al.,* 2017). FR repressed miR159 and

miR319/increased *Azfi*GAMYB in *Azolla* were thus consistent with predicted Arabidopsis wiring. GA signal-

transduction evolved in lycophytes, however, and we have yet to find out how far-red light causes miR319

induction in TL-grown sporophytes. Or, rather, why red-light gated ferns have increased miR319.

Far-red induced RD in the haploid phase of the liverwort *Marchantia polymorpha* (Chiyoda *et al.*, 2008;

Kubota *et al.*, 2014; Yamaoka *et al.*, 2018; Tsuzuki *et al.*, 2019). The mechanism in these liverworts

involved the *Mp*miR529c/*Mp*SPL module which was necessary for development of reproductive branches.

*Mp*PHY and *Mp*PIF were required to mediate the response (Inoue *et al.*, 2019). Our analyses with *A.*

filiculoides reveal induction of AzfimiR529 under far-red light but only insignificant change in SPL targets.

The pathway may differ in the two plants because the specific MpmiR529c sequence involved in

Marchantia was not detected in our ferns.

Control over RD as it switched from the gametophyte, in gametophyte dominated seed-free plants, to the

diploid sporophyte in lycophytes, ferns and seed plants, also resulted from the evolution of MIKC^C

regulons specific for each plant lineage (Leebens-Mack *et al.*, 2019, SI Fig. S**4**). The MIKC^C clade of TF that

specify both the transition to RD in sporophytes, such as *At*SOC1 and *At*FLC, and the A,C, D and E homeotic functions in the inflorescence development of angiosperms are peculiar in that the TF radiated in each megaphyll plant lineage separately. The TF work as hetero-tetramers and may have evolved from an ancient tandem gene duplication through subsequent polyploidy events (Zhao *et al.*, 2017). Sepals are organs formed only in angiosperms. Consistently, TF with A functions specifying sepals occurred solely in angiosperms. MIKC[C] specifying organs that contain the reproductive structures in angiosperms (C,D and E homeotic genes), had direct homologues in gymnosperms, but not in ferns. Instead, ferns radiated a sister clade to that containing AtSOC1, and A, C, D and E MIKC[C], in which the *Azolla* MIKC[C] responsive to far-red light were placed. This fern clade comprises *Ceratopteris thalictoides* CMADS1 reported highly expressed in the sporangial initial (Hasebe *et al.*,1998). At the base of the radiating clade of modern MIKC[C] that specify modern organs of the sporophytes controlling the gametophyte development within, is the LAMB1 protein expressed specifically in sporogenic structures of lycophytes (Svensson *et al.*, 2000). Combined phylogenetic and RNAseq analyses, therefore, suggest that *Azolla* fern sporocarps are likely specified by two MIKC[C] TF from the fern sister-lineage to AtSOC1. The result further suggests that SOC1 control over flowering has its roots in the common ancestor of seed plants and ferns when a MIKC[C] regulon controlled the diploid to haploid phase transition.

Dual RNA sequencing to study coordinate development in assemblages of pro- and eukaryotic organisms

The dual RNAseq method could be applied more generally to study bacteria/plant assemblages common in the seed-free plant lineages, representing an important base of the "tangled tree" of lineage evolution (Quammen D 2018). For *Azolla* where co-evolution of *N. azollae* and fern host genomes demonstrated that fitness selection occurs at the level of the metagenome (Li *et al.*, 2018), it delivered foundational data and opens the way to studying the coordinate development of host and obligate symbiont. Efficient removal of the rRNA was key to the success of reading long RNA. Fern chloroplast rRNA removal could be improved still, using specific DNA probes integrated during synthesis of the stranded sequencing libraries. Transcripts from the fern's mitochondrial genome are likely in the data but we have yet to assemble the *A. filiculoides* mitochondrial genome to confirm this.

N. azollae extracts did not contain photosynthesis pigments absorbing/fluorescing in the far-red spectrum and the *N. azollae* genome did not contain psbA4/Chlf-S like proteins (data not shown), yet it contained phytochromes and cyanobacteriochromes capable of sensing far-red light as other filamentous cyanobacteria do (Wiltbank & Kehoe, 2019). The RNA profiling did not reveal whether the *N. azollae*

trigger fern RD since roles of the far-red responsive transcripts in *N. azollae*, particularly those encoding

the transporters, have yet to be studied. An approach could be to study these in free living filamentous

cyanobacteria amenable to genetic manipulation. Manipulation of the symbiont would, however, be

preferred. Tri-parental mating protocols may be effective on *N. azollae in situ* if combined with high-

frequency RNA-guided integration with Cas12d enzyme complexes (Strecker *et al.,* 2020). *N. azollae* Cas

genes were found to be mostly pseudogenes and CRISPR arrays appeared missing (Cai *et al.,* 2013),

suggesting that incoming cargo plasmid may not be destroyed by Cas complexes. Nevertheless, sRNA

profiles from Fig. **5** show discrete accumulation of 31 and 42 b sRNA of very low complexity and a

functional Cas6 splicing CRISPR precursor RNA was induced on TL with FR (SI Table S**4**, *N. azollae*). CRISPR

arrays may not have been recognized, we conclude, if they contained few repeats due to the sheltered

life-style inside the fern. Rendering *A. filiculoides* and *N. azollae* amenable to genetic manipulation will be

crucial to provide evidence for hypotheses generated by transcript profiling in the future.

Acknowledgements

NWO-ALW EPS grant (ALWGS.2016.020) supported LWD, NWO-TTW grant (Project 16294) supported EG.

We are grateful to Bruno Huettel for advice on dual RNA sequencing library preparation methods. We

thank Bas Dutilh and Berend Snel for access to the Utrecht University bioinformatic computing resources.

Author Contributions

Fern induction experiments were done by BST and PB, fern crosses and comparisons were done by EG,

BST and unnamed graduate students, phylogenetic analyses and miR discovery were carried out by LWD

and NR, sRNA and dual RNA profiling was done by HS. HS conceived the manuscript with all authors

contributing to editing it.

Reference list

Achard P, Herr A, Baulcombe DC, Harberd NP. **2004**. Modulation of floral development by a gibberellin-
regulatedmicroRNA. *Development* **131**: 3357–3365.

Anders S, Huber W. **2010**. Differential expression analysis for sequence count data. *Genome Biology* **11**:
R106.

Anders S, Pyl PT, Huber W. **2015**. HTSeq-a Python framework to work with high-throughput sequencing
data. *Bioinformatics* **31**: 166–169.

Aukerman MJ, Hirschfeld M, Wester L, Weaver M, Clack T, Amasino RM, Sharrock RA. **1997**. A deletion in the PHYD gene of the Arabidopsis Wassilewskija ecotype defines a role for phytochrome D in red/far-red light sensing. *The Plant Cell* **9**: 1317–1326.

Axtell MJ, Meyers BC. **2018**. Revisiting Criteria for Plant MicroRNA Annotation in the Era of Big Data. *The Plant Cell* **30**: 272–284.

Aya K, Hiwatashi Y, Kojima M, Sakakibara H, Ueguchi-Tanaka M, Hasebe M, Matsuoka M. **2011**. The Gibberellin perception system evolved to regulate a pre-existing GAMYB-mediated system during land plant evolution. *Nature Communications* **2**: 544.

Benson-Evans K. **1961**. Environmental Factors and Bryophytes. *Nature* **191**: 255–260.

Berruezo F, de Souza FSJ, Picca PI, Nemirovsky SI, Martínez Tosar L, Rivero M, Mentaberry AN, Zelada AM. **2017**. Sequencing of small RNAs of the fern Pleopeltis minima (Polypodiaceae) offers insight into the evolution of the microrna repertoire in land plants (T Unver, Ed.). *PLOS ONE* **12**: e0177573.

Bonnet E, Wuyts J, Rouze P, Van de Peer Y. **2004**. Evidence that microRNA precursors, unlike other non-coding RNAs, have lower folding free energies than random sequences. *Bioinformatics* **20**: 2911–2917.

Bowman JL, Kohchi T, Yamato KT, Jenkins J, Shu S, Ishizaki K, Yamaoka S, Nishihama R, Nakamura Y, Berger F, *et al.* 2017. Insights into Land Plant Evolution Garnered from the Marchantia polymorpha Genome. *Cell* **171**: 287-304.e15.

Brouwer P, Bräutigam A, Buijs VA, Tazelaar AOE, van der Werf A, Schlüter U, Reichart G-J, Bolger A, Usadel B, Weber APM, *et al.* 2017. Metabolic Adaptation, a Specialized Leaf Organ Structure and Vascular Responses to Diurnal N2 Fixation by Nostoc azollae Sustain the Astonishing Productivity of Azolla Ferns without Nitrogen Fertilizer. *Frontiers in Plant Science* **8**.

Brouwer P, Bräutigam A, Külahoglu C, Tazelaar AOE, Kurz S, Nierop KGJ, van der Werf A, Weber APM, Schluepmann H. **2014**. Azolla domestication towards a biobased economy? *New Phytologist* **202**: 1069–1082.

Brouwer P, Schluepmann H, Nierop KGJ, Elderson J, Bijl PK, van der Meer I, de Visser W, Reichart G-J, Smeekens S, van der Werf A. **2018**. Growing Azolla to produce sustainable protein feed: the effect of differing species and CO2 concentrations on biomass productivity and chemical composition. *Journal of the Science of Food and Agriculture* **98**: 4759–4768.

Chiyoda S, Ishizaki K, Kataoka H, Yamato KT, Kohchi T. **2008**. Direct transformation of the liverwort Marchantia polymorpha L. by particle bombardment using immature thalli developing from spores. *Plant Cell Reports* **27**: 1467–1473.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**: 1972-3.

Coulter JM, Barnes CR, Cowles HC. **1910**. *A Textbook of Botany for Colleges and Universities*. London: American Book Company.

Dai X, Zhuang Z, Zhao PX. **2018**. psRNATarget: a plant small RNA target analysis server (2017 release). *Nucleic Acids Research* **46**: W49–W54.

Dijkhuizen LW, Brouwer P, Bolhuis H, Reichart G-J, Koppers N, Huettel B, Bolger AM, Li F-W, Cheng S, Liu X, *et al.* **2018**. Is there foul play in the leaf pocket? The metagenome of floating fern Azolla reveals endophytes that do not fix N 2 but may denitrify. *New Phytologist* **217**: 453–466.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. **2013**. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

Farahpour-Haghani A, Hassanpour M, Alinia F, Nouri-Ganbalani G, Razmjou J, Agassiz D. **2017**. Water ferns Azolla spp. (Azollaceae) as new host plants for the small China-mark moth, Cataclysta lemnata (Linnaeus, 1758) (Lepidoptera, Crambidae, Acentropinae). *Nota Lepidopterologica* **40**: 1–13.

Gras DE, Vidal EA, Undurraga SF, Riveras E, Moreno S, Dominguez-Figueroa J, Alabadi D, Blázquez MA, Medina J, Gutiérrez RA. **2018**. SMZ/SNZ and gibberellin signaling are required for nitrate-elicited delay of flowering time in Arabidopsis thaliana. *Journal of Experimental Botany* **69**: 619–631.

Gruber AR, Lorenz R, Bernhart SH, Neubock R, Hofacker IL. **2008**. The Vienna RNA Websuite. *Nucleic Acids Research* **36**: W70–W74.

Güngör E, Brouwer P, Dijkhuizen LW, Shaffar D, Nierop K, de Vos R, Toraño J, van der Meer I, Schluepmann H. 2020. *Azolla* ferns testify: seed plants and ferns share a common ancestor for leucoanthocyanidin reductase enzymes. *New Phytologist in press.*

Guo C, Xu Y, Shi M, Lai Y, Wu X, Wang H, Zhu Z, Poethig RS, Wu G. **2017**. Repression of miR156 by miR159 Regulates the Timing of the Juvenile-to-Adult Transition in Arabidopsis. *The Plant Cell* **29**: 1293–1304.

Halliday KJ, Koornneef M, Whitelam GC. **1994**. Phytochrome B and at Least One Other Phytochrome Mediate the Accelerated Flowering Response of Arabidopsis thaliana L. to Low Red/Far-Red Ratio. *Plant Physiology* **104**: 1311–1315.

Hasebe M, Wen CK, Kato M, Banks JA. 1998. Characterization of MADS homeotic genes in the fern Ceratopteris richardii. *Proceedings of the National Academy of Sciences* **95**:6222-6227.

Hisanaga T, Okahashi K, Yamaoka S, Kajiwara T, Nishihama R, Shimamura M, Yamato KT, Bowman JL, Kohchi T, Nakajima K. **2019**. A cis -acting bidirectional transcription switch controls sexual dimorphism in the liverwort. *The EMBO Journal* **38**: e100240.

Hou S, Brenes-Álvarez M, Reimann V, Alkhnbashi OS, Backofen R, Muro-Pastor AM, Hess WR. **2019**. CRISPR-Cas systems in multicellular cyanobacteria. *RNA Biology* **16**: 518–529.

Hyun Y, Vincent C, Tilmes V, Bergonzi S, Kiefer C, Richter R, Martinez-Gallegos R, Severing E, Coupland G. **2019**. A regulatory circuit conferring varied flowering response to cold in annual and perennial plants. *Science* **363**: 409–412.

Inoue K, Nishihama R, Araki T, Kohchi T. **2019**. Reproductive Induction is a Far-Red High Irradiance Response that is Mediated by Phytochrome and PHYTOCHROME INTERACTING FACTOR in Marchantia polymorpha. *Plant and Cell Physiology* **60**: 1136–1145.

Jiang C-K, Rao G-Y. **2020**. Insights into the Diversification and Evolution of R2R3-MYB Transcription Factors in Plants. *Plant Physiology* **183**: 637–655.

Kaneko M, Inukai Y, Ueguchi-Tanaka M, Itoh H, Izawa T, Kobayashi Y, Hattori T, Miyao A, Hirochika H,
Ashikari M, *et al.* 2004. Loss-of-Function Mutations of the Rice GAMYB Gene Impair α-Amylase
Expression in Aleurone and Flower Development. *The Plant Cell* **16**: 33–44.

Katoh K, Rozewicki J, Yamada KD. **2019.** MAFFT online service: multiple sequence alignment, interactive
sequence choice and visualization. *Briefings in bioinformatics* **20**: 1160-6.

Klose C, Nagy F, Schäfer E. **2020**. Thermal Reversion of Plant Phytochromes. *Molecular Plant*.

Kozomara A, Birgaoanu M, Griffiths-Jones S. **2019**. miRBase: from microRNA sequences to function.
Nucleic Acids Research **47**: D155–D162.

Kuang Z, Wang Y, Li L, Yang X. **2019**. miRDeep-P2: accurate and fast analysis of the microRNA
transcriptome in plants (I Birol, Ed.). *Bioinformatics* **35**: 2521–2522.

Kubota A, Kita S, Ishizaki K, Nishihama R, Yamato KT, Kohchi T. **2014**. Co-option of a photoperiodic
growth-phase transition system during land plant evolution. *Nature Communications* **5**: 3668.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. **2004**. Versatile and
open software for comparing large genomes. *Genome biology* **5**: R12.

Labouriau LG. **1958**. Studies on the initiation of sporangia in ferns.

Leebens-Mack JH, Barker MS, Carpenter EJ, Deyholos MK, Gitzendanner MA, Graham SW, Grosse I, Li
Z, Melkonian M, Mirarab S, *et al.* 2019. One thousand plant transcriptomes and the phylogenomics of
green plants. *Nature* **574**: 679–685.

Legris M, Nieto C, Sellaro R, Prat S, Casal JJ. **2017**. Perception and signalling of light and temperature
cues in plants. *The Plant Journal* **90**: 683–697.

Li F-W, Brouwer P, Carretero-Paulet L, Cheng S, de Vries J, Delaux P-M, Eily A, Koppers N, Kuo L-Y, Li Z,
***et al.* 2018**. Fern genomes elucidate land plant evolution and cyanobacterial symbioses. *Nature Plants* **4**:
460–472.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. **2009**. The
Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Li F-W, Melkonian M, Rothfels CJ, Villarreal JC, Stevenson DW, Graham SW, Wong GK-S, Pryer KM,
Mathews S. **2015**. Phytochrome diversity in green plants and the origin of canonical plant
phytochromes. *Nature Communications* **6**: 7852.

Lohse M, Nagel A, Herter T, May P, Schroda M, Zrenner R, Tohge T, Fernie AR, Stitt M, Usadel B. **2014**.
Mercator: a fast and simple web server for genome scale functional annotation of plant sequence data.
Plant, Cell & Environment **37**: 1250–1258.

Love MI, Huber W, Anders S. **2014.** Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome biology* **15**: 550.

Lucero LE, Manavella PA, Gras DE, Ariel FD, Gonzalez DH. **2017**. Class I and Class II TCP Transcription
Factors Modulate SOC1-Dependent Flowering at Multiple Levels. *Molecular Plant* **10**: 1571–1574.

Madeira PT, Hill MP, Dray FA, Coetzee JA, Paterson ID, Tipping PW. **2016**. Molecular identification of

Azolla invasions in Africa: The Azolla specialist, Stenopelmus rufinasus proves to be an excellent taxonomist. *South African Journal of Botany* **105**: 299–305.

Martinez-Zapater JM, Somerville CR. **1990**. Effect of Light Quality and Vernalization on Late-Flowering Mutants of Arabidopsis thaliana. *Plant Physiology* **92**: 770–776.

Millar AA, Lohe A, Wong G. **2019**. Biology and Function of miR159 in Plants. *Plants* **8**: 255.

Nagalingum NS, Schneider H, Pryer KM. **2006**. Comparative Morphology of Reproductive Structures in Heterosporous Water Ferns and a Reevaluation of the Sporocarp. *International Journal of Plant Sciences* **167**: 805–815.

Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular biology and evolution* **32**: 268-74.

Nishihama R, Ishizaki K, Hosaka M, Matsuda Y, Kubota A, Kohchi T. **2015**. Phytochrome-mediated regulation of cell division and growth during regeneration and sporeling development in the liverwort Marchantia polymorpha. *Journal of Plant Research* **128**: 407–421.

Nobori T, Velásquez AC, Wu J, Kvitko BH, Kremer JM, Wang Y, He SY, Tsuda K. **2018**. Transcriptome landscape of a bacterial pathogen under plant immunity. *Proceedings of the National Academy of Sciences* **115**: E3055–E3064.

Olas JJ, Van Dingenen J, Abel C, Działo MA, Feil R, Krapp A, Schlereth A, Wahl V. **2019**. Nitrate acts at the Arabidopsis thaliana shoot apical meristem to regulate flowering time. *New Phytologist* **223**: 814–827.

Olm MR, Brown CT, Brooks B, Banfield JF. **2017**. dRep: a tool for fast and accurate genomic comparisons that enables improved genome recovery from metagenomes through de-replication. *The ISME Journal* **11**: 2864–2868.

Kar P, Mishra S, Singh D. **2002**. *Azolla* sporulation in response to application of some selected auxins and their combination with gibberellic acid. *Biology and Fertility of Soils* **35**: 314–319.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome biology* **5**:R12.

Palatnik JF, Wollmann H, Schommer C, Schwab R, Boisbouvier J, Rodriguez R, Warthmann N, Allen E, Dezulian T, Huson D, *et al.* 2007. Sequence and Expression Differences Underlie Functional Specialization of Arabidopsis MicroRNAs miR159 and miR319. *Developmental Cell* **13**: 115–125.

Perkins SK, Peters GA. **2006**. The Azolla-Anabaena symbiosis: endophyte continuity in the Azolla life-cycle is facilitated by epidermal trichomes †. *New Phytologist* **123**: 53–64.

Plackett ARG, Conway SJ, Hewett Hazelton KD, Rabbinowitsch EH, Langdale JA, Di Stilio VS. **2018**. LEAFY maintains apical stem cell activity during shoot development in the fern *Ceratopteris richardii*. *eLife* **7**.

Possart A, Hiltbrunner A. **2013**. An evolutionarily conserved signaling mechanism mediates far-red light responses in land plants. *The Plant Cell* **25**: 102–114.

Quammen D. 2018. The tangled tree: a radical new history of life. Simon & Schuster, New York.

Ran L, Larsson J, Vigil-Stenman T, Nylander JAA, Ininbergs K, Zheng W-W, Lapidus A, Lowry S,
Haselkorn R, Bergman B. **2010**. Genome erosion in a nitrogen-fixing vertically transmitted
endosymbiotic multicellular cyanobacterium (N Ahmed, Ed.). *PLoS ONE* **5**: e11486.

Rolauffs S, Fackendahl P, Sahm J, Fiene G, Hoecker U. **2012**. Arabidopsis COP1 and SPA genes are
essential for plant elongation but not for acceleration of flowering time in response to a low red light to
far-red light ratio. *Plant Physiology* **160**: 2015–2027.

Sánchez-Retuerta C, Suaréz-López P, Henriques R. **2018**. Under a New Light: regulation of light-
dependent pathways by non-coding RNAs. *Frontiers in Plant Science* **9**.

Sayou C, Nanao MH, Jamin M, Posé D, Thévenon E, Grégoire L, Tichtinsky G, Denay G, Ott F, Peirats
Llobet M, *et al.* 2016. A SAM oligomerization domain shapes the genomic binding landscape of the
LEAFY transcription factor. *Nature Communications* **7**: 11222.

Schwacke R, Ponce-Soto GY, Krause K, Bolger AM, Arsova B, Hallab A, Gruden K, Stitt M, Bolger ME,
Usadel B. **2019**. MapMan4: A Refined Protein Classification and Annotation Framework Applicable to
Multi-Omics Data Analysis. *Molecular Plant* **12**: 879–892.

Sessa EB, Der JP. **2016**. Evolutionary genomics of ferns and lycophytes. In: Advances in Botanical
Research. 215–254.

Shimmura S, Nozoe M, Kitora S, Kin S, Matsutani S, Ishizaki Y, Nakahira Y, Shiina T. **2017**. Comparative
Analysis of Chloroplast psbD Promoters in Terrestrial Plants. *Frontiers in Plant Science* **8**.

Song YH, Kubota A, Kwon MS, Covington MF, Lee N, Taagen ER, Laboy Cintrón D, Hwang DY, Akiyama
R, Hodge SK, *et al.* 2018. Molecular basis of flowering under natural long-day conditions in Arabidopsis.
Nature Plants **4**: 824–835.

Srivastava PK, Moturu T, Pandey P, Baldwin IT, Pandey SP. **2014**. A comparison of performance of plant
miRNA target prediction tools and the characterization of features for genome-wide target prediction.
BMC Genomics **15**: 348.

Strecker J, Ladha A, Gardner Z, Schmid-Burgk JL, Makarova KS, Koonin E V., Zhang F. **2019**. RNA-guided
DNA insertion with CRISPR-associated transposases. *Science* **365**: 48–53.

Sun Z, Li M, Zhou Y, Guo T, Liu Y, Zhang H, Fang Y. **2018**. Coordinated regulation of Arabidopsis
microRNA biogenesis and red light signaling through Dicer-like 1 and phytochrome-interacting factor 4
(L-J Qu, Ed.). *PLOS Genetics* **14**: e1007247.

Svensson ME, Johannesson H, Engström P. **2000**. The LAMB1 gene from the clubmoss, Lycopodium
annotinum, is a divergent MADS-box gene, expressed specifically in sporogenic structures. *Gene* **253**:
31–43.

Theißen G, Rümpler F, Gramzow L. **2018**. Array of MADS-Box Genes: Facilitator for Rapid Adaptation?
Trends in Plant Science **23**: 563–576.

Tsuzuki M, Futagami K, Shimamura M, Inoue C, Kunimoto K, Oogami T, Tomita Y, Inoue K, Kohchi T,
Yamaoka S, *et al.* 2019. An early arising role of the MicroRNA156/529-SPL module in reproductive
development revealed by the liverwort marchantia polymorpha. *Current Biology* **29**: 3307-3314.e5.

Vasco A, Smalls TL, Graham SW, Cooper ED, Wong GK-S, Stevenson DW, Moran RC, Ambrose BA. **2016**.
Challenging the paradigms of leaf evolution: Class III HD-Zips in ferns and lycophytes. *New Phytologist*
212: 745–758.

Vicherová E, Glinwood R, Hájek T, Šmilauer P, Ninkovic V. **2020**. Bryophytes can recognize their
neighbours through volatile organic compounds. *Scientific Reports* **10**: 7405.

de Vries A, Ripley BD. **2016**. Package 'ggdendro': Create Dendrograms and Tree Diagrams Using
'ggplot2' (R package). *R package version 0.1-20*.

Wickham H. **2011**. ggplot2. *Wiley Interdisciplinary Reviews: Computational Statistics* **3**: 180–185.

Wiltbank LB, Kehoe DM. **2019**. Diverse light responses of cyanobacteria mediated by phytochrome
superfamily photoreceptors. *Nature Reviews Microbiology* **17**: 37–50.

Xie Y, Zhou Q, Zhao Y, Li Q, Liu Y, Ma M, Wang B, Shen R, Zheng Z, Wang H. 2020. FHY3 and FAR1
Integrate Light Signals with the miR156-SPL Module-Mediated Aging Pathway to Regulate Arabidopsis
Flowering. *Molecular Plant* **13**:483-98.

Yamaoka S, Nishihama R, Yoshitake Y, Ishida S, Inoue K, Saito M, Okahashi K, Bao H, Nishida H,
Yamaguchi K, *et al.* **2018**. Generative Cell Specification Requires Transcription Factors Evolutionarily
Conserved in Land Plants. *Current Biology* **28**: 479-486.e5.

Yeh K. **1997**. A Cyanobacterial Phytochrome Two-Component Light Sensory System. *Science* **277**: 1505–
1508.

You C, Cui J, Wang H, Qi X, Kuo L-Y, Ma H, Gao L, Mo B, Chen X. **2017**. Conservation and divergence of
small RNA pathways and microRNAs in land plants. *Genome Biology* **18**: 158.

Zhao T, Holmer R, de Bruijn S, Angenent GC, van den Burg HA, Schranz ME. **2017**. Phylogenomic
Synteny Network Analysis of MADS-Box Transcription Factor Genes Reveals Lineage-Specific
Transpositions, Ancient Tandem Duplications, and Deep Positional Conservation. *The Plant Cell* **29**:
1278–1292.

Zhu Q, Fisher S, Shallcross J, Kim J. **2016**. VERSE: a versatile and efficient RNA-Seq read counting tool.

Table 1 Sporeling counts obtained from random crosses of *A. filiculoides* strains collected in The
Netherlands and Spain (Gran Canaria).

Megaspores:	Galgenwaard	Krommerijn	Hoogwoud	Gran Canaria	Den Bosch	Nijmegen	Nieuwebrug	Groningen
Massulae :								
Galgenwaard	>1000	17		1		13		2
Krommerijn	1				16		2	
Hoogwoud			20					2
Gran Canaria*	17				>40	1		
Den Bosch		9		1				
Nijmegen		12		33			2	
Nieuwebrug		6	3			22		
Groningen			1				5	

709 *crosses involving megaspores from Gran Canaria reproducibly germinated late, generally after 5 weeks
710 instead of 2-3 weeks.

715　**Table 2** Transcription factors with highest changes of read counts in sporophytes after a week on TL with

716　versus without far-red LED.

A. filiculoides locus	baseMean	Log2Fold Change	DESeq2 padj	Mercator 4.0 annotation	Closest Arabidopsis Homolog
Azfi_s0028.g024032	**9**	**7.22**	**0.001**	**15.5.14 MADS/AGL**	**AGL20/SOC1, MIKCᶜ**
Azfi_s0113.g045874	155	6.41	0.002	15.5.2 R2R3MYB	AtMYB 32 (AT4G34990.1), R2R3MYB VIII-E*
Azfi_s0083.g038807	59	5.75	0	15.5.32 Basic Helix-Loop-Helix	EDA33, IND1, INDEHISCENT (AT4G00120.1)
Azfi_s0003.g007560	8	5.27	0.011	15.5.32 Basic Helix-Loop-Helix	FIT1 (regulates iron transport)
Azfi_s0003.g007559	34	3.49	0	15.5.32 Basic Helix-Loop-Helix	FIT1 (regulates iron transport)
Azfi_s0096.g043715	31	3.13	0.001	15.5.7 DREB subfamily A-2 of ERF/AP2	AT5G18450.1
Azfi_s0003.g007710	1778	2.82	0	**15.5.14 MADS/AGL**	**AGL6, MIKCᶜ**
Azfi_s0015.g013719	111	2.57	0	15.5.2 G2-like, GARP	HHO2, NIGT1.2
Azfi_s0112.g045798	66	2.2	0	15.5.51.1 NF-Y component NF-YA	AT5G12840.4
Azfi_s0014.g013539	105	2.07	0.005	**15.5.2 R2R3MYB**	**LOF2, R2R3MYB V***
Azfi_s0015.g014012	165	1.99	0.026	15.5.17 NAC domain	NAC025
Azfi_s0004.g008455	**103**	**1.93**	**0.003**	**15.5.2 R2R3MYB**	**miRNA319 controlled GAMYB 33, R2R3MYB VII***
Azfi_s0132.g049213	63	-2.35	0.004	15.5.32 Basic Helix-Loop-Helix	No hits
Azfi_s0182.g056462	10	-4.46	0.049	15.5.3 HD-ZIP I/II	HB16

718　*R2R3 MYB classification according to Jiang & Rao, 2020.

 Figure legends

Figure 1 Far-red light and nitrogen affect the reproductive phase transition in *A. filiculoides*. (a)
Vegetatively growing ferns. (b) Ferns in the reproductive phase with micro (mi) and megaspores (me). (c)
growth habit under TL and TL with far-red LED (FR). (d) A mature sporophyte and (e) its schematic
representation: a root is formed at each branch point, and sporocarp pairs are found close-by on the
branch side of the branch point but not at all branch points. (f) Sporulation frequencies at increasing plant
density (41.7, 128.8 and 208.3 g m^{-2} dry weight) and with increasing red to far-red light ratios (FR0-FR4).
The red to far-red ratios were 19.13 (FR0), 1.16 (FR1), 0.63 (FR2), 0.50 (FR3) and 0.31 (FR4); to obtain the
FR4 ratio, the TL intensity was halved. Each data point represents the average of three independent
continuous growth replicates. (g) Induction of sporulation without nitrogen (-N) or with 2 mM NH_4NO_3
(+N) in the medium, data are averages from three replicates with standard deviations.

Figure 2 Chloroplast marker regions in strains from the Netherlands, Spain and Iran, and the rRNA
intergenic regions of the *A. filiculoides* genome. Phylogenetic trees of chloroplast regions trnL-trnF (a) and
trnG-trnR (b). The maximum likelihood trees were bootstrapped 1000x. All bootstrap values >0.70 are
displayed. In bold: sequences from accessions in this study labelled after their collection site and
sequences from the differing species extracted from genome shot-gun sequencing (Li *et al.,* 2018). In
regular type: sequences from the Madeira et al. (2016) study. (c), Sequence logo of all ITS1 gene regions
in the *A. filiculoides* genome (Li *et al.,* 2018). The gene regions were extracted and aligned (MEGA 7) then
used for *in-silico* PCR. Results were visualized using WebLogo (vs 2.8.2 Crooks *et al.,* 2004): at each position
of the ITS, the height of the stack indicates the sequence conservation, while the height of symbols within
the stack indicates the relative frequency of each base.

Figure 3 Profiling eukaryotic and prokaryotic RNA simultaneously in *A. filiculoides* (dual RNAseq). (a)
Similarity of the *N. azollae* 0708 genome and Metagenome Assembled Genomes (MAGs) from *N. azollae*
strains in *Azolla* species. Average Nucleotide Identity (ANI) was calculated with the dRep implementation
of nucmer using the ANImf preset (Kurtz *et al.*, 2004; Olm *et al.*, 2017). Pairwise average ANI is shown in
an UPGMA dendrogram (de Vries & Ripley, 2016; Wickham, 2011). Aligned MAGS covered over 90% of
the reference *N. azollae* 0708. (b) Proportion (percent) of uniquely aligned PE reads (dark color) and PE
reads aligning at multiple loci (light color) on the genomes of fern (green), fern chloroplast (purple) and
N. azollae 0708 (red). PE reads were from samples of sporophytes grown for a week on TL-light (replicates
T1, T2, T3) or TL light with far-red LED (replicates F1, F2, F3). STAR-aligner was used in PE mode. (c) Yields
of PE uniquely mapped compared with PE read counts on features and PE reads that STAR aligned

ambiguously to several features. Features were specified by current general features files (GFF) of each

of the fern (green), chloroplast (purple) and *N.azollae* 0708 (red) genomes.

Figure 4 sRNA size distribution and diversity for each of the genomes in the *A. filiculoides* symbiosis. sRNA

reads were mapped with 0 mismatches, no overhang, to the concatenated genomes; bam2fasta was then

used to extract sRNA sequences aligning per genome, the fasta files obtained were then collapsed to

compute read counts for each sRNA, the lists obtained were then joined to a read count matrix imposing

at least 1 read per sRNA in each of the three replicate sample types: f (far-red), t (no far-red) and fn (far-

red and nitrogen in the medium). sRNA counts were then normalized to the total read counts per sample

for that genome. Fern nucleus. Chloroplast, included samples from sporophytes without cyanobacteria in

the analysis. *N. azollae*, read count files joined imposing at least 10 read per sRNA in every sample

Figure 5 The miRNA 172α locus in *A.filiculoides*. Alignments at the Azfi_miRNA172α locus are visualized

using the Integrative Genomics Viewer (Thorvaldsdóttir *et al.,* 2013); S, read summary; i, individual reads

(a) Reads from pre-miRNA at the miR172α locus. RNAseq all conditions, all reads pooled from the diel

RNAseq experiment in (Brouwer *et al.,* 2017) . (b) Reads of the miR172 in sRNA libraries including those

of ferns without *N.azollae* (sRNAseq-*N.azollae*), ferns grown with and without far-red LED (sRNAseq+FR,

sRNAseq TL-FR). (c) Reads of the miR172*. (d) Azfi_miRNA172α hairpin folded using Vienna RNAfold

(Gruber *et al.,* 2008; predicts -104,07 kcal mol^{-1} at 22 ^{o}C upon folding); miRNA (purple) and miRNA* (pink)

stem arms are shown with 3'overhangs.

Figure 6 Conserved and far-red light responsive miRNA in the fern *A. filiculoides*. (a) conserved miRNA in

the *A.filiculoides* fern compared to other land plant lineages. miRNA from the fern were predicted

computationally by miRDEEP-P2 using sRNA seq reads then curated manually, the remainder of the table

was from Axtell & Meyers, 2018. (b), miRNA with altered steady state in response to far-red light. Shown

are average read counts per million 20-22 nt reads in samples of sporophytes after one week on TL light

with far-red LED (f) or without (t); raw read counts were submitted to DESeq2 for statistical analyses

yielding P-adjusted values: * indicates Padj <0.1; ^ indicates Padj <0.16.

Figure 7 R2R3MYB phylogenetic analyses and fold change comparing sporophytes with and without FR

LED. Sequences were extracted from the genome browsers of each species, aligned with

MAFFT linsi or einsi, then trimmed with trimAL. Phylogenetic inferences were computed with IQtree and

its internal model fitter; non-parametric bootstrap values are shown. *Chara autralis* (Ca), *Chara braunii*

(Cb), *Chara leptospora* (Cl), *Marchantia polymorpha* (Mp), *Selaginella moellendorfii* (Sm), *Azolla*

filiculoides (Azfi),), *Salvinia cuculata* (Sc), *Picea abies* (Pa), *Amborella trichopoda* (Atri), *Arabidopsis*

thaliana (At). Sequence names are color coded after the R2R3 MYB subfamilies defined in Jiang & Rao

(2020). Fold change in response to FR was calculated by DESeq2, green stars mark significant changes with

Padj <0.1.

Figure 1 Far-red light and nitrogen affect the reproductive phase transition in *A. filiculoides*. (a)
Vegetatively growing ferns. (b) Ferns in the reproductive phase with micro (mi) and megaspores (me). (c)
growth habit under TL and TL with far-red LED (FR). (d) A mature sporophyte and (e) its schematic
representation: a root is formed at each branch point, and sporocarp pairs are found close-by on the
branch side of the branch point but not at all branch points. (f) Sporulation frequencies at increasing plant
density (41.7, 128.8 and 208.3 g m^{-2} dry weight) and with increasing red to far-red light ratios (FR0-FR4).
The red to far-red ratios were 19.13 (FR0), 1.16 (FR1), 0.63 (FR2), 0.50 (FR3) and 0.31 (FR4); to obtain the
FR4 ratio, the TL intensity was halved. Each data point represents the average of three independent
continuous growth replicates. (g) Induction of sporulation without nitrogen (-N) or with 2 mM NH$_4$NO$_3$
(+N) in the medium, data are averages from three replicates with standard deviations.

797 **Figure 2** Chloroplast marker regions in strains from the Netherlands, Spain and Iran, and the rRNA

798 intergenic regions of the *A. filiculoides* genome. Phylogenetic trees of chloroplast regions trnL-trnF (a) and

799 trnG-trnR (b). The maximum likelihood trees were bootstrapped 1000x. All bootstrap values >0.70 are

800 displayed. In bold: sequences from accessions in this study labelled after their collection site and

801 sequences from the differing species extracted from genome shot-gun sequencing (Li *et al.,* 2018). In

802 regular type: sequences from Madeira et al. (2016)*.* (c) Sequence logo of all ITS1 gene regions in the *A.*

803 *filiculoides* genome (Li *et al.,* 2018). The ITS1 sequences were extracted and aligned, then alignments

804 visualized using WebLogo (vs 2.8.2 Crooks *et al.,* 2004): at each position of the ITS1, the height of the stack

805 indicates the sequence conservation, while the height of symbols within the stack indicates the relative

806 frequency of each base.

Figure 3 Profiling eukaryotic and prokaryotic RNA simultaneously in *A. filiculoides* (dual RNAseq). (a)

Similarity of the *N. azollae* 0708 genome and Metagenome Assembled Genomes (MAGs) from *N. azollae*

strains in *Azolla* species. Average Nucleotide Identity (ANI) was calculated with the dRep implementation

of nucmer using the ANImf preset (Kurtz *et al.*, 2004; Olm *et al.*, 2017). Pairwise average ANI is shown in

an UPGMA dendrogram (de Vries & Ripley, 2016; Wickham, 2011). Aligned MAGS covered over 90% of

the reference *N. azollae* 0708. (b) Proportion (percent) of uniquely aligned PE reads (dark color) and PE

reads aligning at multiple loci (light color) on the genomes of fern (green), fern chloroplast (purple) and

N. azollae 0708 (red). PE reads were from samples of sporophytes grown for a week on TL-light (replicates

T1, T2, T3) or TL light with far-red LED (replicates F1, F2, F3). STAR-aligner was used in PE mode. (c) Yields

of PE uniquely mapped compared with PE read counts on features and PE reads that STAR aligned

ambiguously to several features. Features were specified by current general features files (GFF) of each

of the fern (green), chloroplast (purple) and *N.azollae* 0708 (red) genomes.

 Figure 4 sRNA size distribution and diversity for each of the genomes in the *A. filiculoides* symbiosis. sRNA

reads were mapped, imposing zero mismatches and no overhang, to the concatenated genomes;

bam2fasta was then used to extract sRNA sequences aligning per genome, the fasta files obtained were

then collapsed to compute read counts for each sRNA, the lists obtained were then joined to a read count

matrix imposing at least 1 read per sRNA in each of the three replicate sample types: f (far-red), t (no far-

red) and fn (far-red and nitrogen in the medium). sRNA counts were then normalized to the total read

counts per sample for that genome. Fern nucleus. Chloroplast, included samples from sporophytes

without cyanobacteria in the analysis. *N. azollae*, read count files joined imposing at least 10 read per

sRNA in e

Figure 5 The miRNA 172α locus in *A.filiculoides*. Alignments at the Azfi_miRNA172α locus were visualized

using the Integrative Genomics Viewer (Thorvaldsdóttir *et al.,* 2013); S, read summary; i, individual reads

(a) Reads from pre-miRNA at the miR172α locus. RNAseq all conditions, all reads pooled from the diel

RNAseq experiment in (Brouwer *et al.,* 2017) . (b) Reads of the miR172 in sRNA libraries including those

of ferns without *N.azollae* (sRNAseq-*N.azollae*), ferns grown with and without far-red LED (sRNAseq+FR,

sRNAseq TL-FR). (c) Reads of the miR172*. (d) Azfi_miRNA172α hairpin folded using Vienna RNAfold

(Gruber *et al.,* 2008; predicts -104,07 kcal mol^{-1} at 22 $^{\circ}$C upon folding); miRNA (purple) and miRNA* (pink)

stem arms are shown with 3'overhangs.

Figure 6 Conserved and far-red light responsive miRNA in the fern *A. filiculoides*. (a) Conserved miRNA in

the *A.filiculoides* fern compared to other land plant lineages. miRNA from the fern were predicted

computationally by miRDEEP-P2 using sRNA seq reads then curated manually, the remainder of the table

was from Axtell & Meyers, 2018. (b) miRNA in *A. filiculoides* sporophytes responsive to far-red light. Shown

are average read counts per million 20-22 nt reads in samples of sporophytes after one week on TL light

with far-red LED (sRNA +FR) or without (sRNA -FR); raw read counts were submitted to DESeq2 for

statistical analyses yielding P-adjusted values: * indicates Padj <0.1; ^ indicates Padj <0.16.

Figure 7 *Azolla* R2R3MYB phylogenetic analysis and response to far-red light (FR). Sequences extracted from the genome browsers of each species were aligned with MAFFT linsi or einsi (Katoh *et al.*, 2019), then trimmed with trimAL (Capella-Gutiérrez *et al.*, 2009). Phylogenetic inferences were computed with IQtree (Nguyen *et al.*, 2015) and its internal model fitter; non-parametric bootstrap values are shown. Zygnematophyceae: *Spirogyra sp*. Charophyceae: *Chara autralis* (*Chaus*), *Chara braunii* (*Chbrau*), *Chara leptospora* (*Chlep*). Bryophyte: *Marchantia polymorpha* (*Mapoly*). Lycophyte: *Selaginella moellendorfii* (*Selmo*). Pteridophytes: *Azolla filiculoides* (Azfi), *Salvinia cuculata* (Sacu). Gymnosperm: *Picea abies* (Piabi). Angiosperms: *Amborella trichopoda* (Amtr), *Arabidopsis thaliana* (Arath). Sequence names are color coded after the R2R3 MYB subfamilies defined in Jiang & Rao (2020). Foldchange in response to FR was calculated by DESeq2 (Love *et al*., 2014); green stars mark significant changes of Padj <0.1.