LIQA: Long-read Isoform Quantification and Analysis

Yu Hu[1], Li Fang[1], Xuelian Chen[2], Jiang F. Zhong[2], Mingyao Li[3], Kai Wang[1,4]*

[1] Raymond G. Perelman Center for Cellular and Molecular Therapeutics, Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA

[2] Department of Otolaryngology, Keck School of Medicine, University of Southern California, CA 90033, USA

[3] Department of Biostatistics, Epidemiology and Informatics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

[4] Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

* To whom correspondence should be addressed. Email: wangk@email.chop.edu

Abstract

Long-read RNA sequencing (RNA-seq) technologies have made it possible to sequence full-length transcripts, facilitating the exploration of isoform-specific gene expression (isoform relative abundance and isoform-level TPM) over conventional short-read RNA-seq. However, long-read RNA-seq suffers from high per-base error rate, presence of chimeric reads or alternative alignments, and other biases, which require different analysis methods than short-read RNA-seq. Here we present LIQA (Long-read Isoform Quantification and Analysis), an Expectation-Maximization based statistical method to quantify isoform expression and detect differential alternative splicing (DAS) events using long-read RNA-seq data. Rather than summarizing isoform-specific read counts directly as done in short-read methods, LIQA incorporates base-pair quality score and isoform-specific read length information to assign different weights across reads, which reflects alignment confidence. Moreover, LIQA can detect DAS events between conditions using isoform usage estimates. We evaluated LIQA's performance on simulated data and demonstrated that it outperforms other approaches in characterizing isoforms with low read coverage and in detecting DAS events between two groups. We also generated one direct mRNA sequencing dataset and one cDNA sequencing dataset using the Oxford Nanopore long-read platform, both with paired short-read RNA-seq data and qPCR data on selected genes, and we demonstrated that LIQA performs well in isoform discovery and quantification. Finally, we evaluated LIQA on a PacBio dataset on esophageal squamous epithelial cells, and demonstrated that LIQA recovered DAS events that failed to be detected in short-read data. In summary, LIQA leverages the power of long-read RNA-seq and achieves higher accuracy in estimating isoform abundance than existing approaches, especially for isoforms with low coverage and biased read distribution. LIQA is freely available at https://github.com/WGLab/LIQA.

Introduction

RNA splicing is a major mechanism for generating transcriptomic variations, and mis-regulation of splicing is associated with a large array of human diseases caused by hereditary and somatic mutations[1-5]. Over the past decade, RNA sequencing (RNA-seq) has revolutionized transcriptomics studies and facilitated the characterization and understanding of transcriptomic variations in an unbiased fashion. With RNA-seq, we can quantitatively measure isoform-specific gene expression, discover novel and unique transcript isoform signature and detect differential alternative splicing (DAS) events[6-8]. Until now, short-read RNA-seq has been the method of choice for transcriptomics studies due to its high coverage and single nucleotide resolution[8]. However, due to limited read length, it is difficult to accurately characterize transcripts using short reads, as 81% of isoforms have length greater than 500bp in the GENCODE annotation (median = 1,543 bp and mean = 2,121 bp). This fragmented sequencing of the RNA/cDNA molecules results in biases and has become a barrier for short reads to be correctly mapped to the reference genome, which is crucial for gene or isoform expression estimation and novel or unique isoform detection. As a consequence, transcriptome profiling using short-read RNA-seq is highly biased by read coverage heterogeneity across isoforms. To tackle these challenges, a number of computational tools, including RSEM[9], eXpress[10], TIGAR2[11], Salmon[12], Sailfish[13], Kallisto[14], Cufflinks[15], CEM[16], PennSeq[17], IsoEM[18], and RD[19], have been developed to quantify isoform expression from short-read RNA-seq data, but different bias correction algorithms can result in conflicting estimates[17]. Overall, quantifying isoform expression using fragmented short reads remains challenging, especially at complex gene loci[20, 21].

In recent years, long-read RNA sequencing has gained popularity due to its potential to overcome the above-mentioned issues when compared to short-read RNA-seq[22, 23].

Previous studies have utilized both single-molecule long-read PacBio Iso-Seq and synthetic

long-read MOLECULO methods[24-27]. For Oxford Nanopore sequencing, there are two types

of RNA-seq technologies: direct mRNA sequencing and cDNA sequencing. Recently, the

Oxford Nanopore Technologies (ONT) MinION has been used to analyze both full-length cDNA

samples and mRNA samples derived from tissue cells[28]. Nanopore sequencing is able to

generate reads as long as 2Mbp, which allows a large portion or the entire mRNA or cDNA

molecule to be sequenced. Compared to short-reads, this advantage of long-reads greatly

facilitates rare isoform discovery, isoform expression quantification and DAS event detection.

However, there are still a few unique challenges to analyze long-read RNA-seq data because

existing methods developed for Illumina short-read RNA-seq do not have optimal performance

when directly used on long-read RNA-seq. This is because that parametric bias correction of

short-read approaches require high read coverage and isoform-read assignment is not robust to

small range misalignment from long-read data[16, 18, 19, 29]. Methods designed specifically for

isoform expression estimation in long-read RNA-seq have only emerged recently. For example,

Byrne *et al*[30] demonstrated the feasibility of quantifying complex isoform expression using

Nanopore RNA-seq data. Tang *et al*[31] characterized mutated gene *SF3B1* at isoform level in

chronic lymphocytic leukemia cells by leveraging full-length transcript sequencing data

generated by Nanopore. While long-read RNA-seq has great potential, the isoform

quantification accuracy is still constrained by high error rates and sequencing biases[32], which

has yet to be thoroughly accounted for. Specifically, high sequencing error rates (~15%) of

Nanopore data can result in misalignment of sequencing reads, but current methods assume

equal weight for each single molecule read without accounting for error rate differences when

estimating isoform expression. This may complicate isoform usage quantification. In addition,

potential read coverage biases are not explicitly taken into account by existing long-read

transcriptomic tools[32]. In Nanopore direct RNA sequencing protocol, pore block and

fragmentation can result in truncated reads, leading to biased coverage towards the 3' end of a

transcript[32]. These biases are also shown in data sequenced from cDNA. In the presence of

such biases, the accuracy of isoform expression quantification inference can be severely

affected, leading to over estimation of expression for isoforms with short length.

In this article, we present LIQA, a statistical method that allows each read to have its own

weight when quantifying isoform expression. Rather than counting single molecule reads

equally, we give a different weight to each read to account for read-specific error rate and

alignment bias at the gene (Figure 1). To evaluate the performance of LIQA, we simulated long

data with known ground truth and also sequenced two real samples using Oxford Nanopore

sequencing. Our results demonstrate that LIQA is an accurate approach for isoform expression

quantification accounting for read coverage bias and high error rate of long-read data.

Results

Overview of LIQA

Figure 1 shows the workflow of LIQA and highlights the read length bias correction step. LIQA

requires aligned long-read RNA-seq files in BAM or SAM format and isoform annotation file as

input. For estimation steps, LIQA first feeds read alignment information to a complete likelihood

function and corrects biases for each long read by accounting for quality score and read

coverage bias. Second, given that isoform origins are unobserved for some reads, an

Expectation Maximization (EM)-algorithm is utilized to achieve the optimal solution of isoform

relative abundance estimation. The output values of LIQA are isoform expression estimates.

Moreover, LIQA can further detect DAS events given estimated isoform expression values.

To evaluate the performance of LIQA, we compared it with existing long-read based

quantification algorithms, including FLAIR [31], Mandalorion [30], TALON [33] and the Oxford

Nanopore Pipeline (ONP, https://github.com/nanoporetech/pipeline-transcriptome-de). These

methods use long-read RNA-seq data to detect novel isoforms and quantify transcript

expression by counting the number of reads, which give equal weight for each read. To make

the comparisons fair, we ran LIQA, FLAIR. TALON, Mandalorion and ONP in quantification

mode only with isoform annotation information provided by GENCODE. We benchmarked the

performance of each method on both simulated and real data. In addition, we simulated more

data to evaluate the performance of LIQA in detecting DAS events between conditions.

Nanopore RNA-seq data simulation

We conducted a simulation study to evaluate the performance of LIQA and compared it with

other state-of-the-art algorithms for isoform expression estimation and DAS detection based on

GENCODE v24 annotation. To simulate a realistic dataset with known ground truth, we used

NanoSim[34] to generate the ONT RNA-seq data. NanoSim is a fast and scalable read

simulator that captures the technology-specific features of ONT data, and allows for adjustment

upon improvement of Nanopore sequencing technology. This simulator first characterizes

Nanopore reads and models features of the library preparation protocols *in silico* for read

simulation. The human genome sequence (GRCh38), transcriptome sequence and GTF

annotation file were downloaded from GENCODE. To make the simulated data close to real

studies, we assigned abundance values for each isoform obtained from a real human eye RNA-

seq dataset. Using NanoSim, we generated 5 million (5M) Nanopore reads. To evaluate the

impact of sequencing depth on isoform expression quantification, we down-sampled 3 million

(3M), 1 million (1M) and 0.5 million (0.5M) reads for the simulated data, respectively. These

reads were aligned to the reference genome using minimap2 [35]. Then, we selected genes

with 2 or more isoforms (67.2%) to evaluate the performance of LIQA in isoform expression

quantification. For each isoform, we compared it with Mandalorion, FLAIR, TALON and ONP. All

methods were run with the same set of simulated aligned data in BAM format as input.

The characteristics of the simulated data are shown in Figure 2(A) and Supplementary Figure

1(A). The median lengths of ONT reads in the 0.5M, 1M, 3M and 5M datasets are 896, 922,

1,010 and 993 base pairs, respectively. Among the evaluated genes with multiple isoforms

(67.2%) based on GENCODE annotation, 13% have two isoforms, 14% have three isoforms

and 73% have four or more isoforms. The simulated isoforms have a wide range of relative

abundance (interquartile range = (0.002, 0.75), median = 0.041). In addition, by training the

statistical model of NanoSim with a real long-read RNA-seq dataset, the coverage plots of the

simulated data capture the features of real ONT RNA-seq data, demonstrating 3' bias

(Supplementary Figure 1(B)). These simulated data thus provide an ideal basis to evaluate the

performance of LIQA as the ground truth is known.

Gene or Isoform expression quantification accuracy

For each simulated dataset, we computed a set of measurements to evaluate the estimation

accuracy of each method. First, we measured the similarity between the estimated isoform

relative abundance and the ground truth by calculating the Spearman's correlation. Second, we

measured the estimation accuracy by calculating the root mean squared error (RMSE), defined

as $\sqrt{\frac{\sum_g \sum_i (\hat{\theta}_{g,i} - \theta_{g,i})^2}{n}}$, where the summation is taken over all genes and all isoforms within each

gene and n is the total number of isoforms across all genes. Both statistics were computed at

three levels: global gene expression, global isoform expression, and within-gene isoform relative

abundances.

Figure 2(B)(C) (Supplementary Figure 2) show the summary statistics between estimated and

true values of global isoform expression (global isoform expression and isoform relative

abundances) at different read coverages. Spearman's correlation and RMSE were calculated

for all three methods. LIQA , FLAIR and TALON have higher Spearman's correlation than

Mandalorion and ONP across all simulated datasets. Compared to FLAIR and TALON, LIQA

has similar estimation accuracy based on Spearman's correlation. Figure 2(C) gives summary

statistics of relative abundance estimates for the five methods. For relative abundance

estimation, LIQA outperforms FLAIR and TALON with 6.6% and 7% lower RMSE on average,

respectively. Comparison results at gene level reveal similar pattern (Supplementary Figure 4).

The improved performance of LIQA is likely due to its use of the EM-algorithm, which assigns

unequal weight to each read to better account for mapping uncertainty and read mapping bias

(Figure 3(C)(D) and Supplementary Table 3). In contrast, FLAIR, TALON and Mandalorion

provide discrete estimations by directly counting the number of reads aligned to each

corresponding gene or isoform. Due to the limited read coverage of ONT RNA-seq data, it is not

surprising that they yield lower estimation accuracy.

Next, we evaluated the robustness of LIQA to 3' read coverage bias (Figure 3 and

Supplementary Figure 3). First, we compared the accuracy statistics for 5' terminal exon and 3'

terminal exon of each isoform. Isoform expression with non-uniform read coverage is more

challenging to estimate because the 5' end is less likely to be covered by sequencing reads

compared to 3' end. Figure 3(A) shows the comparison of Spearman's correlation for five

methods with 0.5M read coverage. LIQA is more accurate than the other four methods at 5'

terminal exon, especially when sequencing depth is low. The Spearman's correlation coefficient

of LIQA is 11% higher than the second best performing method FLAIR for 5' terminal exons,

while only 6% higher for 3' exons. This improved performance of LIQA in terminal exons

quantification is also demonstrated by RMSE values. LIQA has 8%-15% improvement of RMSE

values compared to other methods. Second, we considered the chi-square statistics that

measures the goodness-of-fit of coverage uniformity. Then, we divided the isoforms into two

categories based on median of the corresponding measure (Chi-square statistic > median, Chi-

square statistic < median) and summarized Spearman's correlation coefficient and RMSE for

each group of isoforms. For isoforms with less non-uniform read coverage, Spearman's

correlations of LIQA and FLAIR are close. However, despite reduced Spearman's correlation

value, LIQA is more accurate than the other four methods for isoforms with more non-uniform

read coverage (Chi-square statistic > median). The improvement of LIQA compared to FLAIR is

5% higher for these isoforms. This is likely because LIQA models potential truncated reads

which result in 3' coverage bias when quantifying isoform expression.

Moreover, we assessed the impact of modeling read length bias and read quality score on the

accuracy of isoform expression estimation. Figure 3(C)(D) show the comparison of isoform

estimation accuracy using different models. For isoforms with more non-uniform read coverage

(Chi-square statistics > median), model with read length bias correction has 9% (Full model vs

read quality model) and 10% (Read length model vs baseline model) higher Spearman's

correlation. For genes with less average read quality, model with read quality score has 6%

higher Spearman's correlation. Overall, isoform estimation accuracy drops noticeably when

using model with no read-length bias correction. This comparison demonstrates the advantage

of LIQA in handling read length bias and 3' bias correction over other approaches.

Differential alternative splicing (DAS) detection

Next, we evaluated the performance of LIQA in DAS detection. More ONT RNA-seq data across

multiple samples (*10* cases and *10* controls) were simulated for 10 times. NanoSim generated 3

million reads based on the GENCODE annotation per sample. To make true DAS events more

realistic, we sampled relative abundances of isoforms from a Dirichlet distribution with mean

and variance parameters estimated from a human eye RNA-seq dataset. Similarly, these

simulated data were mapped to the hg38 human reference genome using minimap2. Isoform

expression and usage difference between conditions were quantified using LIQA, FLAIR,

TALON, Mandalorion and ONP, respectively. We first compared the performance of DAS

detection between these methods using three summary statistics. After FDR control, we

measured the recalls (power) of our method by calculating the proportion of correctly predicted

DAS events among true DAS events. Second, we obtained precisions by calculating the

proportion of correctly predicted DAS events among predicted DAS events. Additionally, F1

score ($\text{F1 score} = 2 \cdot \frac{\text{precision} \cdot \text{recall}}{\text{precision} + \text{recall}}$) was summarized to average the precision and recall values.

As shown in Supplementary Figure 5(B), LIQA, FLAIR and TALON are more powerful than

others for all three evaluation metrics. This is not surprising because Madalorion and ONP have

lower accuracy in isoform expression estimation. For recall value, FLAIR (mean = 0.809, SD =

0.041) gives better and more consistent performance across 10 simulations than LIQA (mean =

0.776, SD = 0.058). However, in terms of precision value, LIQA (mean = 0.915, SD = 0.043)

yields less false positives than FLAIR (mean = 0.884, SD = 0.051). LIQA, FLAIR and TALON

had similar performance in detecting DAS events based on F1 score. Furthermore, we

generated ROC curve and precision-recall curve to compare the performance between methods

at different FDR thresholds (Figure 3(E) and Supplementary Figure 6, 7). As shown in Figure

3(E), LIQA achieved AUC = 0.94 after FDR control (FDR < 0.05). Given FDR threshold equals

to 0.05, LIQA gave the best performance with precision = 0.98 and recall = 0.78. Compared to

LIQA, the second-best performing method FLAIR yields 0.1%, 0.3%, 3.5% less in precision,

recall and AUC respectively. In addition, we examined isoform relative abundance estimation

accuracy from correct and incorrect detected DAS genes by LIQA. After FDR control, we

identified that 537 out of 2,465 genes are significantly differential spliced, which 431 are true

positives and 1,836 are true negatives. For these correctly predicted genes, true isoform relative

abundance is highly correlated with LIQA's estimates (Spearman's correlation= 0.91). For false

positive and negative genes, the Spearman's correlation is 38% lower compared to true positive

and negative. This is not surprising because accurate estimation of isoform expression level

leverages the power of regression model in detecting DAS events.

Application to the Universal Human Reference (UHR) RNA-seq data

As NanoSim generates ONT RNA-seq data based on trained parametric statistical model, we

recognized that simulated data is hardly a full representation of reality. To evaluate the

performance of LIQA in a real setting, we sequenced the Universal Human Reference sample

with Nanopore Direct mRNA sequencing (Supplementary Figure 12). Then, the resulting ONT-

RNA-seq data were analyzed using all five long-read based methods (LIQA, FLAIR,

Mandalorion, TALON, ONP). As quantitative real time polymerase chain reaction (qRT-PCR) is

considered as the most reliable technology for measuring true isoform abundance, we

downloaded the qRT-PCR measurements from the MAQC project under Gene Expression

Omnibus with accession number GSE5350. As part of the MAQC project, the expression levels

of 894 isoforms were measured by TaqMan Gene Expression Assay based qRT-PCR.

Additionally, we downloaded the UHR short-read RNA-seq data generated using the Illumina

platform. This dataset was analyzed using Cufflinks[15], CEM[16], Salmon[36] and Kallisto[14]

to compare the performance in isoform quantification between long reads and short reads.

Specifically, we mapped ONT and Illumina sequenced reads to the reference genome using

Minimap2[35] and STAR[37], respectively, and applied each quantification method to the RNA-

seq data. qRT-PCR measurements were treated as gold standard to compare the performance

across methods. We note that 563 of the 894 transcripts with qRT-PCR measurements are from

genes with a single isoform. Estimation results from these genes were served as positive

controls (Supplementary Figure 8(A)) because estimating isoform-specific expression for these

single-transcript genes is trivial. To compare the performance across different methods, we

considered those transcripts that are derived from genes with two or more isoforms.

To assess the accuracy between estimates and qRT-PCR measurements, we summarized

similarity metrics (Spearman's correlation and Pearson's correlation) of the isoform abundance

values in log-scale. As shown in Figure 4(A)(B), the estimation accuracy of all methods is lower

than simulated data, especially for those transcripts with qRT-PCR measures close to 0.

Nevertheless, we observed consistent results in terms of relative performance of different

methods with simulation data. LIQA is more accurate other methods with stronger linear

relationship between logarithm estimates and qRT-PCR measurements. However, many of the

lowly to moderately expressed isoforms were underestimated using the other methods with their

TPM values being compacted toward 0. For ONT data, the spearman's correlation of LIQA is

0.68, whereas the corresponding values from second best method TALON is 0.57. For Illumina

data, Cufflinks seems to correlate with the qRT-PCR measurements better than others

(Supplementary Figure 8(B)). The main reason for the better performance of LIQA is likely due

to quantifying isoform expression by accounting for isoform length bias and base quality scores.

Moreover, we randomly selected 3 genes and generated sashimi plots in Figure 4(C) to show

the read coverage difference between direct mRNA sequencing and Illumina data. Overall, read

distribution of long-read data is less heterogeneous than short-read. Specifically, for gene

CAPNS1, there is clearly severe 5' degradation in Illumina data, but with full length and even

coverage across the transcripts for long-read data. Terminal exons at 5' end in red square are

crucial informative regions for splicing analysis, which enable us to differentiate read origin from

different isoforms. As shown in Figure 4(C), these exonic regions were captured by Nanopore

reads but missed by Illumina reads, which significantly facilitates isoform expression

quantification using long-read RNA-seq data. Similarly, sashimi coverage plots of other two

genes showed the same pattern, which demonstrates the advantage of long-read data over

short-read in alternative splicing study.

Application to Nanopore cDNA sequencing data on a patient with acute myeloid

leukemia

AML is a type of blood cancer where abnormal myeloblasts are made by bone marrow[38]. In

this study, we sequenced peripheral blood from an acute myeloid leukemia (AML) patient using

GridION Nanopore sequencer with Guppy basecalling (https://denbi-nanopore-training-

course.readthedocs.io/en/latest/basecalling/basecalling.html). In total, we generated 8,061,683

long-reads with 6.6 GB bases (Supplementary Figure 13). We aligned the data against a

reference genome (hg38) using minimap2[35], and 63% long-reads (73% bases) are mapped.

Then, we analyzed this ONT RNA-seq data with LIQA for genes with at least two isoforms.

We considered two ways to benchmark the performance of LIQA. First, we used PennSeq to

analyze short-read sequencing data for the same AML sample and treated the estimates as

gold standard. This dataset included 440M short read with 150bp in length. Figure 5(A) shows

the scatter plots of isoform relative abundance estimates between long- and short-reads data.

Spearman's correlation coefficients were calculated. We found that correlation was improved

significantly for genes with at least 50 reads compared to all genes without filtration. Then, we

examined the major isoforms (with the highest expression level in a gene) inferred by LIQA. As

shown in Figure 5(B), long-read and short-read shared consistent estimates for the major

isoforms. This is not surprising because major isoforms were more likely to be sequenced,

leading to higher read coverage at unique exonic regions. Second, we visually examined the

read coverage plots at unique exonic regions with at least 100 reads to benchmark the

performance of LIQA. We generated sashimi plots for two randomly selected genes, *EOGT* and

RRBP1 (Figure 5(C)). For gene *EOGT*, the read coverage ratio between exons in red and green

squares suggests that isoforms NM_103826 and NM_001278689 expressed much higher than

NM_173654. This is consistent with LIQA' estimates, with relative abundance of NM_173654

less than 0.01. A similar pattern is observed for gene *RRBP1*, where isoform NM_004587

(relative abundance estimates = 0.68) is the major isoform. Results from this AML data

demonstrate the robust performance of LIQA to 3' coverage biases.

Application to PacBio data on esophageal squamous epithelial cell (ESCC)

Next, we evaluated the performance of LIQA in differential alternative splicing (DAS) detection

using an RNA-seq dataset generated from esophageal squamous epithelial cell (ESCC)[39].

This dataset includes PacBio SMRT reads generated from normal immortalized and cancerous

esophageal squamous epithelial cell lines. The RNA-seq data were downloaded from Gene

Expression Omnibus (PRJNA515570). We applied LIQA to detect differential isoform usage

between normal-like and cancer cells. Known splicing differences in existing studies were

treated as ground truth to evaluate LIQA's performance in characterizing isoform usage across

samples. In addition, short-read data from these two samples were sequenced using the

Illumina platform, which allows us to compare the consistency and accuracy of DAS detection

between long-read and short-read data.

Employing LIQA and PennDiff[40], PacBio and Illumina data were analyzed to detect DAS

events, which are classified into different types, such as skipped exon (SE), alternative 5' splice

site (A5SS), alternative 3' splice site (A3SS), mutually exclusive exon (MXE) and retained

intron(RI). Our results showed that SE is the most frequent type of event among detected DAS

between normal-like and cancerous cells, followed by RI, A5SS and A3SS. MXE is the most

infrequent splicing type. As shown in Figure 5(D), detected DAS events by long- and short-read

share strong association at both exon and gene level (Cramer's V > 0.5). Also, the concordance

rate between long and short read data is greater than 98%. Compared to short-read, long-read

data shows preference in detecting more differential splicing events at both exon and gene

level. This is not surprising because read coverage heterogeneity, which might bias DAS

detection, is alleviated in long-read data by capturing full-length transcript in each read.

The expression of alternatively spliced isoforms from gene *CDV3* shows difference in cancerous

ESCC cells compared to non-cancerous[41, 42]. Figure 5(E) provides the sashimi plots of a

DAS exon at gene *CDV3* detected by LIQA, but was missed by PennDiff using short-read data.

From long-read data, it is clear that the relative expression of exon in the red square is lower in

cancerous cells than normal-like ESCC. However, this event is missed by short-read data. The

read coverage difference between normal-like and cancerous ESCC in sashimi plots indicates

the less usage of isoforms (NM_001134422, NM_0011134423, NM_001282765) which include

this exon in ESCC cells, suggesting better performance of long-read data.

Discussion

Accurate estimation of isoform-specific gene expression is a critical step for transcriptome

profiling. The emergence of long-read RNA-seq has made it possible to discover complex novel

isoforms and quantify isoform usage based on full-length sequenced fragments without

amplification bias. However, there are still issues for long-read data, which if not taken into

account, can affect the estimations. The major challenges in the analysis of long-read RNA-seq

data are the presence of high error rate and potential coverage bias. In this article, we propose

LIQA, a statistical method that allows read-specific weight in estimating isoform-specific gene

expression. The central idea of our method is to extract error rate information and model non-

uniformity read coverage distribution of long-read data. LIQA is the first long-read transcriptomic

tool that takes these limitations of long-read RNA-seq data into account. Results of our

simulation study and analyses of real data demonstrated that LIQA is more effective in bias

correction than the limited existing approaches (Supplementary Table 2, 3).

However, we note that there is still room to improve LIQA. LIQA is computationally intensive

because the approximation of non-parametric Kaplan-Meier estimator of function $f(L_r)$ relies on

empirical read length distribution and the parameters are estimated using EM-algorithm. Based

on the analysis of the UHR and AML data, we found that running LIQA is slower than FLAIR

and Mandalorion (Supplementary Table 1). Currently, we are evaluating the impact of possible

parametric functions such as exponential or Weibull distributions for read distribution modeling.

This will sacrifice the robustness of isoform expression estimates but the running time can be

significantly reduced. We believe it may be worth making this trade-off between computational

efficiency and estimation accuracy for LIQA.

We have benchmarked the performance of LIQA with the use of minimap2 for long-read

alignment, while there have been several approaches supporting RNA-seq long-read alignment,

such as STAR[37], GMAP[43], BLAT[44], BBMap (https://sourceforge.net/projects/bbmap/), and

GraphMap2[45]. LIQA can take SAM or BAM files generated from any listed aligner as input.

Nevertheless, we recognize that it is important to evaluate whether LIQA's superior performance

is robust to different aligners. Therefore, we plan to explore more long-read aligner options and

settings to benchmark LIQA in the future.

As LIQA is EM-algorithm-based, the robustness to parameter initialization is a potential issue.

Read-specific weight of LIQA extracts more information from observed data than direct read-

count strategy as implemented in Mandalorion and FLAIR. Especially, more read coverage is

needed for stable approximation of function $f(L_r)$. For genes with limited reads coverage (less

than 5), the likelihood function of LIQA will be flattened, then optimal points are harder to be

reached by EM-algorithm and estimates may be sensitive to initial values of parameter.

Therefore, the sensitivity of LIQA to parameter initialization should be further evaluated and

improved.

With full-length transcript sequencing, long-read RNA-seq data (ONT and PacBio) are expected

to facilitate transcriptomic studies by offering number of advantages over short reads. For

PacBio, HiFi reads are generated with circular consensus sequencing (CCS) using single-

molecule consensus, which increases their accuracy over traditional multi-molecule consensus.

Compared to Nanopore sequencing, this protocol yields much lower per-base error rate

compared to Nanopore sequencing, but potentially shorter reads. Smaller read length may

introduce much larger biases in 5' or 3' coverage ratio, which requires further adjustment for

LIQA to derive more accurate isoform expression estimates. LIQA has custom settings that

allow users to flexibly adjust such parameters to handle these platforms. Compared to PacBio

(either with traditional library or HiFi library preparation protocols), ONT may be a more

promising platform in quantifying isoform expression while generating data with much higher

error rate. This is because ONT is currently more affordable with lower per-based cost of data

generation, and sequencing data with high read coverage can improve estimation accuracy of

isoform usage. For ONT RNA-seq, there are two types: direct mRNA sequencing and cDNA

sequencing. Compared to direct mRNA sequencing, cDNA sequencing allows samples to be

amplified and requires less amounts of starting materials. Our studies showed that the decrease

of read coverage had less impact on LIQA compared to other existing approaches.

In summary, long-read RNA-seq data offer advantages and can help us better understand

transcriptomic variations than short-read data. However, better utilizing informative single

molecule sequencing read is not straightforward. LIQA is a robust and effective computational

tool to estimate isoform-specific gene expression from long-read RNA-seq data. With the

increasing adoption of long-read RNA-seq in biomedical research, we believe LIQA will be well-

suited for various transcriptomics studies and offer additional insights beyond gene expression

analysis in the future.

Methods and materials

Complete likelihood function of LIQA

Given a gene of interest, let R denote the set of reads that are mapped to the gene of interest,

and I denote the set of known isoforms. For a specific isoform $i \in I$, let θ_i denote its relative

abundance, with $0 \leq \theta_i \leq 1$ and $\sum_{i \in I} \theta_i = 1$ and l_i denote its length. For each single-molecule

long-read r, let L_r denote its length. The probability that a read originates from isoform i is

$P(\text{iso.} = i) = \theta_i$. For read-isoform assignment, LIQA accounts for incorrect alignment at splice

site. We define parameter $\mathbf{Z}_{R,I}$ as a $|R| \times |I|$ a read-isoform compatibility matrix with $\mathbf{Z}_{R,I}(r, i) =$

1 if long-read r is generated from a molecule that is originated from isoform i (number of

mismatch base pairs < 20 bp instead of exact match), and $\mathbf{Z}_{R,I}(r, i) = 0$ otherwise. For isoform

quantification, our goal is to estimate $\Theta = \{\theta_i, i \in I\}$ based on RNA-seq long-reads mapped to

the gene.

With the notation above, the complete data likelihood of the RNA-seq data can be written as

$$L\left(\widetilde{\Theta}\big|\mathbf{R}, \mathbf{Z}\right) = \prod_{r \in R} \prod_{i \in I} \left(P(\text{read} = r, \text{read len.} = L_r, \text{iso.} = i)\right)^{\mathbf{Z}_{R,I}(r,i)}$$

$$= \prod_{r \in R} \prod_{i \in I} \left(P(\text{read} = r, \text{read len.} = L_r \mid \text{iso.} = i) \cdot P(\text{iso.} = i)\right)^{\mathbf{Z}_{R,I}(r,i)}$$

$$= \prod_{r \in R} \prod_{i \in I} \left(P(\text{read} = r, \text{read len.} = L_r \mid \text{iso.} = i) \cdot \theta_i\right)^{\mathbf{Z}_{R,I}(r,i)}$$

418   This formula is based the fact that given the isoform origin, the probability of observing read

419   alignment can be inferred. The conditional probability of read r derived from isoform i with

420   length L_r is

$$P(\text{read} = r, \text{read len.} = L_r \mid \text{iso.} = i) = q(r, i) \cdot f(L_r \mid \text{iso.} = i)$$

where $q(r, i)$ is isoform-specific read quality score and $f(L_r \mid \text{iso.} = i)$ is isoform-specific read

length probability. Essentially, we quantify isoform relative abundance with weighted read

assignment. To account for the error-prone manner of Nanopore sequencing data, we consider

isoform-specific read quality score $q(r, i) = \prod_{j=1}^{m} q_j(x_j, y_{(j)})$ where x is the sequence of the long-

read r, y is the sequence of the corresponding isoform i in the reference genome, and $q_j(a, b)$

is the probability that we observe base a at position j of the read given that the true base is b,

which can be calculated as $1 - 10^{-Q_j/10}$, with Q_j being the per-based Phred quality score at

position j.

Estimation of isoform-specific read length probability $f(L_r|\text{ iso.} = i)$

Because read length L_r is not fixed and short prone in Nanopore sequencing, we treat it as a

random variable with right skewed distribution density function $f(\cdot)$. Given an isoform, existing

long-read methods assume fixed read length for all sequenced read, and this is equivalent to

setting $f(L_r)$ at 1. However, this assumption does not hold as recent studies suggest that

potential 3' coverage bias exists in long-read RNA-seq data [24, 32, 46]. To offer flexibility in

modeling read length distribution, we employ a nonparametric approach. For all long-reads

mapped to the genome, we categorize them into two groups: complete reads and truncated

reads. Accounting for misalignment due to high error rate, the read is treated as complete when

the distance between its ending alignment position and any known isoform 5' end is less than a

tolerance threshold (default = 20 bp) (Figure 1(A)). This indicates that this read is completely

sequenced from a known isoform. Otherwise, the read is considered as truncated. The

presence of truncated reads is due to incomplete sequencing or novel isoforms. As known

annotated isoforms are treated as gold standard during estimation, we assume true length of

truncated read is censored. Given the observed lengths of all complete and truncated reads, we

fit them into a survival model, a natural modeling approach for censored data (Supplementary

Figure 9, 10, 11). Function $\hat{F}(l) = P(read\ len. < l)$ can be estimated based on Kaplan-Meier

estimator[47], hence we have $f(l) = \hat{F}(l+1) - \hat{F}(l)$.

Given a gene of interest with $I = \{\text{isoform } i : 1 \leq i \leq I\}$, isoform-specific read length probability

$f(L_r \mid \text{iso.} = i)$ can be written as

$$f(L_r \mid \text{iso.} = i) = \frac{f(L_r) \cdot P(\text{iso.} = i \mid L_r)}{P(\text{iso.} = i)} = \frac{f(L_r) \cdot \theta_i / \sum_{l_j > L_r} \theta_j}{\theta_i} = \frac{f(L_r)}{\sum_{l_j > L_r} \theta_j}$$

449   This isoform-specific read length probability $f(L_r \mid \text{iso.} = i)$ captures the sequencing biases due

450   to fragmentation during library preparation or pore-blocking for nanopore data.

Quantification of isoform expression level

Given that isoform indicators $\mathbf{Z}_{R,I}(r, i)$ for some reads are not observed from read data, Θ are

estimated using EM algorithm. Then, we have isoform relative abundance $\hat{\theta}_i$. In addition to

relative abundance, it is also important to quantify the absolute expression level of an isoform.

At gene level, we consider read per gene per 10K reads (RPG 10K) as the standard for long-

read RNA-seq data. RPG is defined as $\text{RPG} = N/10^4$ where N is the number of reads mapped

to the gene of interest. With this concept, we estimate the expression level of a particular

isoform by replacing N with estimated number of long-reads originated from isoform i ($\text{RPG}_i =$

$N \cdot \hat{\theta}_i/10^4$).

Parameter estimation using the EM algorithm

461   The complete data likelihood is

$$L(\Theta|\mathbf{R}, \mathbf{Z}) = \prod_{r \in R} \prod_{i \in I} (q(r, i) \cdot f(L_r) \cdot \theta_i)^{\mathbf{Z}_{R,I}(r,i)}$$

462   and the update procedure of the EM algorithm is as follows:

463   **E-step:** We calculate function

$$Q\left(\Theta\big|\Theta^{(t)}\right) = E_{Z_{R,I}|\Theta^{(t)}}[\log L(\Theta|\mathbf{R})]$$

$$= \sum_{r \in R} \sum_{i \in I} E_{Z_{R,I}|\Theta^{(t)}}[\mathbf{Z}_{R,I}(r, i)] \cdot \log (q(r, i)f(L_r)\theta_i)$$

464   where $E_{Z_{R,I}|\Theta^{(t)}}[\mathbf{Z}_{R,I}(r, i)] = \frac{q(r,i)f(L_r)\theta_i^{(t)}}{\sum_{u \in I} q(r,u)f(L_r)\theta_u^{(t)}}.$

465   **M-step:** We maximize function $Q\left(\Theta\big|\Theta^{(t)}\right)$ and have

$$\theta_i^{(t+1)} = \frac{\sum_{r \in R} E_{Z_{R,I}|\Theta^{(t)}}[Z_{R,I}(r, i)]}{|R|}$$

467 The EM algorithm consists of alternating between the E- and M-steps until convergence. We

468 start the algorithm with $\Theta^{(0)}$ assuming all isoforms are equally expressed and stop when the log

469 likelihood is no longer increasing significantly.

470 **Detection of differential alternative splicing (DAS) with LIQA**

471 The relative abundance of an isoform takes values between 0 and 1. Therefore, we assume it

472 follows a beta distribution, which is well known as a flexible distribution in modeling proportion

473 because its density can have different shapes depending on the values of the two parameters

474 that characterize the distribution. i.e. $\theta_i \sim \text{Beta}(\mu_i, \phi_i)$. The expected value and variance of θ_i are

$$E(\theta_i) = \mu_i$$

$$Var(\theta_i) = \frac{\mu_i(1 - \mu_i)}{1 + \phi_i}$$

475 To detect splicing difference of isoform i between two groups of samples, we utilized beta

476 regression model with ϕ_i as precision parameter. We apply logit link function and have the

477 model

$$\text{logit}(\theta_i) = \beta_0 + \beta_1 Z$$

478 where Z is the condition indicator (1 for case; 0 for control), β_0 and β_1 are coefficient

479 parameters.

480 Since the isoform relative abundances of isoforms within the same gene are correlated, a robust

481 and flexible model is needed when comparing them between conditions at gene level. To

482 account for this, we utilize Gaussian copula regression model to test splicing difference

483 significance between conditions of correlated isoform relative abundances. The separation of

484 marginal distributions and correlation structure makes Gaussian copula regression versatile in

485 modeling non-normal dependent observations. Therefore, the joint distribution of isoform

486 relative abundances from the same gene is given by

$$\Phi_{I-1}(\Phi^{-1}(F(\theta_1|\beta_0, \beta_1, \varphi_1)), \dots, \Phi^{-1}(F(\theta_{I-1}|\beta_0, \beta_{I-1}, \varphi_{I-1}))|\boldsymbol{\Gamma})$$

where φ_i is the dispersion parameter of the marginal generalized linear model for isoform i.

$\Phi_{I-1}(|\boldsymbol{\Gamma})$ is the cumulative distribution function of multivariate normal random variables with

$I-1$ dimensions and correlation matrix $\boldsymbol{\Gamma}$. We choose to use exchangeable correlation structure

for $\boldsymbol{\Gamma}$. Given regression models above, we can detect DAS both for at isoform level and at gene

level. For isoform i, we test $H_0: \beta_0 = 0$ vs $H_1: \beta_0 \neq 0$ to determine splicing change between

conditions. For gene g, we test $H_0: \beta_0 = \dots = \beta_{I-1} = 0$ vs $H_1: \beta_i \neq 0$ for any $1 \leq i \leq I-1$.

Nanopore direct mRNA sequencing of universal human reference RNA-seq data

Universal human reference (UHR) RNA comprises of mixed RNA molecules by a diverse set of

10 cancer cell lines with equal quantities of DNase-treated RNA from adenocarcinoma in

mammary gland, hepatoblastoma in liver, adenocarcinoma in cervix, embryonal carcinoma in

testis, glioblastoma in brain, melanoma, liposarcoma, histocytic lymphoma in histocyte

macrophage, lymphoblastic leukemia and plasmacytoma in B lymphocyte. This reference

sample from MicroArray Quality Control (MAQC)[48-50] project has been utilized in many

studies. For example, Gao et al[51] sequenced this UHR RNA sample and treated it as

reference to measure the technical variations of scRNA-seq data. Also, the qRT-PCR

measurements of gene or isoform expressions from this sample were used to benchmark and

optimize computational tools[17, 52-55]. In this study, we used GridION Nanopore technique to

sequence mRNA directly, and used Guppy for base calling. In total, we generated 476,000 long-

reads with 557 MB bases. We aligned the UHR RNA-seq data against a reference genome

(hg38) using minimap2[35], and 95% long-reads (89% of total bases) are mapped, demonstrating very high sequencing and basecalling quality. qRT-PCR measurements were downloaded and treated as ground truth to compare the performance between LIQA, FLAIR, Mandalorion, CEM, Cufflinks and RD.

Data availability

The direct mRNA sequencing data on UHR has been deposited and available at BioProject database (PRJNA639366). The cDNA sequencing data on a patient with cancer has been deposited and available at BioProject database (PRJNA640456). The simulation data used in our study can be reproduced using code provided in the LIQA software repository and NanoSim version 2.0.0.

Acknowledgements

We thank the Wang lab members for insightful comments and for testing the software tools. We also thank the developers of the NanoSim software tool, and the generators of the short-read and qRT-PCR results on the UHR datasets for making the data publicly available for benchmarking studies. We thank three anonymous reviewers for their constructive comments and suggestions on benchmarking studies.

Funding

This study is supported by NIH/NIGMS grant GM132713 and the CHOP Research Institute.

References

1.   Han, J., et al., *Pre-mRNA splicing: where and when in the nucleus.* Trends Cell Biol, 2011. **21**(6): p. 336-43.
2.   Scotti, M.M. and M.S. Swanson, *RNA mis-splicing in disease.* Nat Rev Genet, 2016. **17**(1): p. 19-32.
3.   Montes, M., et al., *RNA Splicing and Disease: Animal Models to Therapies.* Trends Genet, 2019. **35**(1): p. 68-87.

4.   Li, Y.I., et al., *RNA splicing is a primary link between genetic variation and disease.* Science, 2016.
352(6285): p. 600-4.
5.   Kim, H.K., et al., *Alternative splicing isoforms in health and disease.* Pflugers Arch, 2018. **470**(7):
p. 995-1016.
6.   Pan, Q., et al., *Deep surveying of alternative splicing complexity in the human transcriptome by*
high-throughput sequencing. Nat Genet, 2008. **40**(12): p. 1413-5.
7.   Wang, E.T., et al., *Alternative isoform regulation in human tissue transcriptomes.* Nature, 2008.
456(7221): p. 470-6.
8.   Wang, Z., M. Gerstein, and M. Snyder, *RNA-Seq: a revolutionary tool for transcriptomics.* Nat Rev
Genet, 2009. **10**(1): p. 57-63.
9.   Li, B. and C.N. Dewey, *RSEM: accurate transcript quantification from RNA-Seq data with or*
without a reference genome. BMC Bioinformatics, 2011. **12**: p. 323.
10.   Roberts, A. and L. Pachter, *Streaming fragment assignment for real-time analysis of sequencing*
experiments. Nat Methods, 2013. **10**(1): p. 71-3.
11.   Nariai, N., et al., *TIGAR2: sensitive and accurate estimation of transcript isoform expression with*
longer RNA-Seq reads. BMC Genomics, 2014. **15 Suppl 10**: p. S5.
12.   Zhang, C., et al., *Evaluation and comparison of computational tools for RNA-seq isoform*
quantification. BMC Genomics, 2017. **18**(1): p. 583.
13.   Patro, R., S.M. Mount, and C. Kingsford, *Sailfish enables alignment-free isoform quantification*
from RNA-seq reads using lightweight algorithms. Nat Biotechnol, 2014. **32**(5): p. 462-4.
14.   Bray, N.L., et al., *Near-optimal probabilistic RNA-seq quantification.* Nat Biotechnol, 2016. **34**(5):
p. 525-7.
15.   Trapnell, C., et al., *Differential gene and transcript expression analysis of RNA-seq experiments*
with TopHat and Cufflinks. Nat Protoc, 2012. **7**(3): p. 562-78.
16.   Li, W. and T. Jiang, *Transcriptome assembly and isoform expression level estimation from biased*
RNA-Seq reads. Bioinformatics, 2012. **28**(22): p. 2914-21.
17.   Hu, Y., et al., *PennSeq: accurate isoform-specific gene expression quantification in RNA-Seq by*
modeling non-uniform read distribution. Nucleic Acids Res, 2014. **42**(3): p. e20.
18.   Nicolae, M., et al., *Estimation of alternative splicing isoform frequencies from RNA-Seq data.*
Algorithms Mol Biol, 2011. **6**(1): p. 9.
19.   Wan, L., et al., *Modeling RNA degradation for RNA-Seq with applications.* Biostatistics, 2012.
13(4): p. 734-47.
20.   Steijger, T., et al., *Assessment of transcript reconstruction methods for RNA-seq.* Nat Methods,
2013. **10**(12): p. 1177-84.
21.   Tilgner, H., et al., *Defining a personal, allele-specific, and single-molecule long-read*
transcriptome. Proc Natl Acad Sci U S A, 2014. **111**(27): p. 9869-74.
22.   Burgess, D.J., *Genomics: Next regeneration sequencing for reference genomes.* Nat Rev Genet,
2018. **19**(3): p. 125.
23.   Pollard, M.O., et al., *Long reads: their purpose and place.* Hum Mol Genet, 2018. **27**(R2): p.
R234-R241.
24.   Sharon, D., et al., *A single-molecule long-read survey of the human transcriptome.* Nat
Biotechnol, 2013. **31**(11): p. 1009-14.
25.   Tilgner, H., et al., *Comprehensive transcriptome analysis using synthetic long-read sequencing*
reveals molecular co-association of distant splicing events. Nat Biotechnol, 2015. **33**(7): p. 736-
42.
26.   Treutlein, B., et al., *Cartography of neurexin alternative splicing mapped by single-molecule long-*
read mRNA sequencing. Proc Natl Acad Sci U S A, 2014. **111**(13): p. E1291-9.

27. Vollmers, C., et al., *Novel exons and splice variants in the human antibody heavy chain identified*
by single cell and single molecule sequencing. PLoS One, 2015. **10**(1): p. e0117050.

28. Oikonomopoulos, S., et al., *Benchmarking of the Oxford Nanopore MinION sequencing for*
quantitative and qualitative assessment of cDNA populations. Sci Rep, 2016. **6**: p. 31602.

29. Roberts, A., et al., *Improving RNA-Seq expression estimates by correcting for fragment bias.*
Genome Biol, 2011. **12**(3): p. R22.

30. Byrne, A., et al., *Nanopore long-read RNAseq reveals widespread transcriptional variation*
among the surface receptors of individual B cells. Nat Commun, 2017. **8**: p. 16027.

31. Tang, A.D., et al., *Full-length transcript characterization of SF3B1 mutation in chronic*
lymphocytic leukemia reveals downregulation of retained introns. Nat Commun, 2020. **11**(1): p.
1438.

32. Amarasinghe, S.L., et al., *Opportunities and challenges in long-read sequencing data analysis.*
Genome Biol, 2020. **21**(1): p. 30.

33. Wyman, D., et al., *A technology-agnostic long-read analysis pipeline for transcriptome discovery*
and quantification. BioRxiv, 2020: p. 672931.

34. Hafezqorani, S., et al., *Trans-NanoSim characterizes and simulates nanopore RNA-sequencing*
data. Gigascience, 2020. **9**(6).

35. Li, H., *Minimap2: pairwise alignment for nucleotide sequences.* Bioinformatics, 2018. **34**(18): p.
3094-3100.

36. Patro, R., et al., *Salmon provides fast and bias-aware quantification of transcript expression.* Nat
Methods, 2017. **14**(4): p. 417-419.

37. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-21.

38. De Kouchkovsky, I. and M. Abdul-Hay, *'Acute myeloid leukemia: a comprehensive review and*
2016 update'. Blood Cancer J, 2016. **6**(7): p. e441.

39. Cheng, Y.W., et al., *Long Read Single-Molecule Real-Time Sequencing Elucidates Transcriptome-*
Wide Heterogeneity and Complexity in Esophageal Squamous Cells. Front Genet, 2019. **10**: p.
915.

40. Hu, Y., et al., *PennDiff: detecting differential alternative splicing and transcription by RNA*
sequencing. Bioinformatics, 2018. **34**(14): p. 2384-2391.

41. Xie, Z.C., et al., *Prognostic alternative splicing signatures and underlying regulatory network in*
esophageal carcinoma. Am J Transl Res, 2019. **11**(7): p. 4010-4028.

42. Ueno, N., et al., *Enhanced Expression of Fibroblast Growth Factor Receptor 3 IIIc Promotes*
Human Esophageal Carcinoma Cell Proliferation. J Histochem Cytochem, 2016. **64**(1): p. 7-17.

43. Wu, T.D. and C.K. Watanabe, *GMAP: a genomic mapping and alignment program for mRNA and*
EST sequences. Bioinformatics, 2005. **21**(9): p. 1859-75.

44. Kent, W.J., *BLAT--the BLAST-like alignment tool.* Genome Res, 2002. **12**(4): p. 656-64.

45. Maric, J., et al., *Graphmap2-splice-aware RNA-seq mapper for long reads.* bioRxiv, 2019: p.
720458.

46. Kellner, S., J. Burhenne, and M. Helm, *Detection of RNA modifications.* RNA Biol, 2010. **7**(2): p.
237-47.

47. Kaplan, E.L. and P. Meier, *Nonparametric estimation from incomplete observations.* Journal of
the American statistical association, 1958. **53**(282): p. 457-481.

48. Consortium, M., et al., *The MicroArray Quality Control (MAQC) project shows inter- and*
intraplatform reproducibility of gene expression measurements. Nat Biotechnol, 2006. **24**(9): p.
1151-61.

49. Shi, L., et al., *The MicroArray Quality Control (MAQC)-II study of common practices for the*
development and validation of microarray-based predictive models. Nat Biotechnol, 2010. **28**(8):
p. 827-38.

50.   Sun, P., et al., *Expression of estrogen receptor-related receptors, a subfamily of orphan nuclear*
receptors, as new tumor biomarkers in ovarian cancer cells. Journal of Molecular Medicine,
2005. **83**(6): p. 457-467.
51.   Gao, F., et al., *Evaluation of biological and technical variations in low-input RNA-Seq and single-*
cell RNA-Seq. International Journal of Computational Biology and Drug Design, 2018. **11**(1-2): p.
5-22.
52.   Xu, J., et al., *Cross-platform ultradeep transcriptomic profiling of human reference RNA samples*
by RNA-Seq. Sci Data, 2014. **1**: p. 140020.
53.   Garcia-Alonso, L., et al., *Benchmark and integration of resources for the estimation of human*
transcription factor activities. Genome Res, 2019. **29**(8): p. 1363-1375.
54.   Teng, M., et al., *A benchmark for RNA-seq quantification pipelines.* Genome Biol, 2016. **17**: p. 74.
55.   Hayer, K.E., et al., *Benchmark analysis of algorithms for determining and quantifying full-length*
mRNA splice forms from RNA-seq data. Bioinformatics, 2015. **31**(24): p. 3938-45.

Figures

Figure 1. Framework of LIQA. (A) The flowchart to illustrate how LIQA works. The inputs to

LIQA are long-read RNA-seq data and isoform annotation file. LIQA models observed splicing

information, high error rate of data and read length bias. The output of LIQA are isoform

expression estimates and detected DAS events. (B) Quantification of potential 3' bias of long-

read RNA-seq data. Complete and degraded RNA transcript are indicated by orange and blue.

Complete (orange) and truncated (blue) long reads are jointly modeled to correct read length

bias by estimating read length distribution.

Figure 2. Simulation study results. (A) Characteristics of simulated data with 5M. Read length

distribution (left) and read count distribution by genes in log-scale (right). (B-C) Summary

statistics between true and estimated isoform expressions using LIQA (blue), FLAIR (red),

Mandalorion (gray), TALON (green) and Oxford Nanopore Pipeline (ONP) (yellow) at different

read coverages. (B) Spearman's correlations (left) and RMSE (right) between estimated and

true TPM. (C) Spearman's correlations (left) and RMSE (right) between estimated and true

relative abundance.

Figure 3. Evaluation of robustness to read length bias and low read quality (A-D) and comparison of DAS events detection (E-F) using LIQA (blue), FLAIR (red), Mandalorion (gray), TALON (green) and ONP (yellow). (A) Spearman's correlations between estimated and true TPM for 3' and 5' terminal exons. (B) Spearman's correlations between estimated and true TPM. Isoforms are stratified by the chi-squared goodness of fit statistic for uniformity. The left panel is for those isoforms in which the chi-squared statistic is less than the median, and the right panel is for the remaining isoforms. (C-D) Spearman's correlation comparison between different models (full model, read length model, read quality model and baseline model) based

674   on isoforms with chi-squared statistic less than the median (C) and isoforms with average read

675   quality score less the median (D). (E) ROC curve and precision recall curve of different methods

676   for DAS gene detection. Threshold FDR<0.05 is highlighted using black dot. (F) Scatter plot of

677   true and estimated isoform relative abundance from the first DAS simulation dataset. LIQA's

678   prediction and ground truth are marked for each isoform.

Figure 4. UHR data analysis results. (A-B) Comparison of different methods. (A) Scatter plots

of estimated isoform-specific expression versus qRT-PCR measurements in log scale. Black

line represents the local polynomial regression line fitted with estimates and qRT-PCR in log

scale (B) Pearson's correlation coefficients (left) and spearman's correlation coefficient (right)

between estimated isoform-specific expression versus qRT-PCR measurements in log scale.

(C) Examination of read coverage difference between Illumina and Nanopore data at 3 genes.

Informative exonic regions were in red square.

Figure 5. Performance of LIQA using AML data (A-C) and ESCC data (D, E). (A) Scatter
plots of estimated isoform relative abundances using long-read data (LIQA) versus short-read
data (PennSeq) for all genes (left) and genes with at least 50 read coverage (right). (B) Scatter
plot of estimated isoform-specific expression using long-read data (LIQA) versus short-read
data (PennSeq) in log-scale for all major isoforms. (C) Examination of isoform usage inferred by
LIQA. Sashimi plots of gene *EOGT* and *RRBP1*. Informative exonic regions were in green and
red squares. (D) DAS detections between long- and short-read data. Consistency of detected
DAS events between long- and short-read data were quantified using Cramer's V and
concordance rate. (E) Examination of AS exon usage inferred by LIQA (long-read) but missed
by PennDiff (short-read). Sashimi plots of gene *CDV3*. Informative exonic regions were red
squares.