Novel regulatory and transcriptional networks associated with resistance to platinum-based chemotherapy in ovarian cancer

Danai G. Topouza[a], Jihoon Choi[a], Sean Nesdoly[b], Anastasiya Tarnouskaya[b], Christopher J.B. Nicol[a,c,d], Qing Ling Duan[a,b]

[a]Department of Biomedical and Molecular Sciences, Queen's University, Kingston, Ontario, Canada; [b]School of Computing, Queen's University, Kingston, Ontario, Canada; [c]Department of Pathology and Molecular Medicine, Queen's University, Kingston, Ontario, Canada; [d]Division of Cancer Biology and Genetics, Queen's University Cancer Research Institute, Queen's University, Kingston, Ontario, Canada.

Correspondence: Qing Ling Duan

Computational Genomics Laboratory

Botterell Hall, Room 530

18 Stuart St, Kingston, ON, Canada

K7L 3N6

Email : qingling.duan@queensu.ca

Telephone: 613-533-6356

Abstract

Background

High-grade serous ovarian cancer (HGSOC) is a highly lethal gynecologic cancer, in part due to

resistance to platinum-based chemotherapy reported among 20% of patients. This study aims to

elucidate the biological mechanisms underlying chemotherapy resistance, which remain poorly

understood.

Methods

Sequencing data (mRNA and microRNA) from HGSOC patients were analyzed to identify

differentially expressed genes and co-expressed transcript networks associated with chemotherapy

response. Initial analyses used datasets from The Cancer Genome Atlas and then replicated in two

independent cancer cohorts. Moreover, transcript expression datasets and genomics data (i.e.

single nucleotide polymorphisms) were integrated to determine potential regulation of the

associated mRNA networks by microRNAs and expression quantitative trait loci (eQTLs).

Results

In total, 196 differentially expressed mRNAs were enriched for adaptive immunity and translation,

and 21 differentially expressed microRNAs were associated with angiogenesis. Moreover, co-

expression network analysis identified two mRNA networks associated with chemotherapy

response, which were enriched for ubiquitination and lipid metabolism, as well as three associated

microRNA networks enriched for lipoprotein transport and oncogenic pathways. In addition,

integrative analyses revealed potential regulation of the mRNA networks by the associated

microRNAs and eQTLs.

Conclusion

We report novel transcriptional networks and pathways associated with resistance to platinum-based chemotherapy among HGSOC patients. These results aid our understanding of the effector networks and regulators of chemotherapy response, which will improve drug efficacy and identify novel therapeutic targets for ovarian cancer.

Keywords

Drug Resistance, Neoplasm; Carcinoma, Ovarian Epithelial; High-Throughput RNA Sequencing; Gene Regulatory Networks; MicroRNAs; Quantitative Trait Loci

Abbreviations

AOCS: Australian Ovarian Cancer Study

BLCA: Bladder Urothelial Carcinoma

eQTL: Expression Quantitative Trait Locus

HGSOC: High-Grade Serous Ovarian Cancer

OC: Ovarian Cancer

OS: Overall Survival

PFS: Progression-Free Survival

TCGA: The Cancer Genome Atlas

WGCNA: Weighted Gene Co-Expression Network Analysis

1. Introduction

High-grade serous ovarian cancer (HGSOC) is a highly lethal gynecologic cancer, in part due to resistance to first-line, platinum-based chemotherapy treatment among 20% of patients [1]. Chemotherapy resistant patients have a significantly shorter overall survival (OS) than sensitive patients, and many experience tumor recurrence within six months of completing chemotherapy [2]. There is currently no strategy for predicting response to platinum-based chemotherapy, which reflects our limited understanding of the underlying molecular mechanisms of chemoresistance [3].

Earlier studies reporting gene expression signatures associated with chemoresistance in ovarian cancer patients had used univariate analysis methods, which assume that chemotherapy response is driven by a single gene [4]. However, it is well established that chemotherapy response, like other drug response outcomes, is a complex multifactorial trait modulated by multiple genes contributing to one or more biological pathways [5].

In this study, we apply both univariate and multivariate analysis methods to provide the first evidence of novel mRNA and microRNA (miRNA) signalling pathways and networks associated with chemotherapy response in HGSOC patients. We further unveil miRNAs and expression quantitative trait loci (eQTLs) correlated with the expression of significant transcripts. These findings are validated using two independent cancer cohorts and provide important therapeutic strategies for regulating platinum-based chemotherapy response in HGSOC patients.

2. Materials and Methods

2.1 Chemotherapy response classification

Sequencing of mRNA and miRNA was derived from chemotherapy-naïve tumors of 191 and 205 HGSOC patients of TCGA, respectively [6]. Patients who received platinum-based adjuvant chemotherapy were selected and classified for chemotherapy response based on their platinum-free interval. Sensitive patients remained cancer-free for at least 12 months after chemotherapy completion, whereas resistant patients experienced cancer recurrence within 6 months (**Table 1**; **Supplementary Methods**).

2.2 Processing of sequencing data

Sequencing reads from mRNA were downloaded as FASTQ files (i.e. level 1 data from TCGA), filtered for base-quality, aligned, and quantified (detailed in **Supplementary Methods**). Sequencing of miRNA data were downloaded as quantified expression files (i.e. level 3 data from TCGA). Both mRNA and miRNA datasets underwent outlier detection, normalization, and non-specific filtering, resulting in 49,116 mRNA and 4,479 miRNA transcripts for further analyses.

2.3 Differential expression analysis

Differentially expressed mRNA and miRNA transcripts were detected between sensitive and resistant patients using a negative binomial generalized linear model (GLM) in the DESeq2 R package [7]. This analysis controlled for patients' ages at diagnosis, as resistant patients were significantly older (**Table 1**). The Benjamini-Hochberg method corrected for multiple testing.

2.4 Weighted co-expression network analysis

The weighted gene co-expression network analysis (WGCNA) R package [8] was used to identify

modules of co-expressed mRNA and miRNA transcripts using an unsupervised machine learning

approach. The first principal component of expression values was calculated for each transcript

module, resulting in an eigengene value. Module eigengenes were used to determine association

with chemotherapy response using a GLM, adjusted for patients' age as a covariate

(**Supplementary Methods**).

2.5 Pathway enrichment analysis

Pathway enrichment analysis was used to determine over-representation of biological pathways

from lists of differentially expressed transcripts (mRNA and miRNAs) and co-expression networks

(**Supplementary Methods**).

2.6 eQTL analysis

Germline single nucleotide polymorphisms (SNPs) from TCGA-HGSOC patients were imputed

as described by Choi *et al.* [9] before undergoing quality control and linkage disequilibrium-based

pruning, retaining 1,722,608 common SNPs for analysis. SNPs were integrated with patient

mRNA-seq data (n=167) and miRNA-seq data (n=178) to identify correlations with transcript

expression (eQTLs) using the MatrixEQTL R package [10] (**Supplementary Methods**).

2.7 mRNA-microRNA integration

Potential regulation of mRNA networks by miRNAs was tested using data from 165 patients by

applying the Spearman correlation of module eigengenes from the mRNA and miRNA co-

expression networks. Results were validated using miRNet [11], a database of experimentally

validated mRNA-miRNA interactions, and miRGate [12], a database of predicted mRNA-miRNA

interactions based on ribonucleotide motif matching.

 2.8 Replication cohort and analysis

Replication of differentially expressed transcripts used the Kaplan-Meier analysis tool KM Plotter

[13] to test the association of each transcript with progression-free survival (PFS) or OS in two

independent cancer cohorts. First, mRNA transcripts were replicated with PFS from the Australian

Ovarian Cancer Study (AOCS; GSE9891; n=285) [14]. Next, miRNA isoforms were replicated

with OS in the TCGA bladder urothelial carcinoma cohort (BLCA) (n=409) [15]. Validation of

transcript networks used the Prognostic Index estimation method [16] using data from these two

cohorts (**Supplementary Methods**).

3. Results

Differential expression analysis identified 196 mRNAs associated with chemotherapy response

(adjusted p<0.05) that map to 190 unique genes (**Figure 1A, Table S1**). Pathway enrichment

analysis of these associated transcripts indicated enrichment of 41 annotation terms, including B-

cell receptor regulation, complement activation, and peptide chain elongation (**Table 2**).

Differential miRNA expression analysis revealed 21 differentially expressed miRNA

isoforms (adjusted p<0.05), which map to 16 unique miRNAs (**Figure 1B, Table S2**). Pathway

enrichment analysis of these miRNA isoforms revealed 16 pathways, such as blood vessel

morphogenesis and negative regulation of autophagy (**Table 3**).

WGCNA of the mRNA transcripts resulted in 58 co-expression modules (**Table S3**), of

which two were associated with platinum-based chemotherapy. First, the *lavenderblush3* module

is negatively associated with chemotherapy resistance (p-value=0.016; **Figure 2A**). This module

contains 39 transcripts, mapping to 31 unique genes (**Figure 2B**). Pathway analysis indicates

enrichment of biological pathways related to protein ubiquitination, and the binding motif for

transcription factor GABP-alpha (**Table 4**). Second, the *darkolivegreen* module is significantly

upregulated in chemo-resistant patients (p-value=0.032) (**Figure 2A**) and contains 82 transcripts

mapping to 80 unique genes (**Figure 2B**). This module is significantly enriched for the protein-

containing complex term, as well as for 8 pathways involved in fatty acid metabolism with nominal

significance (**Table 4**).

WGCNA analysis of the miRNA dataset constructed 100 co-expression modules (**Table**

S4), of which three are associated with chemotherapy response. The *ivory* (p-value=0.0097) and

lightcoral (p-value=0.042) modules are negatively associated with chemotherapy resistance, while

the third *plum* network (p-value = 0.045) is positively associated with chemotherapy resistance

(**Figure 2C**). The *ivory* network consisted of 25 miRNA isoforms mapping to 11 unique miRNAs,

which are enriched for 7 pathways and functions, including regulation of lipoprotein transport and

cholesterol efflux (**Table 5**). The *lightcoral* module consists of 17 isoforms of miR-187 and the

plum network consists of 17 isoforms of miR-221 and miR-222 (**Figure 2D**). While no pathway

annotations were derived for the *lighcoral* module, the *plum* module is enriched for 18 pathways

and oncogenic functions, such as inhibition of the TRAIL-activated apoptotic pathway and

inflammatory cytokine production, and up-regulation of protein kinase B signaling (**Table 5**).

Integrative analysis with germline SNP data identified 268 unique cis-eQTLs associated

with the expression of mRNAs and miRNAs which were correlated with chemotherapy response

in differential expression and network analysis. A total of 248 SNPs are associated with the

expression of 55 significant mRNAs, and 20 SNPs are associated with the expression of 7

significant miRNAs (**Table S5**). Of the 268 eQTLs, 118 are novel whereas 126 are previously

known, and 24 are not yet recorded in the annotation database. The majority (227) are predicted

to alter regulatory motifs, and 67 are associated with 94 human phenotypes from published

genome-wide association studies. The most common phenotypes are related to triglycerides, high-

density lipoprotein (HDL), and low-density lipoprotein (LDL) cholesterol.

170   Integration of the associated mRNA and miRNA networks determine that the *plum* miRNA

171   network significantly correlates with the *lavenderblush3* mRNA network (Spearman's ρ = -0.26,

172      $p < 0.001$), and the *ivory* miRNA network significantly correlates with the *darkolivegreen* mRNA

173   network (Spearman's ρ = -0.17, $p = 0.023$). Annotations using miRNet and miRGate determined

174   that 20 of these mRNA-miRNA interactions are experimentally validated, while 15 others are

175   supported by *in silico* predictions (**Table 6, Figure 3, Supplementary Data 1**).

176   Replication analysis of the mRNA results used RNA microarray data from the AOCS

177   cohort. In total, 140 microarray probes in the AOCS dataset (73.68%) matched to differentially

178   expressed genes from TCGA-HGSOC, 28 (20%) of which replicated with the same direction of

179   effect (**Table S6**). In addition, the *lavenderblush3* and *darkolivegreen* mRNA network modules

180   replicated in the AOCS cohort ($p < 0.0001$) (**Figure 4A, B**).

181   Replication of the miRNA findings used miRNA-seq data from the BLCA cohort of

TCGA. Of the 16 differentially expressed miRNAs isoforms in the HGSOC data, 13 (81.25%)

were available in the TCGA-BLCA data and 7 (53.85%) replicated with the same direction of

effect (**Table S7**). In addition, the *ivory* miRNA network module replicated in the TCGA-BLCA

cohort ($p = 0.012$), while the *lightcoral* and *plum* modules did not replicate (**Figure 4C-E**).

However, differential expression of miR-221 and miR-222, which make up the *plum* network,

replicated in this cohort using Kaplan-Meier analysis ($p = 0.022$ and $p = 0.037$) (**Figure 4F, G**).

4. Discussion

In this study, we analyzed mRNA and miRNA sequencing data from chemotherapy-naïve tumors

of HGSOC patients to identify transcripts and networks associated with chemotherapy response.

Our findings implicate novel and known biological pathways that replicated in independent cancer

cohorts. In addition, we identified potential interactions among miRNAs and mRNAs, as well as

eQTLs that potentially regulate the associated transcripts. Thus, our results provide an integrative,

multi-omics view of biological networks associated with chemotherapy response.

We identified one mRNA co-expression network (*lavenderblush3*) significantly

upregulated in platinum sensitive patients, which replicated in the AOCS. This module consists of

genes involved in ubiquitin-mediated proteolysis in the endoplasmic reticulum (ER). We also

detected a significant downregulation of genes responsible for translation initiation in sensitive

patients. These findings suggest that the unfolded protein response (UPR), a cellular process

responsible for resolving ER stress, may be increasingly activated in sensitive patients compared

to resistant cases. The UPR alleviates ER stress through several pathways, including increased ER-

associated protein degradation (ERAD) to remove misfolded proteins, and inhibition of translation

to reduce protein load in the ER [17]. ER stress promotes cisplatin resistance in OC cell lines [18]

and the upregulation of ERAD genes such as *VCP* in the *lavenderblush3* module is associated with

longer OS and platinum sensitivity in HGSOC cohorts [9, 19]. Eighteen of 31 module genes were

also reported in an earlier study of microarray data from the same TCGA-HGSOC cohort [9].

We identified a second mRNA co-expression network associated with chemotherapy

resistance in our HGSOC cohort that replicated in the AOCS. The *darkolivegreen* module included

genes associated with fatty acid metabolism (*SREBF1*, *ACAA1*, *ACADV*L), and the protein kinase

B oncogene (*AKT1*), which promotes de-novo lipid biosynthesis in cancer [20]. SREBF1 is a key

enzyme for cholesterol and fatty acid synthesis, and an essential gene for OC tumor growth [21].

Specifically, *SREBF1* is activated by *AKT1*, promoting fatty acid synthesis [22], which favors cell

proliferation in OC [23]. Expression of *ACADVL*, involved in the β-oxidation of long-chain fatty

acids, is linked to OC metastasis and cell survival [24]. Our findings indicate the upregulation of

these lipid metabolism genes among chemotherapy resistant patients. Lipid metabolism

dysregulation activates the UPR, which triggers lipid metabolism-based adaptations in the cell

through several pathways, including *SREBF1* regulation [25]. The interaction of these pathways

may present a link between our two gene co-expression modules and warrants further study.

Differential expression analysis identified a downregulation of mRNA transcripts involved

in the adaptive immune system, which is associated with chemoresistance. Previous studies

reported that a high tumor immune score is a strong predictor of chemosensitivity in HGSOC [26].

In addition, there are potential links between this immune response activation, UPR and lipid

metabolism. ER stress can induce pro-inflammatory cytokine production and UPR activation in

tumor cells [27], which can disrupt dendritic cell function in the OC tumor microenvironment [28].

Moreover, dendritic cell function can also be inhibited by increased lipid uptake in various cancers

[29].

The *ivory* miRNA co-expression network module, associated with chemotherapy

sensitivity in HGSOC and BLCA, is involved in the negative regulation of lipid transport. This

enrichment is mainly mediated by miR-128-1 and miR-128-2, which play a key role in cholesterol

and lipid homeostasis through their suppression of the ABCA1 cholesterol efflux transporter and

the low-density lipoprotein receptor (LDLR) [30, 31]. MiR-148a is also a regulator of these key

genes [30], which is significantly downregulated in resistant patients. The overexpression of

ABCA1 is associated with reduced survival in OC patients [32], and levels of LDLR are increased

in chemoresistant OC cell lines [33]. In addition, overexpression of miR-128 promotes sensitivity

to cisplatin in previously resistant OC cells [34]. Our results are consistent with the

chemosensitivity-promoting role of miR-128 and its potential activity in cholesterol efflux

inhibition alongside miR-148a in this cohort.

The *plum* miRNA co-expression network consists of miR-221 and miR-222 isoforms,

which have been implicated in the development of chemotherapy resistance in OC. Expression of

miR-221/miR-222 transcripts is high in cisplatin-resistant OC cell lines, and their inhibition

increases cellular sensitivity [35]. Over-expression of miR-221 and miR-222 promotes

proliferation of OC cell lines [36, 37] and reduced disease-free and overall survival [36]. Thus, our

findings are consistent with earlier studies showing increased activity of miR-221 and miR-222 in

chemoresistant tumors.

Integrative analysis of mRNA-seq and miRNA-seq datasets identified potential

interactions of the associated transcript co-expression modules. The overexpression of miR-

221/222 in resistant patients may be inhibiting the chemosensitivity-associated *lavenderblush3*

mRNA network, revealing a novel potential mechanism of chemotherapy resistance. This finding,

combined with the accumulating evidence of miR-221/miR-222 involvement in chemoresistance,

may point to a promising avenue for therapeutic intervention. However, overexpression of miR-

221/miR-222 promotes UPR-induced apoptosis in hepatocellular carcinoma (HCC) cells [38].

Additionally, ER stress suppresses miR-221/miR-222 in HCC, promoting resistance to apoptosis.

The contribution of this mechanism to chemotherapy response in HGSOC is currently unclear and

presents an area for future investigation.

We also identified potential regulation of the *darkolivegreen* mRNA module by the *ivory*

miRNA network, which may inhibit lipid metabolism in chemotherapy sensitive patients. As

increased lipid metabolism by cancer cells is a known mechanism of chemoresistance in HGSOC,

this miRNA-mediated inhibition may present a novel mechanism of chemotherapy sensitivity.

Finally, cis-eQTL analysis identified known and novel genomic variants correlated with

the expression of mRNAs and miRNAs, which are associated with lipid-related phenotypes. High

HDL and triglyceride levels have been correlated with increased cancer stage at diagnosis in OC

patients [39]. In addition, advanced-stage OC patients with high LDL levels have a shorter PFS

than patients with normal levels [40]. Further investigation of these eQTLs is necessary to further

elucidate their role in platinum-based chemotherapy resistance and HGSOC prognosis.

5. Conclusion

Our study provides novel insight of the underlying mechanisms modulating resistance to platinum-

based chemotherapy in HGSOC. Specifically, we conducted whole-transcriptome analysis of

mRNA-seq and miRNA-seq data to generate novel mechanistic hypotheses using both univariate

and network methods. Moreover, we integrated this data with miRNA-seq and genome-wide SNPs

to determine potential regulation of the associated transcripts and networks. Our findings implicate

novel and known signaling pathways and networks associated with chemotherapy response in

HGSOC as well as regulators, which could become novel drug targets. Further studies are needed

to validate these findings in other cancers.

CRediT authorship contribution statement

Danai Georgia Topouza: Investigation, Formal analysis, Software, Visualization, Validation,

Data curation, Writing – original draft, Writing – review & editing. **Jihoon Choi:** Formal analysis,

Software, Data curation, Writing – review & editing. **Sean Nesdoly:** Software, Data curation.

Anastasiya Tarnouskaya: Data curation. **Christopher J.B. Nicol:** Writing – review & editing.

Qing Ling Duan: Conceptualization, Supervision, Methodology, Resources, Funding acquisition,

Writing – review & editing.

Acknowledgements

Computations in this manuscript were performed on resources and with support provided by the

Centre for Advanced Computing (CAC) at Queen's University in Kingston, Ontario. The CAC is

funded by the Canada Foundation for Innovation, the Government of Ontario, and Queen's

University.

Funding sources

D.G.T. is funded by Queen's University and internal awards in the Department of Biomedical and

Molecular Sciences. Q.L.D. receives funding from the Canadian Institutes of Health Research and

Queen's University National Scholar Award. The funders did not play any role in the study design,

data collection, data analysis, data interpretation, writing of the manuscript or decision to publish

results.

Data statement

Transcriptomics, genomics, and clinical data used for the analysis of the TCGA-OV and TCGA-

BLCA cohorts can be accessed and downloaded from the Genomic Data Commons (GDC) Data

Portal (https://portal.gdc.cancer.gov/). Gene expression and clinical data of the AOCS cohort can

be accessed and downloaded from the Gene Expression Omnibus (GEO) database

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE9899).

Ethics declarations

Controlled access to datasets from The Cancer Genome Atlas (TCGA) was provided by the

National Institute of Health (NIH). The terms of access are outlined in the Data Use Certification

Agreement: https://dbgap.ncbi.nlm.nih.gov/aa/wga.cgi?view_pdf&wlid=10654&tlsid=274.

Details of the Human subject protection and data access policies implemented by TCGA are as

described: https://www.cancer.gov/about-nci/organization/ccg/research/structural-

genomics/tcga/history/policies/tcga-human-subjects-data-policies.pdf. Local ethics clearance for

human subject research was granted by Queen's University.

Declaration of interest statement

None declared.

References

[1] Dion L, Carton I, Jaillard S, Nyangoh Timoh K, Henno S, Sardain H, et al. The Landscape
and Therapeutic Implications of Molecular Profiles in Epithelial Ovarian Cancer. J Clin Med.
2020;9.
[2] Davis A, Tinker AV, Friedlander M. "Platinum resistant" ovarian cancer: what is it, who to
treat and how to measure benefit? Gynecol Oncol. 2014;133:624-31.
[3] Freimund AE, Beach JA, Christie EL, Bowtell DDL. Mechanisms of Drug Resistance in
High-Grade Serous Ovarian Cancer. Hematol Oncol Clin North Am. 2018;32:983-96.
[4] Lloyd KL, Cree IA, Savage RS. Prediction of resistance to chemotherapy in ovarian cancer: a
systematic review. BMC Cancer. 2015;15:117.
[5] Kalow W. Pharmacogenetics and pharmacogenomics: origin, status, and the hope for
personalized medicine. Pharmacogenomics J. 2006;6:162-5.
[6] Cancer Genome Atlas Research N. Integrated genomic analyses of ovarian carcinoma.
Nature. 2011;474:609-15.
[7] Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15:550.
[8] Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis.
BMC Bioinformatics. 2008;9:559.
[9] Choi J, Topouza DG, Tarnouskaya A, Nesdoly S, Koti M, Duan QL. Gene networks and
expression quantitative trait loci associated with adjuvant chemotherapy response in high-grade
serous ovarian cancer. BMC Cancer. 2020;20:413.
[10] Shabalin AA. Matrix eQTL: ultra fast eQTL analysis via large matrix operations.
Bioinformatics. 2012;28:1353-8.

[11] Chang L, Zhou G, Soufan O, Xia J. miRNet 2.0: network-based visual analytics for miRNA
functional analysis and systems biology. Nucleic Acids Res. 2020;48:W244-W51.
[12] Andres-Leon E, Gomez-Lopez G, Pisano DG. Prediction of miRNA-mRNA Interactions
Using miRGate. Methods Mol Biol. 2017;1580:225-37.
[13] Gyorffy B, Lanczky A, Szallasi Z. Implementing an online tool for genome-wide validation
of survival-associated biomarkers in ovarian-cancer using microarray data from 1287 patients.
Endocr Relat Cancer. 2012;19:197-208.
[14] Tothill RW, Tinker AV, George J, Brown R, Fox SB, Lade S, et al. Novel molecular
subtypes of serous and endometrioid ovarian cancer linked to clinical outcome. Clin Cancer Res.
2008;14:5198-208.
[15] Cancer Genome Atlas Research N. Comprehensive molecular characterization of urothelial
bladder carcinoma. Nature. 2014;507:315-22.
[16] Aguirre-Gamboa R, Gomez-Rueda H, Martinez-Ledesma E, Martinez-Torteya A, Chacolla-
Huaringa R, Rodriguez-Barrientos A, et al. SurvExpress: an online biomarker validation tool and
database for cancer gene expression data using survival analysis. PLoS One. 2013;8:e74250.
[17] Kim H, Bhattacharya A, Qi L. Endoplasmic reticulum quality control in cancer: Friend or
foe. Semin Cancer Biol. 2015;33:25-33.
[18] Tian J, Liu R, Qu Q. Role of endoplasmic reticulum stress on cisplatin resistance in ovarian
carcinoma. Oncol Lett. 2017;13:1437-43.
[19] Bastola P, Neums L, Schoenen FJ, Chien J. VCP inhibitors induce endoplasmic reticulum
stress, cause cell cycle arrest, trigger caspase-mediated cell death and synergistically kill ovarian
cancer cells in combination with Salubrinal. Mol Oncol. 2016;10:1559-74.
[20] Koundouros N, Poulogiannis G. Reprogramming of fatty acid metabolism in cancer. Br J
Cancer. 2020;122:4-22.
[21] Nie LY, Lu QT, Li WH, Yang N, Dongol S, Zhang X, et al. Sterol regulatory element-
binding protein 1 is required for ovarian tumor growth. Oncol Rep. 2013;30:1346-54.
[22] Porstmann T, Griffiths B, Chung YL, Delpuech O, Griffiths JR, Downward J, et al.
PKB/Akt induces transcription of enzymes involved in cholesterol and fatty acid biosynthesis via
activation of SREBP. Oncogene. 2005;24:6465-81.
[23] Wang HQ, Altomare DA, Skele KL, Poulikakos PI, Kuhajda FP, Di Cristofano A, et al.
Positive feedback regulation between AKT activation and fatty acid synthase expression in
ovarian carcinoma cells. Oncogene. 2005;24:3574-82.
[24] Wheeler LJ, Watson ZL, Qamar L, Yamamoto TM, Sawyer BT, Sullivan KD, et al. Multi-
Omic Approaches Identify Metabolic and Autophagy Regulators Important in Ovarian Cancer
Dissemination. iScience. 2019;19:474-91.
[25] Pinto BAS, Franca LM, Laurindo FRM, Paes AMA. Unfolded Protein Response: Cause or
Consequence of Lipid and Lipoprotein Metabolism Disturbances? Adv Exp Med Biol.
2019;1127:67-82.
[26] Hao D, Liu J, Chen M, Li J, Wang L, Li X, et al. Immunogenomic Analyses of Advanced
Serous Ovarian Cancer Reveal Immune Score is a Strong Prognostic Factor and an Indicator of
Chemosensitivity. Clin Cancer Res. 2018;24:3560-71.
[27] Mahadevan NR, Rodvold J, Sepulveda H, Rossi S, Drew AF, Zanetti M. Transmission of
endoplasmic reticulum stress and pro-inflammation from tumor cells to myeloid cells. Proc Natl
Acad Sci U S A. 2011;108:6561-6.

[28] Cubillos-Ruiz JR, Silberman PC, Rutkowski MR, Chopra S, Perales-Puchalt A, Song M, et
al. ER Stress Sensor XBP1 Controls Anti-tumor Immunity by Disrupting Dendritic Cell
Homeostasis. Cell. 2015;161:1527-38.
[29] Herber DL, Cao W, Nefedova Y, Novitskiy SV, Nagaraj S, Tyurin VA, et al. Lipid
accumulation and dendritic cell dysfunction in cancer. Nat Med. 2010;16:880-6.
[30] Wagschal A, Najafi-Shoushtari SH, Wang L, Goedeke L, Sinha S, deLemos AS, et al.
Genome-wide identification of microRNAs regulating cholesterol and triglyceride homeostasis.
Nat Med. 2015;21:1290-7.
[31] Adlakha YK, Khanna S, Singh R, Singh VP, Agrawal A, Saini N. Pro-apoptotic miRNA-
128-2 modulates ABCA1, ABCG1 and RXRalpha expression and cholesterol homeostasis. Cell
Death Dis. 2013;4:e780.
[32] Hedditch EL, Gao B, Russell AJ, Lu Y, Emmanuel C, Beesley J, et al. ABCA transporter
gene expression and poor outcome in epithelial ovarian cancer. J Natl Cancer Inst. 2014;106.
[33] Zheng L, Li L, Lu Y, Jiang F, Yang XA. SREBP2 contributes to cisplatin resistance in
ovarian cancer cells. Exp Biol Med (Maywood). 2018;243:655-62.
[34] Li B, Chen H, Wu N, Zhang WJ, Shang LX. Deregulation of miR-128 in ovarian cancer
promotes cisplatin resistance. Int J Gynecol Cancer. 2014;24:1381-8.
[35] Amini-Farsani Z, Sangtarash MH, Shamsara M, Teimori H. MiR-221/222 promote
chemoresistance to cisplatin in ovarian cancer cells by targeting PTEN/PI3K/AKT signaling
pathway. Cytotechnology. 2018;70:203-13.
[36] Li J, Li Q, Huang H, Li Y, Li L, Hou W, et al. Overexpression of miRNA-221 promotes cell
proliferation by targeting the apoptotic protease activating factor-1 and indicates a poor
prognosis in ovarian cancer. Int J Oncol. 2017.
[37] Sun C, Li N, Zhou B, Yang Z, Ding D, Weng D, et al. miR-222 is upregulated in epithelial
ovarian cancer and promotes cell proliferation by downregulating P27(kip1.). Oncol Lett.
2013;6:507-12.
[38] Dai R, Li J, Liu Y, Yan D, Chen S, Duan C, et al. miR-221/222 suppression protects against
endoplasmic reticulum stress-induced apoptosis via p27(Kip1)- and MEK/ERK-mediated cell
cycle regulation. Biol Chem. 2010;391:791-801.
[39] Zhang Y, Wu J, Liang JY, Huang X, Xia L, Ma DW, et al. Association of serum lipids and
severity of epithelial ovarian cancer: an observational cohort study of 349 Chinese patients. J
Biomed Res. 2018;32:336-42.
[40] Li AJ, Elmore RG, Chen IY, Karlan BY. Serum low-density lipoprotein levels correlate
with survival in advanced stage epithelial ovarian cancers. Gynecol Oncol. 2010;116:78-81.
411

Table 1. Characteristics of the HGSOC patient cohorts with mRNA-Seq and miRNA-Seq data from TCGA.

	mRNA-Seq cohort				miRNA-Seq cohort			
	Sensitive	Resistant	P value	All cases	Sensitive	Resistant	P value	All cases
Age								
Mean	57.15	61.98	**0.0043** [a]	59.22	57.76	61.52	**0.018** [a]	59.26
Range	30 - 87	38 - 87		30 - 87	30 - 87	38 - 87		30 - 87
Grade								
G2	18 (15.93 %)	8 (10.26 %)	0.27 [b]	26 (13.61 %)	23 (18.70 %)	8 (9.76 %)	0.14 [b]	31 (15.12 %)
G3/4	95 (84.07 %)	69 (88.46 %)		164 (85.86 %)	99 (80.49 %)	73 (89.02 %)		172 (83.90 %)
Ungraded	0 (0.00 %)	1 (1.28 %)		1 (0.52 %)	1 (0.81 %)	1 (1.22 %)		2 (0.98 %)
Stage								
II	7 (6.19 %)	2 (2.56 %)	0.48 [b]	9 (4.71 %)	8 (6.50 %)	2 (2.44 %)	0.41 [b]	10 (4.88 %)
III	91 (80.53 %)	67 (85.9 %)		158 (82.72 %)	99 (80.49 %)	67 (81.71 %)		166 (80.98 %)
IV	15 (13.27 %)	9 (11.54 %)		24 (12.57 %)	16 (13.01 %)	13 (15.85 %)		29 (14.15 %)
Cytoreductive surgery outcome								
Optimal (≤ 10mm)	73 (64.60 %)	54 (69.23 %)	0.23 [c]	127 (66.49 %)	77 (62.60 %)	56 (68.29 %)	0.21 [c]	133 (64.88 %)
Suboptimal (> 10mm)	26 (23.01 %)	20 (25.64 %)		46 (24.08 %)	29 (23.58 %)	21 (25.61 %)		50 (24.39 %)
Unknown	14 (12.39 %)	4 (5.13 %)		18 (9.42 %)	17 (13.82 %)	5 (6.10 %)		22 (10.73 %)
Overall survival								
Number of mortality events	48 (42.48 %)	60 (76.92 %)	**2.35E-06** [c]	108 (56.54 %)	55 (44.72 %)	66 (80.49 %)	**0.0002** [c]	121 (59.02 %)
Median months	52.04	28.20	**6.97E-10** [a]	39.87	51.91	28.20	**1.13E-10** [a]	39.50
95% CI	50.07 - 54.01	26.22 - 30.18		37.90 - 41.85	49.94 - 53.88	26.23 - 30.18		37.53 - 41.47
Adjuvant chemotherapy regimen								
Platinum agent only	4 (3.54 %)	3 (3.85 %)	1 [b]	7 (3.66 %)	4 (3.25 %)	4 (4.88 %)	0.71 [b]	8 (3.90 %)
Platinum/Taxane combination	109 (96.46 %)	75 (96.15 %)		184 (96.34 %)	119 (96.75 %)	78 (95.12 %)		197 (96.10 %)
Primary tumor mRNA subtype								
Differentiated	37 (32.74 %)	22 (28.21 %)	0.49 [b]	59 (30.89 %)	42 (34.15 %)	24 (29.27 %)	0.68 [c]	66 (32.20 %)
Immunoreactive	23 (20.35 %)	10 (12.82 %)		33 (17.28 %)	24 (19.51 %)	13 (15.85 %)		37 (18.05 %)
Mesenchymal	22 (19.47 %)	18 (23.08 %)		40 (20.94 %)	24 (19.51 %)	18 (21.95 %)		42 (20.49 %)
Proliferative	30 (26.55 %)	27 (34.62 %)		57 (29.84 %)	33 (26.83 %)	27 (32.93 %)		60 (29.27 %)
Subtype unknown	1 (0.88 %)	1 (1.28 %)		2 (1.05 %)	0 (0.00 %)	0 (0.00 %)		0 (0.00 %)
Total patients	113	78		191	123	82		205

[a] p-value based on t-test; [b] p-value based on Fisher's Exact test; [c] p-value based on Chi-Squared test; CI, Confidence Interval

Table 2. Significantly enriched annotation terms for differentially expressed mRNA transcripts.

Term name	Adjusted P value	Number of query genes in term
CD22 mediated BCR regulation	4.42E-45	30
Complement activation, classical pathway	3.07E-44	36
Antigen binding	1.74E-42	39
Scavenging of heme from plasma	1.03E-38	28
Immune response-regulating cell surface receptor signaling pathway involved in phagocytosis	3.89E-34	30
Fc-gamma receptor signaling pathway involved in phagocytosis	3.89E-34	30
Serine-type peptidase activity	7.94E-28	28
Cell surface interactions at the vascular wall	2.63E-27	30
Blood microparticle	2.57E-21	25
Leukocyte migration	3.36E-20	33
Immunoglobulin complex	2.24E-09	10
Cell periphery	3.36E-09	80
Plasma membrane	1.37E-08	78
Extracellular exosome	5.22E-07	43
Immunoglobulin receptor binding	5.98E-07	6
Phagocytosis, recognition	1.33E-06	6
External side of plasma membrane	1.48E-06	14
Regulation of B cell activation	8.13E-06	11
Phagocytosis, engulfment	8.91E-06	6
Ribosome	9.44E-05	6
Cytosolic large ribosomal subunit	2.91E-04	6
Ribosome, cytoplasmic	5.57E-04	6
60S ribosomal subunit, cytoplasmic	8.99E-04	5
Formation of a pool of free 40S subunits	9.17E-04	7
Peptide chain elongation	1.39E-03	6
Viral mRNA Translation	1.39E-03	6
Selenocysteine synthesis	1.79E-03	6
Eukaryotic Translation Termination	1.79E-03	6
Abnormal leukocyte morphology	2.02E-03	7
Nonsense Mediated Decay (NMD) independent of the Exon Junction Complex (EJC)	2.02E-03	6
SRP-dependent cotranslational protein targeting to membrane	3.09E-03	6
Nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	4.14E-03	7
Abnormal immunoglobulin level	4.44E-03	5
SRP-dependent cotranslational protein targeting to membrane	5.18E-03	6
Structural constituent of ribosome	6.70E-03	7
Factor: ZNF491; motif: NGKGGKCAGT	1.13E-02	7
Agammaglobulinemia	1.32E-02	2
Verrucae	2.39E-02	2
Cellulitis	4.10E-02	2
TRBP containing complex (DICER, RPL7A, EIF6, MOV10 and subunits of the 60S ribosomal particle)	4.33E-02	4
Nop56p-associated pre-rRNA complex	4.57E-02	5

Table 3. Significantly enriched annotation terms for differentially expressed miRNA isoforms.

Term name	Adjusted P value	Number of query miRNAs in term
mRNA binding involved in posttranscriptional gene silencing	4.98E-16	10
Gene silencing by miRNA	7.12E-14	10
MicroRNAs in cancer	1.09E-12	10
Negative regulation of cardiac muscle cell proliferation	5.21E-08	4
Positive regulation of cardiac muscle tissue regeneration	2.10E-07	3
Re-entry into mitotic cell cycle	5.25E-07	3
Negative regulation of cell growth involved in cardiac muscle cell development	8.65E-06	3
Positive regulation of lung blood pressure	2.05E-04	2
Negative regulation of cardiac muscle cell apoptotic process	3.12E-04	3
Regulation of vasculature development	3.32E-03	5
Negative regulation of autophagy	8.70E-03	3
Extracellular space	1.03E-02	9
Positive regulation of connective tissue replacement	2.14E-02	2
Negative regulation of nitric oxide metabolic process	2.45E-02	2
Blood vessel morphogenesis	3.25E-02	5
Positive regulation of protein kinase B signaling	3.97E-02	3

Table 4. Significantly enriched terms in significant mRNA transcript co-expression networks.

Network name	Term name	P value	Number of query genes in term
lavenderblush3	Protein modification by small protein conjugation or removal	2.40E-02[a]	9
	Protein modification by small protein conjugation	2.71E-03[a]	9
	Protein ubiquitination	1.60E-03[a]	9
	Factor: GABP-alpha; motif: CTTCCK	6.72E-03[a]	26
darkolivegreen	Protein-containing complex	4.60E-02[a]	37
	Biosynthesis of unsaturated fatty acids	5.20E-03[b]	2
	Fatty acid degradation	1.34E-02[b]	2
	Notch signaling pathway	1.58E-02[b]	2
	Non-alcoholic fatty liver disease (NAFLD)	2.19E-02[b]	3
	Ribosome	2.35E-02[b]	3
	RNA transport	2.85E-02[b]	3
	Synthesis and degradation of ketone bodies	3.93E-02[b]	1
	Peroxisome	4.37E-02[b]	2

[a] Benjamini-Hochberg adjusted p value (g:Profiler); [b] non-adjusted p value (Enrichr)

Table 5. Significantly enriched terms in significant miRNA isoform co-expression networks.

Network name	Term name	Adjusted P value	Number of query miRNAs in term
ivory	mRNA binding involved in posttranscriptional gene silencing	1.16E-09	7
	Gene silencing by miRNA	3.61E-08	7
	Extracellular space	4.24E-05	10
	MicroRNAs in cancer	9.55E-05	5
	Regulation of lipoprotein transport	1.22E-03	2
	Negative regulation of cholesterol efflux	2.69E-02	2
	Negative regulation of lipoprotein particle clearance	4.27E-02	2
plum	Positive regulation of Schwann cell proliferation involved in axon regeneration	2.55E-06	2
	Negative regulation of hematopoietic stem cell proliferation	2.55E-06	2
	Positive regulation of Schwann cell migration	7.66E-06	2
	Negative regulation of TRAIL-activated apoptotic signaling pathway	7.66E-06	2
	Negative regulation of cell adhesion molecule production	1.53E-05	2
	Negative regulation by host of viral genome replication	5.36E-05	2
	Negative regulation of leukocyte adhesion to vascular endothelial cell	1.40E-04	2
	Negative regulation of cytokine production involved in inflammatory response	2.32E-04	2
	Positive regulation of erythrocyte differentiation	1.19E-03	2
	Positive regulation of G1/S transition of mitotic cell cycle	1.61E-03	2
	Positive regulation of vascular smooth muscle cell proliferation	2.53E-03	2
	Positive regulation of epithelial to mesenchymal transition	3.00E-03	2
	Negative regulation of muscle cell differentiation	5.15E-03	2
	Negative regulation of translation, ncRNA-mediated	6.17E-03	2
	Negative regulation of sprouting angiogenesis	8.07E-03	2
	Positive regulation of epithelial cell migration	3.33E-02	2
	Positive regulation of protein kinase B signaling	3.58E-02	2
	mRNA binding involved in posttranscriptional gene silencing	4.25E-02	2

Table 6. Predicted and validated mRNA - miRNA network interactions.

Integrated Networks	Gene ID	mRNA ID	miRNA ID	Spearman's ρ	P value	Interaction type
lavenderblush3 - plum	*PSIP1*	ENST00000397519	hsa-mir-222-3p	-0.32	0.000021	Experimental validation
	CDC37L1	ENST00000381854	hsa-mir-222-3p	-0.31	0.000042	Computational prediction
	CDC37L1	ENST00000381854	hsa-mir-221-3p	-0.27	0.00034	Computational prediction
	CDC37L1	ENST00000381854	hsa-mir-221-5p	-0.27	0.00034	Computational prediction
	NFIB	ENST00000380959	hsa-mir-221-3p	-0.27	0.00039	Experimental validation
	SMU1	ENST00000397149	hsa-mir-221-5p	-0.26	0.00082	Experimental validation
	VCP	ENST00000358901	hsa-mir-222-3p	-0.25	0.00089	Experimental validation
	CAAP1	ENST00000333916	hsa-mir-222-5p	-0.22	0.0041	Computational prediction
	CAAP1	ENST00000333916	hsa-mir-221-5p	-0.21	0.0076	Computational prediction
	TOPORS	ENST00000360538	hsa-mir-221-3p	-0.19	0.014	Experimental validation
	SLC25A51	ENST00000496760	hsa-mir-222-3p	-0.19	0.016	Experimental validation
	PPAPDC2	ENST00000381883	hsa-mir-222-5p	-0.18	0.021	Computational prediction
	VCP	ENST00000358901	hsa-mir-221-3p	-0.17	0.023	Experimental validation
	VCP	ENST00000358901	hsa-mir-221-5p	-0.17	0.023	Experimental validation
	NFX1	ENST00000379540	hsa-mir-221-5p	-0.17	0.027	Computational prediction
	PPAPDC2	ENST00000381883	hsa-mir-221-5p	-0.17	0.03	Computational prediction
	DNAJA1	ENST00000330899	hsa-mir-221-5p	-0.16	0.035	Experimental validation
	AK3	ENST00000381809	hsa-mir-222-5p	-0.16	0.037	Computational prediction
darkolivegreen - ivory	*RPL18*	ENST00000552347	hsa-mir-212-3p	-0.29	0.00017	Experimental Validation
	MRPL55	ENST00000411464	hsa-miR-1306-5p	-0.25	0.001	Computational prediction
	FASTK	ENST00000466855	hsa-mir-140-3p	-0.25	0.0013	Experimental Validation
	RPL18	ENST00000552347	hsa-mir-361-3p	-0.24	0.0021	Experimental Validation
	ACBD4	ENST00000321854	hsa-miR-128-2-5p	-0.22	0.0041	Computational prediction
	OAZ1	ENST00000581150	hsa-mir-361-3p	-0.22	0.0047	Experimental Validation
	ACBD4	ENST00000321854	hsa-miR-128-1-5p	-0.21	0.0059	Computational prediction
	MYL12A	ENST00000578611	hsa-mir-103a-3p	-0.17	0.023	Experimental Validation
	MYL12A	ENST00000578611	hsa-mir-107	-0.17	0.023	Experimental Validation
	HTRA2	ENST00000484352	hsa-mir-1306-5p	-0.17	0.029	Experimental Validation
	ACADVL	ENST00000579425	hsa-mir-103a-3p	-0.17	0.03	Experimental Validation
	ACADVL	ENST00000579425	hsa-mir-107	-0.17	0.03	Experimental Validation
	SOD1	ENST00000470944	hsa-mir-140-3p	-0.16	0.037	Experimental Validation
	RHOT2	ENST00000569675	hsa-mir-1306-5p	-0.16	0.044	Experimental Validation
	TSSK6	ENST00000360913	hsa-miR-212-5p	-0.15	0.048	Computational prediction
	RCC1	ENST00000373832	hsa-miR-1306-3p	-0.15	0.048	Computational prediction
	RCC1	ENST00000373832	hsa-miR-1306-5p	-0.15	0.048	Computational prediction

Figure Legends

Figure 1. Differential expression analysis of mRNA and miRNA transcripts. (A) Significant differentially expressed mRNA transcripts (n=196) are shown in red. The Benjamini-Hochberg adjusted significance threshold is indicated by the dashed line. Transcripts with a positive fold change are upregulated in resistant patients, whereas transcripts with a negative fold change are downregulated in resistant patients. (B) Significant differentially expressed miRNA isoforms (n=21) are shown in red. The Benjamini-Hochberg adjusted significance threshold is indicated by the dashed line. Transcripts with a positive fold change are upregulated in resistant patients, whereas transcripts with a negative fold change are downregulated in resistant patients.

Figure 2. Weighted transcript co-expression network analysis of mRNA and miRNA transcripts. (A) Heatmap of the correlation between the 58 mRNA co-expression modules with platinum-based chemotherapy resistance, red indicating a positive effect and blue a negative effect. P-values are indicated for each module. Two mRNA co-expression modules are significantly associated with chemotherapy resistance (*lavenderblush3*, *darkolivegreen*). (B) The mRNA modules *lavenderblush3* and *darkolivegreen* are visualized in their respective colors. Each node represents a transcript, and each edge represents a connection or co-expression. The distance between nodes is the connection weight, where more similar transcripts are plotted closer. (C) Heatmap of the correlation between the 100 miRNA modules with platinum-based chemotherapy resistance, red indicating a positive effect and blue a negative effect. P-values are indicated for each module. Three miRNA co-expression modules are significantly associated with chemotherapy resistance (*ivory*, *lightcoral*, and *plum*). (D) The miRNA modules *lightcoral*, *plum*, and *ivory* are visualized in their respective colors. Each node represents one miRNA isoform, and

each edge represents a connection or co-expression. The distance between nodes is the connection

weight, where more similar transcripts are plotted closer.

Figure 3. Integration of mRNA - miRNA networks. The mRNA transcripts (oval nodes) from

the *lavenderblush3* and *darkolivegreen* modules are arranged based on co-expression similarity

(grey edges). Four common miRNA isoforms (rectangular nodes) were validated (orange solid

edges) or predicted (orange dashed edges) to regulate the expression of genes in the *lavenderblush3*

network module, while 10 miRNA isoforms were validated or predicted to regulate genes in the

darkolivegreen module. Genes targets of miRNAs indicated by orange border.

Figure 4. Replication analysis of the gene co-expression networks. (A-B) Prognostic Index

Kaplan-Meier analysis of patient PFS association with the prognostic index of mRNA networks

lavenderblush3 (A) and *darkolivegreen* (B). (C-E) Prognostic Index Kaplan-Meier analysis of

patient OS association with the prognostic index of miRNA networks *ivory* (C), *plum* (D), and

lightcoral (E). (F-G) Kaplan-Meier analysis of patient OS with the expression of miR-221 (F) and

miR-222 (G). PFS, progression-free survival; OS, overall survival.

Supporting Information

Supplementary Methods

Supplementary Data 1. mRNA-microRNA interaction annotations.

Table S1. mRNA differential expression analysis results.

Table S2. microRNA differential expression analysis results.

Table S3. Significant WGCNA module mRNAs and correlation to chemotherapy resistance.

Table S4. Significant WGCNA module microRNAs and correlation to chemotherapy resistance.

Table S5. eQTL analysis results and annotation.

Table S6. Kaplan-Meier validation results for differentially expressed mRNAs.

Table S7. Kaplan-Meier validation results for differentially expressed microRNAs.

Figure 1

Figure 2

Figure 3

Figure 4