Integrated biological networks associated with platinum-based chemotherapy response in ovarian cancer

Danai Georgia Topouza[a], Jihoon Choi[a], Sean Nesdoly[b], Anastasiya Tarnouskaya[b], Christopher J.B. Nicol[a,c,d], Qing Ling Duan[a,b]

[a]Department of Biomedical and Molecular Sciences, Queen's University, Kingston, Ontario, Canada; [b]School of Computing, Queen's University, Kingston, Ontario, Canada; [c]Department of Pathology and Molecular Medicine, Queen's University, Kingston, Ontario, Canada; [d]Division of Cancer Biology and Genetics, Queen's University Cancer Research Institute, Queen's University, Kingston, Ontario, Canada.

Correspondence: Qing Ling Duan (qingling.duan@queensu.ca)

Abstract

Ovarian cancer is a highly lethal gynecologic cancer, partly due to resistance to platinum-based chemotherapy reported among 20-30% of patients. This study aims to elucidate the biological mechanisms underlying chemotherapy resistance, which remain poorly understood. Using mRNA and microRNA sequencing data from high-grade serous ovarian cancer (HGSOC) patients from The Cancer Genome Atlas, we identified transcripts and networks associated with chemotherapy response. In total, 196 differentially expressed mRNAs were enriched for adaptive immunity and translation, and 21 differentially expressed microRNAs were associated with angiogenesis. Moreover, co-expression network analysis identified two mRNA networks associated with chemotherapy response, which were enriched for ubiquitination and lipid metabolism, as well as three associated microRNA networks enriched for lipoprotein transport and oncogenic pathways. In addition, integrative analyses of these sequence datasets revealed potential regulation of the mRNA networks by the associated microRNAs and single nucleotide polymorphisms. Thus, we report novel transcriptional networks and pathways associated with resistance to platinum-based chemotherapy among HGSOC patients. These results aid our understanding of the effector networks and regulators of chemotherapy response, which will improve drug efficacy and identify novel therapeutic targets for ovarian cancer.

Keywords

Chemotherapy resistance, high-grade serous ovarian cancer (HGSOC), RNA-sequencing, co-expression network analysis, expression quantitative trait loci (eQTL)

Highlights

- mRNA and miRNA transcript networks associated with adjuvant chemotherapy response

- Integrative omics data analysis identified regulatory miRNAs and eQTLs

- Protein ubiquitination and immune activation associated with chemosensitivity

- Increased translation and lipid metabolism may promote chemoresistance

Abbreviations

AOCS: Australian Ovarian Cancer Study

BLCA: Bladder Urothelial Carcinoma

eQTL: Expression Quantitative Trait Locus

ER: Endoplasmic Reticulum

ERAD: Endoplasmic-Reticulum-Associated Protein Degradation

GLM: Generalized Linear Model

HCC: Hepatocellular Carcinoma

HDL: High-Density Lipoprotein

HGSOC: High-Grade Serous Ovarian Cancer

LDL: Low-Density Lipoprotein

miRNA: microRNA

mRNA: Messenger RNA

OC: Ovarian Cancer

OS: Overall Survival

PFI: Platinum-Free Interval

PFS: Progression-Free Survival

PI: Prognostic Index

RNA-Seq: RNA-Sequencing

SNP: Single Nucleotide Polymorphism

69   TCGA: The Cancer Genome Atlas

70   TF: Transcription Factor

71   UPR: Unfolded Protein Response

72   WGCNA: Weighted Gene Co-Expression Network Analysis

1. Introduction

Ovarian cancer (OC) is a highly lethal gynecologic cancer accounting for 225,000 new cases and 140,000 deaths worldwide each year, the majority of which are high-grade serous ovarian cancer (HGSOC) cases [1]. The first-line standard of care for HGSOC is debulking surgery and platinum-based adjuvant chemotherapy [2]. However, 20-30% of HGSOC patients are resistant to chemotherapy, while the majority of patients who are initially sensitive relapse with chemoresistant disease within three years [2]. Resistant patients have a significantly shorter overall survival (OS) than sensitive patients, and many experience tumor recurrence within six months of completing chemotherapy [3].

Currently, there is no strategy for predicting response to platinum-based chemotherapy, which reflects our limited understanding of the underlying molecular mechanisms of chemotherapy resistance as reviewed elsewhere [4, 5]. Several studies have reported molecular signatures of chemotherapy resistance in OC patients, with little consensus in the results [6]. The majority of these earlier studies used univariate analysis methods such as differential expression analysis, univariate Cox regression, and t-tests that report single gene associations with chemotherapy response in HGSOC [7-10]. The common assumption of these methods is that chemotherapy response is driven by a single gene. However, it is well established that chemotherapy response, like other pharmacogenomics traits, is a complex multifactorial trait modulated by multiple genes contributing to one or more biological pathways [11, 12].

In this study, we apply both univariate and multivariate analysis methods to identify genes and gene networks associated with chemotherapy response in HGSOC. Using mRNA and microRNA (miRNA) sequencing datasets from HGSOC patients of The Cancer Genome Atlas (TCGA) [13], we identify novel transcriptomic differences between patients we define as

sensitive or resistant to platinum-based chemotherapy. Here, we provide the first evidence of

novel mRNA and miRNA signalling pathways and networks associated with chemotherapy

response in HGSOC patients. We further unveil miRNAs and expression quantitative trait loci

(eQTLs) that correlate with the expression of significant transcripts and networks. These findings

are validated against two independent cancer cohorts and may provide important therapeutic

strategies for regulating platinum-based chemotherapy response in HGSOC patients.

2. Materials and Methods

2.1 Chemotherapy response classification

Sequencing of mRNA and miRNA from the tumors of 191 and 205 HGSOC patients,

respectively, prior to chemotherapy were obtained from TCGA [13]. Patients who received

platinum-based adjuvant chemotherapy following cytoreductive surgery were selected and

separated into sensitive or resistant groups based on their platinum-free interval (PFI) (**Table 1**;

see details in **Supplemental Methods**). Sensitive patients remained cancer-free for at least 12

months after completion of chemotherapy, whereas resistant patients experienced cancer

recurrence within 6 months of completing chemotherapy. Patients with a recurrence between six

months to one year were excluded from analysis in order to enrich for genetic differences

between the response groups.

2.2 Processing of RNA sequencing datasets

Raw mRNA-sequence reads were obtained from the TCGA database [13] on August 31st, 2017,

as FASTQ files (i.e. level 1 data), filtered for base-quality, aligned, and quantified as described

in **Supplemental Methods**. MiRNA-sequencing reads from the same TCGA-HGSOC cohort

were obtained as pre-aligned and quantified files (i.e. level 3 data) from the Broad Institute

Genome Data Analysis Center Firehose [14] on September 17th, 2018 (TCGA data version

2016_01_28 for OV). Both mRNA and miRNA expression datasets underwent outlier control,

normalization, and non-specific filtering, resulting in 49,116 mRNA transcripts and 4,479

miRNA transcripts for inclusion in further analysis (details in **Supplemental Methods**).

2.3 Differential expression analysis

Differentially expressed mRNA transcripts and miRNA isoforms between sensitive and resistant

patients were detected using a negative binomial generalized linear model (GLM) in the DESeq2

R package [15]. This analysis was performed for each individual transcript and controlled for

patients' ages at time of diagnosis, as resistant patients were significantly older (**Table 1**). The

Benjamini-Hochberg method was applied to correct for multiple testing.

2.4 Weighted co-expression network analysis

The weighted gene co-expression network analysis (WGCNA) R package [16] was used, which

applies an unsupervised machine learning approach to identify modules of co-expressed mRNA

and miRNA transcripts. Principal component analysis of the expression of all transcripts within

each module was calculated, resulting in a value known as the eigengene. Module eigengenes

were then used to determine association with chemotherapy response using a GLM, adjusted for

patients' age as a covariate (**Supplemental Methods**). Cytoscape was used for visualisation of

the gene networks [17].

2.5 Pathway enrichment analysis

Pathway enrichment analysis softwares g:Profiler [18] (rev 1760, database versions Ensembl 93

and Ensembl Genomes 40) and Enrichr [19] were used to determine over-representation of

biological pathways from lists of differentially expressed transcripts (mRNA and miRNAs), and

co-expressed networks. In addition, these bioinformatics tools identified enrichment of

regulatory motifs, known protein complexes and human phenotype ontologies.

2.6 eQTL analysis

Single nucleotide polymorphisms (SNPs) from TCGA-HGSOC patients were genotyped using

the Affymetrix SNP Array 6.0. SNP data was phased and imputed as described by Choi et al.

[20]. The imputed data underwent quality control and linkage disequilibrium-based pruning,

retaining 1,722,608 common SNPs (minor allele frequency > 0.05) for analysis (**Supplemental**

Methods, Figure A.3). This genomics data was integrated with mRNA-seq data of 167 patients

and with miRNA-seq data of 178 patients to identify SNPs correlated with transcript expression

(i.e. eQTLs) using the MatrixEQTL R package [21]. We tested common SNPs located within 1

Mb of recorded transcripts [22], which included 1,676,365 proximal common SNPs tested for

association with mRNAs, and 356,189 proximal common SNPs tested for association with

miRNAs (**Supplemental Methods**). Functional annotation of resulting eQTLs was performed

using HaploReg v4.1 [23].

2.7 mRNA-microRNA data integration

Potential regulation of mRNA networks by miRNAs was tested using data from 165 patients

with both mRNA and miRNA data available. We used the Spearman correlation of module

eigengenes from mRNA and miRNA co-expression networks. Results were validated using

miRnet [24, 25], a database of experimentally validated mRNA-miRNA interactions, and

miRGate [26], a database of predicted mRNA-miRNA interactions based on ribonucleotide motif

matching.

2.8 Replication cohort and analysis

Replication of differentially expressed transcripts used the Kaplan-Meier analysis tool KM

Plotter [27] to test the association of each transcript with progression-free survival (PFS) or OS

in two independent cancer cohorts. First, mRNA data generated from microarrays were analyzed

from tumor samples from the Australian Ovarian Cancer Study (AOCS; GSE9891; n=285) [28].

Next, a miRNA dataset from an independent ovarian cancer cohort was not available; we opted

to use the bladder urothelial carcinoma cohort (BLCA) (n=409) from TCGA, given that bladder

cancer is also typically treated with platinum-based chemotherapy [29]. OS was used as the

target phenotype, as PFS was not available for this cohort. Replication of transcript networks

used the Prognostic Index (PI) estimation method as described by Aguirre-Gamboa et al. [30]

using data from the above cohorts (**Supplemental Methods**).

3. Results

3.1 Differentially expressed mRNAs/miRNAs between sensitive and resistant patients are

involved in immune response, translation, and vasculature development

Differential expression analysis identified 196 mRNAs associated with platinum-based

chemotherapy response in HGSOC patients (adjusted $p<0.05$) that map to 190 unique genes

(**Supplementary Table 2**). A total of 88 transcripts are upregulated in resistant patients, while

108 are downregulated (**Figure 1A**). Pathway enrichment analysis of these associated transcripts

using g:Profiler indicated enrichment of 41 annotation terms, including CD22 mediated B-cell

receptor regulation, complement activation, and peptide chain elongation (**Table 2,**

Supplementary Data 1).

Differential miRNA expression analysis revealed 21 differentially expressed miRNA

isoforms (adjusted $p<0.05$), which map to 16 unique miRNAs (**Figure 1B, Supplementary**

Table 3). Pathway enrichment analysis of these miRNA isoforms using g:Profiler revealed 16

pathways and cellular functions such as blood vessel morphogenesis and negative regulation of

autophagy (**Table 3, Supplementary Data 1**).

3.2 Two mRNA co-expression networks involved in protein ubiquitination and fatty acid

metabolism are associated with chemotherapy response

WGCNA resulted in 58 co-expression modules from the mRNA transcripts (**Supplementary**

Table 4), of which two were associated with platinum-based chemotherapy response

(**Supplementary Table 5**). First, the *lavenderblush3* module is downregulated or negatively

associated with chemotherapy resistance (p-value=0.016; **Figure 2A**). This module contains 39

transcripts, mapping to 31 unique genes (**Figure 2B**). Pathway analysis using g:Profiler indicates

enrichment of three biological pathways related to protein ubiquitination, and the transcription

factor (TF) GABP-alpha (GA Binding Protein Transcription Factor Subunit Alpha) binding

motif (**Table 4**). Second, the *darkolivegreen* module is significantly upregulated in

chemoresistant patients (p-value=0.032) (**Figure 2A**) and contains 82 transcripts mapping to 80

unique genes (**Figure 2B**). Enrichr pathway analysis shows that this module is significantly

enriched for the protein-containing complex term, as well as for 8 pathways involved in fatty

acid metabolism with nominal significance (**Table 4**). Moreover, the *darkolivegreen* module is

enriched for 41 TF motifs (**Supplementary Data 1**), with c-Ets-2 (ETS Proto-Oncogene 2) being

the most significantly enriched TF motif.

3.3 Three miRNA co-expression networks involved in lipid transport and oncogenic pathways

are associated with chemotherapy response

WGCNA analysis of the miRNA dataset constructed 100 co-expression network modules

(**Supplementary Table 6**), of which three are associated with chemotherapy response

(**Supplementary Table 7**). The *ivory* (p-value=0.0097) and *lightcoral* (p-value=0.042) modules

are downregulated or negatively associated with chemotherapy resistance, while the third *plum*

network (p-value = 0.045) is positively associated with chemotherapy resistance (**Figure 2C**).

The *ivory* network consisted of 25 miRNA isoforms mapping to 11 unique miRNAs, which are

enriched for 7 pathways and biological functions, including regulation of lipoprotein transport

and negative regulation of cholesterol efflux (**Table 5**). The *lightcoral* module consists of 17

isoforms of miR-187 and the *plum* network consists of 17 isoforms of miR-221 and miR-222

(**Figure 2D**). While no pathway annotations were derived for the *lighcoral* module, the *plum*

module is enriched for 18 pathways with a variety of oncogenic functions, such as negative

regulation of the TRAIL-activated apoptotic pathway, negative regulation of cell adhesion

molecule production, negative regulation of cytokine production involved in inflammatory

response, and positive regulation of protein kinase B signaling (**Table 5, Supplementary Data**

1).

3.4 Germline eQTLs may regulate the expression of associated mRNAs, miRNAs, and networks

We identified 268 unique germline cis-eQTLs associated with the expression of mRNAs and

miRNAs that are differentially expressed or members of significant co-expression networks. 248

SNPs are associated with the expression of 55 significant mRNAs, and 20 SNPs are associated

with the expression of 7 significant miRNAs (**Supplementary Table 8**). Of the 268 eQTLs, 126

are previously known whereas 118 are novel, while 24 are yet not recorded in the HaploReg

database. The majority (227) of these eQTLs alter regulatory motifs. Finally, 67 eQTLs are

associated with 94 human phenotypes from published Genome-Wide Association Studies

(GWAS). The most common phenotypes associated with these SNPs are related to lipid

metabolism, such as high-density lipoprotein (HDL) cholesterol, triglyceride change with statins,

and low-density lipoprotein (LDL) cholesterol.

3.5 Network integration reveals novel miRNA-mediated regulation of chemotherapy response

mechanisms

Next, we tested for potential interactions between the two significant mRNA modules and the

three miRNA modules. The *plum* miRNA network significantly correlates with the

lavenderblush3 mRNA network (Spearman's ρ = -0.26, p < 0.001), and the *ivory* miRNA

network significantly correlates with the *darkolivegreen* mRNA network (Spearman's ρ = -0.17,

p = 0.023) (**Supplementary Table 9**). Annotations using miRnet and miRbase determined that

20 of these mRNA-miRNA interactions are experimentally validated, while 18 others are

supported by *in silico* predictions (**Table 6, Figure 3**).

3.6 Replication of study results in two independent cancer cohorts

Kaplan-Meier analysis was used to replicate the association of differentially expressed mRNAs

from the TCGA cohort in the second AOCS ovarian cancer cohort. A total of 140 microarray

probes in the AOCS dataset (73.68%) matched to differentially expressed genes from TCGA. Of

these, 28 (20%) replicated with the same direction of effect (**Supplementary Table 10**). The

same analysis was used to test the association of differentially expressed miRNAs from the

TCGA-HGSOC cohort in the BLCA cohort from TCGA. Out of the 16 differentially expressed

miRNAs, 13 (81.25%) were available in the bladder cancer miRNA-Seq dataset and 7 (53.85%)

replicated with the same direction of effect (**Supplementary Table 11**).

Our network analysis results were replicated by applying the PI estimation method [30].

The *lavenderblush3* and *darkolivegreen* mRNA network modules replicated in the AOCS cohort

(p < 0.0001) (**Figure 4A, B**). In addition, the *ivory* miRNA network module replicated in the

TCGA-BLCA cohort (p = 0.012), while the *lightcoral* and *plum* modules did not replicate

(**Figure 4C-E**). However, differential expression of miR-221 and miR-222, which make up the

plum network, replicated in this second cohort using Kaplan-Meier analysis (p = 0.022 and p =

0.037, respectively) (**Figure 4F, G**).

4. Discussion

In this study, we analyzed mRNA and miRNA sequencing data from chemotherapy-naïve tumors

of HGSOC patients to identify transcripts and transcript networks associated with chemotherapy

response. Our findings implicate novel and known biological pathways that replicated in

independent cancer cohorts. In addition, we identified potential interactions among miRNAs and

mRNAs, as well as the potential regulation of gene transcripts by SNPs (i.e. expression

quantitative trait loci or QTLs). Thus, our results provide an integrative, multi-omics view of

gene networks associated with chemotherapy response. This study identifies novel biological

targets for further investigation, which will improve our understanding of the mechanisms

underlying chemotherapy resistance in HGSOC patients.

We identified one mRNA co-expression network (*lavenderblush3*) significantly

upregulated in platinum sensitive patients, which replicated in the AOCS ovarian cancer cohort.

The genes within this module are critical for protein processing in the endoplasmic reticulum

(ER) and ubiquitin-mediated proteolysis. We also detected a significant downregulation of genes

responsible for translation initiation within sensitive patients. These findings suggest that the

activation of the unfolded protein response (UPR), a cellular process responsible for resolving

ER stress, may be important in chemotherapy response. The UPR alleviates ER stress through

several pathways, including increased endoplasmic-reticulum-associated protein degradation

(ERAD) to remove misfolded proteins, and inhibition of translation to reduce protein load in the

ER [31]. Our findings indicate that the UPR may be increasingly activated in sensitive patients

compared to resistant cases. ER stress, and the ER stress response involving ERAD, promote

chemotherapy resistance in ovarian and other cancers [32, 33]. However, the upregulation of

ERAD genes such as *VCP* (Valosin Containing Protein), a member of our *lavenderblush3*

network, is associated with longer OS and platinum sensitivity in HGSOC patient cohorts [20,

34]. Eighteen of 31 module genes were replicated in an earlier study of the microarray dataset

from the same TCGA HGSOC cohort [20].

We identified a second mRNA co-expression network associated with chemotherapy

resistance in our HGSOC cohort and replicated in the AOCS cohort. The *darkolivegreen* network

included genes associated with fatty acid and cholesterol metabolism (*SREBF1*, *ACAA1*,

*ACADV*L), and the Protein Kinase B oncogene (*AKT1*), which promotes de-novo lipid

biosynthesis in cancer [35]. Sterol Regulatory Element-Binding Transcription Factor 1

(SREBF1) is a key enzyme for cholesterol and fatty acid synthesis, and an essential gene for

ovarian cancer tumor growth whose expression was correlated with the expression of lipogenic

genes in ovarian cancer cell lines [36]. *SREBF1* is activated by *AKT1*, promoting fatty acid

synthesis [37], which favors cell growth and proliferation in ovarian cancer [38]. Expression of

Acyl-CoA Dehydrogenase Very Long Chain (*ACADVL*), which is involved in the β-oxidation of

long-chain fatty acids, has been linked to ovarian cancer metastasis and cell survival in

suspension [39]. Increased expression of fatty acid synthase, a key enzyme in fatty acid

biosynthesis, was an independent prognostic factor for ovarian cancer recurrence [40], while its

inhibition sensitized resistant ovarian cancer cell lines to platinum [41]. Our findings indicate the

upregulation of these lipid metabolism genes among chemotherapy resistant patients in the

TCGA-HGSOC cohort. Finally, lipid metabolism dysregulation activates the UPR, and the UPR

triggers lipid metabolism-based adaptations in the cell through several pathways, including

SREBF1 regulation [42]. The interaction of these two systems may present a link between our

gene co-expression networks and warrants further study.

Moreover, our differential expression analysis identified a downregulation of mRNA

transcripts encoding for immunoglobulins, B and T lymphocyte factors, and interferon regulatory

factors in chemotherapy-resistant patients. The involvement of the immune response is well-

known in OC prognosis. Previous studies confirm that a high tumor immune score is a strong

predictor of chemosensitivity [43], and that a pre-existing inflammatory tumor

microenvironment is associated with increased PFS in HGSOC [44]. There are potential links

between this differential immune response activation and our UPR and lipid metabolism-related

gene co-expression networks. ER stress can induce pro-inflammatory cytokine production

alongside UPR activation in tumor cells [45], but may also be responsible for the disruption of

dendritic cell function in the ovarian cancer tumor microenvironment [46]. Finally, increased

lipid uptake and fatty acid oxidation inhibit tumor-infiltrating myeloid cell function in various

cancers [47-49].

The *ivory* miRNA co-expression network module is enriched for negative regulation of

lipoprotein transport, cholesterol efflux and sterol transport. This enrichment is mainly mediated

by miR-128-1 and miR-128-2, which play a key role in cholesterol and lipid homeostasis

through their suppression of the ABCA1 (ATP Binding Cassette Subfamily A Member 1)

cholesterol efflux transporter and the Low-Density Lipoprotein Receptor (LDLR) [50, 51]. MiR-

148a is also a regulator of these key genes [52] which is significantly downregulated in resistant

patients. The overexpression of *ABCA1* is associated with reduced survival in OC patients [53],

and levels of LDLR are increased in chemoresistant OC cell lines [54]. Finally, overexpression

of miR-128 resensitizes resistant OC cells to cisplatin by inhibiting ABCC5 (ATP Binding

Cassette Subfamily C Member 5) and BMI-1 (BMI1 ProtoOncogene, Polycomb Ring Finger)

[55]. Our results are consistent with the platinum resistance-inhibiting role of miR-128, and its

potential activity in cholesterol efflux inhibition alongside miR-148a in this cohort. Elevated

cholesterol levels directly contribute to chemotherapy resistance in OC ascites by upregulating

drug efflux pump proteins ABCG2 (ATP Binding Cassette Subfamily G Member 2) and ABCB1

(ATP Binding Cassette Subfamily B Member 1) [56], providing a potential mechanism that miR-

128 and miR-148a may be intercepting to provide their protective effect in our cohort.

The *plum* miRNA co-expression network consists of multiple isoforms of miR-221 and

miR-222, which have been implicated in the development of chemotherapy resistance in OC.

MiR-221/miR-222 expression is high in cisplatin-resistant OC cell lines, and its inhibition

sensitizes them [57]. Over-expression of miR-221 is associated with reduced disease-free and

overall survival, and promotes cell proliferation in OC cell lines through suppression of *APAF1*

(Apoptosis Protease Activating Factor 1) [58]. In addition, miR-222 promotes OC cell

proliferation by targeting the tumor suppressor $p27^{Kip1}$ (Cyclin-Dependent Kinase Inhibitor 1B)

[59]. Our findings are consistent with the increased activity of miR-221 and miR-222 in

chemotherapy-resistant HGSOC patient tumors, as previously demonstrated *in vitro*.

Correlation analysis of mRNA-seq and miRNA-seq datasets identified potential

interactions of *lavenderblush3* transcripts with miRNAs in the *plum* co-expression module, and

between *darkolivegreen* transcripts and miRNAs in the *ivory* co-expression module. The

negative correlation of the *plum* miRNA network with the UPR-associated *lavenderblush3*

mRNA network reveals a novel potential mechanism of chemotherapy resistance, as miR-221

and miR-222 may be responsible for the downregulation of this sensitivity-associated gene

network in platinum-resistant patients. This finding, combined with the accumulating evidence

of miR-221/miR-222 involvement in OC chemoresistance, may point to a promising avenue of

therapeutic intervention. However, overexpression of miR-221/miR-222 promotes UPR-induced

apoptosis in hepatocellular carcinoma (HCC) cells [60]. While the miR-222-mediated inhibition

of $p27^{Kip1}$ promotes cell proliferation in OC, it may also promote apoptosis under ER stress in

HCC. ER stress is also responsible for suppressing miR-221/miR-222 in HCC, rendering the

cells more resistant to apoptosis. The contribution of this mechanism to chemotherapy response

in HGSOC is currently unclear and presents an area for future investigation.

We also identified a negative correlation between the *darkolivegreen* mRNA network and

the *ivory* miRNA network. MiRNAs in the *ivory* network, which inhibits lipid transport, may

also inhibit lipid metabolism in chemotherapy-sensitive patients through targeting genes in the

resistance-associated *darkolivegreen* network. As increased lipid metabolism by cancer cells is a

known mechanism of chemotherapy resistance in HGSOC, our finding may present a novel

mechanism of chemotherapy sensitivity through miRNA-mediated inhibition of this lipid

metabolism-associated network.

Cis-eQTL analysis identified known and novel genomic variants associated with the

expression of mRNAs and miRNAs, which are associated with HDL and LDL cholesterol as

well as triglyceride levels. High HDL and triglyceride levels correlate with increased cancer

stage at diagnosis in epithelial OC patients [61]. In addition, advanced-stage epithelial OC

patients with high LDL levels have a significantly shorter PFS compared to patients with normal

LDL levels [62]. Further investigation of these eQTLs is necessary, as they may regulate the

expression of transcripts associated with chemotherapy response and may be useful as potential

predictors of platinum-based chemotherapy resistance and ovarian cancer prognosis.

Our study improves our understanding of the underlying mechanisms modulating

resistance to platinum-based chemotherapy, especially among HGSOC patients, and identifies

novel signalling pathways, networks, and regulators. Together, our data may serve as a basis to

identify personalized gene signatures to optimize clinical management of patients diagnosed with

HGSOC, as well as develop improved therapeutic strategies.

CRediT authorship contribution statement

Danai G. Topouza: Formal analysis, Software, Validation, Writing – Original Draft, Writing –

Review & Editing. **Jihoon Choi:** Formal analysis, Software, Writing – Review & Editing. **Sean**

Nesdoly: Software, Data Curation. **Anastasiya Tarnouskaya:** Data Curation. **Christopher J.B.**

Nicol: Writing – Review & Editing. **Qing Ling Duan:** Conceptualization, Supervision,

Methodology, Resources, Funding acquisition, Writing – Review & Editing.

Acknowledgements

Computations in this manuscript were performed on resources and with support provided by the

Centre for Advanced Computing (CAC) at Queen's University in Kingston, Ontario. The CAC is

funded by the Canada Foundation for Innovation, the Government of Ontario, and Queen's

University.

Funding sources

394 D.G.T. is funded by Queen's University and internal awards in the Department of Biomedical

395 and Molecular Sciences. Q.L.D. receives funding from the Canadian Institutes of Health

396 Research and Queen's University National Scholar Award. The funders did not play any role in

397 the study design, data collection, data analysis, data interpretation, writing of the manuscript or

398 decision to publish results.

Data statement

401 Transcriptomics, genomics, and clinical data used for the analysis of the TCGA-OV and TCGA-

402 BLCA cohorts can be accessed and downloaded from the Genomic Data Commons (GDC) Data

403 Portal (https://portal.gdc.cancer.gov/). Gene expression and clinical data of the AOCS cohort can

404 be accessed and downloaded from the Gene Expression Omnibus (GEO) database

405 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE9899).

Ethics declarations

Controlled access to datasets from The Cancer Genome Atlas (TCGA) was provided by the

National Institute of Health (NIH). The terms of access are outlined in the Data Use Certification

Agreement: https://dbgap.ncbi.nlm.nih.gov/aa/wga.cgi?view_pdf&wlid=10654&tlsid=274.

Details of the Human subject protection and data access policies implemented by TCGA are as

described: https://www.cancer.gov/about-nci/organization/ccg/research/structural-

genomics/tcga/history/policies/tcga-human-subjects-data-policies.pdf. Local ethics clearance for

human subject research was granted by Queen's University.

Table 1. Characteristics of the HGSOC patient cohorts with mRNA-Seq and miRNA-Seq data from TCGA

	mRNA-Seq cohort				miRNA-Seq cohort			
	Sensitive	Resistant	P value	All cases	Sensitive	Resistant	P value	All cases
Age								
Mean (range)	57.15 (30-87)	61.98 (38-87)	**0.0043** [a]	59.22 (30-87)	57.76 (30-87)	61.52 (38-87)	**0.018** [a]	59.26 (30-87)
Grade								
G2	18 (15.93 %)	8 (10.26 %)	0.27 [b]	26 (13.61 %)	23 (18.70 %)	8 (9.76 %)	0.14 [b]	31 (15.12 %)
G3/4	95 (84.07 %)	69 (88.46 %)		164 (85.86 %)	99 (80.49 %)	73 (89.02 %)		172 (83.90 %)
Ungraded	0 (0.00 %)	1 (1.28 %)		1 (0.52 %)	1 (0.81 %)	1 (1.22 %)		2 (0.98 %)
Stage								
II	7 (6.19 %)	2 (2.56 %)	0.48 [b]	9 (4.71 %)	8 (6.50 %)	2 (2.44 %)	0.41 [b]	10 (4.88 %)
III	91 (80.53 %)	67 (85.9 %)		158 (82.72 %)	99 (80.49 %)	67 (81.71 %)		166 (80.98 %)
IV	15 (13.27 %)	9 (11.54 %)		24 (12.57 %)	16 (13.01 %)	13 (15.85 %)		29 (14.15 %)
Cytoreductive surgery outcome								
Optimal (≤ 10mm)	73 (64.60 %)	54 (69.23 %)	0.23 [c]	127 (66.49 %)	77 (62.60 %)	56 (68.29 %)	0.21 [c]	133 (64.88 %)
Suboptimal (> 10mm)	26 (23.01 %)	20 (25.64 %)		46 (24.08 %)	29 (23.58 %)	21 (25.61 %)		50 (24.39 %)
Unknown	14 (12.39 %)	4 (5.13 %)		18 (9.42 %)	17 (13.82 %)	5 (6.10 %)		22 (10.73 %)
Overall survival								
Number of mortality events	48 (42.48 %)	60 (76.92 %)	**2.35E-06** [c]	108 (56.54 %)	55 (44.72 %)	66 (80.49 %)	**0.0002** [c]	121 (59.02 %)
Median months (95% CI)	52.04 (50.07-54.01)	28.2 (26.22-30.18)	**6.97E-10** [a]	39.87 (37.90-41.85)	51.91 (49.94-53.88)	28.2 (26.23-30.18)	**1.13E-10** [a]	39.5 (37.53-41.47)
Adjuvant chemotherapy regimen								
Platinum agent only	4 (3.54 %)	3 (3.85 %)	1 [b]	7 (3.66 %)	4 (3.25 %)	4 (4.88 %)	0.71 [b]	8 (3.90 %)
Platinum/Taxane combination	109 (96.46 %)	75 (96.15 %)		184 (96.34 %)	119 (96.75 %)	78 (95.12 %)		197 (96.10 %)

Primary tumor mRNA subtype

Differentiated	37 (32.74 %)	22 (28.21 %)	0.49 [b]	59 (30.89 %)	42 (34.15 %)	24 (29.27 %)	0.68 [c]	66 (32.20 %)
Immunoreactive	23 (20.35 %)	10 (12.82 %)		33 (17.28 %)	24 (19.51 %)	13 (15.85 %)		37 (18.05 %)
Mesenchymal	22 (19.47 %)	18 (23.08 %)		40 (20.94 %)	24 (19.51 %)	18 (21.95 %)		42 (20.49 %)
Proliferative	30 (26.55 %)	27 (34.62 %)		57 (29.84 %)	33 (26.83 %)	27 (32.93 %)		60 (29.27 %)
Subtype unknown	1 (0.88 %)	1 (1.28 %)		2 (1.05 %)	0 (0.00 %)	0 (0.00 %)		0 (0.00 %)
Total patients	113	78		191	123	82		205

[a] p-value based on t-test; [b] p-value based on Fisher's Exact test; [c] p-value based on Chi-Squared test

Table 2. Significantly enriched annotation terms for differentially expressed mRNA transcripts

Term name	Adjusted P value	Number of query genes in term
CD22 mediated BCR regulation	4.42E-45	30
Complement activation, classical pathway	3.07E-44	36
Antigen binding	1.74E-42	39
Scavenging of heme from plasma	1.03E-38	28
Immune response-regulating cell surface receptor signaling pathway involved in phagocytosis	3.89E-34	30
Fc-gamma receptor signaling pathway involved in phagocytosis	3.89E-34	30
Serine-type peptidase activity	7.94E-28	28
Cell surface interactions at the vascular wall	2.63E-27	30
Blood microparticle	2.57E-21	25
Leukocyte migration	3.36E-20	33
Immunoglobulin complex	2.24E-09	10
Cell periphery	3.36E-09	80
Plasma membrane	1.37E-08	78
Extracellular exosome	5.22E-07	43
Immunoglobulin receptor binding	5.98E-07	6
Phagocytosis, recognition	1.33E-06	6
External side of plasma membrane	1.48E-06	14
Regulation of B cell activation	8.13E-06	11
Phagocytosis, engulfment	8.91E-06	6
Ribosome	9.44E-05	6
Cytosolic large ribosomal subunit	2.91E-04	6
Ribosome, cytoplasmic	5.57E-04	6
60S ribosomal subunit, cytoplasmic	8.99E-04	5
Formation of a pool of free 40S subunits	9.17E-04	7
Peptide chain elongation	1.39E-03	6
Viral mRNA Translation	1.39E-03	6
Selenocysteine synthesis	1.79E-03	6
Eukaryotic Translation Termination	1.79E-03	6
Abnormal leukocyte morphology	2.02E-03	7
Nonsense Mediated Decay (NMD) independent of the Exon Junction Complex (EJC)	2.02E-03	6
SRP-dependent cotranslational protein targeting to membrane	3.09E-03	6
Nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	4.14E-03	7
Abnormal immunoglobulin level	4.44E-03	5
SRP-dependent cotranslational protein targeting to membrane	5.18E-03	6
Structural constituent of ribosome	6.70E-03	7

Factor: ZNF491; motif: NGKGGKCAGT	1.13E-02	7
Agammaglobulinemia	1.32E-02	2
Verrucae	2.39E-02	2
Cellulitis	4.10E-02	2
TRBP containing complex (DICER, RPL7A, EIF6, MOV10 and subunits of the 60S ribosomal particle)	4.33E-02	4
Nop56p-associated pre-rRNA complex	4.57E-02	5

Table 3. Significantly enriched annotation terms for differentially expressed miRNA isoforms

Term name	Adjusted P value	Number of query miRNAs in term
mRNA binding involved in posttranscriptional gene silencing	4.98E-16	10
Gene silencing by miRNA	7.12E-14	10
MicroRNAs in cancer	1.09E-12	10
Negative regulation of cardiac muscle cell proliferation	5.21E-08	4
Positive regulation of cardiac muscle tissue regeneration	2.10E-07	3
Re-entry into mitotic cell cycle	5.25E-07	3
Negative regulation of cell growth involved in cardiac muscle cell development	8.65E-06	3
Positive regulation of lung blood pressure	2.05E-04	2
Negative regulation of cardiac muscle cell apoptotic process	3.12E-04	3
Regulation of vasculature development	3.32E-03	5
Negative regulation of autophagy	8.70E-03	3
Extracellular space	1.03E-02	9
Positive regulation of connective tissue replacement	2.14E-02	2
Negative regulation of nitric oxide metabolic process	2.45E-02	2
Blood vessel morphogenesis	3.25E-02	5
Positive regulation of protein kinase B signaling	3.97E-02	3

Table 4. Significantly enriched terms in significant mRNA transcript co-expression networks

Network name	Term name	P value	Number of query genes in term
lavenderblush3	Protein modification by small protein conjugation or removal	2.40E-02[a]	9
	Protein modification by small protein conjugation	2.71E-03[a]	9
	Protein ubiquitination	1.60E-03[a]	9
	Factor: GABP-alpha; motif: CTTCCK	6.72E-03[a]	26
darkolivegreen	Protein-containing complex	4.60E-02[a]	37
	Biosynthesis of unsaturated fatty acids	5.20E-03[b]	2
	Fatty acid degradation	1.34E-02[b]	2
	Notch signaling pathway	1.58E-02[b]	2
	Non-alcoholic fatty liver disease (NAFLD)	2.19E-02[b]	3
	Ribosome	2.35E-02[b]	3
	RNA transport	2.85E-02[b]	3
	Synthesis and degradation of ketone bodies	3.93E-02[b]	1
	Peroxisome	4.37E-02[b]	2

[a] Benjamini-Hochberg adjusted p value; [b] non-adjusted p value

Table 5. Significantly enriched terms in significant miRNA isoform co-expression networks

Network name	Term name	Adjusted P value	Number of query miRNAs in term
ivory	mRNA binding involved in posttranscriptional gene silencing	1.16E-09	7
	Gene silencing by miRNA	3.61E-08	7
	Extracellular space	4.24E-05	10
	MicroRNAs in cancer	9.55E-05	5
	Regulation of lipoprotein transport	1.22E-03	2
	Negative regulation of cholesterol efflux	2.69E-02	2
	Negative regulation of lipoprotein particle clearance	4.27E-02	2
plum	Positive regulation of Schwann cell proliferation involved in axon regeneration	2.55E-06	2
	Negative regulation of hematopoietic stem cell proliferation	2.55E-06	2
	Positive regulation of Schwann cell migration	7.66E-06	2
	Negative regulation of TRAIL-activated apoptotic signaling pathway	7.66E-06	2
	Negative regulation of cell adhesion molecule production	1.53E-05	2
	Negative regulation by host of viral genome replication	5.36E-05	2
	Negative regulation of leukocyte adhesion to vascular endothelial cell	1.40E-04	2
	Negative regulation of cytokine production involved in inflammatory response	2.32E-04	2
	Positive regulation of erythrocyte differentiation	1.19E-03	2
	Positive regulation of G1/S transition of mitotic cell cycle	1.61E-03	2
	Positive regulation of vascular smooth muscle cell proliferation	2.53E-03	2
	Positive regulation of epithelial to mesenchymal transition	3.00E-03	2
	Negative regulation of muscle cell differentiation	5.15E-03	2
	Negative regulation of translation, ncRNA-mediated	6.17E-03	2
	Negative regulation of sprouting angiogenesis	8.07E-03	2
	Positive regulation of epithelial cell migration	3.33E-02	2
	Positive regulation of protein kinase B signaling	3.58E-02	2
	mRNA binding involved in posttranscriptional gene silencing	4.25E-02	2

Table 6. Predicted and validated mRNA - miRNA network interactions

Integrated Networks	Gene ID	mRNA ID	miRNA ID	Spearman's ρ	P value	Interaction type	Source
lavenderblush3 - plum	PSIP1	ENST00000397519	hsa-mir-222-3p	-0.32	0.000021	Experimental validation	Xue et al. (2013) [1]
	CDC37L1	ENST00000381854	hsa-mir-222-3p	-0.31	0.000042	Computational prediction	Miranda|7mer-m8|889|909|1.11303779449042|;
	CDC37L1	ENST00000381854	hsa-mir-221-3p	-0.27	0.00034	Computational prediction	Miranda|7mer-m8|887|909|1.0718982884376|;
	CDC37L1	ENST00000381854	hsa-mir-221-5p	-0.27	0.00034	Computational prediction	Miranda|8mer|1115|1134|1.3187353247545|;
	CDC37L1	ENST00000381854	hsa-mir-221-5p	-0.27	0.00034	Computational prediction	Miranda|N/A|145|167|1.0718982884376|;
	NFIB	ENST00000380959	hsa-mir-221-3p	-0.27	0.00039	Experimental validation	Grosswendt et al. (2014) [2]
	SMU1	ENST00000397149	hsa-mir-221-5p	-0.26	0.00082	Experimental validation	Kameswaran et al. (2014) [3]
	VCP	ENST00000358901	hsa-mir-222-3p	-0.25	0.00089	Experimental validation	Helwak et al. (2013) [4]
	CAAP1	ENST00000333916	hsa-mir-222-5p	-0.22	0.0041	Computational prediction	Miranda|7mer-m8|1097|1123|1.20902997528032|;
	CAAP1	ENST00000333916	hsa-mir-221-5p	-0.21	0.0076	Computational prediction	Miranda|N/A|694|718|1.14046413185896|;
	TOPORS	ENST00000360538	hsa-mir-221-3p	-0.19	0.014	Experimental validation	Balakrishnan et al. (2014) [5]
	SLC25A51	ENST00000496760	hsa-mir-222-3p	-0.19	0.016	Experimental validation	Helwak et al. (2013) [4]
	PPAPDC2	ENST00000381883	hsa-mir-222-5p	-0.18	0.021	Computational prediction	Microtar|8mer|173|178|-0.863350042828295|;Miranda|7mer-1A|155|179|1.0718982884376|;
	PPAPDC2	ENST00000381883	hsa-mir-222-5p	-0.18	0.021	Computational prediction	Microtar|8mer|1836|1841|-0.87475584746292|;
	VCP	ENST00000358901	hsa-mir-221-3p	-0.17	0.023	Experimental validation	Lipchina et al. (2011) [6]
	VCP	ENST00000358901	hsa-mir-221-5p	-0.17	0.023	Experimental validation	Balakrishnan et al. (2014) [5]
	NFX1	ENST00000379540	hsa-mir-221-5p	-0.17	0.027	Computational prediction	Microtar|7mer-m8|763|768|-0.872619157510717|;Miranda|7mer-m8|748|769|1.09932462580615|;
	PPAPDC2	ENST00000381883	hsa-mir-221-5p	-0.17	0.03	Computational prediction	Miranda|Offset 3-8 6mer|916|937|1.12675096317469|;
	PPAPDC2	ENST00000381883	hsa-mir-221-5p	-0.17	0.03	Computational prediction	Miranda|N/A|132|153|1.05818511975333|;Microtar|7mer-m8|147|152|-0.870278223624131|;
	DNAJA1	ENST00000330899	hsa-mir-221-5p	-0.16	0.035	Experimental validation	Balakrishnan et al. (2014) [5]

	Gene	Transcript	miRNA	Corr	p-value	Evidence	Source
	AK3	ENST00000381809	hsa-mir-222-5p	-0.16	0.037	Computational prediction	Miranda|7mer-m8|111|131|1.11303779449042|;
darkolivegreen - ivory	*RPL18*	ENST00000552347	hsa-mir-212-3p	-0.29	0.00017	Experimental Validation	Karginov et al. (2013) [7]
	MRPL55	ENST00000411464	hsa-miR-1306-5p	-0.25	0.001	Computational prediction	Rnahybrid|Offset 3-9 7mer-m8|35|41|-0.874295427824345|;
	FASTK	ENST00000466855	hsa-mir-140-3p	-0.25	0.0013	Experimental Validation	Kishore et al. (2011) [8]
	RPL18	ENST00000552347	hsa-mir-361-3p	-0.24	0.0021	Experimental Validation	Karginov et al. (2013) [7]
	ACBD4	ENST00000321854	hsa-miR-128-2-5p	-0.22	0.0041	Computational prediction	Miranda|7mer-m8|501|523|1.08561145712188|;Rnahybrid|Offset 1-7 7mer-m8|515|522|-0.873597482390991|;
	OAZ1	ENST00000581150	hsa-mir-361-3p	-0.22	0.0047	Experimental Validation	Karginov et al. (2013); Kameswaran et al. (2014) [3, 7]
	ACBD4	ENST00000321854	hsa-miR-128-1-5p	-0.21	0.0059	Computational prediction	Miranda|7mer-m8|500|523|1.0718982884376|;Rnahybrid|Offset 1-7 7mer-m8|515|522|-0.872932832531201|;
	MYL12A	ENST00000578611	hsa-mir-103a-3p	-0.17	0.023	Experimental Validation	Balakrishnan et al. (2014); Karginov et al. (2013) [5, 7]
	MYL12A	ENST00000578611	hsa-mir-107	-0.17	0.023	Experimental Validation	Balakrishnan et al. (2014); Karginov et al. (2013); Nelson et al. (2011) [5, 7, 9]
	HTRA2	ENST00000484352	hsa-mir-1306-5p	-0.17	0.029	Experimental Validation	Karginov et al. (2013) [7]
	ACADVL	ENST00000579425	hsa-mir-103a-3p	-0.17	0.03	Experimental Validation	Balakrishnan et al. (2014) [5]
	ACADVL	ENST00000579425	hsa-mir-107	-0.17	0.03	Experimental Validation	Balakrishnan et al. (2014) [5]
	SOD1	ENST00000470944	hsa-mir-140-3p	-0.16	0.037	Experimental Validation	Xue et al. (2013) [1]
	RHOT2	ENST00000569675	hsa-mir-1306-5p	-0.16	0.044	Experimental Validation	Karginov et al. (2013) [7]
	TSSK6	ENST00000360913	hsa-miR-212-5p	-0.15	0.048	Computational prediction	Microtar|7mer-m8|674|679|-0.874694700443757|;Miranda|N/A|658|680|1.05818511975333|;
	RCC1	ENST00000373832	hsa-miR-1306-3p	-0.15	0.048	Computational prediction	Miranda|7mer-m8|675|693|1.30502215607023|;
	RCC1	ENST00000373832	hsa-miR-1306-5p	-0.15	0.048	Computational prediction	Miranda|7mer-m8|785|805|1.11303779449042|;

[1] Y. Xue, K. Ouyang, J. Huang, Y. Zhou, H. Ouyang, H. Li, G. Wang, Q. Wu, C. Wei, Y. Bi, L. Jiang, Z. Cai, H. Sun, K. Zhang, Y. Zhang, J. Chen, X.D. Fu, Direct conversion of fibroblasts to neurons by reprogramming PTB-regulated microRNA circuits, Cell, 152 (2013) 82-96.
[2] S. Grosswendt, A. Filipchyk, M. Manzano, F. Klironomos, M. Schilling, M. Herzog, E. Gottwein, N. Rajewsky, Unambiguous identification of miRNA:target site interactions by different types of ligation reactions, Mol Cell, 54 (2014) 1042-1054.
[3] V. Kameswaran, N.C. Bramswig, L.B. McKenna, M. Penn, J. Schug, N.J. Hand, Y. Chen, I. Choi, A. Vourekas, K.J. Won, C. Liu, K. Vivek, A. Naji, J.R. Friedman, K.H. Kaestner, Epigenetic regulation of the DLK1-MEG3 microRNA cluster in human type 2 diabetic islets, Cell Metab, 19 (2014) 135-145.

[4] A. Helwak, G. Kudla, T. Dudnakova, D. Tollervey, Mapping the human miRNA interactome by CLASH reveals frequent noncanonical binding, Cell, 153 (2013) 654-665.

[5] I. Balakrishnan, X. Yang, J. Brown, A. Ramakrishnan, B. Torok-Storb, P. Kabos, J.R. Hesselberth, M.M. Pillai, Genome-wide analysis of miRNA-mRNA interactions in marrow stromal cells, Stem Cells, 32 (2014) 662-673.

[6] I. Lipchina, Y. Elkabetz, M. Hafner, R. Sheridan, A. Mihailovic, T. Tuschl, C. Sander, L. Studer, D. Betel, Genome-wide identification of microRNA targets in human ES cells reveals a role for miR-302 in modulating BMP response, Genes Dev, 25 (2011) 2173-2186.

[7] F.V. Karginov, G.J. Hannon, Remodeling of Ago2-mRNA interactions upon cellular stress reflects miRNA complementarity and correlates with altered translation rates, Genes Dev, 27 (2013) 1624-1632.

[8] S. Kishore, L. Jaskiewicz, L. Burger, J. Hausser, M. Khorshid, M. Zavolan, A quantitative analysis of CLIP methods for identifying binding sites of RNA-binding proteins, Nat Methods, 8 (2011) 559-564.

[9] P.T. Nelson, W.X. Wang, G. Mao, B.R. Wilfred, K. Xie, M.H. Jennings, Z. Gao, X. Wang, Specific sequence determinants of miR-15/107 microRNA gene group targets, Nucleic Acids Res, 39 (2011) 8163-8172.

 Figure Legends

Figure 1. Differential expression analysis of mRNA and miRNA transcripts. (A) Significant

differentially expressed mRNA transcripts (n=196) are shown in red. The Benjamini-Hochberg

adjusted significance threshold is indicated by the dashed line. Transcripts with a positive fold

change are upregulated in resistant patients, whereas transcripts with a negative fold change are

downregulated in resistant patients. (B) Significant differentially expressed miRNA isoforms

(n=21) are shown in red. The Benjamini-Hochberg adjusted significance threshold is indicated

by the dashed line. Transcripts with a positive fold change are upregulated in resistant patients,

whereas transcripts with a negative fold change are downregulated in resistant patients. These

figures were generated using the R package ggplot2 (v. 3.3.2) [63].

Figure 2. Weighted transcript co-expression network analysis of mRNA and miRNA

transcripts. (A) Heatmap of the correlation between the 58 mRNA co-expression modules with

platinum-based chemotherapy resistance, red indicating a positive effect and blue a negative

effect. P-values are indicated for each module. Two mRNA co-expression modules are

significantly associated with chemotherapy resistance (*lavenderblush3*, *darkolivegreen*). (B) The

mRNA modules *lavenderblush3* and *darkolivegreen* are visualized in their respective colors.

Each node represents a transcript, and each edge represents a connection or co-expression. The

distance between nodes is the connection weight, where more similar transcripts are plotted

closer. (C) Heatmap of the correlation between the 100 miRNA modules with platinum-based

chemotherapy resistance, red indicating a positive effect and blue a negative effect. P-values are

indicated for each module. Three miRNA co-expression modules are significantly associated

with chemotherapy resistance (*ivory*, *lightcoral*, and *plum*). (D) The miRNA modules *lightcoral*,

plum, and *ivory* are visualized in their respective colors. Each node represents one miRNA

isoform, and each edge represents a connection or co-expression. The distance between nodes is

the connection weight, where more similar transcripts are plotted closer.

Figure 3. Integration of mRNA - miRNA networks. The mRNA transcripts (oval nodes) from

the *lavenderblush3* and *darkolivegreen* modules are arranged based on co-expression similarity

(grey edges). Four common miRNA isoforms (rectangular nodes) were validated (orange solid

edges) or predicted (orange dashed edges) to regulate the expression of genes in the

lavenderblush3 network module, while 10 miRNA isoforms were validated or predicted to

regulate genes in the *darkolivegreen* module. Genes targets of miRNAs indicated by orange

border.

Figure 4. Replication analysis of the gene co-expression networks. (A-B) Prognostic Index

Kaplan-Meier analysis of patient PFS association with the prognostic index of mRNA networks

lavenderblush3 (A) and *darkolivegreen* (B). (C-E) Prognostic Index Kaplan-Meier analysis of

patient OS association with the prognostic index of miRNA networks *ivory* (C), *plum* (D), and

lightcoral (E). (F-G) Kaplan-Meier analysis of patient OS with the expression of miR-221 (F)

and miR-222 (G). Figures 4A-E were generated using the R package survminer (v. 0.4.8) [64].

Figures 4F,G were generated by the KM Plotter tool. PFS, progression-free survival; OS, overall

survival.

References

[1] R.J. Kurman, International Agency for Research on Cancer., World Health Organization., WHO classification of tumours of female reproductive organs, 4th ed., International Agency for Research on Cancer, Lyon, 2014.

[2] S. Bouberhan, L. Philp, S. Hill, L.F. Al-Alem, B. Rueda, Exploiting the Prevalence of Homologous Recombination Deficiencies in High-Grade Serous Ovarian Cancer, Cancers (Basel), 12 (2020).

[3] A. Davis, A.V. Tinker, M. Friedlander, "Platinum resistant" ovarian cancer: what is it, who to treat and how to measure benefit?, Gynecol Oncol, 133 (2014) 624-631.

[4] A.E. Freimund, J.A. Beach, E.L. Christie, D.D.L. Bowtell, Mechanisms of Drug Resistance in High-Grade Serous Ovarian Cancer, Hematol Oncol Clin North Am, 32 (2018) 983-996.

[5] M. Binju, M.A. Padilla, T. Singomat, P. Kaur, Y. Suryo Rahmanto, P.A. Cohen, Y. Yu, Mechanisms underlying acquired platinum resistance in high grade serous ovarian cancer - a mini review, Biochim Biophys Acta Gen Subj, 1863 (2019) 371-378.

[6] K.L. Lloyd, I.A. Cree, R.S. Savage, Prediction of resistance to chemotherapy in ovarian cancer: a systematic review, BMC Cancer, 15 (2015) 117.

[7] S. Marchini, R. Fruscio, L. Clivio, L. Beltrame, L. Porcu, I. Fuso Nerini, D. Cavalieri, G. Chiorino, G. Cattoretti, C. Mangioni, R. Milani, V. Torri, C. Romualdi, A. Zambelli, M. Romano, M. Signorelli, S. di Giandomenico, M. D'Incalci, Resistance to platinum-based chemotherapy is associated with epithelial to mesenchymal transition in epithelial ovarian cancer, Eur J Cancer, 49 (2013) 520-530.

[8] M. Koti, R.J. Gooding, P. Nuin, A. Haslehurst, C. Crane, J. Weberpals, T. Childs, P. Bryson, M. Dharsee, K. Evans, H.E. Feilotter, P.C. Park, J.A. Squire, Identification of the IGF1/PI3K/NF kappaB/ERK gene signalling networks associated with chemotherapy resistance and treatment response in high-grade serous epithelial ovarian cancer, BMC Cancer, 13 (2013) 549.

[9] M. Zhang, S.C. Luo, Gene expression profiling of epithelial ovarian cancer reveals key genes and pathways associated with chemotherapy resistance, Genet Mol Res, 15 (2016).

[10] Y. Liu, Z. Zhang, T. Li, X. Li, S. Zhang, Y. Li, W. Zhao, Y. Gu, Z. Guo, L. Qi, A Qualitative Transcriptional Signature for Predicting Recurrence Risk for High-Grade Serous Ovarian Cancer Patients Treated With Platinum-Taxane Adjuvant Chemotherapy, Front Oncol, 9 (2019) 1094.

[11] G. Tapia, I. Diaz-Padilla, Molecular Mechanisms of Platinum Resistance in Ovarian Cancer, Ovarian Cancer - A Clinical and Translational Update, IntechOpen2013.

[12] M. Oti, H.G. Brunner, The modular nature of genetic diseases, Clin Genet, 71 (2007) 1-11.

[13] N. Cancer Genome Atlas Research, Integrated genomic analyses of ovarian carcinoma, Nature, 474 (2011) 609-615.

[14] N. Gehlenborg, M.S. Noble, G. Getz, L. Chin, P.J. Park, Nozzle: a report generation toolkit for data analysis pipelines, Bioinformatics, 29 (2013) 1089-1091.

[15] M.I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2, Genome Biol, 15 (2014) 550.

[16] P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation network
analysis, BMC Bioinformatics, 9 (2008) 559.
[17] P. Shannon, A. Markiel, O. Ozier, N.S. Baliga, J.T. Wang, D. Ramage, N. Amin, B.
Schwikowski, T. Ideker, Cytoscape: a software environment for integrated models of
biomolecular interaction networks, Genome Res, 13 (2003) 2498-2504.
[18] U. Raudvere, L. Kolberg, I. Kuzmin, T. Arak, P. Adler, H. Peterson, J. Vilo, g:Profiler: a
web server for functional enrichment analysis and conversions of gene lists (2019 update),
Nucleic Acids Res, 47 (2019) W191-W198.
[19] M.V. Kuleshov, M.R. Jones, A.D. Rouillard, N.F. Fernandez, Q. Duan, Z. Wang, S. Koplev,
S.L. Jenkins, K.M. Jagodnik, A. Lachmann, M.G. McDermott, C.D. Monteiro, G.W. Gundersen,
A. Ma'ayan, Enrichr: a comprehensive gene set enrichment analysis web server 2016 update,
Nucleic Acids Res, 44 (2016) W90-97.
[20] J. Choi, D.G. Topouza, A. Tarnouskaya, S. Nesdoly, M. Koti, Q.L. Duan, Gene networks
and expression quantitative trait loci associated with adjuvant chemotherapy response in high-
grade serous ovarian cancer, BMC Cancer, 20 (2020) 413.
[21] A.A. Shabalin, Matrix eQTL: ultra fast eQTL analysis via large matrix operations,
Bioinformatics, 28 (2012) 1353-1358.
[22] B.E. Stranger, S.B. Montgomery, A.S. Dimas, L. Parts, O. Stegle, C.E. Ingle, M. Sekowska,
G.D. Smith, D. Evans, M. Gutierrez-Arcelus, A. Price, T. Raj, J. Nisbett, A.C. Nica, C. Beazley,
R. Durbin, P. Deloukas, E.T. Dermitzakis, Patterns of cis regulatory variation in diverse human
populations, PLoS Genet, 8 (2012) e1002639.
[23] L.D. Ward, M. Kellis, HaploReg v4: systematic mining of putative causal variants, cell
types, regulators and target genes for human complex traits and disease, Nucleic Acids Res, 44
(2016) D877-881.
[24] Y. Fan, J. Xia, miRNet-Functional Analysis and Visual Exploration of miRNA-Target
Interactions in a Network Context, Methods Mol Biol, 1819 (2018) 215-233.
[25] L. Chang, G. Zhou, O. Soufan, J. Xia, miRNet 2.0: network-based visual analytics for
miRNA functional analysis and systems biology, Nucleic Acids Res, 48 (2020) W244-W251.
[26] E. Andres-Leon, G. Gomez-Lopez, D.G. Pisano, Prediction of miRNA-mRNA Interactions
Using miRGate, Methods Mol Biol, 1580 (2017) 225-237.
[27] B. Gyorffy, A. Lanczky, Z. Szallasi, Implementing an online tool for genome-wide
validation of survival-associated biomarkers in ovarian-cancer using microarray data from 1287
patients, Endocr Relat Cancer, 19 (2012) 197-208.
[28] R.W. Tothill, A.V. Tinker, J. George, R. Brown, S.B. Fox, S. Lade, D.S. Johnson, M.K.
Trivett, D. Etemadmoghadam, B. Locandro, N. Traficante, S. Fereday, J.A. Hung, Y.E. Chiew, I.
Haviv, G. Australian Ovarian Cancer Study, D. Gertig, A. DeFazio, D.D. Bowtell, Novel
molecular subtypes of serous and endometrioid ovarian cancer linked to clinical outcome, Clin
Cancer Res, 14 (2008) 5198-5208.
[29] F. Petrelli, A. Coinu, M. Cabiddu, M. Ghilardi, I. Vavassori, S. Barni, Correlation of
pathologic complete response with survival after neoadjuvant chemotherapy in bladder cancer
treated with cystectomy: a meta-analysis, Eur Urol, 65 (2014) 350-357.
[30] R. Aguirre-Gamboa, H. Gomez-Rueda, E. Martinez-Ledesma, A. Martinez-Torteya, R.
Chacolla-Huaringa, A. Rodriguez-Barrientos, J.G. Tamez-Pena, V. Trevino, SurvExpress: an
online biomarker validation tool and database for cancer gene expression data using survival
analysis, PLoS One, 8 (2013) e74250.

[31] H. Kim, A. Bhattacharya, L. Qi, Endoplasmic reticulum quality control in cancer: Friend or
foe, Semin Cancer Biol, 33 (2015) 25-33.
[32] E. Bahar, J.Y. Kim, H. Yoon, Chemotherapy Resistance Explained through Endoplasmic
Reticulum Stress-Dependent Signaling, Cancers (Basel), 11 (2019).
[33] J. Tian, R. Liu, Q. Qu, Role of endoplasmic reticulum stress on cisplatin resistance in
ovarian carcinoma, Oncol Lett, 13 (2017) 1437-1443.
[34] P. Bastola, L. Neums, F.J. Schoenen, J. Chien, VCP inhibitors induce endoplasmic
reticulum stress, cause cell cycle arrest, trigger caspase-mediated cell death and synergistically
kill ovarian cancer cells in combination with Salubrinal, Mol Oncol, 10 (2016) 1559-1574.
[35] N. Koundouros, G. Poulogiannis, Reprogramming of fatty acid metabolism in cancer, Br J
Cancer, 122 (2020) 4-22.
[36] L.Y. Nie, Q.T. Lu, W.H. Li, N. Yang, S. Dongol, X. Zhang, J. Jiang, Sterol regulatory
element-binding protein 1 is required for ovarian tumor growth, Oncol Rep, 30 (2013) 1346-
1354.
[37] T. Porstmann, B. Griffiths, Y.L. Chung, O. Delpuech, J.R. Griffiths, J. Downward, A.
Schulze, PKB/Akt induces transcription of enzymes involved in cholesterol and fatty acid
biosynthesis via activation of SREBP, Oncogene, 24 (2005) 6465-6481.
[38] H.Q. Wang, D.A. Altomare, K.L. Skele, P.I. Poulikakos, F.P. Kuhajda, A. Di Cristofano,
J.R. Testa, Positive feedback regulation between AKT activation and fatty acid synthase
expression in ovarian carcinoma cells, Oncogene, 24 (2005) 3574-3582.
[39] L.J. Wheeler, Z.L. Watson, L. Qamar, T.M. Yamamoto, B.T. Sawyer, K.D. Sullivan, S.
Khanal, M. Joshi, V. Ferchaud-Roucher, H. Smith, L.A. Vanderlinden, S.W. Brubaker, C.M.
Caino, H. Kim, J.M. Espinosa, J.K. Richer, B.G. Bitler, Multi-Omic Approaches Identify
Metabolic and Autophagy Regulators Important in Ovarian Cancer Dissemination, iScience, 19
(2019) 474-491.
[40] P.L. Alo, P. Visca, M.L. Framarino, C. Botti, S. Monaco, V. Sebastiani, D.E. Serpieri, U. Di
Tondo, Immunohistochemical study of fatty acid synthase in ovarian neoplasms, Oncol Rep, 7
(2000) 1383-1388.
[41] E. Papaevangelou, G.S. Almeida, C. Box, N.M. deSouza, Y.L. Chung, The effect of FASN
inhibition on the growth and metabolism of a cisplatin-resistant ovarian carcinoma model, Int J
Cancer, 143 (2018) 992-1002.
[42] B.A.S. Pinto, L.M. Franca, F.R.M. Laurindo, A.M.A. Paes, Unfolded Protein Response:
Cause or Consequence of Lipid and Lipoprotein Metabolism Disturbances?, Adv Exp Med Biol,
1127 (2019) 67-82.
[43] D. Hao, J. Liu, M. Chen, J. Li, L. Wang, X. Li, Q. Zhao, L.J. Di, Immunogenomic Analyses
of Advanced Serous Ovarian Cancer Reveal Immune Score is a Strong Prognostic Factor and an
Indicator of Chemosensitivity, Clin Cancer Res, 24 (2018) 3560-3571.
[44] M. Koti, A. Siu, I. Clément, M. Bidarimath, G. Turashvili, A. Edwards, K. Rahimi, A.M.
Mes-Masson, J.A. Squire, A distinct pre-existing inflammatory tumour microenvironment is
associated with chemotherapy resistance in high-grade serous epithelial ovarian cancer, Br J
Cancer, 112 (2015) 1215-1222.
[45] N.R. Mahadevan, J. Rodvold, H. Sepulveda, S. Rossi, A.F. Drew, M. Zanetti, Transmission
of endoplasmic reticulum stress and pro-inflammation from tumor cells to myeloid cells, Proc
Natl Acad Sci U S A, 108 (2011) 6561-6566.
[46] J.R. Cubillos-Ruiz, P.C. Silberman, M.R. Rutkowski, S. Chopra, A. Perales-Puchalt, M.
Song, S. Zhang, S.E. Bettigole, D. Gupta, K. Holcomb, L.H. Ellenson, T. Caputo, A.H. Lee, J.R.

Conejo-Garcia, L.H. Glimcher, ER Stress Sensor XBP1 Controls Anti-tumor Immunity by Disrupting Dendritic Cell Homeostasis, Cell, 161 (2015) 1527-1538.

[47] D.L. Herber, W. Cao, Y. Nefedova, S.V. Novitskiy, S. Nagaraj, V.A. Tyurin, A. Corzo, H.I. Cho, E. Celis, B. Lennox, S.C. Knight, T. Padhya, T.V. McCaffrey, J.C. McCaffrey, S. Antonia, M. Fishman, R.L. Ferris, V.E. Kagan, D.I. Gabrilovich, Lipid accumulation and dendritic cell dysfunction in cancer, Nat Med, 16 (2010) 880-886.

[48] F. Hossain, A.A. Al-Khami, D. Wyczechowska, C. Hernandez, L. Zheng, K. Reiss, L.D. Valle, J. Trillo-Tinoco, T. Maj, W. Zou, P.C. Rodriguez, A.C. Ochoa, Inhibition of Fatty Acid Oxidation Modulates Immunosuppressive Functions of Myeloid-Derived Suppressor Cells and Enhances Cancer Therapies, Cancer Immunol Res, 3 (2015) 1236-1247.

[49] R. Ramakrishnan, V.A. Tyurin, F. Veglia, T. Condamine, A. Amoscato, D. Mohammadyani, J.J. Johnson, L.M. Zhang, J. Klein-Seetharaman, E. Celis, V.E. Kagan, D.I. Gabrilovich, Oxidized lipids block antigen cross-presentation by dendritic cells in cancer, J Immunol, 192 (2014) 2920-2931.

[50] A. Wagschal, S.H. Najafi-Shoushtari, L. Wang, L. Goedeke, S. Sinha, A.S. deLemos, J.C. Black, C.M. Ramirez, Y. Li, R. Tewhey, I. Hatoum, N. Shah, Y. Lu, F. Kristo, N. Psychogios, V. Vrbanac, Y.C. Lu, T. Hla, R. de Cabo, J.S. Tsang, E. Schadt, P.C. Sabeti, S. Kathiresan, D.E. Cohen, J. Whetstine, R.T. Chung, C. Fernandez-Hernando, L.M. Kaplan, A. Bernards, R.E. Gerszten, A.M. Naar, Genome-wide identification of microRNAs regulating cholesterol and triglyceride homeostasis, Nat Med, 21 (2015) 1290-1297.

[51] Y.K. Adlakha, S. Khanna, R. Singh, V.P. Singh, A. Agrawal, N. Saini, Pro-apoptotic miRNA-128-2 modulates ABCA1, ABCG1 and RXRalpha expression and cholesterol homeostasis, Cell Death Dis, 4 (2013) e780.

[52] L. Goedeke, N. Rotllan, A. Canfran-Duque, J.F. Aranda, C.M. Ramirez, E. Araldi, C.S. Lin, N.N. Anderson, A. Wagschal, R. de Cabo, J.D. Horton, M.A. Lasuncion, A.M. Naar, Y. Suarez, C. Fernandez-Hernando, MicroRNA-148a regulates LDL receptor and ABCA1 expression to control circulating lipoprotein levels, Nat Med, 21 (2015) 1280-1289.

[53] E.L. Hedditch, B. Gao, A.J. Russell, Y. Lu, C. Emmanuel, J. Beesley, S.E. Johnatty, X. Chen, P. Harnett, J. George, G. Australian Ovarian Cancer Study, R.T. Williams, C. Flemming, D. Lambrechts, E. Despierre, S. Lambrechts, I. Vergote, B. Karlan, J. Lester, S. Orsulic, C. Walsh, P. Fasching, M.W. Beckmann, A.B. Ekici, A. Hein, K. Matsuo, S. Hosono, T. Nakanishi, Y. Yatabe, T. Pejovic, Y. Bean, F. Heitz, P. Harter, A. du Bois, I. Schwaab, E. Hogdall, S.K. Kjaer, A. Jensen, C. Hogdall, L. Lundvall, S.A. Engelholm, B. Brown, J. Flanagan, M.D. Metcalf, N. Siddiqui, T. Sellers, B. Fridley, J. Cunningham, J. Schildkraut, E. Iversen, R.P. Weber, A. Berchuck, E. Goode, D.D. Bowtell, G. Chenevix-Trench, A. deFazio, M.D. Norris, S. MacGregor, M. Haber, M.J. Henderson, ABCA transporter gene expression and poor outcome in epithelial ovarian cancer, J Natl Cancer Inst, 106 (2014).

[54] L. Zheng, L. Li, Y. Lu, F. Jiang, X.A. Yang, SREBP2 contributes to cisplatin resistance in ovarian cancer cells, Exp Biol Med (Maywood), 243 (2018) 655-662.

[55] B. Li, H. Chen, N. Wu, W.J. Zhang, L.X. Shang, Deregulation of miR-128 in ovarian cancer promotes cisplatin resistance, Int J Gynecol Cancer, 24 (2014) 1381-1388.

[56] S. Kim, M. Lee, D.N. Dhanasekaran, Y.S. Song, Activation of LXRa/beta by cholesterol in malignant ascites promotes chemoresistance in ovarian cancer, BMC Cancer, 18 (2018) 1232.

[57] Z. Amini-Farsani, M.H. Sangtarash, M. Shamsara, H. Teimori, MiR-221/222 promote chemoresistance to cisplatin in ovarian cancer cells by targeting PTEN/PI3K/AKT signaling pathway, Cytotechnology, 70 (2018) 203-213.

[58] J. Li, Q. Li, H. Huang, Y. Li, L. Li, W. Hou, Z. You, Overexpression of miRNA-221
promotes cell proliferation by targeting the apoptotic protease activating factor-1 and indicates a
poor prognosis in ovarian cancer, Int J Oncol, (2017).
[59] C. Sun, N. Li, B. Zhou, Z. Yang, D. Ding, D. Weng, L. Meng, S. Wang, J. Zhou, D. Ma, G.
Chen, miR-222 is upregulated in epithelial ovarian cancer and promotes cell proliferation by
downregulating P27(kip1.), Oncol Lett, 6 (2013) 507-512.
[60] R. Dai, J. Li, Y. Liu, D. Yan, S. Chen, C. Duan, X. Liu, T. He, H. Li, miR-221/222
suppression protects against endoplasmic reticulum stress-induced apoptosis via p27(Kip1)- and
MEK/ERK-mediated cell cycle regulation, Biol Chem, 391 (2010) 791-801.
[61] Y. Zhang, J. Wu, J.Y. Liang, X. Huang, L. Xia, D.W. Ma, X.Y. Xu, P.P. Wu, Association of
serum lipids and severity of epithelial ovarian cancer: an observational cohort study of 349
Chinese patients, J Biomed Res, 32 (2018) 336-342.
[62] A.J. Li, R.G. Elmore, I.Y. Chen, B.Y. Karlan, Serum low-density lipoprotein levels
correlate with survival in advanced stage epithelial ovarian cancers, Gynecol Oncol, 116 (2010)
78-81.
[63] H. Wickham, Ggplot2 : elegrant graphics for data analysis, Use R!, Springer,, Switzerland,
2016, pp. 1 online resource.
[64] A. Kassambara, M. Kosinski, P. Biecek, survminer: Drawing Survival Curves using
'ggplot2', 2020.