Modulating the activities of chloroplasts and mitochondria promotes ATP production and plant growth

Chia Pao Voon[1,†], Yee-Song Law[1,†], Xiaoqian Guan[1,†], Shey-Li Lim[1], Zhou Xu[1], Wing-Tung Chu[1], Renshan Zhang[1], Feng Sun[1], Mathias Labs[2, &], Dario Leister[2], Mathias Pribil[3], Marie Hronková[4], Jiří Kubásek[4], Yong Cui[5,9], Liwen Jiang[5,9], Michito Tsuyama[6], Per Gardeström[7], Mikko Tikkanen[8], Boon Leong Lim[1,9*]

[1]School of Biological Sciences, the University of Hong Kong, Pokfulam, Hong Kong, China.

[2]Plant Molecular Biology, Department of Biology, Ludwig-Maximilians-University Munich (LMU), D-82152 Planegg-Martinsried, Germany.

[3]Copenhagen Plant Science Centre, Department of Plant and Environmental Sciences, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark

[4]Faculty of Science, University of South Bohemia, Branisovska 31, 370 05 Ceske Budejovice, Czech Republic.

[5]School of Life Sciences, Centre for Cell and Developmental Biology, The Chinese University of Hong Kong, Shatin, Hong Kong, China.

[6]Department of Agriculture, Kyushu University, Fukuoka 812-8581, Japan.

[7]Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, SE-901 87 Umeå, Sweden.

[8]Molecular Plant Biology, Department of Life Technologies, University of Turku, FI-20014 Turku, Finland

[9]State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong, China.

[†] These authors contributed equally to this work.

[*]Corresponding author: Boon Leong Lim (bllim@hku.hk); Tel.: (852) 22990826

[&] Current address: KWS SAAT SE, Gateway Research Center, St. Louis, MO, USA.

Running title: Cooperation of chloroplasts and mitochondria

Abstract

Efficient photosynthesis requires a balance of ATP and NADPH production/consumption in chloroplasts and the exportation of reducing equivalents from chloroplasts is important for balancing stromal ATP/NADPH ratio. Here we showed that the overexpression of purple acid phosphatase 2 on the outer membranes of chloroplasts and mitochondria can streamline the production and consumption of reducing equivalents in these two organelles, respectively. A higher capacity of consumption of reducing equivalents in mitochondria can indirectly help chloroplasts to balance the ATP/NADPH ratio in stroma and recycle $NADP^+$, the electron acceptors of the linear electron flow. A higher rate of ATP and NADPH production from the linear electron flow, a higher capacity of carbon fixation by the Calvin-Benson-Bassham cycle and a greater consumption of NADH in mitochondria, enhance photosynthesis in the chloroplasts, ATP production in the mitochondria, sucrose synthesis in the cytosol, and eventually boosting plant growth and seed yields in the overexpression lines.

Keywords

Arabidopsis thaliana, ATP, AtPAP2, chloroplasts, mitochondria, NADPH, photosynthesis, photosystem

1. Introduction

In plant cells, chloroplasts convert light energy into chemical energy, and mitochondria consume the chemical energy to produce ATP. The optimal carbon fixation and plant growth require these two energy-transforming organelles to perform strictly coordinated actions. Both organelles utilize protein complexes to construct electron transport chains (ETC), which are responsible for the formation (chloroplasts) or consumption (mitochondria) of reducing equivalents, translocation of protons, and the build-up of the proton gradient as a driving force for ATP synthesis. The cooperation between the chloroplasts and mitochondria, which was invented over 1.5 billion years ago, is nowadays much more complicated than the relationship between a supplier and a consumer in modern plant cells (Dutilleul et al., 2003; Noguchi and Yoshida, 2008). In chloroplasts, the linear electron flow (LEF) generates ATP/NADPH at a ratio of approximately 1.28, and the fixation of carbon dioxide consumes ATP/NADPH at a ratio of 1.5 (Allen, 2003; Foyer et al., 2012). Therefore, the photosynthetic efficiency requires the production and consumption of ATP and reductants at appropriate ratios in the chloroplasts, and this process is complicated by their fluxes across the chloroplast inner membrane. Our recent study showed that, in order to limit energy expenditure of chloroplasts in the dark, the importation of cytosolic ATP into mature chloroplasts is negligible (Voon et al., 2018; Voon and Lim, 2019). Hence, during photosynthesis the ATP/NADPH ratio can either be balanced by extra ATP production from the cyclic electron flow (CEF) or the export of excess reductants to the cytosol (Sato et al., 2019; Selinski and Scheibe, 2019). In addition, photorespiration also generates a large amount of NADH in mitochondria and surplus reducing equivalents are exported to the cytosol through the malate-oxaloacetate (OAA) shuttle (Lim et al., 2020). Hence, surplus reducing equivalents generated during photosynthesis have to be stored as malate in the vacuole (Gerhardt et al., 1987). Light-dependent production of reducing equivalents in mitochondria is supported by the observation that illumination can cause a rapid pH change in the mitochondrial matrix, which was unseen when DCMU was applied (Voon et al., 2018). Furthermore, when the mitochondrial ETC was interfered by inhibitors, photosynthetic ATP production in stroma was also affected (Voon et al., 2018). Here, by studying the physiology of transgenic lines that overexpress *Arabidopsis thaliana* purple acid phosphatase 2 (AtPAP2), we showed that efficient cooperation of chloroplasts and mitochondria in optimizing reductant production in chloroplasts and consumption in chloroplasts and mitochondria is important for enhancing photosynthesis and productivity.

AtPAP2 is anchored on the outer membranes of chloroplasts and mitochondria and plays

a role in protein import into these two organelles (Sun et al., 2012a; Law et al., 2015; Zhang et

al., 2016). The overexpression of AtPAP2 in *Arabidopsis thaliana* resulted in earlier bolting

(Supplementary Figure S1; Supplementary Movie S1), a higher seed yield (+40-50%), and

higher leaf sugar and ATP levels (Sun et al., 2012b; Sun et al., 2013; Liang et al., 2015). The

overexpression of AtPAP2 also promotes plant growth and seed yield of the biofuel crop

Camelina sativa (Zhang et al., 2012). Similarly to Toc33/34 and Tom20s, AtPAP2 is anchored

onto the outer membranes of chloroplasts and mitochondria via its hydrophobic *C*-terminal

motif (Sun et al., 2012a). AtPAP2 interacts with the precursor of the small subunit of RuBisCO

(pSSU) (Zhang et al., 2016) and the presequences of a number of multiple organellar RNA

editing factor (pMORF) proteins (Law et al., 2015) and plays a role in their import into

chloroplasts (Zhang et al., 2016) and mitochondria (Law et al., 2015), respectively. Here, we

examined how AtPAP2 overexpression affects the physiology of chloroplasts and mitochondria

and how these two energy-generating organelles orchestrate to produce more sugars and ATP

in leaf cells. Our data suggest that the efficiency of photosynthesis is dependent on the activities

of the mitochondria. Surplus reducing equivalents generated from the LEF have to be exported

and dissipated; and mitochondria that more actively dissipate the reducing equivalents can

supply more ATP to the cytosol, thereby simultaneously relieving the pressure of over-

reduction of ETC in the chloroplast (Scheibe et al., 2005). This streamlined cooperation enables

a higher efficacy in carbon fixation, sucrose synthesis and ATP production in leaf cells.

74   2. Methods

75   *2.1. Plant growth conditions*

The WT *Arabidopsis thaliana* ecotype Columbia-o (WT), the *Arabidopsis thaliana* AtPAP2-

overexpressing lines (OE7, OE21) and the *pap2* T-DNA insertion mutant (Salk_013567) (Sun

et al., 2012b) were used in this study. *Arabidopsis* seeds were plated for 10 days on Murashige

and Skoog (MS) medium, which was supplemented with 2% (w/v) sucrose. Seedlings of

similar sizes were transferred to soil under growth chamber conditions (120-150 µmol m^{-2} s^{-1},

16-h photoperiod, 22 °C).

83   *2.2 Chlorophyll a fluorescence measurement*

The chlorophyll *a* fluorescence levels of the *Arabidopsis* leaves (20 days old) were monitored

with IMAGING-PAM M-Series Maxi Version (WALZ, Germany). The plants were dark-

adapted for at least 1 hour before the measurements. After the maximum and initial

fluorescence (F_m and F_0, respectively) were determined with a delay of 40 s, the plants were

illuminated using the following light intensities: 0, 81, 145, 186, 281, 335, 461, 701, and 926

µmol photon m^{-2} s^{-1}. The duration of the illumination of each light intensity was 3 min. A

saturation pulse (800 ms, 2700 µmol photon m^{-2} s^{-1}) was applied at the end of the 3-min

illumination. For each light intensity, the photosynthesis parameters were calculated using the

formulas: Y(II) was calculated as (Fm' – F) /Fm' ; qP is calculated as (Fm' – F) /(Fm' – Fo');

NPQ was calculated as (Fm – Fm') /Fm' ; ETR was calculated as Y(II) x light intensity x 0.5

x absorptivity. Fm' is the maximal fluorescence of light-illuminated plant. F is the current

fluorescence yield. Fo' was estimated by Fo /(Fv /Fm + Fo /Fm') (Oxborough and Baker, 1997).

2.3 Measurements of P700 oxidation

P700 oxidation (Y(ND) = P/PM) was recorded with Dual-PAM-100 (Heinz Walz GmbH,

Effeltrich, Germany). P700 was determined by the difference of 875-nm and 830-nm

measuring lights (Klughammer and Schreiber, 2008). 3-week-old plants were illuminated using

the following light intensities: 0, 25, 50, 125, 450 and 150 µmol photon m-2 s-1. The duration

of the illumination of each light intensity was 5 min and saturating pulse (800 ms, 8000 µmol

photon m^{-2} s^{-1}) was applied in 1-min intervals.

2.4 Measurements of CO$_2$ assimilation

The carbon dioxide assimilation in the 20-day-old plants was measured using the portable

photosynthesis system LI-6400XT (LiCor, USA) and whole plant *Arabidopsis* chamber 6400-

17. The plants were cultivated hydroponically in four- times diluted Hoagland nutrient solution

(plants were fixed in "Grodan"-mineral wool cubes) under growth chamber conditions

(irradiation 120-150 µmol m^{-2} s^{-1}, 16/8 h day/night photoperiod, temperature 22°C). The plants

were transferred immediately before the measurement into conical pots dedicated to the 6400-

17 chamber, and the boundary between the substrate (roots) and rosette was established using

aluminium foil to minimize non-leaves transpiration. The light response of CO$_2$ assimilation

was measured under ambient CO$_2$ concentration 400 µmol mol^{-1} and PAR levels 2000, 1000,

500, 250, 120, 60, 30, 15 and 0 µmol photon m^{-2} s^{-1}. Stable temperature at 25 °C and the relative

air humidity between 50 and 70% were kept in the chamber during measurement.

2.5 Measurements of the post-illumination chlorophyll fluorescence transient

The modulated chlorophyll fluorescence was measured using a PAM 101 fluorometer (H. Walz,

Germany). Before the minimal fluorescence (F_0) determination, the leaves were kept in

darkness for at least 1 hour. The F_0 was induced by a red-modulated measuring light with a

photon flux density (PFD) of approx. 0.2 µmol photon m^{-2} s^{-1}. The post-illumination Chl

fluorescence increase is attributed to the back flow of electrons to the PQ pool from NADPH

in the stroma via the cyclic electron flow around photosystem I (CEF-PSI), which depends on

the NDH complex (Gotoh et al., 2010; Yamamoto et al., 2011).

2.6 Enzyme assays of the Calvin-Benson-Bassham cycle and TCA cycle

The chloroplast proteins were used to assay the fructose 1,6-bisphosphate aldolase (Schaeffer

et al., 1997) and $NADP^+$-glyceraldehyde-3-phosphate dehydrogenase activities. Enzyme

assays of the TCA cycle were performed as described previously (Huang et al., 2015). The

activity of the mitochondrial proteins (1-25 µg protein) were assayed spectrophotometrically

using a Multiskan™ GO Microplate Spectrophotometer (Thermo Scientific, USA).

2.7 Measurements of oxygen consumption rate

The respiration rate of 21-day-old fully expanded leaf was measured with a Clark-type

electrode (Oxygraph plus, Hansatech) at room temperature in the dark. Leaves were infiltrated

for 5 min in half strength MS medium and incubated in the dark for 10 min before measurement.

The respiration rate was measured in the first 10 min. 10 mM SHAM and 4 mM KCN were

then added at the 10th min and the 20th min. AOX pathway respiration rate was defined as the

sensitivity of O_2 uptake to 10 mM SHAM and the COX pathway respiration rate was defined

as the sensitivity of O_2 uptake to 4 mM KCN in the presence of 10 mM SHAM. The total

respiration rate, AOX respiration rate and COX respiration rate were calculated as nanomoles

of O_2 per min per gram of fresh weight as described previously (Zhang et al., 2017b).

Mitochondria were isolated (Law et al., 2015) and their integrity (Jacoby et al., 2015) was

checked as described previously. The respiration rate of isolated mitochondria was measured

at 25 °C in a chamber containing 1 ml of the respiration buffer (0.3 M Suc, 5 mM KH_2PO_4, 10

mM TES, 10 mM NaCl, 2 mM $MgSO_4$, and 0.1% [w/v] BSA, pH 6.8 or pH 7.2). The capacity

of complex I, internal and external NADH dehydrogenases, and complex II were measured in

the presence of deamino-NADH (1 mM), malate (10 mM), glutamate (10 mM), and NADH (1

mM) with or without ADP (100 µM) and/or rotenone (5 µM) as previously described (Meyer

et al., 2009; Jacoby et al., 2015).

3. Results

3.1 Overexpression of AtPAP2 changes the composition of the thylakoid membrane

Using transmission electron microscopy (TEM), alterations in the ultrastructure of the thylakoid membrane of mesophyll chloroplasts were observed in the OE lines (Figure 1). The average diameters of the grana stacks were approximately 0.43 μm in all four lines, which is consistent with a previous report (Armbruster et al., 2013). Conversely, the average heights of the grana stack in the OE7 and OE21 chloroplasts were smaller than the average heights of the grana stack in the wild-type (WT) and *pap2* lines (Figure 1). The decreased stacking may be due to the decreased amount of PSII (Figure 2). An analysis of the photosynthetic pigments revealed that chlorophyll a, chlorophyll b, lutein and β-carotene were reduced in the two OE lines. The reduced chlorophyll content in the OE lines correlates with the reduction in the grana stacking. The level of violaxanthin was, however, unaltered, suggesting that the xanthophyll cycle in the OE plants was suppressed to minimize the thermal loss of the excitation energy (Supplementary Table S1).

3.2 Overexpression of AtPAP2 changes the compositions of the photosystems and respiratory chain

2D BN/SDS-PAGE DIGE revealed that the abundances of specific photosystem components were altered in the OE7 chloroplasts (Figure 2a). The protein levels of the photosystem II (PSII) core proteins PsbC and PsbB were significantly reduced in OE7, while the abundances of the photosystem I (PSI) core proteins PsaA, PsaB, and PsaD were unaltered, implying that the PSI to PSII complex ratio is higher in the OE7 chloroplasts than that in the WT (Supplementary Table S2). The protein abundances of PsbO1 and RbcL (RuBisCO large subunit) were increased by at least 3-fold in the chloroplasts in the OE7 line. The ATP synthase complex subunits were significantly reduced in the OE7 line, including ATPA, ATPB, and ATPC (Supplementary Table S2). In the mitochondria, the protein abundances of the ATP synthase subunits alpha, beta and gamma were also lower in OE7 (Figure 2b, Supplementary Table S3).

3.3 LEF, but not the CEF, was enhanced in the OE lines

To investigate the impact of the altered thylakoid composition on the photosynthetic performance, the photosynthetic electron transport in 20-day-old leaves was assessed using chlorophyll fluorescence analysis (Figure 3, Supplementary Figure S2). Both AtPAP2 OE lines displayed an increased PSII quantum yield Y(II), PSII photochemical capacity (qP), and

electron transport rate (ETR) compared to the WT (Figure 3a), which suggests that the OE lines exhibited a higher LEF efficiency. No difference in terms of absorptivity was observed between the WT and OE line, implying that the altered photosynthesis parameters in OE line was not due to any changes in chlorophyll absorption capacity (Supplementary Figure S3). An analysis of the P700 redox state showed that the PSI in the OE lines was highly oxidized under low light conditions ($25 - 125$ µmol photon m^{-2} s^{-1}), suggesting that in the OE lines, the number of electrons that are consumed from the PSI by downstream pathways are higher than the number of electrons that are supplied by PSII to PSI (Figure 3b). This phenomenon was not observed when the electron flow was larger at the higher light intensities. Although the ETR and P700 redox analysis suggested that the OE lines have a higher electron transfer capacity, the NAD(P)H dehydrogenase- and Fd- dependent CEF were not significantly changed in the OE lines according to the post-illumination chlorophyll fluorescence transient and ruptured chloroplast assay, respectively (Supplementary Figure S4). In summary, the LEF, but not the CEF, was enhanced in the OE lines, and PSI played a key role in this enhancement by relaying more electrons to $NADP^+$ in the transfer chain.

3.4 Photosynthesis rate was enhanced in the chloroplasts of the AtPAP2 OE lines

To study the effect of the altered thylakoid composition on the photosynthesis performance, the photosynthetic parameters were characterized in 20-day-old leaves by an analysis of gas exchange. Both OE lines exhibited a higher light-saturated-carbon fixation rate under ambient CO_2 concentration than the WT (Figure 3c). This increased rate correlates with the higher abundance of RbcL in the OE chloroplasts (Figure 2a). Assays of the Calvin-Benson-Bassham (CBB) cycle enzymes showed that the capacities of fructose bisphosphate aldolase and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) were increased by 30 - 40% in the chloroplasts in the OE7 line (Table 1).

3.5 The mitochondria in the AtPAP2 OE line had a higher capacity in dissipating reducing equivalents

The OE lines were shown to exhibit a higher total respiratory rate than the WT in the dark, with a significantly higher COX pathway respiratory rate (Supplementary Figure S5). Reductant equivalents can be exported from the chloroplasts in the form of malate or DHAP (Shameer et al., 2019), which can be used for sucrose synthesis in the cytosol or converted to pyruvate and fed into the TCA cycle. Enzyme assays of the TCA cycle suggested that the OE7 mitochondria had an increased capacity of mitochondrial NAD^+-dependent malate dehydrogenase (mMDH)

but not in the capacities of pyruvate dehydrogenase, citrate synthase or aconitase (Table 1), whereas the capacities of chloroplast $NADP^+$-dependent malate dehydrogenase, mMDH, 2-oxoglutarate dehydrogenase (2-OGDH), and succinyl-CoA synthetase were increased by 1.26-fold, 1.15-fold, 2.86-fold and 10.3-fold in the OE7 line, respectively. Our recent studies showed that glycine decarboxylase generates a large amount of NADH in mitochondria during photorespiration, which exceeds the NADH-dissipating capacity of the mitochondria. Consequently, the surplus NADH must be exported to the cytosol through the mitochondrial malate-OAA shuttle (Lim et al., 2020). The oxygraph studies showed that the electron-feeding capacities of complex II and internal NADH dehydrogenase, but not of complex I or external NADH dehydrogenase, were enhanced in the OE7 mitochondria (Supplementary Table S4). If the OE line had a higher capacity in dissipating NADH, less surplus reducing equivalents will be exported to the cytosol through the malate-OAA shuttle.

3.6 Higher mitochondrial activity is responsible for the higher cytosolic ATP in the OE line

Our previous study showed that the leaf in the AtPAP2 OE lines contained a significantly higher level of ATP and a higher ATP/NADPH ratio in both dark and light conditions (Liang et al., 2015). Here, we introduced a FRET $MgATP^{2-}$ sensor (Imamura et al., 2009; Voon et al., 2018) into the chloroplasts and the cytosol and a pH sensor (Schwarzländer et al., 2011) to the mitochondrial matrix and compared the responses of the OE7 and WT lines (Voon et al., 2018). Illumination led to an increased ATP concentration in the stroma (Supplementary Figure S6a) and alkalization of the mitochondrial matrix in both lines (Supplementary Figure S6b). The extent of the alkalization of the mitochondrial matrix was greater in the OE7 line than that in the WT, implying that under the same light intensity, the reducing power harvested by the chloroplasts may lead to a stronger proton translocation activity across the mitochondrial inner membrane (Supplementary Figure S6b). By contrast, the rate of increase in the ATP concentration in the stroma is lower in the OE7 line than that in the WT, which is likely due to a higher ATP consumption rate in the OE7 chloroplasts because the CO_2 fixation rate (Figure 3c) and the capacities of certain CBB cycle enzymes are higher in the OE7 line (Table 1). We then examined the contribution of complex I, complex II and complex V in the OE7 mitochondria to cytosolic ATP. These inhibitors lower the cytosolic ATP in both lines (Supplementary Figure S6c). When complex I was inhibited by rotenone, the cytosolic ATP concentration was significantly higher in the OE7 line, suggesting that non-complex I activities in the OE7 mitochondria, such as complex II and internal NADH dehydrogenase, can

compensate more effectively than those in the WT (Supplementary Table S4). This difference was not observed when complex II was inhibited by TTFA. After one hour of incubation with oligomycin, cytosolic ATP in WT dropped to an undetectable level, whereas a higher level of cytosolic ATP was remained in OE7, suggesting that the intrinsic cytosolic ATP level was higher in OE7 than WT (Supplementary Figure S6c).

3.7 AtPAP2 selectively interacts with a number of chloroplast and mitochondrial proteins

A yeast two-hybrid (Y2H) library screening was carried out to identify other AtPAP2-interacting proteins. Forty nuclear-encoded AtPAP2-interacting proteins were identified; of these proteins, 32 and 3 proteins have been experimentally verified in chloroplasts and mitochondria, respectively (Supplementary Table S5). The other 5 AtPAP2-interacting proteins were predicted to be targeted to chloroplasts and/or mitochondria. AtPAP2 is exposed to the cytosolic side of the outer membrane of both organelles and its role in the importation of organellar proteins may account for the enrichment of nuclear-encoded organellar proteins. A systematic Y2H assay was then carried out to examine the interactions between AtPAP2 and various nucleus-encoded photosystem proteins (Supplementary Figure S7A). AtPAP2 could specifically interact with Fd1, Fd2, PsaE2 (but not PsaE1), ferredoxin/thioredoxin reductase subunit A2 (FTRA2) (but not FTRA1), FTRB, and photosynthetic NDH subcomplexes L1, L2, and L3 (PsbQ-like1, PsbQ-like2, and PsbQ-like3). These interactions were verified by *in vivo* bimolecular fluorescence complementation (BiFC) assays (Supplementary Figure S7B and Supplementary Figure S8). Hence, AtPAP2 may play a role in the importation of these precursor proteins, particularly the proteins at the acceptor side of PSI, into chloroplasts.

4. Discussion

Maintenance of optimal photosynthetic efficiency requires the production and consumption of ATP and reductants in chloroplasts at appropriate ratios. Our recent study showed that the importation of cytosolic ATP into mature chloroplasts of *Arabidopsis thaliana* is negligible and therefore could not supply additional ATP to the CBB cycle (Voon et al., 2018). Hence, the export of surplus reducing equivalents to extrachloroplast compartments is essential for balancing the ATP/NADPH ratio (Voon and Lim, 2019). Under illumination, the chloroplast is the producer of reducing equivalents and ATP, and the mitochondria consume the excess reducing equivalents to supply ATP to the cytosol (Gardeström and Igamberdiev, 2016; Voon et al., 2018; Shameer et al., 2019). The results presented here provide an example on how the

cooperation of chloroplasts and mitochondria boosts plant productivity.

4.1. Simultaneous alteration of chloroplast and mitochondrial activities leads to higher carbon

fixation, higher leaf ATP but lower NADPH contents in the OE lines.

Illumination greatly elevates the amount of NADPH and the NADPH/ATP ratio in WT leaves

(Liang et al., 2016), which represents a high reduction state in plant cells. Our previous studies

showed that in the middle-of-day, the leaf sucrose and ATP contents are higher in the OE lines

than those in the WT line in 20-day-old plants, whereas the leaf NADPH contents,

NADPH/NADP$^+$ and NADPH/ATP ratios are lower in the OE lines than that in the WT line

(Liang et al., 2015). Here, our data (Figure 4, Tables 1 and Supplementary Table S4) indicated

that the mitochondria of the OE line may have a higher capacity to consume extra reducing

equivalents to generate ATP (Figure 4). This mechanism could account for the high ATP, high

sucrose, but low NADPH content in the leaves of the OE lines in the middle-of-day (Sun et al.,

2012b; Liang et al., 2015). In addition, the RuBisCO content (Figure 2a) and the capacities of

the CBB cycle enzymes FBA and GAPDH (Table 1) were higher in the OE chloroplasts. These

could lead to a higher rate of CO_2 fixation (Figure 3c) and a rapid recycling of ADP, NADP$^+$

and RuBP in the stroma. Hence, a faster recycling of ribulose-1,5-bisphosphate (RuBP) and an

enhanced output of dihydroxyacetone phosphate (DHAP) from the OE chloroplasts could be

achieved. This, together with a higher consumption rate of reductants by the mitochondria

(Supplementary Table S4), might cause a lower reduction state in the stroma, thereby leading

to an enhancement of the LEF in the OE lines. This hypothesis is supported by our data from

the P700 redox analysis (Figure 3b). At low light conditions (25 – 125 µmol photon m^{-2} s^{-1}), in

which the electron flow is smaller, PSI in the OE lines was highly oxidized, which was likely

due to a high demand of electrons for the NADP$^+$ reduction or a higher PSI/PSII ratio (Figure

2a and Supplementary Table S2). This oxidized state was relieved under higher light conditions

(>125 µmol photon m^{-2} s^{-1}) when the electron flow was larger (Figure 3b).

4.2. The OE line has a higher PSI/PSII ratio, a higher LEF but indifferent CEF.

In pea (*Pisum sativum*), the PSI/PSII ratios were adjusted under different light conditions as

follows: sunlight-grown, 0.55; yellow light (preferentially excites PSII), 0.40; and red light

(preferentially excites PSI), 0.91 (Chow et al., 1990). Compared to plants with a lower PSI/PSII

ratio, plants with a higher PSI/PSII exhibit a higher rate of oxygen evolution under illumination

intensities that exceed 200 µmol photon m^{-2} s^{-1} (Chow et al., 1990). We also observed a higher

PSI/PSII ratio in the OE chloroplasts than that in the WT chloroplasts in the 2D-DIGE analysis (Figure 2a and Supplementary Table S2). The OE lines exhibited a higher ETR than the WT when the light intensity exceeded 200 µmol photon m^{-2} s^{-1} (Figure 3). A higher PSI/PSII ratio allows for a faster throughput of electrons from the photosynthetic ETC via the PSI acceptor side and results in a lower possibility of an over-reduction in ETC (Tikkanen et al., 2014). Interestingly, the OE lines had also less stacked thylakoid membranes compared to the WT (Figure 1). It is likely that the loose stacking increases the share of the grana margin domain that is composed of all components of the photosynthetic electron transfer chain in relation to the PSII-enriched grana stacks and PSI-enriched stroma lamellae. The importance of grana stacking for photosynthetic efficiency is not understood. Based on the results presented here, the loose stacking may be essential for the efficient interactions among the components of the photosynthetic electron transfer chain.

Consistently with this hypothesis, the analysis of the P700 (PSI reaction centre) redox state revealed that the P700 in the OE lines was relatively oxidized compared to that in the WT when the electron supply from PSII was limited (25 – 125 µmol photon m^{-2} s^{-1}) (Figure 3b). Y2H showed that AtPAP2 preferably interacts with the downstream components of PSI, including PsaE2, Fd1, Fd2, FTRA2 and FTRB (Supplementary Figure S7). Altogether, we hypothesize that the LEF rate in the OE lines was increased because of the higher capacity of electron transfer in PSI. LEF generates 1.28 mole ATP per mole of NADPH, but a minimum of 1.5 mole of ATP per mole of NADPH is required for carbon fixation (Allen, 2003; Foyer et al., 2012). It is generally believed that CEF pathways are needed to supplement ATP and modulate the ATP/NADPH ratio to meet the demand of metabolism (Foyer et al., 2012; Ishikawa et al., 2016). However, our studies showed that an enhancement of CEF is not required to support a more robust CBB cycle in the OE line (Supplementary Figure S4).

4.3. The mitochondrial capacity in dissipating surplus reducing equivalents is important for photosynthesis.

A recent study in diatom showed that the ATP generated from mitochondria can be transported to chloroplasts for carbon fixation (Bailleul et al., 2015), and it was believed that this also occurs in higher plants. However, our recent study showed that in the mesophyll of *Arabidopsis thaliana*, cytosolic ATP does not enter mature chloroplast to support carbon fixation (Voon et al., 2018). Altogether, because the NADPH/ATP demand for carbon dioxide fixation is approximately 0.67, and a 0.77 molecule of NADPH is generated from each ATP molecule in

LEF (Allen, 2003; Foyer et al., 2012), one could expect that a surplus reducing power is
generated from LEF under illumination. Other metabolisms in chloroplasts, such as
transcription and translation, also consume ATP molecules; therefore, the surplus of reducing
equivalents would be more evident. The surplus of reducing equivalents must be exported and
dissipated to recycle $NADP^+$ for efficient photosynthesis.

The reducing power generated from LEF can be exported as DHAP and malate (Figure 4)
(Noctor and Foyer, 2000; Scheibe, 2004). While modelling predicted that only 5% or less of
the reducing equivalents from LEF can be exported through the malate-OAA shuttle (Fridlyand
et al., 1998), the higher capacity of the $NADP^+$-MDH in the OE chloroplasts might allow for
the export of more reducing equivalents from the chloroplasts to the cytosol through the malate-
OAA shuttle. Illumination promotes the association between mitochondria and chloroplasts
and peroxisomes (Oikawa et al., 2015), and, therefore, the malate-OAA shuttle can be a very
efficient energy and reductant transfer pathway between the chloroplasts, mitochondria and
peroxisomes. Malate and OAA in the stroma and cytosol were estimated to have concentrations
ranging from 1 – 3 mM and 0.025 – 0.098 mM, respectively (Heineke et al., 1991). This
concentration ratio is maintained by the large positive value of free energy change, $\Delta G^{o\prime}$, of
malate dehydrogenation. This equilibrium can drive malate synthesis in the chloroplasts upon
illumination because the chloroplast $NADP^+$-dependent MDH is activated by light (Scheibe,
1987; Zhao et al., 2018). The reducing power, thus, can be readily channelled to the
peroxisomes for hydroxypyruvate reduction through the malate-OAA shuttle. As more
reductants are consumed by the more active OE mitochondria, less exportation of reductants
from mitochondria is expected (Lim et al., 2020). Hence, more reductants from the chloroplasts
can be dissipated indirectly (Figure 4). This is in line with our recent observation that the
increase in stromal NADPH during illumination disappeared when photorespiration was absent
(Abdel-Ghany, 2009; Lim et al., 2020).

4.4. The abundance of ATP synthases is not necessarily correlated with the cellular ATP levels.
While the protein abundance of the ATP synthase subunits in the chloroplasts (Figure 2a,
Supplementary Table S2) and mitochondria (Figure 2b, Supplementary Table S3) were
significantly reduced in the OE7 line, the leaves of the OE lines contain higher ATP levels than
the WT (Sun et al., 2012b; Sun et al., 2013; Liang et al., 2015). In Arabidopsis cotyledon, the
cytosolic ATP concentration (> 1.4 mM) is substantially higher than the stromal ATP
concentration (~0.2 mM). Hence, the higher leaf ATP concentration of the OE line is likely to

be contributed by the cytosolic ATP generated from the mitochondrial ATP synthase, which is supported by the cytosolic ATP sensor data (Supplementary Fig. 6c) and oxygraph data (Supplementary table 4). A lower abundance of ATP synthase does not contradict with a higher ATP content in the OE7 line. The output of ATP synthase is dependent on the magnitude of the proton motive force (Schmidt and Graber, 1987; Zubareva et al., 2020), and may also be affected by the fuel supply (NADH) to the mETC and ADP turnover, etc. Despite of a lower abundance, the mitochondrial ATP synthase of the OE7 line may have a higher output than that of the WT.

4.5. Summary

In summary, the overexpression of AtPAP2 modulates the importation of selected nucleus-encoded proteins into chloroplasts (Zhang et al., 2016) and mitochondria (Law et al., 2015), which leads to stronger sinks of reductants, including higher activities in the CBB cycle and mitochondrial ETC, in both organelles. A higher output of reducing equivalents from the LEF of chloroplasts (Figure 3) and higher mitochondrial activities in utilizing the reducing equivalents for ATP production, thus, contribute to the higher ATP content and ATP/NADPH ratio in the OE lines (Liang et al., 2015). The simultaneous activation of chloroplasts and mitochondria is required for the production of a surplus of ATP and sucrose; the transgenic lines that overexpress AtPAP2 solely in the chloroplasts grew similarly to the WT (unpublished data), and the transgenic lines that overexpress AtPAP2 solely in the mitochondria exhibited early senescence and a lower seed yield (Law et al., 2015). These results provide an example of how the efficient cooperation of chloroplasts and mitochondria can enhance ATP and sucrose production in leaf cells. While AtPAP2 can enhance yield and growth in *Arabidopsis thaliana*, it has negative impacts to plant survival. The OE lines are less resistant to drought and to *P. syringae* infection (Zhang et al., 2017a). Therefore, while AtPAP2-like genes were evolved from green algae during evolution (Sun et al., 2012b), it may not be fully used by higher plants to promote growth in natural environment, as survival is more important than high yield. It may be more 'useful' for unicellular algae as water is never a constraint, and algae need to control their growth rate and carbon level subject to the availability of P/N/Fe.

Over the past few decades, a controversy has emerged in the field regarding whether the shortfall of ATP for the CBB cycle is fulfilled by the CEF or the importation of ATP from the cytosol. Combining the findings of our recent study (Voon et al., 2018) and this study, we concluded that the efficient carbon fixation in the OE line is not dependent on an enhanced

CEF or the importation of ATP from the cytosol to fulfil the shortfall of ATP generated from

the LEF. Instead, an efficient carbon fixation is dependent on an enhanced LEF, enhanced

capacities of CBB enzymes, the efficient export of surplus reducing equivalents from

chloroplasts through the malate-OAA shuttle and a higher reductant-dissipating activity of

mitochondria. The ability of mitochondria to dissipate reducing equivalents is important for

relieving the built-up of surplus reducing power in the stroma, which, in excess, will limit the

LEF due to an insufficient supply of $NADP^+$, the major electron acceptor of the LEF. Our recent

studies showed that the reducing equivalents from photorespiration are the major fuel for

mitochondria in *Arabidopsis thaliana* (Lim et al., 2020). The optimal use of reducing

equivalents by mitochondria not only generates more ATP for sucrose synthesis, but also

reduces the export of reducing equivalents from the mitochondria and therefore the overall

redox status of the photosynthetic cells. This provides a good example to illustrate the

relationship between chloroplasts and mitochondria in bioenergetics.

Acknowledgements

We thank Prof. Wah Soon Chow, Dr. Boris Feniouk and Dr. Maurice Cheung for their comments on this manuscript. We thank Dr. Huang Shaobai and Dr. Richard Jacoby for training of YL in enzyme assays, oxygraph and 2D-DIGE. We thank Prof. Hans-Peter Braun and Dr. Holger Eubel for technical advice in the 2D BN/SDS-PAGE DIGE analysis.

Financial Support. This project was supported by the Seed Funding Program for Basic Research (201311159043) of the University of Hong Kong, the General Research Fund (772012M) and the Area of Excellence Scheme (AoE/M-403/16) of the Hong Kong Research Grants Council, the Innovation and Technology Fund (Funding Support to State Key Laboratory of Agrobiotechnology) of the HKSAR, China. Access to LI-6400 was supported by the Czech research infrastructure for systems biology C4SYS (project no LM2015055).

Conflicts of Interest. AtPAP2 is the subject of US patent number 9,476,058.

Authorship Contributions. CPV, YL, and BLL wrote the manuscript with PG's input. CPV produced some of the transgenic sensor lines, performed the fluorescent sensor measurements, carried out the Maxi-Pam measurements and 2D BN-PAGE of the chloroplasts. YL carried out the 2D BN-PAGE of the mitochondria, the enzyme assays with WC, and the mitochondrial oxygraph assays with ZX. XG screened the yeast two-hybrid library and carried out the BiFC. SL carried out respiration, AOX and COX oxygraph assays. RZ and FS produced the overexpression lines. FS measured the chlorophyll content. ML, MP and DL carried out the chloroplast rupture assay, MTik carried out the P700 assay, and MTsu carried out the post-illumination chlorophyll fluorescence. JK and MH carried out the CO_2 exchange measurements. YC and LJ carried out the TEM analysis. BLL coordinated this study. All authors read and approved the manuscript.

Data and Coding Availability Statement. The authors declare that the main data supporting the findings of this study are available within the article and its Supplementary Information files. Extra data are available from the corresponding author upon request.

Supplementary Materials. To view Supplementary material for this article, please visit

Supplementary Figure S1. The OE lines grew faster under both SD (8h/16h) and LD (16h/8h)

conditions.

Supplementary Figure S2. Fluorescence kinetics of WT, *pap2,* OE7, and OE21 lines.

Supplementary Figure S3. Leaf absorptivity is similar between WT line and AtPAP2

overexpression line (OE7).

Supplementary Figure S4. *In vivo* analysis of the cyclic electron flow rate.

Supplementary Figure S5. Respiratory rate of 21-day-old leaves.

Supplementary Figure S6. Assessing organelle activities using fluorescent sensors.

Supplementary Figure S7. AtPAP2 selectively interacts with certain photosystem proteins.

Supplementary Figure S8. Results of BiFC negative controls.

Supplementary Table S1. Photosynthetic pigment content.

Supplementary Table S2. Chloroplast proteins identified in 2D BN PAGE.

Supplementary Table S3. Summary of identified mitochondrial protein spots.

Supplementary Table S4. Respiration rates of different complexes of WT and OE7

mitochondria.

Supplementary Table S5. AtPAP2-interacting proteins identified by Y2H library screening.

Supplementary Table S6. Primer list.

References

References

Abdel-Ghany, S.E. (2009). Contribution of plastocyanin isoforms to photosynthesis and
copper homeostasis in Arabidopsis thaliana grown at different copper regimes. Planta **229,** 767-
779.

Allen, J.F. (2003). Cyclic, pseudocyclic and noncyclic photophosphorylation: new links
in the chain. Trends in plant science **8,** 15-19.

Armbruster, U., Labs, M., Pribil, M., Viola, S., Xu, W., Scharfenberg, M., Hertle,
A.P., Rojahn, U., Jensen, P.E., Rappaport, F., Joliot, P., Dormann, P., Wanner, G., and
Leister, D. (2013). Arabidopsis CURVATURE THYLAKOID1 proteins modify thylakoid
architecture by inducing membrane curvature. Plant Cell **25,** 2661-2678.

Bailleul, B., Berne, N., Murik, O., Petroutsos, D., Prihoda, J., Tanaka, A., Villanova,
V., Bligny, R., Flori, S., Falconet, D., Krieger-Liszkay, A., Santabarbara, S., Rappaport,
F., Joliot, P., Tirichine, L., Falkowski, P.G., Cardol, P., Bowler, C., and Finazzi, G. (2015).
Energetic coupling between plastids and mitochondria drives CO_2 assimilation in diatoms.
Nature **524,** 366-369.

Chow, W.S., Melis, A., and Anderson, J.M. (1990). Adjustments of photosystem
stoichiometry in chloroplasts improve the quantum efficiency of photosynthesis. Proc Natl
Acad Sci U S A **87,** 7502-7506.

Dutilleul, C., Driscoll, S., Cornic, G., De Paepe, R., Foyer, C.H., and Noctor, G. (2003).
Functional mitochondrial complex I is required by tobacco leaves for optimal photosynthetic
performance in photorespiratory conditions and during transients. Plant Physiol **131,** 264-275.

Foyer, C.H., Neukermans, J., Queval, G., Noctor, G., and Harbinson, J. (2012).
Photosynthetic control of electron transport and the regulation of gene expression. J Exp Bot
63, 1637-1661.

Fridlyand, L.E., Backhausen, J.E., and Scheibe, R. (1998). Flux control of the malate
valve in leaf cells. Arch Biochem Biophys **349,** 290-298.

Gardeström, P., and Igamberdiev, A.U. (2016). The origin of cytosolic ATP in
photosynthetic cells. Physiol Plantarum **157,** 367-379.

Gerhardt, R., Stitt, M., and Heldt, H.W. (1987). Subcellular metabolite levels in spinach
leaves : regulation of sucrose synthesis during diurnal alterations in photosynthetic partitioning.
Plant Physiol **83,** 399-407.

Gotoh, E., Matsumoto, M., Ogawa, K., Kobayashi, Y., and Tsuyama, M. (2010). A
qualitative analysis of the regulation of cyclic electron flow around photosystem I from the
post-illumination chlorophyll fluorescence transient in Arabidopsis: a new platform for the in
vivo investigation of the chloroplast redox state. Photosynthesis research **103,** 111-123.

Heineke, D., Riens, B., Grosse, H., Hoferichter, P., Peter, U., Flugge, U.I., and Heldt,
H.W. (1991). Redox transfer across the inner chloroplast envelope membrane. Plant Physiol
95, 1131-1137.

Huang, S., Lee, C.P., and Millar, A.H. (2015). Activity assay for plant mitochondrial
enzymes. Methods Mol Biol **1305,** 139-149.

Imamura, H., Nhat, K.P., Togawa, H., Saito, K., Iino, R., Kato-Yamada, Y., Nagai, T.,
and Noji, H. (2009). Visualization of ATP levels inside single living cells with fluorescence
resonance energy transfer-based genetically encoded indicators. Proc Natl Acad Sci U S A **106,**
15651-15656.

Ishikawa, N., Takabayashi, A., Sato, F., and Endo, T. (2016). Accumulation of the
components of cyclic electron flow around photosystem I in C4 plants, with respect to the
requirements for ATP. Photosynthesis research **129,** 261-277.

Jacoby, R.P., Millar, A.H., and Taylor, N.L. (2015). Assessment of respiration in isolated
plant mitochondria using clark-type electrodes. In Plant Mitochondria: Methods and Protocols,
J. Whelan and M.W. Murcha, eds (New York, NY: Springer New York), pp. 165-185.

Klughammer, C., and Schreiber, U. (2008). Saturation pulse method for assessment of
energy conversion in PS I   PAM Appl. Notes **1,** 11-14.

Law, Y.S., Zhang, R., Guan, X., Cheng, S., Sun, F., Duncan, O., Murcha, M., Whelan,
J., and Lim, B.L. (2015). Phosphorylation and dephosphorylation of the presequence of
pMORF3 during import into mitochondria from *Arabidopsis thaliana*. Plant Physiology **169,**
1-12.

Liang, C., Zhang, Y., Cheng, S., Osorio, S., Sun, Y., Fernie, A.R., Cheung, C.Y.M.,
and Lim, B.L. (2015). Impacts of high ATP supply from chloroplasts and mitochondria on the
leaf metabolism of *Arabidopsis thaliana*. Frontiers in plant science **6,** 922.

Liang, C., Cheng, S., Zhang, Y., Sun, Y., Fernie, A.R., Kang, K., Panagiotou, G., Lo,
C., and Lim, B.L. (2016). Transcriptomic, proteomic and metabolic changes in Arabidopsis
thaliana leaves after the onset of illumination. BMC Plant Biol **16,** 43.

Lim, S.L., Voon, C.P., Guan, X., Yang, Y., Gardestrom, P., and Lim, B.L. (2020). *In*
planta study of photosynthesis and photorespiration using NADPH and NADH/NAD$^+$
fluorescent protein sensors. Nat Commun **11,** 3238.

Meyer, E.H., Tomaz, T., Carroll, A.J., Estavillo, G., Delannoy, E., Tanz, S.K., Small,
I.D., Pogson, B.J., and Millar, A.H. (2009). Remodeled respiration in ndufs4 with low
phosphorylation efficiency suppresses Arabidopsis germination and growth and alters control
of metabolism at night. Plant Physiol **151,** 603-619.

Noctor, G., and Foyer, C.H. (2000). Homeostasis of adenylate status during
photosynthesis in a fluctuating environment. J Exp Bot **51,** 347-356.

Noguchi, K., and Yoshida, K. (2008). Interaction between photosynthesis and respiration
in illuminated leaves. Mitochondrion **8,** 87-99.

Oikawa, K., Matsunaga, S., Mano, S., Kondo, M., Yamada, K., Hayashi, M., Kagawa,
T., Kadota, A., Sakamoto, W., Higashi, S., Watanabe, M., Mitsui, T., Shigemasa, A., Iino,
T., Hosokawa, Y., and Nishimura, M. (2015). Physical interaction between peroxisomes and

chloroplasts elucidated by in situ laser analysis. Nature Plants **1,** 15035.

Oxborough, K., and Baker, N.R. (1997). Resolving chlorophyll a fluorescence images
of photosynthetic efficiency into photochemical and non-photochemical components –
calculation of qP and Fv-/Fm-; without measuring Fo. Photosynthesis Research **54,** 135-142.

Sato, R., Kawashima, R., Trinh, M.D.L., Nakano, M., Nagai, T., and Masuda, S.
(2019). Significance of PGR5-dependent cyclic electron flow for optimizing the rate of ATP
synthesis and consumption in Arabidopsis chloroplasts. Photosynthesis research **139,** 359-365.

Schaeffer, G.W., Sharpe, F.T., and Sicher, R.C. (1997). Fructose 1,6-bisphosphate
aldolase activity in leaves of a rice mutant selected for enhanced lysine. Phytochemistry **46,**
1335-1338.

Scheibe, R. (1987). $NADP^+$-malate dehydrogenase in C3 plants - regulation and role of a
light-activated enzyme. Physiol Plantarum **71,** 393-400.

Scheibe, R. (2004). Malate valves to balance cellular energy supply. Physiol Plantarum
120, 21-26.

Scheibe, R., Backhausen, J.E., Emmerlich, V., and Holtgrefe, S. (2005). Strategies to
maintain redox homeostasis during photosynthesis under changing conditions. J Exp Bot **56,**
1481-1489.

Schmidt, G., and Graber, P. (1987). The rate of ATP synthesis catalyzed by reconstituted
CF0F1-liposomes - dependence on delta-pH and delta-psi. Biochimica Et Biophysica Acta **890,**
392-394.

Schwarzländer, M., Logan, D.C., Fricker, M.D., and Sweetlove, L.J. (2011). The
circularly permuted yellow fluorescent protein cpYFP that has been used as a superoxide probe
is highly responsive to pH but not superoxide in mitochondria: implications for the existence
of superoxide 'flashes'. Biochem J **437,** 381-387.

Selinski, J., and Scheibe, R. (2019). Malate valves: old shuttles with new perspectives.
Plant Biol (Stuttg) **21,** 21-30.

Shameer, S., Ratcliffe, R.G., and Sweetlove, L.J. (2019). Leaf energy balance requires
mitochondrial respiration and export of chloroplast NADPH in the light. Plant Physiology **180,**
1947-1961.

Sun, F., Liang, C., Whelan, J., Yang, J., Zhang, P., and Lim, B.L. (2013). Global
transcriptome analysis of AtPAP2-overexpressing *Arabidopsis thaliana* with elevated ATP.
Bmc Genomics **14,** 752.

Sun, F., Carrie, C., Law, S., Murcha, M.W., Zhang, R., Law, Y.S., Suen, P.K., Whelan,
J., and Lim, B.L. (2012a). AtPAP2 is a tail-anchored protein in the outer membrane of
chloroplasts and mitochondria. Plant signaling & behavior **7,** 927-932.

Sun, F., Suen, P.K., Zhang, Y., Liang, C., Carrie, C., Whelan, J., Ward, J.L., Hawkins,
N.D., Jiang, L., and Lim, B.L. (2012b). A dual-targeted purple acid phosphatase in
Arabidopsis thaliana moderates carbon metabolism and its overexpression leads to faster plant

growth and higher seed yield. The New phytologist **194,** 206-219.

Tikkanen, M., Mekala, N.R., and Aro, E.M. (2014). Photosystem II photoinhibition-
repair cycle protects Photosystem I from irreversible damage. Bba-Bioenergetics **1837,** 210-
215.

Voon, C.P., and Lim, B.L. (2019). ATP translocation and chloroplast biology National
Science Review **6,** 1073-1076.

Voon, C.P., Guan, X., Sun, Y., Sahu, A., Chan, M.N., Gardeström, P., Wagner, S.,
Fuchs, P., Nietzel, T., Versaw, W.K., Schwarzländer, M., and Lim, B.L. (2018). ATP
compartmentation in plastids and cytosol of *Arabidopsis thaliana* revealed by fluorescent
protein sensing. Proc Natl Acad Sci U S A **115,** 10778-10787.

Yamamoto, H., Peng, L., Fukao, Y., and Shikanai, T. (2011). An Src homology 3
domain-like fold protein forms a ferredoxin binding site for the chloroplast NADH
dehydrogenase-like complex in Arabidopsis. Plant Cell **23,** 1480-1493.

Zhang, R., Qi, H., Sun, Y., Xiao, S., and Lim, B.L. (2017a). Transgenic *Arabidopsis*
thaliana containing increased levels of ATP and sucrose is more susceptible to *Pseudomonas*
syringae. PLoS One **12,** e0171040.

Zhang, R., Guan, X., Law, Y.S., Sun, F., Chen, S., Wong, K.B., and Lim, B.L. (2016).
AtPAP2 modulates the import of the small subunit of Rubisco into chloroplasts. Plant signaling
& behavior **11,** e1239687.

Zhang, Y., Yu, L., Yung, K.F., Leung, D.Y., Sun, F., and Lim, B.L. (2012). Over-
expression of AtPAP2 in *Camelina sativa* leads to faster plant growth and higher seed yield.
Biotechnology for biofuels **5,** 19.

Zhang, Z.S., Liu, M.J., Scheibe, R., Selinski, J., Zhang, L.T., Yang, C., Meng, X.L.,
and Gao, H.Y. (2017b). Contribution of the alternative respiratory pathway to PSII
photoprotection in C3 and C4 plants. Molecular plant **10,** 131-142.

Zhao, Y.N., Luo, L.L., Xu, J.S., Xin, P.Y., Guo, H.Y., Wu, J., Bai, L., Wang, G.D.,
Chu, J.F., Zuo, J.R., Yu, H., Huang, X., and Li, J.Y. (2018). Malate transported from
chloroplast to mitochondrion triggers production of ROS and PCD in *Arabidopsis thaliana*.
Cell Res **28,** 448-461.

Zubareva, V.M., Lapashina, A.S., Shugaeva, T.E., Litvin, A.V., and Feniouk, B.A.
(2020). Rotary Ion-Translocating ATPases/ATP Synthases: Diversity, Similarities, and
Differences. Biochemistry-Moscow+ **85,** 1613-1630.

Mean ± SEM	Diameter (nm)	Height (nm)
WT	432 ± 19 [a]	125 ± 8 [a]
pap2	444 ± 11 [a]	102 ± 5 [b]
OE7	431 ± 16 [a]	78 ± 4 [c]
OE21	430 ± 12 [a]	79 ± 3 [c]

Fig. 1. Comparison of the thylakoid architecture between the WT and OE plants. TEM micrograph of ultrathin sections of 20-day-old leaves from the WT, OE (OE7 and OE21) and AtPAP2 knock-out (*pap2*) lines. Average value (n>30) of the diameter (nm) and height (nm) of the thylakoid in the WT and OE lines were shown. Values marked by different letters in the same column are significantly different ($p < 0.05$) by Student's t-test. The diameter (nm) and height (nm) of WT chloroplasts isolated from 28-day-old leaves in another study (15) were 448 $\pm$ 16 nm and 113 $\pm$ 5 nm, respectively.

Fig. 2. Comparative analysis of the chloroplast and mitochondrial proteomes of WT and OE7

plants. (a) Chloroplasts; (b) Mitochondria. Proteins from the WT fraction were pre-labelled
with Cy3; proteins from the OE fraction were pre-labelled with Cy5. Combined protein
fractions were separated by 2D Blue-native/SDS-PAGE, and the protein visualization was
carried out by laser scanning at the respective wavelengths using the Typhoon laser scanner.
On the resulting overlay image, Cy3 is represented by red, and Cy5 is represented by green.
Proteins with a reduced abundance in the OE7 line are shown in red; proteins with an increased
abundance in the OE7 line are shown in green; and proteins of equal abundance in the two
compared fractions are shown in yellow. Protein spots were extracted and identified by an
MS/MS analysis. Identified proteins with the highest unused score and at least 2 unique
peptides (95%) were labelled with the corresponding spot ID (Supplementary Tables S2 & S3).
The representative gel images of 3 biological replicates are presented.

Fig. 3. *In vivo* analysis of the electron transport activity and CO_2 assimilation rate. (a). Light intensity-dependent NPQ, qP, ETR and Y(II). The 3-week-old plants were dark-acclimated for 1 hour before the measurement. Data are presented as the mean $\pm$ SE (n > 10 per line). (b) P700 Oxidation. Data are presented as the mean $\pm$ SD (n = 3 per line). Light intensity was increased in a step-wise manner as stated in the graph. (c) Light response of CO_2 assimilation rates (An) under ambient CO_2 concentration. Data are presented as the mean $\pm$ SE (n = 5 per line). The asterisks indicate significant differences between the WT and both OE lines by one-way ANOVA with post-hoc Tukey HSD test (p < 0.05)

Fig. 4. A model on how efficient collaboration between chloroplasts and mitochondria promote ATP and sucrose production. ① and ② AtPAP2 on the outer membranes of chloroplasts and mitochondria promotes the import of certain proteins into these two organelles via the Toc or the Tom complexes. ③ Higher PSI/PSII ratio and higher LEF generate more NADPH and ATP at a ratio of 0.78, which are consumed at a ratio of 0.67 by the enhanced CBB enzymes in the OE chloroplasts for CO_2 fixation. The surplus reducing equivalents are exported from the chloroplasts via the malate/OAA shuttle to recycle $NADP^+$ as the electron acceptors of the LEF. ④ Higher reductant-dissipating activities of OE mitochondria reduce the needs for mitochondria to export reductants from photorespiration in the form of malate. ⑤ OE chloroplasts with enhanced rate of carbon fixation export more carbon skeletons to the cytosol. ⑥ OE mitochondria with higher reductant-dissipating activities generate more ATP through the respiratory electron transfer (RET) chain. ⑦ Higher ATP production in OE mitochondria and higher output of carbon skeletons from chloroplasts enhance sucrose synthesis in the cytosol. Red and blue lines indicate upregulated and downregulated pathways/metabolites in the OE lines, respectively.

674 **Table 1.** Enzyme activities of the CBB cycle and the Krebs cycle in 20-day-old plants.

Enzyme	Source	WT	OE7	Change
Fructose bisphosphate aldolase	Chloroplast	92 ± 8	$123 \pm 14^{**}$	1.34X
$NADP^+$-GAPDH	Chloroplast	97 ± 11	$138 \pm 4^{**}$	1.42X
$NADP^+$-MDH	Chloroplast	328 ± 20	$413 \pm 40^{**}$	1.26X
$NADP^+$-MDH	Leaf	26 ± 4	28 ± 4	N.S.
NAD^+-MDH	Leaf	1525 ± 25	1439 ± 39	N.S.
NAD^+-MDH	Mitochondria	4657 ± 206	$5354 \pm 46^{**}$	1.15X
NAD^+-malic enzyme, ME	Mitochondria	40 ± 3	42 ± 3	N.S.
Pyruvate dehydrogenase, PDC	Mitochondria	45 ± 6	48 ± 4	N.S.
Citrate synthase, CS	Mitochondria	84 ± 12	90 ± 12	N.S.
Aconitase, ACN	Mitochondria	314 ± 18	322 ± 18	N.S.
NAD^+-ICDH	Mitochondria	43 ± 3	$63 \pm 8^{**}$	1.47X
$NADP^+$-ICDH	Leaf	18.5 ± 2.2	$6.7 \pm 0.5^{**}$	0.36X
2-oxoglutarate dehydrogenase, 2OGDH	Mitochondria	1.4 ± 0.6	$4.0 \pm 0.4^{**}$	2.86X
Succinate dehydrogenase, SDH	Mitochondria	7.6 ± 0.3	$10.3 \pm 1.0^{**}$	1.35X
Succinyl-CoA synthetase	Mitochondria	29 ± 1.4	$300 \pm 63^{**}$	10.3X
Fumarase, FUM	Mitochondria	334 ± 55	$209 \pm 34^{**}$	0.62X

675 *GAPDH*, glyceraldehyde-3-phosphate dehydrogenase; *ICDH*, isocitrate dehydrogenase; *MDH*, malate

676 dehydrogenase. All enzyme unit is presented as nmol/min/mg protein.

677 Independent sample t-test was carried out. Significant differences between WT and OE7 are showed by asterisks.

678 * $p < 0.05$, ** $p < 0.01$, n=4. N.S.: Not significant.