**Efficient cooperation of chloroplasts and mitochondria enhances ATP and sucrose**
**production**

Chia Pao Voon[1,†], Yee-Song Law[1,†], Xiaoqian Guan[1,†], Zhou Xu[1], Wing-Tung Chu[1], Renshan

Zhang[1], Feng Sun[1], Mathias Labs[2, &], Mathias Pribil[2, ‡], Dario Leister[2], Marie Hronková[3], Jiří

Kubásek[3], Yong Cui[4,8], Liwen Jiang[4,8], Michito Tsuyama[5], Per Gardeström[6], Mikko Tikkanen[7],

Boon Leong Lim[1,8*]

[1]School of Biological Sciences, the University of Hong Kong, Pokfulam, Hong Kong, China.

[2]Plant Molecular Biology, Department of Biology, Ludwig-Maximilians-University Munich

(LMU), D-82152 Planegg-Martinsried, Germany.

[3]Faculty of Science, University of South Bohemia, Branisovska 31, 370 05 Ceske Budejovice,

Czech Republic.

[4]School of Life Sciences, Centre for Cell and Developmental Biology, The Chinese

University of Hong Kong, Shatin, Hong Kong, China.

[5]Department of Agriculture, Kyushu University, Fukuoka 812-8581, Japan.

[6]Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, SE-901 87

Umeå, Sweden.

[7]Department of Biochemistry and Food Chemistry, Molecular Plant Biology, University of

Turku, FI-20014 Turku, Finland.

[8]State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin,

Hong Kong, China.

[†] These authors contributed equally to this work.

[*]Corresponding author: Boon Leong Lim (bllim@hku.hk); Tel.: (852) 22990826

[&] Current address: KWS SAAT SE, Gateway Research Center, St. Louis, MO, USA.

[‡] Current address: Copenhagen Plant Science Centre (CPSC), Department of Plant and

Environmental Sciences, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg

C, Denmark.

**Running title:** Cooperation of chloroplasts and mitochondria

**ABSTRACT**

Efficient photosynthesis requires a balance of ATP and NADPH production/consumption in chloroplasts as the linear electron flow generates a higher NADPH/ATP ratio than that is consumed by the Calvin-Benson-Bassham cycle. Recent works suggested that ATP importation into mature chloroplasts of *Arabidopsis thaliana* is negligible, and therefore the exportation of reducing equivalents from chloroplasts is important for balancing stromal ATP/NADPH ratio. Here we showed that the overexpression of purple acid phosphatase 2 on the outer membranes of chloroplasts and mitochondria can streamline the production and consumption of reducing equivalents in these two organelles, respectively. A higher capacity of consumption of reducing equivalents in mitochondria can indirectly help chloroplasts to balance the ATP/NADPH ratio in stroma and recycle $NADP^+$, the electron acceptors of the linear electron flow. A higher rate of ATP and NADPH production from the linear electron flow, a higher capacity of carbon fixation by the Calvin-Benson-Bassham cycle and a greater consumption of NADH in mitochondria, enhance photosynthesis in the chloroplasts, ATP production in the mitochondria, sucrose synthesis in the cytosol, and eventually boosting plant growth and seed yields in the overexpression lines.

**Keywords**

ATP, chloroplasts, mitochondria, NADPH, photosynthesis

**One-Sentence Summary**

This study demonstrates the importance of chloroplast-mitochondria cooperation in redox balance and illustrates that an optimized function of mitochondria can enhance the efficiency of photosynthesis.

## Introduction

In plant cells, chloroplasts convert light energy into chemical energy, and mitochondria consume the chemical energy to produce ATP. The optimal carbon fixation and plant growth require these two energy-transforming organelles to perform strictly coordinated actions. Both organelles utilize protein complexes to construct electron transport chains (ETC), which are responsible for the formation (chloroplasts) or consumption (mitochondria) of reducing equivalents, translocation of protons, and the build-up of the proton gradient as a driving force for ATP synthesis. The cooperation between the chloroplasts and mitochondria, which was invented over 1.5 billion years ago, is nowadays much more complicated than the relationship between a supplier and a consumer in modern plant cells (Dutilleul et al., 2003; Noguchi and Yoshida, 2008). In chloroplasts, the linear electron flow (LEF) generates ATP/NADPH at a ratio of approximately 1.28, and the fixation of carbon dioxide consumes ATP/NADPH at a ratio of 1.5 (Allen, 2003; Foyer et al., 2012). Therefore, the photosynthetic efficiency requires the production and consumption of ATP and reductants at appropriate ratios in the chloroplasts, and this process is complicated by their fluxes across the chloroplast inner membrane. Our recent study showed that, in order to limit energy expenditure of chloroplasts in the dark, the import of cytosolic ATP into mature chloroplasts is negligible (Voon et al., 2018; Voon and Lim, 2019). Hence, during photosynthesis the ATP/NADPH ratio can either be balanced by extra ATP production from the cyclic electron flow (CEF) or the export of excess reductants to the cytosol (Sato et al., 2019; Selinski and Scheibe, 2019). In addition, photorespiration also generates a large amount of NADH in mitochondria and surplus reducing equivalents are exported to the cytosol through the malate-oxaloacetate (OAA) shuttle (Lim et al., 2020). Hence, surplus reducing equivalents generated during photosynthesis have to be stored as malate in the vacuole (Gerhardt et al., 1987). Light-dependent production of reducing equivalents in mitochondria is supported by the observation that illumination can cause a rapid pH change in the mitochondrial matrix, which was unseen when DCMU was applied (Voon et al., 2018). Furthermore, when the mitochondrial ETC was interfered by inhibitors, photosynthetic ATP production in stroma was also affected (Voon et al., 2018). Here, by studying the physiology of transgenic lines that overexpress *Arabidopsis thaliana* purple acid phosphatase 2 (AtPAP2), we showed that efficient cooperation of chloroplasts and mitochondria in optimizing reductant production in chloroplasts and consumption in chloroplasts and mitochondria is important for enhancing photosynthesis and productivity.

AtPAP2 is anchored on the outer membranes of chloroplasts and mitochondria and plays a

role in protein import into these two organelles (Sun et al., 2012; Law et al., 2015; Zhang et al.,

2016). The overexpression of AtPAP2 in *Arabidopsis thaliana* resulted in earlier bolting

(Supplemental Fig. S1; Supplemental Video S1), a higher seed yield (+40-50%), and higher

leaf sugar and ATP levels (Sun et al., 2012; Sun et al., 2013; Liang et al., 2015). Similarly to

Toc33/34 and Tom20s, AtPAP2 is anchored onto the outer membranes of chloroplasts and

mitochondria via its hydrophobic *C*-terminal motif (Sun et al., 2012). AtPAP2 interacts with

the precursor of the small subunit of RuBisCO (pSSU) (Zhang et al., 2016) and the

presequences of a number of multiple organellar RNA editing factor (pMORF) proteins (Law

et al., 2015) and plays a role in their import into chloroplasts (Zhang et al., 2016) and

mitochondria (Law et al., 2015), respectively. Here, we examined how AtPAP2 overexpression

affects the physiology of chloroplasts and mitochondria and how these two energy-generating

organelles orchestrate to produce more sugars and ATP in leaf cells. Our data suggest that the

efficiency of photosynthesis is dependent on the activities of the mitochondria. Surplus

reducing equivalents generated from the LEF have to be exported and dissipated; and

mitochondria that more actively dissipate the reducing equivalents can supply more ATP to

the cytosol, thereby simultaneously relieving the pressure of over-reduction of ETC in the

chloroplast (Scheibe et al., 2005). This streamlined cooperation enables a higher efficacy in

carbon fixation, sucrose synthesis and ATP production in leaf cells.

**Results**

***Overexpression of AtPAP2 changes the composition of the thylakoid membrane.*** Using

transmission electron microscopy (TEM), alterations in the ultrastructure of the thylakoid

membrane of mesophyll chloroplasts were observed in the OE lines (Fig. 1). The average

diameters of the grana stacks were approximately 0.43 μm in all four lines, which is consistent

with a previous report (Armbruster et al., 2013). Conversely, the average heights of the grana

stacks in the OE7 and OE21 chloroplasts were smaller than the average heights of the grana

stacks in the wild-type (WT) and *pap2* lines (Fig. 1). The decreased stacking may be due to the

decreased amount of PSII (Fig. 2). An analysis of the photosynthetic pigments revealed that

chlorophyll a, chlorophyll b, lutein and β-carotene were reduced in the two OE lines. The

reduced chlorophyll content in the OE lines correlates with the reduction in the grana stacking.

The level of violaxanthin was, however, unaltered, suggesting that the xanthophyll cycle in the

OE plants was suppressed to minimize the thermal loss of the excitation energy (Supplemental

Table S1).

***Overexpression of AtPAP2 changes the compositions of the photosystems and respiratory chain.*** 2D BN/SDS-PAGE DIGE revealed that the abundances of specific photosystem components were altered in the OE7 chloroplasts (Fig. 2A). The protein levels of the photosystem II (PSII) core proteins PsbC and PsbB were significantly reduced in OE7, while the abundances of the photosystem I (PSI) core proteins PsaA, PsaB, and PsaD were unaltered, implying that the PSI to PSII complex ratio is higher in the OE7 chloroplasts than that in the WT (Supplemental Table S2). The protein abundances of PsbO1 and RbcL (RuBisCO large subunit) were increased by at least 3-fold in the chloroplasts in the OE7 line. The ATP synthase complex subunits were significantly reduced in the OE7 line, including ATPA, ATPB, and ATPC (Supplemental Table S2). In the mitochondria, the protein abundances of the ATP synthase subunits alpha, beta and gamma were also lower in OE7 (Fig. 2B, Supplemental Table S3).

***LEF, but not the CEF, was enhanced in the OE lines.*** To investigate the impact of the altered thylakoid composition on the photosynthetic performance, the photosynthetic electron transport in 20-day-old leaves was assessed using chlorophyll fluorescence analysis (Fig. 3). Both AtPAP2 OE lines displayed an increased PSII quantum yield Y(II), PSII photochemical capacity (qP), and electron transport rate (ETR) compared to the WT (Fig. 3A), which suggests that the OE lines exhibited a higher LEF efficiency. No difference in terms of absorptivity was observed between the WT and OE line, implying that the altered photosynthesis parameters in OE line was not due to any changes in chlorophyll absorption capacity (Supplemental Fig. S2). An analysis of the P700 redox state showed that the PSI in the OE lines was highly oxidized under low light conditions ($25 - 125$ µmol photon $m^{-2}$ $s^{-1}$), suggesting that in the OE lines, the number of electrons that are captured from the PSI by downstream recipients (e.g., $NADP^+$) is higher than the number of electrons that is supplied by PSII to PSI (Fig. 3B). This phenomenon was not observed when the electron flow was larger at the higher light intensities. Although the ETR and P700 redox analysis suggested that the OE lines have a higher electron transfer capacity, the NAD(P)H dehydrogenase- and Fd- dependent CEF were not significantly changed in the OE lines according to the post-illumination chlorophyll fluorescence transient and ruptured chloroplast assay, respectively (Supplemental Fig. S3). In summary, the LEF, but not the CEF, was enhanced in the OE lines, and PSI played a key role in this enhancement by relaying more electrons to $NADP^+$ in the transfer chain.

***Photosynthesis rate was enhanced in the chloroplasts of the AtPAP2 OE lines.*** To study the
effect of the altered thylakoid composition on the photosynthesis performance, the
photosynthetic parameters were characterized in 20-day-old leaves by an analysis of gas
exchange. Both OE lines exhibited a higher light-saturated-carbon fixation rate under ambient
$CO_2$ concentration than the WT (Fig. 3C). This increased rate correlates with the higher
abundance of RbcL in the OE chloroplasts (Fig. 2A). Assays of the Calvin-Benson-Bassham
(CBB) cycle enzymes showed that the capacities of fructose bisphosphate aldolase and
glyceraldehyde-3-phosphate dehydrogenase (GAPDH) were increased by 30 - 40% in the
chloroplasts in the OE7 line (Table 1).

***The mitochondria in the AtPAP2 OE line had a higher capacity in dissipating reducing
equivalents.*** Reductant equivalents can be exported from the chloroplasts in the form of
malate or DHAP (Shameer et al., 2019), which can be used for sucrose synthesis in the
cytosol or converted to pyruvate and fed into the TCA cycle. Enzyme assays of the TCA
cycle suggested that the OE7 mitochondria had an increased capacity of mitochondrial
$NAD^+$-dependent malate dehydrogenase (mMDH) but not in the capacities of pyruvate
dehydrogenase, citrate synthase or aconitase (Table 1), whereas the capacities of chloroplast
$NADP^+$-dependent malate dehydrogenase, mMDH, 2-oxoglutarate dehydrogenase (2-OGDH),
and succinyl-CoA synthetase were increased by 1.26-fold, 1.15-fold, 2.86-fold and 10.3-fold
in the OE7 line, respectively. Our recent studies showed that glycine decarboxylase generates
a large amount of NADH in mitochondria during photorespiration, which exceeds the
NADH-dissipating capacity of the mitochondria. Consequently, the surplus NADH must be
exported to the cytosol through the mitochondrial malate-OAA shuttle (Lim et al., 2020). The
oxygraph studies showed that the electron-feeding capacities of complex II and internal
NADH dehydrogenase, but not of complex I or external NADH dehydrogenase, were
enhanced in the OE7 mitochondria (Table 2). If the OE line had a higher capacity in
dissipating NADH, less surplus reducing equivalents will be exported to the cytosol through
the malate-OAA shuttle.

***Higher mitochondrial activity is responsible for the higher cytosolic ATP in the OE line.*** Our
previous study showed that the leaf in the AtPAP2 OE lines contained a significantly higher
level of ATP and a higher ATP/NADPH ratio in both dark and light conditions (Liang et al.,
2015). Here, we introduced a FRET $MgATP^{2-}$ sensor (Imamura et al., 2009; Voon et al., 2018)

into the chloroplasts and the cytosol and a pH sensor (Schwarzländer et al., 2011) to the

mitochondrial matrix and compared the responses of the OE7 and WT lines (Voon et al., 2018).

Illumination led to an increased ATP concentration in the stroma (Fig. 4A) and alkalization of

the mitochondrial matrix in both lines (Fig. 4B). The extent of the alkalization of the

mitochondrial matrix was greater in the OE7 line than that in the WT, implying that under the

same light intensity, the reducing power harvested by the chloroplasts may lead to a stronger

proton translocation activity across the mitochondrial inner membrane (Fig. 4B). By contrast,

the rate of increase in the ATP concentration in the stroma is lower in the OE7 line than that in

the WT, which is likely due to a higher ATP consumption rate in the OE7 chloroplasts because

the $CO_2$ fixation rate (Fig. 3C) and the capacities of certain CBB cycle enzymes are higher in

the OE7 line (Table 1). We then examined the contribution of complex I, complex II and

complex V in the OE7 mitochondria to cytosolic ATP. These inhibitors lower the cytosolic ATP

in both lines (Fig. 4C). When complex I was inhibited by rotenone, the cytosolic ATP

concentration was significantly higher in the OE7 line, suggesting that non-complex I activities

in the OE7 mitochondria, such as complex II and internal NADH dehydrogenase, can

compensate more effectively than those in the WT (Table 2). This difference was not observed

when complex II was inhibited by TTFA. After one hour of incubation with oligomycin,

cytosolic ATP in WT dropped to an undetectable level, whereas a higher level of cytosolic ATP

was remained in OE7, suggesting that the intrinsic cytosolic ATP level was higher in OE7 than

WT (Fig. 4C).

***AtPAP2 selectively interacts with a number of chloroplast and mitochondrial proteins.*** A

yeast two-hybrid (Y2H) library screening was carried out to identify other

AtPAP2-interacting proteins. Forty nuclear-encoded AtPAP2-interacting proteins were

identified; of these proteins, 32 and 3 proteins have been experimentally verified in

chloroplasts and mitochondria, respectively (Supplemental Table S4). The other 5

AtPAP2-interacting proteins were predicted to be targeted to chloroplasts and/or

mitochondria. AtPAP2 is exposed to the cytosolic side of the outer membrane of both

organelles and its role in import of organellar proteins may account for the enrichment of

nuclear-encoded organellar proteins. A systematic Y2H assay was then carried out to examine

the interactions between AtPAP2 and various nucleus-encoded photosystem proteins

(Supplemental Fig. S4A). AtPAP2 could specifically interact with Fd1, Fd2, PsaE2 (but not

PsaE1), ferredoxin/thioredoxin reductase subunit A2 (FTRA2) (but not FTRA1), FTRB, and

photosynthetic NDH subcomplexes L1, L2, and L3 (PsbQ-like1, PsbQ-like2, and PsbQ-like3).

These interactions were verified by *in vivo* bimolecular fluorescence complementation (BiFC)

assays (Supplemental Fig. S4B and Supplemental Fig. S5). Hence, AtPAP2 may play a role in

the import of these precursor proteins, particularly the proteins at the acceptor side of PSI, into

chloroplasts.

**Discussion**

Maintenance of optimal photosynthetic efficiency requires the production and consumption of ATP and reductants in chloroplasts at appropriate ratios. Our recent study showed that the import of cytosolic ATP into mature chloroplasts of *Arabidopsis thaliana* is negligible and therefore could not supply additional ATP to the CBB cycle (Voon et al., 2018). Hence, the export of surplus reducing equivalents to extrachloroplast compartments is essential for balancing the ATP/NADPH ratio (Voon and Lim, 2019). Under illumination, the chloroplast is the producer of reducing equivalents and ATP, and the mitochondria consume the excess reducing equivalents to supply ATP to the cytosol (Gardeström and Igamberdiev, 2016; Voon et al., 2018; Shameer et al., 2019). The results presented here provide an example on how chloroplast function is highly dependent on the mitochondria. Illumination greatly elevates the amount of NADPH and the NADPH/ATP ratio in WT leaves (Liang et al., 2016), which represents a high reduction state in plant cells. Our previous studies showed that in the middle-of-day, the leaf sucrose and ATP contents are higher in the OE lines than those in the WT line in 20-day-old plants, whereas the leaf NADPH contents, $NADPH/NADP^+$ and NADPH/ATP ratios are lower in the OE lines than that in the WT line (Liang et al., 2015). Here, our data (Fig. 4, tables 1 and 2) indicated that the OE mitochondria may have a higher capacity to consume extra reducing equivalents to generate ATP (Fig. 5). This mechanism could account for the high ATP, high sucrose, but low NADPH content in the leaves of the OE lines in the middle-of-day (Sun et al., 2012; Liang et al., 2015). In addition, the RuBisCO content (Fig. 2A) and the capacities of the CBB cycle enzymes FBA and GAPDH (Table 1) were higher in the OE chloroplasts. These could lead to a higher rate of $CO_2$ fixation (Fig. 3C) and a rapid recycling of ADP, $NADP^+$ and RuBP in the stroma. Hence, a faster recycling of ribulose-1,5-bisphosphate (RuBP) and an enhanced output of dihydroxyacetone phosphate (DHAP) from the OE chloroplasts could be achieved. This, together with a higher consumption rate of reductants by the mitochondria (Table 2), might cause a lower reduction state in the stroma, thereby leading to an enhancement of the LEF in the OE lines. This hypothesis is supported by our data from the P700 redox analysis (Fig. 3B). At low light conditions (25 – 125 $\mu$mol photon $m^{-2}$ $s^{-1}$), in which the electron flow is smaller, PSI in the OE lines was highly oxidized, which was likely due to a high demand of electrons for the $NADP^+$ reduction or a higher PSI/PSII ratio (Fig. 2A and Supplemental Table S2). This oxidized state was relieved under higher light conditions (>125 $\mu$mol photon $m^{-2}$ $s^{-1}$) when the electron flow was larger

(Fig. 3B).

In pea (*Pisum sativum*), the PSI/PSII ratios were adjusted under different light conditions

as follows: sunlight-grown, 0.55; yellow light (preferentially excites PSII), 0.40; and red light

(preferentially excites PSI), 0.91 (Chow et al., 1990). Compared to plants with a lower

PSI/PSII ratio, plants with a higher PSI/PSII exhibit a higher rate of oxygen evolution under

illumination intensities that exceed 200 μmol photon $m^{-2}$ $s^{-1}$ (Chow et al., 1990). We also

observed a higher PSI/PSII ratio in the OE chloroplasts than that in the WT chloroplasts in the

2D-DIGE analysis (Fig. 2A and Supplemental Table S2). The OE lines exhibited a higher ETR

than the WT when the light intensity exceeded 200 μmol photon $m^{-2}$ $s^{-1}$ (Fig. 3). A higher

PSI/PSII ratio allows for a faster throughput of electrons from the photosynthetic ETC via the

PSI acceptor side and results in a lower possibility of an over-reduction in ETC (Tikkanen et al.,

2014). Interestingly, the OE lines had also less stacked thylakoid membranes compared to the

WT (Fig. 1). It is likely that the loose stacking increases the share of the grana margin domain

that is composed of all components of the photosynthetic electron transfer chain in relation to

the PSII-enriched grana stacks and PSI-enriched stroma lamellae. The importance of grana

stacking for photosynthetic efficiency is not understood. Based on the results presented here,

the loose stacking may be essential for the efficient interactions among the components of the

photosynthetic electron transfer chain.

Consistently with this hypothesis, the analysis of the P700 (PSI reaction centre) redox

state revealed that the P700 in the OE lines was relatively oxidized compared to that in the WT

when the electron supply from PSII was limited (25 – 125 μmol photon $m^{-2}$ $s^{-1}$) (Fig. 3B). Y2H

showed that AtPAP2 preferably interacts with the downstream components of PSI, including

PsaE2, Fd1, Fd2, FTRA2 and FTRB (Supplemental Fig. 4). Altogether, we hypothesize that

the LEF rate in the OE lines was increased because of the higher capacity of electron transfer in

PSI. LEF generates 1.28 mole ATP per mole of NADPH, but a minimum of 1.5 mole of ATP

per mole of NADPH is required for carbon fixation (Allen, 2003; Foyer et al., 2012). It is

generally believed that CEF pathways are needed to supplement ATP and modulate the

ATP/NADPH ratio to meet the demand of metabolism (Foyer et al., 2012; Ishikawa et al.,

2016). However, our studies showed that an enhancement of CEF is not required to support a

more robust CBB cycle in the OE line (Supplemental Fig. S3). A recent study in diatom

showed that the ATP generated from mitochondria can be transported to chloroplasts for carbon

fixation (Bailleul et al., 2015), and it was believed that this also occurs in higher plants.

However, our recent study showed that in the mesophyll of *Arabidopsis thaliana*, cytosolic

ATP does not enter mature chloroplast to support carbon fixation (Voon et al., 2018).

Altogether, because the NADPH/ATP demand for carbon dioxide fixation is approximately

0.67, and a 0.77 molecule of NADPH is generated from each ATP molecule in LEF (Allen,

2003; Foyer et al., 2012), one could expect that a surplus reducing power is generated from

LEF under illumination. Other metabolisms in chloroplasts, such as transcription and

translation, also consume ATP molecules; therefore, the surplus of reducing equivalents would

be more evident. The surplus of reducing equivalents must be exported and dissipated to

recycle $NADP^+$ for efficient photosynthesis.

The reducing power generated from LEF can be exported as DHAP and malate (Fig. 5)

(Noctor and Foyer, 2000; Scheibe, 2004). While modelling predicted that only 5% or less of

the reducing equivalents from LEF can be exported through the malate-OAA shuttle

(Fridlyand et al., 1998), the higher capacity of the $NAD^+$-MDH in the OE line might allow

for the export of more reducing equivalents from the mitochondria to the cytosol through the

malate-OAA shuttle. Illumination promotes the association between mitochondria and

chloroplasts and peroxisomes (Oikawa et al., 2015), and, therefore, the malate-OAA shuttle

can be a very efficient energy and reductant transfer pathway between the chloroplasts,

mitochondria and peroxisomes. Malate and OAA in the stroma and cytosol were estimated to

have concentrations ranging from $1 - 3$ mM and 0.025-0.098 mM, respectively (Heineke et

al., 1991). This concentration ratio is maintained by the large positive value of free energy

change, $\Delta_f G$, of malate dehydrogenation. This equilibrium can drive malate synthesis in the

chloroplasts upon illumination because the chloroplast $NADP^+$-dependent malate

dehydrogenase is activated by light (Scheibe, 1987; Zhao et al., 2018). The reducing power,

thus, can be readily channelled to the peroxisomes for hydroxypyruvate reduction through the

malate-OAA shuttle. As more reductants are consumed by the more active OE mitochondria,

less exportation of reductants from mitochondria is expected. Hence, more reductants from

the chloroplasts can be dissipated indirectly (Fig. 5). This is in line with our recent

observation that the increase in stromal NADPH during illumination disappeared when

photorespiration was absent (Abdel-Ghany, 2009; Lim et al., 2020).

In summary, the overexpression of AtPAP2 modulates the import of selected

nucleus-encoded proteins into chloroplasts (Zhang et al., 2016) and mitochondria (Law et al.,

2015), which leads to stronger sinks of reductants, including higher activities in the CBB

cycle and mitochondrial ETC, in both organelles. A higher output of reducing equivalents

from the LEF of chloroplasts (Fig. 3) and higher mitochondrial activities in utilizing the

reducing equivalents for ATP production, thus, contribute to the higher ATP content and ATP/NADPH ratio in the OE lines (Liang et al., 2015). The simultaneous activation of chloroplasts and mitochondria is required for the production of a surplus of ATP and sucrose; the transgenic lines that overexpress AtPAP2 solely in the chloroplasts grew similarly to the WT (unpublished data), and the transgenic lines that overexpress AtPAP2 solely in the mitochondria exhibited early senescence and a lower seed yield (Law et al., 2015). These results provide an example of how the efficient cooperation of chloroplasts and mitochondria can enhance ATP and sucrose production in leaf cells. While AtPAP2 can enhance yield and growth in *Arabidopsis thaliana*, it has negative impacts to plant survival. The OE lines are less resistant to drought and to *P. syringae* infection (Zhang et al., 2017). Therefore, while AtPAP2-like genes were evolved from green algae during evolution (Sun et al., 2012), it may not be fully used by higher plants to promote growth in natural environment, as survival is more important than high yield. It may be more 'useful' for unicellular algae as water is never a constraint, and algae need to control their growth rate and carbon level subject to the availability of P/N/Fe.

Over the past few decades, a controversy has emerged in the field regarding whether the shortfall of ATP for the CBB cycle is fulfilled by the CEF or the import of ATP from the cytosol. Combining the findings of our recent study (Voon et al., 2018) and this study, we concluded that the efficient carbon fixation in the OE line is not dependent on an enhanced CEF or the import of ATP from the cytosol to fulfil the shortfall of ATP generated from the LEF. Instead, an efficient carbon fixation is dependent on an enhanced LEF, enhanced capacities of CBB enzymes, the efficient export of surplus reducing equivalents from chloroplasts through the malate-OAA shuttle and a higher reductant-dissipating activity of mitochondria. The ability of mitochondria to dissipate reducing equivalents is important for relieving the built-up of surplus reducing power in the stroma, which, in excess, will limit the LEF due to an insufficient supply of $NADP^+$, the major electron acceptor of the LEF. Our recent studies showed that the reducing equivalents from photorespiration are the major fuel for mitochondria in *Arabidopsis thaliana* (Lim et al., 2020). The optimal use of reducing equivalents by mitochondria not only generates more ATP for sucrose synthesis, but also reduces the export of reducing equivalents from the mitochondria and therefore the overall redox status of the photosynthetic cells. This provides a good example to illustrate the relationship between chloroplasts and mitochondria in bioenergetics.

**Methodology**

**Plant growth conditions.** The WT *Arabidopsis thaliana* ecotype Columbia-o (WT), the *Arabidopsis thaliana* AtPAP2-overexpressing lines (OE7, OE21) and the *pap2* T-DNA insertion mutant (Salk_013567) (Sun et al., 2012) were used in this study. *Arabidopsis* seeds were plated for 10 days on Murashige and Skoog (MS) medium, which was supplemented with 2% (w/v) sucrose. Seedlings of similar sizes were transferred to soil under growth chamber conditions (120-150 µmol m$^{-2}$ s$^{-1}$, 16-h photoperiod, 22 °C).

**Transmission electron microscopy (TEM).** Plant materials (5th to 7th true leaves of 19-day-old plants at the beginning of the light period) were fixed with 2.5% (v/v) glutaraldehyde in 75 mM sodium cacodylate and 2 mM MgCl$_2$, pH 7.0, for 1 hour at 25 °C and post-fixed for 2 hours with 1% (w/v) osmium tetroxide in a fixative buffer at 25 °C. After two washing steps with distilled water, the pieces were dehydrated and then embedded in Spurr's low-viscosity resin. Ultrathin sections of 50–70 nm were cut with a diamond knife. The sections were post-stained with aqueous uranyl acetate/lead citrate. The TEM examination was performed under a Hitachi H-7650 transmission electron microscope with a charge-coupled device camera (Hitachi High-Technologies, Japan) operating at 80 kV.

**Determination of the chlorophyll content using an HPLC analysis.** Leaves (30-50 mg) were freeze-dried and extracted 3 times using 100% acetone (500 µl, 250 µl and 250 µl). The mixture was combined and centrifuged twice to eliminate the insoluble substances prior to the HPLC analysis. The pigment extract was separated and analysed (20 µL aliquots) on a Waters Spherisorb 5 µm ODS2 (4.6 × 250 mm) analytical column (Waters, USA) at room temperature. The fluids were eluted at a flow rate of 1.2 ml min$^{-1}$ with a linear gradient from 100% solvent A [acetonitrile/methanol/0.1 M TRIS–HCl (pH 8.0), 84:2:14, v/v/v] to 100% solvent B (methanol/ethyl acetate, 68:32, v/v) over a 15-min period, followed by 10 min of 100% solvent B. The tridimensional chromatogram was recorded from 250 to 700 nm. The chromatographic peaks were identified by comparing the retention times and spectra to known standards provided by Sigma (USA) and Wako (Japan). The pigments were finally quantified by integrating the peak areas and converting them to concentrations according to the comparisons. The amounts of lutein, β−carotene, violaxanthin, chlorophyll a (Ca) and chlorophyll b (Cb) were calculated as follows: lutein: $Y = 2.56 \times 10^8 X - 153665$, $R = 0.99577$, $p < 0.0001$; β-carotene: $Y = 6.57 \times 10^8 X - 61818$; violaxanthin: $Y = 2.68 \times 10^8 X +$

236.6; Chl a: $Y = 3.30 \times 10^7 X + 38832$; and Chl b: $Y = 7.31 \times 10^7 X + 29646$. Y = area, X =

mg/ml.

**2D Blue Native PAGE DIGE.** Leaves from 20-day-old soil-grown WT and OE7 plants at

the middle of the day (T = 8) were used as the starting materials. The chloroplast

(Lamkemeyer et al., 2006) and mitochondria isolation (Lee et al., 2008) were performed as

previously described. Cy Dye labelling of the chloroplast or mitochondrial proteins was

carried out as previously described with modifications (Heinemeyer et al., 2009). In total, 100

µg chloroplast/mitochondrial proteins were centrifuged for 10 min at 4 °C at 1000 g using an

Eppendorf centrifuge. The sedimented chloroplasts or mitochondria were re-suspended in a

10 µl solubilization buffer containing 30 mM HEPES, pH 7.4, 150 mM potassium acetate, 10%

[v/v] glycerol, supplemented with 2% [w/v] beta-dodecyl maltoside (chloroplasts) or 5% [w/v]

digitonin (mitochondria). The solubilized proteins were centrifuged for 20 min at 4 °C at full

speed to remove the insoluble material. The supernatant containing the solubilized protein

complexes was supplemented with 10 µl of the solubilization buffer containing 30 mM

HEPES, pH 10, 150 mM potassium acetate, 10% [v/v] glycerol to adjust the pH value of the

protein solution to approximately 8.5, which is a prerequisite for efficient labelling. Cy2, Cy3

and Cy5 NHS ester minimal dyes (Lumiprobe Corporation, USA) were reconstituted into 400

pmol/µl in DMF according to the manufacturer's instructions, and 2 µl of the Cy dye were

added to the solubilized chloroplast or mitochondrial proteins. The WT was labelled using a

Cy3 dye, and the OE was labelled with the Cy5 dye, while the internal standard (consisting of

a pooled sample comprising an equal amount of all samples in the experiment) was labelled

with Cy2. The labelling occurred for 30 min in the dark on ice and was stopped by adding 1

µl of a Lysine solution (10 mM). Finally, the sample was supplemented with 3 µl 5% (w/v)

Serva Blue G (750 mM aminocaproic acid, 5% [w/v] Coomassie 250 G) and directly

transferred into the slot of a blue native gel. The solubilized proteins were separated by

two-dimensional Blue native/SDS PAGE as described previously by Wittig et al. (2006). The

image acquisition was performed with a Typhoon Scanner, and the quantification of the Cy

Dye labelled proteins resolved by the 2D Blue native PAGE was carried out using the

SAMESPOTS software (Totallab, Tyne, England) based on the gel-images acquired from 3

biological replicates.

**Chlorophyll *a* fluorescence measurement.** The chlorophyll *a* fluorescence levels of the

*Arabidopsis* leaves (20 days old) were monitored with IMAGING-PAM M-Series Maxi

Version (WALZ, Germany). The plants were dark-adapted for at least 1 hour before the

measurements. After the maximum and initial fluorescence ($F_m$ and $F_0$, respectively) were

determined with a delay of 40 s, the plants were illuminated using the following light

intensities: 0, 81, 145, 186, 281, 335, 461, 701, and 926 µmol photon $m^{-2}$ $s^{-1}$. The duration of

the illumination of each light intensity was 3 min. A saturation pulse (800 ms, 2700 µmol

photon $m^{-2}$ $s^{-1}$) was applied at the end of the 3-min illumination. For each light intensity, the

photosynthesis parameters were calculated using the formulas: Y(II) was calculated as (Fm' –

F) /Fm' ; qP is calculated as (Fm' – F) /(Fm' – Fo'); NPQ was calculated as (Fm – Fm') /Fm' ;

ETR was calculated as Y(II) x light intensity x 0.5 x absorptivity. Fm' is the maximal

fluorescence of light-illuminated plant. F is the current fluorescence yield. Fo' was estimated

by Fo /(Fv /Fm + Fo /Fm') (Oxborough and Baker, 1997).

**Measurements of $CO_2$ assimilation.** The carbon dioxide assimilation in the 20-day-old plants

was measured using the portable photosynthesis system LI-6400XT (LiCor, USA) and whole

plant *Arabidopsis* chamber 6400-17. The plants were cultivated hydroponically in four- times

diluted Hoagland nutrient solution (plants were fixed in "Grodan"-mineral wool cubes) under

growth chamber conditions (irradiation 120-150 µmol $m^{-2}$ $s^{-1}$, 16/8 h day/night photoperiod,

temperature 22°C). The plants were transferred immediately before the measurement into

conical pots dedicated to the 6400-17 chamber, and the boundary between the substrate (roots)

and rosette was established using aluminium foil to minimize non-leaves transpiration. The

light response of $CO_2$ assimilation was measured under ambient $CO_2$ concentration 400 µmol

$mol^{-1}$ and PAR levels 2000, 1000, 500, 250, 120, 60, 30, 15 and 0 µmol photon $m^{-2}$ $s^{-1}$. Stable

temperature at 25 °C and the relative air humidity between 50 and 70% were kept in the

chamber during measurement.

**Measurements of the post-illumination chlorophyll fluorescence transient.** The modulated

chlorophyll fluorescence was measured using a PAM 101 fluorometer (H. Walz, Germany).

Before the minimal fluorescence ($F_0$) determination, the leaves were kept in darkness for at

least 1 hour. The $F_0$ was induced by a red-modulated measuring light with a photon flux density

(PFD) of approx. 0.2 µmol photon $m^{-2}$ $s^{-1}$. The post-illumination Chl fluorescence increase is

attributed to the back flow of electrons to the PQ pool from NADPH in the stroma via the cyclic

electron flow around photosystem I (CEF-PSI), which depends on the NDH complex (Gotoh et

al., 2010).

**Ruptured chloroplast assay**. The cyclic electron flow was measured in ruptured chloroplasts

through an Fd-dependent plastoquinone reduction as previously described (Endo et al., 1998;

Munekage et al., 2002). Briefly, the isolated chloroplasts (10 µg Chl) were osmotically

ruptured in the analysis buffer (7 mM $MgCl_2$, 1 mM $MnCl_2$, 2 mM EDTA, 30 mM KCl, 0.25

mM $KH_2PO_4$, and 50 mM HEPES, pH 7.6) for 10 min prior to the analysis. One min after the

initiation of the measurement, NADPH (Roche, Basel, Switzerland) was added to a final

concentration of 0.25 mM. Maize ferredoxin (Sigma Aldrich, St. Louis, MO, USA) was added

after an additional 30 s.

**Enzyme assays of the Calvin-Benson-Bassham cycle and TCA cycle.** The chloroplast

proteins were used to assay the fructose 1,6-bisphosphate aldolase (Schaeffer et al., 1997) and

$NADP^+$-glyceraldehyde-3-phosphate dehydrogenase activities. Enzyme assays of the TCA

cycle were performed as described previously (Huang et al., 2015). The activity of the

mitochondrial proteins (1-25 µg protein) were assayed spectrophotometrically using a

Multiskan™ GO Microplate Spectrophotometer (Thermo Scientific, USA).

**Measurements of oxygen consumption.** A Clark-type oxygen electrode (Hansatech

Instrument, UK) was used to measure oxygen consumption. The respiration rate of isolated

mitochondria was measured at 25 °C in a chamber containing 1 ml of the respiration buffer

(0.3 M Suc, 5 mM $KH_2PO_4$, 10 mM TES, 10 mM NaCl, 2 mM $MgSO_4$, and 0.1% [w/v] BSA,

pH 6.8 or pH 7.2). The capacity of complex I, internal and external NADH dehydrogenases,

and complex II were measured in the presence of deamino-NADH (1 mM), malate (10 mM),

glutamate (10 mM), and NADH (1 mM) with or without ADP (100 µM) and/or rotenone (5

µM) as previously described (Meyer et al., 2009; Jacoby et al., 2015).

**FRET-based ATP sensor and pH-dependent fluorescent probe.** In the ATP sensor AT1.03,

an ATP-binding epsilon-subunit of *Bacillus subtilis* $F_oF_1$-ATP synthase was fused between

mseCFP and mVenus (Imamura et al., 2009). The binding of ATP changes the distance between

mseCFP and mVenus and modulates the FRET signal (Imamura et al., 2009). The use of this

sensor in plant systems was recently verified, and the chlorophyll fluorescence does not

interfere with the FRET ratio in chloroplasts (Voon et al., 2018). To exclude the intensity

contributed from sources other than the FRET signal, correction coefficients for bleed-through

and cross-excitation were determined separately in different plant compartments (Broussard et

al., 2013). Sensitized FRET was used in all subsequent measurements involved AT1.03. To

estimate the ATP concentration *in vivo*, sensitized FRET/ CFP ratios of purified recombinant

AT1.03 sensor (6 μM) were calibrated with increasing ATP concentrations (0 – 6 mM) as

previously described (Voon et al., 2018). A pH-dependent fluorescent probe with circularly

permuted yellow fluorescent protein (cpYFP) was introduced into the matrix of the

mitochondria (Nagai et al., 2001). cpYFP has a bimodal absorption spectrum with two peak

maxima, 395 and 475 nm, and a single emission peak at 528 nm (Nagai et al., 2001). The

absorption intensity of these two spectra is dependent on whether the chromophore of cpYFP is

protonated or deprotonated. Protonated cpYFP displays a higher absorption at 395 nm, which

emits a higher signal at 528 nm if excited at 395 nm. cpYFP was shown to response specifically

to the pH value (Schwarzländer et al., 2011).

**Yeast two-hybrid assay.** A normalized yeast two-hybrid (Y2H) library was prepared from

mRNAs that were isolated from eleven *Arabidopsis* tissues and was employed for the

screening (Clontech Laboratories, Japan). The coding sequence of the mature bait protein

AtPAP2 (25-613 a.a.), which lacks its signal peptide and C-terminal transmembrane motif, was

fused to the C-terminus of the GAL4 DNA-binding domain (BD) in the pGBKT7 vector. The

GAL4-based Y2H library screening was performed using the Matchmaker Gold Yeast

Two-Hybrid System following the manufacturer's instructions (Clontech Laboratories, Japan).

The mated culture was concentrated by centrifugation and spread onto agar plates (90 mm)

containing triple dropout medium without Leu, Trp and His (TDO). Putative positive colonies

(> 2 mm) were cultured on TDO liquid medium, and pGADT7 plasmids were extracted from

the yeast cells for the *E. coli* transformation. The plasmids (pGADT7-prey) were re-extracted

and sequenced. To identify the interactions between AtPAP2 and various components of the

photosynthetic apparatus, the coding sequences of prey proteins were amplified from

*Arabidopsis* leaf cDNA by Platinum[TM] *Pfx* polymerase (Thermo Scientific, USA) and fused

downstream of the DNA-activating domain (AD) in the pGADT7 vector. The Y2HGold yeast

cells were co-transformed with both the bait and prey protein constructs using the lithium

acetate method and plated onto double dropout without Leu and Trp (DDO), TDO, and

quadruple dropout without Ade, His, Leu and Trp (QDO) agar plates (Clontech Laboratories,

Japan).

**Bimolecular fluorescence complementation (BiFC)**. The full-length coding sequence of the bait protein AtPAP2 was fused to the *C*-terminus of YFP[C] in the pSPYCE vector, while the coding sequences of the photosynthetic apparatus subunits were fused to the N-terminus of the pSPYNE containing YFP[N] (Kerppola, 2006). Both the prey and bait plasmids were transformed into the *Agrobacterium tumefaciens* strain GV3101, and the bacteria were infiltrated into the epidermal cell layers of tobacco leaves as previously described (Schweiger and Schwenkert, 2014). The transfected regions of the leaves were used for fluorescence detection after a 48-hour incubation in the dark under an LSM710 confocal laser scanning microscope (Zeiss). The primers that were used in this study are listed in Supplemental Table S5.

**Acknowledgements**

This project was supported by the Seed Funding Program for Basic Research (201311159043) of the University of Hong Kong, the General Research Fund (772012M) and the Area of Excellence Scheme (AoE/M-403/16) of the Hong Kong Research Grants Council, the Innovation and Technology Fund (Funding Support to State Key Laboratories in Hong Kong) of the HKSAR, China. Access to LI-6400 was supported by the Czech research infrastructure for systems biology C4SYS (project no LM2015055). We thank Prof. Wah Soon Chow and Dr. Maurice Cheung for their comments on this manuscript. We thank Dr. Huang Shaobai and Dr. Richard Jacoby for training of YL in enzyme assays, oxygraph and 2D-DIGE. We thank Prof. Hans-Peter Braun and Dr. Holger Eubel for technical advice in the 2D BN/SDS-PAGE DIGE analysis.

**Authors' contributions**

CPV, YL, and BLL wrote the manuscript with PG's input. CPV produced some of the transgenic sensor lines, performed the fluorescent sensor measurements, carried out the Maxi-Pam measurements and 2D BN-PAGE of the chloroplasts. YL carried out the 2D BN-PAGE of the mitochondria, the enzyme assays with WC, and the oxygraph assays with ZX. XG screened the yeast two-hybrid library and carried out the BiFC. RZ and FS produced the overexpression lines. FS measured the chlorophyll content. ML, MP and DL carried out the chloroplast rupture assay, MTik carried out the P700 assay, and MTsu carried out the post-illumination chlorophyll fluorescence. JK and MH carried out the $CO_2$ exchange

555    measurements. YC and LJ carried out the TEM analysis. BLL coordinated this study. All
556    authors read and approved the manuscript.

557    **Competing financial interests.**

558    AtPAP2 is the subject of US patent number 9,476,058.

**References**

**Abdel-Ghany SE** (2009) Contribution of plastocyanin isoforms to photosynthesis
and copper homeostasis in Arabidopsis thaliana grown at different copper regimes.
Planta **229:** 767-779

**Allen JF** (2003) Cyclic, pseudocyclic and noncyclic photophosphorylation: new
links in the chain. Trends Plant Sci **8:** 15-19

**Armbruster U, Labs M, Pribil M, Viola S, Xu W, Scharfenberg M, Hertle AP,**
**Rojahn U, Jensen PE, Rappaport F, Joliot P, Dormann P, Wanner G, Leister D**
(2013) Arabidopsis CURVATURE THYLAKOID1 proteins modify thylakoid
architecture by inducing membrane curvature. Plant Cell **25:** 2661-2678

**Bailleul B, Berne N, Murik O, Petroutsos D, Prihoda J, Tanaka A, Villanova**
**V, Bligny R, Flori S, Falconet D, Krieger-Liszkay A, Santabarbara S, Rappaport F,**
**Joliot P, Tirichine L, Falkowski PG, Cardol P, Bowler C, Finazzi G** (2015)
Energetic coupling between plastids and mitochondria drives $CO_2$ assimilation in
diatoms. Nature **524:** 366-369

**Broussard JA, Rappaz B, Webb DJ, Brown CM** (2013) Fluorescence
resonance energy transfer microscopy as demonstrated by measuring the activation of
the serine/threonine kinase Akt. Nat Protoc **8:** 265-281

**Chow WS, Melis A, Anderson JM** (1990) Adjustments of Photosystem
Stoichiometry in Chloroplasts Improve the Quantum Efficiency of Photosynthesis.
Proc Natl Acad Sci U S A **87:** 7502-7506

**Dutilleul C, Driscoll S, Cornic G, De Paepe R, Foyer CH, Noctor G** (2003)
Functional mitochondrial complex I is required by tobacco leaves for optimal
photosynthetic performance in photorespiratory conditions and during transients. Plant
Physiol **131:** 264-275

**Endo T, Shikanai T, Sato F, Asada K** (1998) NAD(P)H
dehydrogenase-dependent, antimycin A-sensitive electron donation to plastoquinone in
tobacco chloroplasts. Plant and Cell Physiology **39:** 1226-1231

**Foyer CH, Neukermans J, Queval G, Noctor G, Harbinson J** (2012)
Photosynthetic control of electron transport and the regulation of gene expression. J
Exp Bot **63:** 1637-1661

**Fridlyand LE, Backhausen JE, Scheibe R** (1998) Flux control of the malate

valve in leaf cells. Archives of Biochemistry and Biophysics **349:** 290-298

  **Gardeström P, Igamberdiev AU** (2016) The origin of cytosolic ATP in photosynthetic cells. Physiol. Plant. **157:** 367-379

  **Gerhardt R, Stitt M, Heldt HW** (1987) Subcellular Metabolite Levels in Spinach Leaves : Regulation of Sucrose Synthesis during Diurnal Alterations in Photosynthetic Partitioning. Plant Physiol **83:** 399-407

  **Gotoh E, Matsumoto M, Ogawa K, Kobayashi Y, Tsuyama M** (2010) A qualitative analysis of the regulation of cyclic electron flow around photosystem I from the post-illumination chlorophyll fluorescence transient in Arabidopsis: a new platform for the in vivo investigation of the chloroplast redox state. Photosynth Res **103:** 111-123

  **Heineke D, Riens B, Grosse H, Hoferichter P, Peter U, Flugge UI, Heldt HW** (1991) Redox transfer across the inner chloroplast envelope membrane. Plant Physiol **95:** 1131-1137

  **Heinemeyer J, Scheibe B, Schmitz UK, Braun HP** (2009) Blue native DIGE as a tool for comparative analyses of protein complexes. J Proteomics **72:** 539-544

  **Huang S, Lee CP, Millar AH** (2015) Activity assay for plant mitochondrial enzymes. Methods Mol Biol **1305:** 139-149

  **Imamura H, Nhat KP, Togawa H, Saito K, Iino R, Kato-Yamada Y, Nagai T, Noji H** (2009) Visualization of ATP levels inside single living cells with fluorescence resonance energy transfer-based genetically encoded indicators. Proc Natl Acad Sci U S A **106:** 15651-15656

  **Ishikawa N, Takabayashi A, Sato F, Endo T** (2016) Accumulation of the components of cyclic electron flow around photosystem I in C4 plants, with respect to the requirements for ATP. Photosynth Res **129:** 261-277

  **Jacoby RP, Millar AH, Taylor NL** (2015) Assessment of Respiration in Isolated Plant Mitochondria Using Clark-Type Electrodes. *In* J Whelan, MW Murcha, eds, Plant Mitochondria: Methods and Protocols. Springer New York, New York, NY, pp 165-185

  **Kerppola TK** (2006) Design and implementation of bimolecular fluorescence complementation (BiFC) assays for the visualization of protein interactions in living cells. Nature Protocols **1:** 1278-1286

  **Lamkemeyer P, Laxa M, Collin V, Li W, Finkemeier I, Schottler MA, Holtkamp V, Tognetti VB, Issakidis-Bourguet E, Kandlbinder A, Weis E, Miginiac-Maslow M, Dietz KJ** (2006) Peroxiredoxin Q of *Arabidopsis thaliana* is attached to the thylakoids and functions in context of photosynthesis. Plant Journal **45:** 968-981

  **Law YS, Zhang R, Guan X, Cheng S, Sun F, Duncan O, Murcha M, Whelan J, Lim BL** (2015) Phosphorylation and Dephosphorylation of the Presequence of

pMORF3 During Import into Mitochondria from Arabidopsis thaliana. Plant Physiol **169:** 1-12

**Lee CP, Eubel H, O'Toole N, Millar AH** (2008) Heterogeneity of the mitochondrial proteome for photosynthetic and non-photosynthetic Arabidopsis metabolism. Mol Cell Proteomics **7:** 1297-1316

**Liang C, Cheng S, Zhang Y, Sun Y, Fernie AR, Kang K, Panagiotou G, Lo C, Lim BL** (2016) Transcriptomic, proteomic and metabolic changes in Arabidopsis thaliana leaves after the onset of illumination. BMC Plant Biol **16:** 43

**Liang C, Zhang Y, Cheng S, Osorio S, Sun Y, Fernie AR, Cheung CYM, Lim BL** (2015) Impacts of high ATP supply from chloroplasts and mitochondria on the leaf metabolism of *Arabidopsis thaliana*. Front. Plant Sci. **6:** 922

**Lim SL, Voon CP, Guan X, Yang Y, Gardestrom P, Lim BL** (2020) *In planta* study of photosynthesis and photorespiration using NADPH and NADH/NAD$^+$ fluorescent protein sensors. Nature communications **11**

**Meyer EH, Tomaz T, Carroll AJ, Estavillo G, Delannoy E, Tanz SK, Small ID, Pogson BJ, Millar AH** (2009) Remodeled respiration in ndufs4 with low phosphorylation efficiency suppresses Arabidopsis germination and growth and alters control of metabolism at night. Plant Physiol **151:** 603-619

**Munekage Y, Hojo M, Meurer J, Endo T, Tasaka M, Shikanai T** (2002) PGR5 is involved in cyclic electron flow around photosystem I and is essential for photoprotection in Arabidopsis. Cell **110:** 361-371

**Nagai T, Sawano A, Park ES, Miyawaki A** (2001) Circularly permuted green fluorescent proteins engineered to sense Ca$^{2+}$. Proc Natl Acad Sci U S A **98:** 3197-3202

**Noctor G, Foyer CH** (2000) Homeostasis of adenylate status during photosynthesis in a fluctuating environment. J Exp Bot **51:** 347-356

**Noguchi K, Yoshida K** (2008) Interaction between photosynthesis and respiration in illuminated leaves. Mitochondrion **8:** 87-99

**Oikawa K, Matsunaga S, Mano S, Kondo M, Yamada K, Hayashi M, Kagawa T, Kadota A, Sakamoto W, Higashi S, Watanabe M, Mitsui T, Shigemasa A, Iino T, Hosokawa Y, Nishimura M** (2015) Physical interaction between peroxisomes and chloroplasts elucidated by in situ laser analysis. Nature Plants **1:** 15035

**Oxborough K, Baker NR** (1997) Resolving chlorophyll a fluorescence images of photosynthetic efficiency into photochemical and non-photochemical components – calculation of qP and Fv-/Fm-; without measuring Fo. Photosynthesis Research **54:** 135-142

**Sato R, Kawashima R, Trinh MDL, Nakano M, Nagai T, Masuda S** (2019) Significance of PGR5-dependent cyclic electron flow for optimizing the rate of ATP

synthesis and consumption in Arabidopsis chloroplasts. Photosynth Res **139:** 359-365

**Schaeffer GW, Sharpe FT, Sicher RC** (1997) Fructose 1,6-bisphosphate aldolase activity in leaves of a rice mutant selected for enhanced lysine. Phytochemistry **46:** 1335-1338

**Scheibe R** (1987) Nadp+-Malate Dehydrogenase in C-3-Plants - Regulation and Role of a Light-Activated Enzyme. Physiol Plantarum **71:** 393-400

**Scheibe R** (2004) Malate valves to balance cellular energy supply. Physiol. Plant. **120:** 21-26

**Scheibe R, Backhausen JE, Emmerlich V, Holtgrefe S** (2005) Strategies to maintain redox homeostasis during photosynthesis under changing conditions. J Exp Bot **56:** 1481-1489

**Schwarzländer M, Logan DC, Fricker MD, Sweetlove LJ** (2011) The circularly permuted yellow fluorescent protein cpYFP that has been used as a superoxide probe is highly responsive to pH but not superoxide in mitochondria: implications for the existence of superoxide 'flashes'. Biochem J **437:** 381-387

**Schweiger R, Schwenkert S** (2014) Protein-protein Interactions visualized by bimolecular fluorescence complementation in tobacco protoplasts and leaves. JoVe-Journal of Visualized Experiments **85:** e51327

**Selinski J, Scheibe R** (2019) Malate valves: old shuttles with new perspectives. Plant Biol (Stuttg) **21:** 21-30

**Shameer S, Ratcliffe RG, Sweetlove LJ** (2019) Leaf energy balance requires mitochondrial respiration and export of chloroplast NADPH in the light. Plant Physiology **180:** 1947-1961

**Sun F, Carrie C, Law S, Murcha MW, Zhang R, Law YS, Suen PK, Whelan J, Lim BL** (2012) AtPAP2 is a tail-anchored protein in the outer membrane of chloroplasts and mitochondria. Plant Signal Behav **7:** 927-932

**Sun F, Liang C, Whelan J, Yang J, Zhang P, Lim BL** (2013) Global transcriptome analysis of AtPAP2--overexpressing *Arabidopsis thaliana* with elevated ATP. BMC Genomics **14:** 752

**Sun F, Suen PK, Zhang Y, Liang C, Carrie C, Whelan J, Ward JL, Hawkins ND, Jiang L, Lim BL** (2012) A dual-targeted purple acid phosphatase in *Arabidopsis thaliana* moderates carbon metabolism and its overexpression leads to faster plant growth and higher seed yield. New Phytol **194:** 206-219

**Tikkanen M, Mekala NR, Aro EM** (2014) Photosystem II photoinhibition-repair cycle protects Photosystem I from irreversible damage. Biochimica Et Biophysica Acta-Bioenergetics **1837:** 210-215

**Voon CP, Guan X, Sun Y, Sahu A, Chan MN, Gardeström P, Wagner S, Fuchs P, Nietzel T, Versaw WK, Schwarzländer M, Lim BL** (2018) ATP

compartmentation in plastids and cytosol of *Arabidopsis thaliana* revealed by
fluorescent protein sensing. Proc Natl Acad Sci U S A **115:** 10778-10787

**Voon CP, Lim BL** (2019) ATP translocation and chloroplast biology National
Science Review **6:** 1073-1076

**Wittig I, Braun HP, Schagger H** (2006) Blue native PAGE. Nat Protoc **1:**
418-428

**Zhang R, Guan X, Law YS, Sun F, Chen S, Wong KB, Lim BL** (2016) AtPAP2
modulates the import of the small subunit of Rubisco into chloroplasts. Plant Signal
Behav **11:** e1239687

**Zhang R, Qi H, Sun Y, Xiao S, Lim BL** (2017) Transgenic Arabidopsis thaliana
containing increased levels of ATP and sucrose is more susceptible to *Pseudomonas*
*syringae*. PLoS One **12:** e0171040

**Zhao YN, Luo LL, Xu JS, Xin PY, Guo HY, Wu J, Bai L, Wang GD, Chu JF,**
**Zuo JR, Yu H, Huang X, Li JY** (2018) Malate transported from chloroplast to
mitochondrion triggers production of ROS and PCD in *Arabidopsis thaliana*. Cell
Research **28:** 448-461

| Mean ± SEM | Diameter (nm) | Height (nm) |
|---|---|---|
| WT | $432 \pm 19$ [a] | $125 \pm 8$ [a] |
| *pap2* | $444 \pm 11$ [a] | $102 \pm 5$ [b] |
| OE7 | $431 \pm 16$ [a] | $78 \pm 4$ [c] |
| OE21 | $430 \pm 12$ [a] | $79 \pm 3$ [c] |

**Figure 1. Comparison of the thylakoid architecture between the WT and OE plants.**
TEM micrograph of ultrathin sections of 20-day-old leaves from the WT, OE (OE7 and OE21) and AtPAP2 knock-out (*pap2*) lines. Average value (n>30) of the diameter (nm) and height (nm) of the thylakoid in the WT and OE lines were shown. Values marked by different letters in the same column are significantly different ($p < 0.05$) by Student's t-test. The diameter (nm) and height (nm) of WT chloroplasts isolated from 28-day-old leaves in another study (15) were 448 $\pm$ 16 nm and 113 $\pm$ 5 nm, respectively.

**Figure 2. Comparative analysis of the chloroplast and mitochondrial proteomes of WT**

**and OE7 plants**. **A**. Chloroplasts; **B**. Mitochondria. Proteins from the WT fraction were

pre-labelled with Cy3; proteins from the OE fraction were pre-labelled with Cy5. Combined

protein fractions were separated by 2D Blue-native/SDS-PAGE, and the protein visualization

was carried out by laser scanning at the respective wavelengths using the Typhoon laser

scanner. On the resulting overlay image, Cy3 is represented by red, and Cy5 is represented by

green. Proteins with a reduced abundance in the OE7 line are shown in red; proteins with an

increased abundance in the OE7 line are shown in green; and proteins of equal abundance in

the two compared fractions are shown in yellow. Protein spots were extracted and identified by

an MS/MS analysis. Identified proteins with the highest unused score and at least 2 unique

peptides (95%) were labelled with the corresponding spot ID (Supplemental Tables S2 & S3).

The representative gel images of 3 biological replicates are presented.

741
**Figure 3.** *In vivo* **analysis of the electron transport activity and $CO_2$ assimilation rate. A**.
Light intensity-dependent NPQ, qP, ETR and Y(II). The 3-week-old plants were
dark-acclimated for 1 hour before the measurement. Data are presented as the mean ± SE (n >
10 per line). **B**. P700 Oxidation. Data are presented as the mean ± SD (n = 3 per line). Light
intensity was increased in a step-wise manner as stated in the graph. **C**. Light response of $CO_2$
assimilation rates (An) under ambient $CO_2$ concentration. Data are presented as the mean ± SE
(n = 5 per line). The asterisks indicate significant differences between the WT and both OE
lines by one-way ANOVA with post-hoc Tukey HSD test (p < 0.05).

**Figure 4. Assessing organelle activities using fluorescent sensors. A**. Change in apparent

MgATP$^{2-}$ concentration in the chloroplast stroma. 10-day-old WT and OE7 cotyledon

expressing TKTP-AT1.03 were illuminated (296 μmol photon m$^{-2}$ s$^{-1}$) for 3 mins, followed by

5 mins in the dark. **B**. Changes in the mitochondrial matrix pH sensor ratio (Ex 488/ Ex 405).

10-day-old WT and OE7 cotyledon expressing matrix-localized pH sensor, mt-cpYFP were

illuminated (296 μmol photon m$^{-2}$ s$^{-1}$) for 30 s, followed by 1 min in the dark. Black and white

bars indicate the dark and illumination periods, respectively. Average value of 6 independent

measurements and SE are shown. The asterisks indicate significant differences between the

WT and the OE line by one-way ANOVA with post-hoc Tukey HSD test ($p < 0.05$). Graphs

were normalized to the initial point. **C**. Effects of the inhibitors on cytosolic ATP concentration.

Inhibitors of complex I (50 μM rotenone), complex II (100 μM TTFA) and complex V (10μM

oligomycin) were applied to 10-day-old WT and OE7 seedling expressing C-AT1.03 seedlings

by vacuum infiltration for 5 min. Before the measurements, the seedlings were incubated in the

dark for 1 hour. Average value of 3 replicates is presented. Error bars are the standard error of

the average MgATP$^{2-}$ concentration. Groups with significant difference by one-way ANOVA

with post-hoc Tukey HSD test ($p < 0.05$) are indicated by different letters. The dashed line

indicates the maximal MgATP$^{2-}$ concentration (1.4 mM) that can be reported by AT1.03 (Voon

et al., 2018). The groups marked by asterisks are significantly different from the groups marked

by letters but whether they have intergroup differences are not known.

**Figure 5. A model on how efficient collaboration between chloroplasts and mitochondria promote ATP and sucrose production.** ① and ② AtPAP2 on the outer membranes of chloroplasts and mitochondria promotes the import of certain proteins into these two organelles via the Toc or the Tom complexes. ③ Higher PSI/PSII ratio and higher LEF generate more NADPH and ATP at a ratio of 0.78, which are consumed at a ratio of 0.67 by the enhanced CBB enzymes in the OE chloroplasts for $CO_2$ fixation. The surplus reducing equivalents are exported from the chloroplasts via the malate/OAA shuttle to recycle $NADP^+$ as the electron acceptors of the LEF. ④ Higher reductant-dissipating activities of OE mitochondria reduce the needs for mitochondria to export reductants from photorespiration in the form of malate. ⑤ OE chloroplasts with enhanced rate of carbon fixation export more carbon skeletons to the cytosol. ⑥ OE mitochondria with higher reductant-dissipating activities generate more ATP through the respiratory electron transfer (RET) chain. ⑦ Higher ATP production in OE mitochondria and higher output of carbon skeletons from chloroplasts enhance sucrose synthesis in the cytosol. Red and blue lines indicate upregulated and downregulated pathways/metabolites in the OE lines, respectively.

788    **Table 1. Enzyme activities of the CBB cycle and the Krebs cycle**

| Enzyme | Source | WT | OE | Change |
|---|---|---|---|---|
| Fructose bisphosphate aldolase | Chloroplast | $92 \pm 8$ | $123 \pm 14$** | 1.34X |
| NADP$^+$-GAPDH | Chloroplast | $97 \pm 11$ | $138 \pm 4$** | 1.42X |
| NADP$^+$-MDH | Chloroplast | $328 \pm 20$ | $413 \pm 40$** | 1.26X |
| NADP$^+$-MDH | Leaf | $26 \pm 4$ | $28 \pm 4$ | N.S. |
| NAD$^+$-MDH | Leaf | $1525 \pm 25$ | $1439 \pm 39$ | N.S. |
| NAD$^+$-MDH | Mitochondria | $4657 \pm 206$ | $5354 \pm 46$** | 1.15X |
| NAD$^+$-malic enzyme, ME | Mitochondria | $40 \pm 3$ | $42 \pm 3$ | N.S. |
| Pyruvate dehydrogenase, PDC | Mitochondria | $45 \pm 6$ | $48 \pm 4$ | N.S. |
| Citrate synthase, CS | Mitochondria | $84 \pm 12$ | $90 \pm 12$ | N.S. |
| Aconitase, ACN | Mitochondria | $314 \pm 18$ | $322 \pm 18$ | N.S. |
| NAD$^+$-ICDH | Mitochondria | $43 \pm 3$ | $63 \pm 8$** | 1.47X |
| NADP$^+$-ICDH | Leaf | $18.5 \pm 2.2$ | $6.7 \pm 0.5$** | 0.36X |
| 2-oxoglutarate dehydrogenase, 2OGDH | Mitochondria | $1.4 \pm 0.6$ | $4.0 \pm 0.4$** | 2.86X |
| Succinate dehydrogenase, SDH | Mitochondria | $7.6 \pm 0.3$ | $10.3 \pm 1.0$** | 1.35X |
| Succinyl-CoA synthetase | Mitochondria | $29 \pm 1.4$ | $300 \pm 63$** | 10.3X |
| Fumarase, FUM | Mitochondria | $334 \pm 55$ | $209 \pm 34$** | 0.62X |

789    *GAPDH*, glyceraldehyde-3-phosphate dehydrogenase; *ICDH*, isocitrate dehydrogenase; *MDH*, malate

790    dehydrogenase. All enzyme unit is presented as nmol/min/mg protein.

791    Independent sample t-test was carried out. Significant differences between WT and OE7 are showed by asterisks.

792    * $p < 0.05$, ** $p < 0.01$, n=4. N.S.: Not significant.

794  **Table 2 Respiration rates of different complexes of WT and OE7 mitochondria.**

| Enzyme | Substrate | ADP | Rotenone | WT | OE7 | Change |
|--------|-----------|-----|----------|-----|-----|--------|
| Complex I | deamino-NADH | - | - | 30 ± 6 | 33 ± 5 | ns |
| | | - | + | 8 ± 3 | 8 ± 5 | ns |
| Complex II | Succinate | - | - | 49 ± 3 | 48 ± 23 | ns |
| | | + | - | 75 ± 4 | 92 ± 10 * | 1.23X |
| NdEx | NADH | - | - | 45 ± 4 | 45 ± 12 | ns |
| | | + | - | 65 ± 4 | 68 ± 4 | ns |
| NdIn | Malate + Glutamate | - | + | 39 ± 4 | 46 ± 6 | ns |
| | | + | + | 64 ± 8 | 85 ± 4 * | 1.32X |

795  Oxygen consumption rates of complex I (rotenone-sensitive deamino-NADH oxidation), complex II, external

796  NADH dehydrogenase, internal NADH dehydrogenase (rotenone-insensitive NADH oxidation) were expressed

797  as nmol $min^{-1}$ $mg^{-1}$ protein.

798  * indicates statistically significant difference (p <0.05, n≥3) and ns indicates not significant difference between

799  WT and OE7 by Student's t test.

# Parsed Citations

Abdel-Ghany SE (2009) Contribution of plastocyanin isoforms to photosynthesis and copper homeostasis in Arabidopsis thaliana grown at different copper regimes. Planta 229: 767-779

Google Scholar: Author Only Title Only Author and Title

Allen JF (2003) Cyclic, pseudocyclic and noncyclic photophosphorylation: new links in the chain. Trends Plant Sci 8: 15-19

Google Scholar: Author Only Title Only Author and Title

Armbruster U, Labs M, Pribil M, Viola S, Xu W, Scharfenberg M, Hertle AP, Rojahn U, Jensen PE, Rappaport F, Joliot P, Dormann P, Wanner G, Leister D (2013) Arabidopsis CURVATURE THYLAKOID1 proteins modify thylakoid architecture by inducing membrane curvature. Plant Cell 25: 2661-2678

Google Scholar: Author Only Title Only Author and Title

Bailleul B, Berne N, Murik O, Petroutsos D, Prihoda J, Tanaka A, Villanova V, Bligny R, Flori S, Falconet D, Krieger-Liszkay A, Santabarbara S, Rappaport F, Joliot P, Tirichine L, Falkowski PG, Cardol P, Bowler C, Finazzi G (2015) Energetic coupling between plastids and mitochondria drives CO2 assimilation in diatoms. Nature 524: 366-369

Google Scholar: Author Only Title Only Author and Title

Broussard JA, Rappaz B, Webb DJ, Brown CM (2013) Fluorescence resonance energy transfer microscopy as demonstrated by measuring the activation of the serine/threonine kinase Akt. Nat Protoc 8: 265-281

Google Scholar: Author Only Title Only Author and Title

Chow WS, Melis A, Anderson JM (1990) Adjustments of Photosystem Stoichiometry in Chloroplasts Improve the Quantum Efficiency of Photosynthesis. Proc Natl Acad Sci U S A 87: 7502-7506

Google Scholar: Author Only Title Only Author and Title

Dutilleul C, Driscoll S, Cornic G, De Paepe R, Foyer CH, Noctor G (2003) Functional mitochondrial complex I is required by tobacco leaves for optimal photosynthetic performance in photorespiratory conditions and during transients. Plant Physiol 131: 264-275

Google Scholar: Author Only Title Only Author and Title

Endo T, Shikanai T, Sato F, Asada K (1998) NAD(P)H dehydrogenase-dependent, antimycin A-sensitive electron donation to plastoquinone in tobacco chloroplasts. Plant and Cell Physiology 39: 1226-1231

Google Scholar: Author Only Title Only Author and Title

Foyer CH, Neukermans J, Queval G, Noctor G, Harbinson J (2012) Photosynthetic control of electron transport and the regulation of gene expression. J Exp Bot 63: 1637-1661

Google Scholar: Author Only Title Only Author and Title

Fridlyand LE, Backhausen JE, Scheibe R (1998) Flux control of the malate valve in leaf cells. Archives of Biochemistry and Biophysics 349: 290-298

Google Scholar: Author Only Title Only Author and Title

Gardeström P, Igamberdiev AU (2016) The origin of cytosolic ATP in photosynthetic cells. Physiol. Plant. 157: 367-379

Google Scholar: Author Only Title Only Author and Title

Gerhardt R, Stitt M, Heldt HW (1987) Subcellular Metabolite Levels in Spinach Leaves : Regulation of Sucrose Synthesis during Diurnal Alterations in Photosynthetic Partitioning. Plant Physiol 83: 399-407

Google Scholar: Author Only Title Only Author and Title

Gotoh E, Matsumoto M, Ogawa K, Kobayashi Y, Tsuyama M (2010) A qualitative analysis of the regulation of cyclic electron flow around photosystem I from the post-illumination chlorophyll fluorescence transient in Arabidopsis: a new platform for the in vivo investigation of the chloroplast redox state. Photosynth Res 103: 111-123

Google Scholar: Author Only Title Only Author and Title

Heineke D, Riens B, Grosse H, Hoferichter P, Peter U, Flugge UI, Heldt HW (1991) Redox transfer across the inner chloroplast envelope membrane. Plant Physiol 95: 1131-1137

Google Scholar: Author Only Title Only Author and Title

Heinemeyer J, Scheibe B, Schmitz UK, Braun HP (2009) Blue native DIGE as a tool for comparative analyses of protein complexes. J Proteomics 72: 539-544

Google Scholar: Author Only Title Only Author and Title

Huang S, Lee CP, Millar AH (2015) Activity assay for plant mitochondrial enzymes. Methods Mol Biol 1305: 139-149

Google Scholar: Author Only Title Only Author and Title

Imamura H, Nhat KP, Togawa H, Saito K, Iino R, Kato-Yamada Y, Nagai T, Noji H (2009) Visualization of ATP levels inside single living cells with fluorescence resonance energy transfer-based genetically encoded indicators. Proc Natl Acad Sci U S A 106: 15651-15656

Google Scholar: Author Only Title Only Author and Title

Ishikawa N, Takabayashi A, Sato F, Endo T (2016) Accumulation of the components of cyclic electron flow around photosystem I in C4 plants, with respect to the requirements for ATP. Photosynth Res 129: 261-277

Google Scholar: Author Only Title Only Author and Title

Jacoby RP, Millar AH, Taylor NL (2015) Assessment of Respiration in Isolated Plant Mitochondria Using Clark-Type Electrodes. In J Whelan, MW Murcha, eds, Plant Mitochondria: Methods and Protocols. Springer New York, New York, NY, pp 165-185
  Google Scholar: Author Only Title Only Author and Title

Kerppola TK (2006) Design and implementation of bimolecular fluorescence complementation (BiFC) assays for the visualization of protein interactions in living cells. Nature Protocols 1: 1278-1286
  Google Scholar: Author Only Title Only Author and Title

Lamkemeyer P, Laxa M, Collin V, Li W, Finkemeier I, Schottler MA, Holtkamp V, Tognetti VB, Issakidis-Bourguet E, Kandlbinder A, Weis E, Miginiac-Maslow M, Dietz KJ (2006) Peroxiredoxin Q of Arabidopsis thaliana is attached to the thylakoids and functions in context of photosynthesis. Plant Journal 45: 968-981
  Google Scholar: Author Only Title Only Author and Title

Law YS, Zhang R, Guan X, Cheng S, Sun F, Duncan O, Murcha M, Whelan J, Lim BL (2015) Phosphorylation and Dephosphorylation of the Presequence of pMORF3 During Import into Mitochondria from Arabidopsis thaliana. Plant Physiol 169: 1-12
  Google Scholar: Author Only Title Only Author and Title

Lee CP, Eubel H, O'Toole N, Millar AH (2008) Heterogeneity of the mitochondrial proteome for photosynthetic and non-photosynthetic Arabidopsis metabolism. Mol Cell Proteomics 7: 1297-1316
  Google Scholar: Author Only Title Only Author and Title

Liang C, Cheng S, Zhang Y, Sun Y, Fernie AR, Kang K, Panagiotou G, Lo C, Lim BL (2016) Transcriptomic, proteomic and metabolic changes in Arabidopsis thaliana leaves after the onset of illumination. BMC Plant Biol 16: 43
  Google Scholar: Author Only Title Only Author and Title

Liang C, Zhang Y, Cheng S, Osorio S, Sun Y, Fernie AR, Cheung CYM, Lim BL (2015) Impacts of high ATP supply from chloroplasts and mitochondria on the leaf metabolism of Arabidopsis thaliana. Front. Plant Sci. 6: 922
  Google Scholar: Author Only Title Only Author and Title

Lim SL, Voon CP, Guan X, Yang Y, Gardestrom P, Lim BL (2020) In planta study of photosynthesis and photorespiration using NADPH and NADH/NAD+ fluorescent protein sensors. Nature communications 11
  Google Scholar: Author Only Title Only Author and Title

Meyer EH, Tomaz T, Carroll AJ, Estavillo G, Delannoy E, Tanz SK, Small ID, Pogson BJ, Millar AH (2009) Remodeled respiration in ndufs4 with low phosphorylation efficiency suppresses Arabidopsis germination and growth and alters control of metabolism at night. Plant Physiol 151: 603-619
  Google Scholar: Author Only Title Only Author and Title

Munekage Y, Hojo M, Meurer J, Endo T, Tasaka M, Shikanai T (2002) PGR5 is involved in cyclic electron flow around photosystem I and is essential for photoprotection in Arabidopsis. Cell 110: 361-371
  Google Scholar: Author Only Title Only Author and Title

Nagai T, Sawano A, Park ES, Miyawaki A (2001) Circularly permuted green fluorescent proteins engineered to sense Ca2+. Proc Natl Acad Sci U S A 98: 3197-3202
  Google Scholar: Author Only Title Only Author and Title

Noctor G, Foyer CH (2000) Homeostasis of adenylate status during photosynthesis in a fluctuating environment. J Exp Bot 51: 347-356
  Google Scholar: Author Only Title Only Author and Title

Noguchi K, Yoshida K (2008) Interaction between photosynthesis and respiration in illuminated leaves. Mitochondrion 8: 87-99
  Google Scholar: Author Only Title Only Author and Title

Oikawa K, Matsunaga S, Mano S, Kondo M, Yamada K, Hayashi M, Kagawa T, Kadota A, Sakamoto W, Higashi S, Watanabe M, Mitsui T, Shigemasa A, Iino T, Hosokawa Y, Nishimura M (2015) Physical interaction between peroxisomes and chloroplasts elucidated by in situ laser analysis. Nature Plants 1: 15035
  Google Scholar: Author Only Title Only Author and Title

Oxborough K, Baker NR (1997) Resolving chlorophyll a fluorescence images of photosynthetic efficiency into photochemical and non-photochemical components – calculation of qP and Fv-/Fm-; without measuring Fo. Photosynthesis Research 54: 135-142
  Google Scholar: Author Only Title Only Author and Title

Sato R, Kawashima R, Trinh MDL, Nakano M, Nagai T, Masuda S (2019) Significance of PGR5-dependent cyclic electron flow for optimizing the rate of ATP synthesis and consumption in Arabidopsis chloroplasts. Photosynth Res 139: 359-365
  Google Scholar: Author Only Title Only Author and Title

Schaeffer GW, Sharpe FT, Sicher RC (1997) Fructose 1,6-bisphosphate aldolase activity in leaves of a rice mutant selected for enhanced lysine. Phytochemistry 46: 1335-1338
  Google Scholar: Author Only Title Only Author and Title

Scheibe R (1987) Nadp+-Malate Dehydrogenase in C-3-Plants - Regulation and Role of a Light-Activated Enzyme. Physiol Plantarum 71: 393-400
  Google Scholar: Author Only Title Only Author and Title

Scheibe R (2004) Malate valves to balance cellular energy supply. Physiol. Plant. 120: 21-26
Google Scholar: Author Only Title Only Author and Title

Scheibe R, Backhausen JE, Emmerlich V, Holtgrefe S (2005) Strategies to maintain redox homeostasis during photosynthesis under changing conditions. J Exp Bot 56: 1481-1489
Google Scholar: Author Only Title Only Author and Title

Schwarzländer M, Logan DC, Fricker MD, Sweetlove LJ (2011) The circularly permuted yellow fluorescent protein cpYFP that has been used as a superoxide probe is highly responsive to pH but not superoxide in mitochondria: implications for the existence of superoxide 'flashes'. Biochem J 437: 381-387
Google Scholar: Author Only Title Only Author and Title

Schweiger R, Schwenkert S (2014) Protein-protein Interactions visualized by bimolecular fluorescence complementation in tobacco protoplasts and leaves. JoVe-Journal of Visualized Experiments 85: e51327
Google Scholar: Author Only Title Only Author and Title

Selinski J, Scheibe R (2019) Malate valves: old shuttles with new perspectives. Plant Biol (Stuttg) 21: 21-30
Google Scholar: Author Only Title Only Author and Title

Shameer S, Ratcliffe RG, Sweetlove LJ (2019) Leaf energy balance requires mitochondrial respiration and export of chloroplast NADPH in the light. Plant Physiology 180: 1947-1961
Google Scholar: Author Only Title Only Author and Title

Sun F, Carrie C, Law S, Murcha MW, Zhang R, Law YS, Suen PK, Whelan J, Lim BL (2012) AtPAP2 is a tail-anchored protein in the outer membrane of chloroplasts and mitochondria. Plant Signal Behav 7: 927-932
Google Scholar: Author Only Title Only Author and Title

Sun F, Liang C, Whelan J, Yang J, Zhang P, Lim BL (2013) Global transcriptome analysis of AtPAP2--overexpressing Arabidopsis thaliana with elevated ATP. BMC Genomics 14: 752
Google Scholar: Author Only Title Only Author and Title

Sun F, Suen PK, Zhang Y, Liang C, Carrie C, Whelan J, Ward JL, Hawkins ND, Jiang L, Lim BL (2012) A dual-targeted purple acid phosphatase in Arabidopsis thaliana moderates carbon metabolism and its overexpression leads to faster plant growth and higher seed yield. New Phytol 194: 206-219
Google Scholar: Author Only Title Only Author and Title

Tikkanen M, Mekala NR, Aro EM (2014) Photosystem II photoinhibition-repair cycle protects Photosystem I from irreversible damage. Biochimica Et Biophysica Acta-Bioenergetics 1837: 210-215
Google Scholar: Author Only Title Only Author and Title

Voon CP, Guan X, Sun Y, Sahu A, Chan MN, Gardeström P, Wagner S, Fuchs P, Nietzel T, Versaw WK, Schwarzländer M, Lim BL (2018) ATP compartmentation in plastids and cytosol of Arabidopsis thaliana revealed by fluorescent protein sensing. Proc Natl Acad Sci U S A 115: 10778-10787
Google Scholar: Author Only Title Only Author and Title

Voon CP, Lim BL (2019) ATP translocation and chloroplast biology National Science Review 6: 1073-1076
Google Scholar: Author Only Title Only Author and Title

Wittig I, Braun HP, Schagger H (2006) Blue native PAGE. Nat Protoc 1: 418-428
Google Scholar: Author Only Title Only Author and Title

Zhang R, Guan X, Law YS, Sun F, Chen S, Wong KB, Lim BL (2016) AtPAP2 modulates the import of the small subunit of Rubisco into chloroplasts. Plant Signal Behav 11: e1239687
Google Scholar: Author Only Title Only Author and Title

Zhang R, Qi H, Sun Y, Xiao S, Lim BL (2017) Transgenic Arabidopsis thaliana containing increased levels of ATP and sucrose is more susceptible to Pseudomonas syringae. PLoS One 12: e0171040
Google Scholar: Author Only Title Only Author and Title

Zhao YN, Luo LL, Xu JS, Xin PY, Guo HY, Wu J, Bai L, Wang GD, Chu JF, Zuo JR, Yu H, Huang X, Li JY (2018) Malate transported from chloroplast to mitochondrion triggers production of ROS and PCD in Arabidopsis thaliana. Cell Research 28: 448-461
Google Scholar: Author Only Title Only Author and Title