A new mechanism of fibronectin fibril assembly revealed by live imaging and super-resolution microscopy.

Darshika Tomer[1], Cecilia Arriagada[1], Sudipto Munshi[2,‡], Brianna E. Alexander[1,3], Brenda French[2], Pavan Vedula[4], Valentina Caorsi[5], Andrew House[6], Murat Guvendiren[6], Anna Kashina[4], Jean E. Schwarzbauer[7], and Sophie Astrof[1,*].

1. Department of Cell Biology and Molecular Medicine, Cardiovascular Research Institute, Rutgers Biomedical and Health Sciences, 185 South Orange Ave, Newark, NJ, 07103, USA.

2. Sidney Kimmel Medical College of Thomas Jefferson University, Philadelphia, PA, 19107, USA

3. Multidisciplinary Ph.D. Program in Biomedical Sciences. Cell Biology, Neuroscience and Physiology track, Rutgers Biomedical and Health Sciences, Newark, NJ, 07103, USA.

4. Department of Biomedical Sciences, University of Pennsylvania, Philadelphia, PA 19104, USA

5. Abbelight, 191 Avenue Aristide Briand, 94230 Cachan, France

6. Otto H. York Chemical and Materials Engineering, Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ 07102, USA

7. Department of Molecular Biology, Princeton University, Princeton, New Jersey, USA

27   ‡Current Address: Department of Chemical and Structural Biology, Weizmann Institute of

28   Science, Rehovot, Israel

30   *Author for correspondence: Sophie Astrof, Ph.D.

31   Phone: 617-429-8295

32   E-mail: sophie.astrof@rutgers.edu

Abstract

34• Fn1 fibrils have long been viewed as continuous fibers composed of extended, periodically

aligned Fn1 molecules. However, our live imaging and single-molecule localization microscopy

(SMLM) are inconsistent with this traditional view and show that Fn1 fibrils are composed of

roughly spherical nanodomains containing 6-11 Fn1 dimers. As they move toward the cell

center, Fn1 nanodomains become organized into linear arrays, wherein nanodomains are

spaced at the average periodicity of 105±17 nm. Periodical Fn1 nanodomain arrays are bona

fide fibrils: they are resistant to deoxycholate treatment and retain nanodomain periodicity in

the absence of cells. The nanodomain periodicity in fibrils remained constant when probed with

antibodies recognizing distinct Fn1 epitopes or combinations of antibodies recognizing

epitopes spanning the length of Fn1. FUD, a bacterial peptide that binds Fn1 N-terminus and

disrupts Fn1 fibrillogenesis does not disrupt the formation of Fn1 nanodomains, instead, it

blocks the organization of Fn1 nanodomains into periodical arrays. These studies establish a

new paradigm of Fn1 fibrillogenesis.

Introduction

Fibronectin (Fn1) is a requisite component of the extracellular matrix (ECM) necessary for embryogenesis and homeostasis (Schwarzbauer and DeSimone, 2011). In the absence of Fn1 fibrillogenesis, the binding of Fn1 to cells is not sufficient to regulate key biological processes such as embryonic development, angiogenesis, vascular remodeling, or cartilage condensation (Chiang et al., 2009; Rozario et al., 2009; Singh and Schwarzbauer, 2014; Zhou et al., 2008). Therefore, understanding the mechanisms by which Fn1 proteins assemble into macromolecular fibrils is essential to gain insights into the various functions of Fn1. Fn1 is secreted as a dimer wherein two Fn1 molecules are held in an anti-parallel orientation by two disulfide bonds close to their C-termini (Skorstengaard et al., 1986; Wagner and Hynes, 1979). Fn1 fibrillogenesis is a cell-dependent process (McKeown-Longo and Mosher, 1983), occurs following the binding of Fn1 dimers to cell-surface integrins (Schwarzbauer and DeSimone, 2011). Following integrin binding, intracellular cytoskeletal forces such as actomyosin contractility acting through integrins generate pulling forces on Fn1 dimers, exposing epitopes that promote Fn1 fibrillogenesis (Chernousov et al., 1987; Hocking et al., 1994; Smith et al., 2007; Zhang et al., 1994; Zhang et al., 1997; Zhong et al., 1998). At a cellular level, the process of Fn1 fibrillogenesis is correlated with the formation of fibrillar adhesions, whereby small mobile adhesions containing Fn1 and integrin $\alpha5\beta1$ somehow elongate while translocating toward the nucleus, first giving rise to focal adhesions (< 1 μm), and then as the translocation continues, to longer filaments, termed fibrillar adhesions ($\geq$1 μm) containing Fn1 and intracellular cytoplasmic effectors linking Fn1 and actin cytoskeleton (Geiger et al., 2001; Geiger and Yamada, 2011; Lu et al., 2020; Pankov et al., 2000; Zamir et al., 1999; Zamir et al., 2000).

It has been thought that Fn1 fibrils resemble ropes, in which extended Fn1 dimers align such that regions containing overlapping N-termini alternate with regions containing C-termini

(Chen et al., 1997; Dzamba and Peters, 1991; Fruh et al., 2015), also illustrated in **Sup. Fig.**

8A. To understand how the process of fibrillogenesis occurs in real-time and at a nanoscale

level, we adopted a CRISPR/Cas9-mediated mutagenesis approach to generate fluorescently-

labeled Fn1, which was subject to the physiological regulation of expression and splicing. This

approach has enabled visualization of Fn1 fibrillogenesis over an extended time. Using live

imaging and super-resolution microscopy, we uncovered an unexpected mechanism of Fn1

fibrillogenesis.

Our data demonstrate that Fn1 fibrils form as a result of centripetally-moving Fn1

nanodomains originating at the cell periphery. As Fn1 nanodomains move toward the cell

center, they assemble into arrays of periodically-spaced nanodomains. The arrays become

longer as the movement towards the cell center continues and as more nanodomains are

added. We show that these nanodomain arrays are in bona fide fibrils and that each Fn1

nanodomain contains multiple Fn1 dimers. Our live imaging and SMLM reveal a new role by

which the N-terminal region of Fn1 protein regulates fibril assembly, and show that interactions

mediated by the N-terminal Fn1 assembly region are not required for the formation of Fn1

nanodomains or their centripetal translocation; Instead, the N-terminus of Fn1 regulates the

organization of Fn1 nanodomains into nanodomain arrays. Our model integrates the process

of fibrillogenesis with the process of adhesion elongation and provides significant new insights

into the mechanisms of Fn1 ECM formation, remodeling, and signaling.

Results

Beaded structure of Fn1 fibrils revealed by diffraction-limited microscopy

While examining Fn1[+] ECM in mid-gestation mouse embryos by confocal

immunofluorescence microscopy using an Abcam monoclonal anti-Fn1 antibody which binds

an epitope within a central region of Fn1 (**Table M2** in Methods), we observed that Fn1 fibrils

in the pharyngeal arches and the heart appeared dotted, with regularly-spaced regions of high

and low fluorescence intensity (**Fig. 1, Movie 1**). The dotted appearance of embryonic Fn1

suggested that the distribution of Fn1 molecules in Fn1 fibrils is not homogenous. To

understand how Fn1 fibrils form, we employed a CRISPR/Cas9 knock-in strategy to modify the

endogenous Fn1 locus by replacing the termination codon of Fn1 with a sequence encoding a

fluorescent protein. We used this strategy to generate cell lines and $Fn1^{mEGFP/mEGFP}$

homozygous knock-in mice expressing Fn1 fused to monomeric enhanced green fluorescent

protein (mEGFP) or other monomeric fluorescent proteins (**Sup. Figs. 1-2**). Homozygous

$Fn1^{mEGFP/mEGFP}$ mice were obtained at the correct Mendelian ratio (**Sup. Fig. 1B,** panels 4 and

5), and are viable and fertile, indicating that Fn1-mEGFP supports all functions of Fn1

necessary for embryonic development, fetal viability, and adult homeostasis. Examination of

Fn1 expression patterns in knock-in mice showed that Fn1-mEGFP is expressed in the same

pattern as the unmodified Fn1 in embryos (Peters and Hynes, 1996), i.e., there were no

regions that were GFP+ but Fn1-negative and vice versa in $Fn1^{mEGFP/+}$ embryos expressing

one wild-type allele of Fn1 (**Sup. Fig. 2D**). In addition, we used CRISPR/Cas9 mutagenesis to

generate five independent lines of mouse embryo fibroblasts (MEFs) expressing Fn1-mEGFP,

Fn1-mScarlet-I, Fn1-Neon Green, or Fn1-tdTomato fluorescent proteins (FP). Western blots

showed that FP fusions to Fn1 were specific: FPs were only fused to Fn1 as no other FP

fusions were detected either by western blotting or immunofluorescence (IF) (**Sup. Fig. 2A-B**).

Measuring deoxycholate (DOC) insolubility of Fn1 ECM is a classical biochemical assay

to assay stable incorporation Fn1 proteins into the ECM (Choi and Hynes, 1979; McKeown-

Longo and Mosher, 1983; Schwarzbauer, 1991; Singh et al., 2010; Wierzbicka-Patynowski et

al., 2004). For these assays, we carefully controlled the number of cells plated, as cell density

affects the extent of Fn1 fibrillogenesis (Hynes, 1990). We performed DOC insolubility assays

using 5 independently-generated cells lines expressing Fn1-FP proteins. These experiments

demonstrated that the incorporation of Fn1-FPs into the ECM was indistinguishable from wild-

type, untagged Fn1 (**Sup. Fig. 2C**). Together with the viability of $Fn1^{mEGFP/mEGFP}$ homozygous

knock-in mice, these data demonstrate that Fn1-FP fusion proteins are suitable reagents for

investigating mechanisms of Fn1 fibrillogenesis.

To visualize the process of fibrillogenesis in real-time, we plated Fn1$^{mEGFP/+}$ MEFs on

the gelatin-coated cover glass and imaged cells 16 hours after plating using total internal

reflection (TIRF) microscopy at the critical angle of incidence. These experiments showed that

Fn1 fibrillogenesis initiated at cell periphery as distinct, brightly-fluorescent Fn1 densities that

moved centripetally in parallel with F-actin and became aligned into linear arrays of "beads"

(arrows in **Movie 2**). TIRF imaging also showed that domains of higher fluorescence intensity

of Fn1 co-localized with integrin $\alpha5\beta1$ both in non-fibrillar (arrows in **Fig. 2A-A2**) and in fibrillar

adhesions (arrows in **Fig. 2B-B2**), and that both Fn1 and $\alpha5\beta1$ fibrillar adhesions appeared

beaded (**Fig. 2B-B2**, arrows). Beaded architecture of Fn1 fibrils was also observed by an

independent imaging method using Zeiss Airyscan technology (**Sup. Fig. 3**). To test whether

the beaded appearance of Fn1 fibrils depended on antibody epitope availability, we stained

wild-type MEFs or mouse endothelial cells using three different antibodies 1) 297.1 polyclonal

antibody raised against the rat plasma Fn1 protein and recognizing multiple Fn1 epitopes

(**Sup. Fig. 4**); 2) 3E2 monoclonal antibody recognizing the alternatively spliced EIIIA exon of

Fn1, and 3) an Abcam monoclonal antibody recognizing an epitope within the central region of

Fn1 (see **Table M2**, in Methods). The beaded appearance of Fn1 fibrils did not depend on the

type of antibody used (**Sup. Fig. 5**). Moreover, staining with a polyclonal antibody to Fn1,

297.1, which recognizes multiple epitopes along the Fn1 molecule resulted in the beaded

appearance of Fn1 fibrils (**Sup. Fig. 5A-A1**). Fn1 fibrils formed by cells plated on uncoated

glass, as well as on glass coated with gelatin, laminin 111, or vitronectin, also appeared

beaded **Sup. Figs. 5-6**). Similarly, Fn1 fibrils in cell-free areas (**Sup. Fig. 5A, A1**) and between

cells were beaded (**Sup Fig. 5B, B1, C, C1**). Fn1 fibrils produced by cells plated on soft

substrata such as hydrogels of variable stiffness also appeared beaded (**Sup. Fig. 6C-D, 6C1-**

D1). In this latter experiment, Fn1 was detected by imaging the fluorescence of Fn1-mEGFP

protein, indicating that the beaded appearance of Fn1 fibrils is independent of antibody

staining. To determine whether Fn1 fibrils in cell-free, fibrillar ECM were beaded, cultures were

treated with 2% DOC (Lu et al., 2020). 2% DOC treatment dissolves cell membranes and

cytoplasmic components, leaving behind ECM devoid of cell contact (**Movie 3** for time-lapse of

dissolution of cellular components, F-actin and DNA). This experiment showed that Fn1 fibrils

retained their beaded architecture in the absence of cell contact (**Sup. Fig. 5D**). Taken

together, these studies indicated that the beaded appearance of Fn1 fibrils is a general feature

of Fn1 ECM that can be seen in multiple contexts: in DOC-resistant cell-free fibrils, in different

embryonic tissues *in vivo*, and when cells are cultured on various substrata *in vitro*. Moreover,

the beaded appearance of Fn1 fibrils is independent of the type of antibody and a method of

detection: It is seen by indirect detection methods such as immunofluorescence and by direct

detection of fluorescent Fn1-mEGFP protein.

Nanoarchitecture of Fn1 fibrils revealed by standardized single-molecule localization

microscopy.

To examine the structure of Fn1 fibrils at a nanoscale level, we adopted single-molecule

localization microscopy (SMLM) using the direct stochastic optical reconstruction microscopy

(dSTORM) method (Heilemann et al., 2008; Rust et al., 2006). To optimize SMLM imaging

conditions and determine the effective labeling efficiency of our reagents, we used the

methodology and the reference cell line NUP96-mEGFP, as gold-standard tools to optimize

image quality, measure effective labeling efficiency, and to quantify the number of GFP

molecules in our structure(s) of interest (Lelek et al., 2021; Thevathasan et al., 2019). In this

reference cell line, both copies of the *NUP96* gene are modified by CRISPR/Cas9

mutagenesis to generate NUP96-mEGFP protein, a component of the nuclear pore complex,

NPC (Thevathasan et al., 2019). Thirty-two copies of NUP96 protein are evenly distributed

among the eight corners of the NPC at known distances (see schematic in **Sup. Fig. 7**) (Bui et

al., 2013; Thevathasan et al., 2019; von Appen et al., 2015). To optimize our SMLM imaging,

we were guided by the current best SMLM practices (Lelek et al., 2021; Mund and Ries, 2020),

the methodology, and the SMAP software developed by (Diekmann et al., 2020; Ries, 2020;

Thevathasan et al., 2019). Measurements using the Fourier ring correlation method

(Nieuwenhuizen et al., 2013) implemented in the SMAP software indicated that the resolution

of our images ranged between 14 and 28 nm. Localization precision was 9.1±1.9 nm, on

average. Using antibodies to detect NUP96-mEGFP and the SMAP software, we analyzed

images from 3 independent experiments, 8 nuclei and 4571 NPCs, and determined the NPC

radius to be 63.6±0.86 nm (**Fig. 3A-C** and **Table 1**). This radius is consistent with the reported

NPC radius of 64.3+/-2.6 nm, measured using this cell line and a combination of commercial

anti-GFP primary and Alexa-647-conjugated secondary antibodies (Thevathasan et al., 2019).

We then used NUP96-mEGFP cells to optimize the effective labeling efficiency (ELE) of our

reagents. ELE summarily measures how well one's reagents and methods, including the

SMLM imaging protocol, generate images that reflect the actual structure under the study

(Thevathasan et al., 2019). SMLM imaging of NUP96 in the x-y plane should produce 8-fold

symmetrical ring-like structures wherein 4 copies of the NUP96 protein are positioned at each

of the NPC's 8 vertices at known intervals and with known dimensions (Thevathasan et al.,

2019). Using NUP96-mEGFP as a reference cell line, anti-GFP primary antibodies,

commercially-labeled Alexa-647-conjugated secondary antibodies, and the SMAP software

(Thevathasan et al., 2019), we analyzed 4,571 NPCs from 8 cells and 3 independent

experiments, and determined the ELE to be 79.6±5.4% (**Table 1**), reflecting that the majority

of NPCs have eight NUP96[+] corners in our images (**Fig. 3A-C**). This ELE and the standard

deviation of our measurements (<10%) are comparable with the best ELE measured for this

system ~74% (Thevathasan et al., 2019). Together, these experiments indicate that our

reagents and SMLM imaging conditions are within the accepted SMLM standards (Lelek et al.,

2021).

To attain a comparable ELE to that measured with NUP96-mEGFP, we stained

homozygous Fn1$^{mEGFP/mEGFP}$ MEFs at the same time as NUP96-mEGFP cells, using aliquots of

the same mixtures of reagents. Furthermore, pairs of stained Fn1$^{mEGFP/mEGFP}$ MEFs and

NUP96-mEGFP cells were imaged on the same day, using aliquots of the same imaging

buffer, and the same off-switching and imaging protocols (see Methods). Similar to (Fruh et al.,

2015), we saw that Fn1 fibrils appeared as arrays of nanodomains situated along Fn1 fibrils

with a regular periodicity (**Figs. 3D-F, Fig. 5D-D2, Fig. 6D-D"**). As in (Fruh et al., 2015),

nanodomain periodicity was ascertained using autocorrelation, wherein the position of the first

autocorrelated peak marks the nanodomain periodicity (see discussion and simulations in

(Fruh et al., 2015)). Autocorrelation analysis of 14 fibrils from 7 cells and 3 independent

experiments indicated that the nanodomain periodicity of the GFP epitopes in Fn1-mEGFP

fibrils was 102±29 nm (**Figs. 6D-D", 6F, Table 2**). This periodicity is comparable to the

nanodomain periodicity measured by (Fruh et al., 2015), which was 99±17 nm, and the

periodicity determined by immuno-electron microscopy using anti-EIIIA antibody, which was on

average 84 nm and ranged between 40 and 280 nm (Dzamba and Peters, 1991).

Localization of Fn1 epitopes within periodical nanodomain arrays

Fn1 is a large, multi-domain, ~250 kDa glycoprotein secreted as a dimer, wherein two

Fn1 subunits are linked in an anti-parallel orientation by two di-sulfide bonds at their C-termini

(**Fig. 4A**), (Schwarzbauer and DeSimone, 2011; Skorstengaard et al., 1986; Wagner and

Hynes, 1979). To investigate the relationship between the protein domains of Fn1 and the

nanodomain architecture of Fn1 fibrils, we first used antibodies to different regions of Fn1

protein (**Figs. 4B** and **Table M2**). For these experiments, Fn1$^{mEGFP/mEGFP}$ MEFs were plated on

uncoated glass coverslips. Cells were then fixed and stained with antibodies recognizing

different Fn1 epitopes: 1) R457 rabbit polyclonal antibodies raised to the N-terminal 70 kDa

domain of Fn1 (Aguirre et al., 1994; Sechler et al., 2001), 2) R184 rabbit polyclonal antibodies

raised to recognize the domain immediately following the 70 kDa N-terminal domain and

containing the first six type III repeats of Fn1 (Fn1 III$_{1-6}$) (Raitman et al., 2018), 3) a rabbit

monoclonal antibody from Abcam recognizing an epitope within the central region of Fn1

(**Table M2** for further description of this antibody), and 4) an anti-GFP antibody recognizing the

C-terminus of Fn1-mEGFP protein. In our analyses, we focused on long fibrillar adhesions that

were over 1 μm in length, characteristic of mature assembled Fn1 fibrils (Lu et al., 2020).

Independent of the antibodies used, Fn1 fibrils appeared as arrays of periodically-

spaced nanodomains (**Figs. 5-6**). To ensure that the beaded appearance of fibrils was not due

to under-sampling, we tested 2 – 4 dilutions of each antibody, followed by SMLM imaging and

autocorrelation analysis, as in (Fruh et al., 2015). Our analyses showed that nanodomain

periodicity remained constant at all antibody dilutions tested (**Fig. 5, 6A"-E', summarized in**

6F). The periodicity of nanodomains detected with either R457, R184, or Abcam monoclonal

antibodies was similar to that measured with the anti-GFP antibody, suggesting that each of

these antibodies was used at the ELE comparable with that of GFP-labeling reagents. The

finding that nanodomain periodicity was independent of antibody specificity supports the

conclusion that the beaded appearance of Fn1 fibrils is not a result of a particular Fn1 protein

conformation, since staining using antibodies to three different regions of Fn1 in addition to

antibodies to GFP resulted in the same pattern. In all cases, nanodomain periodicity was

independent of the antibody concentrations, arguing against under-sampling (Fruh et al.,

2015).

Analysis using DBSCAN (Caetano et al., 2015) discovered clusters corresponding with

nanodomains, when the radius of the neighborhood (ε) was set to 14 nm, the average

apparent radius of an Fn1 nanodomain (**Table 2**, compare (**Figs. 7A and B**). Unbiased

DBSCAN clustering using the neighborhood radius that was automatically estimated by the

SMAP software discovered larger clusters of Fn1 nanodomains (**Fig. 7C**). This suggests that

Fn1 nanodomains are organized in a higher-order structure within long fibrils, consistent with

the observation of "beads" in lower-resolution diffraction-limited images of Fn1 (**Figs. 1, 2** and

Sup. Figs. 3, 5, 6).

Fn1 nanodomains contain multiple full-length Fn1 molecules.

It was previously-thought that Fn1 nanodomains detected by immunoelectron

microscopy or by immunofluorescence SMLM corresponded with particular regions in Fn1

protein sequence (Dzamba and Peters, 1991; Fruh et al., 2015), illustrated in **Sup. Fig. 8A-B**.

Surprisingly, staining using 297.1 polyclonal anti-Fn1 antibody raised against full-length

plasma Fn1 and recognizing multiple epitopes along Fn1 protein resulted in the same

periodicity of nanodomains as staining with antibodies to distinct regions of Fn1 protein (**Figs.**

6E, 6F, Figs. 8A, 8D). Since 297.1 polyclonal antibody recognizes multiple epitopes along

Fn1, including an epitope in the 70 kDA N-terminal assembly region of Fn1 (**Sup. Fig.4**), these

experiments suggested that each Fn1 nanodomains contained at least one entire Fn1 dimer.

To determine the number of Fn1 molecules per nanodomain, we used the methodology

developed by (Thevathasan et al., 2019), and identical labeling reagents and imaging

conditions for the detection of mEGFP in NUP96-mEGFP and Fn1-mEGFP cells (Thevathasan

et al., 2019). This analysis indicates that each Fn1 nanodomain contains 16.85 +/- 5.1 mEGFP

proteins on average (**Table 2**). Since Fn1 assembled in ECM fibrils is an obligate dimer of two

disulfide-bonded Fn1 molecules (Schwarzbauer, 1991), these findings show that Fn1

nanodomains contain 6 – 11 Fn1-mEGFP dimers, on average.

These results are not consistent with the model that thinnest Fn1 fibrils are made of

extended single Fn1 molecules that are periodically aligned in a regular, end-to-end fashion

with regions containing N-termini alternating with regions containing C-termini (Chen et al.,

1997; Dzamba and Peters, 1991; Fruh et al., 2015) and illustrated in **Sup.** **Fig. 8A-B**. Such

periodic alignment of Fn1 dimers necessitates that staining using antibodies recognizing

multiple epitopes along the length of Fn1 molecule would result in a uniform labeling of Fn1

fibrils, (**Sup**. **Fig. 8C**). However, this is not what we observed: staining with a polyclonal 297.1

antibody showed that Fn1 fibrils were composed of periodically-spaced nanodomains (**Fig. 8A-**

A", 8D). Autocorrelation analysis showed that the spacing was periodical at multiple

concentrations of 297.1 antibody (**Figs. 6E-E", 6F,** and **8D**). Importantly, the periodicity of

nanodomains detected using 297.1 antibody was statistically indistinguishable from

periodicities seen with any regions-specific antibody tested, including that of anti-GFP

antibodies (**Fig. 6F**). The latter suggested that 297.1 polylconal antibodies were used at an

ELE that was at least as high as that for GFP labeling reagents. Finally, the periodical

distribution of 297.1 epitopes (average periodicity 121±11) and the presence of multiple Fn1

dimers per nanodomain, does not fit with the model proposed by (Dzamba and Peters, 1991),

illustrated in **Sup**. **Fig. 8A**, and suggest a different model of Fn1 fibrillogenesis (e.g., **Fig. 8G**).

To further evaluate the hypothesis that Fn1 nanodomains contain full-lengths Fn1

dimers, we used combinations of antibodies and the imaging parameters resulting in the high

ELE (**Fig. 6F** and Methods). We first stained Fn1$^{mEGFP/mEGFP}$ MEFs using a combination of

R457 and anti-GFP antibodies, detecting the N- and C- termini of Fn1-mEGFP, respectively.

Consistent with the staining using 297.1 antibody, the periodicity of nanodomains detected with

a mixture of R457 and anti-GFP antibodies was 101±45 nm, which is statistically

indistinguishable from the periodicities when either of these antibodies were used alone (**Figs.**

6F, 8B-B", 8E).

Next, we stained Fn1$^{mEGFP/mEGFP}$ MEFs with a cocktail of five antibodies recognizing

epitopes distributed along the length of the Fn1 molecule: 1) the N-terminal 70 kDa assembly

domain of Fn1 (R457), 2) the sequence immediately following the 70 kDA domain (R184), 3)

an epitope in the middle of Fn1 (Abcam monoclonal), 4) multiple epitopes along the Fn1

protein (297.1), and 5) the anti-GFP antibody marking the C-terminus of Fn1-mEGFP (αGFP)

(Fig. 8C-C", 8F). Each antibody was used at a dilution resulting in a periodical staining (**Fig.**

6F and legend). The binding of all primary antibodies in the cocktail was detected by a cocktail

of Alexa 647-conjugated secondary antibodies. If Fn1 fibrils were indeed composed of

continuous, linear arrays of extended and periodically aligned Fn1 molecules, this cocktail of

antibodies recognizing epitopes from the N- to the C-termini of Fn1 would uniformly label Fn1

fibrils. Even in the scenario in which R457 epitopes did not extend to the N-terminal-most

sequence of Fn1, the staining using this mixture of five antibodies would not be expected to

produce periodical staining. However, we observed that the cocktail of 5 antibodies labeled

nanodomains with the average periodicity of 94±47 nm at the spatial resolution of 23 nm in thin

Fn1 fibrils (**Fig. 8C-8C", 8F**). This periodicity was statistically indistinguishable from the

periodicities of individual antibodies at all the dilutions tested (**Fig. 6F**).

Unbiased clustering using the Voronoi tessellation method (Andronov et al., 2016b)

implemented in SMAP showed that Fn1 fibrils stained with the 5-antibody cocktail can be

segmented into clusters (**Fig. 8C", 8C"-1**). This analysis shows that the nanodomain structure

of Fn1 fibrils can be discovered using an unbiased computational approach. The size of the

nanodomains detected with the 5-antibody cocktail was larger than when nanodomains were

detected by the use of any single antibody (e.g., compare **Fig. 8A"** with **Fig. 8C"**). This type of

behavior is expected in SMLM when molecules are clustered (Baumgart et al., 2016).

In agreement with the results described above, periodically-spaced nanodomains were

detected in Fn1 fibrils using two other cell lines, heterozygous Fn1^mEGFP/+ MEFs and wild-type

MEFs (**Sup. Fig. 9**). The presence of the periodical nanodomains in fibrils produced by wild-

type cells indicates that the periodical structure of Fn1 fibrils is not an artifact of GFP labeling.

Nanodomain spacing in Fn1 fibrils treated with 2% DOC, which removes cells and cellular

components, was similar to that of untreated fibrils (**Sup. Fig. 9,** columns **4** and **8,** and **Sup.**

Fig. 9F). Together, these data indicate that the nanodomain architecture of Fn1 fibrils is

independent of antibody(ies) used for staining, is seen in cells expressing wild-type Fn1 and in

the absence of cell contact, and that periodically-spaced nanodomains within Fn1 fibrils

contain multiple Fn1 dimers (e.g., model in **Fig. 8G**).

To further test the hypothesis that Fn1 nanodomains contain full-length Fn1 molecules,

we performed double-color dSTORM experiments, wherein the N-terminus of Fn1 was labeled

with the R457 primary antibody and anti-rabbit CF680-conjugated secondary immunoglobulins,

and the C-terminus of Fn1-mEGFP was labeled with the anti-GFP primary antibody and Alexa-

647-conjugated anti-chicken secondary antibodies. As a control, we stained Fn1$^{mEGFP/mEGFP}$

MEFs using two different anti-GFP antibodies, one made in rabbit (detected using CF680-

conjugated secondary antibodies) and the other, in chicken (detected with Alexa-647-

conjugated anti-chicken secondary antibodies). The majority of nanodomains detected using

the two anti-GFP antibodies contained overlapping CF680 and AF647 localizations, suggesting

a high ELE for both reagents (**Fig. 9A-C,C'**; Gray lines in **Fig. 9A', 9B'** mark overlapping

staining in nanodomains). Nanodomains in which the two labels did not overlap were marked

by purple lines (**Fig. 9A', 9B'**). To determine the extent of co-localization between the two

labels, we performed coordinate-based colocalization (CBC) analysis, a widely-accepted

method for detecting co-localization in SMLM data (Malkusch et al., 2012). The CBC

coefficient was calculated by counting the number of CF680 localizations within a radius *r* of

each of the AF647 localizations and normalizing by the number of localizations within the area

πR^2_{max} (Malkusch et al., 2012) (**Fig. 9D,F**). Conversely, we also determined the CBC coefficient

by counting the number of AF647 localizations within a radius *r* of each of the CF680

localizations and normalizing by the number of localizations within the area πR^2_{max} (**Fig. 9E,F**).

The CBC coefficients were calculated using algorithms implemented either by the Abbelight

software or by the publicly-available ThunderSTORM software (Ovesny et al., 2014). The

coefficient "+1" indicates a high probability of co-localization, while CBC ≤ 0 indicates low

probability of co-localization. CBC analysis using the Abbelight software and color-coding

according to the CBC coefficient, showed that the parameters chosen for these analyses,

r=30nm and R_{max}=300nm, reflect the overlap (e.g., yellow color in **Fig. 9D-E** and gray lines in

Fig. 9A'-B') and the lack of overlap (purple color and arrows in **Fig. 9D-E** and purple lines and

arrows in **Fig. 9A'-B'**) seen in SMLM images. These figures also show that we can detect non-

overlapping signals in nanodomains located next to one another (**Fig. 9D-F**). CBC analyses

performed on twenty-two regions containing long (≥1μm) Fn1 fibrils from Fn1$^{mEGFP/mEGFP}$ MEFs

showed extensive overlap among localizations arising from rabbit and chicken anti-GFP

antibodies (**Fig. 9F**).

CBC analyses of twenty regions containing long fibrils from Fn1$^{mEGFP/mEGFP}$ MEFs

stained with R457 and anti-GFP antibodies showed extensive overlap between R457 and anti-

GFP localizations, detecting the N- and the C- termini of Fn1-mEGFP, respectively (**Fig. 9G-**

L). Using ThunderSTORM software, we varied the parameters of CBC analysis, e.g., setting

the radius r from 2 or 5 nm and R_{max} from 20 or 50nm, respectively. These analyses had the

same outcome as above, i.e., CBC coefficients were close to +1, indicating a high probability

of overlap among localizations resulting from R457 and anti-GFP antibody binding (**Sup. Fig.**

10). Together, these studies show that the N- and the C-termini of Fn1 are contained within

each Fn1 nanodomain and support the model wherein Fn1 nanodomains in Fn1 fibrils contain

multiple full-length Fn1 molecules (**Fig. 8G**).

Nanoarchitecture of Fn1 fibrils formed by ectopically-added Fn1

It is known that ectopic Fn1 added to cells becomes assembled into fibrils (McKeown-

Longo and Mosher, 1983). To determine whether fibrils incorporating ectopically-added Fn1

contain nanodomains, we first used live imaging to film the assembly of ectopic Fn1. In these

experiments, MEFs producing Fn1-mEGFP were cultured on the uncoated glass inside an

insert while a confluent monolayer of MEFs producing Fn1-tdTomato was grown in the space

surrounding the insert. Prior to imaging, the insert was lifted, and Fn1-mEGFP+ MEFs were

imaged at ~17 min intervals for 16 hours. These movies show that Fn1-tdTomato fibrils

become clearly visible on the surface of Fn1-mEGFP cells by about 3 hours of co-culture

without first accumulating inside cells, suggesting that ectopic Fn1 is assembled into fibrils at

the cell surface (**Movie 4**). To determine the nanoarchitecture of Fn1 fibrils formed by

ectopically-added Fn1, we cultured Fn1$^{mEGFP/mEGFP}$ MEFs in the presence of 10 μg of Fn1-

tdTomato for 24 hours. Cells were fixed and stained without permeabilization using rabbit anti-

Cherry primary antibodies and anti-rabbit Alexa647-conjugated secondary antibodies. ECM in

regions between cells was imaged using TIRF to detect Fn1-mEGFP and Fn1-tdTomato

fluorescence, and dSTORM to detect Fn1-tdTomato due to the Alexa 647 label. Fn1-tdTomato

extensively co-assembled with Fn1-mEGFP (**Sup. Fig. 11A-B**), and high-resolution SMLM

reconstructions of thin and thick fibrils contained nanodomain arrays composed of Fn1-

tdTomato proteins (**Sup. Fig. 11C-E, 11F**). Together, these experiments demonstrate that 1)

Fn1 fibrils deposited into the intercellular ECM space are composed of nanodomain arrays and

2) ectopic Fn1 assembles into fibrils comprised of nanodomain arrays.

Fibrillogenesis inhibitor FUD disrupts the organization of Fn1 nanodomains into

periodical arrays

To understand the relationship between the nanodomain architecture of Fn1 fibrils and

the process of fibrillogenesis, we adopted a live imaging approach using Fn1$^{mEGFP/+}$ MEFs and

inhibitors of fibrillogenesis. One such inhibitor is a 49-amino acid peptide derived from

Streptococcus pyogenes adhesin F1, termed the functional upstream domain (FUD), a highly

potent inhibitor of Fn1 fibrillogenesis (Tomasini-Johansson et al., 2001). Fn1 fibrillogenesis

critically depends on the interactions mediated by the N-terminal 70 kDa assembly domain of

Fn1 (**Fig. 4A**), and FUD is one of the inhibitors that interferes with these interactions (Filla et

al., 2017; Morla et al., 1994; Schwarzbauer, 1991; Sechler et al., 2001; Sechler et al., 1996;

Tomasini-Johansson et al., 2001).

To further investigate the mechanism of Fn1 fibrillogenesis and the role of the N-

terminal domain of Fn1 in this process, Fn1$^{mEGFP/+}$ MEFs were plated on uncoated glass for 4

hours, and then imaged for 15 – 18 hours either in the imaging medium alone, or in the

medium containing 225 nM FUD. We also imaged cells incubated with 274 nM 11-IIIC peptide,

a 68 amino-acid control peptide that does not interfere with Fn1 fibrillogenesis (Morla et al.,

1994; Sottile and Chandler, 2005). Untreated cells or cells treated with the control peptide

developed and accumulated long Fn1 fibrils (**Movie 5**). In contrast, treatment with FUD led to

a) the dismantling of the pre-existing Fn1 fibrils and b) inhibiting the formation of new Fn1

fibrils (**Movie 6**). Instead of fibrils, cells cultured in the presence of FUD mainly contained

centripetally-moving Fn1-mEGFP fluorescent "beads" that only rarely assembled into fibrils

(**Movie 6**). These experiments suggested that FUD inhibits fibrillogenesis by interfering with

the process by which Fn1 "beads" become arranged or connected into linear arrays. To test

this hypothesis, Fn1$^{mEGFP/+}$ MEFs were plated for 16 hours in the continuous presence of either

225 nM FUD or 274 nM III-11C control peptides or were left untreated. Cells were then washed

with PBS, fixed, and stained without permeabilization using monoclonal anti-Fn1 antibodies

and Alexa Fluor 647-conjugated secondary antibodies, and imaged at the critical angle of

incidence by STORM (Jimenez et al., 2020). This approach maximizes the detection of cell-

surface Fn1 due to: a) the absence of a detergent during fixation, staining and washing and b)

imaging at the critical angle of incidence to detect fluorescence in close proximity to the

plasma membrane. These experiments demonstrated that the organization of Fn1

nanodomains into linear arrays was lost upon incubation with FUD (compare **Fig. 10A, B, A1,**

B1 with **Fig. 10C, C1**). Non-fibrillar (NF) Fn1 nanodomains in cells treated with FUD had a

similar number of Fn1 localizations per nanodomain, and were of similar sizes compared with

fibrillar or non-fibrillar Fn1 nanodomains in untreated cells, or cells incubated with the control

peptide (**Fig. 10A2, 10B2, 10C2**, **10D**). Taken together, these data indicate that FUD does not

interfere with the formation of Fn1 nanodomains but inhibits the organization of Fn1

nanodomains into linear arrays. Since Fn1 proteins lacking the N-terminal assembly domain do

not form fibrils (Schwarzbauer, 1991), our experiments suggest that interactions mediated by

the N-terminal assembly domain of Fn1 with a yet unidentified factor(s), are critical for the

organization of Fn1 nanodomains into fibrillar arrays (**Fig. 10E** and *vide infra*).

Discussion

In this manuscript, we describe data supporting a novel mechanism underlying the process of

Fn1 fibrillogenesis. In this model, 6-11 Fn1 dimers assemble into nanodomains containing

integrin $\alpha5\beta1$ at the cell periphery, move rearward with actin flow, and become organized into

linear arrays of periodically-spaced nanodomains. Joining of the additional Fn1 nanodomains

to these arrays leads to the generation of longer and longer fibrils as nanodomain assemblies

move toward the cell center. We show that these periodical nanodomain assemblies are bona

fide Fn1 fibrils.

The N-terminus of Fn1 is essential for Fn1 fibrillogenesis (Schwarzbauer, 1991). FUD

binds tightly to the N-terminal domain of Fn1 (K_D<2.6 nM), with a fast k_{on} and a slow k_{off}, and

acts as a competitive inhibitor of Fn1-Fn1 interactions (Ma et al., 2015; Maurer et al., 2010;

Tomasini-Johansson et al., 2001). Although in the absence of FUD, individual fibrils in an

established matrix are stable and can be tracked for over 16 hours (S.A., unpublished

observations), when FUD is added to cells, it localizes with Fn1 fibrils and dismantles the

mature Fn1 ECM (Filla et al., 2017; Tomasini-Johansson et al., 2001). These findings suggest

that the fibrillar Fn1[+] ECM is maintained through dynamic interactions mediated by the 70 kDa

Fn1 N-terminal domain. Our live imaging experiments demonstrated that in addition to

dismantling pre-existing fibrils, FUD effectively blocks their de-novo formation. SMLM showed

that FUD does not affect the formation of Fn1 nanodomains, instead, it blocks the organization

of Fn1 nanodomains into linear arrays. Together, these data suggest that FUD blocks the

dynamic interactions between the N-terminal Fn1 assembly domain and factor(s) linking Fn1

nanodomains into fibrils.

Previously, the periodical distribution of domain-specific antibody epitopes in EM and

SMLM images of Fn1 fibrils was explained by a model stipulating that Fn1 fibrils are made of

aligned and extended Fn1 molecules (Dzamba and Peters, 1991) (**Sup. Fig. 8A-B**). In EM

micrographs of Fn1 molecules sprayed onto mica surface, one can see extended Fn1 dimers

that are on average 120-140 nm in length (Engel et al., 1981; Erickson et al., 1981). Therefore,

it was hypothesized that spacing of ~84 nm seen in EM with domain-specific antibodies

(Dzamba and Peters, 1991) or spacing of ~100 nm in SMLM images (Fruh et al., 2015) was

due to the overlap between the extended Fn1 molecules at their N-termini (Dzamba and

Peters, 1991), e.g., (**Sup. Fig. 8B,** overlap marked by a bubble). The findings that the N-

terminal 70 kDa domain is required for Fn1 fibrillogenesis, that this domain can mediate

intermolecular Fn1-Fn1 interactions, and that inhibitors of Fn1 fibrillogenesis bind to this

domain were interpreted in favor of this model. However, these biochemical data are also

consistent with a very different model of Fn1 fibrillogenesis (**Sup. Fig. 8**), one that was

proposed by (Tomasini-Johansson et al., 2006) and discussed below.

A canonical model poses that Fn1 fibrillogenesis entails intermolecular Fn1-Fn1

interactions mediated by the 70 kDa N-terminal domain of Fn1 (Hocking et al., 1994; Zhong et

al., 1998), summarized in (Singh et al., 2010). These interactions were hypothesized to be

important for linking extended Fn1 dimers into fibrils (e.g., **Sup. Fig. 8A**). It was stipulated that

FUD blocks Fn1 fibrillogenesis because of its ability to compete with Fn1-Fn1 intermolecular

interactions and disassemble fibrils by inhibiting intermolecular interactions between Fn1

dimers (**Sup**. **Fig. 8D**). However, these biochemical and cell biological experiments are also

consistent with a different model, wherein Fn1 fibrillogenesis is mediated by the binding of the

70 kDa N-terminal domain to large apparent molecular mass cell-surface complexes (LAMMs)

(Tomasini-Johansson et al., 2006; Zhang and Mosher, 1996) (**Sup**. **Fig. 8E**), and that FUD can

disrupt fibrillogenesis by disrupting the interactions between the 70 kDa Fn1 assembly domain

and LAMMs (**Sup**. **Fig. 8E**). Biochemical evidence supporting this model is multifold: 1)

isolated 70 kDa N-terminal domain can bind to Fn1-null cells in the absence of Fn1, forming

short linear arrays; this binding is time- and dose-dependent, and is detectable with as little as

5 nM of 70 kDa N-terminal segment of Fn1 (Tomasini-Johansson et al., 2006); 2) Incubation of

the 70 kDa domain of Fn1 with cells precipitates LAMMs in pull-down assays (Zhang and

Mosher, 1996); 3) LAMMs contain trypsin-sensitive protein(s) (Zhang and Mosher, 1996); and

4) FUD inhibits the binding of 70 kDa N-terminal fragment to Fn1-null cells (Tomasini-

Johansson et al., 2006). These findings suggest that Fn1 nanodomains in fibrils can be held

together by the interactions between the 70 kDa N-terminal assembly domain of Fn1 and

LAMMs (**Sup. Fig. 8F1**). However, at this time, we cannot rule out a possibility that Fn1

nanodomains are held by an extended Fn1 molecule via N-N-terminal interactions (**Sup. Fig.**

8F2), or that a combination of the latter two models may be true.

Our model in which Fn1 fibrils are composed of periodically-spaced nanodomains

containing multiple Fn1 dimers is also supported by electron microscopy and atomic force

microscopy studies done in the past: 1) periodical staining of Fn1 fibrils was observed in EM

with polyclonal antibodies raised to entire human plasma Fn1 (Furcht et al., 1980a); 2) grapes-

on-a-vine appearance of immuno-gold complexes detecting plasma Fn1 in fibrils in electron

micrographs (Peters et al., 1990); 3) bulbous appearance of Fn1 fibrils in cryo-scanning

transmission electron tomography images (Lansky et al., 2019); and 4) beaded appearance of

Fn1 fibrils detected by atomic force microscopy (Gudzenko and Franz, 2015). The latter two

methods did not rely on antibody staining to detect Fn1 nanodomains, supporting the notion

that the beaded architecture of Fn1 fibrils is not an artifact of antibody labeling.

Together with our data, these experiments suggest a new model of Fn1 fibrillogenesis

(**Fig. 10E**). In this model, Fn1 nanodomains containing multiple Fn1 dimers form at cell

periphery. These nanodomains may be similar to focal complexes or adhesion-related particles

seen by cryo-electron tomography by (Patla et al., 2010). The centripetal translocation of Fn1

nanodomains is coordinated with their organization into linear arrays, giving rise to focal

adhesions composed of ~3-5 Fn1 nanodomains and long Fn1 fibrillar adhesions composed of

dozens of nanodomains. Fn1 fibrils generated in this process are then incorporated into Fn1

ECM (Mautner and Hynes, 1977; Pankov et al., 2019; Sivakumar et al., 2006). Two pieces of

evidence suggest that Fn1 nanodomains in fibrils are linked: 1) the preservation of the linear

organization and the nanoarchitecture of Fn1 fibrils after the treatment of cells with DOC which

dissolves cell membranes, and 2) the presence of fibrous material between immunogold

densities in electron micrographs (Chen et al., 1997; Dzamba and Peters, 1991; Furcht et al.,

1980a; Furcht et al., 1980b; Furcht et al., 1980c; Lansky et al., 2019; Peters et al., 1990). The

sparsity of Fn1 localizations between Fn1 nanodomains in fibrils suggests that molecules other

than or in addition to Fn1 participate in the potential linking of Fn1 nanodomains into periodical

nanodomain arrays.

The beaded architecture of Fn1 ECM has important implications for the mechanisms of

ECM formation, remodeling, and signal transduction. The tensile strength of knotted strings is

significantly lower than that of strings with uniformly-aligned fibers (Arai et al., 1999; Saitta et

al., 1999), thus the beaded architecture of Fn1 fibrils may facilitate their rupture under strain

(Ohashi et al., 1999). The non-uniform, nanodomain architecture of Fn1 may facilitate the

accessibility of Fn1 fibrils to matrix metalloproteases. In this model, degradation of Fn1 fibrils

by metalloproteases may be accomplished by the cleavage of LAMMs between Fn1

nanodomains facilitating dynamic ECM remodeling. Finally, Fn1 is known to bind growth

factors (Martino and Hubbell, 2010; Saunders and Schwarzbauer, 2019; Wijelath et al., 2002), and cell adhesion to ECM is known to orchestrate growth factor signaling (Hynes, 2009). Thus, Fn1 nanodomains could serve as platforms for the binding and presentation of concentrated packets of growth factors to cells, the organization of Fn1 nanodomains into closely-spaced arrays could further facilitate clustering and signaling by growth factor receptors.

Acknowledgments

We thank Richard Hynes and Nathan Astrof for insightful discussions and careful reading of the manuscript, Sydney Astrof for encouragement and help with data entry, Patrick Murphy for endothelial cells, Richard Hynes for the gift of 297.1 antibody, Tung Chan for help with setting up Western Blotting using ProteinSimple, and Jonas Ries for advice that helped us improve SMLM resolution of our imaging, and for his help with adopting the SMAP software. Epitope mapping of 297.1 antibody was performed by PEPperPRINT GmbH, Heidelberg, Germany.

Sources of Funding This work was supported by the funding from the National Heart, Lung, and Blood Institute of the NIH R01 HL103920, R01 HL134935, R01 HL158049, American Heart Association Transformative Project Award 20TPA35490074 to SA, and by the NIH Office of the Director R21 OD025323-01 to SA, by the pre-doctoral fellowship F31HL151046 to BEA, by the National Institute of General Medicine R35GM122505 to AK, by the National Institute of Arthritis and Musculoskeletal and Skin Diseases R01 AR073236 to JES, by the Faculty Seed Grant from the Center for Engineering MechanoBiology (CEMB), an NSF Science and Technology Center, under grant agreement CMMI: 15-48571. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

Materials and Methods

Generation Fn1-fluorescent protein targeting constructs Sequences of monomeric (m)

green fluorescent protein (GFP), mNeonGreen, mScarlet-I, and tdTomato were obtained from

FPbase (https://www.fpbase.org). The sequence encoding one of the above fluorescent

proteins (FPs) was knocked into the Fn1 locus following the last coding exon of mouse Fn1,

and separated from the last coding amino acid by a flexible, proline-rich linker, PPPELLGGP

(Snitkovsky and Young, 1998). Targeting was achieved by CRISPR/Cas9 (Ran et al., 2013).

The sequence of the guide RNA was chosen and off-target sites were identified using

GuideScan and Off-Spotter software (Perez et al., 2017; Pliatsika and Rigoutsos, 2015). The

guide RNA (gRNA) sequence 5'-AGC GGC ATG AAG CAC TCA AT-3' targeting the last

coding exon of *Fn1* was subcloned downstream the U6 promoter into the PX459 vector

(Addgene, cat # 62988) encoding the Cas9-2A-Puromycin cassette (Ran et al., 2013). The

homology-directed repair (HDR) template was constructed using pBS-KS vector (Sup. Fig. 1a).

The sequence of the last coding exon of *Fn1* 5'-

AACGTAAATTGCCCCATTGAGTGCTTCATGCCGCTAGATGTGCAAGCTGACAGAGACGAT

TCTCGAGAG-3' was modified to 5'-

AACGTAAATTGCCCCAT<u>c</u>GA<u>a</u>TGCTTCATGCCGCTAGATGTGCAAGCTGACAGAGACGATT

CTCGAGAG-3' in the HDR template by introducing silent mutations (underlined) to prevent

targeting of the template by the gRNA. Homology arm 1 contained 677 bp encoding exon #45,

intron, and a portion of the last exon (#46), of the transcript *ENSMUST00000055226.12*.

Homology arm 2 encoded 1739 bp immediately downstream of the *Fn1* termination codon and

included the unmodified 3'UTR of Fn1. Knockin Fn1^{mEGFP/+} mice were generated by

Biocytogen using the same HDR construct and a longer gRNA, 5'-<u>T</u>AG CGG CAT GAA GCA

CTC AAT <u>G</u>G-3',

targeting the same sequence in the last coding exon (differences between the two gRNAs are

underlined). Targeting was confirmed by sequencing and Southern Blotting

(Sub. Fig. 1b). 500 bp around each of the top ten predicted off-target sites were sequenced

and no mutations were found in the founder mice. Mice containing correctly-targeted *Fn1* locus

were used to establish living colonies of Fn1$^{mEGFP/mEGFP}$ animals. Wild-type, Fn1$^{mEGFP/+}$, and

Fn1$^{mEGFP/mEGFP}$ mice were genotyped using the following primers Fn1-WT-Fwd 5'-

TCCCCGAAACACACACTTTTGGT-3', Fn1-WT-Rev 5'

GTCACCCTGTTCTGCTTCAGGGTTT-3', and Fn1GFP-Rev 5'-

GACCCGCGCCGAGGTGAAG-3'. Wild type band is Fn1-WT-Fwd and Fn1-WT-Rev give rise

to 372 bp for the wild-type allele; Fn1-WT-Fwd and Fn1GFP-Rev primer located in the GFP

sequence give rise to 512 bp if the targeted allele is present. Mice were housed in an

AAALAC-approved barrier facility. All experimental procedures were approved by the

Institutional Animal Care and Use Committee of Rutgers University and conducted in

accordance with the Federal guidelines for the humane care of animals.

Cell Culture

Table M1. Media

	Cells	Composition of Medium
Complete medium	MEFs	High-glucose Dulbecco's Modified Eagle Medium (DMEM, Corning, cat # 10-013-CV) supplemented with 10% v/v fetal bovine serum (Gemini Biosciences, cat # 100-106), 1% v/v penicillin/streptomycin solution (GE Healthcare, cat #SV30010), 1% v/v L-glutamine (Gibco, cat # 35050-061).

Imaging medium	MEFs	FluoroBrite DMEM (Thermo Fisher Scientific, catalog # A1896701) supplemented with 2% v/v fetal bovine serum (Gemini Biosciences 100-106), 1% v/v penicillin/streptomycin solution (GE Healthcare, SV30010), 1% v/v L-glutamine (Gibco 35050-061), pH8.14.
McCoy's 5a	Nup96-mEGFP U2OS cells	McCoy's 5a medium containing 1xGlutaMax (ThermoFisher, cat #35050061), 10% v/v fetal bovine serum (Gemini Biosciences, cat # 100-106), and 1x penicillin/streptomycin (GE Healthcare, cat #SV30010).

Cells Wild-type mouse embryonic fibroblasts (MEFs) were isolated from embryonic day (E)

13.5 embryos derived from the C57BL/6J strain (Jackson Labs, stock # 664) according to

established protocols (Behringer et al., 2014) and cultured in complete medium. For

propagation, MEFs were plated in flasks pre-coated with 0.1% gelatin solution and grown in

complete medium at 37°C, 5% CO_2.

$Fn1^{flox/+}$;$Rosa^{mTmG/+}$ MEFs were isolated from E13.5 embryos, as above, and express

membrane-bound tdTomato and wild-type Fn1 in the absence of Cre recombinase (Muzumdar

et al., 2007). Fn1-null MEFs were generated by treating $Fn1^{flox/-}$;$Rosa^{mTmG/+}$ cells with Ad-Cre-

IRED-GFP virus according to the manufacturer's recommendations (Vector Biolabs, #1710). In

this experiment, Cre-recombinase is expressed transiently and mediates site-specific

recombination between a pair of loxP sites flanking the 1st exon of Fn1 (Sakai et al., 2001) and

another pair of loxP sites flanking the STOP cassette in the mTmG reporter (Muzumdar et al.,

2007). Three days following infection with Ad-Cre-IRED-GFP, GFP+ cells were sorted resulting

in a pure population of Fn1-null MEFs (confirmed by Western Blotting, data not shown, and

immunofluorescence, **Sup. Fig. 12**). Fn-null MEFs were culture on gelatin-coated dishes in

complete medium.

Fn1-FP cells (except Fn1-mEGFP cells) were generated by CRISPR/Cas9 mutagenesis

of wild-type MEFs. CRISPR/Cas9 targeting was performed by transfecting wild-type MEFs

using the PX459 plasmid encoding Fn1 gRNA and the HDR template using lipofectamine

3000, as described (Ran et al., 2013). MEFs expressing Fn1-mEGFP proteins used for live

imaging and STORM were generated from homozygous E13.5 $Fn1^{mEGFP/mEGFP}$ or, when noted

from $Fn1^{mEGFP/+}$ embryos. $Fn1^{mEGFP/mEGFP}$ MEFs were used to quantify Fn1 molecule number in

nanodomains, measure nanodomain spacing, and nanodomain diameter, as well as in

experiments to label both the N- and the C-termini of Fn1. For live imaging, $Fn1^{mEGFP/+}$ were

used. Fibronectins from $Fn1^{mEGFP/mEGFP}$ and $Fn1^{mEGFP/+}$ cells behave equivalently in Fn1 matrix

assembly assays (data not shown).

Nup96-mEGFP cells ((Thevathasan et al., 2019), clone #195, Cell Line Services, CLS,

clsgmbh.de, catalogue no. 300174)) were cultured according to the vendor's specifications in

McCoy's 5a medium containing 1xGlutaMax (ThermoFisher, cat #35050061), 10% v/v fetal

bovine serum (Gemini Biosciences, cat # 100-106), and 1x penicillin/streptomycin (GE

Healthcare, cat #SV30010).

Reagents and buffers

FUD and III-11C peptides were generated as described (Sottile and Chandler, 2005; Tomasini-

Johansson et al., 2001) and stored in PBS at -80° C. 4% DOC solution was prepared by

dissolving 0.4 g deoxycholate salt (Sigma, catalog # D6750) in 10 ml of imaging medium; the

solution was then vortexed and filter sterilized. The pH of the final solution was 8.01. 16%

paraformaldehyde (PFA) (Electron microscopy Sciences; catalog # 50-980-487) was diluted in

1x PBS to prepare 4% PFA. The 4% PFA solution was aliquoted into 1 ml microfuge tubes,

stored at -80° C, and thawed at 37° C immediately before use.

1X phosphate buffered saline pH 7.5 (PBS) was prepared from 10X PBS (VWR, catalog #

76180-740). PBST was prepared using Triton X-100 (Sigma-Aldrich, catalog # T-8787) and

contained 0.1% Triton for all stainings except those involving Nup96-mGEFP cells, as detailed

below. Blocking buffer was prepared by adding 10% Donkey serum (Sigma-Aldrich, catalog #

D9663) to 1X PBST; 5 mg/ml stock of DAPI (Fisher Scientific, cat #D3571) was prepared in

water and used at 1:300 dilution. Stain Buffer (cat # 554656 BD Pharmingen) was used for

antibody dilutions and washing of cells that were stained without permeabilization.

Hoechst 33342 Trihydrochloride (Thermo Fisher, catalog # H1399, 10mg/ml) was used for

labelling live MEFs at 1:300 dilution. In live MEFs, F-actin was labelled using SiR actin (cat#

CY-SC001 used at 1 μM final concentration). mCardinal-Lifeact-7 was a gift from Michael

Davidson (Addgene plasmid # 54663 ; http://n2t.net/addgene:54663 ; RRID:Addgene_54663).

Vectashield antifade mounting medium (Vectorlabs, catalog # H-1000) was used for cover

slipping.

For STORM imaging, we used 25 mm high precision glass cover slips #1.5H (Marienfeld ref#

0117650, obtained from Azer Scientific, PA, USA, cat # ES0117650) without coating. Prior to

their use, glass cover slips were cleaned using concentrated nitric acid, washed in water, air

dried, and autoclaved, as described in (Kaech and Banker, 2006). Clean coverslips were

stored in sealed 6-well plates for no longer than a week prior to their use.

GLOX/BME STORM buffer contained 50 mM Tris-HCl (Fisher Scientific, catalog # T-395-1),

pH 8.0, 10 mM NaCl (Sigma- Aldrich, catalog # S-7653), 10% glucose (Sigma- Aldrich, catalog

G8270), 0.5 mg/ml glucose oxidase (Sigma- Aldrich, catalog # G2133), 40 μg/ml catalase

(Sigma- Aldrich, catalog # C40), and 143 mM β-mercaptoethanol (βME, Sigma-Aldrich, cat#

444203) (Thevathasan et al., 2019). Stocks of enzyme solutions were prepared and stored at -

20ºC as described in (Jimenez et al., 2020). GLOX/BME buffer was used for STORM imaging

of cells plated on coverslips.

GLOX/MEA STORM buffer was used for STORM imaging of cells plated in Ibidi glass bottom

8-well chambers (catalog # 80827). This buffer was prepared as above, but instead of βME, it

contained 50 mM mercaptoethylamine (MEA, Sigma-Aldrich, catalog # 30070) (Jimenez et al.,

2020). For double-color STORM imaging, we used the SMART Kit buffer (Abbelight).

Antibodies

All primary antibodies were checked for specificity on cells that were genetically-null for the

antigen (e.g., **Sup. Fig. 12**) and tissues: Fn1-null tissue sections obtained from Fn1-null

embryos were used to assay the specificity of each of the anti-Fn1 antibodies; Tissues isolated

from GFP-null, Itga5-null, and mCherry-null embryos were used to authenticate the specificity

of anti-GFP, anti-Itga5, and anti-mCherry antibodies. For each of the antibodies, staining of

control tissues resulted in no more fluorescent signal than the background fluorescence

produced by the use of secondary antibodies only. Rabbit polyclonal antibody R457 was raised

against the 70 kDa N-terminal domain of Fn1 (Aguirre et al., 1994; Sechler et al., 2001) and

rabbit polyclonal R184 was raised against the first six type III repeats of Fn1 (Raitman et al.,

2018).

Table M2. Antibodies

Primary Antibodies (1° Ab)	Specificity	Source/ reference

Abcam recombinant rabbit monoclonal	Epitope recognized by this antibody is located within 301 aa between amino acids 1350- 1651 of mouse Fn1, transcript variant 2, mRNA, RefSeq NM_001276408	Abcam, cat # 199056
297.1 rabbit polyclonal	Raised to rat plasma Fn1. Recognizes multiple epitopes, see **Sup. Fig. 4**	Richard Hynes lab (Rickelt and Hynes, 2018)
R457 rabbit polyclonal	70 kDa N-terminal domain	Jean Schwarzbauer's lab (Aguirre et al., 1994; Sechler et al., 2001)
R184 rabbit polyclonal	First six type III repeats of Fn1	Jean Schwarzbauer's lab (Raitman et al., 2018)
3E2 mouse monoclonal	EIIIA	Sigma, cat #SAB4200784-100UL
GFP, chicken polyclonal	GFP	Aves, cat# GFP-1010;
GFP, rabbit polyclonal	GFP	MBL International; cat #598
Itgα5, rat monoclonal	Integrin alpha 5	BD biosciences, cat # 553319
mCherry, rabbit polyclonal	Cherry; tdTomato is composed of two mCherry	Abcam, cat # ab167453

	molecules in tandem (Shaner et al., 2005)	

Secondary Antibodies (2° Abs)	Source, catalog #, concentration	Dilution
AffiniPure F(ab')₂ Fragment Donkey Anti-Rabbit IgG (H+L)	Jackson Immunoresearch, cat # 711-606-152, 1.5 mg/ml	1:300
Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 555	ThermoFisher Scientific, A-31570, 2 mg/ml	1:300
Cy™3 AffiniPure F(ab')₂ Fragment Donkey Anti-Rat IgG (H+L)	Jackson Immunoresearch, 712-166-150, 1.5mg/ml	1:300
Alexa Fluor® 488 AffiniPure F(ab')₂ Fragment Donkey Anti-Chicken IgY (IgG) (H+L)	Jackson Immunoresearch, 703-546-155, 1.5mg/ml	1:300
Alexa Fluor® 647 AffiniPure F(ab')₂ Fragment Donkey Anti-Chicken IgY (IgG) (H+L)	Jackson Immunoresearch, 703-606-155, 1.5mg/ml	1:300

CF680 Donkey anti-rabbit, highly cross-adsorbed	Biotium, cat # 20418-50ul), 2mg/ml	1:200

Epitope mapping of rabbit 297.1 polyclonal antibody. Epitopes recognized by the

polyclonal 297.1 antibody were mapped by generating custom overlapping peptide arrays

(PEPperPRINT GmbH, Heidelberg). Fn1 protein sequence including alternatively-spliced

exons was encoded by 15-amino acid peptides with a peptide-peptide overlap of 13 amino

acids. The resulting Fn1 peptide microarrays contained 1,239 different peptides printed in

duplicate (2,478 peptide spots), and were framed by additional HA-tag (YPYDVPDYAG, 106

spots) control peptides. HA-tag peptide was used to monitor array quality and served as a

positive control for clean antibody binding detected with anti-HA-tag antibody (mouse

monoclonal anti-HA DyLight800 used at 0.5 µg/ml concentration). Prior to staining, arrays were

incubated with blocking buffer (Rockland, cat # MB-070) for 30 min at RT. To measure

background antibody binding, arrays were first incubated with secondary goat anti-rabbit IgG

(Fc) DyLight680 at 0.2 µg/ml concentration diluted in PBS, pH 7.4, containing 0.05% Tween 20

and 10% blocking buffer, and imaged. To determine specific 297.1 antibody binding, arrays

were incubated for 16 hours with shaking at 4°C with two dilutions of 297.1 antibody (1:300

and 1:1000). Antibody dilutions were made in PBS, pH 7.4 containing 0.05% Tween 20 and

10% blocking buffer (Rockland, cat # MB-070). Arrays were then washed with PBS, pH 7.4

containing 0.05% Tween 20 and incubated with goat anti-rabbit IgG (Fc) DyLight680 (0.2

µg/ml) for 45 min at RT. Arrays were then washed and imaged using LI-COR Odyssey Imaging

System; scanning offset 0.65 mm, resolution 21 µm, scanning intensities of 7/7 (red = 700

nm/green = 800 nm). Quantification of spot intensities and peptide annotation were done with

PepSlide® Analyzer.

Embryo Staining E9.5 embryos were isolated either from matings of wild-type C57BL6/J mice

or from mating Fn1$^{mEGFP/mEGFP}$ mice with wild-type mice to obtain Fn1$^{mEGFP/+}$ embryos.

Embryos were fixed using cold 4% PFA overnight at 4°C, washed 3 x 5 min in 1xPBS, and

blocked overnight at 4°C in blocking buffer containing PBS, 0.1% Triton-X, and 10% donkey

serum. Embryos were stained either with Abcam monoclonal anti-Fn1 antibodies at 1:300

dilution in blocking buffer or with both Abcam monoclonal anti-Fn1 antibodies and anti-GFP

antibodies at 1:300 dilution. The staining procedure was executed exactly as described in

(Ramirez and Astrof, 2020).

Analysis of Fn1 matrix assembly Matrix Assembly was performed according to the

established protocols (Wierzbicka-Patynowski et al., 2004). MEFs were plated in 6-well dishes

(9 cm^2 growth area) at a density of 2×10^5 cells per well for 48 h, in complete medium and

incubated under sterile conditions at 37°C, 5% CO2. Cells were washed twice with ice cold

PBS (supplemented with Mg^{2+} and Ca^{2+}), scraped with a cell scraper and lysed with either 500

µl RIPA lysis buffer pH 8.0 (50 mM Tris-Cl, 150 mM NaCl, 2 mM EDTA, 1% v/v NP-40, 0.5%

w/v sodium deoxycholate, 0.1% w/v SDS, 1X protease inhibitor cocktail (Cell Signaling

Technology, cat # 5871), or DOC lysis buffer, pH 8.8 (20 mM Tris-Cl, 2 mM EDTA, 2% w/v

sodium deoxycholate, 1X protease inhibitor cocktail (Cell Signaling Technology, 5871).

Extracts were carefully transferred to Eppendorf tubes containing 1 µl (250 units) Benzonase®

Nuclease (Sigma-Aldrich, E1014), mixed by inverting a few times and incubated at 37 °C for

15 mins. The samples were then centrifuged at 16,000 × g for 15 min at 4 °C. For cells lysed

with DOC lysis buffer, the supernatant containing DOC-soluble material was carefully

removed, and the pellet containing the DOC-insoluble material was resuspended in 100 µl

SDS solubilization buffer, pH8.8 (20 mM Tris-Cl, 2 mM EDTA, 1% w/v SDS, 1X protease

inhibitor cocktail (Cell Signaling Technology, 5871). The DOC-insoluble pellet was thoroughly

dissolved by heating the sample to 95 °C and vortexing. All samples were aliquoted and stored

at -80 °C until further use. Prior to quantification of Fn1 in the samples, the total protein

concentration of the RIPA and DOC lysates was determined using the BCA protein assay

(Pierce™ BCA Protein Assay Kit, 23225). Fn1 and Fn1-FP fusion proteins were resolved using

66-440 kDa Wes separation module (ProteinSimple, SM-W007). Primary antibodies were used

at the following dilutions: anti-total Fn1 – 1:1000 (Abcam, ab199056), anti-GFP – 1:1000

(Roche, 11814460001), anti-mCherry – 1:1000 (Abcam, 167453). Primary antibodies were

detected using horseradish peroxidase-conjugated secondary antibodies (anti-Rabbit

Detection Module ProteinSimple, DM-001), and chemiluminescence was quantified using the

Compass for SW software (v3.1.8). Prior to running experimental samples, care was taken to

optimize the dilutions of lysates to be within the linear range of the detection.

Coating of coverslips with different ECM proteins #1.5 round glass coverslips (Electron

Microscopy Sciences. Catalog # 72230-01) were coated with the following ECM proteins when

noted: gelatin (Sigma Aldrich, catalog # G2500) (0.1% (w/v) of gelatin was prepared in Milli-Q

water and autoclaved to dissolve), vitronectin (Sigma Aldrich, catalog # SRP3186; stock

solution was prepared as 200µg/ml in 0.1% BSA and water) and laminin (R&D systems,

catalog # 3400-010-02, stock 1 mg/ml was pipetted into 10ul aliquots and stored at -80^{O}C). To

coat with gelatin, glass surfaces were incubated with the 0.1% gelatin solution for 5 min at

room temperature (RT). To coat with vitronectin or laminin, glass surfaces were incubated at

37° C for 1 hr in 20 µg/ml of either vitronectin or laminin, excess liquid was removed, cover

slips were rinsed once with 1X PBS, and blocked with 10 µg/ml heat denatured BSA for 30 min

before plating cells (Lu et al., 2020).

MEFs plated on different substrata MEFs were grown either on #1.5 round glass coverslips

in 24-well dishes or in 8-well glass Ibidi dishes depending on the experiment for the times

indicated in figure legends. MEFs were then rinsed with 1X PBS (warmed to 37^0 C) for 5 min,

fixed with freshly thawed 4% PFA pre-warmed to 37^0 C for 20 min, and washed three times

with 1X PBS (warmed to 37^0 C) with mild shaking. All subsequent washing steps were done

with shaking. For permeabilization, cells were washed once in 1X PBS containing 0.1% Triton-

X 100 (PBST). Blocking was done for 30 min in 10% Donkey serum prepared in PBST

(blocking solution). After blocking, cells were incubated in primary antibodies diluted in

blocking solution overnight at 4^0 C, as specified in **Table M2**. This was followed by 3 washes in

PBST for 10 min each. Cells were then incubated with secondary antibodies diluted in PBST

for 60 min at RT. Finally, cells were washed three times with PBST for 10 min each. DAPI

(1:300) was added to the second wash. Cells were mounted using Vectashield.

Hydrogels

Methacrylated Alginate Synthesis: Methacrylated alginate (MeAlg) was synthesized

according to a previously established protocol (Khetan et al., 2013). In brief, alginic acid sodium

salt from brown algae (Sigma-Aldrich, USA) (3% w/v) was fully dissolved in Dulbecco's

phosphate buffered saline (dPBS, Sigma-Aldrich, USA). Then, methacrylic anhydride (Sigma-

Aldrich, USA) (8% v/v) was added drop-wise to the alginate solution and stirred for 12 h at 4°C,

using 2M NaOH (Sigma-Aldrich, USA) to ensure that the pH remained between 8 and 9 for the

duration of the reaction. The resulting solution was passed through filter paper (GE Whatman)

and poured into Spectra/Por dialysis membrane with a 6–8 kDa molecular weight cutoff (Fischer

Scientific) and kept in DIW under stirring for 7 days to eliminate the unreacted MA and salts.

Dialyzed solution was then freeze-dried for 4 days to obtain MeAlg foam.

Fabrication of the Hydrogel Substrates: MeAlg substrates were fabricated using a previously established protocol (Guvendiren and Burdick, 2012). Briefly, petri dishes with glass bottoms were treated with UV/ozone (UVO) for 30 minutes, immediately followed by a coating of 3-(trimethoxysilyl)propyl methacrylate (TMS) (Sigma-Aldrich, USA) to methacrylate the glass surfaces (Guvendiren et al., 2009). The dishes were left in a desiccator overnight. The hydrogels were fabricated using Michael-type addition polymerization. First, 2-hydroxy-4'-(2-hydroxyethoxy)-2-methylpropiophenone (I2959) (Sigma-Aldrich, USA), a photoinitiatior (0.5% w/v) was completely dissolved in Dulbecco's PBS (dPBS), followed by the lyophilized MeAlg (3% w/v) synthesized previously. This was kept at room temperature until a clear solution was achieved. Crosslinking occurs with the introduction of DL-Dithiothreitol (DTT) (Sigma-Aldrich, USA) to the solution, along with 0.2M triethanolamine (Sigma-Aldrich, USA) at pH 10. To form 3kPa and 12 kPa gels, 20% and 30% (w/v) DTT are used, respectively. To promote cell adhesion, GRGDSPC peptide (1% w/v) (Genscript) was added to the solution. After all contents were thoroughly mixed, 5μL of MeAlg solution was pipetted onto the surface of the dish before being covered with a glass coverslip in order to create gels less than 30μm thick. These were left at room temperature for an hour to crosslink before being submerged in dPBS to remove the coverslip.

Atomic Force Microscopy For stiffness measurements, hydrogel samples were submerged in PBS and placed in a Dimension Icon AFM with ScanAsyst (Bruker). Using the PeakForce-QNM mode, hydrogel samples were indented using an MLCT-Bio probe tip with pyramidal geometry (Bruker, CA) and a nominal spring constant of 0.03 N/m, checked by thermal calibration.

Treatment of cells with Deoxycholate (DOC) 10^4 Fn1$^{mEGFP/+}$ MEFs were plated for 48 hrs in 8-well glass bottom Ibidi dishes in complete medium and incubated at 37° C and 5% CO_2. Two hours before imaging SiR-actin was added at 1 μM final concentration. SiR-actin contains a

far-red dye, silicon rhodamine, conjugated to jasplakinolide that labels F-actin in live and fixed

cells (Lukinavicius et al., 2014). Just before imaging, complete medium was replaced by 150 μl

imaging medium containing 33 μg/ml of Hoechst 33342. Positions were marked in each well

and live imaging was initiated at 37^{o}C and 5 % CO_2 humidified chamber. After 15 min, 150 μl

4% DOC solution prepared in imaging medium containing 33 μg/ml Hoechst was added to the

experimental well (final pH 8.01) and 150 μl imaging medium containing 33 μg/ml Hoechst but

without DOC was added to the control well. Cells were imaged at 50 sec intervals until F-actin

and DNA disappeared (see Movie 4). The medium was then removed, cells were rinsed for 1

min with 1X PBS pre-warmed to 37^{o}C, fixed with 4% PFA pre-warmed to 37^{o}C. For staining,

cells were permeabilized and blocked as above, and then incubated with Abcam monoclonal

anti-Fn1 antibody diluted 1:300 dilution in the blocking solution at 4^{o}C overnight. Primary

antibodies were detected with anti-rabbit secondary antibodies conjugated to AlexaFluor647.

Enhanced-resolution imaging was used to image the DOC-treated fibrils, as described below.

Confocal settings for enhanced resolution imaging Confocal images of fixed samples were

recorded using Nikon A1-HD25 inverted confocal microscope equipped with CFI

Apochromat TIRF 100xC Oil objective with the pinhole set to 0.8 Airy units, and imaged

through 2 – 4 microns with step size of 0.125 μm - 0.15 μm at a sampling of 40 nm per pixel

and 180 nm optical resolution. Deconvolution was done using Nikon 3D deconvolution

software (v5.11.01). Airyscan imaging was performed using Zeiss LSM 880 fitted with a 32

array AiryScan GaAsP-PMT detector and the Plan Apochromat 63X Oil (NA 1.4) objective.

Deconvolution and pixel reassignment were done using Zeiss LSM software.

Quantification of effective labeling efficiency and the number of Fn1 molecules in Fn1

nanodomains. To quantify the number of Fn1 molecules per nanodomain and to measure

effective labeling efficiency, we used Nup96-mEGFP U2OS cells as a reference cell line and

SMAP software (Ries, 2020; Thevathasan et al., 2019). In Nup96-mEGFP cells, two copies of

the nucleoporin Nup96 gene are tagged with the monomeric enhanced green fluorescent

protein (mEGFP) at the C-terminus of the Nup96 gene, generating Nup96-mEGFP fusion

protein (Thevathasan et al., 2019). The stoichiometry of Nup96-mEGFP in nucleopores is well

characterized, and careful measurements, imaging methodology, and software have been

developed (Diekmann et al., 2020; Ries, 2020; Thevathasan et al., 2019). Together, these

tools allow the use of Nup96-mEGFP cell line as a reference to assess the quality of SMLM

imaging protocol, measure effective labeling efficiency, and determine the number of Fn1-

mEGFP molecules in Fn1 nanodomains.

Nup96-mEGFP cells (clone #195, Cell Line Services, CLS, clsgmbh.de, catalogue no. 300174)

were cultured according to the vendor's specifications in McCoy's 5a medium. Fn1-mEGFP

MEFs were cultured in complete DMEM as described above. 3×10^5 Nup96-mEGFP cells and

5×10^4 Fn1-mEGFP cells were plated in their specified culture medium on 25 mm high precision

glass cover slips #1.5H (Marienfeld ref# 0117650, obtained from Azer Scientific, PA, USA, cat

ES0117650) positioned in 6-well plates (Corning, cat # 353046). Prior to their use, glass

cover slips were cleaned using concentrated nitric acid, washed in water, and autoclaved, as

described in (Kaech and Banker, 2006). Coverslips were used without coating. 24 hours after

plating, cells were fixed and stained as described in (Thevathasan et al., 2019) with minor

modifications. Cover slips with Nup96-mEGFP cells and Fn1-mEGFP cells were handled

contemporaneously in pairs at each step. For fixation, permeabilization and washing,

coverslips were kept in 6-well plates. Cells were fixed in PBS containing 2.4% PFA for 20 min

at RT. PFA was aspirated and cells were incubated with 100mM NH_4Cl in PBS for 5 min at RT,

and then washed 3 x 5 min in PBS with agitation. Cells were permeabilized for 20 min at RT

using PBS containing either 0.2% Triton-X for Nup96-mEGFP cells or 0.1% Triton-X for Fn1-

mEGFP. Cells were then washed in PBS 3 x 5 min at RT, and washed once more for 5 min in

PBS containing 0.1% Triton-X, and then either used immediately for staining, or stored at 4°C

in PBS containing 0.1% Triton-X (PBST) until further use. For each experiment, sufficient

amounts of each solution were prepared such that Nup96- mEGFP and Fn1-mEGFP cells

were incubated with the same mixtures. Solutions containing blocking reagents and antibodies

were spun for 5 min at top speed using tabletop centrifuges to get rid of particulates.

Humidified chambers were prepared from empty pipet tip boxes with water placed in the lower

chamber and parafilm partially covering the surface of the tip rack. Drops of solutions were

placed on the parafilm, and cells were incubated with various solutions by inverting coverslips

on top of the droplets. To block non-specific antibody binding, cells were first incubated with a

blocking solution (10% donkey serum in PBST) for 30 min at RT. Cover slips were then gently

lifted, drained, and incubated with anti-GFP antibody (Aves lab, cat# GFP-1010) diluted 1:100

in the blocking solution overnight at 4°C. Coverslips were then placed into 6-well plates and

washed 3 x 5 min in PBST with agitation. Cells were then incubated with Alexa-647-conjugated

anti-chicken F(ab)$_2$ (Jackson ImmunoResearch 703-606-155) diluted 1:300 in the blocking

solution for 4 hours at RT, and then washed 3 x 5 min in PBST with agitation. Stained

coverslips were stored at 4°C until imaging.

Preparation of cover slips for SMLM imaging Coverslips with plated cells were rinsed with

GLOX/BME buffer and mounted onto single cavity glass slides (VWR, cat #10118-600), pre-

filled with 80 μl GLOX/BME buffer immediately prior to the placement of coverslip. Excess

buffer was fully adsorbed from the sides and the top of the coverslip with Whatman paper,

taking care to keep the coverslip centered on top of the cavity. Coverslips were sealed onto the

cavity slide by pipetting Acid-Free Elmer's No-Wrinkle Rubber Cement (Amazon.com,

https://www.amazon.com/Elmers-No-Wrinkle-Rubber-Cement-Acid-

Free/dp/B014JUDMBA/ref=sr_1_2?keywords=Elmer%27s+No-

Wrinkle+Rubber+Cement%2C+Acid-Free&qid=1637633240&sr=8-2) around the edge, and

allowing the rubber cement to cure for ~15 min. After imaging, rubber cement was gently

peeled off, and the slides were soaked in PBST for 5 min at RT to remove cover slips which

were then stored in 6-well plates filed with PBST containing 0.02% NaN_3 at 4°C until further

use. In this set up, the GLOX/BME buffer remained at pH 8 for at least 12 hours.

SMLM Imaging protocol to quantify the effective labeling efficiency and the number of

mEGFP molecules in Fn1-mEGFP nanodomains. For the following experiments, 2D imaging

was used to maximize the resolution in the plane of imaging. To minimize fluorophore

bleaching and maximize the number of collected photons we followed the protocol protocol

developed by (Diekmann et al., 2020). Stained Nup96-mEGFP and Fn1-mEGFP cells were

imaged in pairs using the same preparation of the GLOX/BME buffer and the same imaging

conditions (described below). For each independent experiment (n=4) and for each round of

measurements, images of Nup96-mEGFP and Fn1-mEGFP cells taken on the same day were

analyzed. SMLM was performed using Nikon A1-HD25 Ti2E microscope equipped with

motorized TIRF illumination, 125mW 640 nm solid-state laser, Perfect Focus, and CFI

Apochromat TIRF 100xC Oil objective with numerical aperture 1.49 (cat # MRD01905). All

images were acquired at the critical angle of incidence (57°) and recorded using a 512 x 512

EMCCD camera (Princeton Instruments), using 128 x 128 central region on the camera. Gain

was set to 3, multiplication gain amplifier was set to 20 MHz. Prior to acquisition, the centered

ROI (128 x 128) was bleached at 57° angle, and 10% laser power for 500 frames followed by

additional 500 frames at 20% laser power, with the exposure of 60 ms per frame. Images were

then acquired at 57° angle, 20% laser power, for 60,000 frames at 60 ms per frame. Images

were processed, rendered, and quantified using SMAP and Matlab version 2020a (Ries,

2020), as described (Diekmann et al., 2020) and the documentation found on

https://github.com/jries/SMAP. Settings for peak finding in the SMAP software were set

according to our camera's manufacture's specifications and were the following: EM was set to

"on", camera pixel size was set to 0.162 μm, EM gain was set to 300, e-/ADU conversion factor

was set to 2.35. Data was fitted using a Gaussian PSF model, the cutoff parameter for peak

finding was set to 2, and ROI size was set to 7 pixels. Localizations were then grouped and

rendered, using default parameters in SMAP. Drift correction was performed on rendered

localizations in timepoint blocks of 10 or 20, as recommended (Thevathasan et al., 2019) and

https://github.com/jries/SMAP. Following satisfactory drift correction judged by the overlapping

cross-correlations, localizations were filtered according to the recommended settings for Alexa-

647 fluorophore (Thevathasan et al., 2019), in brief, localizations with poor precision were

filtered out by limiting localization precision to 0 – 15 nm, out-of-focus localizations were

excluded by setting the PSF range to 0 – 150 nm in the x-y plane, poorly fitted localizations

were further filtered out by setting the LLrel parameter (relative log-likelihood) in SMAP to a

negative cut-off value leaving the majority of the peak intact, and the first 1000 frames were

excluded from the analyses. All the remaining grouped localization were rendered according to

$\sqrt{photons}$ and images were color-coded using LUTs set according to localization density.

Image resolution was measured using the Fourier Ring Correlation (FRC) method described

by (Nieuwenhuizen et al., 2013) and implemented in SMAP. Voronoi cluster discovery was

performed according to (Andronov et al., 2016a) and their algorithm implemented in SMAP.

DBSCAN cluster analysis was performed according to (Caetano et al., 2015; Ester et al., 1996)

implemented in SMAP; For DBSCAN, the minimum number of points in the neighborhood (k)

was set to 4, as recommended for all 2-dimentional data (Ester et al., 1996). The

neighborhood radius ε was set either to 14 nm (the average apparent radius of Fn1

nanodomains determined as described below) or automatically estimated by the DBSCAN

algorithm in SMAP (Ries, 2020).

NPC radius, effective labeling efficiency and the number of gourped localizations per

fluorophore were determined using the algorithm in SMAP following the published automated

workflow to segment and analyze nucleopore complexes (NPCs), using the parameters

recommended by (Thevathasan et al., 2019) without modifications. In brief, NPCs in focus

(mean fitted PFS size of each NPC was less then 145 nm) were automatically segmented

using circular ROI of 220 nm in diameter. The number of grouped localizations per NPC, NPC

radius, and effective labeling efficiency (ELE) were determined using published algorithms

implemented in SMAP without modifications, as described in (Diekmann et al., 2020;

Thevathasan et al., 2019) and the SMAP user manual "Analysis of NPCs using SMAP". Since

we used identical staining, imaging, and processing parameters for Nup96-mEGFP and Fn1-

mEGFP cells, the number of Fn1-mEGFP molecules per nanodomain was calibrated to the

number of grouped localizations per NUP96-mEGFP protein, as described (Thevathasan et al.,

2019). In brief, we first determined the number of merged localizations per NPC in the ROI

(L_{ref}). Each NPC contains 32 NUP96-mEGFP proteins, therefore, the number of merged

localizations per Nup96-mEGFP (N_{ref}) is $N_{ref} = L_{ref}/32$. To determine the number of Fn1-mEGFP

proteins per Fn1 nanodomain, we segmented Fn1 nanodomains manually using a circular ROI

of 60 nm in diameter and the number of merged localizations per ROI (L_t) was determined by

using the *countingStatistics plugin* in SMAP, as outlined in the SMAP manual "Analysis of

NPCs using SMAP". The number of Fn1-mEGFP proteins per nanodomains (N_t) is L_t / N_{ref}.

Since Fn1 is only secreted as a dimer, the number of Fn1 dimers per nanodomain is $N_t /2$.

Nanodomain periodicity in Fn1 fibrils stained by a variety of different antibodies was assayed

and quantified according to (Fruh et al., 2015). In brief, MEFs were stained using a variety of

antibodies at different dilutions, or using combinations of antibodies, as noted in **Figs. 5, 6, 8**,

according to the staining protocol described above and imaged using GLOX/BME buffer and

the exact imaging settings as described for imaging Nup96-mEGFP and Fn1-mEGFP cells

stained with anti-GFP antibodies (see above). Images were processed and rendered in SMAP

according to the parameters described above. Gaussian rendering of the localizations (min σ

Gaussian was set to 3nm) were saved as uncompressed tiff files and opened in Fiji version

2.1.0/1.53c, intensity line profiles along each fibril were generated and imported into Matlab

2021a. To assay the periodicity, we used the autocorrelation function implemented in the

Matlab's 2021a Econometrics toolbox and the criteria outlined in (Fruh et al., 2015). In brief,

the presence of at least four regularly-spaced peaks in the autocorrelation profile was

considered to reflect the periodical nature of Fn1 nanodomains, and the position of the first

autocorrelation maximum was taken as a quantitative measure of nanodomain periodicity, as

extensively discussed and computationally modelled in the Supplemental Data section of (Fruh

et al., 2015). Ljung-Box Q-test for residual autocorrelation was performed using Matlab's

2021a Econometrics toolbox to assess statistical significance of autocorrelation.

To determine the apparent diameter of Fn1 nanodomains, we imported fibril intensity

line profiles obtained from SMLM images of Fn1$^{mEGFP/mEGFP}$ MEFs stained for GFP into Matlab

2021a. To automate the analyses, we used Signal Processing toolbox in Matlab 2021a to fit

each intensity peak in the line profile with a Gaussian curve and calculate full width at half

height (FWHT) for each peak. Altogether, 1292 nanodomains in 27 long (> 1 μm in length)

fibrils from six cells and 3 independent experiments were assessed. We have also performed

this analysis manually on a subset of Fn1 nanodomains (n=248) in long fibrils by fitting a

Gaussian to an intensity profile of each individual nanodomain and calculating FWHT. The

results were the same.

Assembly of exogenously-added Fn1: Part I, Live Imaging 3-well culture insert (Ibidi, cat #

80366) was placed in the middle of the 35 mm glass-bottom dish (Ibidi, cat # 81158) and then

0.8 x 10^6 Fn1-tdTomato-expressing MEFs were plated surrounding the inserts and cultured for

24 hours to reach confluency and establish Fn1-tdTomato matrix. At a 24-hour time point, Fn1-

mEGFP-expressing MEFs were plated inside the inserts on glass without coating for 5 hours.

Prior to imaging, the culture medium was removed and replaced with the imaging medium.

Live imaging was performed using Plan Fluor 40x Oil (numerical aperture 1.3). Positions

containing Fn1-mEGFP-expressing MEFs were imaged at ~17-18 min intervals for ~16 hours

in humidified Tokai Hit stage-top incubator at 5% CO_2. Each position was imaged 40-43

confocal slices at 0.5 μm thickness, the pinhole was set to 1 Airy unit.

Assembly of exogenously-added Fn1: Part II, SMLM The medium containing secreted Fn1-

tdTomato fusion proteins was collected after a 72-hour culture of confluent Fn1-tdTomato

MEFs generated by CRISPR/Cas9 mutagenesis, as described above. The medium was spun

at 300 x g for 5 min to pellet debris. The concentration of Fn1-tdTomato in the supernatant was

quantified using ELISA kit using antibodies specific to mouse Fn1, which does not cross react

with bovine Fn1 (Abcam, cat # ab21097). Fn1-tdTomato-containing supernatant was diluted 4-

fold with fresh MEF culture medium to the final concentration of 5 μg/ml of Fn1-tdTomato, and

2 ml of this supernatant were added to Fn1-mEGFP MEFs plated on 25 mm 1.5H glass

coverslips the day before at 10^5 cells per well in a 6-well plate. Following 24-hours of

incubation at 37°C and 5% CO_2, cells were washed with PBS and fixed in 4% PFA in PBS for

20 min at RT. PFA was quenched with 100mM NH_4Cl in PBS, cell were washed 3x5 min in

PBS, and permeabilized with PBS containing 0.1% Triton-X (PBST). Cells were then incubated

with a blocking solution containing 10% donkey serum in PBST for 30 min at RT. To detect

Fn1-tdTomato, cells were incubated at 4°C overnight with rabbit anti-mCherry antibody

(Abcam, cat #ab167453) diluted 1:100 in the blocking solution. Primary antibodies were

detected with Alexa-647-conjugated donkey anti-rabbit F(ab)$_2$ secondary antibodies (Jackson

laboratories, cat # 711-606-152) diluted 1:300 in blocking solution and incubated for 4 hrs at

RT. Cells were then washed 3x5min in PBST and stored at 4°C until imaging. dSTORM

imaging was performed using GLOX/BME buffer. Cell were imaged at 57° angle using the

SMLM imaging protocol, as described above. Images were processed and reconstructed using

SMAP using the same parameters, as described above.

1014

Live imaging using TIRF $Fn1^{mEGFP/+}$ MEFs were plated on 35-mm round glass bottom Mattek

dishes (catalog # P35G-1.5-14-C), complete medium was switched to imaging medium prior to

filming. Live TIRF microscopy was performed using Nikon A1-HD25 inverted confocal

microscope equipped with 4 laser lines of 100mW per line at 405, 488, and 561nm and

125mW at 640nm, and motorized TIRF illumination. CFI Apochromat TIRF 100xC Oil objective

and EMCCD camera were used. Before imaging lasers were aligned and the critical angle of

incidence for imaging was determined by the software. The exposure time was 20 ms and

readout speed was set at 10 MHz.

1023

Live imaging using confocal point-scanning Live cell imaging was performed using Ibidi

glass bottom 8-well chambers (catalog # 80827). $0.6*10^4$ wild-type or $Fn1^{mEGFP/+}$ MEFs were

plated on glass in each well of Ibidi glass bottom 8-well chambers and allowed to grow

overnight prior to staining and imaging by direct Stochastic Optical Reconstruction Microscopy

(dSTORM). For FUD and III-11C treatment, $Fn1^{mEGFP/+}$ MEFs were plated in 8-well glass Ibidi

dishes (1 cm^2 growth area) without coating at a density of $0.6x10^4$ cells/well in complete

medium. After 5 hours, DMEM was removed and cells were rinsed once with 1X PBS.

Subsequently, the medium was changed to imaging medium. For FUD experiments, imaging

medium was supplemented either with 225 nM FUD or 274 nM of control III-11C peptide.

Untreated wells contained cells incubated with imaging medium. Following the addition of the

imaging medium (with or without the peptides), the chamber was immediately set up for

imaging in the humidified Tokai Hit stage-top incubator at 37°C, 5% CO_2. Live imaging was

performed using Nikon A1-HD25 inverted confocal microscope with the DUG 4-Channel

Detector and 2 GaAsP, 2 high-sensitivity PMTs, and a motorized XYZ stage with Nikon's

Perfect Focus 4 system, and Plan Fluor 40x Oil (numerical aperture 1.3, cat # MRH01401).

mEGFP was excited using 488 nm laser at 1% power and pinhole set to 1 Airy unit. An optical

zoom of 2 and Z step size of 0.5 μm were used, and stack size was set to 10-15 microns

allowing to image the entire cell. For overnight movies, each position was filmed every 1.5 min

– 4 min, as noted in Movie legends. Movies in the mp4 format were generated using Imaris

9.5.1 (Bitplane), titles and arrows were added using Adobe Premiere Elements Editor 2020.

1044

SMLM imaging protocol used to image cells plated on Ibidi glass-bottom dishes. This

protocol was used to image cells following the live DOC assay or overnight live imaging (**Fig.**

10 and **Sup. Fig. 9**). Following fixation with 4% PFA, cells were washed with PBS, incubated

with blocking buffer containing PBS, 0.1% Triton-X, and 10% donkey serum for 30 min at RT.

Cells were then incubated with Abcam monoclonal anti-Fn1 antibody diluted at 1:300 in

blocking buffer overnight at 4°C. Following three washes at 5 min each in PBST (PBS with

0.1% TritonX), cells were incubated with Alexa-647-conjugated anti-rabbit secondary

antibodies diluted 1:300 in blocking buffer for 4 hours at RT, and washed three washes at 5

min each in PBST. Prior to imaging, freshly prepared GLOX/MEA buffer was added and the

chamber was immediately sealed using parafilm. STORM was performed using Nikon A1-

HD25 Ti2E microscope equipped with motorized TIRF illumination, 125mW 640 nm solid-state

laser, Perfect Focus, and a 100x/1.49NA objective. Images were acquired at the critical angle

of incidence and recorded using a 512 x 512 EMCCD camera (Princeton Instruments).

Calibration was obtained by imaging of 100 nm Tetraspeck beads (Life technologies, catalog #

T-7279) using the same glass surface and buffer conditions. To drive Alexa-647 into the dark

state, samples were pre-bleached by the illumination at 640 nm for 10 seconds at 100 % laser

power. 40,000 frames were acquired at 8.4 ms exposure. Blinking events were fitted using the

Nikon N-STORM localization software. Images were analyzed using Nikon software (Nikon

Elements AR Software v5.11.01). Localization events with fewer than 800 or more than 50000

photons were filtered out to remove blinking evens that were either too faint or too bright. In

addition, blinking events were filtered out if they occurred in more than 5 consecutive frames or

where outside the z-range determined by the calibration using 100 nm Tetraspeck beads.

Images in which z-rejection was below 50% were used for further analyses.

Analysis of localization numbers in fibrillar and non-fibrillar nanodomains In order to

enrich for non-fibrillar nanodomains, Fn1$^{mEGFP/+}$ MEFs were plated in 8 well glass Ibidi dishes

(1 cm^2 growth area) without coating. Cell were plated at the density of 0.6x10^4 cells/well in

imaging medium with or without FUD (225 nM) or III-11C (274 nM), and incubated in at 37°C,

5% CO$_2$ for 1 hr. Subsequently, MEFs were rinsed once in warm 1X PBS and fixed using pre-

warmed 4% PFA for 20 min. After fixation, wells were rinsed three times, 5 min each with Stain

Buffer (cat # 554656 BD Pharmingen), blocked for 30 min at room using 5% Donkey serum

prepared in Stain Buffer, and incubated with the monoclonal anti-Fn1 (Abcam, cat # 199056)

overnight at 4° C. Cell were then washed with Stain Buffer three times, 10 min each, and

incubated with anti-rabbit antibodies conjugated with Alexa-647 for 1 hour at rt. Cell were then

rinsed again with Stain Buffer three times, 10 min each, and stored at 4° C in 1X PBS. STORM

imaging was performed, as above. To quantify the number of grouped localizations per

nanodomain, we used the free-hand ROI tool in the STORM window (Nikon Elements AR

Software v5.11.01) to determine the number of localizations within non-fibrillar and fibrillar Fn1

nanodomains. Fn1 nanodomains were analyzed in 5 random regions from 3 independently

acquired images (a total of 15 fields) for each sample. To determine the number of

localizations in Fn1 nanodomains within fibrils, we analyzed more than 20 from 3 or more

1086   independently acquired images. All the counts were plotted in Prism 8.2.1 (GraphPad

1087   Software, USA), and compared using either one-way ANOVA test with Tukey's correction or

1088   Kruskal-Wallis test with Dunn's correction for multiple testing.

1089

Double-color dSTORM acquisition Samples were mounted with the SMART Kit buffer

(Abbelight). 2D or 3D STORM images were acquired using a SAFe360 module (Abbelight)

coupled to an inverted bright-field Olympus IX83 microscope, equipped with a 100X oil-

immersion objective (1.49 NA). This dual-cam system (sCMOS cameras, Orcaflashv4,

Hamamatsu) allows to perform multicolor STORM by spectral demixing strategy coupled to the

use of far-red dyes (BioOptics World, 2021, Caorsi and Karlsson).

Briefly, a dichroic at 700nm is used to separate the fluorescence emission from AF647 and

CF680 dyes. The Point Spread Function (PSF) of each detection can be retrieved on both

cameras and the measured photon numbers is related to the spectral separation of the

Figure M1. Spectral de-mixing optical scheme. A. A 640 nm laser
is sent through the objective lens to excite AF647 and CF680
fluorophores. The emitted light is separated by a 700 nm long pass
dichroic filter that reflects light below 700 nm into the Reflected
camera (Cam R) and transmits light above 700 nm into the
Transmitted camera (Cam T). **B.** Emission spectra of AF647 and
CF680 dyes together with a 700nm dichroic filter showing spectral
separation into the two cameras Cam R and Cam T.

1099   fluorophore (**Fig. M1**).

1100

1101

1102

1103

1104

1105

1106

Typically, acquisition of 60000 frames at 20ms exposure time on a camera sensor size of

1024x1024 pixel were acquired to collect single-molecule detections. The irradiation at the

sample could be properly tuned thanks to ASTER technology (Mau et al. 2021 Nature

Communications, https://www.nature.com/articles/s41467-021-23405-4) implemented on the

SAFe360 Abbelight module. Resulting coordinate tables and images were processed and

analyzed using NEO software (Abbelight).

1113

1114   As the PSF is captured on both cameras, transmitted and reflected, a ratiometric analysis is

1115   applied: a ratio for each detection is calculated and the final ratio distribution is used for

1116   lambda assignment:

1117
$$R_{A,i} = \frac{I_{R,i}}{I_{R,i} + I_{T,i}}$$

1118

1119   where the suffix A is the fluorophore, i the localisation and I_R and I_T the intensity measured on

1120   camera R and camera T (i.e. the number of photons emitted per molecule).

1121 Average ratio distributions obtained from measurements are shown in **Fig.M2**:

Figure M2. Average ratio distributions for Fn1$^{mEGFP/mEGFP}$ cells stained using chicken anti-GFP and rabbit anti-GFP antibodies (Fn1GG, dark gray) or chicken anti-GFP and rabbit R457 antibodies (Fn1GG_R457, lighter gray). The two fluorophore populations can be clearly distinguished for lambda assignment.

1122

1123

Following the ratio distributions measured on the samples the following parameters have been

used for separation: detections with ratios between 0-0.45 were assigned to CF680 and

detections with ratios between 0.5-1 were assigned to AF647. On average 3% of detections

were rejected while keeping crosstalk below 1%. Following de-mixing, colocalization analysis

was performed on Neo software following the CBC algorithm (Malkusch et al., 2012).

Parameters set for CBC analysis were R_{max} at 300 nm and a number of steps equal to 10.

1130

Legends for Movies

Movie 1. Rotational views through the Fn1+ ECM in the cardiac jelly. The whole E9.5

embryo was stained using Abcam rabbit monoclonal anti-Fn1 antibody and imaged using 100x

objective, N.A. 1.49, with the pinhole set at 0.8 Airy units, and sampling of 40 nm per pixel in x,

y. The movie shows 3D reconstruction through 3.4 μm of tissue sampled every 0.121 μm in z.

Fn1 is in white, DAPI is in blue. Arrows point to examples of beaded Fn1 fibrils.

1137

Movie 2. Fn1 fibrillogenesis imaged by TIRF microscopy. Fn1^{mEGFP} MEFs were transiently transfected with mCardinal-Lifeact, plated on gelatin-coated glass coverslips, and imaged 48 hours later. Filming was done every 2 min for 30 min using TIRF and 100x objective, N.A. 1.49. The first set shows the Fn1-mEGFP channel. Yellow arrows point to centripetally-moving Fn1 nanodomains organized into an elongating linear fibril. The second set is an overlay between Fn1-mEGFP and mCardinal-Lifeact.

Movie 3. 2% DOC dissolved cytoplasm and nucleus in under 13 min leaving Fn1 fibrils. MEFs expressing Fn1-mNeonGreen were plated on glass-bottom slides without coating and labeled with SiRActin (magenta) to visualize F-actin, and Hoechst (blue) to visualize DNA. Time-lapse was recorded every 54 sec immediately following the addition of the DOC solution pH 8.01 to live cells. The presence of 2% DOC dissolves actin cytoskeleton and nuclei and leaves Fn1 ECM fibrils (green). Fn1 fibrils collapse following the dissolution of the actin cytoskeleton due to the loss of tension.

Movie 4. Fibrillogenesis of ectopic Fn1. 3-well culture insert was placed in the middle of the 35 mm glass-bottom dish and 0.8×10^6 Fn1-tdTomato-expressing MEFs were plated surrounding the inserts. 24 hours later, Fn1-mEGFP-expressing MEFs were plated inside the inserts on glass without coating for 5 hours. Confocal live imaging of areas containing Fn1-mEGFP-expressing MEFs was performed using Plan Fluor 40x Oil (NA 1.3). Positions containing Fn1-mEGFP-expressing MEFs were imaged at 17-18 min intervals for ~16 hours. This movie contains maximum intensity projections composed of forty-three confocal slices at 0.5 μm thickness, the pinhole was set to 1 Airy unit. The first still panel in this movie is a montage of the plate to show Fn1-tdTomato and Fn1-mEGFP-expressing cells prior to the start of the time-lapse.

1163

Movie 5. Cells incubated with 11-IIIC, show robust fibrillogenesis. Fn1$^{mEGFP/+}$ MEFs were

plated on uncoated glass in 8-well Ibidi chambers for 4 hours. Medium containing 11-IIIC

control peptide was then added and cells were filmed every 90 sec for about 15 hours, as

described in Methods. The movie begins approximately 30 min after the 11-IIIC-containing

medium was added, the time it takes to set up time-lapse recording. Arrows point to the cell

periphery and examples of centripetally moving Fn1 fibrils.

1170

Movie 6. FUD interferes with linking centripetally moving Fn1+ nanodomains into fibrils.

Fn1$^{mEGFP/+}$ MEFs were plated on glass in 8-well Ibidi chambers for 4 hours. Medium containing

FUD peptide was then added and cells were filmed every 3 min for about 15 hours, as

described in Methods. The movie begins approximately 30 min after the FUD-containing

medium was added, the time it takes to set up time-lapse recording. Note the dismantling of

pre-existing fibrils at the beginning of the movie. Yellow and red arrows point to the cell

periphery. Note the presence of centripetally moving Fn1-mEGFP "beads" and the scarcity of

Fn1 fibrils for the majority of the duration of the movie.

1179

References

Aguirre, K.M., R.J. McCormick, and J.E. Schwarzbauer. 1994. Fibronectin self-association is mediated by complementary sites within the amino-terminal one-third of the molecule. *J Biol Chem*. 269:27863-27868.

Andronov, L., Y. Lutz, J.L. Vonesch, and B.P. Klaholz. 2016a. SharpViSu: integrated analysis and segmentation of super-resolution microscopy data. *Bioinformatics*. 32:2239-2241.

Andronov, L., I. Orlov, Y. Lutz, J.L. Vonesch, and B.P. Klaholz. 2016b. ClusterViSu, a method for clustering of protein complexes by Voronoi tessellation in super-resolution microscopy. *Sci Rep*. 6:24084.

Arai, Y., R. Yasuda, K. Akashi, Y. Harada, H. Miyata, K. Kinosita, Jr., and H. Itoh. 1999. Tying a molecular knot with optical tweezers. *Nature*. 399:446-448.

Baumgart, F., A.M. Arnold, K. Leskovar, K. Staszek, M. Folser, J. Weghuber, H. Stockinger, and G.J. Schutz. 2016. Varying label density allows artifact-free analysis of membrane-protein nanoclusters. *Nat Methods*. 13:661-664.

Behringer, R., M. Gertsenstein, K. Vintersten, and A. Nagy. 2014. Manipulating the Mouse Embryo: A Laboratory Manual. Cold Spring Harbor Laboratory Press, New York. 814 pp.

Bui, K.H., A. von Appen, A.L. DiGuilio, A. Ori, L. Sparks, M.T. Mackmull, T. Bock, W. Hagen, A. Andres-Pons, J.S. Glavy, and M. Beck. 2013. Integrated structural analysis of the human nuclear pore complex scaffold. *Cell*. 155:1233-1243.

Caetano, F.A., B.S. Dirk, J.H. Tam, P.C. Cavanagh, M. Goiko, S.S. Ferguson, S.H. Pasternak, J.D. Dikeakos, J.R. de Bruyn, and B. Heit. 2015. MIiSR: Molecular Interactions in Super-Resolution Imaging Enables the Analysis of Protein Interactions, Dynamics and Formation of Multi-protein Structures. *PLoS Comput Biol*. 11:e1004634.

Chen, Y., L. Zardi, and D.M. Peters. 1997. High-resolution cryo-scanning electron microscopy study of the macromolecular structure of fibronectin fibrils. *Scanning*. 19:349-355.

Chernousov, M.A., A.I. Faerman, M.G. Frid, O. Printseva, and V.E. Koteliansky. 1987. Monoclonal antibody to fibronectin which inhibits extracellular matrix assembly. *FEBS Lett*. 217:124-128.

Chiang, H.Y., V.A. Korshunov, A. Serour, F. Shi, and J. Sottile. 2009. Fibronectin is an important regulator of flow-induced vascular remodeling. *Arterioscler Thromb Vasc Biol*. 29:1074-1079.

Choi, M.G., and R.O. Hynes. 1979. Biosynthesis and processing of fibronectin in NIL.8 hamster cells. *J Biol Chem*. 254:12050-12055.

Diekmann, R., M. Kahnwald, A. Schoenit, J. Deschamps, U. Matti, and J. Ries. 2020. Optimizing imaging speed and excitation intensity for single-molecule localization microscopy. *Nat Methods*. 17:909-912.

Dzamba, B.J., and D.M. Peters. 1991. Arrangement of cellular fibronectin in noncollagenous fibrils in human fibroblast cultures. *J Cell Sci*. 100 (Pt 3):605-612.

Engel, J., E. Odermatt, A. Engel, J.A. Madri, H. Furthmayr, H. Rohde, and R. Timpl. 1981.
Shapes, domain organizations and flexibility of laminin and fibronectin, two multifunctional
proteins of the extracellular matrix. *J Mol Biol*. 150:97-120.

Erickson, H.P., N. Carrell, and J. McDonagh. 1981. Fibronectin molecule visualized in electron
microscopy: a long, thin, flexible strand. *J Cell Biol*. 91:673-678.

Ester, M., H.-P. Krigel, J. Sander, and X. Xu. 1996. A density-based algorithm for discovering
clusters in large spatial databases with noise. *Proc Second Int Conf Knowl Discov Data*
Min.:226–231.

Filla, M.S., K.D. Dimeo, T. Tong, and D.M. Peters. 2017. Disruption of fibronectin matrix affects
type IV collagen, fibrillin and laminin deposition into extracellular matrix of human trabecular
meshwork (HTM) cells. *Exp Eye Res*. 165:7-19.

Fruh, S.M., I. Schoen, J. Ries, and V. Vogel. 2015. Molecular architecture of native fibronectin
fibrils. *Nat Commun*. 6:7275.

Furcht, L.T., D. Smith, G. Wendelschafer-Crabb, D.F. Mosher, and J.M. Foidart. 1980a.
Fibronectin presence in native collagen fibrils of human fibroblasts: immunoperoxidase and
immunoferritin localization. *J Histochem Cytochem*. 28:1319-1333.

Furcht, L.T., G. Wendelschafer-Crabb, D.F. Mosher, and J.M. Foidart. 1980b. Ascorbate-
induced fibroblast cell matrix: reaction of antibodies to procollagen I and III and fibronectin in
an axial periodic fashion. *Prog Clin Biol Res*. 41:829-843.

Furcht, L.T., G. Wendelschafer-Crabb, D.F. Mosher, and J.M. Foidart. 1980c. An axial periodic
fibrillar arrangement of antigenic determinants for fibronectin and procollagen on ascorbate
treated human fibroblasts. *J Supramol Struct*. 13:15-33.

Geiger, B., A. Bershadsky, R. Pankov, and K.M. Yamada. 2001. Transmembrane crosstalk
between the extracellular matrix--cytoskeleton crosstalk. *Nature reviews. Molecular cell*
biology. 2:793-805.

Geiger, B., and K.M. Yamada. 2011. Molecular architecture and function of matrix adhesions.
Cold Spring Harb Perspect Biol. 3.

Gudzenko, T., and C.M. Franz. 2015. Studying early stages of fibronectin fibrillogenesis in
living cells by atomic force microscopy. *Mol Biol Cell*. 26:3190-3204.

Guvendiren, M., and J.A. Burdick. 2012. Stiffening hydrogels to probe short- and long-term
cellular responses to dynamic mechanics. *Nat Commun*. 3:792.

Guvendiren, M., S. Yang, and J.A. Burdick. 2009. Swelling-Induced Surface Patterns in
Hydrogels with Gradient Crosslinking Density. *Advanced Functional Materials*. 19:3038-3045.

Heilemann, M., S. van de Linde, M. Schuttpelz, R. Kasper, B. Seefeldt, A. Mukherjee, P.
Tinnefeld, and M. Sauer. 2008. Subdiffraction-resolution fluorescence imaging with
conventional fluorescent probes. *Angew Chem Int Ed Engl*. 47:6172-6176.

Hocking, D.C., J. Sottile, and P.J. McKeown-Longo. 1994. Fibronectin's III-1 module contains a conformation-dependent binding site for the amino-terminal region of fibronectin. *J Biol Chem*. 269:19183-19187.

Hynes, R.O. 1990. Fibronectins. Springer-Verlag, New York.

Hynes, R.O. 2009. The extracellular matrix: not just pretty fibrils. *Science*. 326:1216-1219.

Jimenez, A., K. Friedl, and C. Leterrier. 2020. About samples, giving examples: Optimized Single Molecule Localization Microscopy. *Methods*. 174:100-114.

Kaech, S., and G. Banker. 2006. Culturing hippocampal neurons. *Nat Protoc*. 1:2406-2415.

Khetan, S., M. Guvendiren, W.R. Legant, D.M. Cohen, C.S. Chen, and J.A. Burdick. 2013. Degradation-mediated cellular traction directs stem cell fate in covalently crosslinked three-dimensional hydrogels. *Nature Materials*. 12:458-465.

Lansky, Z., Y. Mutsafi, L. Houben, T. Ilani, G. Armony, S.G. Wolf, and D. Fass. 2019. 3D mapping of native extracellular matrix reveals cellular responses to the microenvironment. *J Struct Biol X*. 1:100002.

Lelek, M., M.T. Gyparaki, G. Beliu, F. Schueder, J. Griffié, S. Manley, R. Jungmann, M. Sauer, M. Lakadamyali, and C. Zimmer. 2021. Single-molecule localization microscopy. *Nature Reviews Methods Primers*. 1:39.

Lu, J., A.D. Doyle, Y. Shinsato, S. Wang, M.A. Bodendorfer, M. Zheng, and K.M. Yamada. 2020. Basement Membrane Regulates Fibronectin Organization Using Sliding Focal Adhesions Driven by a Contractile Winch. *Dev Cell*. 52:631-646 e634.

Ma, W., H. Ma, and D.F. Mosher. 2015. On-Off Kinetics of Engagement of FNI Modules of Soluble Fibronectin by beta-Strand Addition. *PLoS One*. 10:e0124941.

Malkusch, S., U. Endesfelder, J. Mondry, M. Gelleri, P.J. Verveer, and M. Heilemann. 2012. Coordinate-based colocalization analysis of single-molecule localization microscopy data. *Histochem Cell Biol*. 137:1-10.

Martino, M.M., and J.A. Hubbell. 2010. The 12th-14th type III repeats of fibronectin function as a highly promiscuous growth factor-binding domain. *FASEB J*. 24:4711-4721.

Maurer, L.M., B.R. Tomasini-Johansson, W. Ma, D.S. Annis, N.L. Eickstaedt, M.G. Ensenberger, K.A. Satyshur, and D.F. Mosher. 2010. Extended binding site on fibronectin for the functional upstream domain of protein F1 of Streptococcus pyogenes. *J Biol Chem*. 285:41087-41099.

Mautner, V., and R.O. Hynes. 1977. Surface distribution of LETS protein in relation to the cytoskeleton of normal and transformed cells. *J Cell Biol*. 75:743-768.

McKeown-Longo, P.J., and D.F. Mosher. 1983. Binding of plasma fibronectin to cell layers of human skin fibroblasts. *J Cell Biol*. 97:466-472.

Morla, A., Z. Zhang, and E. Ruoslahti. 1994. Superfibronectin is a functionally distinct form of fibronectin. *Nature*. 367:193-196.

Mund, M., and J. Ries. 2020. How good are my data? Reference standards in superresolution microscopy. *Mol Biol Cell*. 31:2093-2096.

Muzumdar, M.D., B. Tasic, K. Miyamichi, L. Li, and L. Luo. 2007. A global double-fluorescent Cre reporter mouse. *Genesis*. 45:593-605.

Nieuwenhuizen, R.P., K.A. Lidke, M. Bates, D.L. Puig, D. Grunwald, S. Stallinga, and B. Rieger. 2013. Measuring image resolution in optical nanoscopy. *Nat Methods*. 10:557-562.

Ohashi, T., D.P. Kiehart, and H.P. Erickson. 1999. Dynamics and elasticity of the fibronectin matrix in living cell culture visualized by fibronectin-green fluorescent protein. *Proc Natl Acad Sci U S A*. 96:2153-2158.

Ovesny, M., P. Krizek, J. Borkovec, Z. Svindrych, and G.M. Hagen. 2014. ThunderSTORM: a comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution imaging. *Bioinformatics*. 30:2389-2390.

Pankov, R., E. Cukierman, B.Z. Katz, K. Matsumoto, D.C. Lin, S. Lin, C. Hahn, and K.M. Yamada. 2000. Integrin dynamics and matrix assembly: tensin-dependent translocation of alpha(5)beta(1) integrins promotes early fibronectin fibrillogenesis. *J Cell Biol*. 148:1075-1090.

Pankov, R., A. Momchilova, N. Stefanova, and K.M. Yamada. 2019. Characterization of stitch adhesions: Fibronectin-containing cell-cell contacts formed by fibroblasts. *Exp Cell Res*. 384:111616.

Patla, I., T. Volberg, N. Elad, V. Hirschfeld-Warneken, C. Grashoff, R. Fassler, J.P. Spatz, B. Geiger, and O. Medalia. 2010. Dissecting the molecular architecture of integrin adhesion sites by cryo-electron tomography. *Nat Cell Biol*. 12:909-915.

Perez, A.R., Y. Pritykin, J.A. Vidigal, S. Chhangawala, L. Zamparo, C.S. Leslie, and A. Ventura. 2017. GuideScan software for improved single and paired CRISPR guide RNA design. *Nat Biotechnol*. 35:347-349.

Peters, D.M., L.M. Portz, J. Fullenwider, and D.F. Mosher. 1990. Co-assembly of plasma and cellular fibronectins into fibrils in human fibroblast cultures. *J Cell Biol*. 111:249-256.

Peters, J.H., and R.O. Hynes. 1996. Fibronectin isoform distribution in the mouse. I. The alternatively spliced EIIIB, EIIIA, and V segments show widespread codistribution in the developing mouse embryo. *Cell Adhes Commun*. 4:103-125.

Pliatsika, V., and I. Rigoutsos. 2015. "Off-Spotter": very fast and exhaustive enumeration of genomic lookalikes for designing CRISPR/Cas guide RNAs. *Biol Direct*. 10:4.

Raitman, I., M.L. Huang, S.A. Williams, B. Friedman, K. Godula, and J.E. Schwarzbauer. 2018. Heparin-fibronectin interactions in the development of extracellular matrix insolubility. *Matrix Biol*. 67:107-122.

Ramirez, A., and S. Astrof. 2020. Visualization and Analysis of Pharyngeal Arch Arteries using Whole-mount Immunohistochemistry and 3D Reconstruction. *J. Vis. Exp.* 157:e60797.

Ran, F.A., P.D. Hsu, J. Wright, V. Agarwala, D.A. Scott, and F. Zhang. 2013. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc*. 8:2281-2308.

Rickelt, S., and R.O. Hynes. 2018. Antibodies and methods for immunohistochemistry of extracellular matrix proteins. *Matrix Biol*. 71-72:10-27.

Ries, J. 2020. SMAP: a modular super-resolution microscopy analysis platform for SMLM data. *Nat Methods*. 17:870-872.

Rozario, T., B. Dzamba, G.F. Weber, L.A. Davidson, and D.W. DeSimone. 2009. The physical state of fibronectin matrix differentially regulates morphogenetic movements in vivo. *Dev Biol*. 327:386-398.

Rust, M.J., M. Bates, and X. Zhuang. 2006. Sub-diffraction-limit imaging by stochastic optical reconstruction microscopy (STORM). *Nat Methods*. 3:793-795.

Saitta, A.M., P.D. Soper, E. Wasserman, and M.L. Klein. 1999. Influence of a knot on the strength of a polymer strand. *Nature*. 399:46-48.

Sakai, T., K.J. Johnson, M. Murozono, K. Sakai, M.A. Magnuson, T. Wieloch, T. Cronberg, A. Isshiki, H.P. Erickson, and R. Fassler. 2001. Plasma fibronectin supports neuronal survival and reduces brain injury following transient focal cerebral ischemia but is not essential for skin-wound healing and hemostasis. *Nat Med*. 7:324-330.

Saunders, J.T., and J.E. Schwarzbauer. 2019. Fibronectin matrix as a scaffold for procollagen proteinase binding and collagen processing. *Mol Biol Cell*. 30:2218-2226.

Schwarzbauer, J.E. 1991. Identification of the fibronectin sequences required for assembly of a fibrillar matrix. *J Cell Biol*. 113:1463-1473.

Schwarzbauer, J.E., and D.W. DeSimone. 2011. Fibronectins, their fibrillogenesis, and in vivo functions. *Cold Spring Harbor perspectives in biology*. 3.

Sechler, J.L., H. Rao, A.M. Cumiskey, I. Vega-Colon, M.S. Smith, T. Murata, and J.E. Schwarzbauer. 2001. A novel fibronectin binding site required for fibronectin fibril growth during matrix assembly. *J Cell Biol*. 154:1081-1088.

Sechler, J.L., Y. Takada, and J.E. Schwarzbauer. 1996. Altered rate of fibronectin matrix assembly by deletion of the first type III repeats. *J Cell Biol*. 134:573-583.

Shaner, N.C., P.A. Steinbach, and R.Y. Tsien. 2005. A guide to choosing fluorescent proteins. *Nat Methods*. 2:905-909.

Singh, P., C. Carraher, and J.E. Schwarzbauer. 2010. Assembly of fibronectin extracellular matrix. *Annu Rev Cell Dev Biol*. 26:397-419.

Singh, P., and J.E. Schwarzbauer. 2014. Fibronectin matrix assembly is essential for cell condensation during chondrogenesis. *J Cell Sci*. 127:4420-4428.

Sivakumar, P., A. Czirok, B.J. Rongish, V.P. Divakara, Y.P. Wang, and S.L. Dallas. 2006. New insights into extracellular matrix assembly and reorganization from dynamic imaging of extracellular matrix proteins in living osteoblasts. *J Cell Sci*. 119:1350-1360.

Skorstengaard, K., M.S. Jensen, P. Sahl, T.E. Petersen, and S. Magnusson. 1986. Complete primary structure of bovine plasma fibronectin. *Eur J Biochem*. 161:441-453.

Smith, M.L., D. Gourdon, W.C. Little, K.E. Kubow, R.A. Eguiluz, S. Luna-Morris, and V. Vogel.
2007. Force-induced unfolding of fibronectin in the extracellular matrix of living cells. *PLoS*
Biol. 5:e268.

Snitkovsky, S., and J.A. Young. 1998. Cell-specific viral targeting mediated by a soluble
retroviral receptor-ligand fusion protein. *Proc Natl Acad Sci U S A*. 95:7063-7068.

Sottile, J., and J. Chandler. 2005. Fibronectin matrix turnover occurs through a caveolin-1-
dependent process. *Mol Biol Cell*. 16:757-768.

Thevathasan, J.V., M. Kahnwald, K. Cieslinski, P. Hoess, S.K. Peneti, M. Reitberger, D. Heid,
K.C. Kasuba, S.J. Hoerner, Y. Li, Y.L. Wu, M. Mund, U. Matti, P.M. Pereira, R. Henriques, B.
Nijmeijer, M. Kueblbeck, V.J. Sabinina, J. Ellenberg, and J. Ries. 2019. Nuclear pores as
versatile reference standards for quantitative superresolution microscopy. *Nat Methods*.
16:1045-1053.

Tomasini-Johansson, B.R., D.S. Annis, and D.F. Mosher. 2006. The N-terminal 70-kDa
fragment of fibronectin binds to cell surface fibronectin assembly sites in the absence of intact
fibronectin. *Matrix Biol*. 25:282-293.

Tomasini-Johansson, B.R., N.R. Kaufman, M.G. Ensenberger, V. Ozeri, E. Hanski, and D.F.
Mosher. 2001. A 49-residue peptide from adhesin F1 of Streptococcus pyogenes inhibits
fibronectin matrix assembly. *J Biol Chem*. 276:23430-23439.

von Appen, A., J. Kosinski, L. Sparks, A. Ori, A.L. DiGuilio, B. Vollmer, M.T. Mackmull, N.
Banterle, L. Parca, P. Kastritis, K. Buczak, S. Mosalaganti, W. Hagen, A. Andres-Pons, E.A.
Lemke, P. Bork, W. Antonin, J.S. Glavy, K.H. Bui, and M. Beck. 2015. In situ structural
analysis of the human nuclear pore complex. *Nature*. 526:140-143.

Wagner, D.D., and R.O. Hynes. 1979. Domain structure of fibronectin and its relation to
function. Disulfides and sulfhydryl groups. *J Biol Chem*. 254:6746-6754.

Wierzbicka-Patynowski, I., Y. Mao, and J.E. Schwarzbauer. 2004. Analysis of fibronectin
matrix assembly. *Current protocols in cell biology / editorial board, Juan S. Bonifacino ... [et*
al.]. Chapter 10:Unit 10 12.

Wijelath, E.S., J. Murray, S. Rahman, Y. Patel, A. Ishida, K. Strand, S. Aziz, C. Cardona, W.P.
Hammond, G.F. Savidge, S. Rafii, and M. Sobel. 2002. Novel Vascular Endothelial Growth
Factor Binding Domains of Fibronectin Enhance Vascular Endothelial Growth Factor Biological
Activity. *Circulation research*. 91:25-31.

Zamir, E., B.Z. Katz, S. Aota, K.M. Yamada, B. Geiger, and Z. Kam. 1999. Molecular diversity
of cell-matrix adhesions. *J Cell Sci*. 112 (Pt 11):1655-1669.

Zamir, E., M. Katz, Y. Posen, N. Erez, K.M. Yamada, B.Z. Katz, S. Lin, D.C. Lin, A.
Bershadsky, Z. Kam, and B. Geiger. 2000. Dynamics and segregation of cell-matrix adhesions
in cultured fibroblasts. *Nat Cell Biol*. 2:191-196.

Zhang, Q., W.J. Checovich, D.M. Peters, R.M. Albrecht, and D.F. Mosher. 1994. Modulation of
cell surface fibronectin assembly sites by lysophosphatidic acid. *J Cell Biol*. 127:1447-1459.

Zhang, Q., M.K. Magnusson, and D.F. Mosher. 1997. Lysophosphatidic acid and microtubule-
destabilizing agents stimulate fibronectin matrix assembly through Rho-dependent actin stress
fiber formation and cell contraction. *Mol Biol Cell*. 8:1415-1425.

Zhang, Q., and D.F. Mosher. 1996. Cross-linking of the NH2-terminal region of fibronectin to
molecules of large apparent molecular mass. Characterization of fibronectin assembly sites
induced by the treatment of fibroblasts with lysophosphatidic acid. *J Biol Chem*. 271:33284-
33292.

Zhong, C., M. Chrzanowska-Wodnicka, J. Brown, A. Shaub, A.M. Belkin, and K. Burridge.
1998. Rho-mediated contractility exposes a cryptic site in fibronectin and induces fibronectin
matrix assembly. *J Cell Biol*. 141:539-551.

Zhou, X., R.G. Rowe, N. Hiraoka, J.P. George, D. Wirtz, D.F. Mosher, I. Virtanen, M.A.
Chernousov, and S.J. Weiss. 2008. Fibronectin fibrillogenesis regulates three-dimensional
neovessel formation. *Genes Dev*. 22:1231-1243.
1417

Table 1

Sample	Buffer	ELE (%)	Average # Localizations per NPC	# Localizations per protein	Radius of NPC, nm
U2OS-Nup96	GLOX/BME	79.6+/- 5.4	280 +/- 49.2	8.75 +/- 1.54	63.6 +/- 0.86

NUP96$^{mEGFP/mEGFP}$ cells were imaged using αGFP 1º antibody and AF647-conjugated 2º antibody detecting NUP96-mEGFP; Data is from 3 independent experiments, eight cells, and 4571 NPCs

Table 2

	Buffer	Average PSF (nm)	Average # Localizations per nanodomain	Median Localization Precision, nm	# of mEGFP molecules per nanodomain in Fn1-GFP fibrils	Diameter of Fn1 nanodomain, nm	Nanodomain periodicity, nm
Fn1$^{mEGFP/mEGFP}$ MEFs; αGFP 1° antibody; AF647-conjugated 2° antibody	GLOX/BME	143.8 +/- 2.35	159 +/- 77.92	9.1 +/- 1.9	16.85 +/- 5.1	28.26+/-9.7	102±29

αGFP 1° antibody; AF647-conjugated 2° antibody were used to detect Fn1-mEGFP. Data is from 3 independent experiments, 6 cells, 52 fibrils and 833 nanodomains were used to calculate the number of GFP molecules per nanodomain
Data from 3 independent experiments, 15 cells, 27 fibrils, and 1292 nanodomains were used to quantify nanodomain diameter

Figure 1

Figure 1. Beaded architecture of Fn1 fibrils in embryonic ECM. Wild-type E9.5 mouse embryos were fixed and stained with the Alexa-monoclonal antibody to Fn1 (white) and DAPI (blue), and imaged using 100x oil objective, N.A. 1.46, pinhole 0.8, and sampling rate of 40 nm/pixel. **A – A'.** Sagittal optical section through the first pharyngeal arch and **B**, the cardiac jelly, an ECM-rich region between the myocardial and endocardial layers of the cardiac outflow tract. Large arrowheads in **A – A'** point to the ECM at the ectoderm-mesenchyme boundary of the 1st pharyngeal arch. The box in **(A)** is expanded in **A1** to show Fn1 microarchitecture. Arrow in **A1** points to a beaded Fn1 fibril within the 1st arch mesenchyme. **B.** Beaded architecture of Fn1 fibrils in cardiac jelly, e.g., arrow **C**, intensity profile plot of a Fn1 fibril shows a regularly-undulating profile, with troughs well above the noise, e.g., dotted box.

Figure 2. Integrin α5 and Fn1 co-localize in beaded adhesions. Wild-type MEFs were plated for 16 hours on glass coverslips without coating, then fixed, stained with Abcam monoclonal antibodies to Fn1 and anti-integrin α5 (Itga5) antibodies. Cells were imaged at the critical angle of incidence using 100x oil objective, NA 1.49. **A – A2** panels show examples of non-fibrillar Fn1 adhesions. **B – B2** medial portion of a cell containing beaded fibrillar adhesions (arrows). Magnifications in all panels are the same.

Figure 3

Figure 3. Standardized SMLM imaging results in high resolution and high effective labeling efficiency (CLE). NUP96[mEGFP/mEGFP] homozygous knock-in cells (A–C) and Fπ1[mMaison/mMaison] homozygous knock-in cells (D–F) were imaged using the same staining and imaging solutions containing anti-GFP 1:1 at 1:100 dilution and Alexa-647-conjugated 2° antibodies, the same GLOX/BME buffer, and the same imaging protocol (see *Methods*). **A–C.** NUP96[mEGFP/mEGFP] cells. The majority of nuclear pore complexes (NPCs) show 8-fold symmetry with 8 corners labeled by NUP96. See *Table 1* for quantification of NPC dimensions and the ELE. The resolution in (C) measured by the Fourier ring curve method is 18.3+/-4.7 nm. **D–F.** Fπ1[mMaison/mMaison] cells. Nanodomain architecture of Fπ1-mEGFP fibrils is clearly visible in subsequent magnifications of the fibril boxed in (D). The boxed region in **D** is shown in **E**. Boxed region in **E** is shown in **F**. Arrows in **F** point to Fπ1 nanodomains. The resolution in (F) measured by Fourier ring curve method is 14.5+/-1.1 nm. Note that nanodomain size is similar to the size of an NUP96+ NPC corner, which is ~12 nm in actual diameter. N1 is the number of grouped localizations in each panel. See *Table 2* for quantifications of the numbers of mEGFP molecules per nanodomain. Vertical bar in **F** shows color-coding according to localization density for all panels in this figure.

Figure 4

A Schematic of Fn1 dimer

B Antibodies used in those studies and their epitopes

Figure 4. Schematic representations of Fn1 domains and antibody binding sites. A. Fn1 is secreted as a dimer. Two di-sulfide bonds at the C-terminus hold Fn1 molecules in an anti-parallel orientation. 70 kDA Fn1 assembly domain is marked in blue and by a double-headed arrow. Fn1 type III repeats are numbered. Splice variants characterized in mice are shown. B. Schematic of binding sites for the antibodies used in this manuscript. All antibodies except one are polyclonal, see Table M2 in the Methods section. R457 is a rabbit polyclonal antibody raised to the 70 kDA domain of Fn1. R184 is a rabbit polyclonal raised against the first six type III repeats (purple). 297.1 is a rabbit polyclonal antibody raised to the full-length (all repeats) Fn1. The binding sites of 297.1 (red arrows) were mapped in this study (see diag. 6). A monoclonal anti-Fn1 antibody recognizing the epitope within the marked region.

Figure 5. Nanodomain architecture of Fn1 fibrils detected with four different antibodies. Fn1[mEGFP/mEGFP] MEFs were plated on glass cover slips without coating, cultured over night, and stained with the following antibodies: **A.** R457 (1:200 dilution). **B.** R184 (1:100 dilution). **C.** Abcam monoclonal antibody (1:200 dilution). **D.** anti-GFP antibody (1:100 dilution). Boxes in **A**, **B**, **C**, and **D** are expanded to show fibrils in **A1**, **B1**, **C1**, and **D1**, respectively. Brackets in **A1**, **B1**, **C1**, and **D1** are expanded in **A2**, **B2**, **C2**, and **D2**, respectively. Scale bars in **A2**, **B2**, **C2**, and **D2** are 100 nm. Fourier ring correlation (FRC) was used to determine image resolution. Bar in **D** shows color-coding according to localization density for all panels in this figure.

Figure 6

Figure 6. Periodical labeling of Fn1 fibrils by multiple antibodies, their dilutions and combinations. Fn1 $^{EGFP/EGFP}$ MEFs were plated on glass cover slips without coating, cultured over night, and stained with antibodies indicated. **A-E.** SMLM images resulting from the highest dilutions of antibodies tested. Brackets in **A - B** are expanded in **A', B'.** Autocorrelation traces for fibrils shown in **A-E.** Ljung-Box Q-test for autocorrelation at the peaks indicates marked by the red arrows resulted in *n*=1, *p*=0 indicating strong evidence of autocorrelation at the first peak. *h* (the null hypothesis for this test) is that the first *h* autocorrelations are jointly zero. *h*=1 rejects the hypothesis. **F.** Summary of autocorrelation analysis of multiple fibrils (each fibrils analyzed is marked by a square) stained with different antibody dilutions, and combinations of antibodies. Experiments for each antibody dilution were repeated three independent times, and each time, at least three cells were imaged. One-way ANOVA test with Tukey's correction for multiple testing showed no significant differences among the means. $^{1:400}$ and R657 were both used at 1:100 dilution. * The 5 antibodies used and their dilutions were the following: R457(1:400), R184(1:200), Abcam (1:400), uGFP (1:100, 297.1(1:400)).

Figure 7

Figure 7. Clustering using DBSCAN suggests a higher order organization of first nanodomains in fibrils. First nanodomains (B) ... MSF4 were plotted on glass cover slips without coating, contrast over right, and stained with antibody indicated. The first A is also analyzed by DBSCAN clustering a (B and C). The minimum number of points per neighborhood [n] was set to 4, as recommended for all-dimensional datasets. B. The size of the neighborhood ε, was set to 44 nm an apparent radius of first nanodomains. C. The first nanodomains were clustered from clusters are labeled with different colors (C). Clustering within nanodomains by the DBSCAN analysis. Clusters first A: the nanodomains are given the same color. Adjacent clusters are labeled with different colors. DBSCAN clustering in (B) using ε of 44 nm [coordinates (compare DBSCAN image above]. Clustering in (C) using an unbiased estimate of ε suggests a higher order organization of first nanodomains. Scale bar is 1 micron.

Figure 8

Figure 8 Polyclonal antibody to full-length Fv1 and combinations of antibodies recognising epitopes along the length of Fv1 reveal periodical nanodomain architecture of Fv1 fibrils.

A, B7.1 rabbit polyclonal antibody raised to full-length, plasma Fv1 recognising multiple epitopes along Fv1 molecule (red arrows). 1,000 ×, left; colocalising with DAPI staining (blue) in a schematic above the panel. For dilutions of 5 antibodies see legend in Fig. 6F.

B, C, D, E, are equivalent in A, B, C and D in panel luar left as well as in D–F respectively.

D, E, F, graphs of autocorrelation values for all panels in this figure, except E and panels in A and B respectively, plotted along the length of respectively. Bars represent mean ± SD.

Scale bars = 10 µm.

Legend:
- Fv1 nanodomain containing multiple Fv1 dimers
- Large apparent molecular mass complexes

Discussion cont (top, Fig. 6F–F).

Normal crystal density

Figure 9. Double-color dSTORM shows that N- and C-termini of Fe1 overlap within Fe1 nanodomains. A–P, Control stainings. Fe1 fibrils were detected in 8 cryoEM/PAPP-pHL plates (at/or stained using N-terminal rabbit and chicken anti-GFP antibodies (anti-rabbit IgG and anti-chicken IgG and a C-terminal rabbit and chicken-GFP antibodies (anti-chicken CY-GFP antibodies were conjugated to CY90 and anti-chicken JY antibodies were conjugated to AF645. Panels A–B were also displayed as A–B' with vertical lines marking overlapping (gray lines) and non-overlapping (pink lines and arrows) subdomains. C–E' Overlay of two channels in panels C, D and E (respectively). F–H, Intensity profiles plotted along lines... Arrowheads in A', B', D, and E point to an overlapping staining, and white arrows next to the arrowhead in B' and E point to an adjacent non-overlapping staining. Pink arrows in A'/B' are control over B or B', respectively to point out non-overlapping localizations. I–L, Images in A–B are color-coded according to the coordinate-based colocalization (CBC) coefficient. J–L', Enlargements of areas marked by rectangles in J and K. Asterisks and/or arrows indicate nanodomains in which the signal channel is higher than the other or... M–P, Quantification of CBC coefficients... regions containing long fibrils (gray boxes) and liver overlap (blue areas) are plotted. Q–L, Fe1 fibrils were detected in 8 cryoEM/PAPP-pHL cells plated with anti-C-terminal N-terminal rabbit and chicken anti-GFP (anti-rabbit IgG) and a C-terminal rabbit and chicken GFP antibody (chicken-GFP antibodies... C-terminal anti-GFP antibody (anti-rabbit IgG)... Q–R, Overlays and non-overlapping... stainings, respectively, as seen in A–B'. J–Q, Images in Q–R are color coded according to the coordinate-based colocalization... S–U, Intensity profiles plotted along the lines in the fibrils marked in Q and R by arrows and arrowheads. V–Y, Images in Q–R are color-coded according to the coordinate-based colocalization (CBC) coefficient. X–Y', Enlargements of areas in X and Y respectively. Asterisks and/or arrows indicate nanodomains in which the signal channel is higher than in the other... AA–BB, Quantification of CBC coefficients... regions containing long fibrils (gray boxes) and their overlap (blue areas)... of antibody (C-terminal chicken and N-terminal rabbit and chicken anti-GFP)... a control for antibodies to... N-terminus... Scale bars, 2 μm (panels A–B, I–K, Q–R, V–W); 500 nm (panels J–L', X–Y'). Merged data from 3 independent experiments.

Figure 10. The N-terminal Fn1 assembly domain regulates the organization of Fn1 nanodomains into linear fibrillar arrays. Fn1[mEOS3Pm] MEFs were plated on glass and were either left untreated (**A-A2**), incubated with the control 11-IIIC peptide (**B-B2**), or the FUD peptide (**C-C2**) for 16 hrs. Cells were then washed with PBS, fixed and stained with the Abcam monoclonal antibody to Fn1 followed by Alexa 647-conjugated secondary antibodies. Cells were then imaged by dSTORM. **A – A2**. Untreated, non-permeabilized cells. **B – B2**. Cells incubated with control 11-IIIC peptide, non-permeabilized. **C – C2**. FUD-treated, non-permeabilized cells. Boxes marked 1 in **A-B** are expanded in **A1-B1**. Boxes marked 2 in **A-B**, are expanded in **A2 – B2**. The box in C is expanded in **C1** and **C2**. Arrows in **A1-B1** point to Fn1 nanodomains (NDs) in fibrils. Arrows in **C1** points to non-fibrillar (NF) nanodomains expanded in **C2**. **D.** Quantification of the number of grouped Fn1 localizations in nanodomains in fibrils and in non-fibrillar nanodomains after various conditions. Red lines mark medians. Differences are not statistically significant, Kruskal-Wallis test with Dunn's correction for multiple testing. **E.** Model of fibril formation: 6-11 Fn1 dimers assemble into nanodomains containing integrin α5β1 at cell periphery, more rearward with actin flow, and become organized into linear arrays of nanodomains. Joining of the additional Fn1 nanodomains to these arrays leads to the generation of longer fibrils as assemblies move toward the cell center. FUD does not interfere with the formation of Fn1 nanodomains; instead, it blocks the organization of Fn1 nanodomains into linear arrays. LAMMs – large apparent molecular mass complexes, defined in Zhang et al., 1996.