Specific breeding system and ploidy interactions increase niche adaptability in potatoes

Nathan Fumia[1], Daniel Rubinoff[2], Rosana Zenil-Ferguson[3], Colin K. Khoury[4,5], Samuel Pironon[6], Michael A. Gore[7], Michael B. Kantar[1]

[1] Department of Tropical Plant and Soil Science, University of Hawaii at Manoa, Honolulu, HI, USA

[2] Department of Plant and Environmental Protection Science, University of Hawaii at Manoa, Honolulu, HI, USA

[3] School of Life Sciences, University of Hawaii at Manoa, Honolulu, HI, USA

[4] International Center for Tropical Agriculture (CIAT), Cali, Colombia

[5] Department of Biology, Saint Louis University, St. Louis, MO, USA

[6] Royal Botanic Gardens, Kew, Richmond, UK

[7] Plant Breeding and Genetics Section, School of Integrative Plant Science, Cornell University, Ithaca, NY 14853

Abstract

Understanding the role of biotic and abiotic factors that drive ecological and evolutionary interactions of economically important plant species is key for sustainability. Wild potatoes (*Solanum* section *Petota*), given their global importance for food security and the large amount of diversity, were chosen as a model system to better understand these interactions. This study aims to characterize the distinct climatic niches of potato wild relatives, and to elucidate the relationships that exist between biological, evolutionary, and climatic attributes for potato wild relatives. We used a combination of ordinary least-squares (OLS) and phylogenetic generalized least-squares (PGLM) analyses to identify divergent discrete climate classes that wild potato species inhabit. We found that self-incompatible diploid and self-compatible polyploid species were able to colonize new habitats and sustain populations across greater climatic breadth more readily than other types. Self-incompatible diploid or self-compatible polyploid species significantly increased the number of discrete niches expected. This result was sustained when correcting for phylogenetic non-independence in the linear models. Overall, our results support the idea that specific breeding systems and ploidy combinations increase niche divergence through the decoupling of geographical range and niche diversity, and therefore, these species should be prioritized for exploration focused on potato crop adaptation to the changing climate.

Keywords: time-calibrated phylogeny, phylogenetic regression, Köppen-Geiger climate class, ecological niche, crop wild relatives

Introduction

Potato (*Solanum tuberosum* L.) is the most important tuber crop worldwide and is the fourth most important crop internationally (Castañeda-Álvarez et al. 2015). However, there is a lack of genetic diversity among many crops, including *S. tuberosum* (Jansky et al. 2013, Khoury et al. 2014), placing increased pressure upon crop management protocols and food security. A proven approach to increasing genetic diversity in crop species is through the utilization of wild relatives for crop improvement (Jansky et al. 2013; Mehrabi et al. 2019). Cultivated potato has 199 known wild relatives, forming the *Solanum* section *Petota*, inhabiting 16 countries in the Americas, and ranging from 38° N to 41° S (Hijmans 2001); 72 of the most threatened and useful species to humans have recently been prioritized for conservation (Castañeda-Álvarez et al. 2015). These 72 species are most commonly found in tropical highlands at 600 to 1200 m in elevation and possess phenotypes similar to cultivated potato through the production of a starchy tuber (Hijmans 2002).

Given the importance of maintaining the crop's productivity, many attributes of the wild relatives of *S. tuberosum* have been defined, including their ploidy, breeding system, germplasm classification, endosperm balance number, single and multi-gene phylogenies, and geographic ranges (Hijmans 2001, Spooner 2001, Spooner 2007, Castañeda-Álvarez 2015, Robertson et al. 2011). These data can be used to discover novel beneficial characteristics present within the wild relative germplasm such as biotic and abiotic resistances as well as to quantify trait introgression. Furthermore, research has identified potato as one of the crops in Sub-Saharan Africa with the highest potential to benefit from crop wild relatives for climate change adaptation, however, these results were not integrated with biological (e.g. breeding system and ploidy) and evolutionary (e.g. phylogenetic tree) information (Pironon et al. 2019). Despite the wide array of

information surrounding the wild relatives of potato, one attribute continues to be under-defined - the discrete climate zones (e.g. niche) each species inhabits and the factors involved (e.g., breeding system, ploidy) in driving the evolution of the highly dynamic climatic diversity in *Solanum* section *Petota*.

Individually exploring life history traits (Wendel et al. 2003, Hijmans et al. 2007, Köhler 2010, Sessa 2019) such as the breeding system has led to contradictory conclusions regarding these traits' influence on ecological niche (Peterson et al. 1999, Husband et al. 2008, Robertson et al. 2010, Campbell 2013, Grossenbacher et al. 2016, Park 2017, Grant 2020), while exploring other traits such as ploidy (Baniaga et al. 2020) has shown a consistent influence. For example, diversification models Zenil-Ferguson et al. (2019) showed that ploidy is the most probable pathway to evolve self-compatibility across Solanaceae. Therefore, there exists an important interaction between ploidy and breeding system (Barringer 2007, Husband et al. 2008, Robertson et al 2010) that might impact evolutionary and ecological processes (Sessa 2019). Furthermore, polyploidization facilitates self-compatibility because whole genome duplication provides security against inbreeding depression (Barringer 2007, Husband et al. 2008, Robertson et al. 2010, Zenil-Ferguson et al. 2019); whereas, self-compatible diploid populations often suffer from large inbreeding depression (Barringer and Geber 2008, Husband and Schemske 2017). As a result, diploid populations are more reliant on self-incompatibility to drive adaptive changes. In *Solanaceae*, polyploid species show higher rates of self-compatibility (Barringer 2007, Husband et al. 2008, Robertson et al 2010). This clear interaction between ploidy and breeding systems provides us with the opportunity to test two key hypotheses: first, that self-compatible species rely on polyploidy in order to generate the variation they need to colonize diverse ecological

niche space; and second, that diploid species rely on outcrossing to increase niche breadth

through gene flow.

To identify the driving factors of ecological diversity in potato wild relatives, we

investigated two biological aspects of ecological diversity - breeding system and ploidy in 72

wild relatives of potato. We combined species' occurrence, climatic, biological (e.g. breeding

system and ploidy), and phylogenetic tree of *Solanum* taxa to test whether the niche diversity of

a given species is guided by a specific breeding system and ploidy interaction. To account for the

potential decoupling of geographical range and niche breadth (Randel et al. 2009), the measure

of climatic diversity is through the use of discrete climate-classification of each occurrence of

these wild relative species. This work supports classic ecological theory of niche divergence

without the requirement of inferring continuous species distributions from point-based climate

descriptions by featuring the relationship between two common intrinsic factors of niche

expansion: (1) decreased reliance on outcrossing reproduction of polyploid variants; and (2)

increased reliance on outcrossing reproduction of diploid variants (Roughgarden 1972, Barton

1996, De Bodt et al. 2005, Johnson et al. 2014).

Materials and Methods

Data Collection

Data organization and analyses were conducted using the R (R Core Team 2020)

packages "raster" (Hijmans 2020) and "tidyverse" (Wickham et al 2019). We obtained 49,165

occurrence records of the 72 *Solanum* species sourced from Castañeda-Álvarez et al (2015).

These occurrences represent the most threatened and useful wild relatives of *Solanum*

tuberosum, the previously cleaned points were further filtered for those lacking latitudinal and/or

longitudinal information, resulting in a total of 37,032 total occurrence points (Castañeda-

Álvarez et al. 2015). Next, the Köppen-Geiger three-tier climate class system was acquired from Rubel and Kottek (2010). The Köppen-Geiger climate class system divides climates into five main groups that are subdivided based on seasonal precipitation and temperature that result in 30 potential discrete classes globally (reviewed in Rubel and Kottek, 2010). Three-tier climate classes were extracted at each occurrence point. The total number of climate classes per species was counted for each species and climate classes with three or fewer occurrences were removed in order to avoid "by-chance" occurrences. Using discrete climate classes allows for a single measure of both niche diversity and breadth. See github repository "https://github.com/Nfumia/Potato_nichediversity_drivers" for code and data files.

Linear Models for Climate Classes

A linear model was fitted using R package "stats" (R Core Team 2020) to identify which interaction of biological factors is correlated with niche diversity in *Solanum* section *Petota*. We used the number of discrete climate classes in which each taxon can occur as a proxy for niche breadth, as these niches vary spatially within the five broad descriptors of tropical, dry, temperate, continental, and polar each of which possessing 2-12 subclassifications. For example, *S. stoloniferum* has fifteen discrete niches in which it occurs, but *S. albornozii* has only one, a temperate oceanic environment. The number of discrete climate classes is the response variable for the model. The predictor variables were combinations of ploidy (Castañeda-Álvarez et al. 2015) and breeding system (Robertson et al. 2010, Zenil-Ferguson et al. 2019) for each species, which were coded as dummy variable interaction terms: self-incompatible diploid, self-compatible diploid, self-compatible asexually propagating polyploid, and unknown breeding system asexually propagating diploid.

Phylogenetic Tree and Phylogenetic Linear Models

A Bayesian molecular clock phylogeny with time-calibration of section *Petota* to outgroups of domesticated tomato (*Solanum lycopersicum*) and domesticated eggplant (*Solanum melongena*) was estimated using 32 plastid genomes and compared to the most recent time-calibrated phylogeny of Särkinen et al. (2013). Due to a lack of plastid genome availability for some species in *Solanum* section *Petota*, only 27 of the 72 prioritized wild relative species were present in our subsequent analyses. Furthermore, 32 species (27 potato wild relatives, 2 domesticated potato, 1 domesticated tomato, 1 tomato wild relative, 1 domesticated eggplant) were aligned using the software MAFFT (multiple alignment using fast Fourier transform) via maxiterate version (Katoh 2009). MrBayes (Huelsenbeck 2001) as implemented in the Geneious software package (Kearse et al., 2012) was used to conduct an initial phylogenetic analysis (Vallejo-Marín 2006, Newton 1999). We used a chain length of 10 million generations with 25% (or 2.5 million) burn in and a subsampling frequency every 1,000 generations. The General Time Reversible (GTR) substitution model was employed for the Bayesian analysis with rate variation of gamma, including 4 categories.

We used the Bayesian uncorrelated relaxed clock-model dating method as implemented in BEAST2 (Bouckaert et al 2019). The uncorrelated relaxed clock-model allows for rate variation across branches and measures for rate autocorrelation between lineages. Node ages are estimated simultaneously in BEAST2, and, therefore, uncertainty is incorporated into the node-age estimation. Our Bayesian MCMC tree output was used as a starting phylogeny. The Hasegawa-Kishino-Yano (HKY) model for DNA base pair substitution was used to better estimate the substitution rates of transition versus transversion as well as the Felsenstein (F81) proposed four-parameter model. A Kappa of 2.0, as estimated by BEAUti2 (Bouckaert et al 2019), was employed. Calibration points for the node-age estimation were sampled from

Särkinen et al. (2013) to create calibration priors: (1) tomato – potato split circa 8 mya (95% HPD 7—10), and (2) eggplant – tomato/potato split circa 14.3 mya (95% HPD 13-16). These calibration points reflect a normal distribution with standard deviations of 0.85 and 1.10 million years, respectively. Yule tree prior with uniform distribution was used given all ingroup and outgroup species in this study currently persist *ex-situ* and/or *in-situ*. Priors were manually generated for each monophyletic clade showing greater than 85% posterior probability from the MrBayes MCMC analysis. Default priors were used for all other parameters. A total of 100 million generations, 10 runs with 10 million generations each, were run in BEAST2 (Bouckaert et al 2019).

Using the time-calibrated phylogeny, we estimated the phylogenetic generalized linear models version of the OLS models proposed in the previous section to account for potential phylogenetic signals in the errors (Felsenstein 1985, Hansen 1997). This is an important step, since it is possible that our explanatory variables are not tracking the evolutionary history of the *Petota* section, and can incorrectly conclude strong correlations between the climatic classes and the life history traits (Uyeda et al. 2018).

These phylogenetic linear models were estimated using a maximum likelihood PGLM with the R package "phylolm" (Ho and Ane 2014). For all the PGLMs we assumed a Brownian motion model of evolution (Grafen 1989, Martins and Hansen 1997, Revell and Harmon 2008). Outgroup species and cultivated potato were removed at this point due to the inability to differentiate between cultivated and wild occurrence of the given species. This resulted in retention of 27 potato wild relative species, comprising the four major monophyletic clades of section *Petota* (Spooner et al. 2014), for use in the PGLMs analysis.

Results

Climate Regression

The 72 prioritized species in the *Solanum* section *Petota* examined here occurred in 17 distinct climates with individual species distributions ranging from a single climate (e.g. *S. albornozii*, *S. chilliasense*, *S. lesteri*) to 15 distinct climates (e.g. *S. stoloniferum*). Within this range exists a spectrum of breeding system and ploidy combinations between and within these species and their populations, exhibiting different extents of climate niche diversity (Figure 1). This analysis showed that distinct breeding system and ploidy combinations existed in a different number of niches ($p = 3.4 \times 10^{-7}$), described as the number of discrete Köppen-Geiger climate classes. Species that possess populations that are self-incompatible diploid and self-compatible polyploid show the greatest mean climate diversity with 11 discrete climate classes (Figure 1). Self-incompatible diploid species exhibit a greater average niche diversity when compared to self-compatible diploid species (Figure 1). Furthermore, diploid species possessing populations showing polyploidization demonstrate greater sustained ecological divergence.

The maximum likelihood intercept value of ecological niche diversity is 2.81±1.01 climate classes. Species existing as self-incompatible diploid or self-compatible polyploid have significantly (p-value < 0.01) larger climatic niches by 3.13±0.73 and 3.62±0.79 discrete climate classes, respectively (Table 2). However, other predictor (self-compatible diploid, asexually propagating unknown breeding system diploid) variable slope values are not significantly different from zero, and, therefore, they exert no measurable influence on niche diversity within *Solanum* section *Petota*. Overall, the model explained a moderate amount of variance with an adjusted R-square of 0.39.

Evolutionary Climate Regression

In the PGLMs fitted (Supplemental Table 2) using our estimated time-calibrated phylogeny (Supplemental Figure 2), we found an estimated intercept value of 6.43±1.67 (Table 2). The PGLMs confirmed the correlations of OLS models, with self-incompatible diploid (3.98±1.04) and self-compatible polyploid (2.57±0.98) significantly increasing ecological diversity (Table 3). As with OLS, the other predictor variables in PGLMs are not significantly different from zero.

Discussion

Clarifying the impacts of plant traits on niche divergence is important to understanding the structure of global patterns of biodiversity and evolution in plant lineages (Cavender-Bares et al. 2009). Furthermore, life history traits can provide clues about the potential resiliency of plants with increased development of wild areas leading to changes in habitat and climate for many species. However, resilience may be tightly linked with other characteristics. In *Solanum* section *Petota*, the interaction of two specific characters, breeding system and ploidy, explain a large portion of the variation in niche divergence. The models presented here, OLS and PGLMs, explain 39% and 44% (R-squared), respectively, of the climatic niche variation present within *Solanum* section *Petota* with two alternate ends of the biological spectrum serving as the most significant predictors. On one end, self-incompatible diploid species exhibit the greatest significant correlation to climatic niche diversity within potato wild relatives. Such sustained diversity is likely the result of constant capacity for outcrossing between these species and their subsequent heterogenous design, fashioning an adaptive and resilient population through long-distance gene flow (Loveless and Hamrick 1984). Due to the interaction between ploidy and breeding system, self-incompatible diploid species show niche diversity similar to self-

compatible polyploid species, confirming the dynamic nature of the *Solanaceae* system (Barringer 2007, Husband et al. 2008, Robertson et al. 2010, Zenil-Ferguson et al. 2019). However, self-fertilizing polyploid species have a short-term advantage as they can colonize new environments with very few individuals.

For all the Solanaceae family self-incompatible diploid has been shown to be the ancestral state (Zenil-Ferguson et al. 2019), they also have faster net diversification compared to all self-compatibles, both diploid and polyploid (Wright et al. 2013). The expectation given the success of these lineages in diversification is that self-incompatible diploids should have broader niches, an unexpected result was that self-compatible polyploids diversified in a similar way. Evolutionarily this may be a temporal effect, polyploids are successful in short time scales and this may explain the success in diversification identified here, however, this study does not disentangle evolutionary time-scales. Our results suggest that self-compatible diploids appear evolutionarily transient, and the evolution of self-compatibility appears to occur very rarely without a polyploidy event in Solanaceae. This suggests that polyploidy is just an evolutionary byproduct of trying to become self-compatible, allowing for rapid establishment in many new environments.

Self-compatible polyploid species have increased climatic niche diversity which, given their increased genetic variation and plasticity through additional sets of chromosomes, make them capable of adaptive and resilient population generation (Soltis and Soltis 1999). Polyploidy allows self-fertilizing section *Petota* species to maintain and derive novel diversity typically observed in outcrossing/self-incompatible diploid populations. These differences between breeding system and ploidy with niche diversity provide support for the use of these variable

combinations as driving evolutionary forces, with qualitative results (Figure 1) being supported by OLS (Table 1) and PGLMs (Table 2).

Our results suggest the potential to use ecologically plastic species to enhance the adaptability of cultivated potato lines in the face of climate change. The ultimate goal of this investigation is increased beneficial genetic variation among cultivated potato varieties developed through introgression of the various wild adaptations. However, the wild species have limited cross-compatibility with *S. tuberosum*, as evidenced in their endosperm balance numbers. Therefore, time is needed in order to operationalize this diversity in agricultural fields, so that favorable environmental adaptations from a subset of ecologically plastic species, can be introgressed while breaking linkages to agronomically unfavorable traits.

The impact of breeding system on the evolution of climatic niche diversity amongst plants is still unclear and the *Solanum* section *Petota* system contributes important evidence for a multilayered role where breeding system and ploidy interact synergistically with one another. In one case, self-incompatible breeding systems play a large role in sustaining niche diversity over time (Park et al. 2017) when species are diploid, possessing limited reproductive barriers. In contrast, self-compatible breeding systems comparatively increase niche diversity when species are polyploid, by enhancing their ability to reach, reproduce, establish, and adapt (Campbell 2013) with the biological safety net of increased "buffering capacity" through genetic variation (Wendel et al. 2003). Further investigations could focus on the decoupling of breeding system and ploidy; however, due to the self-incompatibility conferred by S-RNases found in polyploid populations of *Solanaceae* this is challenging (Robertson 2010, Barringer 2007, Husband et al. 2008). Furthermore, this study was not able to completely decouple ploidy and breeding system interactions due to lack of data on particular species' breeding systems, exemplifying the need

for more than DNA collection. Additionally, a limitation of this analysis is the limited number of species available for PGLMs, which was due to a lack of publicly available plastid genome sequence data.

Increasing effective genetic diversity through polyploidization has the potential to increase the number of niches to a similar extent as would occur with an outcrossing diploid population. The breeding system is the main driver of niche divergence in the self-incompatible diploid populations, while only a secondary contributor in the self-compatible polyploid populations. Despite the biological differences, the resulting niche diversity is not seen in a difference of preferred climate type but rather the extent of climatic diversity (Supplemental Figure 1). Through decoupling geographical range size and niche breadth (Randel et al. 2009), this study tests classic theory by utilizing a highly diverse, economically important section of plants. Our findings lend credence to the hypothesis that these ecologically plastic responses evolved over millions of years in species with populations of self-incompatible diploids and self-compatible polyploids, and, therefore, these species should be prioritized for conservation and for use to adapt our cultivated varieties to a changing climate.

Acknowledgements

We would like to the Hawaii Agriculture Research Center and the University of Hawaii Office of Sustainability for their support of Nathan Fumia through the Sustainable Agriculture Fellowship, the Information Technology Systems at the University of Hawai'i at Manoa for computer processing support, and access to data via the Centro Internacional de la Papa. We would like to thank Cornell University for supporting the sabbatical of Dr. Michael A. Gore to contribute to this manuscript.

Figures and Tables

Figure 1. Boxplot of niche diversity by breeding system and ploidy interaction in potato wild relative species. Many species exist containing multiple subpopulations with differing biological factors, as seen by combination of such factors on the x-axis.

Figure 2. Dual figure with time-calibrated molecular clock phylogeny (left) with climatic niche diversity (i.e. number of climate classes occupied) (right). On the left side, the x-axis scale bars represent millions of years and the background coloration of the phylogenetic tree highlights widely accepted clades of *Solanum* section *Petota*. On the right side, the number of climate classes a species occurs in is represented by the size of the horizontal bar and measured with the x-axis scale bar and the coloration of the horizontal bars represent species biological attributes as breeding system with ploidy.

Table 1. Results from the linear model for climatic niche diversity following Gaussian distribution. The number of discrete climate classes in which each taxon can occur (i.e. a proxy for niche breadth) is the response variable, *Climate Niche Diversity*. The predictor variables are combinations of ploidy and breeding system for each species, which were coded as dummy variable interaction terms: self-incompatible diploid, self-compatible diploid, self-compatible asexually propagating polyploid, and unknown breeding system asexually propagating diploid.

Ordinary Least-Squares Results

	Dependent variable:
	Climatic Niche Diversity
Self-Incompatible Diploid	3.134***
	(0.734)
Self-Compatible Diploid	0.569
	(1.021)
Self-Compatible Polyploid (Asexual)	3.624***
	(0.789)
Asexual Diploid	0.883
	(0.992)
Constant	2.813***
	(1.006)
Observations	72
R^2	0.426
Adjusted R^2	0.392
Residual Std. Error	2.493 (df = 67)
F Statistic	12.424*** (df = 4; 67)
Note:	*p **p ***p<0.01

Table 2. Results from phylogenetic linear models for climatic niche diversity following Brownian motion. The number of discrete climate classes in which each taxon can occur (i.e. a proxy for niche breadth) is the response variable, *Climate Niche Diversity*. The predictor variables are combinations of ploidy and breeding system for each species, which were coded as dummy variable interaction terms: self-incompatible diploid, self-compatible diploid, self-compatible asexually propagating polyploid, and unknown breeding system asexually propagating diploid.

Phylogenetic Least-Squares Results

	Dependent variable:
	Climatic Niche Diversity
Self-Incompatible Diploid	3.984^{***}
	(1.036)
Self-Compatible Diploid	1.856
	(1.536)
Self-Compatible Polyploid (Asexual)	2.574^{*}
	(0.975)
Asexual Diploid	-2.332
	(1.620)
Constant	6.426^{***}
	(1.670)
$Sigma^2$	8.088e-09
	(2.967e-09,1.067e-08)
$Sigma^2$ Error	4.497
	(1.650,5.933)
Observations	27
R^2	0.527
Adjusted R^2	0.441
Residual Std. Error	4.761 (df = 22)
Parametric Bootstraps	100
Note:	$^{*}p^{**}p^{***}p<0.01$

References

Adams, K.L. & Wendel, J.F. (2005). Polyploidy and genome evolution in plants. *Current Opinion in Plant Biology*, 8, 135-141.

Baniaga, A. E., Marx, H. E., Arrigo, N., & Barker, M. S. (2020). Polyploid plants have faster rates of multivariate climatic niche evolution than their diploid relatives. *Ecology Letters*, 23, 68–78.

Barringer, BC. (2007). Polyploidy and self-fertilization in flowering plants. *American Journal of Botany*, 94, 1527–1533.

Barringer, B.C., Geber, M.A. (2008). Mating System and Ploidy Influence Levels of Inbreeding Depression in Clarkia (Onagraceae). *Evolution*, 62: 1040-1051.

Barton, N.H. & Kirkpatrick, M. (1996). Evolution of a Species' Range. *The American Naturalist*, 150(1), 1-23.

Bouckaert R., Vaughan T.G., Barido-Sottani J., Duchêne S., Fourment M., Gavryushkina A., et al. (2019) BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLoS computational biology*, 15(4), e1006650.

Campbell, S.A. & Kessler, A. (2013). Plant Mating System Transitions Drive the Macroevolution of Defense Strategies. *PNAS*, 110.10, 3973–3978.

Castañeda-Álvarez, N.P., de Haan, S., Juárez, H., Khoury, C.K., Achicanoy, H.A., Sosa, C.C., et al. (2015). Ex Situ Conservation Priorities for the Wild Relatives of Potato (Solanum L. Section Petota). *PLoS ONE*, 10(4), e0122599.

Cavender-Bares, J., Kozak, K. H., Fine, P. V., Kembel, S. W. (2009). The merging of community ecology and phylogenetic biology. *Ecology letters*, 12(7), 693-715.

De Boyt, S., Maere, S., Van de Peer, Y. (2005). Genome duplication and the origin of angiosperms. *Trends in Ecology & Evolution*, 20(11), 591-597.

Dempewolf, H., Eastwood, R.J., Guarino, L., Khoury, C.K., Müller, J.V., & Toll, J. (2014). Adapting Agriculture to Climate Change: A Global Initiative to Collect, Conserve, and Use Crop Wild Relatives. *Agroecology and Sustainable Food Systems*, 38:4, 369-377.

Grafen, A. (1989). The Phylogenetic Regression. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences, 326*(1233), 119-157.

Grant, A.G. & Kalisz, S. (2020). Do Selfing Species Have Greater Niche Breadth? Support from Ecological Niche Modeling. *Evolution*, 74.1, 73–88.

Grossenbacher, D. et al. (2016). No Association Between Plant Mating System and Geographic Range Overlap. *American Journal of Botany*, 103.1, 110–117.

Hijmans, R. (2020). raster: Geographic Data Analysis and Modeling. R package version 3.1-5. https://CRAN.R-project.org/package=raster

Hijmans, R., Gavrilenko, T., Stephenson, S., Bamberg, J., Salas, A., & Spooner, D.M. (2007). Geographical and environmental range expansion through polyploidy in wild potatoes (Solanum section Petota). *Global Ecology and Biogeography*, 16, 485-495.

Hijmans, R. & Spooner, D.M. (2001). Geographic Distribution of Wild Potato Species. *American Journal of Botany*, 88, 2101-2112.

Hijmans, R., Spooner, D.M., Salas, A.R., Guarino, L, de la Cruz, J. (2002) Atlas of wild potatoes. 130 p. ISBN: 978-92-9043-518-1.

Hlavec, M. (2018). Stargazer: Well-Formatted Regression and Summary Statistics Tables. R package version 5.2.2. https://CRAN.R-project.org/package=stargazer

Ho, L. S. T. & Ane, C. (2014) A linear-time algorithm for Gaussian and non-Gaussian trait evolution models. *Systematic Biology*, 63(3), 397-408.

Huang, B., Ruess, H., Liang, Q., Colleoni, C., & Spooner, D. M. (2019). Analyses of 202 plastid genomes elucidate the phylogeny of Solanum section Petota. *Scientific Reports*, 9(1).

Huelsenbeck, J. P. & Ronquist, F. (2001). MRBAYES: Bayesian inference of phylogeny. *Bioinformatics*, 17, 754-755.

Husband, B., Ozimec, B., Martin, S. & Pollock, L. (2008). Mating Consequences of Polyploid Evolution in Flowering Plants: Current Trends and Insights from Synthetic Polyploids. *International Journal of Plant Sciences*, 169.1, 195-206.

Husband, B.C. and Schemske, D.W. (2017), The Effect of Inbreeding in Diploid and Tetraploid Populations of *Epilobium Angustifolium* (Onagraceae): Implications for the Genetic Basis of Inbreeding Depression. *Evolution*, 51: 737-746.

Jansky, S., Dempewolf, H., Camadro, E., Simon, R., Zimnoch-Guzowska, E., Bisognin, D., & Bonierbale, M. (2013). A Case for Crop Wild Relative Preservation and Use in Potato. *Crop Science*. 53, 746-754.

Johnson, A., Govindarajulu, R., Ashman, T.L. (2014). Bioclimatic evaluation of geographical range in *Fragaria* (Rosaceae): consequences of variation in breeding system, ploidy and species age, *Botanical Journal of the Linnean Society*, 176(1), 99–114.

Katoh, K., Asimenos, G., Toh, H. (2009). Multiple alignment of DNA sequences with MAFFT. *Methods in Molecular Biology*, 537, 39-64.

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., & Drummond, A. (2012). Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*, 28(12), 1647-1649.

Khoury, C.K., Bjorkman, A.D., Dempewolf, H., Ramirez-Villegas, J., Guarino, L., Jarvis, A., Rieseberg, L., and Struik, P.C. (2014). Increasing homogeneity in global food supplies and the implications for food security. *Proceedings of the National Academy of Sciences*, 111(11), 4001-4006.

Köhler, C., Scheid, O., Erilova, A. (2010). The impact of the triploid block on the origin and evolution of polyploid plants. Trends in Genetics. 26(3): 142-148.

Loveless, M.D & Hamrick, J.L. (1984). Ecological Determinants of Genetic Structure in Plant Populations. *Annual Review of Ecology and Systematics*, 15(1984), 65-95.

Martins, E.P. and Hansen, T.F. (1997). Phylogenies and the Comparative Method: A General Approach to Incorporating Phylogenetic Information into the Analysis of Interspecific Data. *The American Naturalist*, 149(4), 646-667.

Mehrabi, Z., Pironon, S., Kantar, M.B., Ramankutty, N., Rieseberg, L. (2019). Shifts in the abiotic and biotic environment of cultivated sunflower under future climate change. *OCL,* 26(9).

Mundry R. (2014) Statistical Issues and Assumptions of Phylogenetic Generalized Least Squares. In: Garamszegi L. (eds) Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology. Springer, Berlin, Heidelberg

Newton, M., Mau, B., & Larget, B. (1999). Markov chain Monte Carlo for the Bayesian analysis of evolutionary trees from aligned molecular sequences. In Statistics in molecular biology (F. Seillier-Moseiwitch, T. P. Speed, and M. Waterman, eds.). Monograph Series of the Institute of Mathematical Statistics.

Paradis E. & Schliep K. (2018). ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. Bioinformatics, 35, 526-528.

Park, D., Ellison, A., & Davis, C. (2017). Selfing species exhibit diminished niche breadth over time. BioRxiv, 157974.

Peterson, A., Soberón, J., & Sánchez-Cordero, V. (1999). Conservatism of Ecological Niches in Evolutionary Time. *Science*, 285(5431), 1265-1267.

Pironon, S., Etherington, T.R., Borrell, J.S., Kühn, N., Macias-Fauria, M., Ondo, I., Tovar, C., Wilkin, P. & Willis, K.J. (2019). Potential adaptive strategies for 29 sub-Saharan crops under future climate change. *Nature Climate Change,* 9, 758–763.

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Randel, A., Slyder, J., Kalisz, S. (2009). Can differences in autonomous selfing ability explain differences in range size among sister-taxa pairs of Collinsia (Plantiginaceae)? An extension of Baker's Law. *New Phytologist*, 183, 618-629.

Revell, L.J. (2012). phytools: An R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution*, 3, 217-223.

Robertson, K., Goldberg, E.E. and Igić, B. (2010). Comparative Evidence for the Correlated Evolution of Polyploidy and Self-Compatibility in Solanaceae. *Evolution*, 65: 139-155.

Roughgarden, J. (1972). Evolution of Niche Width. *The American Naturalist*, 106(952), 683-718.

Rubel, F., & Kottek, M. (2010). Observed and projected climate shifts 1901-2100 depicted by world maps of the Köppen-Geiger climate classification. *Meteorol*, 19, 135-141.

Särkinen, T., Bohs, L., Olmstead, R. G., & Knapp, S. (2013). A phylogenetic framework for evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree. *BMC Evolutionary Biology*, 13(1), 214.

Sessa, E. (2019). Polyploidy as a mechanism for surviving global change. *New Phytologist*, 221, 5-6.

Soltis, D. & Soltis, P. (1999). Polyploidy: recurrent formation and genome evolution. *Trends in Ecology & Evolution*, 14(9), 348-352.

Spooner, D.M., Ghislain, M., Simon, R., Jansky, S., Gavrilenko, T. (2014) Systematics, Diversity, Genetics, and Evolution of Wild and Cultivated Potatoes. *The Botanical Review*, 80.

Spooner, D.M., Hetterscheid, W. (2007). Origins, Evolution, and Group Classification of Cultivated Potatoes. Chapter 13, 285-307.

Spooner, D.M., Hijmans, R. (2001). Potato Systematics and Germplasm Collecting, 1989-2000. *American Journal of Potato Research*, 78, 237-268.

Uyeda, J.C., Zenil-Ferguson, R. and Pennell, M.W. (2018). Rethinking phylogenetic comparative methods. *Systematic Biology*, 67(6), 1091-1109.

Vallejo-Marín, M. & O'Brien, H. E. (2006). Correlated evolution of self-incompatibility and clonal reproduction in Solanum (Solanaceae). *New Phytologist*, 173(2), 415–421.

Wendel, J.F. & Cronn, R.C. (2003). Polyploidy and the Evolutionary History of Cotton. *Botany Publication and Papers*. 23.

Wickham et al. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, 4(43), 1686.

Wright, S. I., Kalisz, S., & Slotte, T. (2013). Evolutionary consequences of self-fertilization in plants. *Proceedings of the Royal Society B: Biological Sciences*, 280(1760), 20130133.

Zenil-Ferguson, R., Burleigh, J.G., Freyman, W.A., Igić, B., Mayrose, I. and Goldberg, E.E. (2019). Interaction among ploidy, breeding system and lineage diversification. *New Phytologist*, 224, 1252-1265.

Supplemental Figures

Supplemental Figure 1. Pie chart of climate class proportions by breeding system/ploidy combinations

Supplemental Figure 2. Time-calibrated uncorrelated relaxed molecular clock phylogeny with outgroups and cultivated varieties of potato included as generated with BEAST2.