1   **The evolution of thermal performance in native and invasive populations of *Mimulus***

2   ***guttatus***

4   Aleah Querns[1,2*], Rachel Wooliver[1,3], Mario Vallejo-Marín[4], Seema Nayan Sheth[1*†]

6   [1] Department of Plant and Microbial Biology, North Carolina State University

7   [2] Division of Biology, Kansas State University

8   [3] Department of Biosystems Engineering & Soil Science, University of Tennessee Knoxville

9   [4] Biological and Environmental Sciences, University of Stirling

12   *Co-corresponding authors

13   E-mails: akquerns@ksu.edu, ssheth3@ncsu.edu

14   †Present address: North Carolina State University, Department of Plant and Microbial Biology

15   Box 7612, NCSU Campus, Raleigh, NC 27695

Abstract

1. The rise of globalization has spread organisms beyond their natural range, allowing further opportunity for species to adapt to novel environments and potentially become invaders. Yet, the role of climatic niche evolution in promoting the success of invasive species remains poorly understood. Here, we use thermal performance curves (TPCs) to test hypotheses about thermal adaptation during the invasion process. First, if novel temperature regimes impose strong selection in the introduced range, invasive populations may evolve distinct TPCs relative to native populations. Second, invasive species may not exhibit specialist-generalist tradeoffs and instead may be "masters-of-all" with high maximum performance and broad TPCs. Third, with sufficient time, standing genetic variation, and temperature-mediated selection, TPCs of native and invasive populations may show parallel evolution in response to thermal gradients.

2. To test these hypotheses, we built TPCs for 18 native (United States) and 13 invasive (United Kingdom) populations of the yellow monkeyflower, *Mimulus guttatus*. We grew clones of multiple genotypes per population across six temperature regimes in growth chambers.

3. Invasive populations have not evolved different thermal optima or performance breadths, providing evidence for evolutionary stasis of thermal performance between the native and invasive ranges after over 200 years post-introduction. Further, both native and invasive populations experienced similar specialist-generalist tradeoffs whereby broad TPCS come at the cost of lower peak performance. Inconsistent with the idea that the degree of thermal specialization varies across spatial or temperature gradients, native and invasive populations did not exhibit adaptive clines in thermal performance breadth with latitude or temperature seasonality. However, thermal optimum increased with mean annual temperature in the native range, indicating some adaptive differentiation among native populations

4. *Synthesis:* These findings suggest that thermal niches were static during the invasion process,

and that general-purpose genotypes, rather than rapid evolution in the introduced range, may

promote invasion.

Key-words: adaptive divergence, evolutionary ecology, invasion ecology, latitudinal gradient,

niche conservatism, phenotypic cline, specialist-generalist tradeoff, thermal performance curve,

thermal tolerance

Introduction

Invasive species are one of the greatest threats to biodiversity in the 21st century, and

while the importance of managing their spread is widely recognized, the factors that contribute to

a species becoming invasive are still debated (Sala et al., 2000; Lee, 2002; van Kleunen,

Dawson, et al., 2015). Baker's classic description of "ideal weeds" suggested that invasive

species originate from "general-purpose" genotypes with broad climatic tolerance (Baker, 1965).

However, it is unknown whether climatic tolerance in a species invasive range is pre-existing,

which predicts niche conservatism, or evolved, which predicts niche lability between the native

and invasive range (Petitpierre et al., 2012; Atwater et al., 2018; Liu et al., 2020). Upon

introduction to a new geographic region, a species may expand, contract, or maintain its climatic

niche space depending on how the set of environmental conditions required to support non-

negative population growth changes from the native to the invasive range (Broennimann et al.,

2012). Similarly, a species average niche conditions (often referred to as niche "centroid" or

"position") may or may not shift in the invasive range (Guisan et al., 2014). If climatic niches are

often conserved between species native and invasive ranges, then building climatic niche models

based on native occurrences to identify areas with high invasion risk holds great promise for

predicting and managing biological invasions (Thuiller et al., 2005; Chapman et al., 2017; Da Re

et al., 2020). Numerous studies have used correlative niche models to assess climatic niche

conservatism across species native and invasive ranges (Peterson et al., 2003; Kriticos et al.,

2013; Da Re et al., 2020), but most have neglected among-population variation in niche

properties and variation in physiological tolerances across climatic gradients resulting from

evolutionary change (some exceptions include Ebeling et al., 2008; Hill et al., 2013). We use an

experimental approach to evaluate the evolution of physiological tolerances within and between

a species native and invasive ranges.

One dimension of a species climatic niche that influences its ability to invade new areas

is temperature. Thermal performance curves (TPCs, Fig. 1A) describe the performance of a

genotype, individual, population, or species across a temperature gradient (Huey & Stevenson,

1979). Though TPCs are not strictly equivalent to thermal niches unless the performance metric

is total fitness, they provide a powerful means of experimentally approximating a species

fundamental thermal niche. Comparisons of TPCs may not explicitly test thermal niche

conservatism between a species native and introduced ranges, but they shed light on whether the

evolution of physiological tolerance plays a role in species abilities to invade new areas. Some

TPC parameters that have been hypothesized to influence adaptation to temperature gradients

include maximum performance, thermal optimum, thermal performance breadth, and area under

the curve (Huey & Stevenson, 1979; Angert et al., 2011; Fig. 1A). Maximum performance is the

highest level of performance achievable, corresponding to the peak of the TPC. Thermal

optimum details the temperature at which maximum performance is achieved (Huey &

Stevenson, 1979), analogous to thermal niche position (or centroid). Comparable to thermal

niche breadth, thermal performance breadth describes the span of temperatures across which a

specified percentage of the maximum performance is achieved (Huey & Stevenson, 1979), with

specialists having narrow breadth, and generalists having wide breadth. Finally, the area under

the curve refers to the total amount of area between a TPC and the temperature axis, analogous to

total thermal niche space.

Evolutionary divergence of TPC parameters between invasive and native populations of

the same species could arise through thermal niche contraction, expansion, and/or shifts in the

introduced range. Consistent with thermal niche shifts, changes in thermal optima may

accompany divergence in local thermal regimes between the native and invasive range

(Angiletta, 2009). Nonetheless, if populations possess broad thermal performance breadths,

and/or if temperature regimes are similar in both ranges, then native and invasive populations

may exhibit similar thermal optima (Guisan et al., 2014). Consistent with niche expansion,

invasive populations could evolve a broader TPC upon introduction to novel thermal regimes

(Zerebecki & Sorte, 2011; Bates et al., 2013). Populations can achieve broad TPCs through a

combination of phenotypic plasticity and standing genetic variation (Sheth & Angert, 2014),

particularly if there is admixture of multiple source populations introduced to the novel range

(Kolbe et al., 2004; Lavergne & Molofsky, 2007). Though genetic bottlenecks during founding

events may constrain the ability of invasive populations to adapt to novel conditions, or cause

reductions in breadth (consistent with thermal niche contraction) due to genetic drift (Kitayama

& Mueller☐Dombois, 1995; Daehler, 2003), invasive populations rarely undergo significant

reductions of genetic diversity relative to native populations, making this scenario unlikely

(Wares et al., 2005; Dlugosch & Parker, 2008). Alternatively, if source populations represent

general-purpose genotypes, they may be predisposed toward having a broad thermal niche which

may be conserved in the invasive range (Ainsworth & Drake, 2020), leading to similar thermal

performance breadths in native and invasive populations. To date, there are few empirical studies

which compare TPCs of native and invasive populations of a single species (but see Comeault et

al., 2020), particularly in plants.

Evolutionary trajectories in the introduced range may be influenced by energetic

tradeoffs in thermal reaction norms (Angilletta et al., 2003). Commonly referred to as "jack-of-

all-trades is a master of none" and specialist-generalist tradeoffs, the ability to persist over a

wider range of temperatures can come at the cost of lower maximum performance (Huey &

Hertz, 1984; Richards et al., 2006; Le Vinh Thuy et al., 2016). Such tradeoffs should result in a

negative relationship between maximum performance and thermal performance breadth across

populations, whereby populations with broader TPCs have lower peak performance due to

additional energetic costs (Dewitt et al., 1998; Angilletta et al., 2003; Richards et al., 2006). This

tradeoff operates under the assumption that the area under the TPC remains constant (Palaima &

Spitze, 2004), but physiological or biochemical alterations could allow higher performance

across a broad range of temperatures (Angilletta et al., 2003; Richards et al., 2006) and cause a

positive relationship between thermal performance breadth and maximum performance. This

phenomenon, which has been referred to as "jack-and-master", or "master of all" (Richards et al.,

2006; Matesanz & Sultan, 2013), may be one mechanism that advances invasion.

Given a temperature or latitudinal gradient, populations with sufficient time and genetic

variation may evolve clines in TPC parameters in response to temperature-mediated selection

(Endler, 1977; Diamond et al., 2017; Campbell□Staton et al., 2018). Failure to account for the

presence of latitudinal or environmental clines can mask inferences of trait evolution between

native and invasive ranges, as divergent selection may occur among populations (Colautti et al.,

2009). Adaptation to local thermal regimes should result in thermal optimum increasing with

mean annual temperature (Angert et al., 2011), and thermal performance breadth increasing with

temperature seasonality (often referred to as the climate variability hypothesis, CVH;

Dobzhansky, 1950; Janzen, 1967; Stevens, 1989; Gutiérrez-Pesquera et al., 2016). Thus, parallel

phenotypic clines across latitudinal or climatic gradients in native and invasive ranges would

indicate that rapid evolution plays a role in the invasion process (Huey et al., 2000; Hernández et

al., 2019; van Boheemen et al., 2019). However, if invasive populations have not had sufficient

time or standing genetic variation for adaptation to novel temperature gradients, or if

temperature-mediated selection in the introduced range is weak, then phenotypic clines may be

shallower in the invasive range than in the native range, or absent altogether (Bhattarai et al.,

2017).

In this study, we compare TPCs of invasive and native perennial populations of the

yellow monkeyflower, *Mimulus guttatus* (Phrymaceae; Fig. 2A,B)*,* also known as *Erythranthe*

guttata (Fraga, 2018; Lowry et al., 2019), to improve our understanding of species invasive

abilities. *Mimulus guttatus* is an herbaceous plant native to western North America, occupying

wet habitats across a broad latitudinal and climatic gradient from Alaska to Northern Mexico

(Fraga, 2018). In 1812, *M. guttatus* was brought to the United Kingdom (UK) for horticultural

use and has subsequently become widespread across the British Isles and to a lesser extent parts

of continental Europe (Preston et al., 2002; Newman, 2015). *Mimulus guttatus* is now considered

invasive in the UK and New Zealand (Vallejo-Marín & Lye, 2013; van Kleunen, Röckle, et al.,

2015; Fraga, 2018). Genomic data suggest that genotypes of *M. guttatus* in the UK originated

from perennial populations in Alaska (Puzey & Vallejo-Marín, 2014; Pantoja et al., 2017),

followed by multiple introductions of perennial populations from across the native range

(Vallejo-Marín et al., 2020). The invasive UK range thus consists of a highly admixed melting

pot of native populations and is not likely constrained by low genetic diversity. Climatic niche

models indicate niche conservatism in the native and invasive European ranges of *M. guttatus*,

though invasive populations occur in a subset of conditions occupied in the native range (Da Re

et al., 2020). Because the non-native populations are locally dominant invaders in the UK

compared to continental Europe, and served as a bridgehead for invasions worldwide (Vallejo-

Marín et al., 2020), we focused exclusively on the UK portion of the invasive range.

Here, we assess hypotheses about whether thermal performance evolution and the ability

to overcome specialist-generalist tradeoffs contribute to the invasion success of *M. guttatus*.

First, we hypothesized that if the species has undergone changes in thermal regimes and/or

genetic variation in its invasive range, there would be changes in thermal optimum and

performance breadth between the native and invasive ranges, against the alternative hypothesis

that these parameters do not differ between ranges (Fig. 1B,C). Since average temperatures are

slightly higher in the native range (Fig. 2C), invasive populations may have evolved lower

thermal optima relative to native populations (Fig. 1B). Further, while admixture in the invasive

range could lead to broader TPCs relative to the native range, reduced temperature variation (Fig.

2C,D) and/or genetic bottlenecks could result in narrower TPCs in the invasive range (Fig. 1C).

Second, we evaluated the hypothesis that the ability to overcome specialist-generalist tradeoffs

has fostered the invasion of *M. guttatus* (Fig. 1D) such that native and invasive populations are

"masters-of-all". Alternatively, populations may exhibit specialist-generalist tradeoffs, predicting

that "jack-of-all-trades is a master of none". Third, given sufficient time, standing genetic

variation, and temperature-mediated selection, we hypothesized parallel evolution in response to

thermal gradients in the invasive range (Fig. 1E-H). We predicted that thermal optimum would

increase with mean annual temperature (Fig. 1F), which decreases with latitude in both ranges

(Fig. 1E; Fig. 2C), and thermal performance breadth would increase with temperature seasonality

(Fig. 1H), which increases with latitude in the native range but decreases with latitude in the

invasive range (Fig. 1G; Fig. 2D). Alternatively, clines may be weaker or absent in the invasive

range (Fig. 1E-G) due to weak selection across a narrow thermal gradient or evolutionary

constraints imposed by bottlenecks.

Materials and Methods

Plant Propagation

In June 2019, we grew seeds from 18 North American perennial and 13 United Kingdom

populations of *Mimulus guttatus* (Fig. 2A,B) in the North Carolina State University Phytotron.

We used an average of 3 seed families from populations spanning a broad latitudinal and thermal

gradient in the native and invasive ranges (see "Supplementary methods" for information

regarding population selection), totaling 95 unique genotypes (Table S1). We sowed 3-5 seeds

from each seed family into 72-cell plug trays filled with Fafard 4P soil (Fafard® 4P Mix, Sun

Gro Horticulture) and topped with germination mix (Sunshine® Redi-Earth Plug & Seedling,

Sun Gro Horticulture), placed them in a Percival LT-105X chamber, and subjected them to a 4°C

cold, dark treatment with daily misting for one week. After one week, we transferred seedlings to

a growth room, with a 16-hour day and 8-hour night photoperiod and a 20/15°C day/night

temperature regime. Cold treatments were repeated for seed families with poor initial

germination, which were primarily from the invasive range. After cold treatments, replanted

families remained in the Percival chamber with 20°C days and colder (5°C) nights for two weeks

before being moved to the growth room. Throughout seedling establishment and the remainder

of the experiment, we watered plants in the growth room with a nutrient solution (NPK +

micronutrients), dumped stagnant water, and randomized trays to reduce location effects twice

per week. Three weeks after planting, seedlings were thinned to one, central-most individual per

cell. Six weeks after planting, individuals were repotted into 8-inch pots with Fafard 4P. These

large plants served as the source of replicate cuttings for thermal performance experiments and

remained in the growth room for the duration of the experiment. We randomized the pots into

new trays, each of which held 2 plants. Eight weeks after planting, we cut the primary stem of

each plant to its base to encourage branching, which is ideal for taking replicate cuttings.

Thermal Performance Experiment

Ten weeks after planting, plants possessed sufficient vegetative growth to support many

cuttings. To generate thermal performance data for each genotype within each population, we

took cuttings of similar sizes from these plants and grew them in 2.5-inch pots within 32-pot

trays. As perennial *M. guttatus* often reproduces clonally in the native and invasive range

(Truscott et al., 2006; van Kleunen, 2007; Pantoja et al., 2018), clones permit performance

measurements of the same genotypes across a range of temperatures. Another advantage of using

cuttings rather than seedlings is that they are likely less prone to maternal effects (reviewed in

Roach & Wulff, 1987). Prior to temperature treatments, we allowed each set of clones 2 weeks to

establish roots. During this period, cuttings were kept within chambers set to 20°C day/15°C

night and bottom-watered daily with a nutrient solution. We randomized cuttings both within and

among chambers twice per week to reduce location effects. We subsequently transferred clones

into a growth chamber programmed to one of six temperature regimes. Over the course of the

experiment, we randomly assigned six temperature treatments to one of three identical Percival

LT-105X chambers (Percival Scientific, Inc.; Perry, Iowa, USA), resulting in two full rounds of

chamber use. For each temperature regime, we took two replicates (clones) of each genotype and

randomized them among six trays. We subjected clones to 16-hour days and 8-hour nights,

according to procedures previously implemented by Paul et al. (2011), Sheth & Angert (2014),

and Wooliver et al. (2020), with one of the following day/night temperature regimes (°C): 10/0,

20/10, 25/15, 30/20, 40/30, and 45/35. Temperature regimes were replicated once, along with an

additional replicate of the 45°C/35°C regime. In total, we took 1,338 cuttings. However, 86

cuttings failed to establish, resulting in a final dataset of 1,252 individual clones from 18 native

and 13 invasive populations, each comprising 54 and 41 genotypes, respectively. We exposed

clones to a given temperature regime for one week, during which trays were sub-irrigated daily

with water to prevent different rates of nutrient uptake at different temperatures.

We measured relative change in stem length over the 7-day temperature treatment. We

chose this short treatment period because previous work with *M. guttatus* has documented

significant growth responses to temperature in one week (Sheth & Angert, 2014). We quantified

the length of the primary stem and the total length of branches (including stolons and side

branches) before and after each temperature treatment (*stem_{in}* and *stem_{out}*, respectively). Total

length of branches was estimated as the number of branches multiplied by the length of an

average branch approximately halfway down the primary stem. We then calculated relative

growth rate (RGR) as a proxy for fitness as the change in total stem length per initial stem length

per day (Eq. 1). Negative RGR values arising when clones lost stem length or branches due to

dieback at temperature extremes were set to zero, and we excluded negative RGR values

resulting from accidental damage. Although RGR is not a measure of total reproductive output,

we used this metric as a proxy for fitness because size has been related to reproductive output in

other *Mimulus* species (Sheth & Angert, 2018).

$$RGR = \frac{(stem_{out} - stem_{in})}{(stem_{in} * number\ of\ days)}$$ (Eq. 1)

Bayesian TPC Models

We performed all analyses in R v. 3.6.2 (R Core Team, 2019). To generate TPCs for

native and invasive populations of *M. guttatus*, we used a hierarchical Bayesian model (R

package performr v0.2; Tittes et al., 2019) to simultaneously fit curves predicting RGR across

temperature for all populations while estimating uncertainty. As this model does not allow

inclusion of random effects, we first averaged total RGR across clones of each genotype at each

temperature. This model also required scaling RGR values by the mean RGR across all data and

centering daytime temperature around zero. Thus, we rescaled model outputs to reflect actual

RGR values and temperatures. To improve the effectiveness of posterior sampling, we altered

the default settings of the model to include a total of 4,000 iterations per chain and a maximum

tree depth of 15 (Gelman et al., 2014). This model utilizes a derivation of Kumaraswamy's

probability density function to fit performance curves across a continuous environmental

gradient. We assessed the fit of Bayesian models using a posterior predictive *P*-value, which uses

a test variable to give the probability that values drawn from the simulated posterior predictive

distribution will exceed the observed values. *P*-values closest to 0.5 indicate adequate fit

between the modeled and observed data (Gelman et al., 2014). The Bayesian *P*-value for the

overall model was 0.53, and *P*-values for each population ranged from 0.2-0.82 (Table S2).

From these models, we obtained estimates of thermal optimum (T_{opt}), thermal

performance breadth ($T_{breadth}$), maximum performance (P_{max}), and area under the curve (AUC)

for each population (Table S3). We selected critical values for $T_{breadth}$ from 100 equally spaced

points along the temperature axis closest to the temperatures corresponding to 50% of P_{max}.

Although these models also generated estimates of critical upper and critical lower thermal limits

(temperatures at which RGR decreases to zero), these estimates extended beyond our

temperature treatments. We thus excluded these parameters from our analyses.

Statistical Analyses for Hypothesis Testing

We tested for divergence in TPC parameters between native and invasive populations of

M. guttatus independent of latitude by conducting post-hoc comparisons of TPC model

iterations. For all iterations, we calculated the mean and 95% credible interval of the pairwise

difference in each TPC parameter between the native and invasive range. We interpreted a

statistically significant difference if the 95% credible interval did not include zero.

We examined the potential for specialist-generalist tradeoffs within either range using a

general linear model with $T_{breadth}$, range (native or invasive), and their interaction as predictors

and P_{max} as the response. We also ran a reduced linear model which removed the interaction term

to account for the possibility of similar tradeoffs in both ranges. We used the Akaike information

criterion (AIC) to assess model fit. For all linear models, we evaluated statistical significance

using α=0.05. A negative relationship between $T_{breadth}$ and P_{max} would indicate a specialist-

generalist tradeoff, whereas a positive relationship would be consistent with a "master-of-all"

scenario. Given a final model with a significant interaction term, we ran separate models for each

range. To determine whether the assumption that the area under TPCs remained constant was

supported, we conducted pairwise comparisons of population-level mean *AUC* based on model

iterations. We used a 95% credible interval to assess significant differences between populations

within each range and between populations of different ranges.

To determine whether TPCs of invasive populations have rapidly adapted to temperature

and latitudinal gradients in the novel range, we compared latitudinal and thermal clines of TPC

parameters in the native and invasive range using general linear models. To test whether T_{opt}

decreased with latitude or increased with mean annual temperature, we modeled T_{opt} as a

function of either latitude or mean annual temperature, range, and their interaction. Similarly, to

evaluate whether $T_{breadth}$ increased with latitude in the native range, decreased with latitude in the

invasive range, and increased with temperature seasonality in both ranges, we modeled $T_{breadth}$ as

a function of latitude or temperature seasonality, range, and their interaction. We used AIC to

compare models with and without the interaction term to account for the possibility of clines of

similar magnitude and direction in both ranges. If the final model for each TPC parameter

included a significant interaction term, we conducted range-specific models with only latitude or

temperature as predictors. Shallower or nonsignificant slopes within the invasive range relative

to the native range would suggest a lack of parallel evolution in response to thermal gradients. In

contrast, a significant effect of latitude or temperature, along with a nonsignificant or absent

interaction term would indicate parallel evolution of clines across ranges, though we note that a

nonsignificant interaction term could also arise due to a lack of statistical power.

Results

Overall Shifts in TPC Parameters Between the Native and Invasive Ranges

We found no support for the hypothesis that *M. guttatus* has evolved in response to novel

thermal regimes within its invasive range via TPC shifts. Pairwise comparisons revealed that T_{opt}

and $T_{breadth}$ exhibited 95% credible intervals that included zero (pairwise difference for T_{opt}= -

0.654°C, 95% CI= -1.405, 0.108; pairwise difference for $T_{breadth}$=0.018°C, 95% CI= -1.191,

1.239), indicating that neither differed significantly between ranges (Fig. 3, Fig. S3).

Additionally, pairwise comparisons showed that *AUC* and P_{max} did not differ between ranges

(pairwise difference for *AUC*= -3.043, 95% CI= -7.485, 1.344; pairwise difference for P_{max} = -

0.071 cm/cm/day, 95% CI= -0.194, 0.046; Fig. S3).

Specialist-Generalist Tradeoffs

Consistent with specialist-generalist tradeoffs, P_{max} decreased with $T_{breadth}$ (Table 1, Fig.

4). The final model included an interaction between $T_{breadth}$ and range (Table 1). However, this

interaction was not statistically significant, suggesting that the direction and strength of the

relationship between $T_{breadth}$ and P_{max} did not differ between the native and invasive ranges

(Table 1). The main effect of range was also non-significant (Table 1). Failing to meet the

assumption that area is constant when there are specialist-generalist tradeoffs, pairwise

comparisons indicate that AUC significantly differed among populations both within and

between ranges (Palaima & Spitze, 2004; Fig. S2).

Clines of TPC Parameters

Overall, latitudinal and thermal clines varied in strength depending on the TPC parameter

and/or the geographic range. The final models for T_{opt} as a function of latitude or mean annual

temperature (MAT) did not include an interaction between latitude or MAT and range (Table 1).

Failing to support our hypothesis, T_{opt} did not decrease with latitude (Table 1, Fig. 5A). Further,

there was no main effect of range when assessing clines of T_{opt} across latitude (Table 1). T_{opt}

increased with MAT, but there was also no main effect of range on T_{opt} when assessing clines

across MAT (Table 1, Fig. 5B). To further dissect the relatively low variance explained by this

model (Adj. R^2=0.155), we conducted range-level models. We found that the cline of T_{opt}

increasing with MAT in our full model was primarily driven by adaptive differentiation within

the native range (β=0.153, P=0.036, Adj. R^2=0.199), whereas there was no relationship between

MAT and T_{opt} in the invasive range (β=0.038, P=0.899, Adj. R^2=-0.089).

The final model of $T_{breadth}$ as a function of latitude included an interaction between

latitude and range (Table 1). Contrary to the prediction that $T_{breadth}$ would increase with latitude

in the native range and decrease with latitude in the invasive range, $T_{breadth}$ showed no

relationship with latitude (Table 1, Fig. 5C). There was not a significant main effect of range on

$T_{breadth}$, and the relationship between latitude and $T_{breadth}$ did not differ between ranges (Table 1).

The final model of $T_{breadth}$ as a function of temperature seasonality did not include an interaction

term (Table 1). We found no cline of $T_{breadth}$ with temperature seasonality (Table 1; Fig. 5D),

failing to support the climate variability hypothesis. Further, there was no main effect of range

on $T_{breadth}$ when assessing clines of $T_{breadth}$ across temperature seasonality (Table 1).

Discussion

We compared thermal performance curves (TPCs) of 18 native and 13 invasive

populations of *Mimulus guttatus* to test key hypotheses about the role of climatic niche evolution

in facilitating biological invasions. First, our results provided no support for the hypothesis that

thermal optimum and breadth vary between the native and invasive ranges (Fig. S3). Second,

failing to support the hypothesis that *M. guttatus* is a "master-of-all," there were specialist-

generalist tradeoffs in both the native and invasive ranges (Table 1, Fig. 4). Finally, contrary to

the hypothesis that native and invasive ranges show parallel phenotypic clines, there was limited

evidence for parallel clines in TPC parameters between ranges (Table 1, Fig. 5). Rather than

supporting the hypothesis that TPC evolution has played a role in facilitating *M. guttatus*

invasion in the UK, these results provide physiological support for thermal niche conservatism in

the invasive range. Below, we discuss the implications of these findings in light of the

evolutionary processes that could contribute to biological invasion.

Evolution of Thermal Performance Curves

We found no support for the hypothesis that invasive populations have undergone

changes in mean TPC parameters relative to native populations (Fig. 3, Fig. S3). Instead, T_{opt} and

$T_{breadth}$ were similar between native and invasive ranges, consistent with the finding of climatic

niche conservatism between the native North American and invasive European ranges of *M.*

guttatus based on niche models (Da Re et al., 2020). However, like many other invasive species

(Liu et al., 2020), the invasive range of *M. guttatus* has undergone substantial niche unfilling (Da

Re et al., 2020), where invasive populations are found in a comparatively small subset of the

climatic conditions (including temperature; Fig. S1) occupied in the native range. Although the

range of temperatures is narrower in the introduced range (Fig. 2C), $T_{breadth}$ did not differ

between ranges (Fig. S3), suggesting that invasive *M. guttatus* populations are poised to occupy

greater climatic niche space should it become available. Maintenance of broad thermal tolerance

in the invasive range may allow *M. guttatus* populations to withstand rapid changes in

temperatures, increasing risks that invasive populations will proliferate under climate change.

Future TPC comparisons that include introduced populations from additional regions would

further contribute to our understanding of the role of thermal niche evolution in the global range

expansion of *M. guttatus*.

One factor that could contribute to the evolutionary stasis of TPC parameters in the

invasive range is insufficient standing genetic variation (Prentis et al., 2008; Dlugosch et al.,

2015; Colautti et al., 2017). Although lower neutral genetic variation in the introduced range

(Puzey & Vallejo-Marín, 2014; Pantoja et al., 2017) could have constrained the rapid evolution

of T_{opt} and $T_{breadth}$ in *M. guttatus*, the highly admixed nature of invasive populations (Vallejo-

Marín et al., 2020) make this possibility unlikely. Instead, genetic variation resulting from

admixed invasive populations likely contributed to the maintenance of broad TPCs and allowed

these populations to easily tolerate the narrower range of temperatures in the UK. In sum, the

invasive populations have yet to undergo significant divergence in TPCs from native populations

due to overlap in thermal niche space, rather than lacking the genetic variation necessary for

evolution to occur. If invasive populations exhibit high admixture, they may spread further via

thermal niche evolution in the face of climate change.

Specialist-Generalist Tradeoffs

We found that TPCs in both ranges were constrained by specialist-generalist tradeoffs,

such that broad TPCs came at the cost of lower maximum performance (Table 1, Fig. 4).

Considering the similar $T_{breadth}$ across ranges, maintenance of broad thermal tolerance in the

narrow thermal conditions of the invasive range may incur significant performance costs via

specialist-generalist tradeoffs. However, variation in *AUC* among populations within each range

(Fig. S2) suggests that the assumption that area remains constant does not hold for *M. guttatus*

(Palaima & Spitze, 2004). As such, the tradeoff between $T_{breadth}$ and P_{max} is only partial.

Although *AUC* and $T_{breadth}$ were not correlated (ρ=-0.2692, *P*= 0.1431), suggesting that thermal

niche space is not affected by variance in $T_{breadth}$, *AUC* was positively correlated with P_{max},

(ρ=0.9767, *P*<0.001). These results imply that variation in TPCs among populations is partly

explained by vertical shifts across a fast-slow spectrum of performance (Izem & Kingsolver,

2005), where some populations achieve a greater *AUC* by having both a high maximum

performance and a broad TPC (Palaima & Spitze, 2004). Thus, variation in TPCs in native and

invasive populations of *M. guttatus* seems to be controlled by both vertical shifts where high-

performing populations are 'masters-of-all' and specialist-generalist tradeoffs where generalist

populations are 'masters-of-none'.

Evolution of Clines

We hypothesized that phenotypic clines exhibited in the native range would be repeated

in the invasive range. Although temperature varies predictably with latitude in the invasive range

(Fig. 2C,D), and there is ample evidence that rapid adaptation can facilitate invasion success

(Oduor et al., 2016), our results did not support the hypothesis of parallel clines. Rather, T_{opt}

increased with mean annual temperature across both ranges, though the amount of variance

explained was modest (*Adj. R²*=0.155). As native populations were the primary driver of the

cline detected in the full model (Fig. 5B), adaptive differentiation of T_{opt} to mean annual

temperature may contribute to the success of this species in its native range. The absence of this

cline in the invasive range implies weaker selection on T_{opt} or insufficient time for adaptation in

the invasive range. Given that invasive populations of *M. guttatus* have maintained similar

$T_{breadth}$ as native populations, selection in the invasive range with a relatively narrow range of

mean annual temperatures would not likely favor adaptive differentiation in T_{opt} (Fig. 2C).

414   Failing to support the climate variability hypothesis (Dobzhansky, 1950; Janzen, 1967;

Stevens, 1989; Gutiérrez-Pesquera et al., 2016), $T_{breadth}$ was not related to latitude or temperature

seasonality in either range. There are many possible explanations for a lack of clines in $T_{breadth}$ in

native populations. First, since six native populations were inland perennials, differing regimes

of temperature seasonality in coastal and inland environments of perennial *M. guttatus* could

mask latitudinal and macroclimatic trends in $T_{breadth}$ (Hall & Willis, 2006; Lowry et al., 2009).

Second, microclimatic variation could cause populations of *M. guttatus* to occupy areas with

either highly similar or drastically different regimes of temperature seasonality, which could

hinder the detection of clines at broader macroclimatic scales (Franco & Nobel, 2003;

Rashkovetsky et al., 2006; De Frenne et al., 2013). While additional analyses of variation in TPC

parameters within populations would be helpful for dissecting this hypothesis, our model was

unable to fit genotype-level curves due to insufficient replication. Finally, high gene flow among

native populations could swamp adaptation of TPCs to local thermal regimes (Paul et al., 2011).

Rather than evolving TPCs in response to thermal variation, native populations may consist of

general-purpose genotypes with wide TPCs which contributed to their successful establishment

as an invasive species (van Kleunen et al., 2011). This conclusion is consistent with the finding

of high rates of gene flow among coastal populations occupying similar latitudes in the native

range relative to our study populations (Twyford & Friedman, 2015). Nonetheless, our findings

are particularly surprising given the evolution of parallel phenotypic clines in the introduced

ranges of several species (van Boheemen et al., 2019; Hernández et al., 2019; McGoey et al.,

2020). Overall, our results suggest that though native *M. guttatus* populations are adaptively

differentiated by thermal optima, populations are equally tolerant across a broad temperature

gradient in western North America. Such homogenization of $T_{breadth}$ among native populations

may have allowed *M. guttatus* to maintain a broad thermal niche space upon introduction.

Caveats

There are many caveats which may have impacted the results of our study. First, our

temperature regimes maintained constant respective day and night temperatures, but in nature

plants experience temperature fluctuations. Second, although we incorporated multiple

measurements of stem and branch growth, populations may vary in their investment in stem and

branch growth versus other growth parameters (such as leaf number or belowground growth) at

different temperatures. Third, we focused on one performance metric, RGR over a week-long

period, but to fully understand performance tradeoffs and variation in performance across

temperatures, multiple performance metrics (e.g., survival and fecundity) should be measured

over longer experimental periods and across multiple life stages. Ultimately, performance

metrics that better approximate lifetime fitness could yield stronger phenotypic clines.

Nevertheless, short-term RGR of cuttings captures performance of clonally reproducing plants

characteristic of perennial *M. guttatus*. Finally, we focused on temperature, one axis of the

climatic niche, but evolution along other niche axes such as precipitation and edaphic properties

could also contribute to invasion success (Hall & Willis, 2006; van Kleunen & Fischer, 2008).

Further, the biotic environment may also differ in the introduced range (Holeski et al., 2013;

Thawley et al., 2018), implying that many factors beyond abiotic conditions can play a part in

adaptation to novel conditions during invasions.

Conclusions

Plant invasions have been widely studied for land management purposes, but little

consensus exists on the ecological and evolutionary processes that facilitate invasion success.

Our findings that thermal optima and breadth are similar in the invasive and native ranges, and

that populations did not exhibit clines in thermal breadth even in the native range, suggest that

general-purpose genotypes, rather than rapid TPC evolution, may have contributed toward

successful establishment in the invasive range. Our results have important implications toward

discovering what traits are favored for invasive organisms. For example, although having a broad

thermal performance curve may aid in initial establishment of populations in a new environment,

specialist-generalist tradeoffs may constrain the further evolution of broader TPCs. By

comparing the evolution of thermal performance across broad environmental gradients in a

species native and invasive range, we can predict which species hold the potential to become

aggressive invaders and forecast how they may fare under rapid climate change. Species that are

genetically predisposed towards a broad thermal tolerance may rapidly expand in their invasive

range as climatic conditions shift. These predictions will be invaluable in preparing for future

species invasions, strengthening our efforts to manage invasive species in the face of rapid

climate change and globalization.

Acknowledgements

We thank B. Caldwell, E. Coughlin, N. Gold, M. Kidd, D. Ryan, J. Torres, E. Vtipil, and

M. Wiegmann for assistance with data collection and plant care. We also thank NCSU Phytotron

staff members for assistance with plant care and chamber maintenance. We are grateful to B.

Blackman, J. Colicchio, J. Coughlan, J. Friedman, L. Holeski, D. Lowry, and M. Rotter for

generously providing seeds for this work. W. Hoffmann, M. Burford-Reiskind, Associate Editor

Peter Alpert, and two anonymous reviewers provided valuable feedback on earlier drafts, and M.

Flores-Vergara provided guidance about plant propagation. This project was funded by USDA

National Institute of Food and Agriculture Hatch 1016272.

Author Contributions

A.Q. and S.N.S. conceived of and designed the study with feedback from M.V.M. A.Q.

collected the data. A.Q. and R.W. analyzed the data. A.Q. and S.N.S. led the writing, with

substantial contributions from R.W. and M.V.M. All authors contributed critically to the drafts

and gave final approval for publication.

Data Availability

All data and scripts associated with this study are available at

https://github.com/akquerns/Guttatus_EvolTPCClines.

References

Ainsworth, A., & Drake, D. R. (2020). Classifying Hawaiian plant species along a habitat

generalist-specialist continuum: Implications for species conservation under climate

change. *PLOS ONE*, *15*(2), e0228573. https://doi.org/10.1371/journal.pone.0228573

Angert, A. L., Sheth, S. N., & Paul, J. R. (2011). Incorporating Population-Level Variation in

Thermal Performance into Predictions of Geographic Range Shifts. *Integrative and*

Comparative Biology, *51*(5), 733–750. https://doi.org/10.1093/icb/icr048

Angiletta, M. J. (2009). *Thermal Adaptation: A Theoretical and Empirical Synthesis*. Oxford

University Press.

Angilletta, M. J., Wilson, R. S., Navas, C. A., & James, R. S. (2003). Tradeoffs and the evolution

of thermal reaction norms. *Trends in Ecology and Evolution*, *18*(5), 234–240.

https://doi.org/10.1016/S0169-5347(03)00087-9

Atwater, D. Z., Ervine, C., & Barney, J. N. (2018). Climatic niche shifts are common in

introduced plants. *Nature Ecology & Evolution*, *2*(1), 34–43.

https://doi.org/10.1038/s41559-017-0396-z

Baker, H. G. (1965). Characteristics and modes of origin of weeds. In H. G. Baker & G. L.

Stebbins (Eds.), *The genetics of colonizing species* (pp. 147–172). Academic Press.

Bates, A. E., McKelvie, C. M., Sorte, C. J. B., Morley, S. A., Jones, N. A. R., Mondon, J. A.,

Bird, T. J., & Quinn, G. (2013). Geographical range, heat tolerance and invasion success

in aquatic species. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1772).

https://doi.org/10.1098/rspb.2013.1958

Bhattarai, G. P., Meyerson, L. A., Anderson, J., Cummings, D., Allen, W. J., & Cronin, J. T.

(2017). Biogeography of a plant invasion: Genetic variation and plasticity in latitudinal

clines for traits related to herbivory. *Ecological Monographs*, *87*(1), 57–75.

https://doi.org/10.1002/ecm.1233

Broennimann, O., Fitzpatrick, M. C., Pearman, P. B., Petitpierre, B., Pellissier, L., Yoccoz, N.

G., Thuiller, W., Fortin, M.-J., Randin, C., Zimmermann, N. E., Graham, C. H., &

Guisan, A. (2012). Measuring ecological niche overlap from occurrence and spatial

environmental data. *Global Ecology and Biogeography*, *21*(4), 481–497.

https://doi.org/10.1111/j.1466-8238.2011.00698.x

Campbell□Staton, S. C., Bare, A., Losos, J. B., Edwards, S. V., & Cheviron, Z. A. (2018).

Physiological and regulatory underpinnings of geographic variation in reptilian cold

tolerance across a latitudinal cline. *Molecular Ecology*, *27*(9), 2243–2255.

https://doi.org/10.1111/mec.14580

Chapman, D. S., Scalone, R., Štefanić, E., & Bullock, J. M. (2017). Mechanistic species

distribution modeling reveals a niche shift during invasion. *Ecology*, *98*(6), 1671–1680.

https://doi.org/10.1002/ecy.1835

Colautti, R. I., Alexander, J. M., Dlugosch, K. M., Keller, S. R., & Sultan, S. E. (2017).

Invasions and extinctions through the looking glass of evolutionary ecology.

Philosophical Transactions of the Royal Society B: Biological Sciences, *372*.

https://doi.org/10.1098/rstb.2016.0031

Colautti, R. I., Maron, J. L., & Barrett, S. C. H. (2009). Common garden comparisons of native

and introduced plant populations: Latitudinal clines can obscure evolutionary inferences.

Evolutionary Applications, *2*(2), 187–199. https://doi.org/10.1111/j.1752-

4571.2008.00053.x

Comeault, A. A., Wang, J., Tittes, S., Isbell, K., Ingley, S., Hurlbert, A. H., & Matute, D. R.

(2020). Genetic diversity and thermal performance in invasive and native populations of

African fig flies. *Molecular Biology and Evolution*, *37*(7), 1893–1906.

https://doi.org/10.1093/molbev/msaa050

Da Re, D., Olivares, A. P., Smith, W., & Vallejo-Marín, M. (2020). Global analysis of ecological

niche conservation and niche shift in exotic populations of monkeyflowers (*Mimulus*

guttatus, M. luteus) and their hybrid (*M. × robertsii*). *Plant Ecology & Diversity*, *13*(2),

133–146. https://doi.org/10.1080/17550874.2020.1750721

Daehler, C. C. (2003). Performance Comparisons of Co-Occurring Native and Alien Invasive

Plants: Implications for Conservation and Restoration. *Annual Review of Ecology,*

Evolution, and Systematics, *34*(1), 183–211.

https://doi.org/10.1146/annurev.ecolsys.34.011802.132403

De Frenne, P., Rodriguez-Sanchez, F., Coomes, D. A., Baeten, L., Verstraeten, G., Vellend, M.,

Bernhardt-Romermann, M., Brown, C. D., Brunet, J., Cornelis, J., Decocq, G. M.,

Dierschke, H., Eriksson, O., Gilliam, F. S., Hedl, R., Heinken, T., Hermy, M., Hommel,

P., Jenkins, M. A., … Verheyen, K. (2013). Microclimate moderates plant responses to

macroclimate warming. *Proceedings of the National Academy of Sciences*, *110*(46),

18561–18565. https://doi.org/10.1073/pnas.1311190110

Dewitt, T. J., Sih, A., & Wilson, D. S. (1998). Costs and limits of phenotypic plasticity. *Trends*

in Ecology & Evolution, *13*(2), 77–81. https://doi.org/10.1016/s0169-5347(97)01274-3

Diamond, S. E., Chick, L., Perez, A., Strickler, S. A., & Martin, R. A. (2017). Rapid evolution of

ant thermal tolerance across an urban-rural temperature cline. *Biological Journal of the*

Linnean Society, *121*(2), 248–257. https://doi.org/10.1093/BIOLINNEAN/BLW047

Dlugosch, K. M., Anderson, S. R., Braasch, J., Cang, F. A., & Gillette, H. D. (2015). The devil is

in the details: Genetic variation in introduced populations and its contributions to

invasion. *Molecular Ecology*, *24*(9), 2095–2111. https://doi.org/10.1111/mec.13183

Dlugosch, K. M., & Parker, I. M. (2008). Founding events in species invasions: Genetic

variation, adaptive evolution, and the role of multiple introductions. *Molecular Ecology*,

17(1), 431–449. https://doi.org/10.1111/j.1365-294X.2007.03538.x

Dobzhansky, T. (1950). Evolution in the Tropics. *American Scientist*, *38*, 209–221.

Ebeling, S. K., Welk, E., Auge, H., & Bruelheide, H. (2008). Predicting the spread of an invasive

plant: Combining experiments and ecological niche model. *Ecography*, *31*(6), 709–719.

https://doi.org/10.1111/j.1600-0587.2008.05470.x

Endler, J. A. (1977). *Geographic Variation, Speciation, and Clines*. Princeton University Press.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology*, *37*(12), 4302–4315.

https://doi.org/10.1002/joc.5086

Fraga, N. S. (2018). Erythranthe guttata*, in Jepson Flora Project*. Jepson EFlora.

https://ucjeps.berkeley.edu/eflora/eflora_display.php?tid=99131

Franco, A. C., & Nobel, P. S. (2003). Effect of Nurse Plants on the Microhabitat and Growth of

Cacti. *Journal of Ecology*, *77*(3), 870–886. https://doi.org/10.2307/2260991

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2014).

Bayesian data analysis (3rd ed.). Chapman and Hall.

Guisan, A., Petitpierre, B., Broennimann, O., Daehler, C., & Kueffer, C. (2014). Unifying niche

shift studies: Insights from biological invasions. *Trends in Ecology & Evolution*, *29*(5),

260–269. https://doi.org/10.1016/j.tree.2014.02.009

Gutiérrez-Pesquera, L. M., Tejedo, M., Olalla-Tárraga, M. A., Duarte, H., Nicieza, A., & Solé,

582   M. (2016). Testing the climate variability hypothesis in thermal tolerance limits of

583   tropical and temperate tadpoles. *Journal of Biogeography*, *43*(6), 1166–1178.

584   https://doi.org/10.1111/jbi.12700

585   Hall, M. C., & Willis, J. H. (2006). Divergent Selection on Flowering Time Contributes to Local

586   Adaptation in *Mimulus guttatus* Populations. *Evolution*, *60*(12), 2466–2477.

587   https://doi.org/10.1111/j.0014-3820.2006.tb01882.x

588   Hernández, F., Poverene, M., Garayalde, A., & Presotto, A. (2019). Re-establishment of

589   latitudinal clines and local adaptation within the invaded area suggest rapid evolution of

590   seed traits in Argentinean sunflower (*Helianthus annuus* L.). *Biological Invasions*, *21*(8),

591   2599–2612. https://doi.org/10.1007/s10530-019-01998-8

592   Hill, M. P., Chown, S. L., & Hoffmann, A. A. (2013). A predicted niche shift corresponds with

593   increased thermal resistance in an invasive mite, *Halotydeus destructor*. *Global Ecology*

594   *and Biogeography*, *22*(8), 942–951. https://doi.org/10.1111/geb.12059

595   Holeski, L. M., Keefover-Ring, K., Bowers, M. D., Harnenz, Z. T., & Lindroth, R. L. (2013).

596   Patterns of Phytochemical Variation in *Mimulus guttatus* (Yellow Monkeyflower).

597   *Journal of Chemical Ecology*, *39*, 525–536. https://doi.org/10.1007/s10886-013-0270-7

598   Huey, R. B., Gilchrist, G. W., Carlson, M. L., Berrigan, D., & Serra, L. (2000). Rapid Evolution

599   of a Geographic Cline in Size in an Introduced Fly. *Science*, *287*(5451), 308–309.

600   https://doi.org/10.1126/science.287.5451.308

601   Huey, R. B., & Hertz, P. E. (1984). Is a Jack-of-All-Temperatures a Master of None? *Evolution*,

602   *38*(2), 441–444. JSTOR. https://doi.org/10.2307/2408502

Huey, R. B., & Stevenson, R. D. (1979). Integrating thermal physiology and ecology of

ectotherms: A discussion of approaches. *Integrative and Comparative Biology*, *19*(1),

357–366. https://doi.org/10.1093/icb/19.1.357

Izem, R., & Kingsolver, J. G. (2005). Variation in continuous reaction norms: Quantifying

directions of biological interest. *American Naturalist*, *166*(2), 277–289.

https://doi.org/10.1086/431314

Janzen, D. H. (1967). Why Mountain Passes are Higher in the Tropics. *The American Naturalist*,

101(919), 233–249. https://doi.org/10.1086/282487

Kitayama, K., & Mueller☐Dombois, D. (1995). Biological invasion on an oceanic island

mountain: Do alien plant species have wider ecological ranges than native species?

Journal of Vegetation Science, *6*(5), 667–674. https://doi.org/10.2307/3236436

Kolbe, J. J., Glor, R. E., Rodriguez, L., Ada, S., & Lara, C. (2004). Genetic variation increases

during biological invasion by a Cuban lizard. *Nature*, *431*, 177–181.

Kriticos, D. J., Morin, L., Leriche, A., Anderson, R. C., & Caley, P. (2013). Combining a

Climatic Niche Model of an Invasive Fungus with Its Host Species Distributions to

Identify Risks to Natural Assets: *Puccinia psidii* Sensu Lato in Australia. *PLOS ONE*,

8(5), e64479. https://doi.org/10.1371/journal.pone.0064479

Lavergne, S., & Molofsky, J. (2007). Increased genetic variation and evolutionary potential drive

the success of an invasive grass. *Proceedings of the National Academy of Sciences*,

104(10), 3883–3888. https://doi.org/10.1073/pnas.0607324104

Le Vinh Thuy, J., VandenBrooks, J. M., & Angilletta, M. J. (2016). Developmental plasticity

evolved according to specialist-generalist trade-offs in experimental populations of

Drosophila melanogaster. *Biology Letters*, *12*(7). https://doi.org/10.1098/rsbl.2016.0379

Lee, C. E. (2002). Evolutionary genetics of invasive species. *Trends in Ecology and Evolution*,

17(8), 386–391. https://doi.org/10.1016/S0169-5347(02)02554-5

Liu, C., Wolter, C., Xian, W., & Jeschke, J. M. (2020). Most invasive species largely conserve

their climatic niche. *Proceedings of the National Academy of Sciences*.

https://doi.org/10.1073/pnas.2004289117

Lowry, D. B., Hall, M. C., Salt, D. E., & Willis, J. H. (2009). Genetic and physiological basis of

adaptive salt tolerance divergence between coastal and inland *Mimulus guttatus*. *New*

Phytologist, *183*(3), 776–788. https://doi.org/10.1111/j.1469-8137.2009.02901.x

Lowry, D. B., Sobel, J. M., Angert, A. L., Ashman, T. L., Baker, R. L., Blackman, B. K.,

Brandvain, Y., Byers, K. J. R. P., Cooley, A. M., Coughlan, J. M., Dudash, M. R.,

Fenster, C. B., Ferris, K. G., Fishman, L., Friedman, J., Grossenbacher, D. L., Holeski, L.

M., Ivey, C. T., Kay, K. M., … Yuan, Y. W. (2019). The case for the continued use of the

genus name *Mimulus* for all monkeyflowers. *Taxon*, *68*(4), 617–623.

https://doi.org/10.1002/tax.12122

Matesanz, S., & Sultan, S. E. (2013). High-performance genotypes in an introduced plant:

Insights to future invasiveness. *Ecology*, *94*(11), 2464–2474. https://doi.org/10.1890/12-

1359.1

McGoey, B. V., Hodgins, K. A., & Stinchcombe, J. R. (2020). Parallel clines in native and

introduced ragweed populations are likely due to adaptation. *Ecology and Evolution*,

4594–4608. https://doi.org/10.1002/ece3.6163

Newman, J. (2015). Rapid Risk Assessment Summary Sheet: Monkey flower (*Mimulus*

guttatus). *GB Non-Native Species Secretariat*, *September*.

Oduor, A. M. O., Leimu, R., & van Kleunen, M. (2016). Invasive plant species are locally

adapted just as frequently and at least as strongly as native plant species. *Journal of*

Ecology, *104*(4), 957–968. https://doi.org/10.1111/1365-2745.12578

Palaima, A., & Spitze, K. (2004). Is a jack-of-all-temperatures a master of none? An

experimental test with *Daphnia pulicaria* (Crustacea: Cladocera). *Evolutionary Ecology*

Research, *6*, 215–225.

Pantoja, P. O., Paine, C. E. T., & Vallejo-Marín, M. (2018). Natural selection and outbreeding

depression suggest adaptive differentiation in the invasive range of a clonal plant.

Proceedings of the Royal Society B: Biological Sciences, *285*(1882).

https://doi.org/10.1098/rspb.2018.1091

Pantoja, P. O., Simón-Porcar, V. I., Puzey, J. R., & Vallejo-Marín, M. (2017). Genetic variation

and clonal diversity in introduced populations of *Mimulus guttatus* assessed by

genotyping at 62 single nucleotide polymorphism loci. *Plant Ecology and Diversity*,

10(1), 5–15. https://doi.org/10.1080/17550874.2017.1287785

Paul, J. R., Sheth, S. N., & Angert, A. L. (2011). Quantifying the impact of gene flow on

phenotype- environment mismatch: A demonstration with the scarlet monkeyflower

Mimulus cardinalis. *American Naturalist*, *178*(S1), S62–S79.

https://doi.org/10.1086/661781

Peterson, A. T., Papes, M., & Kluza, D. A. (2003). Predicting the potential invasive distributions

of four alien plant species in North America. *Weed Science*, *51*(6), 863–868.

https://doi.org/10.1614/P2002-081

Petitpierre, B., Kueffer, C., Broennimann, O., Randin, C., Daehler, C., & Guisan, A. (2012).

Climatic Niche Shifts Are Rare Among Terrestrial Plant Invaders. *Science*, *335*(6074),

1344–1348.

Prentis, P. J., Wilson, J. R. U., Dormontt, E. E., Richardson, D. M., & Lowe, A. J. (2008).

Adaptive evolution in invasive species. *Trends in Plant Science*, *13*(6), 288–294.

https://doi.org/10.1016/j.tplants.2008.03.004

Preston, C. D., Pearman, D. A., & Dines, T. D. (Eds.). (2002). *New Atlas of the British and Irish*

Flora (Vol. 1). Oxford University Press.

Puzey, J., & Vallejo-Marín, M. (2014). Genomics of invasion: Diversity and selection in

introduced populations of monkeyflowers (*Mimulus guttatus*). *Molecular Ecology*,

23(18), 4472–4485. https://doi.org/10.1111/mec.12875

R Core Team. (2019). *R: a language and environment for statistical computing*. R Foundation

for Statistical Computing.

Rashkovetsky, E., Iliadi, K., Michalak, P., Lupu, A., Nevo, E., Feder, M. E., & Korol, A. (2006).

Adaptive differentiation of thermotolerance in *Drosophila* along a microclimatic

gradient. *Heredity*, *96*(5), 353–359. https://doi.org/10.1038/sj.hdy.6800784

Richards, C. L., Bossdorf, O., Muth, N. Z., Gurevitch, J., & Pigliucci, M. (2006). Jack of all

trades, master of some? On the role of phenotypic plasticity in plant invasions. *Ecology*

Letters, *9*, 981–993. https://doi.org/10.1111/j.1461-0248.2006.00950.x

Roach, D. A., & Wulff, R. D. (1987). Maternal effects in plants. *Annual Review of Ecology and*

Systematics, *18*(1), 209–235. https://doi.org/10.1146/annurev.es.18.110187.001233

Sala, O. E., Chapin, F. S., Armesto, J. J., Bloomfield, J., Dirzo, R., Huber-Sanwald, E.,

Huenneke, L. F., Jackson, R. B., Kinzig, A., Leemans, R., Lodge, D. M., Mooney, H. A.,

Oesterheld, M., Poff, N. L., Sykes, M. T., Walker, B. H., Walker, M., & Wall, D. H.

(2000). Global Biodiversity Scenarios for the Year 2100. *Science*, *287*(5459), 1770–

1774. https://doi.org/10.1126/science.287.5459.1770

Sheth, S. N., & Angert, A. L. (2014). The evolution of environmental tolerance and range size: A

comparison of geographically restricted and widespread *Mimulus*. *Evolution*, *68*(10),

2917–2931. https://doi.org/10.1111/evo.12494

Sheth, S. N., & Angert, A. L. (2018). Demographic compensation does not rescue populations at

a trailing range edge. *Proceedings of the National Academy of Sciences*, *115*(10), 2413–

2418. https://doi.org/10.1073/pnas.1715899115

Stevens, G. C. (1989). The Latitudinal Gradient in Geographical Range: How so Many Species

Coexist in the Tropics. *The American Naturalist*, *133*(2), 240–256.

Thawley, C. J., Goldy-Brown, M., McCormick, G. L., Graham, S. P., & Langkilde, T. (2018).

Presence of an invasive species reverses latitudinal clines of multiple traits in a native

species. *Global Change Biology*, *25*(2), 620–628. https://doi.org/10.1111/gcb.14510

Thuiller, W., Richardson, D. M., Pyšek, P., Midgley, G. F., Hughes, G. O., & Rouget, M. (2005).

Niche-based modelling as a tool for predicting the risk of alien plant invasions at a global

scale. *Global Change Biology*, *11*(12), 2234–2250. https://doi.org/10.1111/j.1365-

2486.2005.001018.x

Tittes, S. B., Walker, J. F., Torres-Martínez, L., & Emery, N. C. (2019). Grow Where You

Thrive, or Where Only You Can Survive? An Analysis of Performance Curve Evolution

in a Clade with Diverse Habitat Affinities. *The American Naturalist*, *193*(4), 530–544.

https://doi.org/10.1086/701827

Truscott, A. M., Soulsby, C., Palmer, S. C. F., Newell, L., & Hulme, P. E. (2006). The dispersal

characteristics of the invasive plant *Mimulus guttatus* and the ecological significance of

increased occurrence of high-flow events. *Journal of Ecology*, *94*(6), 1080–1091.

https://doi.org/10.1111/j.1365-2745.2006.01171.x

Twyford, A. D., & Friedman, J. (2015). Adaptive divergence in the monkey flower *Mimulus*

guttatus is maintained by a chromosomal inversion. *Evolution*, *69*(6), 1476–1486.

https://doi.org/10.1111/evo.12663

Vallejo-Marín, M., Friedman, J., Twyford, A. D., Lepais, O., Ickert-Bond, S. M., Streisfeld, M.

A., Yant, L., van Kleunen, M., Rotter, M. C., & Puzey, J. R. (2020). Population genomic

and historical analysis reveals a global invasion by bridgehead processes in *Mimulus*

guttatus. *BioRxiv*. https://doi.org/10.1101/2020.06.26.173286

Vallejo-Marín, M., & Lye, G. C. (2013). Hybridisation and genetic diversity in introduced

Mimulus (Phrymaceae). *Heredity*, *110*, 111–122. https://doi.org/10.1038/hdy.2012.91

van Boheemen, L. A., Atwater, D. Z., & Hodgins, K. A. (2019). Rapid and repeated local

adaptation to climate in an invasive plant. *New Phytologist*, *222*(1), 614–627.

https://doi.org/10.1111/nph.15564

van Kleunen, M. (2007). Adaptive genetic differentiation in life-history traits between

populations of *Mimulus guttatus* with annual and perennial life-cycles. *Evolutionary*

Ecology, *21*(2), 185–199. https://doi.org/10.1007/s10682-006-0019-7

van Kleunen, M., Dawson, W., & Maurel, N. (2015). Characteristics of successful alien plants.

Molecular Ecology, *24*(9), 1954–1968. https://doi.org/10.1111/mec.13013

van Kleunen, M., & Fischer, M. (2008). Adaptive rather than non-adaptive evolution of *Mimulus*

guttatus in its invasive range. *Basic and Applied Ecology*, *9*(3), 213–223.

https://doi.org/10.1016/j.baae.2007.03.006

van Kleunen, M., Röckle, M., & Stift, M. (2015). Admixture between native and invasive

populations may increase invasiveness of *Mimulus guttatus*. *Proceedings of the Royal*

Society B: Biological Sciences, *282*(1815). https://doi.org/10.1098/rspb.2015.1487

van Kleunen, M., Schlaepfer, D. R., Glaettli, M., & Fischer, M. (2011). Preadapted for

invasiveness: Do species traits or their plastic response to shading differ between invasive

and non-invasive plant species in their native range? *Journal of Biogeography*, *38*(7),

1294–1304. https://doi.org/10.1111/j.1365-2699.2011.02495.x

Wares, J., Hughes, A., & Grosberg, R. (2005). Mechanisms that drive evolutionary change:

Insights from species introductions and invasions. In Dov. F. Sax, J. J. Stachowicz, & S.

D. Gaines (Eds.), *Species Invasions: Insights into Ecology, Evolution, and Biogeography*

(pp. 229–257). Sinauer.

Wooliver, R., Tittes, S. B., & Sheth, S. N. (2020). A resurrection study reveals limited evolution

of thermal performance in response to recent climate change across the geographic range

of the scarlet monkeyflower. *Evolution*, *74*(8), 1699–1710.

https://doi.org/10.1111/evo.14041

Zerebecki, R. A., & Sorte, C. J. B. (2011). Temperature tolerance and stress proteins as

mechanisms of invasive species success. *PLoS ONE*, *6*(4).

https://doi.org/10.1371/journal.pone.0014806

Tables

Table 1. Regression coefficients and *P*-values from full general linear models relating response variables to predictors. $\beta 1$ indicates

the main effect of either latitude, mean annual temperature (MAT), thermal performance breadth ($T_{breadth}$), temperature seasonality

(TS), or performance maximum (P_{max}). $\beta 2$ indicates the main effect of range (invasive vs. native). $\beta 3$ indicates the interaction between

range and the predictor corresponding to $\beta 1$. Based on AIC, we chose bolded models as the final model for each respective cline.

	Full Model					
Response	Predictor ($\beta 1$, $\beta 2$, $\beta 3$)	$\beta 1$, *P*	$\beta 2$, *P*	$\beta 3$, *P*	AIC	*Adj. R^2*
P_{max}	**$T_{breadth}$, R, $T_{breadth}$*Range**	**-0.140, P=0.008****	**-1.963, P=0.146**	**0.088, P=0.159**	**-10.448**	**0.219**
	$T_{breadth}$, R	-0.083, P=0.009**	-0.070, P=0.323	---	-10.131	0.188
T_{opt}	Lat, R, Lat*R	0.045, P=0.712	3.812, P=0.590	-0.087, P=0.504	111.848	-0.002
	Lat, R	**-0.033, P=0.409**	**-0.908, P= 0.123**	---	**110.372**	**0.018**
	MAT, R, MAT*R	0.038, P=0.899	-1.738, P=0.521	0.115, P=0.710	107.545	0.128
	MAT, R	**0.148, P=0.028***	**-0.748, P=0.108**	---	**105.708**	**0.155**
$T_{breadth}$	**Lat, R, Lat*R**	**-0.201, P=0.069**	**-11.209, P=0.077**	**0.204, P=0.079**	**103.385**	**0.020**
	Lat, R	-0.018, P=0.623	-0.120, P=0.820	---	104.993	-0.062
	TS, R, TS*R	0.058, P=0.407	1.265, P=0.689	-0.034, P=0.640	104.236	-0.008
	TS, R	**0.026, P=0.117**	**-0.197, P=0.662**	---	**102.492**	**0.020**

762 **=P<0.05, **=P<0.01, ***=P<0.001*

 Figures

Figure 1. Hypotheses describing the evolution of thermal performance curves (TPCs) and clines

in native and invasive populations of *M. guttatus*. (A) TPC parameters of interest include

maximum performance (P_{max}), thermal optimum (T_{opt}), thermal performance breadth ($T_{breadth}$),

and area under the curve (shaded region) (B) Invasive populations may exhibit a lower T_{opt}

relative to native populations (dotted line) due to lower average mean annual temperatures

(MAT; Fig. 2C; native average MAT= 9.2°C, invasive average MAT=8.6°C). Alternatively,

native and invasive populations may exhibit similar T_{opt} if populations occupy similar thermal

regimes or if $T_{breadth}$ is maintained (solid lines) in the invasive range. (C) Invasive populations

may evolve wider (dashed line) or narrower (dotted line) $T_{breadth}$ relative to native populations

due to admixture or bottlenecks/reduced temperature variation, respectively. Alternatively, if

general-purpose genotypes from native populations facilitate invasion, $T_{breadth}$ may be maintained

(solid line). (D) Within native and invasive populations, there may be specialist-generalist

tradeoffs such that populations with greater $T_{breadth}$ have a lower P_{max} (yellow) and populations

with narrower $T_{breadth}$ have a higher P_{max} (orange). However, some invasive species may not

exhibit these tradeoffs and instead achieve a wide $T_{breadth}$ and a high P_{max} (purple). (E-F) The

native range (solid blue line) may exhibit clines such that T_{opt} decreases with latitude (E) and

increases with MAT (F), and these clines may be parallel (solid red) or weaker/nonexistent

(dashed red) in the invasive range. (G-H) The native range may exhibit clines such that $T_{breadth}$

increases with latitude (G) and temperature seasonality (H). The invasive range, where

temperature seasonality increases with latitude (Fig. 2D), may evolve parallel clines (solid line)

such that $T_{breadth}$ decreases with latitude (G) and increases with temperature seasonality (H), or

may exhibit weaker/no clines (dashed line).

Figure 2. Map of focal populations of *M. guttatus* in (A) native range in North America and (B)

invasive range in the United Kingdom. Relationships between (C) mean annual temperature and

(D) temperature seasonality with latitude for native and invasive populations. Fitted lines

indicate predicted mean annual temperature or seasonality as a function of latitude based on

linear models. The gray shaded area represents a 95% confidence interval for these predictions.

Climate data were obtained from WorldClim v.2 (~1-km resolution, 1970-2000, Fick & Hijmans,

2017).

Figure 3. Thermal performance curves of 18 native and 13 invasive populations of *M. guttatus*.

Vertical lines represent thermal optima (T_{opt}) and horizontal lines denote thermal performance

breadth ($T_{breadth}$). Points represent genotype-level mean relative growth rate (RGR) at a given

daytime temperature.

Figure 4. Specialist-generalist tradeoffs as shown by a relationship between maximum

performance (P_{max}) and thermal performance breadth ($T_{breadth}$) across populations of *M. guttatus*.

Black line indicates the main effect of $T_{breadth}$ ($\alpha=0.05$; Table 1) from the full model with the

interaction term removed, and grey shading shows 95% confidence interval.

Figure 5. Relationships between population-level thermal optimum (T_{opt}) or thermal

performance breadth ($T_{breadth}$) and latitude, mean annual temperature (MAT), or temperature

seasonality. A main effect regression line (black) is shown to indicate an overall cline of MAT

($\alpha=0.05$; Table 1) with T_{opt}. However, this cline was driven by significant adaptive differentiation

810   within the native range (solid blue line; $P<0.05$), rather than the invasive range (dashed red line;

811    $P>0.05$).