Title

Combined protein and transcript single cell RNA sequencing in human peripheral blood mononuclear cells

Authors

Jenifer Vallejo,[1#] Ryosuke Saigusa,[1#] Rishab Gulati,[1] Yanal Ghosheh,[1] Christopher P. Durant,[1] Payel Roy,[1] Erik Ehinger,[1] Vasantika Suryawanshi,[1] Tanyaporn Pattarabanjird,[2] Lindsey E. Padgett,[1] Claire E. Olingy,[1] David B. Hanna,[3] Alan L. Landay,[4] Russell P. Tracy,[5] Jason M. Lazar,[6] Wendy J. Mack,[7,8] Kathleen M. Weber,[9] Adaora A. Adimora,[10] Howard N. Hodis,[7,8] Phyllis C. Tien,[11] Igho Ofotokun,[12] Sonya L. Heath,[13] Rafael Blanco-Dominguez[14], Huy Q. Dinh,[15] Avishai Shemesh,[16] Coleen A. McNamara,[2] Lewis L. Lanier,[16] Catherine C. Hedrick,[1] Robert C. Kaplan,[3,17] Klaus Ley.[1,18*]

Affiliations

[1]La Jolla Institute for Immunology, La Jolla, CA, USA.

[2]Carter Immunology Center, Cardiovascular Division, Department of Medicine, University of Virginia, Charlottesville, VA, USA.

[3]Albert Einstein College of Medicine, Department of Epidemiology and Population Health, Bronx, NY, USA.

[4]Rush University Medical Center, Department of Internal Medicine, Chicago, IL, USA.

[5]University of Vermont Larner College of Medicine, Departments of Pathology & Laboratory Medicine and Biochemistry, Colchester, VT, USA.

[6]SUNY Downstate Health Sciences University, Department of Medicine, Brooklyn, NY, USA.

[7]Keck School of Medicine, University of Southern California, Department of Medicine and Preventive Medicine, Los Angeles, CA, USA.

[8]Atherosclerosis Research Unit, University of Southern California, Los Angeles, CA, USA.

[9]Cook County Health/Hektoen Institute of Medicine, Chicago, IL, USA.

[10]Department of Medicine, University of North Carolina School of Medicine, The University of North Carolina at Chapel Hill, Chapel Hill, NC, USA.

[11]Department of Medicine, University of California, San Francisco, San Francisco, CA and Department of Veterans Affairs Medical Center, San Francisco, CA, USA.

[12]Emory University School of Medicine, Department of Medicine, Infectious Disease Division and Grady Health Care System, Atlanta, GA, USA.

[13]University of Alabama at Birmingham, Department of Medicine, Birmingham, AL, USA.

[14]Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain.

[15]McArdle Laboratory for Cancer Research, University of Wisconsin-Madison School of Medicine and Public Health, Madison, Wisconsin, USA.

[16]Parker Institute for Cancer Immunotherapy, University of California, San Francisco, CA, USA; Department of Microbiology and Immunology, University of California, San Francisco, CA, USA.

[17] Fred Hutchinson Cancer Research Center, Public Health Sciences Division, Seattle, WA, USA.

[18] Department of Bioengineering, University of California San Diego, San Diego, CA, USA.

[#] Contributed equally to this work

Corresponding Author

Klaus Ley, MD

La Jolla Institute for Immunology

9420 Athena Circle

La Jolla, CA 92037, USA

(858) 752-6661 (tel)

(858) 752-6985 (fax)

klaus@lji.org

Abstract

Cryopreserved peripheral blood mononuclear cells (PBMCs) are frequently collected and

provide disease- and treatment-relevant data in clinical studies. Here, we developed

combined protein (40 antibodies) and transcript single cell (sc)RNA sequencing in

PBMCs. Among 31 participants in the WIHS Study, we sequenced 41,611 cells. Using

Boolean gating followed by Seurat UMAPs and Louvain clustering, we identified 58

subsets among CD4 T, CD8 T, B, NK cells and monocytes. This resolution was superior

to flow cytometry, mass cytometry or scRNA-sequencing without antibodies. Since the

transcriptome was not needed for cell identification, combined protein and transcript

scRNA-Seq allowed for the assessment of disease-related changes in transcriptomes and

cell type proportion. As a proof-of-concept, we showed such differences between healthy

and matched individuals living with HIV with and without cardiovascular disease. In

conclusion, combined protein and transcript scRNA sequencing is a suitable and

powerful method for clinical investigations using PBMCs.

MAIN TEXT

Introduction

PBMCs are a rich source of disease- and treatment-relevant information.[3, 9, 22, 30, 38,

53, 60–62] PBMCs can be analyzed without mechanical or enzymatic dissociation, which

are known to alter cell surface markers and transcriptomes.[56] PBMCs can be

cryopreserved without loss of viability. At the most basic level, lymphocytes and

monocyte can be distinguished by morphology using automated cell counters (CBC).[4]

Current practice is to use flow cytometry of between 8 and 16 markers.[40, 43, 52] More

recently, mass cytometry became available,[1, 11, 47, 57] allowing for analysis of up to

40 markers. Single cell RNA-sequencing (scRNA-Seq) allows the interrogation of

expressed genes[8, 28, 35, 48, 63] and surface markers.[48, 64]

In immune cells, the correlation between mRNA and surface expression of any given

surface marker is weak.[25] This is because cell surface expression is not only

determined by gene expression, but also by posttranslational protein modifications,[26]

trafficking to the cell surface, protein stability, and proteolytic modifications. Cell types

in PBMCs have been defined by flow cytometry, and the surface markers of the major

cell types are very well known. Yet, it is difficult to call even major cell types by scRNA-

Seq. For example, CD4 T cells are not resolved from CD8 T cells and natural killer (NK)

cells.[59] To capitalize on the vast flow and mass cytometry literature, it is necessary to

assess cell surface phenotype along with transcriptomes.

Only two publications reported single cell transcriptomes from patients with

atherosclerosis (carotid endarterectomy specimens and matched PBMCs).[9, 58] 1,652

103	PBMCs from one patient were analyzed by 10x Genomics 3' and cellular indexing of
104	transcriptional epitope sequencing (CITE-Seq),[35, 48] using a panel of 21 antibodies.
105	No healthy control PBMCs were studied. A recent study reported the effect of HIV
106	infection on PBMC transcriptomes,[20] focusing on acute HIV infection (before
107	antiretroviral therapy started) and reporting PBMC transcriptomes in four patients at 8
108	defined time points (average of 1,976 PBMC transcriptomes per participant and
109	condition). No scRNA-Seq or CITE-Seq studies of PBMCs of people living with chronic
110	HIV infection have been reported. No single cell studies of the interaction between HIV
111	and CVD are available.
112	Here, we report transcriptomes and cell surface phenotypes of almost 42,000 PBMCs
113	using the targeted scRNA-Seq BD Rhapsody platform[8, 28] that simultaneously
114	provides single cell surface phenotype (40 monoclonal antibodies, mAbs) and
115	transcriptomes (485 immune and inflammatory transcripts) in the same cells. As a proof-
116	of-concept, we show significant differences in cell proportions and cell transcriptomes
117	between healthy subjects and matched subjects living with HIV or cardiovascular disease
118	from the WIHS cardiovascular sub-study. WIHS is an ongoing multi-center, prospective,
119	observational cohort study of women with or at risk of HIV infection. PBMCs were
120	cryopreserved on liquid N_2, following strict standard operating procedures that ensured
121	preservation of cell surface phenotype, viability, and transcriptomes.
122	
123	
124	
125	
126	
127	

128 **Material and Methods**
129
Study characteristics and sample selection. The Women's Interagency HIV Study

(WIHS) was initiated in 1994 at six (now expanded to ten) U.S. locations.[13, 16] It is an

ongoing prospective study of over 4,000 women with or at risk of HIV infection.

Recruitment in the WIHS occurred in four phases (1994-1995, 2001-2002, 2010-2012,

and 2013-2015) from HIV primary care clinics, hospital-based programs, and community

outreach and support groups. Briefly, the WIHS involves semi-annual follow-up visits,

during which participants undergo similar detailed examinations, specimen collection,

and structured interviews assessing health behaviors, medical history, and medication

use. All participants provided informed consent, and each site's Institutional Review

Board approved the studies.

All participants in the current analysis were part of a vascular sub-study nested within the

WIHS.[13, 16, 18] The baseline visit for the vascular sub-study occurred between 2004

and 2006, and a follow-up visit occurred on average seven years later. Participants

underwent high-resolution B-mode carotid artery ultrasound to image six locations in the

right carotid artery: the near and far walls of the common carotid artery, carotid

bifurcation, and internal carotid artery. A standardized protocol was used at all sites,[18]

and measurements of carotid artery focal plaque, a marker of subclinical atherosclerosis,

were obtained at a centralized reading center (U. of Southern California). Subclinical

CVD (sCVD) was defined based on the presence of one or more carotid artery

lesions.[18]

From the initial 1,865 participants in the WIHS vascular sub-study, 32 participants were selected for scRNA-seq analysis. CVD was defined as presence of carotid artery focal plaque at either vascular sub-study visit to define four groups of eight participants each: HIV-, HIV+CVD-, HIV+CVD+, HIV+CVD+ on CRT. Because we were interested in the joint relationships of HIV infection and sCVD with surface marker and RNA expression by different cell subtypes, we selected matched samples based on HIV, CVD and cholesterol-reducing treatment (CRT, mostly statins). The latter was done because we found that CRT had a major impact on monocyte transcriptomes.[7]. HIV infection status was ascertained by enzyme-linked immunosorbent assay (ELISA) and confirmed by Western blot. Non-CVD participants with self-reported coronary heart disease or current lipid-lowering therapy use were excluded. Participants were formed in quartets matched by race/ethnicity (except one quartet), age (± 5 years) at the baseline vascular sub-study (except one quartet where the age difference was more but all the women were post-menopausal), visit number, smoking history, and date of specimen collection (within 1 year).

Demographic, clinical, and laboratory variables were assessed from the same study visit using standardized protocols. The median age at the baseline study visit was 55 years, and 96% of participants were either of Black race or Hispanic ethnicity. Most (86%) reported a history of smoking. Substance use was highly prevalent, with 43% of HIV+ and 50% of HIV- participants reporting either a history of injection drug use; current use of crack, cocaine, or heroin; or alcohol use (≥14 drinks per week). Among HIV+ participants, over 80% reported use of HAART at the time PBMCs were obtained, and

59% reported an undetectable HIV-1 RNA level. The median CD4+ T-cell count was

585 cells/μL (IQR 382-816) in HIV+ women without sCVD and 535 cells/μL (IQR 265-

792) in HIV+ women with sCVD.

Preparation of PBMC samples for combined protein and RNA-seq. To avoid batch

effects, sixteen samples each were processed on the same day. PBMC tubes were thawed

in a 37°C water bath and tubes filled with 8 mL of complete RPMI-1640 solution (**Table**

S1; cRPMI-1640 contains human serum albumin, HEPES, sodium pyruvate, MEM-

NEAA, penicillin-streptomicyn, GlutaMax, and mercaptoethanol). The tubes were

centrifuged at 400 xg for 5 minutes and pellets resuspended in cold staining buffer (2 %

fetal bovine serum (FBS) in phosphate-buffered saline (PBS)). All reagents,

manufacturers, and catalogue numbers are listed in **Table S1**. Manual cell counting was

performed by diluting cell concentration to achieve 100-400 cells per hemocytometer

count. Cells were aliquoted to a count of 1 million cells each and incubated on ice with

Fc Block (BD, **Table S1**) at a 1:20 dilution, centrifuged at 400 xg for 5 minutes,

resuspended in 180 μL of SB and transferred to their respective sample multiplexing kit

tubes (BD). The cells were incubated for 20 minutes at room temperature, transferred to 5

mL polystyrene tubes, washed 3 times and centrifuged at 400 xg for 5 minutes. The cells

were resuspended in 400 μL of staining buffer and 2 μL of DRAQ7 and calcein AM were

added to each tube. The viability and cell count of each tube was determined using the

BD Rhapsody scanner (**Table S2**). Tube contents were pooled in equal proportions with

total cell counts not to exceed 1 million cells. The tubes were then centrifuged at 400 xg

for 5 minutes and resuspended in a cocktail of 40 AbSeq (**Table S3**) antibodies (2 μL

each and 20 µL of staining buffer) on ice for 30-60 minutes per manufacturer's

recommendations. The tubes were then washed with 2 mL of SB followed by

centrifugation at 400 xg for 5 minutes. This was repeated two more times for a total of 3

washes. The cells were then counted again using the scanner.

Library preparation. Cells were loaded at 800-1000 cells/µL into the primed plate per

the BD user guide. The beads were isolated with a magnet and the supernatant removed.

Reverse transcription was performed at 37 °C on a thermomixer at 1200 rpm for 20

minutes. Exonuclease I was incubated at 37 °C on a thermomixer at 1200 rpm for 30

minutes and then immediately placed on a heat block at 80 °C for 20 minutes. The tube

was placed on ice followed by supernatant removal while beads were on a magnet. The

beads were resuspended in BD bead resuspension solution. Then, the tubes were stored at

4 °C until further processing. Per BD's protocol, the reagents for PCR1 including the BD

Human Immune Response Panel and a custom panel of ~100 genes (**Table S4**) were

added to the beads. Samples were aliquoted into four 0.2 mL strip PCR tubes and

incubated for 10 cycles according to BD's protocol for PCR1. A double size selection

was performed to remove high genomic DNA fragments by adding 0.7x volume AMPure

XP SPRI beads to the PCR products. After incubation, the supernatant is recovered and

transferred to a new tube followed by purifying the supernatant with an additional 100 µL

of AMPure XP beads (sample tags and antibodies). The RNA tube was washed twice

with 500 µL of 80 % ethanol. 550 µl of supernatant were removed from the antibody tube

followed by two washes with 500 µL of 80 % ethanol. The cDNA was eluted off the

beads using 30 µL of BD elution buffer and then transferred to a 1.5 mL tube.

Pre-sequencing quality control (QC). QC/ and quantification was performed on the

tube containing AbSeq and Sample Tags using Agilent TapeStation high sensitivity

D1000 screentape. 5 μL from each tube (mRNA and Ab/ST) was then added to their

respective tubes containing the reagents for PCR2. Each tube had 12 cycles of PCR

performed according to BD's user guide. Each tube was cleaned with AMPure XP beads

at 0.8X for mRNA and 1.2X for sample tags. Two 200 μL washes were performed during

the clean-up using 80 % ethanol per sample. The cDNA was eluted off using BD elution

buffer. QC/ and quantification was performed using Agilent TapeStation high sensitivity

D1000 screen tape and Qubit double stranded high sensitivity DNA test kit. The mRNA

was then diluted, if necessary, to a concentration of 1.2-2.7 ng/μL and the antibody and

sample tag libraries from PCR2 were diluted, if needed, to a concentration of 0.5-1.1

ng/μL. From each sample 3 μL were added to a volume of 47 μL of reagents for PCR3 as

described by BD's user guide following the protocol and number of cycles listed, except

for AbSeq, which had 9 cycles of PCR performed as determined by previous

optimization. The three libraries were then cleaned with AMPure XP beads at 0.7X for

AbSeq and 0.8X for sample tags. Samples were washed twice with 200 μL of 80 %

ethanol. The cDNA was eluted off the beads using BD's elution buffer. Final QC and

quantification was performed using TapeStation and Qubit kits and reagents.

Sequencing. The samples were pooled and sequenced to the following nominal depth

recommended by BD: AbSeq: n x 1000 reads per cell, where n is the plexity of AbSeq

used; mRNA: 20,000 reads per cell; Sample Tags: 600 reads per cell. Thus, a total of

60,600 reads per cell were desired for sequencing on the NovaSeq. The samples and

specifications for pooling and sequencing depth, along with number of cells loaded onto

each plate was optimized for S1 and S2 100 cycle kits (Illumina) with the configuration

of 67x8x50 bp. Once sequencing was complete, a FASTA file was generated by BD as a

reference for our AbSeq and genes we targeted with these assays. The FASTA file and

FASTQ files generated by the NovaSeq were uploaded to Seven Bridges Genomics

pipeline, where the data was filtered and matrices and csv files were generated. This

analysis generated draft transcriptomes and surface phenotypes of 54,078 cells (496

genes, 40 antibodies). 11 genes were not expressed, leaving 485 genes for analysis.

Doublet Removal. Based on the 4 sample tags used per plate, 8,359 doublets were

removed. The remaining 45,719 cells were analyzed using the Doublet Finder package on

R (https://github.com/chris-mcginnis-ucsf/DoubletFinder) with the default doublet

formation rate (7.5%). This removed another 3,322 doublets, leaving 42,397 Cells.

Finally, we removed all cells that had less than 128 (2^7) antibody molecules sequenced.

This removed 786 noisy cells, resulting in 41,611 cell transcriptomes. All antibody data

were CLR (centered log-ratio) normalized and converted to $\log_2$ scale. All transcripts

were normalized by total UMIs in each cell and scaled up to 1000.

Thresholding. Preliminary experiments showed that each antibody had both specific and

non-specific binding, as expected. To remove the non-specific signal, each antibody

threshold (**Table S5**) was obtained by determining its expression in a known negative

cell. To identify the thresholds, biaxial plots of mutually exclusive markers were used to

best separate the positive populations from the noise. In combined protein and transcript

panel single cell sequencing, non-specific background staining is caused by incomplete

Fc block and oligonucleotide-tagged antibody being trapped in the nanowell.[48]

Ridgeline plots of the unthresholded and thresholded antibody expressions for each main

cell type are shown in **Supplemental Figure S1**, which indicates how the thresholding

worked on each antibody expression.

Clustering. Clustering was performed using UMAP (Uniform Manifold Approximation

and Projection) and Louvain clustering.[50] UMAP is a manifold learning technique for

dimensionality reduction. It is based on neighborhood graphs, which captures the local

relationship in the data. UMAP is able to maintain local structure and also preserve

global distances in the reduced dimension, i.e. the cells that are similar in the high

dimension remain close-by in the 2 dimensions and the cells that different are apart in the

2 dimensions. The clustering parameters used were: n_neighbors = 100, n_pcs = 50,

min_dist = 1, spread = 1, random state = 42. Louvain resolution was set at 0.8.

Subclustering of each major cell type was based on all non-negative antibodies (**Table**

S6). Gates were overlaid and used in all subsequent UMAP figures (cell numbers in

Table S7)

Cluster Assignment. In CD4 T cells, 4 of the initial clusters were further divided based

on the expression of CD11c, CD56, CD25, CD127, CXCR3, and CCR2. CD8 T cells had

two clusters that were divided based on CD11c, CD16, and CXCR3 surface marker

expression. One cluster from classical monocytes and one cluster from intermediate

290	monocytes were further divided based on CCR7 and CD152 expression, respectively. In
291	non-classical monocytes, one cluster showed differential expression of CD36 and CD152
292	expression and was divided in two. In B cells, one cluster was split because it showed
293	differential expression of CD25 and CXCR3 within the cluster. Finally, two clusters from
294	NK cells were split due to CD16, CD56, and CD11c expression.
295	
296	**Comparing Gene Expression among Participant Types.** To determine differential
297	expression (DE) among the four types of participants, we use the Seurat package [49] in
298	R with no thresholds over avg_logFC, minimum fraction of cells required in the two
299	populations being compared, minimum number of cells and minimum number of cells
300	expressing a feature in either group. We filtered for adjusted $p<0.05$ and compared HIV-,
301	HIV+CVD-, HIV+CVD+, and HIV+CVD+CRT+. From this data, volcano plots were
302	generated using ggplot2 and ggrepel packages in R. Axes were restricted to the range of
303	(-2,2) on the x-axis and (0,20) on the y-axis. Genes outside these ranges were bounded to
304	the corresponding limit of the axes. Exact p-values of differential gene expression in all
305	major cell types and the top 10 highly regulated genes for the main cell types are shown
306	in **Table S8** and **S9**, respectively.
307	
308	**Comparing Cell Proportions.** To find changes in proportions, we identified the cell
309	numbers for each participant in each cluster (**Table S10)**. Statistical differences in cell
310	proportions were calculated by log-odds ratio defined as p/(1-p) where p is the proportion
311	of cells, followed by ANOVA and Tukey's multiple comparison test between the four
312	groups. For clarity, the data are presented as percentage of cells.

Correlation Analysis. We correlated each antibody to its corresponding gene(s) using

Spearman rank correlation and significance (R package). For each combination of gene-

antibody, we discarded cells that had values below the corresponding threshold for that

antibody as well as cells with zero counts for that gene. After this filter, any gene-

antibody combination that had 10 cells or less was deemed insignificant. Finally, all non-

significant (p-value > 0.05) were designated a nominal value of zero as the Spearman

rank correlation coefficient and we selected only those genes or antibodies that had at

least one correlation whose coefficient >= 0.25 or whose coefficient <= -0.25. All

significant non-negative correlations are reported in **Table S11**.

Results

Identification of main cell types based on antibody expression. To identify the major

known cell types, we used biaxial gating on CD3, CD19, CD4, CD8, CD14, CD16, and

CD56. This approach defines (**Figure 1A-E**):

- B cells: CD19+ CD3-

- T cells: CD19- CD3+

- CD4 T cells: CD4+ CD8- T cells

- CD8 T cells: CD8+ CD4- T cells

- Monocytes (M): CD19-CD3-CD56-

- Classical (CM): CD14+CD16-

- Intermediate (INT): CD14+CD16+

- Nonclassical (NCM): CD14-CD16+CD56-

- NK cells (NK): CD4- CD56+ CD14- CD20- CD123- CD206-

CD3 and CD19 expression are mutually exclusive and specific for T and B cells,

respectively. As is standard in the NK cell field,[31] the CD16- immature NK cells were

gated based on higher levels of CD56 as shown in **Figure 1E**. The mature NK cells were

CD19-CD3-CD16+CD56+. One CD16+CD56- cluster was also identified as NK cells.

This resulted in 2,919 B cells, 11,045 CD4 T cells, 12,843 CD8 T cells, 5,145 CM, 1009

INT, 475 NCM and 1,843 NK cells. Each of these major cell types was then re-clustered

separately, using Seurat [49] to construct UMAPs with Louvain clustering (**Figure 1F**).

Like in flow or mass cytometry, we clustered on antibody staining only. This "preserves"

the transcriptomes for investigations into disease- and treatment-related changes. Using

this approach, we identified 16 CD4 T cell subsets, 14 CD8 T cell subsets, 8 CM subsets,

3 NCM subsets, 5 INT subsets, 6 B cell subsets and 6 NK cell subsets (**Figure 1F**). The

corresponding feature maps are shown in **Figure S2**. Trying to find these cell types based

on transcriptomes was unsatisfactory (**Figure S3**).

Cell subsets calling using 40 surface markers. Next, we constructed heat maps for all

antibodies that were significantly differentially expressed in at least one subset (**Figure**

2). This information allowed us to call all CD4 and CD8 T cell subsets in accordance

with published immunology work. Among CD4 T cells, CD2 was expressed in almost all

cells, as expected. The high affinity IL2 receptor IL2RA (CD25) was expressed in about

a third of the CD4 T cells and was strikingly high in cluster 13, which was also low for

IL7 receptor (CD127), defining cluster 13 as regulatory T cells (Tregs). CD45RA and RO

were mutually exclusive, separating naive and antigen-experienced CD4 T cells. CXCR3

(CD183) identifies T-helper-1 (Th1) cells and was highly expressed in clusters 5, 14, and

16. Cluster 14 co-expressed CXCR5 (CD185) with CXCR3. Cluster 7 expressed CXCR5

as the only chemokine receptor, suggesting it may contain follicular helper (TFH) T cells.

Based on surface marker information, all CD4 T cell clusters were called (**Figure 2A**).

All CD8 T cells expressed CD2. Cluster 3 exclusively expressed CD9 and CD36,

identifying these cells as NK-like CD8 cells. Clusters 7 and 13 were identified as NK-like

T cells with a CD45RA+ terminally differentiated memory (EMRA) phenotype (**Figure**

2B).

Among monocytes, we were able to call 5 of the 8 classical monocyte subsets based on

published data.[11] All CM were CD11b+ (**Figure 2C**). There were gradients of CD9,

CD69, CD137, CD142 (tissue factor), and CD163 (hemoglobin-haptoglobin receptor)

expression. The scavenger receptor CD36, the antigen presentation co-receptor CD86 and

the chemokine receptor CCR2 were expressed in all classical monocytes. Based on these

markers, 5 of the 8 CM subsets were called (**Figure 2C**) and related to subsets described

by mass cytometry. INT CD14+CD16+ monocytes have been considered pro-

inflammatory and are known to be increased in people with HIV[12] and with CVD.[41,

51] All INT highly expressed the inflammation-induced costimulatory molecule CD86

(**Figure 2C**). Cluster INT3 highly expressed CD142 (tissue factor), which has previously

been implicated in people living with HIV.[45] Since INT subsets have not been

described before, we propose a provisional naming suggestion (**Figure 2C**). NCM

formed 3 clusters (**Figure 2C**). Strikingly, expression of CD9 and CD36 was limited to

cluster 3, suggesting that this cluster corresponds to the previously described

CD9+CD36+ NCM.[11] CD11c, CD74, CD86, and CD141 were expressed in all NCMs

(**Figure 2C**).

We were able to call all 6 B cell subsets. As expected, CD20 and CD74 (HLA-DR) were

expressed in all B cells (**Figure 2D**). CD27, IgM and IgD are used to identify naïve B

cells (CD27-IgM+IgD+). Clusters 1, 3, and 4 were negative for CD27 with high

transcript expression for IgM and IgD, consistent with naïve B cells. Clusters 3 and 4

expressed CCR6, a subset found in HIV+ subjects.[32] B cell cluster 2 expressed CD25,

which is a known marker for B cell proliferation and exhaustion, and CD27, identifying

cluster 2 as a likely activated memory B cell. Cluster 5 had high CD11c levels, known to

increase in HIV-infected subjects,[19] and expressed some CXCR3 and CCR6, but was

CD27low. These features together with moderate expression of CD22 transcript suggest that cluster 5 may contain CD11c+ pathologic B cells. (**Figure 2D**). Most NK cells were mature (CD56^{dim}/CD16+), as expected (**Figure 2E**). Cluster 3 also contained immature (CD56^{bright}CD16-) NK cells. The CD56^{low}CD16- cells (clusters 4 and 5) expressed CD2 and CD45RA. Cluster 5 was CD56-CD16^{high}, an NK cell subset known to be elevated in chronic HIV infection.[17] Taken together, this demonstrates the power of combined antibody and transcriptome sequencing.

Changes in PBMC subset abundance on disease or treatment. Based on this data, it is possible to address shifts in cell proportion based on disease or treatment. We found significant differences in cell proportions in 3 intermediate monocyte subsets, one CD8 T cell, one B cell and one NK cell subset (**Figure 3**). Strikingly, three subsets of intermediate monocytes (**Figure 3A**) showed significantly different abundances. INT2 (IL7R-) and INT3 (TF^{hi}) were significantly elevated in WIHS participants living with HIV and drastically reduced in those that also had subclinical CVD. INT4 had an opposite pattern: These CD163- cells were rare in WIHS participants living with HIV, but more abundant in those that also had subclinical CVD. Among B cells, activated memory B2 cells (**Figure 3C**) were severely lower in all WIHS participants with HIV with or without subclinical CVD.

Differential gene expression in each of the clusters. Since the transcriptomic information was not used for UMAPs and clustering shown in **Figures 1** and **2**, we were able to compare the gene expression patterns in each cell subset within the same cell type.

We filtered for genes that were significantly differentially expressed in at least one of the

subsets (**Figure 4, Data S1**). This analysis revealed gene signatures for most subsets.

Such gene signatures can then be used to determine the presence of each subset in bulk

transcriptomes, and to determine their proportions using Cibersort.[34] As an example,

we applied the classical monocyte transcriptomes (8 subsets) to bulk transcriptomes of

sorted classical monocytes from 92 subjects.[7] We found 1 of the CM subsets in all

subjects and others at varying proportions (**Figure 5**).

Transcriptomes shift with HIV, CVD and cholesterol control. Transcriptomes may

also change with disease state. To test this possibility, we constructed volcano plots (log

fold change on the x axis versus -log p on the y axis (**Figure 6, Data S2**). To test for

changes with cardiovascular disease (CVD), we plotted genes significantly different

between subjects with and without CVD. All these subjects were HIV+. Many genes in

CD4 and CD8 T cell subsets showed significant differences. Some genes in monocyte

and B cell subsets showed significant differences. To test whether our method could

detect effects of treatment, we interrogated transcriptomes of subjects with CVD that

received CRT. Again, many genes in T cell and monocyte subsets and some in B cell

subsets showed significant differences subjects (**Figure 6**). In CD4T1, 2, and 8, IL-32

was highly significantly increased by CVD, but not in CVD+ women on CRT (**Figure 6**).

IL-32 is an inflammatory cytokine that is known to be important in CVD.[6, 21] In

CD4T2, L-selectin (SELL), PSGL-1 (SELPLG), and CCR7 were also highly significantly

increased in WIHS participants with HIV and CVD. In addition to SELL and SELPLG,

CD4T8 showed strong upregulation of TNFSF10 (TRAIL). In CD8T1 and 2, IL32 was

high in women with CVD, but less so in women receiving CRT. Other genes highly

induced by CVD in CD8T1 included CD52, TRAC and HOPX. Several killer cell lectin

receptors (KLRC4, KLRD1, KLRG1 and KLRK1) were also significantly upregulated in

CVD. In CD8T3, CD52, CCL5, IL32 and CD160 were all significantly higher in CVD+

participants. CCL5 encodes the chemokine RANTES, known to be important in

atherosclerosis.[54] In CD8T4, CVD was associated with significantly increased IL32,

TRAC, HOPX, CCL5 and the killer lectin receptors KLRK1, KLRC4, KLRD1.

In CM1, CVD was associated with significantly increased CCL4, SLC2A3, SOD2, and

SELPLG. CRT was associated with lower expression of these genes. In CM2, TNF,

DUSP1, and 2 were highly associated with CVD (**Figure 6**), as were TNFSF10 (TRAIL),

TNFSF13 (APRIL), and TNFSF13B (BAFF), important B cell regulators. In addition to

CCL3, CCL4, and DUSP2, IL1B, known to be highly relevant in atherosclerosis, was

highly upregulated in CM3 of HIV+CVD+ participants. The Toll-like receptor TLR2,

which is known to be involved in atherosclerosis, was upregulated by CVD in CM3. In

INT3, CCL3, CCL4, TNF, IL1B, and DUSP2 were associated CVD in the participants

that did not receive CRT.

Discussion

In immunology, surface markers are widely used to define and distinguish cell types.[5, 47, 57] Flow cytometry is the discipline-defining method of immunology.[40] Similar to flow cytometry, in CyTOF, single-cell suspensions are stained with antibody panels to detect cellular antigens. Unlike CyTOF, scRNA-seq allows the detection of single-cell transcriptomes. Since the correlation between cell surface protein and mRNA expression is weak in immune cells[25], the transcriptome provides a valuable additional dimension. scRNA-Seq without surface phenotype information has led to much frustration in the field, because the expression of many genes encoding well-known surface markers remains undetected in scRNA-Seq.[27, 57] It is still difficult to call cell types based on gene expression data alone, which emphasizes the need for cell surface phenotypes in addition to transcriptomes. Here, we correlated gene expression with cell surface expression for 41 pairs of genes and proteins. CD74 surface expression was well correlated with the expression of both the *CD74* and the *HLA-DRA* genes. CD4 and CD16 surface and gene expression were reasonably well correlated across all cell types. A few other genes including *CD14*, *CD16*, IL-3 receptor (*CD123*), and *CD27* were somewhat correlated with the surface expression of their proteins in some cell types. For most markers, we confirm weak correlations,[25] which illustrates the value of monitoring cell surface phenotype in scRNA-Seq.

PBMCs can be analyzed without mechanical or enzymatic dissociation, which are known to alter cell surface markers and transcriptomes.[56] PBMC are attractive for single cell RNA sequencing (scRNA-Seq) studies, because they are available in many clinical

studies of specific populations with defined diseases and outcomes. The participants

sampled for the present study were part of a sub-study nested within the WIHS,[13, 16,

18] which provided detailed information on subclinical atherosclerosis. Participants

underwent high-resolution B-mode carotid artery ultrasound to image six locations in the

right carotid artery.[16] Although our study is not definitive, it is suggestive of significant

changes in cell proportions and transcriptomes in subjects with cardiovascular disease.

scRNA-Seq has been applied to human PBMCs in diseases including cancers,[3, 60–62]

inflammatory bowel disease,[30, 53] and autoimmune disease,[22, 38] as well as

atherosclerosis.[9, 58] The foundational paper for the 10x Genomics drop-Seq

method[63] demonstrated the feasibility of using scRNA-Seq on PBMCs. Other studies

reported scPBMC transcriptomes in colorectal cancer,[61] γδ T cells,[36] liver

cancer,[62] in vitro salmonella infection,[2] and memory T cells.[28] Only two

publications reported single cell transcriptomes from patients with atherosclerosis

(carotid endarterectomy specimens and matched PBMCs).[9, 58] 1,652 PBMCs from one

patient were analyzed by 10x Genomics 3' and cellular indexing of transcriptional

epitope sequencing (CITE-Seq),[35, 48] using a panel of 21 antibodies. No healthy

control PBMCs were studied. ScRNA-Seq revealed that the process of smooth muscle

cell phenotypic modulation *in vivo* can be altered by the expression of *Tcf21*, a gene

causally associated with reduced risk of coronary artery disease. The loss of *Tcf21* results

in fewer fibromyocytes in the lesions and the protective fibrous cap.[58] A recent study

reported the effect of HIV infection on PBMC transcriptomes,[20] focusing on acute HIV

infection (before antiretroviral therapy started) and reporting PBMC transcriptomes in

four patients at 8 defined time points (average of 1,976 PBMC transcriptomes per

participant and condition). No scRNA-Seq or CITE-Seq studies of PBMCs of people

living with chronic HIV infection have been reported. No single cell studies of the

interaction between HIV and CVD are available.

Six clusters showed significantly different abundance of cells in the four groups of

participants, three of them intermediate monocyte subsets, which underscores the

extraordinary importance of this cell type in chronic HIV infection[14, 29] and CVD.[10,

24] Intermediate monocyte numbers have previously been found increased in non-HIV

subjects with peripheral artery occlusive disease[55] and significantly predicted

cardiovascular events.[15, 41, 42] Cells in INT1, the largest cluster, shared CD11b,

CD11c, CD9, CD36, CD38, CD56, CD69, CD83, IL-3RA, IL6R, CD137, CD141,

CD142 (tissue factor), CXCR4 and CD74 (HLA-DR) with other intermediate monocytes.

We found no single positive marker that was specific for INT1 and thus refrained from

naming this cluster. We found the INT2 and INT3 increased in women living with HIV.

Both express tissue factor (CD142). Tissue factor expression on monocytes has

previously been shown to be increased in HIV+ subjects.[45] Intermediate monocytes are

considered pro-atherogenic,[44] and tissue factor expression provides a plausible reason

for this. We found that in INT2 and INT3, the inflammatory chemokines CCL3 and

CCL4 and the known pro-atherogenic cytokine IL-1β were significantly upregulated in

participants with CVD, but not in those receiving CRT. INT4 uniquely lack expression of

CD163, the receptor for hemoglobin-haptoglobin complexes. Thus, we call INT4 CD163-

intermediate monocytes. INT5, called (CTLA4+CXCR3hi) uniquely expressed CTLA4

(CD152) and highly expressed CXCR3.

In CD4T cells clusters 1, 2 and 8, IL-32 was highly significantly increased by CVD,

which was reversed by cholesterol lowering in CD4T1 and 2 (**Figure 6**). IL-32 is a 27

kDa cytokine expressed in T cells and monocytes that is secreted after apoptosis.[33] It is

an inflammatory cytokine that drives IL-1β, clinically important in CVD,[39] TNF, IL-6

and IL-8 expression.[6, 21, 33] IL-32 activates the leukocyte surface protease PR3, which

in turn triggers the G-protein coupled receptor PAR2[33] and is known to be important in

viral infections.[23, 33, 37, 46] We found IL-32 highly expressed in most T and NK cell

clusters. Since IL-32 appears to be CVD-specific, we advocate for future prospective

studies in larger cohorts to determine whether IL32 mRNA is a useful biomarker.

Our discovery study will encourage prospective epidemiological studies to address which

PBMC subset and transcriptomes are best suited as clinical biomarkers to stratify risk and

guide treatment in subjects with coronary or peripheral artery disease. The current

findings also present some limitations. They need to be extended to men (the current data

is based on women) and other races and ethnicities (the current data is based on mostly

African American and Hispanic women). Studies of CVD in non-smokers are also

needed (the current data is based on smokers), and the age range needs to be broadened.

In conclusion, we demonstrate the utility of scRNA-Seq with cell surface phenotype

assessment in the same cells. The identification of 58 distinct clusters of CD4 and CD8 T

cells, B cells, NK cells and monocytes helps to gain a deeper understanding of PBMCs, a

rich and readily accessible source of biological and clinical information. The discovery of

subsets of intermediate monocytes calls for identifying such subsets in model organisms

to test their function *in vivo*.

References

1. Bekele Y, Lakshmikanth T, Chen Y, Mikes J, Nasi A, Petkov S, Hejdeman B, Brodin P, Chiodi F (2019) Mass cytometry identifies distinct CD4+ T cell clusters distinguishing HIV-1-infected patients according to antiretroviral therapy initiation. JCI insight 4: e125442. doi: 10.1172/jci.insight.125442.

2. Bossel Ben-Moshe N, Hen-Avivi S, Levitin N, Yehezkel D, Oosting M, Joosten LAB, Netea MG, Avraham R (2019) Predicting bacterial infection outcomes using single cell RNA-sequencing analysis of human immune cells. Nat Commun 10:3266. doi: 10.1038/s41467-019-11257-y.

3. Brown CC, Gudjonson H, Pritykin Y, Deep D, Lavallée V-P, Mendoza A, Fromme R, Mazutis L, Ariyan C, Leslie C, Pe'er D, Rudensky AY (2019) Transcriptional Basis of Mouse and Human Dendritic Cell Heterogeneity. Cell 179:846-863. doi: 10.1016/j.cell.2019.09.035.

4. Buttarello M, Plebani M (2008) Automated blood cell counts: state of the art. Am J Clin Pathol 130:104–116. doi: 10.1309/EK3C7CTDKNVPXVTN.

5. Cole JE, Park I, Ahern DJ, Kassiteridi C, Danso Abeam D, Goddard ME, Green P, Maffia P, Monaco C (2018) Immune cell census in murine atherosclerosis: cytometry by time of flight illuminates vascular myeloid cell diversity. Cardiovasc Res 114:1360–1371. doi: 10.1093/cvr/cvy109.

6. Damen MSMA, Popa CD, Netea MG, Dinarello CA, Joosten LAB (2017) Interleukin-32 in chronic inflammatory conditions is associated with a higher risk of cardiovascular diseases. Atherosclerosis 264:83–91. doi: 10.1016/j.atherosclerosis.2017.07.005.

7. Ehinger E, Ghosheh Y, Bala PA, Durant CP, Lin J, Hanna DB, Mueller K, Baas L, Qi Q, Wang T, Buscher K, Liu Y, Anastos K, Lazar JM, Mack WJ, Tien PC, Cohen MH, Ofotokun I, Gange S, Heath SL, Hodis HN, Tracy RP, Landay AL, Kaplan RC, Ley K (2020) Classical Monocyte Transcriptomes Reveal Significant Anti- Inflammatory Statin Effect in Women with Chronic HIV. Cardiovasc Res 117:1166–1177. doi: 10.1093/cvr/cvaa188.

8. Fan HC, Fu GK, Fodor SPA (2015) Expression profiling. Combinatorial labeling of single cells for gene expression cytometry. Science 347:1258367. doi: 10.1126/science.1258367.

9. Fernandez DM, Rahman AH, Fernandez NF, Chudnovskiy A, Amir ED, Amadori L, Khan NS, Wong CK, Shamailova R, Hill CA, Wang Z, Remark R, Li JR, Pina C, Faries C, Awad AJ, Moss N, Bjorkegren JLM, Kim-schulze S, Gnjatic S, Ma A, Mocco J, Faries P, Merad M, Giannarelli C (2019) Single-cell immune landscape of human atherosclerotic plaques. Nat Med 25:1576–1588. doi: 10.1038/s41591-019-0590-4.

10. Funderburg NT, Zidar DA, Shive C, Lioi A, Mudd J, Musselwhite LW, Simon DI, Costa MA, Rodriguez B, Sieg SF, Lederman MM (2012) Shared monocyte subset phenotypes in HIV-1 infection and in uninfected subjects with acute coronary syndrome. Blood 120:4599–4608. doi: 10.1182/blood-2012-05-433946.

11. Hamers AAJ, Dinh HQ, Thomas GD, Marcovecchio P, Blatchley A, Nakao CS, Kim C, McSkimming C, Taylor AM, Nguyen AT, McNamara CA, Hedrick CC (2019) Human Monocyte Heterogeneity as Revealed by High-Dimensional Mass Cytometry. Arterioscler Thromb Vasc Biol 39:25–36. doi: 10.1161/ATVBAHA.118.311022.

12.   Han J, Wang B, Han N, Zhao Y, Song C, Feng X, Mao Y, Zhang F, Zhao H, Zeng H
(2009) CD14(high)CD16(+) rather than CD14(low)CD16(+) monocytes correlate with
disease progression in chronic HIV-infected patients. J Acquir Immune Defic Syndr
52:553–559. doi: 10.1097/qai.0b013e3181c1d4fe.
13.   Hanna DB, Post WS, Deal JA, Hodis HN, Jacobson LP, Mack WJ, Anastos K, Gange SJ,
Landay AL, Lazar JM, Palella FJ, Tien PC, Witt MD, Xue X, Young MA, Kaplan RC,
Kingsley LA (2015) HIV Infection Is Associated With Progression of Subclinical Carotid
Atherosclerosis. Clin Infect Dis 61:640–650. doi: 10.1093/cid/civ325.
14.   Hearps AC, Maisa A, Cheng W-J, Angelovich TA, Lichtfuss GF, Palmer CS, Landay AL,
Jaworowski A, Crowe SM (2012) HIV infection induces age-related changes to
monocytes and innate immune activation in young men that persist despite combination
antiretroviral therapy. AIDS 26:843–853. doi: 10.1097/QAD.0b013e328351f756.
15.   Heine GH, Ulrich C, Seibert E, Seiler S, Marell J, Reichart B, Krause M, Schlitt A, Köhler
H, Girndt M (2008) CD14(++)CD16+ monocytes but not total monocyte numbers predict
cardiovascular events in dialysis patients. Kidney Int 73:622–629. doi:
10.1038/sj.ki.5002744.
16.   Hodis HN, Mack WJ, Lobo RA, Shoupe D, Sevanian A, Mahrer PR, Selzer RH, Liu Cr
CR, Liu Ch CH, Azen SP (2001) Estrogen in the prevention of atherosclerosis. A
randomized, double-blind, placebo-controlled trial. Ann Intern Med 135:939–953. doi:
10.7326/0003-4819-135-11-200112040-00005.
17.   Hong HS, Eberhard JM, Keudel P, Bollmann BA, Ahmad F, Ballmaier M, Bhatnagar N,
Zielinska-Skowronek M, Schmidt RE, Meyer-Olson D (2010) Phenotypically and
functionally distinct subsets contribute to the expansion of CD56-/CD16+ natural killer
cells in HIV infection. AIDS 24:1823–1834. doi: 10.1097/QAD.0b013e32833b556f.
18.   Kaplan RC, Kingsley LA, Gange SJ, Benning L, Jacobson LP, Lazar J, Anastos K, Tien
PC, Sharrett AR, Hodis HN (2008) Low CD4+ T-cell count as a major atherosclerosis risk
factor in HIV-infected women and men. AIDS 22:1615–1624. doi:
10.1097/QAD.0b013e328300581d.
19.   Karnell JL, Kumar V, Wang J, Wang S, Voynova E, Ettinger R (2017) Role of CD11c(+)
T-bet(+) B cells in human health and disease. Cell Immunol 321:40–45. doi:
10.1016/j.cellimm.2017.05.008.
20.   Kazer SW, Aicher TP, Muema DM, Carroll SL, Ordovas-Montanes J, Miao VN, Tu AA,
Ziegler CGK, Nyquist SK, Wong EB, Ismail N, Dong M, Moodley A, Berger B, Love JC,
Dong KL, Leslie A, Ndhlovu ZM, Ndung'u T, Walker BD, Shalek AK (2020) Integrated
single-cell analysis of multicellular immune dynamics during hyperacute HIV-1 infection.
Nat Med 26:511–518. doi: 10.1038/s41591-020-0799-2.
21.   Kim S-H, Han S-Y, Azam T, Yoon D-Y, Dinarello CA (2005) Interleukin-32: a cytokine
and inducer of TNFalpha. Immunity 22:131–142. doi: 10.1016/j.immuni.2004.12.003.
22.   Kotliarov Y, Sparks R, Martins AJ, Mule MP, Lu Y, Goswami M, Kardava L, Banchereau
R, Pascual V, Biancotto A, Chen J, Schwartzberg PL, Bansal N, Liu CC, Cheung F, Moir
S, Tsang JS (2020) Broad immune activation underlies shared set point signatures for
vaccine responsiveness in healthy individuals and disease activity in patients with lupus.
Nat Med 26:618–629. doi: 10.1038/s41591-020-0769-8.
23.   Li W, Sun W, Liu L, Yang F, Li Y, Chen Y, Fang J, Zhang W, Wu J, Zhu Y (2010) IL-32:
a host proinflammatory factor against influenza viral replication is upregulated by aberrant
epigenetic modifications during influenza A virus infection. J Immunol 185:5056–5065.

doi: 10.4049/jimmunol.0902667.

24.   Liang H, Xie Z, Shen T (2017) Monocyte activation and cardiovascular disease in HIV
infection. Cell Mol Immunol 14:960–962. doi: 10.1038/cmi.2017.109.

25.   Liu Y, Beyer A, Aebersold R (2016) On the Dependency of Cellular Protein Levels on
mRNA Abundance. Cell 165:535–550. doi: 10.1016/j.cell.2016.03.014.

26.   Londino JD, Gulick DL, Lear TB, Suber TL, Weathington NM, Masa LS, Chen BB,
Mallampalli RK (2017) Post-translational modification of the interferon-gamma receptor
alters its stability and signaling. Biochem J 474:3543–3557. doi: 10.1042/BCJ20170548.

27.   Lundberg E, Fagerberg L, Klevebring D, Matic I, Geiger T, Cox J, Algenas C, Lundeberg
J, Mann M, Uhlen M (2010) Defining the transcriptome and proteome in three
functionally different human cell lines. Mol Syst Biol 6:450. doi: 10.1038/msb.2010.106.

28.   Mair F, Erickson JR, Voillet V, Simoni Y, Bi T, Tyznik AJ, Martin J, Gottardo R, Newell
EW, Prlic M (2020) A Targeted Multi-omic Analysis Approach Measures Protein
Expression and Low-Abundance Transcripts on the Single-Cell Level. Cell Rep
31:107499. doi: 10.1016/j.celrep.2020.03.063.

29.   Martin GE, Gouillou M, Hearps AC, Angelovich TA, Cheng AC, Lynch F, Cheng W-J,
Paukovics G, Palmer CS, Novak RM, Jaworowski A, Landay AL, Crowe SM (2013) Age-
associated changes in monocyte and innate immune activation markers occur more rapidly
in HIV infected women. PLoS One 8:e55279. doi: 10.1371/journal.pone.0055279.

30.   Martin JC, Chang C, Boschetti G, Ungaro R, Giri M, Grout JA, Gettler K, Chuang L-S,
Nayar S, Greenstein AJ, Dubinsky M, Walker L, Leader A, Fine JS, Whitehurst CE,
Mbow ML, Kugathasan S, Denson LA, Hyams JS, Friedman JR, Desai PT, Ko HM,
Laface I, Akturk G, Schadt EE, Salmon H, Gnjatic S, Rahman AH, Merad M, Cho JH,
Kenigsberg E (2019) Single-Cell Analysis of Crohn's Disease Lesions Identifies a
Pathogenic Cellular Module Associated with Resistance to Anti-TNF Therapy. Cell
178:1493-1508. doi: 10.1016/j.cell.2019.08.008.

31.   Milush JM, Long BR, Snyder-Cappione JE, Cappione AJ 3rd, York VA, Ndhlovu LC,
Lanier LL, Michaelsson J, Nixon DF (2009) Functionally distinct subsets of human NK
cells and monocyte/DC-like cells identified by coexpression of CD56, CD7, and CD4.
Blood 114:4823–4831. doi: 10.1182/blood-2009-04-216374.

32.   Moir S, Fauci AS (2017) B-cell responses to HIV infection. Immunol Rev 275:33-48. doi:
10.1111/imr.12502.

33.   Nakayama M, Niki Y, Kawasaki T, Takeda Y, Ikegami H, Toyama Y, Miyamoto T (2013)
IL-32-PAR2 axis is an innate immunity sensor providing alternative signaling for LPS-
TRIF axis. Sci Rep 3:2960. doi: 10.1038/srep02960.

34.   Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, Hoang CD, Diehn M,
Alizadeh AA (2015) Robust enumeration of cell subsets from tissue expression profiles.
Nat Methods 12:453–457. doi: 10.1038/nmeth.3337.

35.   Peterson VM, Zhang KX, Kumar N, Wong J, Li L, Wilson DC, Moore R, McClanahan
TK, Sadekova S, Klappenbach JA (2017) Multiplexed quantification of proteins and
transcripts in single cells. Nat Biotechnol 35:936–939. doi: 10.1038/nbt.3973.

36.   Pizzolato G, Kaminski H, Tosolini M, Franchini D-M, Pont F, Martins F, Valle C,
Labourdette D, Cadot S, Quillet-Mary A, Poupot M, Laurent C, Ysebaert L, Meraviglia S,
Dieli F, Merville P, Milpied P, Dechanet-Merville J, Fournie J-J (2019) Single-cell RNA
sequencing unveils the shared and the distinct cytotoxic hallmarks of human TCRVdelta1
and TCRVdelta2 gammadelta T lymphocytes. Proc Natl Acad Sci U S A 116:11906–

11915. doi: 10.1073/pnas.1818488116.

37. Rasool ST, Tang H, Wu J, Li W, Mukhtar MM, Zhang J, Mu Y, Xing HX, Wu J, Zhu Y
(2008) Increased level of IL-32 during human immunodeficiency virus infection
suppresses HIV replication. Immunol Lett 117:161–167. doi: 10.1016/j.imlet.2008.01.007.

38. Reyes M, Vickers D, Billman K, Eisenhaure T, Hoover P, Browne EP, Rao DA, Hacohen
N, Blainey PC (2019) Multiplexed enrichment and genomic profiling of peripheral blood
cells reveal subset-specific immune signatures. Sci Adv 5:eaau9223. doi:
10.1126/sciadv.aau9223.

39. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne C, Fonseca F,
Nicolau J, Koenig W, Anker SD, Kastelein JJP, Cornel JH, Pais P, Pella D, Genest J,
Cifkova R, Lorenzatti A, Forster T, Kobalava Z, Vida-Simiti L, Flather M, Shimokawa H,
Ogawa H, Dellborg M, Rossi PRF, Troquay RPT, Libby P, Glynn RJ (2017)
Antiinflammatory Therapy with Canakinumab for Atherosclerotic Disease. N Engl J Med
377:1119–1131. doi: 10.1056/NEJMoa1707914.

40. Robinson JP, Roederer M (2015) HISTORY OF SCIENCE. Flow cytometry strikes gold.
Science 350:739–740. doi: 10.1126/science.aad6770.

41. Rogacev KS, Cremers B, Zawada AM, Seiler S, Binder N, Ege P, Große-Dunker G,
Heisel I, Hornof F, Jeken J, Rebling NM, Ulrich C, Scheller B, Böhm M, Fliser D, Heine
GH (2012) CD14++CD16+ monocytes independently predict cardiovascular events: a
cohort study of 951 patients referred for elective coronary angiography. J Am Coll Cardiol
60:1512–1520. doi: 10.1016/j.jacc.2012.07.019.

42. Rogacev KS, Seiler S, Zawada AM, Reichart B, Herath E, Roth D, Ulrich C, Fliser D,
Heine GH (2011) CD14++CD16+ monocytes and cardiovascular outcome in patients with
chronic kidney disease. Eur Heart J 32:84–92. doi: 10.1093/eurheartj/ehq371.

43. De Rosa SC, Brenchley JM, Roederer M (2003) Beyond six colors: A new era in flow
cytometry. Nat Med 9:112–117. doi: 10.1038/nm0103-112.

44. SahBandar IN, Ndhlovu LC, Saiki K, Kohorn LB, Peterson MM, D'Antoni ML,
Shiramizu B, Shikuma CM, Chow DC (2020) Relationship between Circulating
Inflammatory Monocytes and Cardiovascular Disease Measures of Carotid Intimal
Thickness. J Atheroscler Thromb 27:441–448. doi: 10.5551/jat.49791.

45. Schechter ME, Andrade BB, He T, Richter GH, Tosh KW, Policicchio BB, Singh A,
Raehtz KD, Sheikh V, Ma D, Brocca-Cofano E, Apetrei C, Tracy R, Ribeiro RM, Sher A,
Francischetti IMB, Pandrea I, Sereti I (2017) Inflammatory monocytes expressing tissue
factor drive SIV and HIV coagulopathy. Sci Transl Med 9:eaam5441. doi:
10.1126/scitranslmed.aam5441.

46. Smith AJ, Toledo CM, Wietgrefe SW, Duan L, Schacker TW, Reilly CS, Haase AT
(2011) The immunosuppressive role of IL-32 in lymphatic tissue during HIV-1 infection.
J Immunol 186:6576–6584. doi: 10.4049/jimmunol.1100277.

47. Spitzer MH, Nolan GP (2016) Mass Cytometry: Single Cells, Many Features. Cell
165:780–791. doi: 10.1016/j.cell.2016.04.019.

48. Stoeckius M, Hafemeister C, Stephenson W, Houck-Loomis B, Chattopadhyay PK,
Swerdlow H, Satija R, Smibert P (2017) Simultaneous epitope and transcriptome
measurement in single cells. Nat Methods 14:865–868. doi: 10.1038/nmeth.4380.

49. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM 3rd, Hao Y,
Stoeckius M, Smibert P, Satija R (2019) Comprehensive Integration of Single-Cell Data.
Cell 177:1888-1902.e21. doi: 10.1016/j.cell.2019.05.031.

50. Subelj L, Bajec M (2011) Unfolding communities in large complex networks: combining defensive and offensive label propagation for core extraction. Phys Rev E Stat Nonlin Soft Matter Phys 83:36103. doi: 10.1103/PhysRevE.83.036103.

51. Tapp LD, Shantsila E, Wrigley BJ, Pamukcu B, Lip GYH (2012) The CD14++CD16+ monocyte subset and monocyte-platelet interactions in patients with ST-elevation myocardial infarction. J Thromb Haemost 10:1231–1241. doi: 10.1111/j.1538-7836.2011.04603.x.

52. Tracy RP, Doyle MF, Olson NC, Huber SA, Jenny NS, Sallam R, Psaty BM, Kronmal RA (2013) T-helper type 1 bias in healthy people is associated with cytomegalovirus serology and atherosclerosis: the Multi-Ethnic Study of Atherosclerosis. J Am Heart Assoc 2:e000117. doi: 10.1161/JAHA.113.000117.

53. Uniken Venema WT, Voskuil MD, Vila AV, van der Vries G, Jansen BH, Jabri B, Faber KN, Dijkstra G, Xavier RJ, Wijmenga C, Graham DB, Weersma RK, Festen EA (2019) Single-Cell RNA Sequencing of Blood and Ileal T Cells From Patients With Crohn's Disease Reveals Tissue-Specific Characteristics and Drug Targets. Gastroenterology 156:812-815. doi: 10.1053/j.gastro.2018.10.046.

54. Virani SS, Nambi V, Hoogeveen R, Wasserman BA, Coresh J, Gonzalez F 2nd, Chambless LE, Mosley TH, Boerwinkle E, Ballantyne CM (2011) Relationship between circulating levels of RANTES (regulated on activation, normal T-cell expressed, and secreted) and carotid plaque characteristics: the Atherosclerosis Risk in Communities (ARIC) Carotid MRI Study. Eur Heart J 32:459–468. doi: 10.1093/eurheartj/ehq367.

55. Wildgruber M, Aschenbrenner T, Wendorff H, Czubba M, Glinzer A, Haller B, Schiemann M, Zimmermann A, Berger H, Eckstein H-H, Meier R, Wohlgemuth WA, Libby P, Zernecke A (2016) The "Intermediate" CD14(++)CD16(+) monocyte subset increases in severe peripheral artery disease in humans. Sci Rep 6:39483. doi: 10.1038/srep39483.

56. Williams JW, Winkels H, Durant CP, Zaitsev K, Ghosheh Y, Ley K (2020) Single Cell RNA Sequencing in Atherosclerosis Research. Circ Res 126:1112–1126. doi: 10.1161/CIRCRESAHA.119.315940.

57. Winkels H, Ehinger E, Vassallo M, Buscher K, Dinh HQ, Kobiyama K, Hamers AAJ, Cochain C, Vafadarnejad E, Saliba AE, Zernecke A, Pramod AB, Ghosh AK, Michel NA, Hoppe N, Hilgendorf I, Zirlik A, Hedrick CC, Ley K, Wolf D (2018) Atlas of the immune cell repertoire in mouse atherosclerosis defined by single-cell RNA-sequencing and mass cytometry. Circ Res 122:1675–1688. doi: 10.1161/CIRCRESAHA.117.312513.

58. Wirka RC, Wagh D, Paik DT, Pjanic M, Nguyen T, Miller CL, Kundu R, Nagao M, Coller J, Koyano TK, Fong R, Woo YJ, Liu B, Montgomery SB, Wu JC, Zhu K, Chang R, Alamprese M, Tallquist MD, Kim JB, Quertermous T (2019) Atheroprotective roles of smooth muscle cell phenotypic modulation and the TCF21 disease gene as revealed by single-cell analysis. Nat Med 25:1280–1289. doi: 10.1038/s41591-019-0512-5.

59. Zernecke A, Winkels H, Cochain C, Williams JW, Wolf D, Soehnlein O, Robbins CS, Monaco C, Park I, McNamara CA, Binder CJ, Cybulsky MI, Scipione CA, Hedrick CC, Galkina E V, Kyaw T, Ghosheh Y, Dinh HQ, Ley K (2020) Meta-Analysis of Leukocyte Diversity in Atherosclerotic Mouse Aortas. Circ Res 127:402–426. doi: 10.1161/CIRCRESAHA.120.316903.

60. Zhang L, Li Z, Skrzypczynska KM, Fang Q, Zhang W, O'Brien SA, He Y, Wang L, Zhang Q, Kim A, Gao R, Orf J, Wang T, Sawant D, Kang J, Bhatt D, Lu D, Li C-M,

Rapaport AS, Perez K, Ye Y, Wang S, Hu X, Ren X, Ouyang W, Shen Z, Egen JG, Zhang
Z, Yu X (2020) Single-Cell Analyses Inform Mechanisms of Myeloid-Targeted Therapies
in Colon Cancer. Cell 181:442-459.e29. doi: 10.1016/j.cell.2020.03.048.
61.   Zhang Y, Zheng L, Zhang L, Hu X, Ren X, Zhang Z (2019) Deep single-cell RNA
sequencing data of individual T cells from treatment-naive colorectal cancer patients. Sci
data 6:131. doi: 10.1038/s41597-019-0131-5.
62.   Zheng C, Zheng L, Yoo J-K, Guo H, Zhang Y, Guo X, Kang B, Hu R, Huang JY, Zhang
Q, Liu Z, Dong M, Hu X, Ouyang W, Peng J, Zhang Z (2017) Landscape of Infiltrating T
Cells in Liver Cancer Revealed by Single-Cell Sequencing. Cell 169:1342-1356. doi:
10.1016/j.cell.2017.05.035.
63.   Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, Ziraldo SB, Wheeler
TD, McDermott GP, Zhu J, Gregory MT, Shuga J, Montesclaros L, Underwood JG,
Masquelier DA, Nishimura SY, Schnall-Levin M, Wyatt PW, Hindson CM, Bharadwaj R,
Wong A, Ness KD, Beppu LW, Deeg HJ, McFarland C, Loeb KR, Valente WJ, Ericson
NG, Stevens EA, Radich JP, Mikkelsen TS, Hindson BJ, Bielas JH (2017) Massively
parallel digital transcriptional profiling of single cells. Nat Commun 8:14049. doi:
10.1038/ncomms14049.
64.   Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, Ziraldo SB, Wheeler
TD, McDermott GP, Zhu J, Gregory MT, Shuga J, Montesclaros L, Underwood JG,
Masquelier DA, Nishimura SY, Schnall-Levin M, Wyatt PW, Hindson CM, Bharadwaj R,
Wong A, Ness KD, Beppu LW, Deeg HJ, McFarland C, Loeb KR, Valente WJ, Ericson
NG, Stevens EA, Radich JP, Mikkelsen TS, Hindson BJ, Bielas JH (2017) Massively
parallel digital transcriptional profiling of single cells. Nat Commun 8:1–12. doi:
10.1038/ncomms14049.
830
831

Declarations

Funding:

National Institutes of Health Grant R35-HL-145241, R01-HL-121697, R01-HL-148094

(K.L.)

National Institutes of Health Grant P01-KL-136275 (C.C.H.)

National Institutes of Health Grant R01-HL-134236 (C.C.H.)

National Institutes of Health Grant R01-HL-126543, 5R01-HL-126543-05, 5R01-HL-

140976-02, R01-HL-148094-01, R01-HL-148094 (R.C.K.)

National Institutes of Health Grant K01-HL-137557 (D.B.H.)

National Institutes of Health Grant U01-AI-103408 (I.O.)

National Institutes of Health Grant U01-AI-103390 (A.A.A.)

National Institutes of Health Grant U01-AI-034989 (P.C.T.)

Cancer Research Institute (CRI) (A.S.)

American Heart Association (AHA) 19POST34450020 (L.E.P)

National Institutes of Health Grant NIAID, NICHD, NCI, NIDA, NIMH, NIDCR,

NIAAA, NIDCD, UL1-TR-000004, P30-AI-050409, P30-AI-050410, and P30-AI-

027767.

Formación de Profesorado Universitario (FPU) 16/02780 (R.B.D.)

Swedish Society for Medical Research (SSMF) (J.V.)

Data in this manuscript were collected by the Women's Interagency HIV study, now the

MACS/WIHS Combined Cohort Study (MWCCS).

Author contributions:

Design of the study: J.V., R.S., Y.G., C.P.D., E.E.

Collection of samples and data. A.L.L., R.P.T., J.M.L., W.J.M., K.M.W, A.A.A.,

H.N.H., P.C.T., I.O., S.L.H., and R.C.K.

Analysis of clinical data: D.B.H.

Design and collection of data for the B mode ultrasound sub-study: H.N.H.

scRNA-Seq experiments: C.P.D., and E.E.

Analysis of the data: J.V., R.S., R.G., Y.G., P.R., T.P., L.E.P., C.E.O., R.B.D.,

H.Q.D., A.S., C.A.M., L.L.L., T.W., C.C.H., and K.L.

Bioinformatics analysis: R.G., Y.G. and H.Q.D.

Writing-original draft: J.V., R.S., and K.L.

Competing interests: There are no conflicts of interest.

Data and material availability: All data are available in the main text or the

supplementary materials.

Code availability: All packages used for the analysis on this data are available in R.

Ethics approval: All participants provided informed consent, and each site's

Institutional Review Board approved the studies.

Consent to participate: All participants provided informed consent, and each site's

Institutional Review Board approved the studies.

Consent for publication: All the listed authors have reviewed the manuscript and agreed

with its submission.

876
877

Figures

Figure 1. Gating scheme (A), biaxial dot plots (B-E) to identify major known cell

types, and (F) antibody- based UMAP clustering of major cell types.

PBMCs from 32 WIHS participants were hash-tagged and stained with 40

oligonucleotide-tagged mAbs (**Table S3**). (**B**) B cells were defined as CD19+CD3- and T

cells as CD19-CD3+. (**C**) T cells were identified as CD4 (CD4+CD8-) or CD8 (CD4-

CD8+). (**D**) All CD19-CD3- cells were gated for CD14 and CD16, with CD14+CD16-

cells being classical (CM) and CD14+CD16+ being intermediate (INT) monocytes. (**E**)

The CD14-CD16+ cells from panel D contain NK cells, which were identified by CD56

and defined as CD56+CD14-CD20-CD123-CD206-. Most of the remaining CD56-

CD16+ cells were nonclassical monocytes (NCM). (**F**) The major known cell types were

UMAP-Louvain-clustered by CD3, CD19, CD14, CD16, and CD56 surface expression

(central panel). Then, each major known cell type was UMAP-Louvain-clustered by all

non-negative surface markers. CD4 T cells formed 16 clusters, cluster numbers indicated;

CD8 T cells formed 14 clusters; Classical monocytes (CM) formed 8, Intermediate

monocytes (INT) 5, and Nonclassical monocytes (NCM) 3 clusters. B cells and NK cells

formed 6 clusters each.

Figure 2. Heatmaps of antibody expression (log$_2$ scale) in each main cell type.

A. CD4 T cells

B. CD8 T cells

C. Monocytes

D. B cells

E. NK cells

910   (**A**) CD4 T cell, (**B**) CD8 T cell, (**C**) Monocytes, (**D**) B cells, and (**E**) NK cells.

911   Immunophenotypes at the bottom. EM, effector memory; EMRA, terminally

912   differentiated effector memory; CM, Classical Monocyte; INT, Intermediate Monocyte;

913   NCM, Nonclassical Monocyte.

Figure 3. Cell proportions in women with HIV, CVD, both or neither.

HIV-CVD- (green), HIV+CVD- (yellow), HIV+CVD+ (blue), and HIV+CVD+ on CRT

(red), from left to right. 8 samples per group except 7 for HIV+CVD+. Proportions of

cells in each cluster calculated as percentage of the parent cell type as indicated in the

title of each panel. Clusters with significant differences (*, $p < 0.05$, **, $p < 0.01$, ***

$p < 0.001$) in cell proportions (by log odds ratio) are shown with individual data points,

means and standard error of the mean (SEM). Violin plots below show expression of all

40 cell surface markers ($\log_2$ scale). INT, intermediate monocytes; CRT, cholesterol-

reducing treatment.

Figure 4. Significantly differentially expressed genes of cells in each cluster.

Expression of 485 transcripts was determined by targeted amplification (BD Rhapsody system). Significant genes defined as adjusted $p<0.05$ and $\log_2$ fold change >0. Dot plot: fraction of cells in cluster expressing each gene shown by size of circle and level of expression shown from white (=0) to dark blue (=max, $\log_2$ scale). Red bars indicate genes that were significantly higher in one cluster compared to all other clusters of the parent cell type. There were no DEGs in NK cell clusters. (**A**) CD4 T cells, (**B**) CD8 T cells and (**C**) monocytes. CM, Classical monocytes; INT, Intermediate monocytes; NCM, Nonclassical monocytes; EM, effector memory; EMRA, terminally differentiated effector memory.

Figure 5. Cell proportions of classical monocyte subsets.

Cell proportions of classical monocytes (CMs, CM1-8) from the present dataset in 92

samples of classical monocyte transcriptomes in women with HIV, CVD, both or

neither.[7] CM, classical monocyte; CRT, cholesterol-reducing treatment.

Figure 6. Volcano plots comparing gene expression in single cells from WIHS

participant types in each cluster.

Gene expression focused on HIV+CVD- vs HIV+CVD+, and HIV+CVD+ vs

HIV+CVD+ with CRT. All clusters in which at least 10 genes were statistically

significant are shown. Colored dots (HIV+CVD- yellow, HIV+CVD+ blue, and

HIV+CVD+ with CRT red) indicate significantly differentiated expressed genes

(FDR<0.05 and |log$_2$FC|>2). 3 CD4 T and 7 CD8 T cell clusters, 5 CM and 1 each INT

and B cell clusters met these criteria. Dashed line indicates adjusted p-value of 0.05. CM,

Classical monocytes; INT, Intermediate monocytes.

954