1   **Title: Distinct place cell dynamics in CA1 and CA3 encode experience in new contexts**

2   **Authors:** Can Dong[1] and Mark E. J. Sheffield[1*]

3   **Affiliations:**

4   [1] Department of Neurobiology and Grossman Institute for Neuroscience, Quantitative Biology

5   and Human Behavior, University of Chicago, Chicago, IL 60637, USA.

6   [*] Correspondence to: sheffield@uchicago.edu.

Abstract: We compared trial-by-trial dynamics of place cells in CA1 and CA3 in new contexts across days. We found that CA1 place fields form early but shift backwards with experience and partially remap across days. In contrast, CA3 place fields develop gradually but remain stable with experience and across days. This suggests distinct plasticity mechanisms drive the formation and dynamics of place fields in CA1 and CA3 to encode distinct features of experience.

Main text: The hippocampus plays a critical role in learning by rapidly forming and continuously updating experience-driven representations in the brain[1,2]. Representations of contextual experience are encoded and recalled by populations of place cells in hippocampal sub-regions CA1, CA3, and dentate gyrus[3,4]. Synaptic plasticity within these sub-regions is the key process that alters and stores representations and comes in many forms[5,6]. Understanding how place cell representations form, evolve, and are recalled within distinct hippocampal sub-regions during novel experiences can therefore provide key insights into the types of synaptic plasticity mechanisms that are at play during learning and memory recall. Further, because synaptic plasticity mechanisms can work on rapid timescales to continually alter place cell activity, it is essential to determine the real-time place cell dynamics on a trial-by-trial basis. However, the trial-by-trial dynamics of hippocampal place cells during the first moments of a novel experience and throughout ongoing experience measured over long timescales (across days) has not been systematically compared between CA1 and CA3.

To address this, we expressed GCaMP6f in either the CA1 or CA3 sub-region of different mice (Fig. 1c, d). The Grik4-cre line[7] was used to restrict expression to CA3 pyramidal neurons (Fig.

1c). Using 2-photon microscopy we then recorded calcium transients from pyramidal cell

populations in both regions (Extended Data Fig. 1)[8,9]. On experimental day 1 mice were exposed

to a familiar (F) context before being switched to a novel context (N1) (Fig. 1a)[9]. On

experimental day 2, mice experienced the same F-to-N switch but to a different N context (N2;

Fig. 1a). N1 and N2 were grouped together and are referred to as N. Mice momentarily slowed

down after the transition between contexts (Fig. 1b), confirming their perception of the switch.

This paradigm led to many repeated traversals in both contexts with matched behavior,

allowing lap-by-lap place field (PF) dynamics to be measured systematically and compared

across F and N contexts without confounds caused by changes in behavior.

Upon exposure to N, both CA1 and CA3 place cells globally remapped (Fig. 1e, f) and displayed

altered PFs compared to F (Extended Data Fig. 2). Crucially, we wanted to observe the real-time

formation of new place cells in CA1 and CA3 to examine potential differences in their formation

dynamics. Therefore, formation of new PFs in N was quantified on a lap-by-lap basis (Fig. 1g-i).

Some PFs formed instantly, i.e. on the first lap (Instant PFs; Fig. 1g; left), while others were

delayed by several laps (Delayed PFs; Fig. 1g; right). Similar to previous observations, many CA1

PFs formed rapidly[6,9], with a high proportion of instant PFs (30%; Fig. 1h, i). Unexpectedly, only

9% of CA3 PFs were instant and the distribution of PF onset laps was more uniform, indicating

CA3 place cell representations form more gradually than CA1 representations (Fig. 1h, i;

Extended Data Fig. 2c, d). Supporting this, CA1 place cell activity decoded position on the first

lap better than CA3 place cell activity (Fig. 1j-l, Extended Data Fig. 3). These data suggest that in

a novel context, CA1 instantly forms a well-organized map, whereas CA3 forms a map gradually

with experience.

What do these observations suggest about synaptic plasticity mechanisms? First, some models

of CA1 place field formation require the presence of spatially tuned presynaptic inputs from

CA3[5,10]. In this scheme, spatially tuned CA3 inputs coincident with inputs from entorhinal cortex

(EC) generate dendritic plateau potentials in CA1 cells that drive synaptic potentiation[11]. The

lack of spatially tuned CA3 cells that we found on lap 1 suggests CA1 instant place fields are not

formed through this mechanism, and may instead form through other direct inputs from EC or

Nucleus Reunions [12]. Second, delayed CA1 PFs have been proposed to form through a process

of local dendritic spikes that occur on the initial laps in the absence of somatic firing. This

potentiates local clusters of synapses that become strong enough to drive firing, and possibly

plateau potentials, after a delay, to form new PFs[6,9-11,13,14]. Because we found the vast majority

of CA3 PFs form after a delay, our data suggest a similar mechanism is at play in CA3.

To examine plasticity mechanisms occurring after PFs have formed in CA1 and CA3, we tracked

new PFs throughout experience in N. We compared the first and second half of N and found

CA3 PFs were more stable than CA1 PFs (Fig. 2a). Next, we computed each PF's center of mass

(COM) on a lap-by-lap basis in N in both regions. Some PFs were stable throughout N, whereas

others shifted systematically throughout the session (Fig. 2b). Averaging across the PF

population, PFs shifted backwards in both regions, but significantly more so in CA1 than CA3

(Fig. 2c, Extended Data Fig. 4).

Asymmetric synaptic plasticity mechanisms, such as spike-timing-dependent plasticity (STDP)[15]

and behavioral time-scale plasticity (BTSP)[5,10], predict such backward shifting[15]. These two

forms of plasticity occur over different timescales (milliseconds for STDP, seconds for BTSP), so

we calculated the timescale of backward shifting CA1 PFs to determine which form of plasticity

may be occurring. Based on the ability of STDP to potentiate synapses activated up to ~20 ms

prior to postsynaptic firing[16], we calculated PFs would shift maximally 0.48 ± 0.15 cm/lap (mean

± SD) through STDP based on the average running speed of our mice (24.1 ± 7.5 cm/s mean ±

SD). On average, backward shifting PFs in CA1 shifted by 0.34 ± 0.34 cm/lap (mean ± SD). This

provides strong support that STDP is the mechanism driving continuous backward shifting.

Further, STDP is known to occur at CA1-CA3 synapses[16] and given the relative stability of CA3

PFs concurrent with the CA1 backward shifting PFs we observed here, it suggests STDP is

occurring at these synapses to drive the backward shifting of CA1 PFs (although other synapses

could also be involved).

Memory recall of spatial contexts is supported by the reactivation of stable PFs upon re-

exposures[4]. We therefore examined the same place cells upon re-exposure to N across days

(Fig. 3a). On day 2, mouse behavior on the first lap of N revealed mice had become more

familiar with N (Extended Data Fig. 5a, b). We then quantified the spatial correlation of mean

PFs identified on N day 1 with N day 2 (Fig. 3b-e), which was on average significantly higher in

CA3 than CA1 (Fig. 3c-e). The bimodal distribution of CA3 PF spatial correlations (Fig. 3c;

bottom), helped categorize PFs as either stable (spatial correlation > 0.5, green) or unstable (<

0.5; light-green). In all mice we found that CA3 had a higher fraction of stable PFs compared to

CA1 in (Fig. 3d).

We then compared the PF onset laps of stable PFs on day 2 (Fig. 3f, left) with PFs that newly

formed on day 2, which included both unstable PFs from day 1 (Fig. 3f, middle) and PFs that

appeared for the first time on day 2 (Fig. 3f, right). We found that stable PFs appeared earlier in

the session than newly formed PFs, and this difference was much more significant in CA3 than

CA1 (Fig. 3g). This also shows that a high proportion of CA1 PFs are continuously forming even

as the context becomes familiar. These findings reveal that place cell representations that form

following exposure to novel contexts are more stable in CA3 than CA1. Stable PFs are instantly

reactivated on the second day of exposure and the CA1 continuously forms a high proportion of

new PFs. The greater reorganization of CA1 place cell representations across exposures

suggests offline plasticity in the CA1, possibly occurring during sleep[17].

Is the reorganization of CA1 PFs across days due to a continuation of the backward shifting that

occurs within the session on day 1, possibly during offline activation, or a process more akin to

partial/global remapping? To answer this, we calculated the difference between each PF's COM

on the last active lap on day 1 and first active lap in day 2 (Fig. 3h). The mean of the

distributions in both CA1 and CA3 were not statistically different from 0 (Fig. 3h), revealing no

evidence for systematic backward shifting occurring offline. This shows that backward shifting is

an experience-dependent process and the reorganization of CA1 PFs across days is a process

akin to partial/global remapping.

Lastly, we determined whether the backward shifting we had observed on day 1 in N continued

upon re-exposure to N on day 2. We found that lap-by-lap the CA1 systemically shifted

backwards throughout the first part of the session, yet the CA3 map remained relatively stable

throughout the session (Fig. 3i). This was also true in F (Extended Data Fig. 6), although the

extent of CA1 backward shifting decreased with familiarity. This shows that asymmetric CA1

plasticity mechanisms are occurring in novel and familiar contexts, but the level of novelty

enhances this process.

What might be the function of these differences in the formation and plasticity of place cell

representations in CA1 and CA3? First, instant CA1 PFs could mediate the ability of the

hippocampus to rapidly learn and store episodes of events on a single trial[18]. The backward

shifting of the PF population we observed during ongoing experience could allow the CA1 to

gradually predict future locations within a context[5,19]. Indeed, asymmetric plasticity rules

automatically produce predictions of the future by strengthening synapses that are activated

earlier along the track with each traversal[15]. The CA1, then, rapidly generates unique

representations of the world that are then continuously shaped by exploratory experience to

predict the near future (where am I going?). In parallel, the CA3 gradually forms

representations that are preserved during experience and encode the present moment (where

am I currently?). This function seemingly extends across time as relatively stable CA3 place cell

representations are rapidly reinstated upon re-exposure to the same context, possibly to

support memory recall. Offline reorganization of CA1 place cell representations across days

may instead serve to separate events occurring in the same context into distinct memories[20].

This framework is depicted in a conceptual model in Extended Data Fig. 7. Given the known

circuitry between CA3 and CA1, the independent nature of CA1 encoding relative to CA3

(instant PF formation and reorganization across days) suggests the formation and recall of CA1

place cell representations is substantially influenced by other inputs (Entorhinal cortex and

Nucleus Reunions[12]).

Acknowledgments: We thank D. Dombeck and A. Madar for comments on the manuscript, C.

Cherian for helpful manuscript edits, and members of Sheffield lab for manuscript comments

and useful discussions. **Funding:** This work was supported by: The Whitehall Foundation, The

Searle Scholars Program, The Sloan Foundation, The University of Chicago Grossman Institute

for Neuroscience start-up funds, and the NIH (1DP2NS111657-01). **Author contributions:** C.D.

performed surgeries, collected the data, wrote the analysis code and analyzed data. M.S. and

C.D. conceived and designed the experiments and interpreted the data. M.S. wrote the

manuscript with feedback from C.D. **Competing interests:** Authors declare no competing

interests and **Data and materials availability:** Scripts used for data analysis are available on

Github and raw data is available upon request.

References:

1   Parisi, G. I., Kemker, R., Part, J. L., Kanan, C. & Wermter, S. Continual lifelong learning

with neural networks: A review. *Neural Netw* **113**, 54-71,

doi:10.1016/j.neunet.2019.01.012 (2019).

2   Kitamura, T. *et al.* Engrams and circuits crucial for systems consolidation of a memory.

Science **356**, 73-78, doi:10.1126/science.aam6808 (2017).

3   Hainmueller, T. & Bartos, M. Parallel emergence of stable and dynamic memory

engrams in the hippocampus. *Nature* **558**, 292-296, doi:10.1038/s41586-018-0191-2

(2018).

4   Dupret, D., O'Neill, J., Pleydell-Bouverie, B. & Csicsvari, J. The reorganization and

reactivation of hippocampal maps predict spatial memory performance. *Nat Neurosci*

13, 995-1002, doi:10.1038/nn.2599 (2010).

5   Magee, J. C. & Grienberger, C. Synaptic Plasticity Forms and Functions. *Annu Rev*

Neurosci, doi:10.1146/annurev-neuro-090919-022842 (2020).

6   Sheffield, M. E. & Dombeck, D. A. Dendritic mechanisms of hippocampal place field

formation. *Curr Opin Neurobiol* **54**, 1-11, doi:10.1016/j.conb.2018.07.004 (2019).

7   Nakazawa, K. *et al.* Requirement for hippocampal CA3 NMDA receptors in associative

memory recall. *Science* **297**, 211-218, doi:10.1126/science.1071795 (2002).

8   Sheffield, M. E. & Dombeck, D. A. Calcium transient prevalence across the dendritic

arbour predicts place field properties. *Nature* **517**, 200-204, doi:10.1038/nature13871

(2015).

9   Sheffield, M. E. J., Adoff, M. D. & Dombeck, D. A. Increased Prevalence of Calcium

Transients across the Dendritic Arbor during Place Field Formation. *Neuron* **96**, 490-504

e495, doi:10.1016/j.neuron.2017.09.029 (2017).

10   Bittner, K. C., Milstein, A. D., Grienberger, C., Romani, S. & Magee, J. C. Behavioral time

scale synaptic plasticity underlies CA1 place fields. *Science* **357**, 1033-1036,

doi:10.1126/science.aan3846 (2017).

11   Bittner, K. C. *et al.* Conjunctive input processing drives feature selectivity in hippocampal

CA1 neurons. *Nat Neurosci* **18**, 1133-1142, doi:10.1038/nn.4062 (2015).

12   Dolleman-van der Weel, M. J. *et al.* The nucleus reuniens of the thalamus sits at the

nexus of a hippocampus and medial prefrontal cortex circuit enabling memory and

behavior. *Learn Mem* **26**, 191-205, doi:10.1101/lm.048389.118 (2019).

13 Cohen, J. D., Bolstad, M. & Lee, A. K. Experience-dependent shaping of hippocampal CA1

intracellular activity in novel and familiar environments. *Elife* **6**, doi:10.7554/eLife.23040

(2017).

14 Diamantaki, M. *et al.* Manipulating Hippocampal Place Cell Activity by Single-Cell

Stimulation in Freely Moving Mice. *Cell Rep* **23**, 32-38, doi:10.1016/j.celrep.2018.03.031

(2018).

15 Mehta, M. R., Quirk, M. C. & Wilson, M. A. Experience-dependent asymmetric shape of

hippocampal receptive fields. *Neuron* **25**, 707-715, doi:10.1016/s0896-6273(00)81072-7

(2000).

16 Dan, Y. & Poo, M. M. Spike timing-dependent plasticity of neural circuits. *Neuron* **44**, 23-

30, doi:10.1016/j.neuron.2004.09.007 (2004).

17 Sadowski, J. H., Jones, M. W. & Mellor, J. R. Sharp-Wave Ripples Orchestrate the

Induction of Synaptic Plasticity during Reactivation of Place Cell Firing Patterns in the

Hippocampus. *Cell Rep* **14**, 1916-1929, doi:10.1016/j.celrep.2016.01.061 (2016).

18 Lee, S. W., O'Doherty, J. P. & Shimojo, S. Neural computations mediating one-shot

learning in the human brain. *PLoS Biol* **13**, e1002137, doi:10.1371/journal.pbio.1002137

(2015).

19 Stachenfeld, K. L., Botvinick, M. M. & Gershman, S. J. The hippocampus as a predictive

map. *Nat Neurosci* **20**, 1643-1653, doi:10.1038/nn.4650 (2017).

20 Clewett, D., DuBrow, S. & Davachi, L. Transcending time in the brain: How event

memories are constructed from experience. *Hippocampus* **29**, 162-183,

doi:10.1002/hipo.23074 (2019).

21   Dombeck, D. A., Harvey, C. D., Tian, L., Looger, L. L. & Tank, D. W. Functional imaging of

hippocampal place cells at cellular resolution during virtual navigation. *Nat Neurosci* **13**,

1433-1440, doi:10.1038/nn.2648 (2010).

22   Heys, J. G., Rangarajan, K. V. & Dombeck, D. A. The functional micro-organization of grid

cells revealed by cellular-resolution imaging. *Neuron* **84**, 1079-1090,

doi:10.1016/j.neuron.2014.10.048 (2014).

23   Aronov, D., Nevers, R. & Tank, D. W. Mapping of a non-spatial dimension by the

hippocampal-entorhinal circuit. *Nature* **543**, 719-722, doi:10.1038/nature21692 (2017).

24   Mukamel, E. A., Nimmerjahn, A. & Schnitzer, M. J. Automated analysis of cellular signals

from large-scale calcium imaging data. *Neuron* **63**, 747-760,

doi:10.1016/j.neuron.2009.08.009 (2009).

25   Tampuu, A., Matiisen, T., Olafsdottir, H. F., Barry, C. & Vicente, R. Efficient neural

decoding of self-location with a deep recurrent network. *PLoS Comput Biol* **15**,

e1006822, doi:10.1371/journal.pcbi.1006822 (2019).

26   Ho, J., Tumkaya, T., Aryal, S., Choi, H. & Claridge-Chang, A. Moving beyond P values: data

analysis with estimation graphics. *Nat Methods* **16**, 565-566, doi:10.1038/s41592-019-

0470-3 (2019).

27   Dragoi, G. & Tonegawa, S. Preplay of future place cell sequences by hippocampal

cellular assemblies. *Nature* **469**, 397-401, doi:10.1038/nature09633 (2011).

Fig. 1: Place field formation in novel contexts is rapid in CA1 but gradual in CA3

a, Top left, depiction of the virtual reality (VR) set up. Top right, the familiar (F) and two novel

(N1 and N2) contexts. Bottom: Scheme of the experimental procedure. **b,** Top: Single mouse

behavior showing track position versus time during an F to N switch. Bottom: Summary data

across all mice of mean lap velocity during F and N (n = 20 sessions in 11 mice). Lap velocity is

normalized to the mean lap velocity in each mouse ± SEM, and each lap is compared to the first

lap in F using a one-way ANOVA with Tukey HSD post hoc test. ***, $P < 0.001$. **c,** Brain slice

showing specific CA3 expression of cre-dependent GCaMP6f (green) and DAPI for nuclei (blue)

from a Girk4-cre transgenic mouse. **d,** Example field of views (FOV) from multiplane imaging in

CA1 (top) and CA3 (bottom). Right: scheme of the position for the objective during imaging.

Scale bar = 50 μm. **e,** Left: mean place fields in F sorted by track position. Middle: mean activity

of the same neurons on the left in N. Right: mean place fields in N sorted by track position. Far

right: Percentage of place cells in F and N or common to both from all active neurons. CA1, top

panels; CA3, bottom panels. ΔF/F activity is normalized to each neurons' maximum transient. *n*

= 7 sessions in 4 mice for CA1, *n* = 13 sessions in 7 mice for CA3. **f,** Pearson's correlation

coefficient of each cell's mean place field within F (blue) and between F and N (red). Estimation

plots of mean difference (Δ) shown on the right of each plot (see Methods). **g,** Example of an

instant place field (left) and delayed place field (right) in N. Arrows indicate the place field onset

lap. **h,** Histograms of place field onset laps in F (left) and N (middle) and cumulative fraction

plots for F and N (right). Wilcoxon rank sum test. ***, $P < 0.001$. **i,** Cumulative fraction plots for

CA1 and CA3 place field onset lap in F (left) and in N (right). Wilcoxon rank sum test. ***, $P <$

0.001. **j,** Example mouse showing actual track position (black) on laps 37-40 and the predicted

position (red) decoded by an LSTM decoder (see Methods). CA1, top; CA3, bottom. **k,**

Confusion matrix between the predicted (x-axis) and the real (y-axis) position for the first lap in

N. **l,** Predicted error on the first lap in N for CA1 and CA3. Each dot represents the decoding

error for one decoder trial built based on the activity of 200 randomly chosen place cells from

CA1 (left) or CA3 (right). n = 20 decoder trials. Estimation plot of mean difference (Δ) shown on

the right.

Fig. 2: CA3 place fields exhibit relative stability while CA1 place fields continuously shift

backwards with experience in novel contexts

a, Spatial correlation (Pearson's correlation coefficient) of mean place field activity within N

(first half versus second half of session in N). Estimation plot of mean difference (Δ) between

CA1 and CA3 shown on the right. **b,** 6 example place cells with stable place fields (top) or

backward shifting place fields (bottom) in N. Place field transients shown lap-by-lap for the first

25 laps. White arrows depict the direction of place fields with experience. **c,** Mean ± SEM place

field shift from all place cells and all mice lap-by-lap in N in CA1 (green; n = 1444 cells from 4

mice;) and CA3 (orange; n = 520 cells from 7 mice;). For each place cell the COM of place field

transients on each lap was compared to the COM of place field transients on lap 12. Regression

lines are depicted on top of data and have statistically different slopes. ***, $P < 0.001$.

Fig. 3: CA3 place fields exhibit higher stability across days than CA1 place fields

a, Experimental setup. Top: mice are recorded for 20 min in N across 2 days. Bottom: example

field of view from one imaging plane from CA1 and CA3 across 2 days showing the same cells.

scale bar = 50 µm. **b,** Left: mean place fields in N on day 1 sorted by track position. Middle:

mean activity of the same neurons on the left on day 2 in N. Right: mean place fields in N on

day 2 sorted by track position. **c,** Histograms of Pearson's correlation coefficient of mean place

field activity across days in N (only place cells with place fields on both days were used). Spatial

correlations greater than 0.5 were considered stable place fields and less than 0.5 unstable. **d,**

Fraction of stable place fields across days per mouse in CA1 and CA3. Unpaired t-test. *P<0.05.

269 **e,** Cumulative fraction plots of Pearson's correlation coefficient in N across days in CA1 and CA3

270 from the same data shown in (**c**). Wilcoxon rank sum test. ***, $P < 0.001$. **f,** Example of a stable

271 place field (left), an unstable place field with a newly formed place field on day 2 (middle) and a

272 newly formed place field on day 2 (right). Place field transients shown lap by lap for all laps in N

273 on day 1 and day 2 from the same 3 cells. **g,** Histograms of place field onset laps in N on day 2

274 for stable place fields (left), and newly formed place fields (middle) in CA1 (top) and CA3

275 (bottom). Right: cumulative fraction plots of histogram data. Wilcoxon rank sum test. ***, $P <$

276 0.001. **h**, Histogram of place field shifts across days for CA1 (left) and CA3 (right). The shifts

277 were calculated by the difference between each place field's COM of the last active lap on day 1

278 and first active lap in day 2. Only place cells with place fields on both days are included.

279 Wilcoxon signed rank test, N.S., $P > 0.05$. **i,** Mean ± SEM place field shift from all place cells and

280 all mice lap-by-lap in N on day 2 in CA1 (green; $n = 530$ cells from 3 mice) and CA3 (orange; $n =$

281 212 cells from 6 mice). For each place cell the COM of place field transients on each lap was

282 compared to the COM of place field transients on lap 12. Regression lines are depicted on top

283 of data. ***, $P < 0.001$.