Title: Distinct place cell dynamics in CA1 and CA3 encode experience in new environments

Authors: Can Dong[1], Antoine D. Madar[1], Mark E. J. Sheffield[1*]

Affiliations:

[1] Department of Neurobiology and Institute for Neuroscience, University of Chicago, Chicago, IL

60637, USA.

[*] Correspondence to: sheffield@uchicago.edu.

Abstract: When exploring new environments animals form spatial memories that are updated

with experience and retrieved upon re-exposure to the same environment. The hippocampus is

thought to support these memory processes, but how this is achieved by different subnetworks

such as CA1 and CA3 remains unclear. To understand how hippocampal spatial representations

emerge and evolve during familiarization, we performed 2-photon calcium imaging in mice

running in new virtual environments and compared the trial-to-trial dynamics of place cells in

CA1 and CA3 over days. We find that place fields in CA1 emerge rapidly but tend to shift

backwards from trial-to-trial and remap upon re-exposure to the environment a day later. In

contrast, place fields in CA3 emerge gradually but show more stable trial-to-trial and day-to-day

dynamics. These results reflect different roles in CA1 and CA3 in spatial memory processing

during familiarization to new environments and constrain the potential mechanisms that

support them.

Introduction:

The hippocampus plays a critical role in episodic memory by rapidly forming, updating, and retrieving patterns of activity that represent specific memories[1,2]. The CA1 subnetwork is considered the main output of the hippocampus that transmits information to the cortex and other regions[3], but the computational role of CA1 in memory processing remains unclear. Theoretical studies generally highlight networks upstream of CA1, such as CA3 and the medial entorhinal cortex (MEC) as being attractor networks that encode and retrieve representations associated with different environments[3-7]. These representations could then simply be inherited by CA1[8-10]. However, given the many forms of synaptic plasticity at CA1 synapses[11,12], the complex dendritic computations performed by CA1 neurons[13-15] and the diversity of CA1 interneurons[16,17], CA1 activity dynamics are unlikely to be purely inherited from upstream regions. Establishing the difference in activity dynamics between CA1 and its inputs will help reveal how information is processed by CA1, which computations are specific to CA1, and what the role of its inputs are.

Spatial memories are thought to be encoded and retrieved in the hippocampus through the activity of place cells[18-22] – cells with spatially selective firing fields called place fields (PFs). All hippocampal subnetworks (CA1, CA2, CA3, and dentate gyrus) express PFs during navigation[10,18]. Importantly, CA1 pyramidal neurons develop PFs rapidly during exploration of a novel environment[23]. Understanding how hippocampal representations emerge so rapidly during one-shot events is critical to refine theories of memory. However, the specific dynamics and underlying mechanisms of PF emergence in CA1 have only recently come into focus, and

even less is known about PF emergence in CA3, the main input region to CA1[11,14,15,24-27]. An

important step forward came from analyzing PFs on a trial-to-trial basis during the very first

moments in a novel environment[15]. Such trial-to-trial resolution showed that many neurons in

CA1 develop a PF on the first trial and engage synaptic plasticity mechanisms in the form of

increased dendritic branch spike prevalence[14,15]. Determining the dynamics of PF emergence in

CA3 under the same conditions in novel environments will generate new insights into the

mechanisms of PF emergence in CA1 and reveal the extent to which CA1 PF emergence is

inherited from CA3 PFs.

In addition to how spatial representations emerge, understanding how they evolve during

familiarization to a novel environment (a form of spatial learning) is also critical to refine

theories of learning and memory. Little is known about trial-to-trial PF dynamics in CA1 or CA3

with most studies focusing on mean PF dynamics across conditions[18,28,29]. In familiar

environments it has been shown that PFs in both CA1 and CA3 tend to shift backwards with

experience and develop negatively skewed PFs to varying degrees depending on the behavioral

paradigm used[29-32]. This is thought to be an experience-dependent process that requires NMDA

receptor-dependent long-term synaptic plasticity[33]. How soon these phenomena appear during

the familiarization process, and whether they are inherited by CA1 from CA3, remains unclear[31].

Tracking large numbers of CA1 and CA3 neurons under the same conditions during

familiarization to novel environments will help better understand these phenomena.

Memory retrieval is thought to be achieved by reinstating the same neuronal activity that

occurred during learning[10,34,35]. However, evidence from recordings of large ensembles of place

cells have shown that many PFs in CA1 are unstable across exposures to the same familiar

environment[13,18,28,36-39]. Again, as with PF emergence and trial-to-trial dynamics, a lot less is

known about the stability of CA1 and CA3 PFs during re-exposure to a novel environment. This

has left unclear how familiarization influences PF stability. CA1 PFs might be expected to be less

stable across days than CA3 PFs as CA1 has been shown to integrate stable spatial information

from CA3 with time signals from CA2 upon re-exposures to the same familiar environment[10].

Further supporting the idea of reactivation of stable PFs in CA3 comes from its hypothesized

role in pattern completion[7], although recent experimental findings tracking PFs in novel

environments across days suggest otherwise[18]. Lastly, trial-to-trial retrieval dynamics of single

PFs during re-exposure to a novel environment are not known because of the technical

difficulty to track the same place cells across days[31]. Overall, how PFs emerge, evolve and

stabilize during familiarization to a novel environment is unclear, both in CA1 and CA3.

In this work we use 2-photon Ca^{2+} imaging to longitudinally record from large populations of

CA1 and CA3 pyramidal neurons in head-fixed mice running unidirectionally on a treadmill to

repeatedly traverse visually enriched virtual linear environments with consistent behavior. We

track neurons from the very first moments in a novel environment and across days to compare

the emergence and ongoing dynamics of PFs. We find that PFs initially emerge much faster in

CA1 than CA3, but CA1 PFs on average continuously shift backwards with experience. CA3 PFs

emerge relatively slowly but are subsequently more reliable than CA1 PFs across trials,

displaying less backward shifting. We also find that PF backward shifting decreases with

familiarization across trials and across days. Upon re-exposure to the novel environment on the

second day, stable PFs in CA3 reactivate rapidly on the first trial, whereas CA1 PFs demonstrate

a higher propensity to remap across days. Our findings demonstrate major differences in the

initial emergence, shifting, and stability of PFs in CA1 and CA3 during familiarization to a novel

environment. The distinct features of CA1 PFs compared to CA3 PFs suggests the CA1 performs

significant computations on its spatial inputs from CA3 to support a distinct role in spatial

memory processing.

Results

Place field emergence in a novel environment in CA1 and CA3

We expressed GCaMP6f in either dorsal CA1 or CA3a (referred to as CA3 from here on) of

different mice (Fig. 1c, d). The Grik4-cre line[40] was used to restrict expression to CA3 pyramidal

neurons (Fig. 1c). Importantly, these mice show CRE expression in nearly 100% of pyramidal

cells in CA3 which means our recordings were not biased to a sub-population of CA3 pyramidal

cells[40]. Using 2-photon microscopy we then recorded calcium transients from pyramidal cell

populations in both regions (Supplementary Fig. 1)[13,15]. On experimental day 1 mice were

exposed to a familiar (F) environment before being switched to a novel environment (N1) (Fig.

1a)[15]. On experimental day 2, mice experienced the same F-to-N switch but to a different N

environment (N2; Fig. 1a). N1 and N2 were grouped together and are referred to as N. Mice

momentarily slowed down after the transition between environments (Fig. 1b), confirming

their perception of the switch to N. Because mice were restricted to running in 1 dimension on

a custom-built treadmill, this paradigm led to many repeated traversals in both environments

with matched behavior, allowing lap-by-lap PF dynamics to be measured systematically and

compared across F and N environments without confounds caused by changes in behavior.

As has previously been reported, both CA1 and CA3 place cells globally remapped upon

exposure to N (Fig. 2a, b) and displayed altered PF properties compared to F (Supplementary

Fig. 2)[4,6,15,18,41-44]. For instance, PF widths were on average larger and PF precision lower in N

than F in both CA1 and CA3. Out-of-field PF firing also increased in N versus F as did PF transient

occupancy. Crucially, we wanted to observe the real-time emergence dynamics of new place

cells in CA1 and CA3 to examine potential differences. Therefore, emergence of new PFs in N

was quantified on a lap-by-lap basis (Fig. 2c-e). Some PFs formed instantly, i.e. on the first lap

(Instant PFs; Fig. 2c; left), while others were delayed by several laps (Delayed PFs; Fig. 2c; right).

Similar to previous observations, many CA1 PFs formed rapidly[14,15,23], with a high proportion of

instant PFs (30%; Fig. 2d, e). Unexpectedly, only 9% of CA3 PFs were instant and the distribution

of PF onset lap number was more uniform, indicating CA3 PFs form more gradually than CA1

PFs (Fig. 2d, e; Supplementary Fig. 2c, d). Supporting this, CA1 place cell activity decoded

position on the first lap better than CA3 place cell activity (Fig. 2f-h, Supplementary Fig. 3).

These data suggest that in a novel environment, CA1 instantly forms a well-organized map,

whereas CA3 forms a map gradually with experience.

Trial-to-trial place field dynamics in a novel environment

To examine and compare how PFs evolve in CA1 and CA3 during spatial learning, we tracked

new PFs throughout familiarization to N. We compared the first and second half of the session

and found CA3 PFs were more stable than CA1 PFs within a session in N (Fig. 3a). Next, we

computed each PF's center of mass (COM) on a lap-to-lap basis (Fig. 3b): in CA1, approximately

half of the PFs significantly shifted during the session, but only a third in CA3 (Fig 3c). The

direction of shift could go backward or forward relative to the direction of the animal's motion

(Fig. 3b-d) but most PFs shifted backwards, with a larger skew in CA1 than CA3 (Fig 3c-d). As a

population, CA1 PFs shifted backwards much faster than CA3 PFs (Fig. 3e). This difference is not

an artifact of a lower sample size of CA3 PFs as shown by downsampling 1000 times the CA1

dataset (Fig. 3e). Removing transients prior to the onset of a robust PF (as we do for defining PF

onset lap, see methods) did not alter our conclusions (Supplementary Fig. 4a-b). Individual PF

shifting dynamics were not related to PF onset lap or velocity (Supplementary Fig. 5 and 6).

Backward and forward shifting PFs occurred at all positions, with large shifts weakly biased

towards the end of the track for CA1 (Supplementary Fig. 7).

Backward shifting PFs have been previously reported and may be accompanied with an increase

in PF width and the development of a negative skew with experience, although reports are

conflicting[29-33,45]. We found that, in a novel environment, PF width tended to increase through

the first 10 laps in CA1 but not in CA3 (Supplementary Fig. 8). PFs also started with a positive

skew that decreased with experience in CA1 but not in CA3 (Supplementary Fig. 8). Note that

under calcium imaging, as compared to electrophysiology, PFs are generally positively skewed

rather than symmetric as an artifact of calcium transient decay times[46]. An evolution towards

less positively skewed and larger PFs suggests that PFs start earlier on the track (see example in

Supplementary Fig. 8a), consistent with some electrophysiological reports[32,45]. However,

changes in skewness and width were not the only cause of COM backward shifting, as the PFs

end position also shifted backwards (Supplementary Fig. 4c-d).

Interestingly, we noticed that in CA1 the population tended to shift forward during the first

laps, with the global backward shift occurring around the 5th lap (Fig. 3e). We found that the

delay in backward shifting was driven by instant PFs that tended to shift forward on early laps

(Supplementary Fig. 9). It is worth restating here that CA3 shows few early-onset PFs on the

initial laps which is when the forward shifting of instant CA1 PFs occurs. This suggests backward

shifting in CA1 might require the presence of established CA3 PFs, which is why instant PFs

initially shift forward.

Place field stability upon re-exposure to a novel environment across days in CA1 and CA3

Memory recall of spatial environments is generally thought to be supported by the reactivation

of stable PFs upon re-exposures[19-21,47]. We therefore examined the same place cells upon re-

exposure to N (specifically N2, see methods) across days (Fig. 4a). On N day 2, increased lap

velocity on the first lap compared to the first lap on N day 1 revealed mice had become more

familiar with N (Fig. 4b). We then quantified the spatial correlation of mean PFs identified on N

day 1 with N day 2, which was on average significantly higher in CA3 (0.70 ± 0.04) than CA1

(0.49 ± 0.02) (Fig. 4c-f). The bimodal distribution of CA3 PF spatial correlations (Fig. 4d; bottom)

helped categorize PFs as either stable (R > 0.5) or unstable (<= 0.5). In all mice we found that

CA3 had a higher fraction of stable PFs compared to CA1 (Fig. 4e). Interestingly, in CA1, we

found a small but significant positive correlation between PF shifting on N day 1 and PF spatial

correlation across days, suggesting that higher day-to-day PF stability is associated with more

stable trial-to-trial PF dynamics upon the first exposure to a novel environment (Supplementary

Fig. 10).

We then compared the PF onset laps of stable PFs on day 2 (re-emergence, Fig. 4g, left) with

PFs that newly formed on day 2, which included both unstable PFs from day 1 (Fig. 4g, middle)

and PFs that appeared for the first time on day 2 (Fig. 4g, right). We found that stable PFs re-

emerged earlier in the session than newly formed PFs, and this difference was much more

apparent in CA3 than CA1 (Fig. 4h, right). Further, a high proportion of CA1 PFs are continuously

forming even as the environment becomes familiar. Lastly, similar to day 1, the vast majority of

new PFs in CA3 emerge gradually after a delay with very few instant PFs (Fig. 4h, bottom-middle

and bottom-right) whereas the CA1 again shows a much higher proportion of instant PFs on day

2 (Fig. 4h, top-middle and top-right). This reveals that stable CA3 PFs are retrieved much faster

than new PFs emerge during re-exposure on day 2, whereas stable and new PFs in CA1 show

similar emergence dynamics.

Lap-by-lap dynamics across days

We first asked if there was any evidence of continuous PF shifting of the population occurring

offline between sessions separated by a day. We found no such evidence as the distribution of

PF shift distance between the end of the session on day 1 and the start of the session on day 2

was not skewed (Fig. 5a). This indicates that COM shifting reflects experience-dependent

processes that occur during ongoing familiarization but do not continue offline. However, when

considering only stable PFs that on day 1 showed statistically significant backward shifting, we

found they tended to reset on day 2 toward their original position on the early laps of day 1

(Fig. 5b, Supplementary Fig. 11). Therefore, although shifting does not continue offline, offline

processes may be involved in resetting PFs back towards their original position.

We then asked whether the PF shifting dynamics we had observed on day 1 in N continued

upon re-exposure on day 2. The few stable PFs that exhibited significant shifting on both days

did not change much on average, but the direction and amplitude of individual PF shifts were

not correlated across days (Fig. 5c). At the population level, which was thus mostly driven by

newly shifting PFs, we found that CA1 PFs on average shifted backwards like on day 1, before

stabilizing after lap 15 (Fig. 5d). The CA3 map shifted less than CA1 (Fig. 5d, right). In contrast to

day 1, we observed a decrease in PF width over the first laps in CA1, but not CA3, and little

change in skewness with population dynamics indistinguishable between CA1 and CA3

(Supplementary Fig. 8).

Formally comparing dynamics on N day 1, N day 2 and very familiar environments (F), we found

that CA1 population backward shifting slowed down and stabilized with familiarization across

days (Fig. 6a, c). Stabilization also tended to occur earlier with the level of familiarity: in

contrast to day 1, a plateau was reached on day 2 after 15 laps, and only after 12 laps in F.

Shifting of the CA3 map showed a similar trend than CA1 although changes with familiarization

across days were less obvious (Fig. 6b, d), in part because shifting dynamics were already slow

on day 1. Resampling the CA3 data revealed extensive overlap in the population shift across

conditions, indicating similar dynamics, (Fig. 6d, right). This analysis also revealed that the

population becomes more homogeneous with familiarization across days (smaller variance in

the slope distribution). Finally, in CA1, PF skewness and width tended to evolve and stabilize

with familiarization, whereas the CA3 population did not show clear dynamics despite

homogenizing across days (Supplementary Fig. 8). These results suggest that lap-wise PF

dynamics are enhanced by novelty and decrease as a function of familiarity in CA1, and these

PF changes are less apparent in CA3.

Discussion

We used 2-photon calcium imaging of large populations of dorsal CA1 and CA3 pyramidal

neurons to measure and compare the emergence, shifting dynamics, and longitudinal stability

of PFs in novel environments with trial-to-trial resolution. We found that PFs emerge faster in

CA1 but place cells are constantly renewed across exposures whereas they emerge later in CA3

with less turnover across exposures. After emergence, the location of the PFs is not always

stable, sometimes showing prominent backward or forward shifting from lap-to-lap. The

average spatial representation in the hippocampus shifts backwards, with a faster shift in CA1

than CA3. This backward shifting slows down with familiarization across days. These findings

support the idea that CA3 and CA1 perform distinct functions during familiarization to a novel

environment, and CA1 does not simply inherit spatial information from CA3 during this form of

spatial learning. They also constrain the potential mechanisms explaining how spatial

representations emerge, evolve, and reactivate in the hippocampus.

Consistent with previous reports, we observed hippocampal PFs that emerged instantly in a

novel environment (on the very first trial) and others that emerged after multiple trials (delayed

onset PFs)[14,15,23]. Interestingly, the proportion of instant PFs was much lower in CA3 than CA1

(Fig 2). Although to our knowledge this is the first time the trial-to-trial emergence of CA3 PFs

has been reported in completely novel environments, a related study did compare the

emergence of PFs in CA3 and CA1 in a familiar linear belt with novel sensory cues added[29]. This

study found 2 types of place cells: 1) landmark vector (LV) cells that have multiple instant PFs

locked to sensory cues and 2) place cells with single PFs with a mix of instant and delayed-onset

PFs that emerge when novel sensory cues are added. Interestingly, they found 10 times more

LV cells in CA1 than CA3. Because PFs in LV cells emerged instantly, the larger proportion of LV

cells in CA1 is consistent with our finding that CA1 has more instant PFs than CA3. We

confirmed that CA1 has a lot more place cells with multiple PFs than CA3 (10-15% in N and F in

CA1 vs 1-2% in CA3), but, in contrast to LV cells, only a fraction of them had instant PFs (31% in

CA1 in N, comparable to the 27% instant PFs of single-PF place cells). There was no difference

between multi- and single-PF place cells in the distribution of onset laps (2-sample KS-test p =

0.4 in N, CA1). This might be due to differences in paradigms. For instance, in contrast to the

global remapping and emergence of PFs distributed across the whole track that we see in new

environments, adding a cue to a familiar environment only induces partial remapping and PF

emergence near cues[29].

Rapid PF emergence in a novel environment in CA1 is thought to rely on a combination of

strong, spatially tuned excitatory inputs that do not need to be potentiated[14] and high neuronal

excitability, either from novelty-induced disinhibition[16] or from intrinsic properties (low firing

threshold, bursting propensity)[26]. CA3 is the main source of excitatory inputs to CA1 and is

known to drive CA1 spatial representations in a majority of neurons, at least in familiar

environments[48]. It was thus surprising to find that instant PFs were much less prevalent in CA3

than CA1 (Fig 2). The simplest explanation is that CA1 instant PFs are not inherited from CA3

inputs during initial exploration. Indeed, not all CA1 place cells are necessarily driven by CA3[48]

as CA1 receives other sources of spatially modulated inputs (entorhinal cortex[4], CA2[10], non-

imaged subareas of CA3[49] , nucleus reuniens[50]), but the emergence dynamics of spatial

representations in these areas are not currently known. Alternatively, CA1 instant PFs could be

driven by the few CA3 neurons with instant PFs if those neurons have a high degree of

divergence to CA1. Low dendritic inhibition in CA1 pyramidal cells upon initial exposure to novel

environments could serve to amplify the low number of CA3 inputs[15,25,51]. Delayed onset CA3

neurons as well as neurons with unstable spatial modulation could also partially contribute to

CA1 instant PFs since some of them are active on early trials.

After PF emergence, PF properties (position, width, shape) evolve with familiarization and their

dynamics have been used as a proxy to study the synaptic plasticity mechanisms supporting

spatial representations in the hippocampus[30,32,45]. Initial reports showed that, in a familiar

environment, the population of CA1 PFs shifts backwards with experience, a phenomenon

consistent with Hebbian rules[32,45] and dependent on NMDA-receptors[33]. Later

electrophysiological studies compared lap-by-lap shifting dynamics in CA1 versus CA3 under

different familiarity levels[29-31]. Despite discrepancies across studies attributable to different

ways of defining novelty, our results are generally consistent with these studies, especially with

Roth et. al. (2012)[31] who used completely new distal and proximal cues. We observed

significant backward shifting of the population of PFs on day 1 of a new virtual environment in

CA1, and to a lesser extent CA3, with the shift slowing down with familiarization across days

(Figs. 3, 5, 6). Consistent with most reports[31,32,45], we found that CA1 was still backward shifting

in very familiar environments, but not CA3 (Fig. 6).

Population backward shifting was initially reported to coincide with increased negative

skewness and enlargement of PFs[32,45]. However, in contrast to backward shifting, skewness and

width dynamics are inconsistent across studies[30,31], suggesting a heterogeneity of mechanisms

leading to backward shifting[30,52]. Our data indicates that population backward shifting in CA1 in

a novel environment is due to a combination of PF expansion, skewness changes and pure

translation of the PF (Supplementary Figs. 4, 8). These dynamics were dependent on familiarity,

with changes in skewness closely matching population backward shifting. We did not detect

clear dynamics in PF width or skewness in CA3, revealing a dissociation in mechanisms

supporting population shifting between CA3 and CA1.

In addition to the population trend, we characterized shifting dynamics in individual PFs,

revealing that half of PFs show significant linear shifts in CA1, mostly backward, compared to

only a third in CA3 (Fig. 3c). PFs with large shifts tended to be less stable across days

(Supplementary Fig. 10), suggesting that the same mechanisms may underly lap-by-lap and day-

to-day stability. On the other hand, PFs that significantly shifted on N day 1 but were stable

across days tended, upon re-exposure on day 2, to reset their location toward their original

position on day 1 (Fig 5b, Supplementary Fig. 11). This reset opposite of the direction of shifting

shows that the plasticity underlying lap-by-lap shifting does not continue offline and is thus

experience-dependent[32]. It also shows that lap-by-lap plasticity does not necessarily cause long-

term changes in PFs. Together, these findings suggest that the effects of weak lap-by-lap

plasticity decay offline (resetting PFs across exposures), whereas strong lap-by-lap plasticity

promotes remapping across sessions.

Interestingly, we found that PFs did not shift backward in CA1 for the first few trials in the novel

environment, they actually shifted forward on average (Supplementary Fig. 9). This fits with the

idea that synapses from CA3 place cells could be necessary to initiate backward shifting in CA1,

as only a very small number of CA3 PFs emerge on initial laps in the novel environment. As

more CA3 PFs emerge those synapses may undergo asymmetric plasticity, such as spike-timing-

dependent plasticity (STDP)[12] and behavioral time-scale plasticity (BTSP)[11,24], which could start

backward shifting in CA1[11,30-33]. Indeed, both STDP and BTSP are known to occur at CA3-CA1

synapses of pyramidal neurons[11,12,24] but other synapses could also be involved.

A classic STDP rule fits well with the extent of backward shifting we observe across laps[32,52] as it

can potentiate synapses activated up to 20 milliseconds prior to postsynaptic firing, which,

based on the average running speed we observed, would maximally shift PFs backwards by ~0.5

cm/lap, approximately what we see in CA1 on N day 1. However, STDP is not always

asymmetric at CA3-CA1 synapses, depending on induction protocols[12]. Moreover, the relevance

of this type of time-dependent plasticity *in vivo* has been called into question as synaptic

changes might just be too weak to yield an effect on single trials[53,54]. On the other hand, BTSP

induces larger synaptic changes[24] and may thus be more likely to produce visible shifting effects

over single trials. However, BTSP requires plateau potentials associated with burst firing, and

those may not happen on most laps[24,55,56], which is likely necessary to see continuous linear

shifts like we observed. BTSP can also lead to potentiation of synapses activated several

seconds prior to burst firing, which could induce larger PF shifting than observed here. In

addition, BTSP has only been found at CA3-CA1 synapses so far and may not occur in CA3

pyramidal neurons even though CA3 shows some backward shifting, albeit less than CA1. Lastly,

there is evidence that cholinergic input from the medial septum could play a role in regulating

backward shifting, at least in CA1[57]. It is therefore not yet clear which mechanisms are driving

backward shifting in CA1 and CA3 place cells. The mechanisms supporting forward shifting in

some neurons is also unknown. Our careful characterization of the shifting phenomenon in

both CA1 and CA3 will constrain future computational models to clarify how hippocampal

physiology supports plasticity of spatial representations.

The reinstatement of the same place code upon re-exposure to the same environment is

generally thought to provide a neural substrate for remembering the environment[19-21,47]. Single

unit recordings supported this idea by showing place cells remain stable over long

periods[58]. However, recent evidence from calcium imaging of large ensembles of place cells has

shown that many place cells in CA1 are unstable across exposures to the same

environment[10,13,18,28,36-39]. Rather than a deterioration in the memory representation of the

environment, such remapping may serve a function and has been proposed to encode distinct

episodes that occur in the same environment separated by time or small changes in

environment[10,37,39,41,59-63]. This remapping might therefore allow the animal to create

independent representations of different episodes associated with the same environment and

favor mnemonic discrimination of those episodes[64,65]. Our data support this idea as the average

PF correlation in CA1 across days in a novel environment was 0.49 ± 0.02 (see methods)

revealing considerable remapping across days (Fig. 4). This value is higher than what is reported

in a recent study using methods similar to ours (~0.2)[18]. This discrepancy could come from

differences in environment complexity which has been shown to influence PF stability and

reliability[18,66], although no differences were observed between moderate and enriched visual

environments, only impoverished environments showed reduced stability[18]. Differences in the

experimental paradigm could also factor in: they switched repeatedly back and forth between

novel and familiar environments within a single recording session, whereas our mice switched

only once, defining only two different episodes (familiar and novel). Increased remapping due

to repetitive switching, which defines several episodes, is consistent with the hypothesis of CA1

tracking distinct episodes.

A constantly changing place code in the hippocampus could nevertheless be problematic as

information about the familiarity of the environment would be lost. CA3 has long been

theorized to have a stable code to support memory recall through pattern completion via

recurrent connections[7]. We show that CA3 place cells are indeed more stable across exposures

to a novel environment than CA1 place cells. To our knowledge the only other study to track

CA3 place cells across days in a *novel* environment did not observe higher stability in CA3 (~0.1

spatial correlation in their study versus ~0.7 reported in Fig 4)[18]. In addition to differences of

task design stated above, this discrepancy could be explained by their inability to distinguish

between CA3 and CA2 cells, as CA2 cells are known to have unstable place coding properties[10].

Another related study did find a similar effect as we report here in familiar environments[39] and

concluded that CA3 provides a highly stable representation of space and context but little

information about time, whereas CA1 is selectively required to integrate CA3 spatial/contextual

and CA2 temporal information over hours and days. We add to this framework by showing that

the reactivation of stable place cells in CA3 occurs rapidly (on the first traversal) upon re-

exposure to the environment. This is consistent with the idea of the CA3 recurrent connectivity

being the support of pattern completion through discrete attractors, CA3 activity rapidly

settling in the correct attractor basin upon re-exposure[5]. Transition dynamics from one

attractor to the next likely depend on specific connectivity and excitability properties:

identifying the characteristics of recurrent neural network models that would fit the fast

transition we observed will be essential to understand the elusive mechanisms of hippocampal

remapping.

Overall, the differences in emergence, shifting, and stability of PFs in CA1 and CA3 suggest

distinct roles and mechanisms at play in these hippocampal subnetworks to support spatial

memories. Instant CA1 PFs could mediate the ability of the hippocampus to rapidly represent

episodes on a single trial, a feature of episodic memory[67]. The backward shifting of the PF

population during ongoing experience could allow CA1 to gradually better predict future

locations within an environment before physically arriving at those locations[11,68,69]. CA1,

therefore, rapidly generates unique representations of the world that are then continuously

updated by exploratory experience to predict the near future (where am I going?). In parallel,

the CA3 gradually forms representations with stable trial-to-trial dynamics, thus encoding

location in the present moment (where am I currently?). This function seemingly extends across

time as relatively stable CA3 spatial representations are rapidly reinstated upon re-exposure to

the same environment, possibly to support memory recall through pattern completion.

Remapping of CA1 spatial representations across days may instead serve to separate events

occurring in the same environment into distinct memory episodes[70]. This framework is depicted

in a conceptual model in Supplementary Fig. 12.

Acknowledgments: We thank D. Dombeck for comments on the manuscript, C. Cherian for

helpful manuscript edits, and members of Sheffield lab for manuscript comments and useful

discussions. **Funding:** This work was supported by: The Whitehall Foundation, The Searle

Scholars Program, The Sloan Foundation, The University of Chicago Grossman Institute for

Neuroscience start-up funds, and the NIH (1DP2NS111657-01). **Author contributions (CRediT):**

Conceptualization: MS, Data curation: CD, Formal Analysis: CD, AM, MS, Funding acquisition:

MS, Investigation: CD, Methodology: MS, CD, AM, Project administration: CD, Software: CD

Supervision: MS, AM, Validation: AM, MS, Visualization: CD, AM, MS, Writing – original draft:

MS, CD, Writing – review & editing: AM, MS, CD.

Competing interests: Authors declare no competing interests

Data and materials availability: Scripts used for data analysis are available on GitHub public

repository (https://github.com/Candong/Distinct_CA1_CA3.git). The raw imaging data are

available from the corresponding author upon reasonable request. Source data for each figure

are provided with this paper.

References:

1   Parisi, G. I., Kemker, R., Part, J. L., Kanan, C. & Wermter, S. Continual lifelong learning
with neural networks: A review. *Neural Netw* **113**, 54-71,
doi:10.1016/j.neunet.2019.01.012 (2019).
2   Kitamura, T. *et al.* Engrams and circuits crucial for systems consolidation of a memory.
Science **356**, 73-78, doi:10.1126/science.aam6808 (2017).
3   Goode, T. D., Tanaka, K. Z., Sahay, A. & McHugh, T. J. An Integrated Index: Engrams,
Place Cells, and Hippocampal Memory. *Neuron* **107**, 805-820,
doi:10.1016/j.neuron.2020.07.011 (2020).
4   Latuske, P., Kornienko, O., Kohler, L. & Allen, K. Hippocampal Remapping and Its
Entorhinal Origin. *Front Behav Neurosci* **11**, 253, doi:10.3389/fnbeh.2017.00253 (2017).
5   Knierim, J. J. & Zhang, K. Attractor dynamics of spatially correlated neural activity in the
limbic system. *Annu Rev Neurosci* **35**, 267-285, doi:10.1146/annurev-neuro-062111-
150351 (2012).
6   Jeffery, K. J. Place cells, grid cells, attractors, and remapping. *Neural Plast* **2011**, 182602,
doi:10.1155/2011/182602 (2011).
7   Kesner, R. P. & Rolls, E. T. A computational theory of hippocampal function, and tests of
the theory: new developments. *Neurosci Biobehav Rev* **48**, 92-147,
doi:10.1016/j.neubiorev.2014.11.009 (2015).
8   Solstad, T., Moser, E. I. & Einevoll, G. T. From grid cells to place cells: a mathematical
model. *Hippocampus* **16**, 1026-1031, doi:10.1002/hipo.20244 (2006).
9   Neher, T., Azizi, A. H. & Cheng, S. From grid cells to place cells with realistic field sizes.
PLoS One **12**, e0181618, doi:10.1371/journal.pone.0181618 (2017).
10   Mankin, E. A., Diehl, G. W., Sparks, F. T., Leutgeb, S. & Leutgeb, J. K. Hippocampal CA2
activity patterns change over time to a larger extent than between spatial contexts.
Neuron **85**, 190-201, doi:10.1016/j.neuron.2014.12.001 (2015).
11   Magee, J. C. & Grienberger, C. Synaptic Plasticity Forms and Functions. *Annu Rev
Neurosci*, doi:10.1146/annurev-neuro-090919-022842 (2020).
12   Buchanan, K. A. & Mellor, J. R. The activity requirements for spike timing-dependent
plasticity in the hippocampus. *Front Synaptic Neurosci* **2**, 11,
doi:10.3389/fnsyn.2010.00011 (2010).
13   Sheffield, M. E. & Dombeck, D. A. Calcium transient prevalence across the dendritic
arbour predicts place field properties. *Nature* **517**, 200-204, doi:10.1038/nature13871
(2015).

14 Sheffield, M. E. & Dombeck, D. A. Dendritic mechanisms of hippocampal place field
formation. *Curr Opin Neurobiol* **54**, 1-11, doi:10.1016/j.conb.2018.07.004 (2019).
15 Sheffield, M. E. J., Adoff, M. D. & Dombeck, D. A. Increased Prevalence of Calcium
Transients across the Dendritic Arbor during Place Field Formation. *Neuron* **96**, 490-504
e495, doi:10.1016/j.neuron.2017.09.029 (2017).
16 Pedrosa, V. & Clopath, C. The interplay between somatic and dendritic inhibition
promotes the emergence and stabilization of place fields. *PLoS Comput Biol* **16**,
e1007955, doi:10.1371/journal.pcbi.1007955 (2020).
17 Freund, T. F. & Buzsaki, G. Interneurons of the hippocampus. *Hippocampus* **6**, 347-470,
doi:10.1002/(SICI)1098-1063(1996)6:4<347::AID-HIPO1>3.0.CO;2-I (1996).
18 Hainmueller, T. & Bartos, M. Parallel emergence of stable and dynamic memory
engrams in the hippocampus. *Nature* **558**, 292-296, doi:10.1038/s41586-018-0191-2
(2018).
19 Dupret, D., O'Neill, J., Pleydell-Bouverie, B. & Csicsvari, J. The reorganization and
reactivation of hippocampal maps predict spatial memory performance. *Nat Neurosci*
13, 995-1002, doi:10.1038/nn.2599 (2010).
20 Kentros, C. Hippocampal place cells: the "where" of episodic memory? *Hippocampus* **16**,
743-754, doi:10.1002/hipo.20199 (2006).
21 Kentros, C. G., Agnihotri, N. T., Streater, S., Hawkins, R. D. & Kandel, E. R. Increased
attention to spatial context increases both place field stability and spatial memory.
Neuron **42**, 283-295, doi:10.1016/s0896-6273(04)00192-8 (2004).
22 Robinson, N. T. M. *et al.* Targeted Activation of Hippocampal Place Cells Drives Memory-
Guided Spatial Behavior. *Cell* **183**, 1586-1599 e1510, doi:10.1016/j.cell.2020.09.061
(2020).
23 Frank, L. M., Stanley, G. B. & Brown, E. N. Hippocampal plasticity across multiple days of
exposure to novel environments. *J Neurosci* **24**, 7681-7689,
doi:10.1523/JNEUROSCI.1958-04.2004 (2004).
24 Bittner, K. C., Milstein, A. D., Grienberger, C., Romani, S. & Magee, J. C. Behavioral time
scale synaptic plasticity underlies CA1 place fields. *Science* **357**, 1033-1036,
doi:10.1126/science.aan3846 (2017).
25 Cohen, J. D., Bolstad, M. & Lee, A. K. Experience-dependent shaping of hippocampal CA1
intracellular activity in novel and familiar environments. *Elife* **6**, doi:10.7554/eLife.23040
(2017).
26 Epsztein, J., Brecht, M. & Lee, A. K. Intracellular determinants of hippocampal CA1 place
and silent cell activity in a novel environment. *Neuron* **70**, 109-120,
doi:10.1016/j.neuron.2011.03.006 (2011).
27 Leutgeb, S., Leutgeb, J. K., Treves, A., Moser, M. B. & Moser, E. I. Distinct ensemble
codes in hippocampal areas CA3 and CA1. *Science* **305**, 1295-1298,
doi:10.1126/science.1100265 (2004).
28 Ziv, Y. *et al.* Long-term dynamics of CA1 hippocampal place codes. *Nat Neurosci* **16**, 264-
266, doi:10.1038/nn.3329 (2013).
29 Geiller, T., Fattahi, M., Choi, J. S. & Royer, S. Place cells are more strongly tied to
landmarks in deep than in superficial CA1. *Nat Commun* **8**, 14531,
doi:10.1038/ncomms14531 (2017).

30 Lee, I., Rao, G. & Knierim, J. J. A double dissociation between hippocampal subfields:
differential time course of CA3 and CA1 place cells for processing changed
environments. *Neuron* **42**, 803-815, doi:10.1016/j.neuron.2004.05.010 (2004).
31 Roth, E. D., Yu, X., Rao, G. & Knierim, J. J. Functional differences in the backward shifts of
CA1 and CA3 place fields in novel and familiar environments. *PLoS One* **7**, e36035,
doi:10.1371/journal.pone.0036035 (2012).
32 Mehta, M. R., Quirk, M. C. & Wilson, M. A. Experience-dependent asymmetric shape of
hippocampal receptive fields. *Neuron* **25**, 707-715, doi:10.1016/s0896-6273(00)81072-7
(2000).
33 Ekstrom, A. D., Meltzer, J., McNaughton, B. L. & Barnes, C. A. NMDA receptor
antagonism blocks experience-dependent expansion of hippocampal "place fields".
Neuron **31**, 631-638, doi:10.1016/s0896-6273(01)00401-9 (2001).
34 Reijmers, L. G., Perkins, B. L., Matsuo, N. & Mayford, M. Localization of a stable neural
correlate of associative memory. *Science* **317**, 1230-1233, doi:10.1126/science.1143839
(2007).
35 Liu, X. *et al.* Optogenetic stimulation of a hippocampal engram activates fear memory
recall. *Nature* **484**, 381-385, doi:10.1038/nature11028 (2012).
36 Kinsky, N. R., Sullivan, D. W., Mau, W., Hasselmo, M. E. & Eichenbaum, H. B.
Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long Timescales.
Curr Biol **28**, 3578-3588 e3576, doi:10.1016/j.cub.2018.09.037 (2018).
37 Rubin, A., Geva, N., Sheintuch, L. & Ziv, Y. Hippocampal ensemble dynamics timestamp
events in long-term memory. *Elife* **4**, doi:10.7554/eLife.12247 (2015).
38 Jeantet, Y. & Cho, Y. H. Evolution of hippocampal spatial representation over time in
mice. *Neurobiol Learn Mem* **98**, 354-360, doi:10.1016/j.nlm.2012.10.004 (2012).
39 Mankin, E. A. *et al.* Neuronal code for extended time in the hippocampus. *Proc Natl*
Acad Sci U S A **109**, 19462-19467, doi:10.1073/pnas.1214107109 (2012).
40 Nakazawa, K. *et al.* Requirement for hippocampal CA3 NMDA receptors in associative
memory recall. *Science* **297**, 211-218, doi:10.1126/science.1071795 (2002).
41 Colgin, L. L., Moser, E. I. & Moser, M. B. Understanding memory through hippocampal
remapping. *Trends Neurosci* **31**, 469-477, doi:10.1016/j.tins.2008.06.008 (2008).
42 Muller, R. U. & Kubie, J. L. The effects of changes in the environment on the spatial firing
of hippocampal complex-spike cells. *J Neurosci* **7**, 1951-1968 (1987).
43 Bostock, E., Muller, R. U. & Kubie, J. L. Experience-dependent modifications of
hippocampal place cell firing. *Hippocampus* **1**, 193-205, doi:10.1002/hipo.450010207
(1991).
44 Fyhn, M., Hafting, T., Treves, A., Moser, M. B. & Moser, E. I. Hippocampal remapping
and grid realignment in entorhinal cortex. *Nature* **446**, 190-194,
doi:10.1038/nature05601 (2007).
45 Mehta, M. R., Barnes, C. A. & McNaughton, B. L. Experience-dependent, asymmetric
expansion of hippocampal place fields. *Proc Natl Acad Sci U S A* **94**, 8918-8921,
doi:10.1073/pnas.94.16.8918 (1997).
46 Dombeck, D. A., Harvey, C. D., Tian, L., Looger, L. L. & Tank, D. W. Functional imaging of
hippocampal place cells at cellular resolution during virtual navigation. *Nat Neurosci* **13**,
1433-1440, doi:10.1038/nn.2648 (2010).

47   McNamara, C. G., Tejero-Cantero, A., Trouche, S., Campo-Urriza, N. & Dupret, D.
Dopaminergic neurons promote hippocampal reactivation and spatial memory
persistence. *Nat Neurosci* **17**, 1658-1660, doi:10.1038/nn.3843 (2014).
48   Davoudi, H. & Foster, D. J. Acute silencing of hippocampal CA3 reveals a dominant role
in place field responses. *Nat Neurosci* **22**, 337-342, doi:10.1038/s41593-018-0321-z
(2019).
49   Hunsaker, M. R., Rosenberg, J. S. & Kesner, R. P. The role of the dentate gyrus, CA3a,b,
and CA3c for detecting spatial and environmental novelty. *Hippocampus* **18**, 1064-1073,
doi:10.1002/hipo.20464 (2008).
50   Dolleman-van der Weel, M. J. *et al.* The nucleus reuniens of the thalamus sits at the
nexus of a hippocampus and medial prefrontal cortex circuit enabling memory and
behavior. *Learn Mem* **26**, 191-205, doi:10.1101/lm.048389.118 (2019).
51   Nitz, D. & McNaughton, B. Differential modulation of CA1 and dentate gyrus
interneurons during exploration of novel environments. *J Neurophysiol* **91**, 863-872,
doi:10.1152/jn.00614.2003 (2004).
52   Yu, X., Knierim, J. J., Lee, I. & Shouval, H. Z. Simulating place field dynamics using spike
timing-dependent plasticity. *Neurocomputing* **69**, 1253-1259 (2006).
53   Lisman, J. & Spruston, N. Questions about STDP as a General Model of Synaptic
Plasticity. *Front Synaptic Neurosci* **2**, 140, doi:10.3389/fnsyn.2010.00140 (2010).
54   Graupner, M., Wallisch, P. & Ostojic, S. Natural Firing Patterns Imply Low Sensitivity of
Synaptic Plasticity to Spike Timing Compared with Firing Rate. *J Neurosci* **36**, 11238-
11258, doi:10.1523/JNEUROSCI.0104-16.2016 (2016).
55   Hussaini, S. A., Kempadoo, K. A., Thuault, S. J., Siegelbaum, S. A. & Kandel, E. R.
Increased size and stability of CA1 and CA3 place fields in HCN1 knockout mice. *Neuron*
72, 643-653, doi:10.1016/j.neuron.2011.09.007 (2011).
56   Bittner, K. C. *et al.* Conjunctive input processing drives feature selectivity in hippocampal
CA1 neurons. *Nat Neurosci* **18**, 1133-1142, doi:10.1038/nn.4062 (2015).
57   Fattahi, M., Sharif, F., Geiller, T. & Royer, S. Differential Representation of Landmark and
Self-Motion Information along the CA1 Radial Axis: Self-Motion Generated Place Fields
Shift toward Landmarks during Septal Inactivation. *J Neurosci* **38**, 6766-6778,
doi:10.1523/JNEUROSCI.3211-17.2018 (2018).
58   Thompson, L. T. & Best, P. J. Long-term stability of the place-field activity of single units
recorded from the dorsal hippocampus of freely behaving rats. *Brain Res* **509**, 299-308,
doi:10.1016/0006-8993(90)90555-p (1990).
59   Wood, E. R., Dudchenko, P. A., Robitsek, R. J. & Eichenbaum, H. Hippocampal neurons
encode information about different types of memory episodes occurring in the same
location. *Neuron* **27**, 623-633, doi:10.1016/s0896-6273(00)00071-4 (2000).
60   Krishnan, S., Cherian, C. & Sheffield, M. E. J. Changing reward expectation transforms
spatial encoding and retrieval in the hippocampus. *bioRxiv*, 2020.2009.2013.295469,
doi:10.1101/2020.09.13.295469 (2020).
61   Anderson, M. I. & Jeffery, K. J. Heterogeneous modulation of place cell firing by changes
in context. *J Neurosci* **23**, 8827-8835 (2003).

62　Hasselmo, M. E. & Eichenbaum, H. Hippocampal mechanisms for the context-dependent
retrieval of episodes. *Neural Netw* **18**, 1172-1190, doi:10.1016/j.neunet.2005.08.007
(2005).
63　Cai, D. J. *et al.* A shared neural ensemble links distinct contextual memories encoded
close in time. *Nature* **534**, 115-118, doi:10.1038/nature17955 (2016).
64　Allegra, M., Posani, L., Gomez-Ocadiz, R. & Schmidt-Hieber, C. Differential Relation
between Neuronal and Behavioral Discrimination during Hippocampal Memory
Encoding. *Neuron* **108**, 1103-1112 e1106, doi:10.1016/j.neuron.2020.09.032 (2020).
65　Madar, A. *Pattern Separation in the Hippocampus, in Health and Epilepsy* PhD thesis,
University of Wisconsin-Madison, (2018).
66　Bourboulou, R. *et al.* Dynamic control of hippocampal spatial coding resolution by local
visual cues. *Elife* **8**, doi:10.7554/eLife.44487 (2019).
67　Lee, S. W., O'Doherty, J. P. & Shimojo, S. Neural computations mediating one-shot
learning in the human brain. *PLoS Biol* **13**, e1002137, doi:10.1371/journal.pbio.1002137
(2015).
68　Stachenfeld, K. L., Botvinick, M. M. & Gershman, S. J. The hippocampus as a predictive
map. *Nat Neurosci* **20**, 1643-1653, doi:10.1038/nn.4650 (2017).
69　Abbott, L. F. & Blum, K. I. Functional significance of long-term potentiation for sequence
learning and prediction. *Cereb Cortex* **6**, 406-416, doi:10.1093/cercor/6.3.406 (1996).
70　Clewett, D., DuBrow, S. & Davachi, L. Transcending time in the brain: How event
memories are constructed from experience. *Hippocampus* **29**, 162-183,
doi:10.1002/hipo.23074 (2019).
71　Heys, J. G., Rangarajan, K. V. & Dombeck, D. A. The functional micro-organization of grid
cells revealed by cellular-resolution imaging. *Neuron* **84**, 1079-1090,
doi:10.1016/j.neuron.2014.10.048 (2014).
72　Aronov, D., Nevers, R. & Tank, D. W. Mapping of a non-spatial dimension by the
hippocampal-entorhinal circuit. *Nature* **543**, 719-722, doi:10.1038/nature21692 (2017).
73　Mukamel, E. A., Nimmerjahn, A. & Schnitzer, M. J. Automated analysis of cellular signals
from large-scale calcium imaging data. *Neuron* **63**, 747-760,
doi:10.1016/j.neuron.2009.08.009 (2009).
74　Tampuu, A., Matiisen, T., Olafsdottir, H. F., Barry, C. & Vicente, R. Efficient neural
decoding of self-location with a deep recurrent network. *PLoS Comput Biol* **15**,
e1006822, doi:10.1371/journal.pcbi.1006822 (2019).
75　Ho, J., Tumkaya, T., Aryal, S., Choi, H. & Claridge-Chang, A. Moving beyond P values: data
analysis with estimation graphics. *Nat Methods* **16**, 565-566, doi:10.1038/s41592-019-
0470-3 (2019).
76　Dragoi, G. & Tonegawa, S. Preplay of future place cell sequences by hippocampal
cellular assemblies. *Nature* **469**, 397-401, doi:10.1038/nature09633 (2011).
77　Madar, A. D., Ewell, L. A. & Jones, M. V. Pattern separation of spiketrains in hippocampal
neurons. *Sci Rep* **9**, 5282, doi:10.1038/s41598-019-41503-8 (2019).
78　Glaser, J. I. *et al.* Machine learning for neural decoding. arXiv:1708.00909 (2017).
<https://ui.adsabs.harvard.edu/abs/2017arXiv170800909G>.
79　Hochreiter, S. & Schmidhuber, J. Long short-term memory. *Neural Comput* **9**, 1735-
1780, doi:10.1162/neco.1997.9.8.1735 (1997).

Fig. 1: Experimental set up.

a, Top left, depiction of the virtual reality (VR) set up. Top right, the familiar (F) and two novel

(N1 and N2) environments. Bottom: Scheme of the experimental procedure. **b,** Top: Single

mouse behavior showing track position versus time during an F to N switch. Bottom: Summary

data across all mice of mean lap velocity during F and N (n = 20 sessions in 11 mice). Lap

velocity is normalized to the mean lap velocity in each mouse ± SEM, and each lap is compared

to the first lap in F using a one-way ANOVA with Tukey HSD post hoc test. ***, $P < 0.001$. **c,**

Brain slice showing specific CA3 expression of cre-dependent GCaMP6f (green) and DAPI for

nuclei (blue) from a Grik4-cre transgenic mouse. **d,** Example field of views (FOV) from

multiplane imaging in CA1 (top) and CA3 (bottom). Right: scheme of the position for the

objective during imaging.

Fig. 2: Place field emergence in novel environments is rapid in CA1 but gradual in CA3

a, Left: mean place fields (PFs) in F sorted by track position. Middle: mean activity of the same

neurons on the left in N. Right: mean place fields in N sorted by track position. Far right:

Percentage of place cells in F and N or common to both from all active neurons. CA1, top

panels; CA3, bottom panels. ΔF/F activity is normalized to each neurons' maximum transient

amplitude. *n* = 7 sessions in 4 mice for CA1, *n* = 13 sessions in 7 mice for CA3. Only place cells

with single PFs are displayed, but place cells with multiple PFs are included in the pie charts and

later analyses.   **b,** Pearson's correlation coefficient of each cell's average activity map within F

(blue, CA1 *n* = 1392, CA3 *n* = 320 neurons with activity in both halves of the F session) and

between F and N (red, CA1 *n* = 2283, CA3 *n* = 698 neurons with activity in both sessions and a

PF in one of them). Bootstrapped mean difference (Δ) with distribution of the mean (gray) and

95% confidence interval (error bar) on the right of each plot (see Methods). **c,** Example of an

instant PF (left) and delayed PF (right) in N. Arrows indicate PF onset lap.   **d,** Histograms of PF

onset laps in F (left, CA1 n = 1643 PFs, CA3 n = 336 PFs) and N (middle, CA1 n = 1712 PFs, CA3 n

= 532 PFs) and cumulative fraction plots for F and N (right). Wilcoxon rank sum test. ***, $P <$

0.001.   **e,** Cumulative fraction plots for CA1 and CA3 PF onset lap in F (left) and in N (right).

Wilcoxon rank sum test. ***, $P < 0.001$.   **f,** Example mouse showing true track position (black)

on laps 37-40 and the predicted position (red) decoded by an LSTM decoder (see Methods).

CA1, top; CA3, bottom.   **g,** Confusion matrix between the predicted (x-axis) and the true (y-axis)

position for the first lap in N. **h,** Predicted error on the first lap in N for CA1 and CA3. Each dot

represents the decoding error for one decoder trial built based on the activity of 200 randomly

chosen place cells from CA1 (left) or CA3 (right). n = 20 decoder trials. Bootstrapped mean

difference (Δ) with 95% CI (error bar) on the right.

Fig. 3: CA3 place fields exhibit relative lap-by-lap stability while CA1 place fields shift with experience in novel environments

a, Pearson's correlation coefficient of all neurons with place fields within N (average activity map in first 10 laps versus next 10 laps of session: CA1 *n* = 1238, CA3 *n* = 255 neurons). Bootstrapped mean difference (Δ) between CA1 and CA3 shown on the right. **b,** Example place cells with stable or shifting place fields (PFs) in N (lap-by-lap activity for the first 25 laps). White arrows depict the direction of shifting. **c,** Linear regression analysis on the center of mass (COM) of individual PFs (CA1 in N) shows a high number of significantly shifting PFs (blue and red dots and bars). The slope measures the direction and amplitude of the PF shift: for significantly shifting PFs, backward is blue, forward is red. CA3 has a lower proportion of significantly shifting PFs than CA1. **d,** Histograms of the COM difference between last and onset

laps (5-lap weighted average, see Methods) normalized by the number of laps for each PFs in N

for CA1 and CA3. Shuffling the lap numbers (grey distributions) reveals an overrepresentation

of backward shifting PFs, as confirmed by the bootstrapped mean difference (Δ, insets). **e,** Left,

Mean ± SEM, over all PFs, of 5-lap sliding average COM difference relative to lap 12 (see

Methods), in N. CA1 (green) and CA3 (orange). Linear regression analysis (on all data points, not

means) shows that the population of PFs shifts backwards significantly, both in CA1 and CA3.

***, $P < 0.001$. Right, The CA1 dataset was resampled 1000x using $n = 175$ PFs to match the

number of CA3 PFs and the slope and intercept of the regression line was measured each time

(green dots). CA1 slopes are always steeper than the CA3 slope indicating that the CA1

population shifts significantly faster than CA3.

Fig. 4: CA3 place fields exhibit higher stability across days than CA1 place fields and rapidly

reappear upon re-exposure to the environment

a, Experimental setup. Top: mice are recorded for 20 min in N across 2 days. Bottom: example

field of view from one imaging plane from CA1 and CA3 across 2 days showing the same cells.

scale bar = 50 μm. **b,** Top, example mouse behavior showing track position for the first lap in F

(blue), N day 1 (red) and N day 2 (purple). Bottom, summary across all mice of first lap velocity

in F, N day1 and N day2. Lap velocities were normalized to the mean velocity in F in each mouse

(n = 11 mice). Bootstrapped mean difference (Δ) between F and N day 1, and F and N day 2.

Note that lap velocity is only different on N day 1. **c,** Left: mean place fields in N on day 1 sorted

by track position. Middle: mean activity of the same neurons on the left on day 2 in N. Right:

mean place fields in N on day 2 sorted by track position. **d,** Histograms of Pearson's correlation

coefficient of average activity map across days in N. Spatial correlations greater than 0.5 were

considered stable place fields and less than 0.5 unstable. CA1 n = 409, CA3 n = 113 neurons with

PFs in both sessions. **e,** Fraction of stable place fields across days per mouse in CA1 and CA3.

Wilcoxon rank sum test. *P < 0.05. **f,** Cumulative fraction plots of Pearson's correlation

coefficient in N across days in CA1 and CA3 from the same data shown in (**c**). Wilcoxon rank

sum test. ***, P < 0.001. **g,** Example of a stable place field (left), an unstable place field with a

newly formed place field on day 2 (middle) and a newly formed place field on day 2 (right).

Place field transients shown lap-by-lap for all laps in N on day 1 and day 2 from the same 3 cells.

h, Histograms of place field onset laps in N on day 2 for stable place fields (left), and newly

formed place fields (middle) in CA1 (top) and CA3 (bottom). Right: cumulative fraction plots of

histogram data. Wilcoxon rank sum test. ***, P < 0.001. CA1 stable n = 251 PFs, newly formed,

n = 333 PFs, CA3 stable n = 72 PFs, newly formed, n = 94 PFs.

Fig. 5: Place field shifting resets and continues upon first re-exposure to N on day 2.

a, Histogram of place field (PF) shifts across days for CA1 (left, n =171 PFs) and CA3 (right, n = 43

PFs). The shifts were calculated as the difference between each PF's center of mass (COM) of

the last active lap on day 1 and first active lap in day 2. Only place cells with PFs on both days

are included. Wilcoxon signed rank test, N.S., *P* > 0.05. **b,** Comparison of COM position at the

end of day 1 and COM position at the start of day 2 for PFs with day-to-day correlation > 0.5

and significant shifting on day 1 (see Fig. 3c). Dashed diagonal: identity line. On day 2, PFs tend

to reset towards the direction opposite of shifting on day 1. Wilcoxon sign-rank test on Day 2-

Day 1 difference from 0 for backward and forward shifting (*P*-values and median in panel). **c,**

Comparison of PF shifting slope on both days. No clear correlation is observed, but the Day 2-

Day 1 difference is not significantly different from 0 (Wilcoxon sign-rank: *P*-values and medians

in panel) and individual PF shifting was slow (close to 0 cm/lap) on both days. **d,** Same as Fig.

3e but for N day 2, using all PFs recorded that day. CA1 (green; *n* = 371 PFs from 3 mice) and

724   CA3 (orange; n = 84 PFs from 6 mice). The resampling analysis (1000 random resamples

725   matching the CA3 sample size) shows that CA1 slopes are always steeper than the CA3 slope,

726   indicating that on day 2 the CA1 population shifts backward faster than CA3.

Fig. 6: Population backward shifting slows down with familiarization across days

a, Top, histograms of COM_{last_lap}-COM_{onset_lap} (session shift defined as in Fig. 3d) in Familiar (F), N

day 1 and N day 2, for CA1. Bottom, Bootstrapped mean difference (Δ) between the three

conditions. **b,** Same as a for CA3. **c,** Left, Same as Fig. 3e, comparing CA1 population shifts in F

(blue), N day 1 (red) and N day 2 (purple). ***, $P < 0.001$. The resampling analysis uses a

number of PFs equal to 80 percent of the number of PFs in the condition with lowest number of

PFs (1000 resamples). The lack of distribution overlap between conditions indicates dynamics

are significantly different. **d,** Same as c for CA3. Overlap in the slope distributions suggests that

population dynamics are similarly flat (slopes close to 0) although CA3 PFs shifting become

more homogeneous (smaller variance) with familiarization.