Estimating the time since admixture from phased and unphased molecular data

Thijs Janzen[1,2,†,*], Verónica Miró Pina[3,4,5,†]

1 Groningen Institute for Evolutionary Life Sciences, University of Groningen, Box 111039700 CC Groningen, The Netherlands

2 Carl von Ossietzky University, Carl-von-Ossietzky-Str. 9–11, 26111, Oldenburg, Germany

3 Instituto de Investigaciones en Matemáticas Aplicadas y Sistemas (IIMAS), Universidad Nacional Autónoma de México (UNAM), México City, México

4 Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Dr. Aiguader 88, Barcelona 08003, Spain

5 Universitat Pompeu Fabra (UPF), Barcelona, Spain

† These authors contributed equally to this work. * Corresponding author: t.janzen@rug.nl

Abstract

After admixture, recombination breaks down genomic blocks of contiguous ancestry. The breakdown of these blocks forms a new 'molecular clock', that ticks at a much faster rate than the mutation clock, enabling accurate dating of admixture events in the recent past. However, existing theory on the break down of these blocks, or the accumulation of delineations between blocks, so called 'junctions', has mostly been limited to using regularly spaced markers on phased data. Here, we present an extension to the theory of junctions using the Ancestral Recombination Graph that describes the expected number of junctions for any distribution of markers along the genome. Furthermore, we provide a new framework to infer the time since admixture using unphased data. We demonstrate both the phased and unphased methods on simulated data and show that our new extensions performs better than previous methods, especially for smaller population sizes and for more ancient admixture times. Lastly, we demonstrate the applicability of our method on an empirical dataset of

labcrosses of yeast (*Saccharomyces cerevisae*) and on two case studies of hybridization in swordtail fish and *Populus* trees.

Keywords

Admixture, hybridization, recombination, junctions, phasing

1 Introduction   1

The traditional view where taxa accumulate incompatibilities over time and gradually become   2
reproductively isolated from each other – speciation – has led to insight into the processes generating   3
and maintaining biodiversity (Coyne and Orr, 2004). However, it has become apparent that   4
reticulate evolution is common: taxa do not necessarily only branch, they can also come back   5
together – hybridisation – (Abbott et al., 2013). This means that, while the ancestry of non-   6
recombining DNA can be traced linearly back in time (e.g. mitochondrial lineages), the ancestry   7
of genomes backward in time will branch at admixture events. In plants, it has been known for   8
quite some time that hybridization is a widespread phenomenon that can not only generate viable   9
offspring, but also potentially lead to the formation of new taxa, and ultimately, species (Grant,   10
1981). It has long been debated whether this process is also common in animals, but over the past   11
few years numerous examples have appeared, including, but not limited to, butterflies (Mavárez   12
et al., 2006; Capblancq et al., 2015), cichlid fishes (Koblmüller et al., 2007; Keller et al., 2013),   13
warblers (Brelsford et al., 2011), fruit flies (Schwarz et al., 2005) and sculpins (Nolte et al., 2005).   14

Understanding the timeline of these hybridization events is paramount in obtaining a full   15
understanding of the process and its impact. A 'recombination clock' is particularly useful for   16
studying recent evolutionary dynamics since detectable recombination events are markers of admixture   17
(Baird, 2006), and the statistical association across loci created by admixture (admixture linkage   18
disequilibrium) decays slowly (Baird, 2015). After admixture of two taxa, contiguous genomic   19
blocks are broken down by recombination over time. The delineations between these blocks were   20
termed 'junctions' by Fisher (1949, 1954), and inheritance of these junctions is similar to that of   21

point-mutations. Further work on the theory of junctions has shown how they accumulate over time   22
for sib-sib mating (Fisher, 1954), self-fertilization (Bennett, 1953), alternate parent-offspring mating   23
(Fisher, 1959; Gale, 1964), a randomly mating population (Stam, 1980; Baird et al., 2003), and for   24
sub-structured populations (Barton, 1983; Baird, 1995; Chapman and Thompson, 2002, 2003).   25

So far, applying the theory of junctions has proved difficult, as it requires extensive genotyping of   26
the admixed individuals, but also of the source taxa. With the current decrease in genotyping costs   27
(Muir et al., 2016), such analyses are coming within reach, and frameworks are being developed that   28
assist in inferring local ancestry and detecting junctions, given molecular data of source and admixed   29
taxa (Paşaniuc et al., 2009; Maples et al., 2013; Guan, 2014; Corbett-Detig and Nielsen, 2017; Medina   30
et al., 2018; Svedberg et al., 2021). Nevertheless, molecular data always paints an imperfect image of   31
ancestry along the genome, and inferring the number of junctions in a chromosome remains limited   32
by the number of diagnostic markers available (see Fig 1, first panel). Previous work on the theory   33
of junctions does not take into account the effect of a limited number of genetic markers, and so   34
far this effect had to be corrected using simulations (MacLeod et al., 2005; Buerkle and Rieseberg,   35
2008). Recent work by Janzen et al. (2018) resolves this issue by extending the theory of junctions   36
with the effect of using a limited number of markers, but they had to assume an evenly spacing of   37
markers. However, molecular markers are rarely evenly spaced. The first result we present here is   38
an extension of the theory of junctions which includes the effect of marker spacing on inferring the   39
number of junctions in a genome and that has the advantage of working well with fewer markers.   40

Furthermore, most existing theory on the accumulation of junctions is developed for the case   41
where ancestry can be determined within a single chromosome (with the exception of ANCESTRY   42
HMM, (Corbett-Detig and Nielsen, 2017)). For diploid species, sequencing data presents itself   43
as the pileup of ancestry across both chromosomes, requiring an additional step to separate the   44
contributions of both chromosomes, called 'phasing' (see Fig 1, second and third panel). Phasing   45
methods can be classified into three main categories, including haplotype-resolved genome sequencing,   46
pedigree-based methods and statistical methods. Haplotype-resolved genome sequencing methods   47
(reviewed in Snyder et al. (2015)), yield accurate results, but are expensive and require a large   48
number of sequences in order to be able to resolve the required haplotypes (but see (Lutgen et al.,   49

2020)). Pedigree based phasing methods do not require resolved haplotypes, but instead rely on   50
accurate genotyping of related individuals to resolve the required phase. These methods often yield   51
good results but their application has been limited to humans, where large pedigree datasets are   52
available (Browning and Browning, 2011; Loh et al., 2016a; Kong et al., 2008). Statistical methods   53
do not require large pedigree datasets, and are based on recombination rate estimates and allele   54
frequencies in a population. While some of these methods make use of a reference genome (for   55
example Eagle (Loh et al., 2016b), Beagle (Browning and Browning, 2007) or ShapeIt (O'Connell   56
et al., 2016)), others allow *de novo* haplotype aware assembly (for example POLYTE (Baaijens and   57
Schönhuth, 2019)). Further improvements include the usage of third-generation sequencing data   58
(Tourdot and Zhang, 2019; Kronenberg et al., 2019; Ebler et al., 2019; Tangherloni et al., 2019).   59
However, data from hybrid populations is not often available in this form. Across these three groups   60
of methods, the overarching theme is that phasing is often costly and accuracy can be left wanting.   61
Yet, inclusion of information from both chromosomes is expected to improve inference of the onset   62
of admixture considerably and hence expansion of the theory of junctions towards a framework that   63
takes into account data from both chromosomes is warranted.   64

Here we provide a framework to estimate the time since admixture using phased or unphased data   65
from two homologous chromosomes, taking into account marker spacing along the chromosome. Our   66
framework is based on modelling the joint genealogy of loci that are located in the same chromosome   67
or in two homologous chromosomes, using the Ancestral Recombination Graph (ARG) (Hudson,   68
1983; Griffiths, 1991; Griffiths and Marjoram, 1997). It has the advantage of being fast since it relies   69
on mathematical computations and does not require simulations. It has been implemented in the   70
R package 'junctions'. In comparison with previous methods (ANCESTRY HMM, Corbett-Detig and   71
Nielsen (2017)), our method has the advantage of working well with small population sizes and   72
larger admixture times.   73

Our paper is organised as follows. In section 2, we introduce our model, which is a simplified   74
version of the ARG and present three maximum-likelihood methods to infer the time since admixture   75
in hybrid populations: the first method uses information from a single chromosome, the second   76
method uses phased data from two homologous data and the third method uses unphased data from   77

two homologous chromosomes. At the end of section (2.4), we validate our methods using simulations. ₇₈

In section 3, we provide a detailed comparison to previous methods. Lastly, in section 4 we apply ₇₉

them to a dataset from experimental evolution in yeast and to two case studies of hybridization in ₈₀

swordtail fish and *Populus* trees. ₈₁

2   Materials and methods ₈₂

2.1   Mathematical model ₈₃

We assume a diploid population that evolves according to Wright-Fisher dynamics, i.e. generations ₈₄

are non-overlapping, mating is random and all individuals are hermaphrodites. We only keep track ₈₅

of one chromosome (or one pair of chromosomes), assuming that the accumulation of junctions on ₈₆

different pairs of chromosomes is independent of each other (see Section 2.2.1). We assume that ₈₇

hybridization occurred T generations ago between two source taxa, $\mathcal{P}$ and $\mathcal{Q}$. The proportion of ₈₈

individuals from source $\mathcal{P}$ in the initial generation is p and the proportion of individuals from source ₈₉

$\mathcal{Q}$ is $q = 1 - p$. We define the initial heterogenicity $H_0 := 2pq$ as the probability that, when sampling ₉₀

two individuals from this generation they are from different source taxa. This also represents the ₉₁

probability that at any given locus, one allele can be traced to source taxon $\mathcal{P}$ and the other allele ₉₂

to $\mathcal{Q}$. Following Janzen et al. (2018), instead of referring to the lesser known term of heterogenicity, ₉₃

we use the term of heterozygosity in the manuscript throughout because these two concepts are ₉₄

mathematically equivalent. ₉₅

  We assume that the length of the chromosome is C Morgan and that there are n molecular ₉₆

markers whose positions are given by $(z_1, \ldots, z_n) \in [0, C]$. For two consecutive markers at sites z_i ₉₇

and z_{i+1}, we define $d_i = z_{i+1} - z_i$, the distance between them in Morgan. We assume that there are ₉₈

enough markers on the chromosome such that the d_i's are small compared to 1. The genealogy of ₉₉

these n (or $2n$ for a diploid genome) loci is given by the Ancestral Recombination Graph (ARG), ₁₀₀

defined in Hudson (1983); Griffiths (1991); Griffiths and Marjoram (1997). The ARG is a branching- ₁₀₁

coalescence process which follows backwards in time the ancestry of loci sampled in the present ₁₀₂

population (time 0). If two loci belong to the same block at time t, they are identical-by-descent ₁₀₃

(IBD) with respect to generation t, i.e. the sampled loci have been inherited from the same individual living t units of time ago.

Although the ARG for many loci has complicated transition rates and is a computationally intensive model, here we consider only two loci (or two pairs of loci for a diploid genome) at a time. This is sufficient since for our maximum likelihood approach we only use the expected number of junctions – and not its variance or higher moments (see e.g. equation 3).

We assume that the number of diploid individuals N is large, so that we can neglect some transitions (double coalescences and simultaneous coalescence and recombination) and that $d_i \ll 1$ so that there is no more than one crossover per generation between two molecular markers and the mutation rates are small enough so that we can neglect mutations that happened between the admixture time and the present.

2.2   Two sites, one chromosome

The aim of this section is to derive a formula for the expected number of observed junctions on one chromosome given the population size N, the distances between the markers $(d_1, \ldots, d_n)$ and the initial heterozygosity $H_0 := 2pq$. We start by considering two consecutive loci z_i and z_{i+1} sampled in the same chromosome in the present population. The ARG for these two sites has two possible states:

- If the loci are in state $(z_i \sim z_{i+1})$ at time t, it means that the two sampled loci have been inherited from *the same individual* living t generations before, i.e. that they are IBD.

- If the loci are in state $(z_i \nsim z_{i+1})$, it means that the two loci have been inherited from different individuals.

To model the observed junctions, we look at identity-by-descent with respect to time T, when admixture took place. Recall that we only model observed junctions, i.e. where adjacent markers have alleles corresponding to different source taxa (which corresponds to an odd number of "true" junctions, see 1 and its caption).

- If the two loci are in state $(z_i \sim z_{i+1})$ at time T, the ancestor can be from source $\mathcal{P}$ (with probability p) or red (with probability q). In both cases, we observe no junction.

- If the two loci are in state $(z_i \not\sim z_{i+1})$ at time T (not IBD), they have two different ancestors at generation T, which can be:

  - Both blue (with probability p^2) or both red (with probability q^2), in this case there is no junction.

  - One blue and one red (with probability $2pq$), in this case, there is a junction in the chromosome sampled in the present.

The dynamics of the ARG are controlled by two types of events:

- **Recombination** $(z_i \sim z_{i+1}) \rightarrow (z_i \not\sim z_{i+1})$ with probability d_i,

- **Coalescence** $(z_i \not\sim z_{i+1}) \rightarrow (z_i \sim z_{i+1})$ with probability $\frac{1}{2N}$.

Since $N \gg 1$ and the $d_i \ll 1$, other events (such as simultaneous coalescence and recombination events) have probabilities that are negligible. This yields the following transition matrix:

$$\bar{M} = \begin{pmatrix} 1 - d_i & d_i \\ \frac{1}{2N} & 1 - \frac{1}{2N} \end{pmatrix}.$$

Let $\bar{P}_t$ be the probability vector at time t for this Markov chain with two states. $(\bar{P}_t)_1$ is the probability of $(z_i \sim z_{i+1})$ at time t and $(\bar{P}_t)_2$ the probability of $(z_i \not\sim z_{i+1})$ at time t. We have $\bar{P}_0 = (1,0)$ (in the present we sample the two loci in the same individual) and $\bar{P}_t = \bar{P}_0 \bar{M}^t$. We denote by $\mathbb{P}(J_T(z_i, z_{i+1}))$ the probability that a junction is observed between z_i and z_{i+1} (conditioning on the fact that hybridization occurred T generations ago). We have

$$\mathbb{P}(J_T(z_i, z_{i+1})) \;=\; H_0 (\bar{P}_t)_2, \tag{1}$$

which corresponds to the probability that the two loci were carried by different individuals T generations ago and they from different source taxa (see Fig 2, left panel).

Solving equation (1) gives

$$\mathbb{P}(J_T(z_i, z_{i+1})) = H_0 \frac{2N d_i}{2N d_i + 1} \left(1 - \left(1 - d_i - \frac{1}{2N} \right)^T \right). \tag{2}$$

Let $\mathbb{E}(J_T)$ be the expected number of observed junctions on one chromosome, we have

$$\mathbb{E}(J_T) \;=\; \sum_{i=1}^{n-1} \mathbb{P}(J_T(z_i, z_{i+1})) \;=\; H_0 \sum_{i=1}^{n-1} \frac{2Nd_i}{2Nd_i + 1} \left(1 - \left(1 - d_i - \frac{1}{2N} \right)^T \right). \tag{3}$$

2.2.1   Maximum Likelihood

For each chromosome, we can calculate the likelihood of observing the data, where the data are $n-1$ pairs of markers ($z_i \sim z_{i+1}$), which is given by

$$\prod_{i=1}^{n-1} \mathbb{P}((z_i, z_{i+1})),$$

where $\mathbb{P}((z_i, z_{i+1}))$ is the probability of observing the pair of markers (z_i, z_{i+1}) given by:

- $\mathbb{P}((z_i, z_{i+1})) = \mathbb{P}(J_T(z_i, z_{i+1}))$, if there is a junction between the two markers. This is given by equation (2).

- $\mathbb{P}((z_i, z_{i+1})) = 1 - \mathbb{P}(J_T(z_i, z_{i+1}))$, if there is no junction between the two markers.

For different chromosomes sampled in the same individual, we can assume independence between chromosomes and the full likelihood of observing the data, given the time since admixture, is given by the product of these likelihoods, e.g.:

$$\mathcal{L} = \prod_{c=1}^{C} \prod_{i=1}^{n_C - 1} \mathbb{P}(z_i, z_{i+1}), \tag{4}$$

where C indicates the number of chromosomes and n_C indicates the number of markers on chromosome C.

When we have data from several individuals (which are not independent, due to coalescence), we first compute the maximum likelihood estimator of T for each individual (using (4)) and then we average across all individuals in the sample. Confidence intervals (CI's) reported are then the 95 % interquartile range in the age estimates across all individuals.

State		n_i	n_{i+1}	n_{tot}
S^1	$(z_i z_{i+1}), (z_i z_{i+1})$	2	2	2
S^2	$(z_i z_{i+1})(z_i)(z_{i+1})$	2	2	3
S^3	$(z_i)(z_i)(z_{i+1})(z_{i+1})$	2	2	4
S^4	$(z_i z_{i+1})(z_i)$ or $(z_i z_{i+1})(z_{i+1})$	2 (or 1)	1 (or 2)	2
S^5	$(z_i)(z_{i+1})(z_{i+1})$ or $(z_{i+1})(z_i)(z_i)$	1 (or 2)	2 (or 1)	3
S^6	$(z_i z_{i+1})$	1	1	1
S^7	$(z_i), (z_{i+1})$	1	1	2

Table 1. States of the reduced ARG. n_i (resp. n_{i+1}) denotes the number of ancestors of site z_i (resp. z_{i+1}) and n_{tot} the total number of ancestors to the sample.

2.3 Two sites, two chromosomes

We consider two consecutive loci z_i and z_{i+1}, which are at distance d_i (in Morgan), that we sample 163

in two homologous chromosomes. The ARG for these 2 sites in 2 chromosomes has 7 states (see 164

Durrett (2008), Chapter 3). To describe them, we borrow the notation from Durrett (2008) and we 165

write $(z_i z_{i+1})$ to indicate that sites z_i and z_{i+1} are IBD, and notation (z_i) (or (z_{i+1})) to indicate 166

that the ancestor to z_i (or (z_{i+1})) is only ancestor to one of the two sites. The resulting 7 states are 167

summarized in Fig 1. An example of realization of this process is shown in Fig 2 (right panel). 168

The initial state is S^1 because in the present time we sample two different loci in two different chromosomes. The transition matrix of the ARG with 2 loci and a sample size 2 can be approximated, when $N \gg 1$ by

$$M^{(i)} = \begin{pmatrix} 1 - \frac{1}{2N} - 2d_i & 2d_i & 0 & 0 & 0 & \frac{1}{2N} & 0 \\ \frac{1}{2N} & 1 - 3\frac{1}{2N} - d_i & d_i & 2\frac{1}{2N} & 0 & 0 & 0 \\ 0 & 2\frac{1}{2N} & 1 - 4\frac{1}{2N} & 0 & 2\frac{1}{2N} & 0 & 0 \\ 0 & 0 & 0 & 1 - \frac{1}{2N} - d_i & d_i & \frac{1}{2N} & 0 \\ 0 & 0 & 0 & 2\frac{1}{2N} & 1 - 3\frac{1}{2N} & 0 & \frac{1}{2N} \\ 0 & 0 & 0 & 0 & 0 & 1 - d_i & d_i \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{2N} & 1 - \frac{1}{2N} \end{pmatrix}.$$

All other potential events (e.g. double crossovers or simultaneous crossover and coalescence events) 169

have probabilities that are negligible compared to $1/2N$. 170

Let $P_t^{(i)}$ be the vector containing the probabilities of observing each of the states $(S^1, \ldots, S^7)$ at time t. $P_t^{(i)}$ satisfies

$$P_t^{(i)} = P_0 (M^{(i)})^t,$$

where $P_0 = (1, 0, 0, 0, 0, 0, 0)$, since at time 0 we sample all loci in two homologous chromosomes. This equation can only be solved numerically. Recall that the stationary distribution of this process $P^{(i)}$ satisfies

$$P^{(i)} = P^{(i)} M^{(i)}$$

and has the analytical expression

$$P^{(i)} \;=\; \left(0, 0, 0, 0, 0, \frac{1}{2d_i N + 1}, \frac{2d_i N}{2d_i N + 1} \right).$$

Thus, for large values of t the system reduces to states S^6 and S^7, which means that each locus has only one ancestor i.e. forwards in time the process has reached fixation (at each locus). Recall that state S^6 is the state where there is one ancestor for the sample thus we observe no junctions on either chromosome. Furthermore, recall that state S^7 is the state where there are two ancestors, one for the first locus and one for the second locus, and with probability $2pq$ each one of them comes from a different source taxa. This is exactly the probability of observing a junction when $t \to \infty$ for one chromosome (equation 2). In other words, when t is very large, fixation is reached and the two sampled chromosomes are homozygous so the problem reduces to the single chromosome case.

2.3.1 Maximum Likelihood, phased data

We first consider the case of phased data. Each pair of homologous markers can be in one of four states:

- PP i.e. both homologous markers carry the allele from source $\mathcal{P}$,

- QQ i.e. both homologous markers carry the allele from source $\mathcal{Q}$,

- PQ i.e. the marker on the first chromosome carries the allele from source $\mathcal{P}$ and the marker on the second chromosome carries the allele from source $\mathcal{Q}$,

- QP i.e. the marker on the first chromosome carries the allele from source $\mathcal{Q}$ and the marker on the second chromosome carries the allele from source $\mathcal{P}$.

The data can then be represented as a sequence $(O_i, 1 \leq i \leq n)$ that takes values in $\{PP, QQ, PQ, QP\}$ such that O_i is the state of the i-th marker. To derive a maximum likelihood formula for the time since admixture T, we compute the probability of each sequence in $\{PP, QQ, PQ, QP\}^n$ given T, N, C, the distances between the n loci and the initial heterozygosity H_0.

We want to compute the probability of our observations $(O_1, \ldots, O_n)$. These n observations are not independent, as there are non-trivial correlations between loci along the chromosome. However, we can neglect long-range dependencies and assume that O_i only depends on O_{i-1}, i.e. that the probability of observing $(O_1, \ldots, O_n)$, t units of time after hybridization is

$$\mathbb{P}_t((O_1, \ldots, O_n)) \; = \; \mathbb{P}_t(O_1, O_2) \prod_{i=2}^{n-1} \mathbb{P}_t(O_{i+1}|O_i).$$

Recall that ignoring long-range dependencies is a natural approximation and it has been used for example by McVean and Cardin (2005) to define the sequentially Markov coalescent. To compute $\mathbb{P}_t(O_{i+1}|O_i)$, we use the ARG for markers at z_i and z_{i+1} denoted by (Γ_t^i) (and to compute $\mathbb{P}(O_1, O_2)$, we use (Γ_t^1)). For example, we can observe $O_1 = PP$ and $O_2 = QQ$ if:

- $\Gamma_t^1 = S^3$ and the two ancestors for locus 1 are from source taxa $\mathcal{P}$ and the two ancestors for locus 2 from $\mathcal{Q}$, which happens with probability $p^2 q^2$ or,

- $\Gamma_t^1 = S^5$, with probability 1/2 there are two ancestors for locus 1 and one for locus 2 and they are from desired source taxa with probability $p^2 q$. With probability 1/2 there is one ancestor for locus 1 and two for locus 2 and they are from the desired source taxa with probability pq^2 or,

- $\Gamma_t^1 = S^7$ and the ancestor to 1 is from source taxa $\mathcal{P}$ and the ancestor to 2 from $\mathcal{Q}$, which happens with probability pq.

To sum up, when $O_1 = PP$ and $O_2 = QQ$,

$$\mathbb{P}_t(O_1, O_2) \;=\; p^2 q^2 (P_t^{(1)})_3 + \frac{1}{2}(pq^2 + qp^2)(P_t^{(1)})_5 + pq(P_t^{(1)})_7.$$

The probabilities for all combinations of O_1 and O_2 are listed in Fig 3. To compute $\mathbb{P}_t(O_{i+1}|O_i)$ we use Bayes' formula:

$$\mathbb{P}_t(O_{i+1}|O_i) = \frac{\mathbb{P}_t(O_i, O_{i+1})}{\mathbb{P}_t(O_i)},$$

where, using the total probability theorem, $\mathbb{P}_t(O_i)$ can be obtained by summing over the appropriate   204
row in Fig 3. Then, the total probability of observing the data, given N and t, i.e.   205

$$\mathbb{P}_t((O_1, \ldots, O_n)) \;=\; \mathbb{P}_t(O_1, O_2) \prod_{i=2}^{n-1} \frac{\mathbb{P}_t(O_i, O_{i+1})}{\mathbb{P}_t(O_i)} \tag{5}$$

can be maximized in order to find the maximum likelihood estimator of t and N. If the data consists   206
of multiple chromosomes from the same individual, we can calculate the joint likelihood of observing   207
the data, given t and N by assuming independence across chromosomes and calculating the likelihood   208
as the product across chromosomes. As in the one chromosome case, we provide estimates of T by   209
averaging across the different individuals of the sample, and the CI's reflect the 95 % interquartile   210
range of the T estimates across the different individuals of the sample.   211

2.3.2   Maximum Likelihood, unphased data   212

If the data is unphased, we cannot distinguish which allele is in which of the two homologous   213
chromosomes. We can observe one of these three states at each marker: 214

- P i.e. we only observe the allele from source $\mathcal{P}$, i.e. both chromosomes carry the allele from   215
  source $\mathcal{P}$,   216

- Q i.e. we only observe the allele from source $\mathcal{Q}$.   217

- x i.e. we observe both alleles, i.e. each one of the two homologous chromosomes carries a   218
  different allele.   219

The data can then be represented as a sequence (O_i) of length n that takes values in $\{P, Q, x\}$ 220
such that O_i is the state of the i-th marker. We can perform exactly the same method, as in the 221
previous section, except that now the probabilities of each state are given by Fig 4. Again, if the 222
data consists of multiple chromosomes from the same individual, we can calculate the joint likelihood 223
of observing the data, given t and N by assuming independence across chromosomes and calculating 224
the likelihood as the product across chromosomes. 225

2.4   Individual based simulations   226

To test the validity of our maximum likelihood approach, we use individual based simulations, 227
as described in (Janzen et al., 2018), i.e. Wright-Fisher type simulations of randomly mating 228
populations of constant size N, with non-overlapping generations. We then recover local ancestry by 229
analyzing ancestry at n markers whose positions are chosen uniformly at random along the genome. 230

Unless otherwise specified, we show how time can be accurately inferred for a population of $10,000$ 231
individuals, for time points between the first generation and 1,000 generations. We use $n = 10,000$ 232
markers, which is considered to be sufficient to detect the majority of accumulated junctions (Janzen 233
et al., 2018). We report our findings across 100 replicates, where in each replicate 10 individuals 234
were randomly selected from the admixed population and used to infer T. We have simulated with 235
three different values of the initial proportion of source taxa $\mathcal{P}$, ($p \in \{0.053, 0.184, 0.5\}$), to vary the 236
initial heterozygosity H_0 in $\{0.1, 0.3, 0.5\}$.   237

We have first compared the methods we have developed here to previous methods based on 238
the theory of junctions (Figure 5). We observe that, when the number of markers is low, previous 239
methods, that do not take into account marker spacing, tend to underestimate the time since 240
admixture, which is not the case for our methods.   241

We have then compared the estimations of the time since admixture, using the method for one 242
chromosome and the method for two chromosomes (phased) (Fig 6). We observe that using data 243
from the two homologous chromosomes allows to infer the time since admixture more accurately, 244
since it reduces uncertainty.   245

Using unphased data instead of phased data might introduce additional error and in Fig 7 we 246

compare the methods that use phased or unphased information of two homologous chromosomes. 247
We observe that both methods yield very similar results in terms of the relative error. This can 248
be due to the fact that homozygous sites have an important contribution to the likelihood and the 249
uncertainty that comes from sites that are of type x (in the unphased case) is well managed by our 250
method. 251

Finally, we explore error in phasing assignment (switching error). We simulate the effect of 252
error in phasing assignment by randomly swapping a fraction of the markers between chromosomes. 253
We explore phasing error in $\{0.0025, 0.005, 0.0075, 0.01, 0.02\}$. These errors are comparable to the 254
switching error rates reported in the literature. For example, Choi et al. (2018) compared different 255
phasing methods and reported switching error rates between 0.1% and 2%. (Notice that these error 256
rates are for human data where there are good quality references and sample sizes are large). More 257
recent reference-free methods (based on third generation sequencing techniques) report switching 258
error rates of 1-2% (see for example Tourdot and Zhang (2019); Ebler et al. (2019); Kronenberg et al. 259
(2019)). Switching error rate error has strong effects on the inferred time since admixture, as shown 260
in the bottom panel in Fig 8. Generally, imposed errors increase the inferred age, by introducing 261
novel junctions due to mis-phased markers. 262

Another important source of error is the lack of coverage, which would have the effect of reducing 263
the number of markers. An analysis of the sensitivity of our method to reducing the number of 264
markers can be found in the S3 Appendix, and shows that a reduction of coverage also tends to lead 265
to an overestimation of the time since admixture. 266

3   Comparison with other methods   267

Our framework provides a methodology to infer the time since admixture, provided that local 268
ancestry is known. Previous methods aiding in inferring the time since admixture, such as ELAI 269
(Guan, 2014) and ANCESTRY HMM (Corbett-Detig and Nielsen, 2017) jointly infer the time since 270
admixture and local ancestry, because they use the time since admixture to correct for the impact of 271
recombination on local ancestry estimates. We have chosen to compare our framework with results 272
obtained using ANCESTRY HMM, which we currently consider the most accurate method available 273

to jointly infer local ancestry and time since admixture. We do so in a two-fold manner: first, we 274
directly compare the ability of both methods to infer the time since admixture, provided data with 275
known local ancestry. Secondly, we provide both packages with imperfect data, where local ancestry 276
is uncertain. Because our framework is not able to infer local ancestry, we directly use the inferred 277
local ancestry of ANCESTRY HMM, which might add an extra layer of error, but is a good reflection 278
of an empirical use-case (see also section 4.3). For both approaches (known and uncertain ancestry), 279
data was simulated using the junctions package, using a population size N of [1000, 10000] and 280
imposing n markers, where n was [1000, 10000, 40000] (in line with the number of markers available 281
in the Swordtail and *Populus* datasets in section 4). Simulations were run for 10,000 generations, 282
using $H_0 = 0.5$ and $C = 1$. At each generation, time since admixture was inferred for 10 individuals 283
sampled randomly from the population. Time since admixture was inferred given the superimposed 284
n markers, whose location was drawn randomly in [0, 1]. For each parameter combination, 10 285
replicate simulations were performed. When the maximum likelihood did not converge to a final 286
value within the given range, the data point was removed from the analysis. 287

3.1 Known ancestry 288

Because ANCESTRY HMM requires information on allele frequencies in the source taxa in order to 289
jointly infer local ancestry and time since admixture, input data for ANCESTRY HMM was created 290
assuming both source taxa were fully separated, e.g. two alleles were differentially fixed across 291
the source taxa. Results show (Figure 9) that in the limit of large N, our method performs as 292
good as ANCESTRY HMM. However, for smaller population sizes, we observe that ANCESTRY HMM 293
becomes increasingly incorrect as the time since admixture increases, in contrast to our method, 294
which remains accurate. 295

3.2 Uncertain ancestry 296

To mimic uncertainty in ancestry, we explored different degrees of uncertainty, indicated by different 297
values of the allele frequency differential $d_x = A_x - A_y$, where A_x is the frequency of allele A in 298
population x and A_y is the frequency of allele A is population y, following Shriver et al. (1997). 299

Higher values of d_x indicate higher ancestry information content in the respective marker. Here, we explored values of d_x in [0.5, 0.7, 0.9]. Because our framework does not infer local ancestry, we used the inferred local ancestry by ANCESTRY HMM, using a threshold of including only markers with at least 95 % certainty of ancestry. For a large population size ($N = 10,000$, Supplementary material S5) we observe that ANCESTRY HMM consistently underestimates the true age when the time since admixture becomes large, in line with our findings with known ancestry (Figure 9). Our method underestimates the true age even further, most likely as a result of a stacking of errors: because our method can not infer local ancestry, we have used the local ancestry estimates of ANCESTRY HMM. However, when population size is small ($N = 1000$, Figure 10), we observe a striking phenomenon (Figure 10): when the time since admixture becomes large, our framework starts inferring the time correctly again, whereas the age estimates obtained using ANCESTRY HMM remain incorrect and seem to reach a plateau. This effect only occurs when there is a sufficient number of markers of reasonable quality, as we observe that when $n = 1000$ and $d \leq 0.7$, our framework is also unable to accurately infer the time since admixture. We believe that the improved performance when time since admixture becomes large is the result of the interplay between two phenomena. Firstly, while our method can work for any value of T, ANCESTRY HMM typically requires $T \ll \log(N)$ (this is the same assumption used in SMC' (Liang and Nielsen, 2014)). This could introduce some error in the ancestry assignment and thus lead to the underestimates obtained using ANCESTRY HMM. Secondly, our method seems to work better with a small number of informative markers. Combined, our method outperforms ANCESTRY HMM in inferring the time since admixture when population size and number of markers is small.

4 Results

4.1 Saccharomyces cerevisiae

Experimental evolution provides an important reference point to verify our findings. Here, we reanalyze data from an Advanced Intercross Line (AIL) experiment, where two highly differentiated yeast (*Saccharomyces cerevisiae*) lines were crossed, and the resulting hybrid offspring was outbred

for 12 generations in order to obtain maximum genetic diversity (Parts et al., 2011; Illingworth   326
et al., 2013). The data consists of sequencing data for 171 individuals, for all 16 chromosomes.   327
There are on average 3271 ancestry informative markers per chromosome (95% CI: [929, 6284]).   328
Local ancestry was certain in the data, as both source taxa were differentially homozygous on each   329
ancestry informative SNP. H_0 was 0.5, reflecting a 50/50 contribution of both strains to the first   330
generation. We used three different recombination rate estimates: firstly, we used the linkage map   331
of Cherry et al. (1997) where the average recombination rate is $1cM/2.7kb$ (1 centimorgan per 2.7   332
kilobases), secondly, we used the average recombination rate of $1cM/2.2kb$ as inferred in Mancera   333
et al. (2008) and lastly, we used the average recombination rate of $1cM/5.8kb$ as inferred for the   334
two-way cross in Illingworth et al. (2013). In the absence of a detailed recombination map, we   335
assume that recombination is constant across the chromosome, ignoring hotspots and coldspots. We   336
assume a large population size ($N = 100,000$), reflecting outbreeding.   337

We find that when using the older recombination rate estimates, we consistently underestimate   338
the age of the hybrids (See Figure 11, panel A). The mean age using the recombination rate from   339
Cherry et al. (1997) corresponds to 9.1 generations (95 % CI across all individuals = [6.2, 11.3])   340
and the mean age using the recombination rate from Mancera et al. (2008) corresponding to 7.0   341
generations (95 % CI: [5.4, 8.6]). These estimates suggest that the true recombination rate is slightly   342
lower than assumed. When using the most recent recombination rate estimate (i.e. $1cM/5.8kb$, from   343
Illingworth et al. (2013)), we slightly overestimate the age (mean age estimate: 18.7 generations,   344
95 % CI: [13.0, 22.5]). Alternatively, we could be overestimating population size, suggesting that   345
perhaps the rate of inbreeding in the experimental design was higher than anticipated. Reducing the   346
population size tends to increase the time since admixture (Supplement S4), which could help bring   347
the age estimates using the two oldest recombination rates closer to the true number of generations.   348
However, for the most recent recombination rate estimate, this would only increase the error made.   349
The likelihood profiles (See Figure 11, panel B) suggest that the maximum likelihood estimates are   350
robust and thus it seems more likely that the recombination rate estimates do not accurately reflect   351
the amount of recombination accumulated in the experiment.   352

4.2   Swordtail Fish   353

Here, we reanalyze data of hybridizing swordtail fish published in (Schumer et al., 2018). Swordtail   354
fish have received considerable attention in the past years, as they have been shown to hybridize   355
readily in nature. We focus here on a hybrid population located in Tlatemaco, Mexico (Schumer   356
et al., 2018, 2014a). The population is the result of a hybridization event between *Xiphophorus*   357
birchmanni and *X. malinche*, approximately 100-200 generations ago (Pers. Comm. M. Schumer and   358
(Schumer et al., 2018; Powell et al., 2020a)). Currently, the hybrid genome consists for 75% of *X.*   359
malinche, suggesting that the initial hybrid swarm was strongly biased towards *X. malinche*, or that   360
strong selection after hybridization has favored genomic material from *X. malinche*. We use ancestry   361
information provided in the data supplement of Schumer et al. (2018), which contains unphased   362
local ancestry estimates based on multiplexed shotgun genotyping (MSG) results (Andolfatto et al.,   363
2011), with on average 38,936 markers per chromosome (95% CI: [22765, 50256]). The MSG pipeline   364
provides a posterior probability of observing local ancestry. Following Schumer et al. (2018), we   365
converted local ancestry probabilities of >95% to hard ancestry calls. To obtain age estimates, we   366
use the estimated population size in Schumer et al. (2014a) of 1830 individuals. We infer the age   367
for 187 individuals from the Tlatemaco population. We use ancestry information from 24 linkage   368
groups, but remove linkage groups 17 and 24, as these are known to include large inversions (Schumer   369
et al., 2018), making them unsuitable for admixture analysis. As a recombination map, we use   370
three approaches. Firstly, we use the average recombination rate of $1cM/378kb$ as used in Schumer   371
et al. (2014a), which is based on the average genome-wide recombination rate in *Xiphophorus*   372
(Walter et al., 2004). Secondly, we use the average recombination rate of $1cM/500kb$ as reported   373
in Powell et al. (2020b). Lastly, we use the high density recombination map reconstructed from   374
Linkage Disequilibrium patterns as presented in Schumer et al. (2018), which represents an average   375
recombination rate of $1cM/485kb$.   376

We find that the distribution of ages inferred for individuals from the Tlatemaco population is   377
overall higher than the previously inferred age but still consistent with those estimates (see Fig 12   378
A). We recover a mean age of 163 generations (95% CI: [81, 201]) when using the recombination rate   379
reported in Schumer et al. (2014b). Using the high density recombination map from Schumer et al.   380

(2018) we obtain a mean age estimate of 210 generations (95% CI: [111, 257]), due to the shorter map   381

length. Alternatively, using the most recent recombination rate estimate of $1cM/500kb$ reported   382

by Powell et al. (2020b), we recover a mean age of 217 generations (95% CI: [115, 265]). Thus, we   383

generally find that the population is perhaps slightly older than expected, but we would like to   384

emphasize that the likelihood profile across all samples (Figure 12 B) is extremely flat, suggesting   385

that the age estimates obtained tend to be uncertain and susceptible to potential inconsistencies in   386

the data (e.g. sequence errors). Exploration of different values for the population size (Supplement   387

S4) shows that if the true population size is larger than reported by Schumer et al. (2014a), this   388

would decrease the time since admixture, but only minimally so (by only a few generations).   389

4.3   *Populus* trees   390

Here, we reanalyze a dataset of *Populus* trees, published by Suarez-Gonzalez et al. (2016). The   391

dataset focuses on two species of trees, *P. trichocarpa*, found mainly in West-America, and *P.*   392

balsamifera, which is found in Northern America. The two species are thought to have diverged   393

relatively recently, around 760k years ago. Where their ranges meet (around the southern tip of   394

Alaska), the two species hybridize, and a hybrid population has been established. The dataset   395

consists of 32 individuals which are mainly *P. balsamifera*, admixed with *P. trichocarpa* and 36   396

individuals that are mainly *P. trichocarpa*, admixed with *P. balsamifera*. Three chromosomes of   397

interest (chromosomes 6, 12 and 15) were Illumina sequenced, and unphased data was available   398

for on average 60071 ancestry informative markers per chromosome (95% CI: [28745, 101425]).   399

Ancestry information in the markers was very high, with on average 40852 markers with an allele   400

frequency differential of at least 0.7 (Shriver et al., 1997), and 21009 markers with an allele frequency   401

differential of at least 0.9. We use three different population level recombination rates recovered from   402

the literature, being $\rho = 0.00219$ (Wang et al., 2016), $\rho = 0.0092$ (Olson et al., 2010) and $\rho = 0.0197$ 403

(Slavov et al., 2012). We converted these population level recombination rates to individual rates   404

using an effective population size of 5106 individuals, as estimated using phylogenetic methods in   405

(Slavov et al., 2012). This yielded three local recombination rates of $1cM/10.4kb$ (Slavov et al.,   406

2012), $1cM/22.2kb$ (Olson et al., 2010) and $1cM/93.3kb$ (Wang et al., 2016). Local ancestry was   407

determined using ANCESTRY HMM (Corbett-Detig and Nielsen, 2017), assuming equal admixture of   408
both source taxa. Because admixture differed strongly across samples, we used the average local   409
ancestry per sample as input for a second run of ANCESTRY HMM in order to obtain accurate local   410
ancestry calls. We converted local ancestry probabilities of >95% to hard ancestry calls. Lastly, we   411
compared the observed variation in ancestry across samples with the expected variation in ancestry   412
for a single panmictic population, as given by equation 8 in (Gravel, 2012).   413

We find that the time since admixture strongly correlates with the recombination rate used   414
(See Figure 13 A), with a mean number of generations since admixture of 6 (95% CI: [3, 13]) when   415
using the highest estimate of recombination ($1cM/10.4kb$ (Slavov et al., 2012)), an intermediate   416
estimate of 12 generations (95% CI: [7, 28]) when using a recombination rate of $1cM/22.2kb$ (Olson   417
et al., 2010) and a much higher age estimate of 50 generations (95% CI: [30, 114]) when using   418
the lowest recombination estimate of $1cM/93.3kb$ (Wang et al., 2016). The likelihood profiles (See   419
Figure 13 B) intersect on multiple occasions, suggesting multiple optima. We measure a variation in   420
ancestry across the three chromosomes of 0.042. However, we obtain expected levels of variation of   421
ancestry of 0.0088, 0.0010 and 0.0008 for the three different recombinatinon rate estimates (and their   422
corresponding age estimates). Observed variation in the data is thus much higher than expected   423
from admixture in a single panmictic population.   424

5 Discussion   425

The aim of this article was to improve the estimation of the time since admixture in hybrid   426
populations. To do so, we have extended the theory of junctions in two directions. First, we have   427
derived a formula for the expected number of observed junctions in one chromosome that takes into   428
account the number of markers and their positions (equation (3)). Second, we have considered the   429
case in which there is sequencing data from two homologous chromosomes. We have developed a   430
maximum likelihood approach that infers the time since admixture, for both phased and unphased   431
data.   432

Firstly, we have used simulations to validate the accuracy of our method. Results from our   433
simulations show that our method for a single chromosome performs better than previous methods   434

that ignore the effect of having a limited number of markers or assume that the markers are even-spaced (see Fig 5). Furthermore, we find that using information from two chromosomes improves accuracy considerably, as expected. This effect is even stronger for small population sizes (see Fig 6 and Fig 2 in S2 Appendix).

Surprisingly, the method based on unphased data performs similarly to our method based on phased data (see Fig 7 and Fig 3 in S2 Appendix). The phased and unphased approaches differ in their treatment of markers that are heterozygous for ancestry, and hence we expected differences between these methods to manifest themselves primarily during the initial stages of admixture, when heterozygosity is still high. We did find that there were slight differences during these stages (Fig 7 and Fig 3 in S2 Appendix), but these were neglibile compared to the overall uncertainty. Furthermore, these errors were much smaller than errors accumulated due to incorrect phasing (see Fig 8). Our findings here are conservative, as we show that the unphased method performs better even for small error rates, comparable to error rates for human data (for example in (Choi et al., 2018)). Human data sets are typically of very high quality, and these error rates represent an extremely favourable scenario. Thus, given the impact of error rates incurred during phasing, we recommend using our unphased framework if possible, to obtain more accurate time estimates.

Apart from sensitivity to phasing error, we have tested the sensitivity of our method to different parameters such as the number of markers n, the population size N, the initial heterozygosity H_0 and the total recombination rate C (see S1 Appendix). Our method seems to be quite sensitive to H_0 but this parameter can easily be estimated from the proportion of markers that come from each source taxa. One advantage of our approach is that age inference is not very sensitive to population size (see Fig 1 in S1 Appendix), which was not true for previous methods that rely on a good estimation of N (see Janzen et al. (2018)). Our method is not very sensitive either to the number of markers (see Fig 4 in S1 Appendix), provided that it is above a certain threshold. Janzen et al. (2018) inferred that when using regularly spaced markers and information for a single chromosome, the number of markers typically needs to be an order of magnitude larger than $\frac{1}{2}Ct$, where t is the admixture time and C the total amount of recombination. We find similar results when using information from a single chromosome with arbitrarily spaced markers or information from both

chromosomes (see S1 Appendix). When analyzing empirical data, it is often impossible to know *a* 463
priori whether the number of ancestry informative markers is much larger than the admixture time. 464
However, our simulation results indicate that when the number of markers is too small, variation 465
in the age estimate across different chromosomes tends to increase. Thus, large variation in the 466
estimate of admixture time, or inferred admixture times that tend to extremely large values, are 467
potentially indicative of an insufficient marker number. 468

The main issue with our method is its sensitivity to the recombination rate. This is shown in Fig 469
2 of S1 Appendix but also exemplified by the varying results in the empirical datasets, dependent on 470
our assumptions about recombination rates. However, it should be noted that this issue is not novel 471
to our approach, but is a general issue with the theory of junctions. Apart from sensitivity to the 472
average recombination rate, local hot-spots or cold-spots of recombination could potentially also 473
influence admixture time estimates. 474

Furthermore, the recombination rate factors into inference of local ancestry. Our methodology 475
assumes local ancestry to be known, and relies on upstream inference of local ancestry. However, 476
methods to infer local ancestry such as MSG (Andolfatto et al., 2011), ELAI (Guan, 2014) and 477
ANCESTRY HMM (Corbett-Detig and Nielsen, 2017) also use recombination in their calculations to 478
infer local ancestry. Thus, by using these local ancestry estimates, our methodology re-uses the 479
recombination rate (both for local ancestry inference, and subsequent age estimation). Any errors in 480
the recombination rate estimate might then propagate through the pipeline and affect age estimates 481
using our framework. Using simulations including ancestry uncertainty, we have looked into this 482
effect of stacking errors and have found that our method performs similarly to ANCESTRY HMM, 483
even if we use the local ancestry inferred by ANCESTRY HMM. Furthermore, our method outperforms 484
ANCESTRY HMM for long admixture times or for small population sizes. ANCESTRY HMM assumes 485
$T \ll \log_2(N)$ (this is the same assumption used in SMC' (Liang and Nielsen, 2014)), whilst our 486
framework does not need to make these assumptions to obtain age estimates. However, when marker 487
numbers dwindle (Supplement S5), our framework can no longer correct for this approximation, 488
and performance of our framework drops below inference using only ANCESTRY HMM. Yet, in the 489
empirical datasets we have analyzed, we have generally found marker densities for which these limits 490

were not reached and with full genome sequencing now within reach for many researchers, we expect 491
this not to be an issue. 492

To validate our approach we have re-analysed three datasets. The first dataset is from a crossing 493
experiment with *S. cerevisiae*. Here, we applied the single chromosome equations, and estimates of 494
the time since the onset of admixture line up well with the experimental design, although assumptions 495
regarding the recombination rate remain of strong influence on the admixture time estimates. 496

The second dataset we reanalyzed is of Swordtail fish (*Xiphophorus*). We infer an admixture 497
time that is older than previous estimates (Schumer et al., 2014b) but that is in line with more 498
recent estimates done by the same authors (M. Schumer, personal communication, (Powell et al., 499
2020a)) using more recent recombination rate estimates (Powell et al., 2020b). However, likelihood 500
profiles are fairly flat, indicating that our maximum likelihood estimates are sensitive to changes 501
in the data, further reflected by the sensitivity of the results to assumptions made regarding the 502
recombination rate. 503

Finally, we have reanalyzed a dataset on *Populus* trees (Suarez-Gonzalez et al., 2016). We 504
infer an admixture time that is in line with previous findings, but the original analysis did not 505
focus on admixture time and only used admixture time to infer local ancestry. However, we find 506
that the time since admixture correlates strongly with the genetic distance to either of the source 507
taxa, with individuals more closely related to the source taxa inferred to be younger. This suggests 508
that the dataset does not consist of a sample from a single, admixed, population and that this 509
invalidates our analysis. Furthermore, variation in ancestry is much higher than expected from the 510
admixture time alone (Gravel, 2012), again indicating that the analyzed population is most likely 511
not a single admixing population. Lastly, admixture mapping analyses have shown that perhaps 512
late generation backcrosses have contributed as well to the hybrid population (Suarez-Gonzalez 513
et al., 2018), suggesting an intermediate form between on the one hand adaptive introgression & 514
back-crossing and on the other hand ongoing hybridization across a spatial gradient. Across these 515
results, it is clear that our assumption of a single admixed population is violated, and hence it seems 516
likely that our age estimates are incorrect and that our framework is not a good fit for this empirical 517
dataset. Yet, we believe that it provides an interesting example on how our methodology can be 518

applied. ⁵¹⁹ 519

Summarizing, we have presented a framework to estimate the time since admixture using phased 520 or unphased data from two homologous chromosomes, taking into account marker spacing along the 521 chromosome. We have shown that using data from two chromosomes improves the estimations of the 522 admixture time compared to the method that uses only one chromosome. This is true whether the 523 data is phased or unphased. In addition we have shown, using simulations, that applying the phased 524 or the unphased method yields very similar results. However, given that even small (unavoidable) 525 phasing errors produce overestimates in the time since admixture, we suggest that, in most cases, 526 using unphased data is the best strategy. In comparison with previous methods, such as ANCESTRY 527 HMM (Corbett-Detig and Nielsen, 2017), our method performs better when the population size is 528 small or the time since admixture is long. With our new framework, we hope to have opened new 529 avenues towards inferring the time since admixture in admixed populations, and primarily hope 530 to have brought this analysis within reach also for datasets where phased data is unavailable or 531 impossible to acquire. Furthermore, we would like to emphasize that our method also works for a 532 relatively small number of SNPs, which opens up avenues towards analysis of closely related taxa, 533 where the number of ancestry informative markers might be low. We have included the derivations 534 and the numerical solution framework in the R package 'junctions' (Janzen, 2021). By providing the 535 code in an easy to use package, we hope to lower the threshold for other users to apply the theory of 536 junctions to their model system. 537

Acknowledgements 538

We thank Gianni Litti, Molly Schumer, Adriana Suarez-Gonzalez and Quentin Cronk for their help 539 and enthusiasm with respect to our re-analysis of their datasets. We also thank Amaury Lambert 540 who helped building this fruitful collaboration and Arno Siri-Jégousse for a careful reading of a 541 previous version of this manuscript. TJ thanks the Carl von Ossietzky University of Oldenburg for 542 use of their computer cluster CARL and the Center for Information Technology of the University of 543 Groningen for their support and access to the Peregrine high performance computing cluster. VMP 544 was funded by the DGAPA-UNAM postdoctoral program. 545

Data Accessibility

We have included the derivations and the numerical solution framework in the R package 'junctions', which can be found on CRAN on `https://CRAN.R-project.org/package=junctions`. All code used in data analysis and visualization for this manuscript can be downloaded from data dryad, from: `https://datadryad.org/stash/share/GCJGziFXtI6CV42rl3mcwa92-1nNNMND3PH94QxZL-k` (please note that upon publication, this link will be replaced with `doi:10.5061/dryad.xwdbrv1c5`)

Author contributions

TJ and VMP jointly designed the research. VMP inferred the ARG based mathematics, TJ verified findings using individual based simulations and analyzed the empirical data. TJ and VMP jointly wrote the paper.

References

J. A. Coyne and H. A. Orr. *Speciation*. Sinauer Associates, Inc, 2004.

R. Abbott, D. Albach, S. Ansell, J. W. Arntzen, et al. Hybridization and speciation. *Journal of Evolutionary Biology*, 26:229–246, 2013.

V. Grant. *Plant speciation*. Columbia University Press, 1981.

J. Mavárez, C. A. Salazar, E. Bermingham, C. Salcedo, et al. Speciation by hybridization in Heliconius butterflies. *Nature*, 441(7095):868–871, 2006.

T. Capblancq, L. Després, D. Rioux, and J. Mavárez. Hybridization promotes speciation in coenonympha butterflies. *Molecular Ecology*, 24(24):6209–6222, 2015.

S. Koblmüller, N. Duftner, K.M. Sefc, M. Aibara, et al. Reticulate phylogeny of gastropod-shell-breeding cichlids from lake tanganyika–the result of repeated introgressive hybridization. *BMC Evolutionary Biology*, 7(1):7, 2007.

I. Keller, C.E. Wagner, L. Greuter, S. Mwaiko, et al. Population genomic signatures of divergent adaptation, gene flow and hybrid speciation in the rapid radiation of lake Victoria cichlid fishes. *Molecular Ecology*, 22(11):2848–2863, 2013.   568
569
570

A. Brelsford, Mila B., and D.E. Irwin. Hybrid origin of Audubon's warbler. *Molecular Ecology*, 20: 2380–2389, 2011.   571
572

D. Schwarz, B. M. Matta, N.L. Shakir-Botteri, and B. A. McPheron. Host shift to an invasive plant triggers rapid animal hybrid speciation. *Nature*, 436(7050):546–549, 2005.   573
574

A.W. Nolte, J. Freyhof, K.C. Stemshorn, and D. Tautz. An invasive lineage of sculpins, Cottus sp. (Pisces, Teleostei) in the rhine with new habitat adaptations has originated from hybridization between old phylogeographic groups. *Proceedings of the Royal Society B*, 272:2379–2387, Oct 2005.   575
576
577

S.J.E. Baird. Phylogenetics - fisher's markers of admixture. *Heredity*, 97(2):81–83, 2006.   578

S.J.E. Baird. Exploring linkage disequilibrium. *Molecular ecology resources*, 15(5):1017–1019, 2015.   579

R. A. Fisher. *The Theory of Inbreeding*. Oliver and Boyd, 1949.   580

R. A. Fisher. A fuller theory of "junctions" in inbreeding. *Heredity*, 8:187–197, 1954.   581

J.H. Bennett. Junctions in inbreeding. *Genetica*, 26(1):392–406, 1953.   582

R. A. Fisher. An algebraically exact examination of junction formation and transmission in parent-offspring inbreeding. *Heredity*, 13:179–186, 1959.   583
584

J. Gale. Some applications of the theory of junctions. *Biometrics*, pages 85–117, 1964.   585

P. Stam. The distribution of the fraction of the genome identical by descent in finite random mating populations. *Genetical Research*, 35(2):131–155, 1980.   586
587

SJE Baird, NH Barton, and AM Etheridge. The distribution of surviving blocks of an ancestral genome. *Theoretical Population Biology*, 64(4):451–471, 2003.   588
589

N. H. Barton. Multilocus clines. *Evolution*, 37(3):454–471, 1983.   590

S.J.E. Baird. A simulation study of multilocus clines. *Evolution*, 49(6):1038–1045, 1995.

N.H. Chapman and E.A. Thompson. The effect of population history on the lengths of ancestral chromosome segments. *Genetics*, 162(1):449–458, 2002.

N.H. Chapman and E.A. Thompson. A model for the length of tracts of identity by descent in finite random mating populations. *Theoretical Population Biology*, 64(2):141–150, 2003.

P. Muir, S. Li, S.e Lou, D. Wang, et al. The real cost of sequencing: scaling computation to keep pace with data generation. *Genome Biology*, 17(1):53, 2016.

B. Paşaniuc, S. Sankararaman, G. Kimmel, and E. Halperin. Inference of locus-specific ancestry in closely related populations. *Bioinformatics*, 25(12):i213–i221, 2009.

B.K. Maples, S. Gravel, E. E. Kenny, and C. D. Bustamante. Rfmix: a discriminative modeling approach for rapid and robust local-ancestry inference. *The American Journal of Human Genetics*, 93(2):278–288, 2013.

Yongtao Guan. Detecting structure of haplotypes and local ancestry. *Genetics*, 196(3):625–642, 2014.

R Corbett-Detig and R. Nielsen. A hidden Markov model approach for simultaneously estimating local ancestry and admixture time using next generation sequence data in samples of arbitrary ploidy. *PLoS Genetics*, 13(1):e1006529, 2017.

Paloma Medina, Bryan Thornlow, Rasmus Nielsen, and Russell Corbett-Detig. Estimating the timing of multiple admixture pulses during local ancestry inference. *Genetics*, 210(3):1089–1107, 2018.

Jesper Svedberg, Vladimir Shchur, Solomon Reinman, Rasmus Nielsen, and Russell Corbett-Detig. Inferring adaptive introgression using hidden markov models. *Molecular biology and evolution*, 38 (5):2152–2165, 2021.

A.K. MacLeod, C.S. Haley, and P. Woolliams, J.A .and Stam. Marker densities and the mapping of ancestral junctions. *Genetical research*, 85(01):69–79, 2005.

C. A. Buerkle and L.H. Rieseberg. The rate of genome stabilization in homoploid hybrid species. 616
Evolution, 62(2):266–275, 2008. 617

T. Janzen, A. Nolte, and A. Traulsen. The breakdown of genomic ancestry blocks in hybrid lineages 618
given a finite number of recombination sites. *Evolution*, 72(4):735–750, 2018. 619

M.W. Snyder, A. Adey, J.O. Kitzman, and J. Shendure. Haplotype-resolved genome sequencing: 620
Experimental methods and applications. *Nature Reviews Genetics*, 16(6):344–358, 5 2015. ISSN 621
1471-0056. doi: 10.1038/nrg3903. 622

D. Lutgen, R. Ritter, R.-A. Olsen, H. Schielzeth, et al. Linked-read sequencing enables haplotype- 623
resolved resequencing at population scale. *bioRxiv*, 2020. doi: 10.1101/2020.01.15.907261. URL 624
`https://www.biorxiv.org/content/early/2020/01/15/2020.01.15.907261`. 625

S.P. Browning and B.L. Browning. Haplotype phasing: existing methods and new developments. 626
Nature Reviews Genetics, 12:703–714, 2011. 627

P.R. Loh, P.F. Palamara, and L.P. Alkes. Fast and accurate long-range phasing in a uk biobank 628
cohort. *Nature Genetics*, 48:811–816, 2016a. 629

A. Kong, G. Masson, M. L. Frigge, A. Gylfason, et al. Detection of sharing by descent, long-range 630
phasing and haplotype imputation. *Nature Genetics*, 40(9):1068–1075, 2008. 631

P.R. Loh, P.F. Palamara, and A.L. Price. Fast and accurate long-range phasing in a uk biobank 632
cohort. *Nature Genetics*, pages 811–816, 2016b. 633

S.R. Browning and B.L. Browning. Rapid and accurate haplotype phasing and missing-data inference 634
for whole-genome association studies by use of localized haplotype clustering. *American journal* 635
of human genetics, 81(5):1084–97, 2007. 636

J. O'Connell, K. Sharp, N. Shrine, L. Wain, I. Hall, M. Tobin, J.F. Zagury, O. Delaneau, and 637
J. Marchini. Haplotype estimation for Biobank-scale data sets. *Nature Genetics*, 48:817–820, 2016. 638

J. A Baaijens and A. Schönhuth. Overlap graph-based generation of haplotigs for diploids and poly- 639
ploids. *Bioinformatics*, 35(21):4281–4289, 04 2019. ISSN 1367-4803. doi: 10.1093/bioinformatics/ 640
btz255. URL `https://doi.org/10.1093/bioinformatics/btz255`. 641

R.W. Tourdot and C.-Z. Zhang. Whole chromosome haplotype phasing from long-range sequencing. 642
bioRxiv, 2019. doi: 10.1101/629337. URL `https://www.biorxiv.org/content/early/2019/05/` 643
`07/629337`. 644

Z.N. Kronenberg, A. Rhie, S. Koren, G. .T. Concepcion, and others. Extended haplotype phasing 645
of de novo genome assemblies with falcon-phase. *bioRxiv*, 2019. doi: 10.1101/327064. URL 646
`https://www.biorxiv.org/content/early/2019/04/19/327064`. 647

J. Ebler, M. Haukness, T. Pesout, T. Marschall, and B. Paten. Haplotype-aware diplotyping from 648
noisy long reads. *Genome Biol.*, 116(20), 2019. 649

A. Tangherloni, S. Spolaor, L. Rundo, M. S. Nobile, et al. Genhap: a novel computational method 650
based on genetic algorithms for haplotype assembly. *BMC Bioinformatics*, 172(20), 2019. 651

R.R. Hudson. Properties of the neutral model with intragenic recombination. *Theor. Popul. Biol.*, 652
23(2):213–201, 1983. 653

R. C. Griffiths. The two-locus ancestral graph. In I.V. Basawa and R. L. Taylor, editors, *Selected* 654
Proceeedings of the Symposium on Applied Probability, pages 100–117. Institute of Mathematical 655
Statistics, 1991. 656

R.C. Griffiths and P. Marjoram. An ancestral recombination graph. In P. Donnelly and S. Tavaré, 657
editors, *Progress in Population Genetics and Human Evolution, IMA Volumes in Mathematics* 658
and its Applications, volume 87, pages 257–270. Springer Verlag, 1997. 659

R. Durrett. *Probability Models for DNA Sequence Evolution*. Springer, 2 edition, 2008. 660

G.A. McVean and N.J. Cardin. Approximating the coalescent with recombination. *Philos Trans R* 661
Soc Lond B Biol Sci, 1459(360):1387–93, 2005. 662

Y. Choi, A. P. Chan, E. Kirkness, A. Telenti, and N. J. Schork. Comparison of phasing strategies for 663
whole human genomes. *PLOS Genetics*, 14(4):1–26, 04 2018. doi: 10.1371/journal.pgen.1007308. 664
URL https://doi.org/10.1371/journal.pgen.1007308. 665

Mark D Shriver, Michael W Smith, Lin Jin, Amy Marcini, Joshua M Akey, Ranjan Deka, and 666
Robert E Ferrell. Ethnic-affiliation estimation by use of population-specific dna markers. *American* 667
journal of human genetics, 60(4):957, 1997. 668

M. Liang and R. Nielsen. The lengths of admixture tracts. *Genetics*, 197(3):953–967, 2014. 669

L. Parts, F.A. Cubillos, J. Warringer, K. Jain, et al. Revealing the genetic structure of a trait 670
by sequencing a population under selection. *Genome Research*, 21(7):1131–1138, 2011. doi: 671
10.1101/gr.116731.110. URL http://genome.cshlp.org/content/21/7/1131.abstract. 672

C.J.R. Illingworth, L. Parts, A. Bergström, G. Liti, and V. Mustonen. Inferring genome-wide 673
recombination landscapes from advanced intercross lines: application to yeast crosses. *PLoS One*, 674
8(5):e62266, 2013. 675

J.M. Cherry, C. Ball, S. Weng, G. Juvik, et al. Genetic and physical maps of Saccharomyces 676
cerevisiae. *Nature*, 387(6632 Suppl):67, 1997. 677

E. Mancera, R. Bourgon, A. Brozzi, W. Huber, and L.M. Steinmetz. High-resolution mapping of 678
meiotic crossovers and non-crossovers in yeast. *Nature*, 454(7203):479–485, 2008. 679

M. Schumer, C. Xu, D. L. Powell, A. Durvasula, et al. Natural selection interacts with recombination 680
to shape the evolution of hybrid genomes. *Science*, 360(6389):656–660, 2018. 681

M. Schumer, R. Cui, D. L. Powell, R. Dresner, et al. High-resolution mapping reveals hundreds of 682
genetic incompatibilities in hybridizing fish species. *Elife*, 3:e02535, 2014a. 683

Daniel L. Powell, Ben Moran, Bernard Kim, Shreya M. Banerjee, Stepfanie M. Aguillon, Paola 684
Fascinetto-Zago, Quinn Langdon, and Molly Schumer. Two new hybrid zones expand the 685
swordtail hybridization model system. *bioRxiv*, 2020a. doi: 10.1101/2020.11.18.389205. URL 686
https://www.biorxiv.org/content/early/2020/11/20/2020.11.18.389205. 687

P. Andolfatto, D. Davison, D. Erezyilmaz, T. T. Hu, et al. Multiplexed shotgun genotyping for rapid and efficient genetic mapping. *Genome research*, 21(4):610–617, 2011.

R.B. Walter, J.D. Rains, J.E. Russell, T.M Guerra, et al. A microsatellite genetic linkage map for xiphophorus. *Genetics*, 168(1):363–372, 2004.

D. L. Powell, M. García-Olazábal, M. Keegan, P. Reilly, et al. Natural hybridization reveals incompatible alleles that cause melanoma in swordtail fish. *Science*, 368(6492):731–736, 2020b.

M. Schumer, G.G. Rosenthal, and P. Andolfatto. How common is homoploid hybrid speciation? *Evolution*, 68(6):1553–1560, 2014b.

A. Suarez-Gonzalez, C.A. Hefer, C. Christe, O. Corea, et al. Genomic and functional approaches reveal a case of adaptive introgression from Populus balsamifera (balsam poplar) in P. trichocarpa (black cottonwood). *Molecular ecology*, 25(11):2427–2442, 2016.

J. Wang, N.R. Street, D.G. Scofield, and P.K. Ingvarsson. Natural selection and recombination rate variation shape nucleotide polymorphism across the genomes of three related populus species. *Genetics*, 202(3):1185–1200, 2016.

M.S. Olson, A.L. Robertson, N. Takebayashi, S. Silim, et al. Nucleotide diversity and linkage disequilibrium in balsam poplar (Populus balsamifera). *New Phytologist*, 186(2):526–536, 2010.

G.T. Slavov, S.P. Difazio, J. Martin, W. Schackwitz, et al. Genome resequencing reveals multiscale geographic structure and extensive linkage disequilibrium in the forest tree populus trichocarpa. *New Phytologist*, 196(3):713–725, 2012.

Simon Gravel. Population genetics models of local ancestry. *Genetics*, 191(2):607–619, 2012.

A. Suarez-Gonzalez, C.A. Hefer, C. Lexer, C.J. Douglas, and Q.C.B. Cronk. Introgression from Populus balsamifera underlies adaptively significant variation and range boundaries in P. trichocarpa. *New Phytologist*, 217(1):416–427, 2018.

Thijs Janzen. *junctions: The Breakdown of Genomic Ancestry Blocks in Hybrid Lineages*, 2021. URL https//github.com/thijsjanzen/junctions. R package version 2.0.0.

Supporting information 713

S1 Appendix.   **Sensitivity analysis.** Using individual based simulations, we test how sensitive 714
our new framework is to variation of the different parameters. 715

S2 Appendix.   **Small population size.** We test the validity of our method using a smaller value 716
of the population size ($N = 1000$). 717

S3 Appendix.   **Phasing error.** We extend the analysis done in the main text to the case of a 718
smaller number of markers. 719

S4 Appendix.   **Population size effect on age estimates in empirical data.** We extend the 720
analyses done in the main text for the yeast and swordtail datasets by exploring the impact of 721
population size on the age estimates. 722

S5 Appendix.   **Comparison with ancestry hmm for a larger population size.** We repeat 723
the analysis for Figure 10, but for a population size of $N = 10,000$. 724

Tables and Figures 725

Fig 1. Visual depiction of the observed data. We show the differences between the type of data generated by the three methods we present in this paper. On each panel, the chromosome in the center is colored according to ancestry (blue represents source taxa $\mathcal{P}$ and red represents source taxa $\mathcal{Q}$). Above the chromosome are indicated the locations of ancestry informative markers z_i. Resulting inferred ancestry on these markers is shown below, where grey indicates heterozygous ancestry. The first panel represents the one chromosome method. There are 7 junctions in the chromosome, but only 3 are observed in the data due to a limited marker coverage. Notice that a junction in the inferred ancestry corresponds to an odd number of junctions in the "true" genome ancestry and conversely, that an even number of junctions in the "true" ancestry yields no inferred junctions. The second and third panels represent the methods that use information from two chromosomes. In the second panel data is phased whereas in the second panel data is unphased.

Fig 2. The ARG with two markers. Each color represents one source taxa ($\mathcal{P}$ and $\mathcal{Q}$). The black and grey lines (or dotted lines) represent the ancestral lineage of each marker. In the left panel, we show the ARG for two markers in one chromosome. In the present, there is an observed junction between the two markers. In the past (t generations ago, when hybridization took place), each lineage is carried by a different individual and these two individuals are from different source taxa. The right panel shows the ARG for two markers in two homologous chromosomes.

Fig 3. $\mathbb{P}_t(O_i, O_{i+1})$ **for phased data.** The allele from source taxa $\mathcal{P}$ is represented in blue and the allele from source taxa $\mathcal{Q}$ is represented in red.

	$O_{i+1} = P$	$O_{i+1} = Q$	$O_{i+1} = x$
$O_i = P$	$p^2((P_t^i)_1 + (P_t^i)_4 + (P_t^i)_7) + p^3((P_t^i)_2 + (P_t^i)_5) + p^4(P_t^i)_3 + p(P_t^i)_6$	$pq(pq(P_t^i)_3 + \frac{1}{2}(P_t^i)_5 + (P_t^i)_7)$	$pq(p(P_t^i)_2 + 2p^2(P_t^i)_3 + \frac{1}{2}(P_t^i)_4 + p(P_t^i)_5)$
$O_i = Q$	$pq(pq(P_t^i)_3 + \frac{1}{2}(P_t^i)_5 + (P_t^i)_7)$	$q^2((P_t^i)_1 + (P_t^i)_4 + (P_t^i)_7) + q^3((P_t^i)_2 + (P_t^i)_5) + q^4(P_t^i)_3 + q(P_t^i)_6$	$pq(q(P_t^i)_2 + 2q^2(P_t^i)_3 + \frac{1}{2}(P_t^i)_4 + q(P_t^i)_5)$
$O_i = x$	$pq(p(P_t^i)_2 + 2p^2(P_t^i)_3 + \frac{1}{2}(P_t^i)_4 + p(P_t^i)_5)$	$pq(q(P_t^i)_2 + 2q^2(P_t^i)_3 + \frac{1}{2}(P_t^i)_4 + q(P_t^i)_5)$	or $pq(2(P_t^i)_1 + (P_t^i)_2 + 4pq(P_t^i)_3)$

Fig 4. $\mathbb{P}_t(O_i, O_{i+1})$ **for unphased data.** The allele from source taxa $\mathcal{P}$ is represented in blue and the allele from source taxa $\mathcal{Q}$ is represented in red.

Fig 5. Comparison to previous methods. Shown are the median estimates for the time since admixture (dots) for 100 replicates, where in each replicate 10 individuals were analyzed. The dashed line indicates the simulated time. 'Evenly spaced markers' corresponds to the method in (Janzen et al., 2018). 'Infinite markers' corresponds to an idealized scenario where ancestry is known for every locus in the chromosome and is there to quantify the amount of randomness in the process. The population size was 10,000 individuals, and 10,000 randomly spaced markers were used.

Fig 6. Accuracy in age estimate using information from one versus two chromosomes.
Inferred time versus simulated time is represented. Shown are the median estimates (dots) for 100
replicates, where in each replicate 10 individuals were analyzed. The solid white line indicates the
observed is equal to expected line and the shaded area indicates the 95% percentile. Shown are
results using junction information from one chromosome (blue) and results using phased information
from two chromosomes (gold). Numbers above the plots indicate the initial heterozygosity. The
population size was 10,000 individuals, and 10,000 randomly spaced markers were used.

Fig 7. Accuracy in age estimate using the unphased framework versus the phased framework. Shown are the median difference across 100 replicates. We represent the results for three different initial heterozygozities, as indicated at the top of each plot. The population size was 10,000 individuals, and 10,000 randomly spaced markers were used. The inset plots show the same results, including the 95% confidence limits, which are far outside the boundaries of the main plot.

Fig 8. Effect of switching error on the estimated time since admixture. Data simulated with $N = 10,000$, $p = 0.5$, $C = 1$ and $n = 10,000$. The solid black line indicates the simulated = estimated time. Dots indicate the median inferred age and the colored area indicates the 95% confidence interval (CI) envelope. Colors reflect different degrees of phasing error, where a phasing error of 0.01 represents a 1% probability of a SNP being phased incorrectly.

Fig 9. Comparison in estimating the time since admixture between ancestry hmm and the method proposed here. The solid black line indicates the simulated = estimated time. Dots indicate the inferred ages, with the green dots representing ages inferred by ANCESTRY HMM and the blue dots indicate ages inferred by the junctions framework. Age estimates are based on simulated data with known ancestry, using $n = 40,000$, $C = 1$, $H_0 = 0.5$.

Fig 10. Comparison in estimating the time since admixture between ancestry hmm and the method proposed here, for a small population of $N = 1000$**.** The solid black line indicates the simulated = estimated time. Dots indicate the inferred ages, with the green dots representing ages inferred by ANCESTRY HMM and the blue dots indicate ages inferred by the junctions framework. Missing dots are caused by a lack of convergence of the maximum likelihood algorithm. Age estimates are based on simulated data with uncertain ancestry, where uncertainty in ancestry is reflected by the allele frequency differential (Shriver et al., 1997). Because the method proposed here does not include ancestry uncertainy, local ancestry as inferred by ANCESTRY HMM was used.

Fig 11. Inferred age for F12 Hybrid Yeast (*Saccharomyces cerevisiae*) individuals. (A)
Inferred age for three different recombination rates: $1cM/2.7kb$ (Cherry et al., 1997), $1cM/2.2lb$
(Mancera et al., 2008) and $1cM/5.8kb$ (Illingworth et al., 2013). Shown is the distribution of
inferred ages across 171 individuals. (B) Loglikelihood profile across all chromosomes, where each
line represents one individual. Shown is the loglikelihood profile for the most recent recombination
rate estimate, from Illingworth et al. (2013). The vertical dotted line indicates the 12 generations
line (e.g. the true age of the F12 Hybrid individuals).

Fig 12. Inferred age for hybrid *Swordtail* fish (A) Inferred age for hybrid *Xiphophorus* fish from Tlatemaco (Mexico). Shown are age inferences based on three different recombination maps: $1cM/370kb$ (Schumer et al., 2014a), $1cM/485kb$ (Schumer et al., 2018) and $1cM/500kb$ (Powell et al., 2020b). Dotted lines indicate the hypothesized age limits of the admixed population. (B) Loglikelihood profile across all chromosomes, where each line represents one individual. Shown is the loglikelihood profile for the most recent recombination rate estimate (Powell et al., 2020b).

Fig 13. Inferred age for hybrid *Populus* trees. (A) Distribution of inferred time since
admixture for all individuals. Colors indicate different recombination rates used: $1cM/10.4kb$
(Slavov et al., 2012), $1cM/22.2kb$ (Olson et al., 2010) and $1cM/93.3kb$ (Wang et al., 2016). (B)
Inferred time since admixture, split out across the average frequency of *P. trichocarpa* in the
admixed individual. (C) Loglikelihood profile across all chromosomes, where each line represents
one individual. Shown is the loglikelihood profile for the most recent recombination rate drestimate
(Wang et al., 2016).