1 **Title:** Induction of AmpC-mediated β-lactam resistance requires a single lytic transglycosylase
2 in *Agrobacterium tumefaciens*
3

Authors:

Wanda M. Figueroa-Cuilan[1*], Matthew Howell[1&], Christopher Richards[1], Amelia Randich[1^],

Akhilesh K. Yadav[2$], Felipe Cava[2], and Pamela J.B. Brown[1#]

Affiliations:

[1]Division of Biological Sciences, University of Missouri-Columbia, Columbia, Missouri, 65202

[2]Laboratory for Molecular Infection Medicine, Department of Molecular Biology, Umeå University,

Umeå, Sweden

Running title: AmpC-mediated β-lactam resistance of *A. tumefaciens*

[#]**Corresponding author:**

brownpb@missouri.edu

(573) 884-0384

Present addresses:

*Department of Biological Chemistry, Johns Hopkins University School of Medicine,

Baltimore, Maryland 21205

&Department of Biology, Westminster College, Fulton, Missouri 65251

$Analytical Chemistry Division, CSIR-Indian Institute of Toxicology Research, Lucknow-226001, India

^Department of Biology, The University of Scranton, Scranton, Pennsylvania,18510

ABSTRACT

The remarkable ability of *Agrobacterium tumefaciens* to transfer DNA to plant cells has

allowed the generation of important transgenic crops. One challenge of *A. tumefaciens*-

mediated transformation is eliminating the bacteria after plant transformation to prevent

detrimental effects to plants and the release of engineered bacteria to the environment. Here

we use a reverse genetic approach to identify genes involved in ampicillin resistance with the

goal of utilizing these antibiotic-sensitive strains for plant transformations. We show that

treating *A. tumefaciens* C58 with ampicillin led to increased β-lactamase production, a

response dependent on the broad-spectrum β-lactamase AmpC and its transcription factor

AmpR. Loss of the putative *ampD* orthologue, *atu2113,* led to constitutive production of

AmpC-dependent β-lactamase activity and ampicillin resistance. Finally, one cell wall

remodeling enzyme, MltB3, was necessary for the AmpC-dependent β-lactamase activity and

its loss elicited ampicillin and carbenicillin sensitivity in the *A. tumefaciens* C58 and GV3101

strains. Furthermore, GV3101 *ΔmltB3* transforms plants with comparable efficiency to

wildtype but can be cleared with sub-lethal concentrations of ampicillin. The functional

characterization of the genes involved in the inducible ampicillin resistance pathway of *A.*

tumefaciens constitutes a major step forward in efforts to reduce the intrinsic antibiotic

resistance of this bacterium.

IMPORTANCE

Agrobacterium tumefaciens, a significant biotechnological tool for transgenic plant lines, is

highly resistant to a wide variety of antibiotics, posing challenges for various applications. One

challenge is the efficient elimination of *A. tumefaciens* from transformed plant tissue without

using levels of antibiotics that are toxic to the plants. Here, we present the functional

characterization of genes involved in β-lactam resistance in *A. tumefaciens.* Knowledge about

proteins that promote or inhibit β-lactam resistance will enable the development of strains to

improve the efficiency of *Agrobacterium*-mediated plant genetic transformations. Effective

removal of *Agrobacterium* from transformed plant tissue has the potential to maximize crop

yield and food production, improving the outlook for global food security.

Keywords

Antibiotic resistance, *Agrobacterium tumefaciens*, β-lactamases, anhydro amidases,

membrane-bound lytic transglycosylases, ampicillin, plant transformation

INTRODUCTION

Rhizobiaceae is a family of bacteria that include soil-dwelling and plant associated bacteria.

While some species of this family have the ability to establish symbiotic relationships with

plants, others are pathogenic such as the genus *Agrobacterium*. Members of this genus are

responsible for a number of diseases, including the cane gall disease (*Agrobacterium rubi*),

hairy root disease (*Agrobacterium rhizogenes*), crown gall disease of grapes (*Agrobacterium*

vitis), and crown gall disease to flowering plants and woody shrubs (*Agrobacterium*

tumefaciens) (1–5). In nature, *A. tumefaciens* causes crown gall by adhering to wounded

plants and injecting a section of a bacterial DNA plasmid (Transfer DNA [T-DNA]) that

integrates into the plant chromosomes (1–3, 6–11). The T-DNA segment encodes for

metabolic and hormone-producing enzymes that hijack the plant to produce custom energy

sources that only *Agrobacterium* can use (9, 10). The increased abundance of energy

sources, such as phytohormones, within plant cells leads to their over-proliferation and

eventual gall formation (6). Gall formation on plants and trees leads to crop damage, and

significant economic losses have been attributed to this issue every year (2, 3).

While the genus *Agrobacterium* exhibits pathogenicity against plants, the natural ability of

Agrobacterium to transfer DNA to plants has been exploited to produce transgenic plants

through genetic engineering (6, 9, 11–13). However, one challenge for *A. tumefaciens*-

mediated plant transformations is the elimination of the bacteria from the transformed plant

tissue. Elimination of recombinant *A. tumefaciens* from plant tissues is crucial to prevent

detrimental effects to plants and to reduce the risk of releasing engineered bacteria into the

environment (14–16). β-lactam antibiotics are frequently applied during plant transformations

to eliminate *A. tumefaciens* from plant tissues and are preferred over other classes of

antibiotics (17–19). This is because β-lactams target cell wall synthesis, a process unique to

bacteria, and therefore are less toxic to eukaryotic plant cells than antibiotics that inhibit

protein or nucleic acid synthesis (20, 21). However, the natural resistance of *A. tumefaciens*

to β-lactams can be only overcome with toxic levels (~200-1000 mg/L), which has been

shown to cause embryogenic tissue necrosis, or to affect plant tissue growth and regeneration

rates in a wide variety of plants (17, 18, 22–28). Moreover, depending on the concentration

and class of β-lactam, clearing *Agrobacterium* from embryos can take up to 60 days, yet, in

some cases, complete elimination of *A. tumefaciens* is not achieved (29). Thus, currently,

there is a need for the identification and understanding of regulatory pathways and enzymes

involved in β-lactam resistance in *A. tumefaciens.* Functional characterization of bacterial

enzymes involved in β-lactam resistance will permit the development of tools that could

improve the efficiency of plant genetic transformations, and therefore maximize crop yields

and food production.

β-lactam antibiotics target the bacterial cell wall by inhibiting the activity of Penicillin

Binding Proteins (PBPs); the enzymes involved in the synthesis of the bacterial peptidoglycan

(PG) cell wall (30–38). The bacterial PG cell wall is an essential polymer consisting of

alternating *N*-acetylglucosamine (Glc*N*Ac) and *N*-acetylmuramic acid (Mur*N*Ac) sugars

crosslinked through peptide bridges (39–44). Because the PG cell wall is a covalently-

enclosed polymer, its expansion not only requires cell wall synthesis but also remodeling. Cell

wall remodeling is mediated by PG degradation enzymes such as the lytic transglycosylases

(LT) (45–48). To allow cell wall expansion, LTs cleave between the Mur*N*Ac and Glc*N*Ac

sugar strands resulting in the formation of 1,6-anhydroMurNAc Glc*N*Ac on glycan strands and

the liberation of 1,6-anhydromuropeptides (AnhMP) cell wall degradation fragments. The

liberated AnhMP fragments are transported to the bacterial cytoplasm for cell wall recycling

(42, 49, 50). In the cytoplasm, the recycling of AnhMP fragments keeps the concentration of

these products low (51, 52). However, cell wall stressors such as treatment with β-lactam

antibiotics, or mutations that inhibit the cell wall recycling pathway result in the accumulation

of AnhMP cell wall degradation fragments and derepression of β-lactamases (34, 50, 53–56).

In bacterial organisms including *Pseudomonas aeruginosa* and *Enterobacter cloaceae,* the

AnhMP cell wall degradation fragments are transcriptional activators of inducible β-

lactamases, which are enzymes that cleave and inactivate β-lactam antibiotics (57–63).

In the soil environment, many soil microorganisms produce antibiotics to compete for

survival, selecting for intrinsic resistance pathways in soil pathogens. For example, the

genomes of many soil bacteria contain β-lactamases, such as the cephalosporinase AmpC

(34, 64). As a cephalosporinase, AmpC is known to destroy β-lactam antibiotics including

monobactams, cephalosporins, and penicillins (34). The AmpC consensus protein sequence

consists of a signal sequence for periplasmic transport and a β-lactamase catalytic domain

(Fig. S1A). The regulation of AmpC expression varies across bacterial organisms. In

Escherichia coli, AmpC is a non-inducible β-lactamase that is expressed at low levels and

regulated by a promoter and a growth rate-dependent attenuator mechanism (65–67). In

contrast, in *P. aeruginosa* and other enterobacteria, AmpC is normally expressed at low

levels, but is inducible and can be derepressed during exposure to β-lactams (34, 60, 68). In

these cases, AmpC expression is regulated by AmpR, a LysR-type transcriptional regulator

found in an operon with AmpC (60). AmpR consists of two domains; a Helix-Turn-Helix DNA-

binding domain (DBD) that binds the intergenic region between AmpC and AmpR, and a LysR

effector-binding domain (EBD), which contains the regulatory region of AmpR (**Fig. 1A, Fig.**

S1A) (57). AmpR is a bifunctional transcriptional regulator that controls both the activation

and repression of AmpC (60). The induction mechanism of AmpC by AmpR in response to β-

lactams is linked to bacterial cell wall synthesis, remodeling, and recycling (34, 50, 53, 57, 62,

69). Indeed, the AnhMur cell wall degradation fragments released by LTs during cell growth,

are AmpR-activating molecules. In contrast, cell wall building blocks such as UDP-Glc*N*Ac

Mur*N*Ac pentapeptides bind to AmpR and repress *ampC* transcription (**Fig. 1A**). A block in

bacterial cell wall synthesis after exposure to β-lactams results in accumulation of AnhMur cell

wall degradation products in the bacterial cytoplasm, displacement of the AmpR repressor

UDP-Glc*N*Ac Mur*N*Ac pentapeptide, activation of AmpR, and transcription of *ampC* (**Fig. 1A**)**.**

The genus *Agrobacterium* is naturally resistant to β-lactams and the molecular basis for

this resistance is poorly understood. A study that screened for β-lactamase production in *A.*

tumefaciens detected cephalosporinase production (70) and the *A. tumefaciens* genome

encodes one putative cephalosporinase, an AmpC homolog (*atu3077)*. In addition, similar to

Enterobacteriaceae and *Pseudomonadales, ampC* is found in an operon with *ampR* in most

Agrobacterium genomes **(Fig. 1A)**. Here, we sought to determine if AmpC is a functional β-

lactamase*,* if AmpC is inducible, and if the natural ampicillin resistance observed in *A.*

tumefaciens is dependent on AmpC.

We present a functional characterization of proteins involved in intrinsic ampicillin

resistance in *A. tumefaciens.* We found that AmpR is required for AmpC-dependent β-

lactamase activity and that loss of the anhydro-amidase AmpD (*atu2113*, misannotated as an

AmiD homolog in the genome (71)) led to increased resistance to ampicillin, a process

dependent on AmpC. We suggest that AmpD is required for proper recycling of cell wall

degradation products and its loss results in the accumulation of cell wall degradation products

and activation of AmpC by AmpR. Furthermore, we found that deletion of a single LT, the

membrane-bound lytic transglycosylase B3 (MltB3), is necessary for AmpC-dependent β-

lactamase activity and its loss led to ampicillin sensitivity in the *A. tumefaciens* C58 and

GV3101 strains. Finally, transformation of *Arabidopsis thaliana* utilizing a *ΔmltB3* GV3101

strain required significantly lower concentrations of ampicillin while exhibiting similar WT

transformation efficiency. This work underscores the significance of understanding the β-

lactam resistance pathway of *A. tumefaciens* with the aim of expanding tools for the *A.*

tumefaciens-mediated transformations.

RESULTS AND DISCUSSION

The AmpC-AmpR operon is responsible for inducible ampicillin resistance in *A.*

tumefaciens **C58**

To begin our characterization*,* we first assessed the susceptibility of *A. tumefaciens* to different concentrations of ampicillin around the minimum inhibitory concentration (MIC) reported for *A. tumefaciens* on solid and liquid media (72). We found that cells grown in LB with 25 or 50 µg/mL ampicillin (AMP 25 or AMP 50, respectively) for 24 h displayed slow growth in liquid medium in comparison to cells grown in LB without ampicillin (No AMP) **(Fig. 1B)**. To better understand the cause of this growth defect, we performed phase-contrast microscopy of cells treated with AMP 25 for 2 h **(Fig. S1B)**. We found that treatment with AMP 25 causes a significant increase in the median cell length **(Fig. S1C)** and 23.5% of were undergoing cell lysis **(Fig. 1C, Fig. S1B)**, confirming that the bactericidal effect of AMP 25 on WT *A. tumefaciens* is the cause of the overall decrease in optical density*.* Similarly, WT cells grown on AMP 25 solid medium for 36 h have a viability defect in comparison to WT cells grown in LB No AMP **(Fig. 1D)**. The increased sensitivity of WT *A. tumefaciens* to ampicillin in the presence of sulbactam, a broad spectrum β-lactamase inhibitor, suggests that β-lactamase production is responsible for the observed ampicillin resistance **(Fig. 1E)**. Finally, to determine if β-lactamase production is induced, we treated WT cells with AMP 25 for 2 h, generated whole cell lysates, and performed nitrocefin assays on total protein content **(Fig. 1F)**. Nitrocefin is a chromogenic substrate related to the cephalosporins that undergoes a color change when is hydrolyzed by β-lactamases (73). After treatment of WT cells with AMP 25 for 2 h, the activity of β-lactamases is readily detected in nitrocefin assays **(Fig. 1F).** Together, these results suggest that *A. tumefaciens* C58 β-lactamase production is induced in the presence of β-lactams such as ampicillin. To assess the contributions of putative enzymes involved in ampicillin resistance, we employed a reverse genetics approach (74).

The *ampC* ortholog of *A. tumefaciens* C58 (*atu3077)* encodes the only putative

inducible β-lactamase in the genome of *A. tumefaciens* C58 (34)*. ampC* is syntenic with

ampR and the *A. tumefaciens* AmpC and AmpR proteins are 74.7% and 85.9% similar to their

respective orthologs from *P. aeruginosa***.** To determine the role of AmpC, we deleted *ampC*

(*atu3077*) from the *A. tumefaciens* C58 genome. Deletion of *ampC* does not impact cell

growth and cell viability **(Fig. 1B, 1D)** or cell morphology **(Fig. S1B)**, beyond a slight increase

in cell length **(Fig. S1C).** To pinpoint the contribution of *ampC* to ampicillin resistance, we

assessed the growth dynamics of *ΔampC* cells in the presence of AMP in liquid **(Fig. 1B)**.

ΔampC cells treated with AMP 25 or AMP 50 for 24 h showed a severe growth defect

indicating that *ΔampC* is required for ampicillin resistance. *ΔampC* cells treated with AMP 25

exhibited cell division defects (34.8%) and cell lysis (15.8%) **(Fig. 1C, Fig. S1B).** The

increased cell length of *ΔampC* treated with AMP 25 is likely due to the observed cell division

defects **(Fig. S1B-C, Fig 1C).**

To confirm that ampicillin resistance is mediated by the AmpC β-lactamase, we used

the disc diffusion assay to compare resistance levels to ampicillin in the presence and

absence of the broad-spectrum β-lactamase inhibitor, sulbactam **(Fig. 1E)**. As expected,

ΔampC led to increased sensitivity to ampicillin and the presence of sulbactam did not result

in large increases in the zone of growth inhibition **(Fig. 1E).** Furthermore, monitoring the rates

of nitrocefin hydrolysis showed that production of β-lactamase is readily detected in WT cells

treated with AMP 25 but is undetectable in *ΔampC* **(Fig. 1F)**. Together, these observations

suggest that the natural resistance to ampicillin depends on the presence of AmpC, which

functions as an inducible β-lactamase.

We hypothesized that if transcription of *ampC* is strictly controlled by AmpR, deletion of

ampR should mimic deletion of *ampC.* To test this hypothesis, we deleted *ampR* (*atu3078*)

from the genome of *A. tumefaciens* C58. Deletion of *ampR* led to normal cell growth **(Fig.**

1B), morphology **(Fig. S1B),** cell length **(Fig. S1C)** and cell viability **(Fig. 1D).** Low

concentrations of ampicillin in either liquid or solid medium was lethal to *ΔampR* and resulted

in similar cell division defects and cell lysis as *ΔampC* **(Fig. 1B-C, Fig. S1B)**. Like *ΔampC,*

ΔampR is sensitive to ampicillin treatment **(Fig. 1D-E)** and fails to produce detectable β-

lactamase activity when treated with AMP 25 **(Fig. 1F, Fig. S1D).** These results suggest that

AmpR is necessary for induction of the AmpC β-lactamase in the presence of ampicillin.

Loss of AmpD derepresses β-lactamases in *A. tumefaciens* C58.

The finding that AmpC and AmpR are necessary for ampicillin resistance in *A. tumefaciens*

C58 led us to explore how the pools of different cell wall fragments would alter the AmpC-

mediated β-lactamase induction. Similar to exposure to β-lactams, loss of cell wall recycling

amidases has been shown to increase the AmpR-activating fragments (cell wall degradation

fragments) in the cytoplasm, resulting in the transcriptional derepression of *ampC* and β-

lactam resistance (48, 75–78). The genome of *A. tumefaciens* contains one 1,6-anhydro

amidase ortholog, *atu2113*, reannotated as AmpD. The domain organization of AmpD

consists of the Amidase_2 (Ami_2) catalytic domain and a PG-binding domain (PBD) that

facilitates the interaction with cell wall products **(Fig. S2A)** (45, 48). *A. tumefaciens* AmpD

exhibits 64.4% sequence similarity to AmpDh2, one of three broad-spectrum 1,6-anhydro

amidase AmpDh paralogs found in *Pseudomonas aeruginosa* (54, 56, 77, 79).

Given that in *A. tumefaciens* ampicillin triggers the AmpC-dependent production of β-

lactamases, we hypothesized that if AmpD was an anhydro amidase involved in the recycling

of cell wall degradation fragments, its loss should result in increased AmpR-activating

fragments in the cytoplasm, β-lactamase induction, and ampicillin resistance. First, we found

that *ΔampD* cells exhibit normal cell viability **(Fig. 2A)**, cell growth **(Fig. 2B),** and morphology

(Fig. S2B-C). *ΔampD* is highly resistant to ampicillin **(Fig. 2A)**. Indeed, WT cells spotted on

AMP 160 were not viable, whereas *ΔampD* spotted on AMP 160 only displayed a ~10-fold

decrease in viability compared to untreated cells. In liquid, *ΔampD* cells continued to grow

normally, even in the presence of AMP 100 **(Fig. 2B)** and ampicillin treatment did not trigger

obvious morphological changes or cell lysis **(Fig. S2B-C, Fig. 2C)**. Nitrocefin assays of

ΔampD indicated readily detectable amounts of β-lactamase production in both the presence

and absence of ampicillin **(Fig. 2D, Fig. S2D).** The increased zone of inhibition observed in

the presence of ampicillin and sulbactam is consistent with the high level of ampicillin

resistance observed in *ΔampD* being mediated by a β-lactamase **(Fig. 2E)**. Together, these

results indicate loss of AmpD leads to derepression and overproduction of β-lactamases. Our

findings are consistent with other bacterial models such as *P. aeruginosa*, where deletion of

1,6-anhydro amidases involved in the recycling of AmpR-activating ligands leads to increased

β-lactamase expression (53–56) due to the build-up of activating ligands in the cytoplasm.

AmpC is constitutively produced in *ΔampD*

We have shown that AmpC and AmpR are required for ampicillin resistance **(Fig. 1)**, and loss

of AmpD leads elevated β-lactamase production and ampicillin resistance in *A. tumefaciens*

C58 **(Fig 2)**. To confirm that AmpC is the β-lactamase produced by the *ΔampD* strain, we

deleted *ampC* or *ampR* in the *ΔampD* background **(Fig. 3)**. In the absence of ampicillin, we

found that *ΔampCΔampD* and *ΔampRΔampD* display normal cell viability and growth **(Fig.**

3A, Fig S3A). We found that treatment with AMP 25, on either solid medium or liquid medium,

is lethal to *ΔampCΔampD* or *ΔampRΔampD* **(Fig. 3A, Fig. S3A).** Treatment of *ΔampCΔampD*

or *ΔampRΔampD* with AMP 25 for 2 h resulted in lysis of 22.9% and 33.8% of the cells in

comparison to untreated cells (No AMP), where lysis was not readily observed **(Fig. 3B, Fig**

S3B). In comparison to *ΔampC* or *ΔampR*, where cell division defects were observed in >20%

of the population, very few *ΔampCΔampD* or *ΔampRΔampD* cells exhibited cell division

defects when treated with AMP 25 **(Fig. 3B).** The low incidence of cell division defects

observed in *ΔampCΔampD* or *ΔampRΔampD* suggest that the activity AmpD contributes to

the inefficient cell division of *ΔampC* and *ΔampR* cells following ampicillin treatment. Finally,

to assess whether the *ΔampD* strain could induce β-lactamase production in the absence of

ampC or *ampR*, we performed nitrocefin assays. *ΔampCΔampD* and *ΔampRΔampD* failed to

produce detectable levels of β-lactamase in the absence or presence of AMP 25 **(Fig. 3D,**

Fig. S3C). Together, these results suggest that induction of AmpC is the main cause for the

elevated resistance to ampicillin observed in *ΔampD*. This data is consistent with the current

model for β-lactam resistance in *P. aeruginosa,* where the loss of anhydro amidases leads to

accumulation of cell wall degradation products that activate AmpR leading to derepression of

AmpC (54–56, 69).

Absence of MltB3 (*Δatu3779*) leads to a failure of AmpC-dependent induction of β-

lactamases

LTs are likely to function as the enzymes that generate the AmpR-activating fragments.

Different families of LTs have been linked to β-lactam resistance in several bacterial

organisms (80, 81). For instance, in *Caulobacter crescentus* deletion of *sdpA*, a soluble LT,

led to reduced sensitivity to ampicillin (82). In *P. aeruginosa,* loss of several *mltBs* and/or *slt*

led to a decrease in the β-lactam MIC, cell viability, and increased outer membrane

permeability (83, 84). Thus, we sought to determine if LTs contribute to β-lactam resistance of

A. tumefaciens. The *A. tumefaciens* genome consists of 8 putative LTs belonging to 3

families: family 1) the soluble lytic transglycosylases (Slt), 2) Membrane-bound lytic

transglycosylase A (MltA), and 3) Membrane-bound lytic transglycosylase B (MltB) **(Fig. S4A)**.

We found that single deletions of LTs did not affect cell viability, suggesting a wide

redundancy of functions between LTs **(Fig. S4B)**.

Despite the potential for functional redundancy, we found that deletion of a single, family 3,

membrane-bound lytic transglycosylase, MltB3 (*atu3779)*, led to ampicillin hypersensitivity

(Fig. S4B, Fig. 4A). Treatment of *ΔmltB3* with AMP 25 for 2 h led to a severe cell lysis defect

(28.8%) **(Fig. 4A-B)**. In addition, *ΔmltB3* cells grown in AMP 25 for 24 h had a severe growth

defect indicating that MltB3 is required for ampicillin resistance **(Fig. 4C)**. Finally, *ΔmltB3*

exhibited reduced production of β-lactamase after AMP 25 treatment for 2 hours **(Fig. 4D,**

Fig. S4C). Together, these results suggest that in *A. tumefaciens,* MltB3 is a specialized

enzyme that functions in the AmpR-AmpC β-lactamase pathway. These observations contrast

with the *P. aeruginosa* model*,* in which the β-lactam sensitivity of LT mutants are due to

increased outer membrane permeability rather than β-lactamase production (83)**.**

The finding that *Δmltb3* led to ampicillin sensitivity and reduced β-lactamase induction in

A. tumefaciens 58C prompted us to ask whether *Δmltb3* could be sensitive to other β-lactams

or if MltB3 performs the same function in other *A. tumefaciens* strains, especially in those

commonly used for plant transformations. To this end, we deleted *mltB3* in *A. tumefaciens*

GV3103 and found that this mutation made the strain susceptible to AMP 25 **(Fig 5A)**. In

addition, deletion of *mltB3* in GV3101 led to sensitivity to carbenicillin 15 (CARB 15), an

antibiotic commonly used in plant tissue culture systems **(Fig. 5A)**. To confirm that the

absence of MltB3 (*ΔmltB3*) prevented the induction of β-lactamase production in the GV3101

strain after ampicillin treatment, we performed nitrocefin assays using lysates of cells treated

with AMP 25 for 2 h. While β-lactamase activity is readily detected from WT GV3101-treated

cells with AMP 25, the *ΔmltB3* mutant produced relatively low levels of inducible β-lactamase

(Fig. 5B, Fig. S5). Together, we conclude that MltB3 is the major LT in *A. tumefaciens* C58

and GV3101 that is required for the *A. tumefaciens* natural resistance to ampicillin and other

β-lactams such as carbenicillin.

Plant transformation with *Δmltb3* cells requires a low concentration of ampicillin for

the elimination of bacteria

We reasoned that ampicillin sensitivity observed in *ΔmltB* GV3101 could prove to be

beneficial during plant transformation. First, we confirmed that *ΔmltB3 A. tumefaciens*

GV3101 could effectively transform *Arabidopsis thaliana* using the standard floral dip

technique (85) with an efficiency comparable to WT **(Fig. 5C)**. Next, we asked whether

elimination of the bacteria could be performed using low concentrations of ampicillin (AMP

25). Seeds transformed with WT *A. tumefaciens* GV3101 contained similar bacterial loads

compared to seeds transformed with *Δmltb3 A. tumefaciens* GV3101 when plated on solid

media without ampicillin **(Fig. 5D)**. When plated on media containing AMP 25, seeds

transformed with *Δmltb3* exhibited a significant drop in bacterial load. In contrast, this low

level of ampicillin did not reduce the bacterial load of seeds transformed with WT GV3101

(Fig. 5D). These results demonstrate that the *Δmltb3* GV3101 strain can be utilized to

transform *Arabidopsis thaliana* and elimination of the bacteria can be performed using lower

concentrations of ampicillin. Together, this shows the significance of understanding the cell

biology of *A. tumefaciens* as a means to improve genetic engineering approaches.

CONCLUDING REMARKS AND FUTURE PERSPECTIVES

The natural ability of *A. tumefaciens* to transform plants has allowed the production of

transgenic crops of incredible economic importance for the past four decades. One challenge

of the *A. tumefaciens*-mediated plant transformation is natural resistance of *A. tumefaciens* to

antibiotics, which requires toxic concentrations of antibiotics to eliminate *A. tumefaciens* from

transformed tissues. Here we show that *A. tumefaciens* induces β-lactamase activity in

response to ampicillin exposure. Indeed, induction of β-lactamase activity upon exposure to

ampicillin is dependent on the β-lactamase AmpC and the transcription factor AmpR.

Moreover, we found that deletion of a single LT, MltB3, sensitizes *A. tumefaciens* to the β-

lactams.

We propose that during *A. tumefaciens* growth and remodeling, there is a delicate balance

between the synthesis and degradation of the bacterial cell wall. PBPs insert precursor

cytoplasmic monomers into the growing cell wall polymer (**Fig 6, steps 1-2**). During

remodeling, cell wall hydrolytic enzymes such as such as endopeptidases and LTs, including

MltB3, liberate cell wall degradation products, which are transported back into the cytoplasm

of *A. tumefaciens* for their recycling (**Fig 6, steps 3-4**). In the cytoplasm, hydrolytic enzymes

including L,D-carboxypeptidases (LD-CPases), amidases such as AmpD, and glycosidases

limit the pool of cell wall fragments by allowing their recycling **(Fig 6, steps 5-8).** When cell

wall degradation fragments accumulate during β-lactam treatment, the AmpR-dependent

production of the AmpC β-lactamase is increased **(Fig 6, step 9)**. Overall, the identification

and contributions of genes conferring ampicillin resistance in *A. tumefaciens* will be beneficial

for improving the design of *A. tumefaciens*-mediated genetic engineering.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *Agrobacterium tumefaciens* C58 and derived

strains were grown in Luria-Bertani (LB) medium (10 g/l tryptone, 10 g/l yeast extract and 5 g/l

NaCl) at 28°C with shaking. *E. coli* DH5α and S17-1 were routinely cultivated in LB medium at

37°C with shaking and the antibiotic kanamycin was added at a concentration of 50 µg/ml and

25 µg/ml to maintain plasmids in DH5α and S17-1 strains, respectively.

Construction of expression plasmids and strains. All strains and plasmids used are listed

in **Table 1**. Synthesized DNA primers are listed in **Table 2**. To construct deletion vectors, the

corresponding coding sequences were amplified from purified *Agrobacterium tumefaciens*

C58 genomic DNA. The amplicons and plasmid pNTPS139 were digested overnight and

ligated overnight at 4°C using NEB T4 DNA ligase.

Construction of deletion constructs and deletion strains. Gene deletions were achieved

by allelic exchange and vectors were constructed as previously described (86). Briefly, 500

base pairs (bp) fragments upstream and downstream of the gene of interest were amplified

using primer pairs P1/P2; which amplify 500 bp upstream of the gene of interest, and P3/P4;

which amplify the 500 bp downstream of the gene of interest. Overlapping PCR was used to

merge and amplify the amplicons generated by P1/P2 and P3/P4, using primer pair P1/P4.

The 1,000 bp amplicon was digested and ligated into the deletion plasmid, pNTPS139. The

deletion plasmids were introduced into *A. tumefaciens* by mating using an *E. coli* S17

conjugation strain to create kanamycin (KAN) resistant, sucrose sensitive primary

exconjugants. Deletion strains were constructed as described previously (74). Briefly, primary

exconjugants were grown overnight at 28°C in minimal medium ATGN with no selection, and

plated in ATGN KAN 300 for 48 h at 28°C. Colonies were screened by patching for KAN

resistance and sucrose sensitivity. Colony PCR using primers P5/P4 were used to confirm

that recombination took place and at the region of interest. Next, positive colonies are grown

in ATGN at 28°C overnight and plated in ATSN 5% sucrose. Secondary recombinants were

screened by patching for sucrose resistance and KAN sensitivity. Colony PCR with primers

P5/P6 for the respective gene target was used to confirm deletion. PCR products from P5/P6

primer sets were sequenced to further confirm deletions.

Cell viability spot assays. The cell viability assay was performed as described (87). For cell

viability spot assays, cultures were grown overnight and diluted to an OD_{600} of 0.05 and

serially diluted in LB up to 10^{-6}. 4 µl of each dilution was spotted onto LB plates and incubated

at 28°C for 36-40 h before imaging. To determine ampicillin antibiotic resistance, *A.*

tumefaciens cultures were grown overnight and diluted to an optical density at 600 nm (OD_{600})

of 0.05 and serially diluted and spotted onto LB plates containing AMP 25, AMP 80 and/or

AMP 160 using the cell viability assay protocol.

Phase-contrast microscopy. For phase-contrast microscopy, 0.8 µl of exponentially-grown

cultures were spotted onto a 1.25% agarose pad as previously described (87) using a Nikon

Eclipse Ti inverted microscope through a Nikon Plan 60X oil Ph3 objective. Cell length

analysis was performed using the MicrobeJ plug-in for Fiji (88). A One-Way analysis of

variance (ANOVA) Kruskal-Wallis test with Dunn's post-test was used to compare the

indicated strains. Images were prepared using Adobe Photoshop, Adobe Illustrator, and

Prism.

Growth curves*.* For growth curves, exponentially-growing cultures were diluted to an OD_{600} =

0.2 and submitted to a plate reader at 28°C with shaking every 5-10 min. Ampicillin was

added to a final concentration of AMP 25 or AMP 100. OD_{600} represents two technical

replicates for each culture measured every 5 minutes for 24-48 h in 96-well plates.

Disc diffusion method. The disc-diffusion assay was used to determine the resistance of

Agrobacterium to various antibiotics. Cells were growth on LB agar plates. Sterile discs (6.5

mm in diameter) were placed on the surface of LB agar plates seeded with overnight culture

of indicated strains. We used three different discs: Blank (sterile disc containing no antibiotic),

AMP 10 (ampicillin 10µg/ml), and AMP 10/SUL 10 (ampicillin 10µg/ml sulbactam 10µg/ml).

The LB agar assay plates used for testing *A. tumefaciens* susceptibility were incubated at

28°C for 24-36 h. The assessment of antibacterial activity was based on the measurement of

diameter of the zone of inhibition formed around the disc minus the size of the disc. Three

independent trials were conducted for each concentration of each antibiotic. A Brown-

Forsythe and Welch analysis of variance (ANOVA) was used to compare the means within

each indicated group.

***Nitrocefin Assay**.* Nitrocefin is a chromogenic substrate for measuring β-lactamase activity.

Nitrocefin has an absorbance maximum of 390 nm. Upon hydrolysis of the β-lactam ring by a

β-lactamase, the absorbance shifts from 390 nm to 486 nm. By monitoring absorbance at 486

nm over time and using Beer's law (A=εlc), we directly measured the rate of β-lactamase

hydrolytic activity. In two separate experiments, indicated strains were grown in LB until

desired optical densities (OD600) of 0.6 were reached. The cells were then pelleted by

centrifugation at 25,900 x g (rotor: Fiberlite F14-6x250y) for 5 minutes and the supernatant

collected and washed three times with PBS. Cell lysates were generated by adding the cell

lysis buffer BugBuster and sonication. No lysozyme or protase inhibitors were added. Cell

lysates were normalized based on total protein content (7.5 μg/ml) and volume before

incubation in 100 μM nitrocefin solution in a 200 μl reaction volume at room temperature in 20

mM HEPES, 300 mM NaCl, pH 7.5. BugBuster lysis buffer was used as blank. *ΔampD* lysates

were normalized based on total protein content (7.5 μg/ml) and subsequently diluted to 1:5 as

the rate of nitrocefin hydrolysis was significantly faster than the controls (WT, WT AMP 25).

Absorbance was immediately measured at 486 nm in 5 second intervals for 300 seconds (5

minutes). The change in absorbance (A) was converted to change in concentration (c) of

hydrolyzed product by using Beer's law (A=εlc) where the molar extinction coefficient (ε) =

20500 M-1 cm-1 and path length (l) = 1 cm. Data represents one biological replicate.

Construction of Agrobacterium strain for plant transformation experiments. *mltB3* was

deleted from the genome of WT *Agrobacterium tumefaciens* GV3101 using the same plasmid

and technique used to delete the gene in the strain C58. The resulting strain,

GV3101 *ΔmltB3*, was next transformed by introduction of the empty binary vector, pUBQ10-

GW (89) via electroporation. This plasmid confers KAN resistance to

the *Agrobacterium* strain, and also contains T-DNA repeat blocks which allow the transfer of

BASTA resistance to transformed into *Arabidopsis thaliana*. pUBQ10-GW was also

electroporated into wildtype GV3101 as a control.

***Arabidopsis thaliana* transformation efficiency**. Plants with a bolt height between 2 and 7

centimeters were transformed via the floral dip method (85). 5 plants were transformed on

separate days from independent colonies for each C58 and GV3101. Plants were grown in

Promix BX (Premier Tech Horticulture) at 23°C, 16 h light/8 h dark, 100–150 µE m^{-2} s^{-1}, and

50–70% humidity until seeds were fully developed. Seeds were collected from fully matured

plants and store at 4°C at low humidity for one week. Seeds were then surface sterilized in

15% bleach containing 0.1% Triton X-100 with gentle rocking for 5 minutes. Seeds were then

washed 3 times using sterile water. Seeds were then imbibed in sterile water at 4°C for 3

days. Seeds were then sown into soil using conditions listed above. Once seeds developed

true leaves BASTA (glufosinate ammonium) was applied via spray at a concentration of

10mg/L. After 4 days BASTA was re-applied ensuring only transformed plants survived.

Determining bacterial load of transformed *Arabidopsis thaliana* seeds. Seeds were

collected from fully matured transformed plants and stored at 4°C for 1 week with low

humidity. Seeds were then surface sterilized in 15% bleach containing 0.1% Triton X-100 with

gentle rocking for 5 minutes. Seeds were then washed 3 times using sterile water. 10mg of

seeds for each experimental condition were ground with a sterile mortar and pestle. Ground

seeds were then suspended in 1mL of sterile water, and serially diluted. 200µl of 10^{-1} dilution

was plated on minimal medium ATGN No AMP and AMP 25. Plates were then incubated at

28°C before colonies were counted.

ACKNOWLEDGEMENTS

We thank the Brown lab for helpful discussions. Research in the Brown lab on *A. tumefaciens*

cell growth is supported by the National Science Foundation (IOS1557806). WFC was

supported by a Gus T. Ridgel Fellowship.

AUTHOR CONTRIBUTIONS

WFC and PJBB conceptualized this work; WFC and CR developed methods; WFC, MH,

CR, AMR and AKY performed the experiments and completed data analysis; WFC, AMR, and

PJB wrote the manuscript; WFC, AMR, AKY, FC, and PJBB edited the manuscript; PJBB

acquired funding for this work.

REFERENCES

1. Valentine L, Crops GM. 2003. *Agrobacterium tumefaciens* and the plant: The David and

Goliath of modern genetics. Plant Physiol 133:948–955.

2. Escobar MA, Dandekar AM. 2003. *Agrobacterium tumefaciens* as an agent of disease.

Trends Plant Sci 8:380–386.

3. Pulawska J. 2010. Crown gall of stone fruits and nuts, economic significance and

diversity of its causal agents: Tumorigenic *Agrobacterium* spp. J Plant Pathol 92:S87–

S98.

4. Otten L, Burr T, Szegedi E. 2008. *Agrobacterium*: A disease-causing bacterium. Tzfira

T, Citovsky V (eds) *Agrobacterium* from Biol to Biotechnol Springer 1–46.

5.   Gelvin SB. 2003. *Agrobacterium*-mediated plant transformation: the biology behind the

"gene-jockeying" tool. Microbiol Mol Biol Rev 67:16–37.

6.   Nester EW. 2015. *Agrobacterium*: Nature's genetic engineer. Front Plant Sci 5:1–16.

7.   Dessaux Y, Faure D. 2018. Niche construction and exploitation by *Agrobacterium*: How

to survive and face competition in soil and plant habitats. Curr Top Microbiol Immunol

Springer 418.

8.   Singer K. 2018. The mechanism of T-DNA integration: Some major unresolved

questions. Curr Top Microbiol Immunol 418:287–317.

9.   Tzfira T, Citovsky V. 2000. From host recognition to T-DNA integration: the function of

bacterial and plant genes in the *Agrobacterium*-plant cell interaction. Mol Plant Pathol

1:201–212.

10.   Gelvin S. 2000. *Agrobacterium* and plant genes involved in T-DNA transfer and

integration. Annu Rev Plant Biol 51:223–256.

11.   Tzfira T, Citovsky V. 2006. *Agrobacterium*-mediated genetic transformation of plants:

biology and biotechnology. Curr Opin Biotechnol 17:147–154.

12.   Gelvin SB. 2003. *Agrobacterium*-mediated plant transformation: the biology behind the

"gene-jockeying" tool. Society 67:16–37.

13.   De La Riva GA, González-Cabrera J, Vázquez-Padrón R, Ayra-Pardo C. 1998.

Agrobacterium tumefaciens: A natural tool for plant transformation. Electron J

Biotechnol 1:25–48.

14.   Leifert C, Cassells AC. 2000. Microbial hazards in plant tissue and cell cultures. Vitr Cell

Dev Biol 37:133–138.

15.   Mogilner N, Zutra D, Gafny R, Bar-Joseph M. 1993. The persistence of engineered

508   *Agrobacterium tumefaciens* in agroinfected plants. Mol plant-microbe Interact MPMI.

509   16.   Matzk A, Mantell S, Schiemann J. 1996. Localization of persisting agrobacteria in

510   transgenic tobacco plants. Mol Plant-Microbe Interact.

511   17.   Hammerschlag FA, Zimmerman RH, Yadava UL, Hunsucker S, Gercheva P. 1997.

512   Effect of antibiotics and exposure to an acidified medium on the elimination of

513   *Agrobacterium tumefaciens* from apple leaf explants and on shoot regeneration. J Am

514   Soc Hortic Sci.

515   18.   Mohana Priya A, Karutha Pandian S, Manikandan R. 2012. The effect of different

516   antibiotics on the elimination of *Agrobacterium* and high frequency *Agrobacterium*-

517   mediated transformation of indica rice (Oryza sativa L.). Czech J Genet Plant Breed

518   48:120–130.

519   19.   Demain AL, Blander RP. 1999. The β-lactam antibiotics: Past, present, and future.

520   Antonie van Leeuwenhoek, Int J Gen Mol Microbiol 75:5–19.

521   20.   Okkels FT, Pedersen MG. 1988. The toxicity to plant tissue and to *Agrobacterium*

522   *tumefaciens* of some antibiotics. Acta Hortic 225:199–208.

523   21.   Shackelford NJ, Chlan CA. 1996. Identification of antibiotics that are effective in

524   eliminating *Agrobacterium tumefaciens*. Plant Mol Biol Report 14:50–57.

525   22.   Estopà M, Marfà V, Melé E, Messeguer J. 2001. Study of different antibiotic

526   combinations for use in the elimination of *Agrobacterium* with kanamycin selection in

527   carnation. Plant Cell Tissue Organ Cult 65:211–220.

528   23.   Holford P, Newbury HJ. 1992. The effects of antibiotics and their breakdown products

529   on the *in vitro* growth of *Antirrhinum majus*. Plant Cell Rep 11:93–96.

530   24.   Minden V, Deloy A, Volkert AM, Leonhardt SD, Pufal G. 2017. Antibiotics impact plant

531   traits, even at small concentrations. AoB Plants 9.

532   25.   Jiang C, Brown PJB, Ducret A, Brun Y V. 2014. Sequential evolution of bacterial

533   morphology by co-option of a developmental regulator. Nature 506:489–493.

534   26.   Wiebke B, Ferreira F, Pasquali G, Bodanese-Zanettini MH, Droste A. 2006. Influence of

535   antibiotics on embryogenic tissue and *Agrobacterium tumefaciens* suppression in

536   soybean genetic transformation. Bragantia 65:543–551.

537   27.   Obembe O, Adesoye A, Machuka J. 2005. Evaluation of the Effectiveness of Selected

538   Antibiotics in the Suppression of *Agrobacterium* from Cowpea (Vigna Unguiculata L.

539   Walp.) Embryo Explants and as Potential Selective Agents in *Agrobacterium*-Mediated

540   Transformation. J Trop Microbiol Biotechnol 1:27–33.

541   28.   Haddadi F, Aziz MA, Abdullah SNA, Tan SG, Kamaladini H. 2015. An efficient

542   *Agrobacterium*-mediated transformation of Strawberry cv. Camarosa by a dual plasmid

543   system. Molecules 20:3647–3666.

544   29.   Tang H, Ren Z, Krczal G, Glimelius K. 2000. An evaluation of antibiotics for the

545   elimination of *Agrobacterium tumefaciens* from walnut somatic embryos and for the

546   effects on the proliferation of somatic embryos and regeneration of transgenic plants.

547   Plant Cell Rep 19:881–887.

548   30.   Waxman DJ, Strominger JL. 1983. Penicillin-binding proteins and the mechanism of

549   action of β-lactam antibiotics. Annu Rev Biochem 52:825–869.

550   31.   Sauvage E, Kerff F, Terrak M, Ayala JA, Charlier P. 2008. The Penicillin-binding

551   proteins: Structure and role in peptidoglycan biosynthesis. FEMS Microbiol Rev 32:234–

552   258.

32.   Spratt BG. 2006. Distinct Penicillin-binding proteins involved in the division, elongation,

and shape of *Escherichia coli* K12. Proc Natl Acad Sci 72:2999–3003.

33.   Rolinson GN. 1980. Effect of β-lactam antibiotics on bacterial cell growth rate.

Microbiology 120:317–323.

34.   Jacoby GA. 2009. AmpC β-lactamases. Clin Microbiol Rev 22:161–182.

35.   Egan AJF, Cleverley RM, Peters K, Lewis RJ, Vollmer W. 2017. Regulation of bacterial

cell wall growth. FEBS J 284:851–867.

36.   Den Blaauwen T, Andreu JM, Monasterio O. 2014. Bacterial cell division proteins as

antibiotic targets. Bioorg Chem 55:27–38.

37.   Egan AJF, Biboy J, van't Veer I, Breukink E, Vollmer W. 2015. Activities and regulation

of peptidoglycan synthases. Philos Trans R Soc B Biol Sci 370.

38.   Typas A, Banzhaf M, Gross CA, Vollmer W. 2012. From the regulation of peptidoglycan

synthesis to bacterial growth and morphology. Nat Rev Microbiol 10:123–136.

39.   Vollmer W, Blanot D, De Pedro MA. 2008. Peptidoglycan structure and architecture.

FEMS Microbiol Rev 32:149–167.

40.   de Pedro MA, Cava F. 2015. Structural constraints and dynamics of bacterial cell wall

architecture. Front Microbiol.

41.   Margolin W. 2009. Sculpting the Bacterial Cell. Curr Biol 19:812–822.

42.   Park JT, Uehara T. 2008. How bacteria consume their own exoskeletons (turnover and

recycling of cell wall peptidoglycan). Microbiol Mol Biol Rev 72:211–227.

43.   Silhavy TJ, Kahne D, Walker S. 2010. The bacterial cell envelope. Cold Spring Harb

Perspect Biol 2:1–16.

44.   Scheffers D, Pinho MG. 2005. Bacterial cell wall synthesis: New insights from

localization studies. Microbiol Mol Bio Rev 69:585–607.

45.   Vermassen A, Leroy S, Talon R, Provot C, Popowska M, Desvaux M. 2019. Cell wall

hydrolases in bacteria: Insight on the diversity of cell wall amidases, glycosidases and

peptidases toward peptidoglycan. Front Microbiol 10:331.

46.   Frirdich E, Gaynor EC. 2013. Peptidoglycan hydrolases, bacterial shape, and

pathogenesis. Curr Opin Microbiol 16:767–778.

47.   Vollmer W, Joris B, Charlier P, Foster S. 2008. Bacterial peptidoglycan (murein)

hydrolases. FEMS Microbiol Rev 32:259–286.

48.   van Heijenoort J. 2011. Peptidoglycan hydrolases of *Escherichia coli*. Microbiol Mol Biol

Rev 75:636–663.

49.   Dhar S, Kumari H, Balasubramanian D, Mathee K. 2018. Cell-wall recycling and

synthesis in *Escherichia coli* and *Pseudomonas aeruginosa* – their role in the

development of resistance. J Med Microbiol 67:1–21.

50.   Dik DA, Fisher JF, Mobashery S. 2018. Cell-wall recycling of the Gram-negative

bacteria and the nexus to antibiotic resistance. Chem Rev. review-article, American

Chemical Society.

51.   Lee TK, Huang KC. 2013. The role of hydrolases in bacterial cell-wall growth. Curr Opin

Microbiol 16:760–766.

52.   Vollmer W. 2012. Bacterial growth does require peptidoglycan hydrolases. Mol

Microbiol 86:1031–1035.

53.   Fisher JF, Mobashery S. 2014. The sentinel role of peptidoglycan recycling in the β-

lactam resistance of the Gram-negative Enterobacteriaceae and *Pseudomonas*

aeruginosa. Bioorg Chem 56:41–48.

54. Jacobs C, Joris B, Jamin M, Klarsov K, van Beeumen J, Mengin-Lecreulx D, van

Heijenoort J, Park JT, Normark S, Frère J-M. 1995. AmpD, essential for both β-

lactamase regulation and cell wall recycling, is a novel cytosolic N-acetylmuramyl-L-

alanine amidase. Mol Microbiol 15:553–559.

55. Langaee TY, Gagnon L, Huletsky A. 2000. Inactivation of the *ampD* gene in

Pseudomonas aeruginosa leads to moderate-basal-level and hyperinducible AmpC β-

lactamase expression. Antimicrob Agents Chemother 44:583–589.

56. Schmidtke AJ, Hanson ND. 2008. Role of *ampD* homologs in overproduction of AmpC

in clinical isolates of *Pseudomonas aeruginosa*. Antimicrob Agents Chemother

52:3922–3927.

57. Jacobs C, Fre J. 1997. Cytosolic intermediates for cell wall biosynthesis and

degradation control inducible β-lactam resistance in Gram-negative bacteria 88:823–

832.

58. Dik DA, Domínguez-Gil T, Lee M, Hesek D, Byun B, Fishovitz J, Boggess B, Hellman

LM, Fisher JF, Hermoso JA, Mobashery S. 2017. Muropeptide binding and the X-ray

structure of the effector domain of the transcriptional regulator AmpR of *Pseudomonas*

aeruginosa. J Am Chem Soc 139:1448–1451.

59. Johnson JW, Fisher JF, Mobashery S. 2013. Bacterial cell-wall recycling. Ann N Y Acad

Sci 1277:54–75.

60. Balasubramanian D, Kumari H, Mathee K. 2014. *Pseudomonas aeruginosa* AmpR: An

acute-chronic switch regulator. Pathog Dis 73:1–14.

61. Vadlamani G, Thomas MD, Patel TR, Donald LJ, Reeve TM, Stetefeld J, Standing KG,

Vocadlo DJ, Mark BL. 2015. The β-lactamase gene regulator AmpR is a tetramer that

recognizes and binds the D-Ala-D-Ala motif of its repressor UDP-N-acetylmuramic acid

(MurNAc)-pentapeptide. J Biol Chem 290:2630–2643.

62. Lindberg F, Normark S. 1987. Common mechanism of ampC β-lactamase induction in

enterobacteria: Regulation of the cloned *Enterobacter cloacae* P99 β-lactamase gene. J

Bacteriol 169:758–763.

63. Torrens, G, Hernández S, Alfonso Ayala J, Moya B, Juan C, Cava F, Oliver A. 2019.

Regulation of AmpC-driven β-lactam resistance in *Pseudomonas aeruginosa*: Different

pathways, different signaling 4:1–14.

64. Juan C, Torrens G, González-Nicolau M, Oliver A. 2017. Diversity and regulation of

intrinsic β-lactamases from non-fermenting and other Gram-negative opportunistic

pathogens. FEMS Microbiol Rev 41:781–815.

65. Bajaj P, Singh NS, Virdi JS. 2016. *Escherichia coli* β-lactamases: What really matters.

Front Microbiol 7:1–14.

66. Jaurin BA, Normark S. 1979. *In vivo* regulation of chromosomal β-lactamase in

Escherichia coli. J Bacteriol 138:896–902.

67. Jaurin B, Grundström T, Edlund T, Normark S. 1981. The *E. coli* β-lactamase attenuator

mediates growth rate-dependent regulation. Nature 290:221–225.

68. Honoré N, Nicolas MH, Cole ST. 1986. Inducible cephalosporinase production in clinical

isolates of *Enterobacter cloacae* is controlled by a regulatory gene that has been

deleted from *Escherichia coli*. EMBO J 5:3709–3714.

69. Jacobs C, Huang LJ, Bartowsky E, Normark S, Park JT. 1994. Bacterial cell wall

recycling provides cytosolic muropeptides as effectors for β-lactamase induction. EMBO

J 13:4684–4694.

70. Ogawa Y, Mii M. 2004. Screening for highly active β-lactam antibiotics against

Agrobacterium tumefaciens. Arch Microbiol 181:331–336.

71. Goodner B, Hinkle G, Gattung S, Miller N, Blanchard M, Qurollo B, Goldman BS, Cao

Y, Askenazi M, Halling C, Mullin L, Houmiel K, Gordon J, Vaudin M, Iartchouk O, Epp A,

Liu F, Wollam C, Allinger M, Doughty D, Scott C, Lappas C, Markelz B, Flanagan C,

Crowell C, Gurson J, Lomo C, Sear C, Strub G, Cielo C, Slater S. 2001. Genome

Sequence of the Plant Pathogen and Biotechnology Agent *Agrobacterium tumefaciens*

C58. Science (80-) 294:2323 LP – 2328.

72. CAO Y, Niimi Y, HU S lian. 2006. Meropenem as an alternative antibiotic agent for

suppression of *Agrobacterium* in genetic transformation of orchid. Agric Sci China

5:839–846.

73. O'Callaghan CH, Morris A, Kirby SM, Shingler AH. 1972. Novel method for detection of

β-lactamases by using a chromogenic cephalosporin substrate. Antimicrob Agents

Chemother 1:283–288.

74. Morton ER, Fuqua C. 2012. Genetic manipulation of *Agrobacterium*. Curr Protoc

Microbiol Chapter 3:Unit-3D.2.

75. Uehara T, Park JT. 2007. An anhydro-N-acetylmuramyl-L-alanine amidase with broad

specificity tethered to the outer membrane of *Escherichia coli*. J Bacteriol 189:5634–

5641.

76. Pennartz A, Généreux C, Parquet C, Mengin-Lecreulx D, Joris B. 2009. Substrate-

induced inactivation of the *Escherichia coli* AmiD N-acetylmuramoyl-L-alanine amidase

highlights a new strategy to inhibit this class of enzyme. Antimicrob Agents Chemother

53:2991–2997.

77.   Langaee TY, Gagnon L, Huletsky A. 2000. Inactivation of the *ampD* gene in

669   P*seudomonas aeruginosa* leads to moderate-basal-level and hyperinducible AmpC β-

670   lactamase expression. Antimicrob Agents Chemother 44:583–589.

671   78.   Mercier F, Zervosen A, Teller N, Frre JM, Herman R, Pennartz A, Joris B, Luxen A.

672   2010. 1,6-AnhMurNAc derivatives for assay development of amidase AmiD. Bioorganic

673   Med Chem 18:7422–7431.

674   79.   Juan C, Moyá B, Pérez JL, Oliver A. 2006. Stepwise upregulation of the *Pseudomonas*

675   *aeruginosa* chromosomal cephalosporinase conferring high-level β-lactam resistance

676   involves three AmpD homologues. Antimicrob Agents Chemother 50:1780–1787.

677   80.   Wu C, Huang Y, Lin Y, Yang T. 2016. Inactivation of lytic transglycosylases increases

678   susceptibility to aminoglycosides and macrolides by altering the outer membrane

679   permeability of *Stenotrophomonas maltophilia*. Antimicrob Agents Chemother 60:3236–

680   3239.

681   81.   Yin J, Sun Y, Sun Y, Yu Z, Qiu J, Gao H, Oliver A. 2018. Deletion of lytic

682   transglycosylases increases β-lactam resistance in *Shewanella oneidensis*. Front

683   Microbiol 9:1–10.

684   82.   Zielińska A, Billini M, Möll A, Kremer K, Briegel A, Izquierdo Martinez A, Jensen GJ,

685   Thanbichler M. 2017. LytM factors affect the recruitment of autolysins to the cell division

686   site in *Caulobacter crescentus*. Mol Microbiol 106:419–438.

687   83.   Lamers RP, Nguyen UT, Nguyen Y, Buensuceso R, Burrows LL. 2015. Loss of

688   membrane-bound lytic transglycosylases increases outer membrane permeability and

689   β-lactam sensitivity in *Pseudomonas aeruginosa*. Microbiology 6:879–895.

690   84.   Cho H, Uehara T, Bernhardt TG. 2014. Beta-lactam antibiotics induce a lethal

691  malfunctioning of the bacterial cell wall synthesis machinery. Cell 159:1300–1311.

692 85. Clough SJ, Bent AF. 1998. Floral dip: A simplified method for *Agrobacterium*-mediated

693  transformation of *Arabidopsis thaliana*. Plant J 16:735–743.

694 86. Howell M, Aliashkevich A, Sundararajan K, Daniel JJ, Lariviere PJ, Goley ED, Cava F,

695  Brown PJB. 2019. *Agrobacterium tumefaciens* divisome proteins regulate the transition

696  from polar growth to cell division. Mol Microbiol 111:1074–1092.

697 87. Howell M, Daniel JJ, Brown PJB. 2017. Live cell fluorescence microscopy to observe

698  essential processes during microbial cell growth. JoVE e56497.

699 88. Ducret A, Quardokus EM, Brun Y V. 2016. MicrobeJ, a tool for high throughput bacterial

700  cell detection and quantitative analysis. Nat Microbiol 1:16077.

701 89. Michniewicz M, Frick EM, Strader LC. 2015. Gateway-compatible tissue-specific vectors

702  for plant transformation Plant Biology. BMC Res Notes 8:6–13.

703 90. Figueroa-Cuilan W, Daniel JJ, Howell M, Sulaiman A, Brown PJB. 2016. Mini-Tn7

704  insertion in an artificial attTn7 site enables depletion of the essential master regulator

705  ctrA in the phytopathogen *Agrobacterium tumefaciens*. Appl Environ Microbiol 82.

706 91. Ikeda RA, An Chi Lin, Clarke J. 1992. Initiation of transcription by T7 RNA polymerase

707  at its natural promoters. J Biol Chem 267:2640–2649.

708 92. Garcia PS, Simorre JP, Brochier-Armanet C, Grangeasse C. 2016. Cell division of

709  Streptococcus pneumoniae: think positive! Curr Opin Microbiol 34:18–23.

 TABLES

712 **Table 1.** Strains and plasmids used in this study.
713

Strain or plasmid	Relevant characteristics	Reference/Source
Source Plasmids		
pSRKKm-Plac-sf*gfp*	pSRKKm vector containing lacl^q and *lac* promoter with sfGFP	(90)
pSRKKm-Pcym-sfGfp	pSRKKm vector containing lac promoter flanked by *cuO* operator site along with the *cymR* repressor gene	This study
pSRKKm-T7-sf*gfp*	pSRKKm vector containing T7 and *lac* promoters with sfGFP	(91)
pNTPS139	Km^r; Suicide vector containing *oriT* and *sacB*	D. Alley
Plasmids		
Deletion plasmids		
pNTPS139 *Δatu3077*	Km^r Suc^s; deletion plasmid for *atu3077*	This study
pNTPS139 *Δatu3078*	Km^r Suc^s; deletion plasmid for *atu3078*	This study
pNTPS139 *Δatu0092*	Km^r Suc^s; deletion plasmid for *atu0092*	This study
pNTPS139 *Δatu2122*	Km^r Suc^s; deletion plasmid for *atu2122*	This study
pNTPS139 *Δatu3779*	Km^r Suc^s; deletion plasmid for *atu3779*	This study

pNTPS139 Δatu2113	Km^r Suc^s; deletion plasmid for *ampD*	This study
E. coli strains		
DH5α	Cloning strain	Life Technologies
S17-1	Sm^r; RP4-2, Tc::Mu,Km-Tn7, for plasmid mobilization	(92)
A. tumefaciens strains		
C58	Nopaline type strain; pTiC58; pAtC58	
GV3101	C58-derived; pTiC58DT-DNA; strain for *Agrobacterium* mediated transformation of dicots	John Walker Lab
FC2452	Deletion strain for *ΔmltA (atu0009)* in C58	This study
FC2444	Deletion strain for *ΔmltB1 (atu0092)* in C58	This study
FC2465	Deletion strain for *ΔmltB2 (atu2122)* in C58	This study
FC2487	Deletion strain for *ΔmltB3 (atu3779)* in C58	This study
FC2446	Deletion strain for *Δslt1 (atu1022)* in C58	This study
FC2448	Deletion strain for *Δslt2 (atu2112)* in C58	This study
FC2450	Deletion strain for *Δslt3 (atu2117)* in C58	This study
C58 *ΔtetRA*::a-*att*Tn7	Replacement of the *ΔtetRA* locus with an artificial *att*Tn7 site	(90)
C58 *ΔtetRA*::a-*att*Tn7 *Δatu3077*	Deletion strain for *ΔampC*	This study

C58 *ΔtetRA*::a-*att*Tn7 *Δatu3078*	Deletion strain for *ΔampR*	This study
C58 *ΔtetRA*::a-*att*Tn7 *ΔampD*	Deletion strain for *ΔampD*	This study
C58 *ΔtetRA*::a-*att*Tn7 *ΔampD ΔampC*	Deletion strain for *ΔampDΔampC*	This study
C58 *ΔtetRA*::a-*att*Tn7 *ΔampD ΔampR*	Deletion strain for *ΔampDΔampR*	This study
C58 GV3101 *Δmltb3*	Deletion strain for *ΔmltB3*	This study

715 **Table 2**. Synthesized DNA primers used in this study.
716

Synthesized DNA	Sequence
atu3077 P1	5'-GCGGCGACTAGTAAACGGATGCCGCTTTTGAAATGC-3'
atu3077 P2	5'-AAGCTTGGTACCGAATTCGCGATTAAATTTCATCTTTCGTGT-3'
atu3077 P3	5'-GAATTCGGTACCAAGCTTGCGCTCGAAAAGGCGCAATAA-3'

atu3077 P4	5'-GTCGTCGGATCCAGATAACTCGGCACACGCCCA-3'
atu3077 P5	5'-CTGCGCCGCCGGTGAAACGCCCGC-3'
atu3077 P6	5'-TGTGCCGGAGGCGCTTGCGATCGC-3'
atu3078 P1	5'-GTCGTCACTAGTGGGTTTTCCCTTCATGGACGG-3'
atu3078 P2	5'-AAGCTTGGTACCGAATTCAAATTGCCGAACCATTCAAGACCT-3'
atu3078 P3	5'-GAATTCGGTACCAAGCTTGAGACCATCGGAACGGCGTGA-3'
atu3078 P4	5'-GTCGTCGGATCCCCTGTTTGATGCTTTTTATCGCGC-3'
atu3078 P5	5'-CCAAGCGCCGGGAAAAGCGTGTCC-3'
atu3078 P6	5'-GGCATCGTCTGCGTGGTGTTCGTC-3'
atu2113 P1	5'-GTCGTCACTAGTCAGATTTTCGATTTCCCGGACGAA-3'
atu2113 P2	5'-AAGCTTGGTACCGAATTCCGAACATTCTTTCATGCGACG-3'
atu2113 P3	5'-GAATTCGGTACCAAGCTTCTGCCGAGATTTTCGGCCGCCTGA-3'
atu2113 P4	5'-GTCGTCGGTACCCCACCGAAACCACGGCATGCGCCA-3'
atu2113 P5	5'-GCGCAATCGGCGACGCGG-3'

atu2113 P6	5'-GTCGTCTGCTGCACACTTCGCCGC-3'
atu0009 P1	5'-AAAAAGCTTAACGCATCTTCTAGCCTTGCG-3'
atu0009 P2	5'-TATTCATTGCTCGGATTCGG-3'
atu0009 P3	5'-GCAATGAATATATCGGCGATGAAAGGC-3'
atu0009 P4	5'-AAAGGATCCGAAAGAACAATTCCTCCGC -3'
atu0009 P5	5'-TTTCGGCGACCTATGACAAGGACGG-3'
atu0009 P6	5'-TTACCAGTTTGCGGAACGCTGGG-3'
atu0092 P1	5'-AAACTGCAGATTCTTGCCCTGATGCCCATTGTCGC-3'
atu0092 P2	5'-AGACCGAATATCGTCTTTTAATGCTGGTCGG-3'
atu0092 P3	5'-ATTCGGTCTCCTCTTGGATGG-3'
atu0092 P4	5'-AAAGGATCCAAGTCGAGATCGACTGAGCCC-3'
atu0092 P5	5'-AATATGTCCGCCACAACCATCGTCGC-3'
atu0092 P6	5'-ATTACATCACAGACCGCCTCTCCG-3'
atu2122 P1	5'-AAAGAATTCAATCATCAGGGTTCCAATGCGG-3'
atu2122 P2	5'-TAGTGCGATTTTCCTCGATAGGTTGTTGGC-3'
atu2122 P3	5'-ATCGCACTACCGGGCCTTTAATCTATCGG-3'

atu2122 P4	5'-AAAAAGCTTATAATGACGTCTTTGAACGC-3'
atu2122 P5	5'-TATACCGCAACCGGCGTCGTACCCG-3'
atu2122 P6	5'-TTTCTGTGATGCGGTGCAGCACGG-3'
atu3779 P1	5'-AAACTGCAGTAGAAATTCGACGGCGCCG-3'
atu3779 P2	5'-TTTCGATTGCGAAAACGCATCGGGCG-3'
atu3779 P3	5'-GCAATCGAAATAATGTGCCGGCGAATTCGG-3'
atu3779 P4	5'-AAAGGATCCTTGGCCGTTCATGTCGTAGCC-3'
atu3779 P5	5'-TTTCGGAACTGCCTTGGTGGCGG-3'
atu3779 P6	5'-ATTCCCGGCCGGAACTACCATCGC-3'
atu2112 P1	5'- AAAAAGCTTTATTTCGTCTTCGAGGATGGG-3'
atu2112 P2	5'- GATGATCGATATGACAGAAACAGTGAAATGGC-3'
atu2112 P3	5'-ATCGATCATCTGCGAAATTGCG-3'
atu2112 P4	5'-AAAGAATTCATATCGTCCTCGGTTTCCGC-3'
atu2112 P5	5'- AAAAAGCTTTATTTCGTCTTCGAGGATGGG-3'
atu2112 P6	5'- TTTGCACCGAAAGATGCCGCG-3'
atu1022 P1	5'-AAAAAGCTTTTATGCGCTTTGACCAGCGCACCC-3'

atu1022 P2	5'-AATCCCCAGACTGTCTTTTTCATGCCG-3'
atu1022 P3	5'-TGGGGATTATTCAGGCACGGGCTAGCC-3'
atu1022 P4	5'-AAAGAATTCAATACGCTCTTCAACTCCATCCG-3'
atu1022 P5	5'- AAAAGCGACGTCGCTCGCG-3'
atu1022 P6	5'-AAACTACTACGACGAAGACGGTCAGG-3'
atu2117 P1	5'-AAAGAATTCAAGAGAATGTCTGGACAGGCGTGGC-3'
atu2117 P2	5'-ATTTTCAATCTAGAGTTGCCGCCTTCGTTATGCC-3'
atu2117 P3	5'- TTGAAAATGCTGCGGCTCCC-3'
atu2117 P4	5'-AAAAAGCTTTAGATGTTCCTGTCGAACACCG-3'
atu2117 P5	5'-ATACAGGCGCATCGCGGCC-3'
atu2117 P6	5'-AAGGCGAGGTGGTCACTGACC -3'
atu3093 P1	5'-GCTGCAACTAGTGCGCCACAGCCCATCTCG-3'
atu3093 P2	5'-AAGCATGGTACCGAATTCGCGGCTTGTTGATATTCCT-3'
atu3093 P3	5'-GAATTCGGTACCATGCTTTAGCAGCGGCGGGCGCCGGA-3'
atu3093 P4	5'-GCACGTAAGCTTGCGGTGATCTTGATGAT-3'
atu3093 P5	5'-GAGCTTCTCGGGCAATTCCG-3'

atu3093 P6	5'-GCCTCACGAAGCCGCACGATC-3'

FIG 1. The AmpC-AmpR operon is responsible for induced ampicillin resistance in

***Agrobacterium tumefaciens* C58. (A)** Operon organization and proposed ampicillin

resistance mechanism (53). Briefly, in the absence of β-lactams such as ampicillin (AMP),

ampC expression is repressed by AmpR. AmpR-mediated repression is maintained as long

as the AmpR-inactivating ligand, UDP-GM-Pentapeptide, is bound to AmpR ("P*ampC* OFF").

In contrast, the presence of ampicillin (AMP) increases the pools of AmpR-activating ligands,

or the cell wall degradation fragments (anhydro modification is depicted by ring), which are

known to displace AmpR-inactivating ligands. As a result, the increase of AmpR-activating

ligands activate AmpR and *ampC* is transcribed ("P*ampC* ON"). **(B)** Growth of WT *A.*

tumefaciens, ΔampC, and *ΔampR* in the absence (No AMP) and presence of ampicillin 25 or

50 µg/ml (AMP 25, AMP 50) for 24 h (n = 1, 2 replicates). **(C)** Quantitative analysis of phase-

contrast microscopy of exponentially-growing strains in absence (No AMP) or presence of

ampicillin 25 ug/ml (AMP 25). % Phenotype was calculated by counting the number of cells

displaying one of the phenotypes indicated (1 cell = 1 phenotype) and dividing it by the total

number of cells per strain. **(D)** Ampicillin susceptibility assay was performed via spotting

dilutions. Briefly, indicated strains were cultured overnight (ON) on LB medium, serially

diluted, spotted on LB solid medium containing no ampicillin (No AMP) or ampicillin 25 µg/ml

(AMP 25), and incubated at 28°C for 36 h before imaging. **(E)** Disc susceptibility was

performed on lawn of indicated strains grown on LB plates for 24 h at 28°C (n = 2). AMP 10 =

disc containing 10 µg/ml ampicillin, AMP 10/SUL 10 = disc containing 10 µg/ml ampicillin and

10 µg/ml sulbactam, a broad-spectrum β-lactamase inhibitor. Data represent the mean (±

standard deviation, SD) of three independent experiments. **** = $P < 0.0001$; *** = $P < 0.001$;

** = $P < 0.01$; * = $P < 0.1$; ns = not significant. **(F)** Determination of β-lactamase production

was performed via a nitrocefin assay using cell lysates. "No AMP" or "AMP 25" indicates cells

untreated or treated, respectively, with ampicillin 25 µg/ml for 2 h before the generation of cell

lysates. Data represents one out of two biological replicates (see **Supplemental Fig. 1D**).

Figure 2

FIG 2. Loss of AmpD results in constitutive β-lactamase activity and elevated ampicillin

resistance. (A) Ampicillin susceptibility assay was performed via spotting dilutions. Briefly,

indicated strains were cultured ON at 28°C, serially diluted, spotted on solid medium

containing no ampicillin (No AMP) or ampicillin 25, 80, or 160 µg/ml (AMP 25, 80, 160), and

incubated at 28°C for ~40 h before imaging. **(B)** Growth of WT *A. tumefaciens* and *ΔampD* in

the absence (No AMP) and presence (AMP) of various concentrations of ampicillin 25, 100

µg/ml (AMP 25, AMP 100) for 24 h (n=1, 2 replicates). **(C)** Quantitative analysis of phase-

contrast microscopy of exponentially-growing strains treated with ampicillin 25 µg/ml (AMP

25). % Phenotype was calculated by counting the number of cells displaying one of the

indicated phenotypes (1 cell = 1 phenotype). **(D)** Determination of β-lactamase production

was performed via a nitrocefin assay using cell lysates. "No AMP" or "AMP 25" indicates cells

untreated or treated, respectively, with ampicillin 25 µg/ml for 2 h before the generation of cell

lysates. *ΔampD* lysates were normalized based on total protein content (7.5 µg/ml) and

subsequently diluted to 1:5 as the rate of nitrocefin hydrolysis was significantly faster than the

controls (WT, WT AMP 25). Finally, data represents one out of two biological replicates (see

Supplemental Fig. 2). **(E)** Disc susceptibility was performed on lawn of indicated strains

grown on LB plates for 24 h at 28°C. Blank = no antibiotic disc, AMP 10 = disc containing 10

µg/ml ampicillin, AMP 10/SUL 10 = disc containing 10 µg/ml ampicillin and 10 µg/ml

sulbactam, a broad-spectrum β-lactamase inhibitor. Data represent the mean (± standard

deviation, SD) of three independent experiments. **** = $P < 0.0001$; *** = $P < 0.001$; ** = $P <$

0.01; * = $P < 0.1$; ns = not significant.

Figure 3

FIG 3. AmpC is the β-lactamase that confers *ΔampD* elevated ampicillin resistance. **(A)**

Ampicillin susceptibility assay was performed via spotting dilutions. Briefly, indicated strains

were cultured ON at 28°C, serially diluted, spotted on solid medium containing no ampicillin

(No AMP) or ampicillin 25 µg/ml (AMP 25) and incubated at 28°C for ~40 h before imaging.

(B) Quantitative analysis of phase-contrast microscopy of exponentially-growing strains

untreated (No AMP) or treated with ampicillin 25 µg/ml (AMP 25). % Phenotype was

calculated by counting the number of cells displaying one of the indicated phenotypes (1 cell =

1 phenotype). **(C)** Determination of β-lactamase production was performed via a nitrocefin

assay using cell lysates. "No AMP" or "AMP 25" indicates cells untreated or treated,

777   respectively, with ampicillin 25 µg/ml for 2 h before the generation of cell lysates. Data

778   represents one out of two biological replicates (see **Supplemental Fig. 3**).

Figure 4

FIG 4. MtlB3 is required for ampicillin resistance in *A. tumefaciens*. (A) Phase-contrast

microscopy of exponentially-growing strains treated with ampicillin 25 µg/ml (AMP 25) for 2 h.

(B) Quantitative analysis of phase-contrast microscopy of exponentially-growing strains

treated with ampicillin 25 µg/ml (AMP 25) for 2 h. % Phenotype was calculated by counting

the number of cells displaying one of the phenotypes indicated (1 cell = 1 phenotype) and

diving it by the total number of cells per strain. **(C)** Growth of WT *A. tumefaciens* and *ΔmltB3*

in the absence (No AMP) and presence of ampicillin 25 µg/ml (AMP 25) for 24 h (n = 1, 2

replicates). **(D)** Determination of β-lactamase production was performed via a nitrocefin assay

using cell lysates. "No AMP" or "AMP 25" indicates cells untreated or treated, respectively,

with ampicillin 25 µg/ml for 2 h before the generation of cell lysates. Data represents one out

of two biological replicates (see **Supplemental Fig. 4**).

Figure 5

FIG 5. *A. tumefaciens* GV3103 *ΔmltB3* can be used to transform plants efficiently and

bacteria can be removed using low concentrations of ampicillin. (A) Ampicillin

susceptibility assay was performed via spotting dilutions. Briefly, indicated strains were

cultured ON at 28°C, serially diluted, spotted on solid medium containing no ampicillin (No

AMP), ampicillin 25 μg/ml (AMP 25), or carbenicillin 15 μg/ml (CARB 15), and incubated at

28°C for ~40 h before imaging. **(B)** Determination of β-lactamase production was performed

802   via a nitrocefin assay using cell lysates. Data represents one out of two biological replicates

803   (see **Supplemental Fig. 5**). "No AMP" or "AMP 25" indicates cells untreated or treated,

804   respectively, with ampicillin 25 µg/ml for 2 h before the generation of cell lysates. **(C)**

805   Transformation efficiency and **(D)** bacterial loads of seeds transformed with WT *A.*

806   *tumefaciens* GV3101 and GV3101 *ΔmltB3* using the floral dip assay technique (85).

Figure 6

 FIG 6. Working model for *A. tumefaciens* ampicillin resistance. Bifunctional PBPs extend

 the cell wall through the transglycosylation (linking of carbohydrates) and transpeptidation

 reactions (linking of peptide stems). β-lactams such as ampicillin (purple stars) target the

transpeptidase domain of Penicillin-Binding Proteins (PBPs) (**Fig S6**, Step 1-2), leading to a

block in bacterial cell growth. In *A. tumefaciens*, treatment with ampicillin leads to induction of

β-lactamases. Inactivation of the lytic transglycosylase MltB3 results inhibition of β-lactamase

derepression and lysis suggesting that MltB3 is likely required for the generation of cell wall

degradation products (**Fig S6**, Step 3). Similarly to treatment with β-lactams, where a block in

cell growth leads to an increase in cell wall degradation products (**Fig S6**, Step 5), inactivation

of anhydro amidases such as AmpD increases the poll of this molecules leading to β-lactam

resistance. In *A. tumefaciens* inactivation of AmpD leads to derepression of β-lactamases and

ampicillin resistance (**Fig S6**, Step 9). Both AmpC, an inducible β-lactamase that is under the

transcriptional control of AmpR, and AmpR seem to be responsible for the derepression

observed in *ΔampD.* Thus, our working model suggests that upon ampicillin exposure, a block

in growth leads to increased activity of MltB3. An increase in cell wall degradation products

leads to induction of AmpC expression by AmpR and the presumed translocation of AmpC to

the periplasm resulting in ampicillin resistance.