1 **Title:** Induction of AmpC-mediated β-lactam resistance requires a single lytic transglycosylase
2 in *Agrobacterium tumefaciens*
3

Authors:

Wanda M. Figueroa-Cuilan[1*], Matthew Howell[1&], Christopher Richards[1], Amelia Randich[1^],

Akhilesh K. Yadav[2$], Felipe Cava[2], and Pamela J.B. Brown[1#]

Affiliations:

[1]Division of Biological Sciences, University of Missouri-Columbia, Columbia, Missouri, 65202

[2]Laboratory for Molecular Infection Medicine, Department of Molecular Biology, Umeå University,

Umeå, Sweden

Running title: AmpC-mediated β-lactam resistance of *A. tumefaciens*

[#]**Corresponding author:**

brownpb@missouri.edu

(573) 884-0384

Present addresses:

*Department of Biological Chemistry, Johns Hopkins University School of Medicine,

Baltimore, Maryland 21205

&Department of Biology, Westminster College, Fulton, Missouri 65251

$Analytical Chemistry Division, CSIR-Indian Institute of Toxicology Research, Lucknow-226001, India

^Department of Biology, The University of Scranton, Scranton, Pennsylvania,18510

ABSTRACT

The remarkable ability of *Agrobacterium tumefaciens* to transfer DNA to plant cells has allowed the generation of important transgenic crops. One challenge of *A. tumefaciens-* mediated transformation is eliminating the bacteria after plant transformation to prevent detrimental effects to plants and the release of engineered bacteria to the environment. Here we use a reverse genetics approach to identify genes involved in ampicillin resistance with the goal of utilizing these antibiotic-sensitive strains for plant transformations. We show that treating *A. tumefaciens* C58 with ampicillin led to increased β-lactamase production, a response dependent on the broad-spectrum β-lactamase AmpC and its transcription factor AmpR. Loss of the putative *ampD* orthologue, *atu2113,* led to constitutive production of AmpC-dependent β-lactamase activity and ampicillin resistance. Finally, one cell wall remodeling enzyme, MltB3, was necessary for the AmpC-dependent β-lactamase activity and its loss elicited ampicillin and carbenicillin sensitivity in the *A. tumefaciens* C58 and GV3101 strains. Furthermore, GV3101 *ΔmltB3* transforms plants with comparable efficiency to wildtype but can be cleared with sub-lethal concentrations of ampicillin. The functional characterization of the genes involved in the inducible ampicillin resistance pathway of *A. tumefaciens* constitutes a major step forward in efforts to reduce the intrinsic antibiotic resistance of this bacterium.

IMPORTANCE

Agrobacterium tumefaciens, a significant biotechnological tool for production of transgenic plant lines, is highly resistant to a wide variety of antibiotics, posing challenges for various applications. One challenge is the efficient elimination of *A. tumefaciens* from transformed

plant tissue without using levels of antibiotics that are toxic to the plants. Here, we present the functional characterization of genes involved in β-lactam resistance in *A. tumefaciens.* Knowledge about proteins that promote or inhibit β-lactam resistance will enable the development of strains to improve the efficiency of *Agrobacterium*-mediated plant genetic transformations. Effective removal of *Agrobacterium* from transformed plant tissue has the potential to maximize crop yield and food production, improving the outlook for global food security.

Keywords

Antibiotic resistance, *Agrobacterium tumefaciens*, β-lactamases, anhydro amidases, membrane-bound lytic transglycosylases, ampicillin, plant transformation

INTRODUCTION

Rhizobiaceae is a family of bacteria that include soil-dwelling and plant associated bacteria. While some species of this family have the ability to establish symbiotic relationships with plants, others are pathogenic such as the genus *Agrobacterium*. Members of this genus are responsible for a number of diseases, including the cane gall disease (*Agrobacterium rubi*), hairy root disease (*Agrobacterium rhizogenes*), crown gall disease of grapes (*Agrobacterium vitis*), and crown gall disease to flowering plants and woody shrubs (*Agrobacterium tumefaciens*) (1–5). In nature, *A. tumefaciens* causes crown gall by adhering to wounded plants and injecting a section of a bacterial DNA plasmid (Transfer DNA [T-DNA]) that integrates into the plant chromosomes (1–3, 6–11). Expression of genes on the T-DNA segment causes the plant to produce custom energy sources that only *Agrobacterium* can

use (9, 10). The increased abundance of energy sources within plant cells leads to their over-

proliferation and eventual gall formation (6). Gall formation on plants and trees leads to crop

damage, and significant economic losses have been attributed to this issue every year (2, 3).

While the genus *Agrobacterium* exhibits pathogenicity against plants, the natural ability

of *Agrobacterium* to transfer DNA to plants has been exploited to produce transgenic plants

through genetic engineering (6, 9, 11–13). However, one challenge for *A. tumefaciens*-

mediated plant transformations is the elimination of the bacteria from the transformed plant

tissue. Elimination of recombinant *A. tumefaciens* from plant tissues is crucial to prevent

detrimental effects to plants and to reduce the risk of releasing engineered bacteria into the

environment (14–16). β-lactam antibiotics are frequently applied during plant transformations

to eliminate *A. tumefaciens* from plant tissues and are preferred over other classes of

antibiotics (17–19). Because β-lactams target cell wall synthesis, a process unique to

bacteria, and they are less toxic to eukaryotic plant cells than antibiotics that inhibit protein or

nucleic acid synthesis (20, 21). However, the natural resistance of *A. tumefaciens* to β-

lactams can be only overcome with toxic levels (~200-1000 mg/L), which has been shown to

cause embryogenic tissue necrosis, or to affect plant tissue growth and regeneration rates in

a wide variety of plants (17, 18, 22–28). Moreover, depending on the concentration and class

of β-lactam, clearing *Agrobacterium* from embryos can take up to 60 days, yet, in some

cases, complete elimination of *A. tumefaciens* is not achieved (29). Thus, currently, there is a

need for the identification and understanding of regulatory pathways and enzymes involved in

β-lactam resistance in *A. tumefaciens.* Functional characterization of bacterial enzymes

involved in β-lactam resistance will permit the development of tools that could improve the

efficiency of plant genetic transformations, and therefore maximize crop yields and food

production.

β-lactam antibiotics target the bacterial cell wall by inhibiting the activity of Penicillin

Binding Proteins (PBPs); the enzymes involved in the synthesis of the bacterial peptidoglycan

(PG) cell wall (30–38). The bacterial PG cell wall is an essential polymer consisting of

alternating N-acetylglucosamine (Glc$N$Ac) and N-acetylmuramic acid (Mur$N$Ac) sugars

crosslinked through peptide bridges (39–44). Because the PG cell wall is a covalently-

enclosed polymer, its expansion not only requires cell wall synthesis but also remodeling. Cell

wall remodeling is mediated by PG degradation enzymes such as the lytic transglycosylases

(LT) (45–48). To allow cell wall expansion, LTs cleave between the Mur$N$Ac and Glc$N$Ac

sugar strands resulting in the formation of 1,6-anhydroMurNAc Glc$N$Ac on glycan strands and

the liberation of 1,6-anhydromuropeptides (AnhMP) cell wall degradation fragments. The

liberated AnhMP fragments are transported to the bacterial cytoplasm for cell wall recycling

(42, 49, 50). In the cytoplasm, the recycling of AnhMP fragments keeps the concentration of

these products low (51, 52). However, cell wall stressors such as treatment with β-lactam

antibiotics or mutations that inhibit the cell wall recycling pathway result in the accumulation of

AnhMP cell wall degradation fragments and derepression of β-lactamases (34, 50, 53–56). In

bacteria including *Pseudomonas aeruginosa* and *Enterobacter cloaceae,* the AnhMP cell wall

degradation fragments are transcriptional activators of inducible β-lactamases, which are

enzymes that cleave and inactivate β-lactam antibiotics (57–63).

In the soil environment, many soil microorganisms produce antibiotics to compete for

survival, selecting for intrinsic resistance pathways in soil pathogens. For example, the

genomes of many soil bacteria contain β-lactamases, such as the cephalosporinase AmpC

(34, 64). As a cephalosporinase, AmpC is known to destroy β-lactam antibiotics including

monobactams, cephalosporins, and penicillins (34). The AmpC consensus protein sequence

consists of a signal sequence for periplasmic transport and a β-lactamase catalytic domain

(Fig. S1A). The regulation of AmpC expression varies across bacteria. In *Escherichia coli*,

AmpC is a non-inducible β-lactamase that is expressed at low levels and regulated by a

promoter and a growth rate-dependent attenuator mechanism (65–67). In contrast, in *P.*

aeruginosa and some enterobacteria, AmpC is normally expressed at low levels, but is

inducible and can be derepressed during exposure to β-lactams (34, 60, 68). In these cases,

AmpC expression is regulated by AmpR, a LysR-type transcriptional regulator found in an

operon with AmpC (60). AmpR consists of two domains; a Helix-Turn-Helix DNA-binding

domain (DBD) that binds the intergenic region between AmpC and AmpR, and a LysR

effector-binding domain (EBD), which contains the regulatory region of AmpR (**Fig. 1A, Fig.**

S1A) (57). AmpR is a bifunctional transcriptional regulator that controls both the activation

and repression of AmpC (60). The induction mechanism of AmpC by AmpR in response to β-

lactams is linked to bacterial cell wall synthesis, remodeling, and recycling (34, 50, 53, 57, 62,

69). Indeed, the AnhMur cell wall degradation fragments released by LTs during cell growth,

are AmpR-activating molecules. In contrast, cell wall building blocks such as UDP-Glc*N*Ac

Mur*N*Ac pentapeptides bind to AmpR and repress *ampC* transcription **(Fig. 1A)**. A block in

bacterial cell wall synthesis after exposure to β-lactams results in accumulation of AnhMur cell

wall degradation products in the bacterial cytoplasm, displacement of the AmpR repressor

UDP-Glc*N*Ac Mur*N*Ac pentapeptide, activation of AmpR, and transcription of *ampC* **(Fig. 1A).**

The genus *Agrobacterium* is naturally resistant to β-lactams and the molecular basis for

this resistance is poorly understood. A study that screened for β-lactamase production in *A.*

tumefaciens detected cephalosporinase production (70) and identified one putative

cephalosporinase, an AmpC homolog (*atu3077),* in the *A. tumefaciens* genome. In addition,

similar to *Enterobacteriaceae* and *Pseudomonadales, ampC* is found in an operon with *ampR*

in most *Agrobacterium* genomes **(Fig. 1A)**. Here, we sought to determine if AmpC is a

functional β-lactamase*,* if AmpC is inducible, and if the natural ampicillin resistance observed

in *A. tumefaciens* is dependent on AmpC.

We present a functional characterization of proteins involved in intrinsic ampicillin

resistance in *A. tumefaciens.* We find that AmpR is required for AmpC-dependent β-

lactamase activity and that loss of the anhydro-amidase AmpD (*atu2113*, misannotated as an

AmiD homolog in the genome (71)) leads to increased resistance to ampicillin, a process

dependent on AmpC. We suggest that AmpD is required for proper recycling of cell wall

degradation products and its loss results in the accumulation of cell wall degradation products

and activation of AmpC by AmpR. Furthermore, we find that a single LT, the membrane-

bound lytic transglycosylase B3 (MltB3), is necessary for AmpC-dependent β-lactamase

activity and its loss leads to ampicillin sensitivity in the *A. tumefaciens* C58 and GV3101

strains. Finally, transformation of *Arabidopsis thaliana* utilizing a *ΔmltB3* GV3101 strain

requires significantly lower concentrations of ampicillin while exhibiting similar WT

transformation efficiency. This work underscores the significance of understanding the β-

lactam resistance pathway of *A. tumefaciens* with the aim of expanding tools for the *A.*

tumefaciens-mediated transformations.

RESULTS AND DISCUSSION

The AmpC-AmpR operon is responsible for inducible ampicillin resistance in *A.*

tumefaciens **C58**

To begin our characterization*,* we first assessed the susceptibility of *A. tumefaciens* to

different concentrations of ampicillin near the minimum inhibitory concentration (MIC) reported

for *A. tumefaciens* on solid and liquid media (72). We found that cells grown in LB with 25 or

50 μg/mL ampicillin (AMP 25 or AMP 50, respectively) for 24 h displayed slow growth in liquid

medium in comparison to cells grown in LB without ampicillin (No AMP) **(Fig. 1B)**. To better

understand the cause of this growth defect, we performed phase-contrast microscopy of cells

treated with AMP 25 for 2 h **(Fig. S1B)**. We found that treatment with AMP 25 causes a

significant increase in the median cell length **(Fig. S1C)** and that 23.5% of the cells were

undergoing cell lysis **(Fig. 1C, Fig. S1B)**, confirming that the bactericidal effect of AMP 25 on

WT *A. tumefaciens* is the cause of the overall decrease in optical density*.* Similarly, WT cells

grown on AMP 25 solid medium for 36 h have a viability defect in comparison to WT cells

grown in LB No AMP **(Fig. 1D)**. The increased sensitivity of WT *A. tumefaciens* to ampicillin in

the presence of sulbactam, a broad spectrum β-lactamase inhibitor, suggests that β-

lactamase production is responsible for the observed ampicillin resistance **(Fig. 1E)**. Finally,

to determine if β-lactamase production is induced, we treated WT cells with AMP 25 for 2 h,

generated whole cell lysates, and performed nitrocefin assays on total protein content **(Fig.**

1F). Nitrocefin is a chromogenic substrate related to the cephalosporins that undergoes a

color change when is hydrolyzed by β-lactamases (73). After treatment of WT cells with AMP

25 for 2 h, the activity of β-lactamases is readily detected in lysates using nitrocefin assays

(Fig. 1F). Together, these results suggest that *A. tumefaciens* C58 β-lactamase production is

induced in the presence of β-lactams such as ampicillin. To assess the contributions of

putative enzymes involved in ampicillin resistance, we employed a reverse genetics approach

(74).

The *ampC* ortholog of *A. tumefaciens* C58 (*atu3077)* is present on the linear

chromosome and encodes the only putative inducible β-lactamase in the genome of *A.*

tumefaciens C58 (34). *ampC* is syntenic with *ampR* and the *A. tumefaciens* AmpC and AmpR

proteins are 74.7% and 85.9% similar to their respective orthologs from *P. aeruginosa***.** To

determine the role of AmpC, we deleted *ampC* (*atu3077*) from the *A. tumefaciens* C58

genome. Deletion of *ampC* does not impact cell growth and cell viability **(Fig. 1B, 1D)** or cell

morphology **(Fig. S1B)**, beyond a slight increase in cell length **(Fig. S1C).** To pinpoint the

contribution of *ampC* to ampicillin resistance, we assessed the growth dynamics of *ΔampC*

cells in the presence of AMP in liquid **(Fig. 1B)**. *ΔampC* cells treated with AMP 25 or AMP 50

for 24 h show a severe growth defect indicating that AmpC contributes to ampicillin

resistance. Similarly, deletion of *ampC* results in severe growth viability defect on solid

medium containing AMP 25 **(Fig 1D).** Production of plasmid-encoded AmpC in *ΔampC*

restores growth and viability in the presence of AMP 25 **(Fig 1D)**. In addition, *ΔampC* cells

treated with AMP 25 exhibit cell division defects (34.8%) and cell lysis (15.8%) **(Fig. 1C, Fig.**

S1B). The increased cell length of *ΔampC* treated with AMP 25 is likely due to the observed

cell division defects **(Fig. S1B-C, Fig 1C).**

To confirm that ampicillin resistance is mediated by the AmpC β-lactamase, we used

the disc diffusion assay to compare resistance levels to ampicillin in the presence and

absence of the broad-spectrum β-lactamase inhibitor, sulbactam **(Fig. 1E)**. As expected,

ΔampC leads to increased sensitivity to ampicillin and the presence of sulbactam does not

result in large increases in the zone of growth inhibition **(Fig. 1E).** Furthermore, monitoring the

rates of nitrocefin hydrolysis show that production of β-lactamase is readily detected in WT

cells treated with AMP 25 but is undetectable in *ΔampC* following AMP25 treatment **(Fig. 1F)**.

Together, these observations suggest that the natural resistance to ampicillin depends on the

presence of AmpC, which functions as an inducible β-lactamase.

We hypothesized that if transcription of *ampC* is strictly controlled by AmpR, deletion of

ampR should mimic deletion of *ampC.* To test this hypothesis, we deleted *ampR* (*atu3078*)

from the genome of *A. tumefaciens* C58. Deletion of *ampR* does not impact cell growth **(Fig.**

1B), morphology **(Fig. S1B),** cell length **(Fig. S1C)** or cell viability **(Fig. 1D)** in LB**.** Low

concentrations of ampicillin in either liquid or solid medium are lethal to *ΔampR* and results in

similar cell division defects and cell lysis as *ΔampC* **(Fig. 1B-C, Fig. S1B)**. Production of

plasmid encoded AmpR restores the viability of *ΔampR* on solid medium with AMP25 **(Fig**

1D). Like *ΔampC, ΔampR* fails to produce detectable β-lactamase activity when treated with

AMP 25 **(Fig. 1F).** Together, these results suggest that AmpR and AmpC contribute to

ampicillin resistance in *A. tumefaciens*. Based on agreement with the general mechanism of

characterized AmpR-AmpC pathways, we hypothesize that AmpR is necessary for induction

of the AmpC β-lactamase in the presence of ampicillin.

Loss of AmpD derepresses β-lactamases in *A. tumefaciens* C58.

The finding that AmpC and AmpR are necessary for ampicillin resistance in *A. tumefaciens*

C58 led us to explore how the pools of different cell wall fragments would alter the AmpC-

mediated β-lactamase induction. Similar to exposure to β-lactams, loss of cell wall recycling

amidases has been shown to increase the AmpR-activating fragments (cell wall degradation

fragments) in the cytoplasm, resulting in the transcriptional derepression of *ampC* and β-

lactam resistance (48, 75–78). The genome of *A. tumefaciens* contains one 1,6-anhydro

amidase ortholog, *atu2113*, reannotated here as AmpD. The domain organization of AmpD

consists of the Amidase_2 (Ami_2) catalytic domain and a PG-binding domain (PBD) that

facilitates the interaction with cell wall products **(Fig. S2A)** (45, 48). *A. tumefaciens* AmpD

exhibits 64.4% sequence similarity to AmpDh2, one of three broad-spectrum 1,6-anhydro

amidase AmpDh paralogs found in *Pseudomonas aeruginosa* (54, 56, 77, 79).

Given that in *A. tumefaciens* ampicillin triggers the AmpC-dependent production of β-

lactamases, we hypothesized that if AmpD was an anhydro amidase involved in the recycling

of cell wall degradation fragments, its loss should result in increased AmpR-activating

fragments in the cytoplasm, β-lactamase induction, and ampicillin resistance. First, we found

that *ΔampD* cells exhibit normal cell viability **(Fig. 2A)**, cell growth **(Fig. 2B),** and morphology

(Fig. S2B-C). *ΔampD* is highly resistant to ampicillin **(Fig. 2A)**. Indeed, WT cells spotted on

AMP 160 are not viable, whereas *ΔampD* spotted on AMP 160 only displays a ~10-fold

decrease in viability compared to untreated cells. In contrast, production of *ampD* from an

IPTG-inducible plasmid (+pAmpD) resulted in a 100,000-fold decrease in viability in the

presence of AMP 100 **(Fig. 2A).** In liquid, *ΔampD* cells continue to grow normally, even in the

presence of AMP 100 **(Fig. 2B)** and ampicillin treatment does not trigger obvious

morphological changes or cell lysis **(Fig. S2B-C, Fig. 2C)**. *ΔampD* produces readily

detectable amounts of β-lactamase production in both the presence and absence of ampicillin

(Fig. S2D). The increased zone of inhibition observed in the presence of ampicillin and

sulbactam is consistent with the high level of ampicillin resistance observed in *ΔampD* being

mediated by a β-lactamase **(Fig. 2D)**. Together, these results indicate loss of AmpD leads to

derepression and increased β-lactamase activity. Our findings are consistent with other

bacterial models such as *P. aeruginosa*, where deletion of 1,6-anhydro amidases involved in

the recycling of AmpR-activating ligands leads to increased β-lactamase expression (53–56)

due to the build-up of activating ligands in the cytoplasm.

AmpC is constitutively produced in *ΔampD*

We have shown that AmpC and AmpR are required for ampicillin resistance **(Fig. 1)**, and loss

of AmpD leads to elevated β-lactamase activity and ampicillin resistance in *A. tumefaciens*

C58 **(Fig 2)**. To confirm that AmpC is the β-lactamase produced by the *ΔampD* strain, we

deleted *ampC* or *ampR* in the *ΔampD* background **(Fig. 3)**. In the absence of ampicillin, we

found that *ΔampCΔampD* and *ΔampRΔampD* display normal cell viability, growth, and

morphology **(Fig. 3A-B, Fig S3A-B).** We found that treatment with AMP 25, on either solid

medium or liquid medium, is lethal to *ΔampCΔampD* or *ΔampRΔampD* **(Fig. 3A, Fig. S3A).**

Treatment of *ΔampCΔampD* or *ΔampRΔampD* with AMP 25 for 2 h results in lysis of 22.9%

and 33.8% of the cells in comparison to untreated cells (No AMP), where lysis is not readily

observed **(Fig. 3B, Fig S3B)**. In comparison to *ΔampC* or *ΔampR*, where cell division defects

are observed in >20% of the population, very few *ΔampCΔampD* or *ΔampRΔampD* cells

exhibit cell division defects when treated with AMP 25 **(Fig. 3B).** The low incidence of cell

division defects observed in *ΔampCΔampD* or *ΔampRΔampD* suggests that the activity of

AmpD contributes to the inefficient cell division of *ΔampC* and *ΔampR* cells following

ampicillin treatment. Finally, to assess whether the *ΔampD* strain could induce β-lactamase

production in the absence of *ampC* or *ampR*, we performed nitrocefin assays. *ΔampCΔampD*

and *ΔampRΔampD* fail to produce detectable levels of β-lactamase in the absence or

presence of AMP 25 **(Fig. 3C).** Together, these results suggest that induction of AmpC is the

main cause for the elevated resistance to ampicillin observed in *ΔampD*. This data is

consistent with the current model for β-lactam resistance in *P. aeruginosa*, where the loss of

anhydro amidases leads to accumulation of cell wall degradation products that activate AmpR

leading to derepression of AmpC (54–56, 69).

Absence of MltB3 (*Δatu3779*) leads to a failure of AmpC-dependent induction of β-

lactamases

Lytic transglycosylases (LTs) are likely to function as the enzymes that generate the AmpR-

activating fragments. Different families of LTs have been linked to β-lactam resistance in

several bacterial organisms (80, 81). For instance, in *Caulobacter crescentus* deletion of

sdpA, a soluble LT, led to increased sensitivity to ampicillin (82). In *P. aeruginosa*, loss of

several *mltBs* and/or *slt* led to a decrease in the β-lactam MIC, cell viability, and increased

outer membrane permeability (83, 84). Thus, we sought to determine if LTs contribute to β-

lactam resistance of *A. tumefaciens*. The *A. tumefaciens* genome consists of 8 putative LTs

belonging to 3 families: family 1) the soluble lytic transglycosylases (Slt), 2) membrane-bound

lytic transglycosylase A (MltA), and 3) membrane-bound lytic transglycosylase B (MltB) **(Fig.**

S4A). We found that single deletions of LTs did not affect cell viability, suggesting a wide

redundancy of functions between LTs **(Fig. S4B)**.

Despite the potential for functional redundancy, we found that deletion of a single, family 3,

membrane-bound lytic transglycosylase, MltB3 (*atu3779)*, causes ampicillin hypersensitivity

(Fig. S4B). Treatment of *ΔmltB3* with AMP 25 for 2 h causes cell lysis defect (28.8%) **(Fig.**

4A-B) and results in a severe growth defect (**Fig. 4C**) indicating that MltB3 is required for

ampicillin resistance. Production MltB3 from an IPTG-inducible plasmid (+pMltB3) in *ΔmltB3*

restores viability in the presence of AMP 25 **(Fig S4B)** confirming that MltB is responsible for

this phenotype. Finally, *ΔmltB3* exhibits reduced production of β-lactamase after AMP 25

treatment for 2 hours **(Fig. 4D).** Together, these results suggest that in *A. tumefaciens,* MltB3

is a specialized enzyme that functions in the AmpR-AmpC β-lactamase pathway. These

observations contrast with the *P. aeruginosa* model*,* in which the β-lactam sensitivity of LT

mutants is due to increased outer membrane permeability rather than β-lactamase production

(83)**.**

Plant transformation with *Δmltb3* cells requires a low concentration of ampicillin for

the elimination of bacteria

Next, we sought to determine if the ampicillin sensitive strains of *A. tumefaciens*

constructed in this work are competent for plant transformation. While the *ΔampC* strain

appears to be the ideal mutant for these studies, we took into consideration the impact of the

mutation on the overall fitness of our ampicillin sensitive strains. *ΔampC* and *ΔampR* growth

dynamics are very similar to WT in liquid; however, these strains exhibit a 10-fold viability

defect on solid medium. In contrast, *ΔmltB3* cells growth dynamics mimic that to WT and lyse

quicky in the presence of low concentrations of ampicillin. Thus, we reasoned that the ΔmltB3

allele would enable us to test the transformation efficiency of an otherwise fit but ampicillin

sensitive *A. tumefaciens* strain. To this end, we deleted *mltB3* in *A. tumefaciens* GV3103 and

find that this mutation causes susceptibility to AMP 25 and carbenicillin 15 (CARB 15) **(Fig.**

5A). To confirm that the absence of MltB3 (*ΔmltB3*) prevented the induction of β-lactamase

production in the GV3101 strain after ampicillin treatment, we performed nitrocefin assays

using lysates of cells treated with AMP 25 for 2 h. While β-lactamase activity is readily

detected from WT GV3101-treated cells with AMP 25, the *ΔmltB3* mutant produces relatively

low levels of inducible β-lactamase **(Fig. 5B)**. Together, we conclude that MltB3 is the major

LT in *A. tumefaciens* C58 and GV3101 that contributes to *A. tumefaciens* natural resistance to

ampicillin and other β-lactam antibiotics.

Next, we confirmed that *ΔmltB3 A. tumefaciens* GV3101 effectively transforms *Arabidopsis*

thaliana using the standard floral dip technique (85) with an efficiency comparable to WT **(Fig.**

5C). Next, we asked whether elimination of the bacteria could be performed using low

concentrations of ampicillin (AMP 25). Seeds transformed with WT *A. tumefaciens* GV3101

contain similar bacterial loads compared to seeds transformed with *Δmltb3 A. tumefaciens*

GV3101 when plated on solid media without ampicillin **(Fig. 5D)**. When plated on media

containing AMP 25, seeds transformed with *Δmltb3* exhibit a significant drop in bacterial load.

In contrast, this low level of ampicillin does not reduce the bacterial load of seeds transformed

with WT GV3101 **(Fig. 5D)**. These results demonstrate that the *Δmltb3* GV3101 strain is

useful for the transformation of *Arabidopsis thaliana* and elimination of the bacteria can be

performed using lower concentrations of ampicillin. While ampicillin or carbenicillin are

occasionally used for clearing *Agrobacterium* after transformation, many labs routinely use

expensive antibiotics such as the proprietary blends of Timentin and Augmentin (86, 87). Our

work highlights the ability to clear *ΔmltB* cells using ampicillin, a cost-effective and readily

available antibiotic. Furthermore, the increased sensitivity of this strain to carbenicillin

suggests that bacterial clearance following plant transformation can likely be achieved using

other β-lactam antibiotics. Overall, these data show the potential impact of improved

understanding the cell biology of *A. tumefaciens* to improve genetic engineering approaches.

CONCLUDING REMARKS AND FUTURE PERSPECTIVES

The natural ability of *A. tumefaciens* to transform plants has allowed the production of transgenic crops of incredible economic importance for the past four decades. One challenge of the *A. tumefaciens*-mediated plant transformation is natural resistance of *A. tumefaciens* to antibiotics, which requires toxic concentrations of antibiotics to eliminate *A. tumefaciens* from transformed tissues. Here we show that *A. tumefaciens* induces β-lactamase activity in response to ampicillin exposure. Indeed, induction of β-lactamase activity upon exposure to ampicillin is dependent on the β-lactamase AmpC and the transcription factor AmpR. Moreover, we found that deletion of a single LT, MltB3, sensitizes *A. tumefaciens* to the β-lactams.

We propose that during *A. tumefaciens* growth and remodeling, there is a delicate balance between the synthesis and degradation of the bacterial cell wall. PBPs insert precursor cytoplasmic monomers into the growing cell wall polymer (**Fig 6, steps 1-2**). During remodeling, cell wall hydrolytic enzymes such as such as endopeptidases and LTs, including MltB3, liberate cell wall degradation products, which are transported back into the cytoplasm of *A. tumefaciens* for their recycling (**Fig 6, steps 3-4**). In the cytoplasm, hydrolytic enzymes including L,D-carboxypeptidases (LD-CPases), amidases such as AmpD, and glycosidases limit the pool of cell wall fragments by allowing their recycling **(Fig 6, steps 5-8).** When cell wall degradation fragments accumulate during β-lactam treatment, the AmpR-dependent production of the AmpC β-lactamase is increased **(Fig 6, step 9)**. Overall, the identification

and contributions of genes conferring ampicillin resistance in *A. tumefaciens* will be beneficial

for improving the design of *A. tumefaciens*-mediated genetic engineering.

MATERIALS AND METHODS

Bacterial strains and growth conditions. Unless otherwise indicated, *Agrobacterium*

tumefaciens GV3101, C58, and derived strains were grown in LB medium (10 g/l tryptone, 10

g/l yeast extract and 5 g/l NaCl) at 28°C with shaking and the antibiotic kanamycin was added

at a concentration of 100 µg/ml to maintain plasmids in complementing strains. For

determining bacterial loads after transformation, *Agrobacterium tumefaciens* GV3101 was

grown in minimal medium ATGN (88). *E. coli* DH5α and S17.1 were routinely cultivated in LB

medium at 37°C with shaking and the antibiotic kanamycin was added at a concentration of

50 µg/ml and 25 µg/ml to maintain plasmids in DH5α and S17-1 strains, respectively.

Construction of plasmids and strains*.* All strains and plasmids used are listed in **Table 1**.

Synthesized DNA primers are listed in **Table 2**.

Gene deletions were achieved by allelic exchange and vectors were constructed as

previously described (89). Briefly, 500 base pairs (bp) fragments upstream and downstream

of the gene of interest were amplified from purified C58 genomic DNA using primer pairs

P1/P2; which amplify 500 bp upstream of the gene of interest, and P3/P4; which amplify the

500 bp downstream of the gene of interest. Overlapping PCR was used to merge and amplify

the amplicons generated by P1/P2 and P3/P4, using primer pair P1/P4. The 1,000 bp

amplicon was digested and ligated into the deletion plasmid, pNTPS139. The deletion

plasmids were introduced into *A. tumefaciens* by mating using an *E. coli* S17.1 conjugation

strain to create kanamycin (KAN) resistant, sucrose sensitive primary exconjugants. Deletion

strains were constructed as described previously (74). Briefly, primary exconjugants were

grown overnight at 28°C in ATGN with no selection and plated in ATGN KAN 300 for 48 h at

28°C. Colonies were screened by patching for KAN resistance and sucrose sensitivity. Colony

PCR using primers P5/P4 were used to confirm that recombination took place and at the

region of interest. Next, positive colonies are grown in ATGN at 28°C overnight and plated in

ATSN 5% sucrose. Secondary recombinants were screened by patching for sucrose

resistance and KAN sensitivity. Colony PCR with primers P5/P6 for the respective gene target

was used to confirm deletion. PCR products from P5/P6 primer sets were sequenced to

further confirm deletions.

For the construction of replicating plasmids, the amplicons and pSRKKM-Plac-*sfgfp,* were

digested overnight and ligated overnight at 4°C using NEB T4 DNA ligase and transformed

into *E. coli* DH5α. Plasmids were sequenced to verify content and were introduced into *A.*

tumefaciens by mating using an *E. coli* S17.1 harboring the appropriate plasmid.

Cell viability spot assays*.* The cell viability assay was performed as described (90). For cell

viability spot assays, cultures were grown overnight and diluted to an OD_{600} of 0.05 and

serially diluted in LB up to 10^{-6}. 4 µl of each dilution was spotted onto LB plates and incubated

at 28°C for 36-40 h before imaging. To determine ampicillin antibiotic resistance, *A.*

tumefaciens cultures were grown overnight and diluted to an optical density at 600 nm (OD_{600})

of 0.05 and serially diluted and spotted onto LB plates containing AMP 25, AMP 100, or AMP

160. IPTG-inducible complementing strains were grown overnight in the absence of IPTG and

diluted to an OD_{600} of 0.05 and serially diluted and spotted onto LB plates containing 1µM

IPTG, Kan 150 µg/ml and AMP 25 or AMP 100.

Phase-contrast microscopy. For phase-contrast microscopy, 0.8 µl of exponentially-grown

cultures were spotted onto a 1.25% agarose pad as previously described (90) using a Nikon

Eclipse Ti inverted microscope and imaged using a Nikon Plan 60X oil Ph3 objective. Cell

length analysis was performed using the MicrobeJ plug-in for Fiji (91). A One-Way analysis of

variance (ANOVA) Kruskal-Wallis test with Dunn's post-test was used to compare the

indicated strains. Images were prepared using Adobe Photoshop, Adobe Illustrator, and

Prism.

Growth curves. For growth curves, exponentially growing cultures were diluted to an $OD_{600} =$

0.2 and 100 µl of dilute culture was added to wells of a 96-well plate. OD_{600} reading were

recording using a plate reader at 28°C with shaking every 5-10 min. When indicated,

ampicillin was added to a final concentration of AMP 25 or AMP 100. Plots of OD_{600} data

represent two technical replicates for each culture measured every 5 minutes for 24-48 h.

Disc diffusion assay. The disc-diffusion assay was used to determine the resistance of

Agrobacterium to various antibiotics. Cells were grown on LB agar plates. Sterile discs (6.5

mm in diameter) were placed on the surface of LB agar plates seeded with overnight culture

of indicated strains. We used three different discs: Blank (sterile disc containing no antibiotic),

AMP 10 (ampicillin 10μg/ml), and AMP 10/SUL 10 (ampicillin 10μg/ml sulbactam 10μg/ml).

The LB agar assay plates used for testing *A. tumefaciens* susceptibility were incubated at

28°C for 24-36 h. The assessment of antibacterial activity was based on the measurement of

diameter of the zone of inhibition formed around the disc minus the size of the disc. Three

independent trials were conducted for each concentration of each antibiotic. A two-way

analysis of variance (ANOVA) was used to compare the means.

Nitrocefin Assay. Nitrocefin is a chromogenic substrate for measuring β-lactamase activity.

Nitrocefin has an absorbance maximum of 390 nm. Upon hydrolysis of the β-lactam ring by a

β-lactamase, the absorbance shifts from 390 nm to 486 nm. By monitoring absorbance at 486

nm over time and using Beer's law ($A=\varepsilon lc$), we directly measured the rate of β-lactamase

hydrolytic activity. In two separate experiments, indicated strains were grown in LB until

desired optical densities (OD600) of 0.6 were reached. The cells were then pelleted by

centrifugation at 25,900 x g (rotor: Fiberlite F14-6x250y) for 5 minutes and the supernatant

collected and washed three times with PBS. Cell lysates were generated by adding the cell

lysis buffer BugBuster and sonicating. No lysozyme or protease inhibitors were added. Cell

lysates were normalized based on total protein content (7.5 µg/ml) and volume before

incubation in 100 µM nitrocefin solution in a 200 µl reaction volume at room temperature in 20

mM HEPES, 300 mM NaCl, pH 7.5. BugBuster lysis buffer was used as a blank. *ΔampD*

lysates were normalized based on total protein content (7.5 µg/ml) and subsequently diluted

to 1:5 as the rate of nitrocefin hydrolysis was significantly faster than the controls (WT, WT

AMP 25). Absorbance was immediately measured at 486 nm in 5 second intervals for 300

seconds (5 minutes). The change in absorbance (A) was converted to change in

concentration (c) of hydrolyzed product by using Beer's law (A=εlc) where the molar extinction

coefficient (ε) = 20500 M-1 cm-1 and path length (l) = 1 cm. Data plots of one experiment

appears in the main figures and the other in the supplementary figures respectively.

Construction of Agrobacterium strain for plant transformation experiments. *mltB3* was

deleted from the genome of WT *Agrobacterium tumefaciens* GV3101 using the same plasmid

and technique used to delete the gene in the strain C58. The resulting strain,

GV3101 *ΔmltB3*, was next transformed by introduction of the empty binary vector, pUBQ10-

GW (92) via electroporation. This plasmid confers KAN resistance to

the *Agrobacterium* strain, and also contains T-DNA repeat blocks which allow the transfer of

BASTA resistance to transformed into *Arabidopsis thaliana*. pUBQ10-GW was also

electroporated into wildtype GV3101 as a control.

Arabidopsis thaliana transformation efficiency. Plants with a bolt height between 2 and 7

centimeters were transformed via the floral dip method (85). 5 plants were transformed on

separate days from independent colonies for each C58 and GV3101. Plants were grown in

Promix BX (Premier Tech Horticulture) at 23°C, 16 h light/8 h dark, 100–150 µE m^{-2} s^{-1}, and

50–70% humidity until seeds were fully developed. Seeds were collected from fully matured

plants and store at 4°C at low humidity for one week. Seeds were then surface sterilized in

15% bleach containing 0.1% Triton X-100 with gentle rocking for 5 minutes. Seeds were then

washed 3 times using sterile water. Seeds were then imbided in sterile water at 4°C for 3

days. Seeds were then sown into soil using conditions listed above. Once seeds developed

true leaves BASTA (glufosinate ammonium) was applied via spray at a concentration of

10mg/L. After 4 days BASTA was re-applied ensuring only transformed plants survived.

Determining bacterial load of transformed *Arabidopsis thaliana* seeds. Seeds were

collected from fully matured transformed plants and stored at 4°C for 1 week with low

humidity. Seeds were then surface sterilized in 15% bleach containing 0.1% Triton X-100 with

gentle rocking for 5 minutes. Seeds were then washed 3 times using sterile water. 10mg of

seeds for each experimental condition were ground with a sterile mortar and pestle. Ground

seeds were then suspended in 1mL of sterile water, and serially diluted. 200µl of 10^{-1} dilution

was plated on minimal medium ATGN No AMP and AMP 25. Plates were then incubated at

28°C before colonies were counted.

ACKNOWLEDGEMENTS

We thank the Brown lab for helpful discussions. Research in the Brown lab on *A. tumefaciens*

cell growth is supported by the National Science Foundation (IOS1557806). WFC was

supported by a Gus T. Ridgel Fellowship.

AUTHOR CONTRIBUTIONS

WFC and PJBB conceptualized this work; WFC and CR developed methods; WFC, MH,

CR, AMR and AKY performed the experiments and completed data analysis; WFC, AMR, and

PJB wrote the manuscript; WFC, AMR, AKY, FC, and PJBB edited the manuscript; PJBB

acquired funding for this work.

REFERENCES

1.   Valentine L, Crops GM. 2003. *Agrobacterium tumefaciens* and the plant: The David and

Goliath of modern genetics. Plant Physiol 133:948–955.

2.   Escobar MA, Dandekar AM. 2003. *Agrobacterium tumefaciens* as an agent of disease.

Trends Plant Sci 8:380–386.

3.   Pulawska J. 2010. Crown gall of stone fruits and nuts, economic significance and

diversity of its causal agents: Tumorigenic *Agrobacterium* spp. J Plant Pathol 92:S87–

S98.

4.   Otten L, Burr T, Szegedi E. 2008. *Agrobacterium*: A disease-causing bacterium. Tzfira

T, Citovsky V (eds) *Agrobacterium* from Biol to Biotechnol Springer 1–46.

5.   Gelvin SB. 2003. *Agrobacterium*-mediated plant transformation: the biology behind the

"gene-jockeying" tool. Microbiol Mol Biol Rev 67:16–37.

6.   Nester EW. 2015. *Agrobacterium*: Nature's genetic engineer. Front Plant Sci 5:1–16.

7.   Dessaux Y, Faure D. 2018. Niche construction and exploitation by *Agrobacterium*: How

to survive and face competition in soil and plant habitats. Curr Top Microbiol Immunol

Springer 418.

8.   Singer K. 2018. The mechanism of T-DNA integration: Some major unresolved

questions. Curr Top Microbiol Immunol 418:287–317.

9.   Tzfira T, Citovsky V. 2000. From host recognition to T-DNA integration: the function of

bacterial and plant genes in the *Agrobacterium*-plant cell interaction. Mol Plant Pathol

1:201–212.

10. Gelvin S. 2000. *Agrobacterium* and plant genes involved in T-DNA transfer and

integration. Annu Rev Plant Biol 51:223–256.

11. Tzfira T, Citovsky V. 2006. *Agrobacterium*-mediated genetic transformation of plants:

biology and biotechnology. Curr Opin Biotechnol 17:147–154.

12. Gelvin SB. 2003. *Agrobacterium*-mediated plant transformation: the biology behind the

"gene-jockeying" tool. Society 67:16–37.

13. De La Riva GA, González-Cabrera J, Vázquez-Padrón R, Ayra-Pardo C. 1998.

Agrobacterium tumefaciens: A natural tool for plant transformation. Electron J

Biotechnol 1:25–48.

14. Leifert C, Cassells AC. 2000. Microbial hazards in plant tissue and cell cultures. Vitr Cell

Dev Biol 37:133–138.

15. Mogilner N, Zutra D, Gafny R, Bar-Joseph M. 1993. The persistence of engineered

Agrobacterium tumefaciens in agroinfected plants. Mol plant-microbe Interact MPMI.

16. Matzk A, Mantell S, Schiemann J. 1996. Localization of persisting agrobacteria in

transgenic tobacco plants. Mol Plant-Microbe Interact.

17. Hammerschlag FA, Zimmerman RH, Yadava UL, Hunsucker S, Gercheva P. 1997.

Effect of antibiotics and exposure to an acidified medium on the elimination of

Agrobacterium tumefaciens from apple leaf explants and on shoot regeneration. J Am

Soc Hortic Sci.

18. Mohana Priya A, Karutha Pandian S, Manikandan R. 2012. The effect of different

antibiotics on the elimination of *Agrobacterium* and high frequency *Agrobacterium*-

mediated transformation of indica rice (Oryza sativa L.). Czech J Genet Plant Breed

48:120–130.

19.   Demain AL, Blander RP. 1999. The β-lactam antibiotics: Past, present, and future.

Antonie van Leeuwenhoek, Int J Gen Mol Microbiol 75:5–19.

20.   Okkels FT, Pedersen MG. 1988. The toxicity to plant tissue and to *Agrobacterium*

tumefaciens of some antibiotics. Acta Hortic 225:199–208.

21.   Shackelford NJ, Chlan CA. 1996. Identification of antibiotics that are effective in

eliminating *Agrobacterium tumefaciens*. Plant Mol Biol Report 14:50–57.

22.   Estopà M, Marfà V, Melé E, Messeguer J. 2001. Study of different antibiotic

combinations for use in the elimination of *Agrobacterium* with kanamycin selection in

carnation. Plant Cell Tissue Organ Cult 65:211–220.

23.   Holford P, Newbury HJ. 1992. The effects of antibiotics and their breakdown products

on the *in vitro* growth of *Antirrhinum majus*. Plant Cell Rep 11:93–96.

24.   Minden V, Deloy A, Volkert AM, Leonhardt SD, Pufal G. 2017. Antibiotics impact plant

traits, even at small concentrations. AoB Plants 9.

25.   Jiang C, Brown PJB, Ducret A, Brun Y V. 2014. Sequential evolution of bacterial

morphology by co-option of a developmental regulator. Nature 506:489–493.

26.   Wiebke B, Ferreira F, Pasquali G, Bodanese-Zanettini MH, Droste A. 2006. Influence of

antibiotics on embryogenic tissue and *Agrobacterium tumefaciens* suppression in

soybean genetic transformation. Bragantia 65:543–551.

27.   Obembe O, Adesoye A, Machuka J. 2005. Evaluation of the effectiveness of selected

antibiotics in the suppression of *Agrobacterium* from Cowpea (*Vigna Unguiculata* L.

Walp.) embryo explants and as potential selective agents in *Agrobacterium*-mediated

transformation. J Trop Microbiol Biotechnol 1:27–33.

28.   Haddadi F, Aziz MA, Abdullah SNA, Tan SG, Kamaladini H. 2015. An efficient

Agrobacterium-mediated transformation of Strawberry cv. Camarosa by a dual plasmid

system. Molecules 20:3647–3666.

29.   Tang H, Ren Z, Krczal G, Glimelius K. 2000. An evaluation of antibiotics for the

elimination of *Agrobacterium tumefaciens* from walnut somatic embryos and for the

effects on the proliferation of somatic embryos and regeneration of transgenic plants.

Plant Cell Rep 19:881–887.

30.   Waxman DJ, Strominger JL. 1983. Penicillin-binding proteins and the mechanism of

action of β-lactam antibiotics. Annu Rev Biochem 52:825–869.

31.   Sauvage E, Kerff F, Terrak M, Ayala JA, Charlier P. 2008. The Penicillin-binding

proteins: Structure and role in peptidoglycan biosynthesis. FEMS Microbiol Rev 32:234–

258.

32.   Spratt BG. 2006. Distinct Penicillin-binding proteins involved in the division, elongation,

and shape of *Escherichia coli* K12. Proc Natl Acad Sci 72:2999–3003.

33.   Rolinson GN. 1980. Effect of β-lactam antibiotics on bacterial cell growth rate.

Microbiology 120:317–323.

34.   Jacoby GA. 2009. AmpC β-lactamases. Clin Microbiol Rev 22:161–182.

35.   Egan AJF, Cleverley RM, Peters K, Lewis RJ, Vollmer W. 2017. Regulation of bacterial

cell wall growth. FEBS J 284:851–867.

36.   Den Blaauwen T, Andreu JM, Monasterio O. 2014. Bacterial cell division proteins as

antibiotic targets. Bioorg Chem 55:27–38.

37.   Egan AJF, Biboy J, van't Veer I, Breukink E, Vollmer W. 2015. Activities and regulation

of peptidoglycan synthases. Philos Trans R Soc B Biol Sci 370.

38.   Typas A, Banzhaf M, Gross CA, Vollmer W. 2012. From the regulation of peptidoglycan

synthesis to bacterial growth and morphology. Nat Rev Microbiol 10:123–136.

39.   Vollmer W, Blanot D, De Pedro MA. 2008. Peptidoglycan structure and architecture.

FEMS Microbiol Rev 32:149–167.

40.   de Pedro MA, Cava F. 2015. Structural constraints and dynamics of bacterial cell wall

architecture. Front Microbiol.

41.   Margolin W. 2009. Sculpting the bacterial cell. Curr Biol 19:812–822.

42.   Park JT, Uehara T. 2008. How bacteria consume their own exoskeletons (turnover and

recycling of cell wall peptidoglycan). Microbiol Mol Biol Rev 72:211–227.

43.   Silhavy TJ, Kahne D, Walker S. 2010. The bacterial cell envelope. Cold Spring Harb

Perspect Biol 2:1–16.

44.   Scheffers D, Pinho MG. 2005. Bacterial cell wall synthesis: New insights from

localization studies. Microbiol Mol Bio Rev 69:585–607.

45.   Vermassen A, Leroy S, Talon R, Provot C, Popowska M, Desvaux M. 2019. Cell wall

hydrolases in bacteria: Insight on the diversity of cell wall amidases, glycosidases and

peptidases toward peptidoglycan. Front Microbiol 10:331.

46.   Frirdich E, Gaynor EC. 2013. Peptidoglycan hydrolases, bacterial shape, and

pathogenesis. Curr Opin Microbiol 16:767–778.

47.   Vollmer W, Joris B, Charlier P, Foster S. 2008. Bacterial peptidoglycan (murein)

hydrolases. FEMS Microbiol Rev 32:259–286.

48.   van Heijenoort J. 2011. Peptidoglycan hydrolases of *Escherichia coli*. Microbiol Mol Biol

Rev 75:636–663.

49.   Dhar S, Kumari H, Balasubramanian D, Mathee K. 2018. Cell-wall recycling and

synthesis in *Escherichia coli* and *Pseudomonas aeruginosa* – their role in the

development of resistance. J Med Microbiol 67:1–21.

50.   Dik DA, Fisher JF, Mobashery S. 2018. Cell-wall recycling of the Gram-negative

bacteria and the nexus to antibiotic resistance. Chem Rev. review-article, American

Chemical Society.

51.   Lee TK, Huang KC. 2013. The role of hydrolases in bacterial cell-wall growth. Curr Opin

Microbiol 16:760–766.

52.   Vollmer W. 2012. Bacterial growth does require peptidoglycan hydrolases. Mol

Microbiol 86:1031–1035.

53.   Fisher JF, Mobashery S. 2014. The sentinel role of peptidoglycan recycling in the β-

lactam resistance of the Gram-negative Enterobacteriaceae and *Pseudomonas*

aeruginosa. Bioorg Chem 56:41–48.

54.   Jacobs C, Joris B, Jamin M, Klarsov K, van Beeumen J, Mengin-Lecreulx D, van

Heijenoort J, Park JT, Normark S, Frère J-M. 1995. AmpD, essential for both β-

lactamase regulation and cell wall recycling, is a novel cytosolic N-acetylmuramyl-L-

alanine amidase. Mol Microbiol 15:553–559.

55.   Langaee TY, Gagnon L, Huletsky A. 2000. Inactivation of the *ampD* gene in

Pseudomonas aeruginosa leads to moderate-basal-level and hyperinducible AmpC β-

lactamase expression. Antimicrob Agents Chemother 44:583–589.

56.   Schmidtke AJ, Hanson ND. 2008. Role of *ampD* homologs in overproduction of AmpC

in clinical isolates of *Pseudomonas aeruginosa*. Antimicrob Agents Chemother

52:3922–3927.

57.   Jacobs C, Fre J. 1997. Cytosolic intermediates for cell wall biosynthesis and

degradation control inducible β-lactam resistance in Gram-negative bacteria 88:823–

832.

58.   Dik DA, Domínguez-Gil T, Lee M, Hesek D, Byun B, Fishovitz J, Boggess B, Hellman

LM, Fisher JF, Hermoso JA, Mobashery S. 2017. Muropeptide binding and the X-ray

structure of the effector domain of the transcriptional regulator AmpR of *Pseudomonas*

aeruginosa. J Am Chem Soc 139:1448–1451.

59.   Johnson JW, Fisher JF, Mobashery S. 2013. Bacterial cell-wall recycling. Ann N Y Acad

Sci 1277:54–75.

60.   Balasubramanian D, Kumari H, Mathee K. 2014. *Pseudomonas aeruginosa* AmpR: An

acute-chronic switch regulator. Pathog Dis 73:1–14.

61.   Vadlamani G, Thomas MD, Patel TR, Donald LJ, Reeve TM, Stetefeld J, Standing KG,

Vocadlo DJ, Mark BL. 2015. The β-lactamase gene regulator AmpR is a tetramer that

recognizes and binds the D-Ala-D-Ala motif of its repressor UDP-N-acetylmuramic acid

(MurNAc)-pentapeptide. J Biol Chem 290:2630–2643.

62.   Lindberg F, Normark S. 1987. Common mechanism of ampC β-lactamase induction in

enterobacteria: Regulation of the cloned *Enterobacter cloacae* P99 β-lactamase gene. J

Bacteriol 169:758–763.

63.   Torrens, G, Hernández S, Alfonso Ayala J, Moya B, Juan C, Cava F, Oliver A. 2019.

Regulation of AmpC-driven β-lactam resistance in *Pseudomonas aeruginosa*: Different

pathways, different signaling 4:1–14.

64.   Juan C, Torrens G, González-Nicolau M, Oliver A. 2017. Diversity and regulation of

intrinsic β-lactamases from non-fermenting and other Gram-negative opportunistic

pathogens. FEMS Microbiol Rev 41:781–815.

65.   Bajaj P, Singh NS, Virdi JS. 2016. *Escherichia coli* β-lactamases: What really matters.

Front Microbiol 7:1–14.

66.   Jaurin BA, Normark S. 1979. *In vivo* regulation of chromosomal β-lactamase in

Escherichia coli. J Bacteriol 138:896–902.

67.   Jaurin B, Grundström T, Edlund T, Normark S. 1981. The *E. coli* β-lactamase attenuator

mediates growth rate-dependent regulation. Nature 290:221–225.

68.   Honoré N, Nicolas MH, Cole ST. 1986. Inducible cephalosporinase production in clinical

isolates of *Enterobacter cloacae* is controlled by a regulatory gene that has been

deleted from *Escherichia coli*. EMBO J 5:3709–3714.

69.   Jacobs C, Huang LJ, Bartowsky E, Normark S, Park JT. 1994. Bacterial cell wall

recycling provides cytosolic muropeptides as effectors for β-lactamase induction. EMBO

J 13:4684–4694.

70.   Ogawa Y, Mii M. 2004. Screening for highly active β-lactam antibiotics against

Agrobacterium tumefaciens. Arch Microbiol 181:331–336.

71.   Goodner B, Hinkle G, Gattung S, Miller N, Blanchard M, Qurollo B, Goldman BS, Cao

Y, Askenazi M, Halling C, Mullin L, Houmiel K, Gordon J, Vaudin M, Iartchouk O, Epp A,

Liu F, Wollam C, Allinger M, Doughty D, Scott C, Lappas C, Markelz B, Flanagan C,

Crowell C, Gurson J, Lomo C, Sear C, Strub G, Cielo C, Slater S. 2001. Genome

Sequence of the Plant Pathogen and Biotechnology Agent *Agrobacterium tumefaciens*

C58. Science (80-) 294:2323 LP – 2328.

72.   CAO Y, Niimi Y, HU S lian. 2006. Meropenem as an alternative antibiotic agent for

suppression of *Agrobacterium* in genetic transformation of orchid. Agric Sci China

5:839–846.

73.   O'Callaghan CH, Morris A, Kirby SM, Shingler AH. 1972. Novel method for detection of

685 β-lactamases by using a chromogenic cephalosporin substrate. Antimicrob Agents

686 Chemother 1:283–288.

687 74. Morton ER, Fuqua C. 2012. Genetic manipulation of *Agrobacterium*. Curr Protoc

688 Microbiol Chapter 3:Unit-3D.2.

689 75. Uehara T, Park JT. 2007. An anhydro-N-acetylmuramyl-L-alanine amidase with broad

690 specificity tethered to the outer membrane of *Escherichia coli*. J Bacteriol 189:5634–

691 5641.

692 76. Pennartz A, Généreux C, Parquet C, Mengin-Lecreulx D, Joris B. 2009. Substrate-

693 induced inactivation of the *Escherichia coli* AmiD N-acetylmuramoyl-L-alanine amidase

694 highlights a new strategy to inhibit this class of enzyme. Antimicrob Agents Chemother

695 53:2991–2997.

696 77. Langaee TY, Gagnon L, Huletsky A. 2000. Inactivation of the *ampD* gene in

697 P*seudomonas aeruginosa* leads to moderate-basal-level and hyperinducible AmpC β-

698 lactamase expression. Antimicrob Agents Chemother 44:583–589.

699 78. Mercier F, Zervosen A, Teller N, Frre JM, Herman R, Pennartz A, Joris B, Luxen A.

700 2010. 1,6-AnhMurNAc derivatives for assay development of amidase AmiD. Bioorganic

701 Med Chem 18:7422–7431.

702 79. Juan C, Moyá B, Pérez JL, Oliver A. 2006. Stepwise upregulation of the *Pseudomonas*

703 *aeruginosa* chromosomal cephalosporinase conferring high-level β-lactam resistance

704 involves three AmpD homologues. Antimicrob Agents Chemother 50:1780–1787.

705 80. Wu C, Huang Y, Lin Y, Yang T. 2016. Inactivation of lytic transglycosylases increases

706 susceptibility to aminoglycosides and macrolides by altering the outer membrane

707 permeability of *Stenotrophomonas maltophilia*. Antimicrob Agents Chemother 60:3236–

3239.

81.   Yin J, Sun Y, Sun Y, Yu Z, Qiu J, Gao H, Oliver A. 2018. Deletion of lytic

transglycosylases increases β-lactam resistance in *Shewanella oneidensis*. Front

Microbiol 9:1–10.

82.   Zielińska A, Billini M, Möll A, Kremer K, Briegel A, Izquierdo Martinez A, Jensen GJ,

Thanbichler M. 2017. LytM factors affect the recruitment of autolysins to the cell division

site in *Caulobacter crescentus*. Mol Microbiol 106:419–438.

83.   Lamers RP, Nguyen UT, Nguyen Y, Buensuceso R, Burrows LL. 2015. Loss of

membrane-bound lytic transglycosylases increases outer membrane permeability and

β-lactam sensitivity in *Pseudomonas aeruginosa*. Microbiology 6:879–895.

84.   Cho H, Uehara T, Bernhardt TG. 2014. Beta-lactam antibiotics induce a lethal

malfunctioning of the bacterial cell wall synthesis machinery. Cell 159:1300–1311.

85.   Clough SJ, Bent AF. 1998. Floral dip: A simplified method for *Agrobacterium*-mediated

transformation of *Arabidopsis thaliana*. Plant J 16:735–743.

86.   Cheng ZM, Schnurr JA, Kapaun JA. 1998. Timentin as an alternative antibiotic for

suppression of *Agrobacterium tumefaciens* in genetic transformation. Plant Cell Rep

17:646–649.

87.   Ieamkhang S, Chatchawankanphanich O. 2005. Augmentin® as an alternative antibiotic

for growth suppression of *Agrobacterium* for tomato (*Lycopersicon esculentum)*

transformation. Plant Cell Tissue Organ Cult 82:213–220.

88.   Morton ER, Fuqua C. 2012. Laboratory Maintenance of *Agrobacterium*. Curr Protoc

Microbiol 6–11.

89.   Howell M, Aliashkevich A, Sundararajan K, Daniel JJ, Lariviere PJ, Goley ED, Cava F,

Brown PJB. 2019. *Agrobacterium tumefaciens* divisome proteins regulate the transition

from polar growth to cell division. Mol Microbiol 111:1074–1092.

90.   Howell M, Daniel JJ, Brown PJB. 2017. Live cell fluorescence microscopy to observe

essential processes during microbial cell growth. JoVE e56497.

91.   Ducret A, Quardokus EM, Brun Y V. 2016. MicrobeJ, a tool for high throughput bacterial

cell detection and quantitative analysis. Nat Microbiol 1:16077.

92.   Michniewicz M, Frick EM, Strader LC. 2015. Gateway-compatible tissue-specific vectors

for plant transformation Plant Biology. BMC Res Notes 8:6–13.

Figure 1

FIG 1. The AmpC-AmpR operon is responsible for induced ampicillin resistance in *Agrobacterium tumefaciens* **C58. (A)** Operon organization and proposed ampicillin resistance mechanism (53). Briefly, in the absence of β-lactams such as ampicillin (AMP), *ampC* expression is repressed by AmpR. AmpR-mediated repression is maintained as long as the AmpR-inactivating ligand, UDP-GM-Pentapeptide, is bound to AmpR ("P*ampC* OFF"). In contrast, the presence of ampicillin (AMP) increases the pools of AmpR-activating ligands, or the cell wall degradation fragments (anhydro modification is depicted by ring), which are known to displace AmpR-inactivating ligands. As a result, the increase of AmpR-activating ligands activates AmpR and *ampC* is transcribed ("P*ampC* ON"). **(B)** Growth of WT *A. tumefaciens, ΔampC,* and *ΔampR* in the absence (No AMP) and presence of ampicillin 25 or 50 µg/ml (AMP 25, AMP 50) for 24 h (n = 1, 2 replicates). **(C)** Quantitative analysis of phase-contrast microscopy of exponentially-growing strains in absence (No AMP) or presence of ampicillin 25 ug/ml (AMP 25). % Phenotype was calculated by counting the number of cells displaying one of the phenotypes indicated (1 cell = 1 phenotype) and dividing by the total number of cells per strain. **(D)** Ampicillin susceptibility assay performed via spotting dilutions. Briefly, indicated strains were cultured overnight (ON) on LB medium, serially diluted, spotted on LB solid medium containing no ampicillin (No AMP) or ampicillin 25 µg/ml (AMP 25), and incubated at 28°C for 36 h before imaging. **(E)** Disc susceptibility was performed on lawns of indicated strains grown on LB plates for 24 h at 28°C (n = 2). AMP 10 = disc containing 10 µg/ml ampicillin, AMP 10/SUL 10 = disc containing 10 µg/ml ampicillin and 10 µg/ml sulbactam, a broad-spectrum β-lactamase inhibitor. Data represent the mean (± standard deviation, SD) of three independent experiments. **** = $P < 0.0001$; *** = $P < 0.001$; ** = $P < 0.01$; * = $P < 0.1$; ns = not significant. **(F)** Determination of β-lactamase production performed via a nitrocefin assay using cell lysates. "No AMP" or "AMP 25" indicates cells untreated or treated, respectively, with ampicillin 25 µg/ml for 2 h before the generation of cell lysates. Data shown represents one out of two biological replicates.

Figure 2

769
FIG 2. Loss of AmpD results in constitutive β-lactamase activity and elevated ampicillin
resistance. (A) Ampicillin susceptibility assay performed via spotting dilutions. Briefly,
indicated strains were cultured ON at 28°C, serially diluted, spotted on solid medium
containing no ampicillin (No AMP) or ampicillin 25, 100, or 160 µg/ml (AMP 25, 100, 160), and
incubated at 28°C for ~40 h before imaging. **(B)** Growth of WT *A. tumefaciens* and *ΔampD* in
the absence (No AMP) and presence (AMP) of various concentrations of ampicillin 25, 100
µg/ml (AMP 25, AMP 100) for 24 h (n=1, 2 replicates). **(C)** Quantitative analysis of phase-
contrast microscopy of exponentially-growing strains treated with ampicillin 25 µg/ml (AMP
25). % Phenotype was calculated by counting the number of cells displaying one of the
indicated phenotypes (1 cell = 1 phenotype) and dividing by the total number of cells per
strain. **(D)** Disc susceptibility performed on lawn of indicated strains grown on LB plates for 24
h at 28°C. Blank = no antibiotic disc, AMP 10 = disc containing 10 µg/ml ampicillin, AMP
10/SUL 10 = disc containing 10 µg/ml ampicillin and 10 µg/ml sulbactam, a broad-spectrum β-
lactamase inhibitor. Data represent the mean (± standard deviation, SD) of three independent

experiments. **** = P < 0.0001; *** = P < 0.001; ** = P < 0.01; * = P < 0.1; ns = not significant.
785

786
FIG 3. AmpC is the β-lactamase that confers *ΔampD* elevated ampicillin resistance. **(A)**
Ampicillin susceptibility assay was performed via spotting dilutions. Briefly, indicated strains
were cultured ON at 28°C, serially diluted, spotted on solid medium containing no ampicillin
(No AMP) or ampicillin 25 µg/ml (AMP 25) and incubated at 28°C for ~40 h before imaging.
(B) Quantitative analysis of phase-contrast microscopy of exponentially-growing strains
untreated (No AMP) or treated with ampicillin 25 µg/ml (AMP 25). % Phenotype was
calculated by counting the number of cells displaying one of the indicated phenotypes (1 cell =
1 phenotype). **(C)** Determination of β-lactamase production was performed via a nitrocefin
assay using cell lysates. "No AMP" or "AMP 25" indicates cells untreated or treated,
respectively, with ampicillin 25 µg/ml for 2 h before the generation of cell lysates. Data shown
represents one out of two biological replicates.
798

799
FIG 4. MtlB3 is required for ampicillin resistance in *A. tumefaciens*. (A) Phase-contrast
microscopy of exponentially-growing strains treated with ampicillin 25 μg/ml (AMP 25) for 2 h.
(B) Quantitative analysis of phase-contrast microscopy of exponentially-growing strains
treated with ampicillin 25 μg/ml (AMP 25) for 2 h. % Phenotype was calculated by counting
the number of cells displaying one of the phenotypes indicated (1 cell = 1 phenotype) and
diving it by the total number of cells per strain. **(C)** Growth of WT *A. tumefaciens* and *ΔmltB3*
in the absence (No AMP) and presence of ampicillin 25 μg/ml (AMP 25) for 24 h (n = 1, 2
replicates). **(D)** Determination of β-lactamase production was performed via a nitrocefin assay
using cell lysates. "No AMP" or "AMP 25" indicates cells untreated or treated, respectively,
with ampicillin 25 μg/ml for 2 h before the generation of cell lysates. Data shown represents
one out of two biological replicates.
811

Figure 5

812
813
FIG 5. *A. tumefaciens* GV3103 *ΔmltB3* can be used to transform plants efficiently and
bacteria can be removed using low concentrations of ampicillin. (A) Ampicillin
susceptibility assay was performed via spotting dilutions. Briefly, indicated strains were
cultured ON at 28°C, serially diluted, spotted on solid medium containing no ampicillin (No
AMP), ampicillin 25 μg/ml (AMP 25), or carbenicillin 15 μg/ml (CARB 15), and incubated at
28°C for ~40 h before imaging. **(B)** Determination of β-lactamase production was performed
via a nitrocefin assay using cell lysates. Data shown represents one out of two biological
replicates. "No AMP" or "AMP 25" indicates cells untreated or treated, respectively, with
ampicillin 25 μg/ml for 2 h before the generation of cell lysates. **(C)** Transformation efficiency
and **(D)** bacterial loads of seeds transformed with WT *A. tumefaciens* GV3101 and GV3101
ΔmltB3 using the floral dip assay technique (85).
825
826

827
FIG 6. Working model for *A. tumefaciens* ampicillin resistance. Bifunctional PBPs extend
the cell wall through the transglycosylation (linking of carbohydrates) and transpeptidation
reactions (linking of peptide stems). β-lactams such as ampicillin (purple stars) target the
transpeptidase domain of Penicillin-Binding Proteins (PBPs) (**Fig S6**, Step 1-2), leading to a
block in bacterial cell growth. In *A. tumefaciens*, treatment with ampicillin leads to induction of
β-lactamases. Inactivation of the lytic transglycosylase MltB3 results inhibition of β-lactamase
derepression and lysis suggesting that MltB3 is likely required for the generation of cell wall
degradation products (**Fig S6**, Step 3). Similar to treatment with β-lactams, where a block in
cell growth leads to an increase in cell wall degradation products (**Fig S6**, Step 5), inactivation
of anhydro amidases such as AmpD increases the poll of this molecules leading to β-lactam

resistance. In *A. tumefaciens* inactivation of AmpD leads to derepression of β-lactamases and
ampicillin resistance (**Fig S6**, Step 9). Both AmpC, an inducible β-lactamase that is under the
transcriptional control of AmpR, and AmpR seem to be responsible for the derepression
observed in *ΔampD.* Thus, our working model suggests that upon ampicillin exposure, a block
in growth leads to increased activity of MltB3. An increase in cell wall degradation products
leads to induction of AmpC expression by AmpR and the presumed translocation of AmpC to
the periplasm resulting in ampicillin resistance.
845