Title:

Single-cell RNA sequencing of Tocilizumab-treated peripheral blood mononuclear cells as an in

vitro model of inflammation

Authors:

Arya Zarinsefat[1], George Hartoularos[2], Chun J. Yee[2], Minnie M. Sarwal[1]

Affiliations:

1. Department of Surgery, University of California, San Francisco, CA

2. Department of Bioengineering and Therapeutic Sciences, University of California, San

Francisco, CA

22
23

Abstract:

COVID-19 has posed a significant threat to global health. Early data has revealed that IL-6, a key regulatory cytokine, plays an important role in the cytokine storm of COVID-19. Multiple trials are therefore looking at the effects of Tocilizumab, an IL-6 receptor antibody that inhibits IL-6 activity, on treatment of COVID-19, with promising findings. As part of a clinical trial looking at the effects of Tocilizumab treatment on kidney transplant recipients with subclinical rejection, we performed single-cell RNA sequencing of comparing stimulated PBMCs before and after Tocilizumab treatment. We leveraged this data to create an in vitro cytokine storm model, to better understand the effects of Tocilizumab in the presence of inflammation. Tocilizumab-treated cells had reduced expression of inflammatory-mediated genes and biologic pathways, particularly amongst monocytes. These results support the hypothesis that Tocilizumab may hinder the cytokine storm of COVID-19, through a demonstration of biologic impact at the single-cell level.

1. Introduction

Coronavirus disease 2019 (COVID-19), caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has posed a significant threat to global health since emerging at the end of 2019. Although the spectrum of symptomatic infection ranges significantly, and most infections are not severe[1–3], the overall global burden of the disease has been significant with up to nearly 20% mortality in certain geographic/demographic groups[4,5]. While notable progress has been made in the understanding the virology and disease process, the abrupt onset and lack of effective vaccination has made treatment of COVID-19 difficult[6,7].

Interleukin (IL)-6 is a key regulatory cytokine for the innate and adaptive immune response and is a growth factor for B cell proliferation and differentiation, an inducer of antibody production, and a regulator of CD4+ T cell differentiation[8,9]. Early data from the COVID-19 outbreak has shown that the complications from the disease are partly due to increases in various cytokines, including IL-6[10–13], and that elevated IL-6 levels may be associated with worse outcomes[13–15]. Tocilizumab is an IL-6 receptor antibody, which binds to both the membrane-bound and soluble forms of the IL-6 receptor (IL-6R), thereby inhibiting the action of the cytokine/receptor complex and interfering with the cytokine's effects[16]. It is a well-studied and accepted therapy for rheumatoid arthritis[17–19], and has also been studied in giant cell arteritis[20] and organ transplantation[9,21,22]. As such, multiple global investigators are currently undertaking clinical trials to further assess the efficacy of Tocilizumab in the treatment of COVID-19 and its complications (ClinicalTrials.gov). Thus far, it has been shown that COVID-19 patient plasma inhibits the expression of HLA-DR which may be partially restored by

Tocilizumab treatment, and that treatment with Tocilizumab may also improve lymphopenia

associated with COVID-19[23]. Preliminary data for Tocilizumab treatment on COVID-19

outcomes has shown improvement in clinical outcomes[24]. While the clinical effects of

Tocilizumab in inflammatory and autoimmune disease has been well-studied, there is a

paucity of data on the mechanistic/biologic impact of the drug on our immune system.

Given the current state of the COVID-19 epidemic and possible efficacy of IL-6/IL-6R inhibition

with the use of Tocilizumab, we believed a deeper analysis of the mechanistic/biologic effects

of Tocilizumab could further elucidate the effects of the drug on our immune system. Herein we

present an analysis of the impact of Tociluzimab on immune cells using single-cell RNA

sequencing (scRNA-seq). We map the response of peripheral blood mononuclear cell (PBMC)

subsets to cellular activation using CD3/CD28 stimulation[25–29]. Relevant to understanding the

impact of Tociluzimab in suppressing immune activation and inflammation, as seen in the

COVID-19 response, we additionally examine the effect of Tociluzimab on unstimulated and

stimulated cells, as part of an investigator-initiated clinical trial in kidney transplant (KT)

recipients with subclinical rejection (*NIAID U01 AI113362-01*). We provide a resource

characterizing the effect of Tociluzimab on immune cells at a single-cell level, and demonstrate

the unique and unexpected impact of Tociluzimab on monocytes, and how its effect on

suppressing inflammation may be further augmented based on the resting versus activated

state of PBMCs before exposing the cells to IL-6R inhibition.

2. Results/Discussion

In order to examine the impact of Tocilizumab on the composition and expression of circulating single cells, we compared scRNA-seq data from anti-CD3/CD28 stimulated cells from control (patients not treated with Tocilizumab) PBMCs, to unstimulated PBMCs after 3 to 6 months of Tociluzimab treatment. After filtering cells, a total of 57,737 cells remained for analysis. These cells were put through our analysis pipeline described (see Methods). After UMAP clustering, there were a total of 21 distinct clusters representing major PBMC groups, inclusive of naïve CD4+ T/CD8+ T, activated CD4+ T/CD8+ T, memory CD4+ T/CD8+ T, B, Natural Killer (NK, both CD56+ dim and bright cells[39]), dendritic (DC) cells, and monocytes. Clusters were annotated according to canonical cell type markers (*Figure 1A*). Clusters 2, 4, and 5 expressed markers of memory T cell expansion (*S100A4, IL7R*[40,41]) while clusters 0, 8, and 16 expressed markers of CD4+ T cell activation (*TNFRS4, CD69*[42]). One cluster of doublets (cluster 20) was removed to give the final annotated clusters (*Figure 1B*).

Feature plots showing the expression of "cytokine storm"[43] related pro-inflammatory genes are cell-type specific, with predominance for expression in T cell and monocyte clusters (*Figure 1C*). Although many genes are known to be involved in the cytokine storm of COVID-19[37,38], we demonstrate that some of the key pro-inflammatory genes (cytokines, interferons, and tumor necrosis factor) are also noted as part of the inflammatory profile in control (no Tocilizumab) patients (*Figure 1C*, control cells). Overall, stimulated PBMCs not exposed to Tociluzimab show a dominant signal for T cell activation. After 6 months of treatment with Tociluzimab there is a shift in peripheral blood subset frequencies observed across no treatment (control) vs.

treatment (Tocilizumab) groups. In comparison to changes in overall cell types, there was little

observed effect on frequencies of naïve CD4+/CD8+ T cells, DC, or NK cells, but with a marked

reduction of activated CD4+ T cells (approximately 12.5% of control PBMCs were activated

CD4+ T cells, while there were essentially no activated CD4+ T cells in the Tocilizumab group,

Figure 1D). Within these different cell subsets, Tociluzimab therapy results in significant

polarization of gene expression based on UMAP presentation (*Figure 1E*), with notable

polarization by treatment status observed in monocytes.

Given Tocilizumab's function as an IL-6R blocker, we looked at the expression of *IL6*, *IL6R*, as

well as *SOCS1* (feedback inhibitor of IL-6 signaling, expressed upon IL-6 pathway activation[44]),

and *PRDM1 (*activated by the *JAK/STAT3* pathway via activation of the IL-6 pathway [45,46]) in

Tocilizumab-treated cells (*Figure 1F*). Tocilizumab-treatment resulted in the expected reduction

of *IL6R*, *SOCS1* and *PRDM1* expression, in CD4+ and CD8+ T cells, and unexpectedly also in

monocytes. *IL6* expression did not appear to be affected by Tocilizumab treatment.

Figure 1: UMAP clustering, cell subset annotation, and expression of inflammatory markers and
IL6R pathway genes in control vs. Tocilizumab-treated PBMCs
a, Dot plot of canonical markers used for annotation of the 20 cell clusters. Average feature
expression represented by color gradient with lower expression represented by light grey, and
higher expression represented by blue. Size of dots represent the percent of cells within that
specific cluster that express the feature of interest **b,** UMAP with final cell type annotations **c,**
Feature plots showing expression of select cytokines involved in SARS-CoV-2 cytokine storm
(*IFNG*, *CCL3*, and *TNF*) based on control vs. Tocilizumab treatment status. Feature expression
represented by color gradient, with low expression represented by blue and high expression
represented by red **d,** Bar plot showing the percentage of each cell type in control vs.
Tocilizumab-treated groups **e,** UMAP with cell clusters identified based on control vs.
Tocilizumab treatment status **f,** Feature plots showing expression of *IL6*, *IL6R,* and downstream
IL6R pathway genes (*SOCS1, PRDM1*) based on control vs. Tocilizumab treatment status. Feature
expression represented by color gradient, with low expression represented by blue and high
expression represented by red
144
145
We then looked at the top 30 most differentially expressed genes (highest log$_2$-fold changes)

for control vs. Tocilizumab amongst all cells (*Figure 2A*), CD4+ T cells (*Figure 2B*), CD8+ T cells

(*Figure 2C*), monocytes (*Figure 2D*), and performed corresponding PA for these genes. PA

showed enrichment of inflammatory pathways such IL and TNF signaling amongst control cells.

We looked at the top 30 most differentially expressed genes (highest log$_2$-fold changes) for

control vs. Tocilizumab monocytes (*Figure 2D*), with some notable differences as would be

expected. Control monocytes were enriched in chemokines such as *CXCL9*, various HLA genes

involved in antigen processing[48] (*HLA-DQB1*, *HLA-DRB5*), *CD40* (member of the TNF-receptor

superfamily[49]), and *SOCS1* (downstream gene activated by IL-6R pathway, as previously

discussed[44]). PA revealed enrichment of many inflammation-related pathways, including

interferon, interleukin, T cell receptor (TCR), and PD-1 signaling in control PBMCs, suggesting

the relative suppression of these pathways in cells exposed to Tocilizumab.

159
160
Figure 2: Differential expression testing and pathway analysis of all cells, CD4+ T cells, CD8+ T
cells, and monocytes
a, Heatmap of top 30 genes with highest log-fold changes in all control and Tocilizumab-treated
cells **b,** Heatmap of top 30 genes with highest log-fold changes in all CD4+ T control and
Tocilizumab-treated cells, with corresponding PA of top 10% most highly differentially expressed
genes (based on $\log_2$-fold change) in control vs. Tocilizumab cells **c,** Heatmap of top 30 genes
with highest log-fold changes in all CD8+ T control and Tocilizumab-treated cells, with
corresponding PA of top 10% most highly differentially expressed genes (based on $\log_2$-fold
change) in control vs. Tocilizumab cells **d,** Heatmap of top 30 genes with highest log-fold
changes in all control and Tocilizumab-treated monocytes, with corresponding PA of top 10%
most highly differentially expressed genes (based on $\log_2$-fold change) in control vs. Tocilizumab
cells. Gene expression level represented by color gradient ranging from purple (low expression)
to yellow (high expression). PA figure x-axis represents the number of genes from each pathway
that was present in the gene list. Adjusted p-values for pathway enrichment are represented as
a color gradient with larger p-values colored blue and smaller p-values colored red
176
177

In addition to the effect of Tociluzimab on T cells, we also observed an unexpected polarization

of monocytes after Tocilizumab treatment (*Figure 1E*). Notably, the Tocilizumab monocyte

cluster was enriched for *CD14*, suggestive of an increased presence of classical monocytes[47],

while *CD16/FCGR3A* expression was more evenly expressed between the two clusters (*Figure*

3A). We then performed cell trajectory analysis of these monocytes for Tociluzimab treatment

effect, utilizing *Monocle*. This revealed six distinct cell trajectory branches, with two of the

branches containing nearly all control cells not exposed to Tocilizumab, and the other four

branches containing nearly all Tocilizumab-exposed PBMCs (*Figure 3B*), supporting the

presence of unique PBMC trajectories after patient exposure to IL6-R blockade. We utilized

Monocle's BEAM function to perform branched expression analysis modeling of the distinct cell

trajectory branches for Tociluzimab-exposed PBMCs (circled branch, *Figure 3B*), which showed

distinct clusters of cells based on treatment status (*Figure 3C*).

190
Figure 3: Differential expression testing and cell trajectory analysis of monocyte subsets
a, Feature plots showing expression of CD14 and CD16 based on control vs. Tocilizumab
treatment status. Feature expression represented by color gradient, with low expression
represented by blue and high expression represented by red, with higher CD14 expression
noted in Tocilizumab cells **b,** Cell trajectory analysis of monocyte clusters showing distinct
lineages of control vs. Tocilizumab cells; blue circle represents branch point used in subsequent
heatmap analysis **c,** Heatmap from branched expression analysis modeling for most
differentially expressed genes between branch points from **b** (analyzed branch point marked by
blue circle), showing clusters of differentially expressed genes between branches. Gene
expression represented as color gradient from blue (low expression) to red (high expression).
Cell type annotation represented by two separate cell fates as seen in **b**, with middle of
heatmap representing the start of pseudotime and clear separation of control vs. Tocilizumab
cell fates
204
205

206   The results of this study showed that in PBMCs undergoing a cytokine storm signal in

207   rejection[50], with overlapping signatures of *IFNG*, *CCL3*, and *TNF* expression, along with TCR

208   signaling also seen in the cytokine storm of COVID-19[37,38], there is suppression of these

209   inflammatory pathways after Tocilizumab treatment. This is inclusive of suppression of

210   downstream signaling of IL6-R pathway genes in both monocytes and T cells.

211
Monocytes have been shown to play a significant role in the pathophysiology of COVID-19[51].

A significant expansion of populations of monocytes producing IL-6 has been observed in the

peripheral blood of patients with COVID-19 in ICUs compared with those patients who did not

require ICU hospitalization[52], with similar findings of increased IL-6 production from monocytes

also seen by scRNA-seq analysis of PBMCs[53]. Our findings are from the first clinical trial utilizing

Tocilizumab for transplant rejection recipients and the first scRNA-seq analysis for such a study.

We show a separation of cell clustering based on treatment status, reduced enrichment of

inflammatory pathways in Tocilizumab patients, and relatively reduced expression of IL-6R

pathway genes in Tocilizumab-treated cells. As would be expected, we did not observe any

differences in IL-6 gene expression between control and Tocilizumab cells (as Tocilizumab is an

IL-6R blocker), but rather only effects on the subsequent function of that cytokine's pathways.

We also show an enrichment of *CD14* expression (associated with classical monocytes) in

Tocilizumab-treated monocytes, which are believed to be phagocytic, but with reduced

inflammatory attributes[47]. This is consistent with our PA described above that shows

enrichment of inflammatory pathways in control cells, but not Tocilizumab-treated cells

(possibly due to the increased presence of non-inflammatory classical monocytes in

Tocilizumab-treated cells).

Our findings, in conjunction with the available data on clinical outcomes of Tocilizumab

treatment[24] and ongoing trials, show promise for the use of Tocilizumab in the treatment of

patients with COVID-19. The results of our study support the belief that Tocilizumab may be

effective in reducing the inflammatory burden that results in the adverse outcomes of COVID-

19. Future studies will need to be undertaken to look at outcomes of Tocilizumab treatment for

COVID-19 in a clinical trial setting, ideally in conjunction with scRNA-seq analysis of these

patient's blood samples to achieve a greater understanding of the transcriptomic effects of

infection and treatment at a single-cell level.

237
238   **3. Materials and Methods**
239
Sample collection

This study was performed as part of an ancillary to a randomized controlled clinical trial of 15

KT recipients that were diagnosed with subclinical rejection on their 6-month post-transplant

protocol biopsy and randomized to either continue standard of care (Tacrolimus,

mycophenolate, and steroid) immunosuppression (control arm, 8 patients) or standard of care

plus Tocilizumab (Tocilizumab treatment arm, 7 patients). Patients in the treatment arm were

given Tocilizumab at a dose of 8 mg/kg IV every 4 weeks, for a total of 6 doses. Patients in both

arms of the study had blood collected at baseline prior to the initiation of Tocilizumab (in the

treatment arm patients), then at 3, 6, and 12 months after the start of the study, for a total of 4

blood samples per all 15 patients in the study. PBMCs were isolated from blood samples by

Ficoll-Paque™ PLUS density gradient centrifugation (GE Healthcare, Chicago, IL) and frozen in

fetal bovine serum (Gibco, Waltham, MA) containing 10% (vol/vol) dimethyl sulfoxide (Sigma-

Aldrich, St. Louis, MS). Cells were frozen and not thawed until the day of the experiment when

they were used directly for in vitro stimulation.

Stimulation with anti-CD3 and anti-CD28 antibodies

Frozen PBMCs were thawed, four vials at a time to ensure maximum cell recovery, in a water

bath at 37 Celsius. Cells were counted using a hematocytometer, split in half, and were then

adjusted to $2x10^5$ cells/well and triplicate plated in multiscreen 96-well plates (Falcon, Corning,

NY). Cells were stimulated with soluble anti-CD3 (5 µg/mL; MabTech, Cincinnati, OH) and anti-

CD28 antibodies (10ug/mL; MabTech, Cincinnati, OH) at 37 Celsius, 5% CO_2 for 24 hours.

Unstimulated PBMCs were incubated under identical conditions to reduce any confounding

from incubation conditions other than stimulation. Since all PBMCs were split in half prior to

any downstream processing, all samples from control and Tocilizumab-treated patients at all

study time points were both stimulated and not stimulated as part of the study design.

Sample processing

After overnight stimulation/incubation, the cells were harvested and counted using a

hematocytometer and orange acridine solution. Any cell suspension that was less than 25

cells/uL was disqualified from multiplexing due to low cell counts. Multiplexing cell pools were

designed such that no pair of stimulated and unstimulated samples from the same patient were

in the same pool and such that no samples from the same collection time point were in the

same pool. The same number of cells from each patient and experimental condition were

multiplexed into their respective pools to make a final total of 300,000 cells per pool. Any

remaining non-pooled cells were resuspended in RNAlater (Thermo-Fisher, West Sacramento,

CA) and saved for SNP array. Cell pools were then centrifuged at 400g for 5 minutes and media

was aspirated. Cell pellet was resuspended in a small volume of Wash Buffer (0.4% BSA in

1XPBS) and the suspension was filtered through a 40uM cell strainer (Falcon, Corning, NY).

Library construction and sequencing

scRNA-seq libraries were prepared using the 10X Chromium Single Cell 3' Reagent Kits

v3, according to the manufacturer's instructions. Briefly, the isolated cells were washed once

with PBS + 0.04% BSA and resuspended in PBS + 0.04% BSA to a final cell concentration

of 1000 cells/μL as determined by hematocytometer. Cells were captured in droplets at a

targeted cell recovery of 4000-8000 cells, resulting in estimated multiplet rates of 0.4-5.4%.

Following reverse transcription and cell barcoding in droplets, emulsions were broken and

cDNA purified using Dynabeads MyOne SILANE (Thermo-Fisher, West Sacramento, CA) followed

by PCR amplification (98°C for 3 min; 12-16 cycles of 98°C for 15 sec, 67°C for 20 sec, 72°C for 1

min; 72°C for 1 min). Amplified cDNA was then used for 3' gene expression library construction.

For gene expression library construction, 2.4-50 ng of amplified cDNA was fragmented and end-

repaired, double-sided size selected with SPRIselect beads (Beckman Coulter, West

Sacramento, CA), PCR amplified with sample indexing primers (98°C for 45 sec; 14-16 cycles of

98°C for 20 sec, 54°C for 30 sec, 72°C for 20 sec; 72°C for 1 min), and double-sided size selected

with SPRIselect beads. Pooled cells were loaded in a 10X chip in three replicate wells such that

each well contained 50,000 cells. Given the large number of cells and large number of patient

samples, the entire experiment and sequencing was performed in 2 separate batches to

prevent cell death during counting. Each day resulted in 4 unique pools with each pool run in

triplicate wells for sequencing. Sequencing single-cell RNA libraries were sequenced on an

Illumina NovaSeq S2 to a minimum sequencing depth of 50,000 reads/cell using the read

lengths 26bp Read1, 8bp i7 Index, 91bp Read2.

Demultiplexing

To assign cells to donors of origin in our multiplexed design, we leveraged the genetic

demultiplexing tools *demuxlet*[30] and *freemuxlet*, both a part of the *popscle* suite of population

genetics tools (https://github.com/statgen/popscle). These tools leverage the genetic

polymorphisms present in transcripts to assign the cells found in each droplet to their donor of

origin. *Demuxlet* uses the genotype calls from a genotyping SNP array to classify cells in

droplets according to their donor of origin, while freemuxlet "learns" the genotypes of a pre-

defined number of donors from the transcripts themselves, and assigns the droplets to a

respective anonymous donor according to those learned genotypes. Upon first receiving

sequencing data, *demuxlet* was run with input genotypes from all the patients in the cohort.

While *demuxlet* was able to assign most droplets to donors of origin, it revealed that two

patients in the genotyping SNP array appeared to have identical genotypes (likely due to human

error) and that cells from some patients seemed to drop out (likely due to low viability cells or

inaccurate cell counting or mixing). Therefore, to validate *demuxlet* results, *freemuxlet* was run

using an independent list of SNP sites: exonic SNPs with a minor allele frequency > 0.05 as

observed in the 1000 Genomes Project. In order to leverage the droplets across multiple

microfluidic reactions, which may enable higher confidence in the learned genotypes, we

merged the BAMs from multiple experiments containing the same patients into a single BAM

and input this merged BAM into *freemuxlet*. The droplet assignments from the anonymous

donors output by *freemuxlet* were then compared to those from *demuxlet*, showing very high

concordance. Moreover, comparing the VCF generated from *freemuxlet* (using the SNPs present

in the droplets) to the VCF generated from the SNP genotyping array yielded a 1:1

correspondence of anonymous individuals to patients, barring those few problematic patients.

Through comparing VCFs and the presence/absence of individuals in each multiplexed

experiment, we were able to definitively assign a detected genotype to all detected individuals.

Droplet barcodes were then filtered to remove heterotypic droplets containing cells from

multiple individuals, and the remaining homotypic droplets were analyzed downstream.

Data analysis

Raw FASTQ files were processed using *CellRanger* (v 3.0.1) to map reads against human genome

38 as a reference, filter out unexpressed genes, and count barcodes and unique molecular

identifiers (UMIs). Subsequent analyses were conducted with *Seurat* (v 3.1.2)[31] in *R* (v 3.6.2).

We compared PBMCs from all anti-CD3/CD28 stimulated cells from the study baseline, to

unstimulated Tocilizumab-treated cells from 3 to 6 months post-treatment with Tocilizumab.

Utilizing *Seurat*, we first filtered cells to only keep those that had less than 10% mitochondrial

genes and cells with numbers of features greater than 200 and less than 2,500. Cells were

assigned patient identification based on the *demuxlet/freemuxlet* output described above, and

once patients were identified, additional treatment/stimulation/time metadata could be

applied. Given that our experiment was divided over 2 days given the high number of

samples/cells, we applied *Seurat's* SCTransform function for data integration to account for any

possible batch effects from experiment days[32,33]. Once the data was integrated, we continued

downstream data processing. We first determined the principal components (PCA), then

constructed a shared nearest neighbor graph (SNN), identified clusters with a resolution of

0.75, and finally visualized the cells using uniform manifold approximate and projection

(UMAP), per the typical Seurat workflow[31]. Clustering was achieved by using 15 components

from the PCA dimensionality reduction.

To identify cluster-specific markers following the creation of UMAP plots, we utilized

normalized RNA counts of all clusters, scaled the data, and performed differential gene

expression (DE) testing by applying the Wilcoxon rank sum test using *Seurat's* FindMarkers

function[31]. We also plotted normalized and scaled gene expression of canonical markers in

conjunction with DE testing to determine identities of each cluster. To compare cell clusters of

stimulated vs. unstimulated cells, or control vs. Tocilizumab-treated cells, we once again utilized

normalized/scaled RNA counts and performed DE testing with FindMarkers.

To perform pathway analysis (PA) for any specific comparison we performed, we filtered for all

differentially expressed genes with an adjusted (based on the Bonferroni correction) p-value <

0.05, and then selected the top 10 percentile of genes with the highest log-fold changes. These

top genes were used to perform the PA utilizing the Reactome database[34] with the

clusterProfiler package[35]. To perform cell trajectory analysis, we first subset our clusters and cell

types of interest from our *Seurat* workflow, then performed dimensionality reduction and cell

ordering with *Monocle*[36] (v 2.14.0). We were then able to plot specific cells by their trajectory

branches based on their pseudotime values assigned by *Monocle*. DE of individual cell trajectory

branches was then performed with *Monocle's* BEAM (branched expression analysis modeling)

function, followed by visualization of these differentially expressed branches with *Monocle's*

heatmap visualization tool.

Acknowledgements:

The authors thank the many individuals without whose enthusiastic participation and help this

study would never have been accomplished. We would like to acknowledge the following: TA

Sigdel who contributed to the study design, JA Liberto who contributed to study design and cell

culture/isolation, P Rashmi who contributed to the study design, and AA Da Silva who

contributed to cell culture/isolation. A Zarinsefat is funded by the NIH: 5 T32 AI 125222.

Competing Interests:

The authors of this study have no financial disclosures or non-financial competing interests to

disclose as described by e*Life*.

References:

1.   Chan JF-W, Yuan S, Kok K-H, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet (London, England)*. 2020;395(10223):514-523

2.   Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet (London, England)*. 2020;395(10223):497-506

3.   Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*. February 2020

4.   Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients Dying in Relation to COVID-19 in Italy. *JAMA*. March 2020

5.   Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. February 2020

6.   Ahmed SF, Quadeer AA, McKay MR. Preliminary Identification of Potential Vaccine Targets for the COVID-19 Coronavirus (SARS-CoV-2) Based on SARS-CoV Immunological Studies. *Viruses*. 2020;12(3)

7.   Mitja O, Clotet B. Use of antiviral drugs to reduce COVID-19 transmission. *Lancet Glob Heal*. March 2020

8.   Hunter CA, Jones SA. IL-6 as a keystone cytokine in health and disease. *Nat Immunol*. 2015;16(5):448-457

9.   Jordan SC, Choi J, Kim I, et al. Interleukin-6, A Cytokine Critical to Mediation of Inflammation, Autoimmunity and Allograft Rejection: Therapeutic Implications of IL-6

Receptor Blockade. *Transplantation*. 2017;101(1):32-44

10.   Yang Y, Shen C, Li J, et al. Exuberant elevation of IP-10, MCP-3 and IL-1ra during SARS-

CoV-2 infection is associated with disease severity and fatal outcome. *medRxiv*. January

2020:2020.03.02.20029975

11.   Vaninov N. In the eye of the COVID-19 cytokine storm. *Nat Rev Immunol*. April 2020

12.   Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ. COVID-19: consider

cytokine storm syndromes and immunosuppression. *Lancet (London, England)*.

2020;395(10229):1033-1034

13.   Chen X, Zhao B, Qu Y, et al. Detectable serum SARS-CoV-2 viral load (RNAaemia) is closely

correlated with drastically elevated interleukin 6 (IL-6) level in critically ill COVID-19

patients. *Clin Infect Dis*. April 2020

14.   Li X, Xu S, Yu M, et al. Risk factors for severity and mortality in adult COVID-19 inpatients

in Wuhan. *J Allergy Clin Immunol*. April 2020

15.   Wan S, Yi Q, Fan S, et al. Relationships among lymphocyte subsets, cytokines, and the

pulmonary inflammation index in coronavirus (COVID-19) infected patients. *Br J*

Haematol. April 2020

16.   Lee DW, Gardner R, Porter DL, et al. Current concepts in the diagnosis and management

of cytokine release syndrome. *Blood*. 2014;124(2):188-195

17.   Smolen JS, Aletaha D. Interleukin-6 receptor inhibition with tocilizumab and attainment

of disease remission in rheumatoid arthritis: the role of acute-phase reactants. *Arthritis*

Rheum. 2011;63(1):43-52

18.   Singh JA, Beg S, Lopez-Olivo MA. Tocilizumab for rheumatoid arthritis: a Cochrane

systematic review. *J Rheumatol*. 2011;38(1):10-20

19.   Campbell L, Chen C, Bhagat SS, Parker RA, Ostor AJK. Risk of adverse events including

serious infections in rheumatoid arthritis patients treated with tocilizumab: a systematic

literature review and meta-analysis of randomized controlled trials. *Rheumatology*

(Oxford). 2011;50(3):552-562

20.   Kwan CC, Thyparampil PJ. Tocilizumab for Giant Cell Arteritis. *Int Ophthalmol Clin*.

2020;60(2):57-62

21.   Shin B-H, Everly MJ, Zhang H, et al. Impact of Tocilizumab (Anti-IL-6R) Treatment on

Immunoglobulins and Anti-HLA Antibodies in Kidney Transplant Patients With Chronic

Antibody-mediated Rejection. *Transplantation*. 2020;104(4):856-863

22.   Choi J, Aubert O, Vo A, et al. Assessment of Tocilizumab (Anti-Interleukin-6 Receptor

Monoclonal) as a Potential Treatment for Chronic Antibody-Mediated Rejection and

Transplant Glomerulopathy in HLA-Sensitized Renal Allograft Recipients. *Am J Transplant*.

2017;17(9):2381-2389

23.   Giamarellos-Bourboulis EJ, Netea MG, Rovina N, et al. Complex Immune Dysregulation in

COVID-19 Patients with Severe Respiratory Failure. *Cell Host Microbe*. April 2020

24.   Xu X, Han M, Li T, et al. Effective treatment of severe COVID-19 patients with

tocilizumab. *Proc Natl Acad Sci U S A*. 2020;117(20):10970-10975

25.   Pizzolato G, Kaminski H, Tosolini M, et al. Single-cell RNA sequencing unveils the shared

and the distinct cytotoxic hallmarks of human TCRVdelta1 and TCRVdelta2 gammadelta T

lymphocytes. *Proc Natl Acad Sci U S A*. 2019;116(24):11906-11915

26.   Szabo PA, Levitin HM, Miron M, et al. Single-cell transcriptomics of human T cells reveals

tissue and activation signatures in health and disease. *Nat Commun*. 2019;10(1):4706

27.   Miragaia RJ, Gomes T, Chomka A, et al. Single-Cell Transcriptomics of Regulatory T Cells

Reveals Trajectories of Tissue Adaptation. *Immunity*. 2019;50(2):493-504.e7

28.   Luo T, Zheng F, Wang K, et al. A single-cell map for the transcriptomic signatures of

peripheral blood mononuclear cells in end-stage renal disease. *Nephrol Dial Transplant*.

December 2019

29.   Cai Y, Dai Y, Wang Y, et al. Single-cell transcriptomics of blood reveals a natural killer cell

subset depletion in tuberculosis. *EBioMedicine*. 2020;53:102686

30.   Kang HM, Subramaniam M, Targ S, et al. Multiplexed droplet single-cell RNA-sequencing

using natural genetic variation. *Nat Biotechnol*. 2018

31.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic

data across different conditions, technologies, and species. *Nat Biotechnol*.

2018;36(5):411-420

32.   Stuart T, Butler A, Hoffman P, et al. Comprehensive Integration of Single-Cell Data. *Cell*.

2019;177(7):1888-1902.e21

33.   Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq

data using regularized negative binomial regression. *bioRxiv*. January 2019:576827

34.   Fabregat A, Sidiropoulos K, Viteri G, et al. Reactome pathway analysis: a high-

performance in-memory approach. *BMC Bioinformatics*. 2017;18(1):142

35.   Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing biological

themes among gene clusters. *OMICS*. 2012;16(5):284-287

36.   Qiu X, Hill A, Packer J, Lin D, Ma Y-A, Trapnell C. Single-cell mRNA quantification and

differential analysis with Census. *Nat Methods*. 2017;14(3):309-315

37.   Ye Q, Wang B, Mao J. The pathogenesis and treatment of the `Cytokine Storm' in COVID-

19. *J Infect*. 2020;80(6):607-613

38.   Chua RL, Lukassen S, Trump S, et al. COVID-19 severity correlates with airway epithelium-

immune cell interactions identified by single-cell analysis. *Nat Biotechnol*. June 2020

39.   Michel T, Poli A, Cuapio A, et al. Human CD56bright NK Cells: An Update. *J Immunol*.

2016;196(7):2923-2931

40.   Salek-Ardakani S, Croft M. Regulation of CD4 T cell memory by OX40 (CD134). *Vaccine*.

2006;24(7):872-883

41.   Martin MD, Badovinac VP. Defining Memory CD8 T Cell . *Front Immunol* . 2018;9:2692

42.   Simms PE, Ellis TM. Utility of flow cytometric detection of CD69 expression as a rapid

method for determining poly- and oligoclonal lymphocyte activation. *Clin Diagn Lab*

Immunol. 1996;3(3):301-304

43.   Tisoncik JR, Korth MJ, Simmons CP, Farrar J, Martin TR, Katze MG. Into the eye of the

cytokine storm. *Microbiol Mol Biol Rev*. 2012;76(1):16-32

44.   Prêle CM, Woodward EA, Bisley J, Keith-Magee A, Nicholson SE, Hart PH. SOCS1 regulates

the IFN but not NFkappaB pathway in TLR-stimulated human monocytes and

macrophages. *J Immunol*. 2008;181(11):8018-8026

45.   Garbers C, Aparicio-Siegmund S, Rose-John S. The IL-6/gp130/STAT3 signaling axis: recent

advances towards specific inhibition. *Curr Opin Immunol*. 2015;34:75-82

46.   Liu J, Liang L, Li D, et al. JAK3/STAT3 oncogenic pathway and PRDM1 expression stratify

clinicopathologic features of extranodal NK/T-cell lymphoma, nasal type. *Oncol Rep*.

2019;41(6):3219-3232

47.   Mukherjee R, Kanti Barman P, Kumar Thatoi P, Tripathy R, Kumar Das B, Ravindran B.

Non-Classical monocytes display inflammatory features: Validation in Sepsis and

Systemic Lupus Erythematous. *Sci Rep*. 2015;5:13886

48.   Yamamoto F, Suzuki S, Mizutani A, et al. Capturing Differential Allele-Level Expression

and Genotypes of All Classical HLA Loci and Haplotypes by a New Capture RNA-Seq

Method. *Front Immunol*. 2020;11:941

49.   Martínez A, Bono C, Gozalbo D, Goodridge HS, Gil ML, Yáñez A. TLR2 and Dectin-1

Signaling in Mouse Hematopoietic Stem and Progenitor Cells Impacts the Ability of the

Antigen Presenting Cells They Produce to Activate CD4 T Cells. *Cells*. 2020;9(5)

50.   Sarwal M, Chua M-S, Kambham N, et al. Molecular heterogeneity in acute renal allograft

rejection identified by DNA microarray profiling. *N Engl J Med*. 2003;349(2):125-138

51.   Merad M, Martin JC. Pathological inflammation in patients with COVID-19: a key role for

monocytes and macrophages. *Nat Rev Immunol*. May 2020:1-8

52.   Zhou Y, Fu B, Zheng X, et al. Pathogenic T-cells and inflammatory monocytes incite

inflammatory storms in severe COVID-19 patients. *Natl Sci Rev*. March 2020

53.   Wen W, Su W, Tang H, et al. Immune cell profiling of COVID-19 patients in the recovery

stage by single-cell sequencing. *Cell Discov*. 2020;6:31