Cell population migration cycle and response to environmental pH in *Bacillus subtilis*

Madoka Nakayama[1], Izumi Takagi[2], Jun-ichi Wakita[3], Wataru Shoji[4], Sohei Tasaki[5*]

[1]Institute for Liberal Arts, Institute of Science Tokyo, Chiba 272-0827, Japan

[2]Mathematical Institute, Tohoku University, Miyagi 980-8579, Japan

[3]Department of Physics, Chuo University, Tokyo 112-8551, Japan

[4]Frontier Research Institute for Interdisciplinary Sciences (FRIS), Tohoku University, Miyagi 980-8579, Japan

[5]Department of Mathematics, Faculty of Science, Hokkaido University, Hokkaido 060-0810, Japan

[*]Corresponding Author: tasaki@math.sci.hokudai.ac.jp

Running title: Cell population migration cycle in *B. subtilis*

Highlights

- We study concentric colonies of *B. subtilis* that embody the migration cycle.
- External pH changes the proportion of migration and growth phases in the cycle.
- We propose a model of cell and cell population type regulation.
- Hysteresis in the model can explain the concentric pattern formation mechanism.
- The model predicts cell population regulation in response to the environment.

Abstract

Microbial populations are ubiquitous and some form a robust, multicellular structure called biofilm that protects the cells from environmental damage. The dispersion cycle is critical for controlling the population of important target microbes (both pathogenic and beneficial). Here, we show that the hysteresis of *Bacillus subtilis* cell-type regulation with respect to auto-inducing signal strength triggers the migration cycle of the cell population. We investigate migration cycle and its dependence on environmental conditions by quantitative analysis of cyclically expanding, concentric circular colonies. Next, we construct an input/output model that controls cell types in response to environmental conditions and signal density. On the basis of this model, we propose a migration cycle model for cell populations. The proposed model will widely predict biofilm-related phenomena and provide the basis for the description of highly self-regulating multicellular systems.

Keywords

Migration cycle; Biofilm life cycle; *Bacillus subtilis*; Concentric colony; pH response

Introduction

Microbial populations form colonies and they often exist stably in a robust system called biofilm [1-8]. The structural robustness of biofilms makes it difficult to kill disease-related microorganisms [9-

11]. In addition, myriads of useful microorganisms are also being discovered, so controlling the

population of target microorganisms is an urgent issue [12-14]. The process of biofilm formation is as

follows: (i) bacteria attach themselves to suitable surfaces available in the environment; (ii) they

proliferate while secreting extracellular polymeric substances (EPS) to form a biofilm; (iii) when the

colony becomes large and the cell density increases, the outer wall is destroyed and mobile cells are

released. These released cells wander around in search for a new environment suitable for growth.

When they identify an environment that satisfies their growth requirements, they become non-

migrating cells and colonize that environment. To summarize, these cells return to the first phase of

biofilm formation as described in (i) above. This cycle of repeating a series of phases is called the life

cycle of a bacterial biofilm. The concept of life cycle was originally used for individual organisms,

whereas in a biofilm, it is used for a group of cells. Therefore, it involves multilevel interactions, such

as information transfer within and between groups. However, the details of the biofilm life cycle

generation mechanism are not well understood.

A phenomenon similar to the life cycle of biofilms is the formation of concentric circular colonies

that embody the migration cycle of bacterial cell populations. (Fig. 1). *Proteus mirabilis* [15-18] and

Bacillus subtilis [18-23] are known to form concentric colonies. During their formation, the dominant

cell population for colony growth switches periodically between migrating and non-migrating states.

In *B. subtilis*, the migrating cell population consists of motile cells, which can swim by bundling and

rotating multiple flagella [24]. Particularly, two-dimensional collective movement, called swarming,

by motile cells plays a central role in rapid space expansion during the migration phase of the life cycle

[25-27]. There are three possible factors that cause the cell population to become non-migrating—

without swarming motility—: (i) switching to an immotile cell type; (ii) cessation of hyperflagellation;

and (iii) insufficient concentration of surface-wetting molecules, such as surfactin. In the case of this

B. subtilis concentric colony, the non-migrating cell population is in the form of matrix-producer-like

chains of proliferating cell populations, and the cells do not move at all [21]. Therefore, the major

mechanism underlying the stoppage of swarming involves the predominance of immotile cell types, most of which are matrix producers. Although the regulation of surface-wetting molecule secretion and hyperflagellation should also be involved in the stabilization of the cycle[21], we will consider this subject in a future research. Thus, here, we focus on cell type switching. A matrix producer does not move, and it proliferates while producing extracellular matrices [28]. The number of cells increases because of cell division, but in this phase, intercellular adhesion is strong and separation is rare [29-31]. Therefore, the cell group looks like a thread that continues to grow, and its growth mode is called chaining or bundling [32]. The colony not only expands two dimensionally but also increases in thickness (grows in the direction perpendicular to the medium plane). Cells called surfactin producers [33, 34] are known to coexist to assist the slow two-dimensional expansion in this case, improving sliding mobility on the medium surface [35], although it is far behind the expansion speed during the migration phase. In this way, the repetition of the two phases, the migration phase in which cells rapidly spread on the surface of the medium and the growth phase in which migration stops and the thickness increases, exactly represents the migration cycle of *B. subtilis* concentric colonies.

Although the process of concentric colony formation is known in this way, the detailed mechanism remains unclear. This is because of the lack of information on environment-dependent control patterns between different cell types, mobile and non-mobile. By focusing on *B. subtilis*, various mutually exclusive cell types have been discovered [24]. The balance between motile cells and matrix producers is particularly important for the migration cycle of the cell population. The genes related to each of these two types are mutually exclusive because of self-activation and mutual repression. When this mutual exclusivity becomes stronger, hysteresis occurs in cell-type regulation (i.e., cell-type selection depends on the history of past cell states) and a migration cycle is induced in the cell population, resulting in the formation of concentric circular colonies. Specifically, whether or not concentric colonies are formed depends on the presence or absence of hysteresis during cell-type control of the cell population and on environmental conditions. Investigating the conditions for forming concentric

colonies is essential for understanding the life cycle generation mechanism of cell populations.

However, the conditions for forming concentric colonies of *B. subtilis* have not yet been clearly

determined, although the need for agar concentrations of about 0.7% in solid medium and high nutrient

conditions is known [19, 21]. Particularly, the agar concentration is required to be in the range of 0.65–

0.75%. In addition, the incubation temperature is preferably 25–35°C, which is lower than the usual

35–37°C [19], and long-term pre-drying of the medium surface is desirable [21]. The situation seems

to be complicated by the effect of temperature and surface moisture other than agar and nutrient

concentrations. Therefore, we introduced environmental pH as a new parameter and attempted to

clarify the growth condition requirements of *B. subtilis*.

Except for concentric colonies, the dominant cell type is often defined as either a motile cell or matrix

producer. First, colonies formed by swarming motile cells are formed under an agar concentration of

0.4–0.7%. These colonies are classified into two types according to the nutrient concentration: disk-

like (high nutrition) [36, 37] and dense branching morphology (DBM)-like (low nutrition) [38-41]. On

the other hand, the slowly expanding colonies dominated by matrix producers are formed on a solid

medium with an agar concentration of 0.7% or more. These colonies are also classified into two types,

namely, Eden cluster-like (high nutrition) [42-44] and diffusion-limited-aggregation (DLA)-like (low

nutrition) [45-48], depending on the nutrient concentration [46, 47]. In general, the classification is

based on these five types. A detailed examination revealed that a small number of motile cells were

mixed even on the surface of a hard medium with an agar concentration of 1%, which affected

macroscopic colony morphology and expansion rate [49]. Particularly, a crater-like pattern in which

cells gather at the tip of the colony is formed by only slightly lowering the environmental pH from the

intracellular pH of 7.4. This is due to chemotaxis toward nutrients, which is a phenomenon that does

not occur during swarming [27]. On the other hand, when the agar concentration is low (0.5%), swift

colony expansion via swarming is observed because the energy required for migration is low due to

the availability of abundant water on the surface of the medium. However, under certain conditions,

i.e., optimum pH of 7.4, the dominant cell types become matrix producers rather than motile cells, and

lacework-patterned colonies may be formed [23]. Thus, other than concentric colonies, different stress

parameters, such as environmental pH, influence the choice between motile cells and matrix producers.

Here, we investigate the effect of environmental pH on cell-type selection and concentric *B. subtilis*

colony formation. By quantitative analysis of culture experiments, we clarify the range of

environmental pHs in which concentric colonies are formed, and the environmental pH dependence of

the temporal and spatial cycles within that range. Further, by combining the mathematical model of

cell-type regulation on the basis of the conventional gene regulatory network and analysis of its

mathematical properties, we propose a model for type regulation of cell population that changes

depending on the environment. Our proposed model provides a reasonable explanation for the

formation of concentric circular colonies and subsequent biofilm life cycle generation.

Results

Concentric colonies embody the biofilm life cycle

We prepared a medium with a high nutrient concentration and an intermediate agar concentration of

0.7% among the range of solid agar media (containing 0.2% or more agar), and inoculated *B. subtilis*

onto the center. When the plate was cultured at about 35°C, concentric colonies were formed on the

medium surface (Fig. 1). This is a simple model for biofilm life cycle generation, and its temporal and

spatial cycles embody that life cycle. We captured two-dimensional transmitted light images of this

colony using a scanner (Fig. 1(a)). The image data were converted to cell density distribution (Fig.

1(b)), using which a 3D structure was reconstructed (Fig. 1(c)), and the cross-section was analyzed

(Fig. 1(d)). By performing such quantitative analyses in a time series, not only the two-dimensional

cell density (colony thickness) but also the dynamics of cell proliferation and migration were analyzed

(Fig. 2). The boundaries of colonies were identified at each time and the depth of each terrace was

measured (Fig. 2(a), (b)). The concentric colonies of *B. subtilis* have distorted boundaries unlike those

Fig. 1. Concentric ring-like colony embodies the life cycle. (a) Raw image of transmitted light obtained by a flat scanner. The center is the inoculation point. The diameter of the dish is 88 mm. (b) Two-dimensional cell density. (c) Three-dimensional reconstruction of the colony. (d) Cross-sectional view.

of *Proteus mirabilis*. Therefore, the colony radius r was defined from area A of the inner area of the

colony: $r = \sqrt{A/\pi}$. Then, from the increase in the colony radius, the migration and growth phases of

B. subtilis biofilm, and their combined life cycle were observed (Fig. 2(c)). During one period, a set

of a migration phase and a growth phase, the rate of increase in total cell number is nearly constant,

and when the cells enter the next cycle and become motile cells, the cell proliferation rate decreases,

which may be due to the large amount of nutrients used in the transition to motile cells (Fig. 2(d)).

Microscopic observation of phase switching in concentric colonies

145   Using phase-contrast microscopy, we investigated how the concentric colony formation occurred, and

146   how the cell state and its spatial distribution changed. First, the termination process of the growth

Fig. 2. Growth dynamics of concentric ring-like colony. (a) Two-dimensional cell density. Color bar, cell density (10^8 cells mm^{-2}). Incubation time, 5, 10, 15, and 20 h after inoculation. (b) Cross-section cell density. The center is the inoculation point. (c) Increasing colony radius. (d) Increasing total number of cells.

147 phase, in which matrix producers are dominant, begins with the appearance of motile cells just before

148 the colony boundary (Fig. 3(a)). There, motile cells continue to grow and form ponds wherein cell

Fig. 3. Microscopic images of phase transition during colony expansion. (a)-(c) Transition from the growth phase to the migration phase. Motile cells appear slightly inside colony boundaries (large arrows in (a) and (b)). The number of motile cells increased to form bacterial ponds where cells organize to undergo swarming collective motion. The motile cells then destroy the boundary walls (small triangles in (a)). Cells flow out from some of the outermost destruction points on the colony boundaries to create rapidly expanding branches ((b) and (c)). (d)-(f) The migration phase. Numerous dense branches grow outward (d). Gradual collapse of the colony boundaries at the roots of branches, and the space around the roots is filled with the outflowing cells ((e) and (f)). (g)-(i) Transition from the migration phase to the growth phase. First, the branches stop growing (g). Then, the swarming regions composed of motile cells (small bubble-like regions in (g) and (h); whole branch region in (i) shrink from the outside to the inside of the colony ((g), (h) to (i); (g) is the outermost boundary, (i) is the innermost). Scale bars, 100 μm (except for (c)), 10 μm (c).

swarming occurs. The surrounding cells are partially separated from each other by autolysins [29, 30,

50, 51], and the pressure of proliferating motile cells is also applied, destroying many points in the

colony border area.

Among those destroyed points, the breakage starts at points on the outer side of the colony boundary,

and motile cells flow out from there to form branches (Fig. 3(b), (c), Supplementary Video S1). The

migration phase where motile cells are dominant is shown in Fig. 3(d)-(f). In this phase, the formed

branches quickly expand outward because of swarming (Fig. 3(d), Supplementary Video S2). Then,

the number of break points near the root increases, and the gap between branches is filled with outflow

cells from the inside (Fig. 3(e), (f), Supplementary Videos S3, S4). The end of the migration phase

begins when the branches stop growing (Fig. 3(g)). Immediately thereafter, the swarming area shrinks

from the tip of the branch toward the inner root (Fig. 3(g)-(i)). When there is no swarming region, the

growth phase begins. In the growth phase, the cells do not move and proliferate, and the cell density

increases. When the concentration exceeds a certain threshold, motile cells reappear, and transition to

the migration phase occurs (Fig. 3(a)). This is one round of the life cycle of *the B. subtilis* cell

population.

Effect of environmental pH on periodically expanding colonies

The conditions for forming concentric colonies of *B. subtilis* have been investigated. Particularly,

under optimal conditions (incubation temperature around 35–37°C), concentric circular colonies are

formed only in an extremely narrow range of agar concentration of 0.65–0.68%, even if nutritional

conditions are suitable [21]. It can be inferred that other stress parameters, such as low temperature

[19], are necessary for the formation of concentric colonies. We investigated the effect of

environmental pH as a stress parameter. Conditions other than environmental pH were fixed, and the

agar concentration was set at an intermediate level of 0.7% (Fig. 4). First, in the neutral pH region, no

periodically growing colonies appeared. Matrix producers were the predominant cell type, and

colonies grew by increasing their volume via proliferation (Fig. 4(a), (b)). As the environmental pH

decreased, many, very-short migration phases appeared around pH 6.8. Under the pH condition close

to this transition point, the variation in space was very strong, the periodicity was not clear, and the

Fig. 4. Effect of environmental pH on periodically expanding colonies. (a) Environmental pH conditions for forming periodically expanding colonies. (b) Colonies under different environmental pH conditions. (c) Circularity of colony ring (defined by circularity of second ring). (d) Spatial period (width of terrace). (e) Time period: growth phase, migration phase, and total period. Data represent mean ± s.d., $n = 6$ ((c) and (d)). All fitting curves are exponential plus constant ((d) and (e)).

pattern was far from that of concentric circles. When the pH decreased below 6.5, concentric circular

colonies that expand periodically were formed. When the pH was lower than 5.3, colony formation

itself became unstable, and the colony formation stopped halfway or no colonies were formed. Finally,

below pH 5.1, colonies were not always formed. In summary, the stable pH range for cyclic growth of

colonies was 5.3–6.5.

Next, we investigated the properties of concentric colonies in this pH range. First, the circularity (4pi

area divided by circumference squared) of the ring was constant, although there was some variation in

the pattern transition region near the neutral pH (Fig. 4(c)). The spatial period (width of the swarm

ring) seemed to increase monotonically as pH decreased (Fig. 4(d)). Specifically, in the pattern

transition region (pH 6.0–6.5), the spatial period sharply increased with decreasing pH, and then it had

a substantially constant period in the range of pH 5.3–6.0.

To clarify how these differences in colony morphology were produced, we examined the periods of

each growth phase and migration phase (Fig. 4(e)). We found that while the total time period was

almost constant, the growth phase duration decreased as the pH decreased; conversely, the duration of

the migration phase increased. Specifically, the finer spatial patterns reflected shorter migration

periods, while the total length of the life cycle of the cell population seemed homeostatic.

Environment-dependent regulation of cells and cell populations

We investigated the mechanism of control of the two different states of the cell population (migration

and growth phases). Cell population control should be understood on the basis of cell-level regulation

in response to environmental conditions The regulation between the two cell types, motile cells and

matrix producers, is well studied, particularly the switch between planktonic and biofilm states [24,

52-55]. The two cell types are designed to have mutually exclusive gene expression states. We created

a simple mathematical model to investigate cellular state dynamics (Fig. 5(a)). In this model, cell states

can be described according to four group variables by grouping coordinated genes and their products:

(1) S: A group represented by Spo0A~P, which also includes phosphorelay of Spo0F, Spo0B, and

Spo0A; (2) H: SigH; (3) A: AbrB; (4) C: A group represented by ComK, driven by the ComX-

ComP-ComA pathway.

The output of this control system is the cell type. In this case, it is a matrix producer when S is high

and a motile cell when S is low. Sporulation is initiated when S is continuously very high.

However, predominant sporulation occurs at a nutrient-depletion stage much later than colonization.

Fig. 5. A model for the response of cells and cell populations to environmental pH. (a) A model for cell-type selection for environmental pH and cell density. (b), (c) Two types of cell-type controls. Anhysteretic (b) and hysteretic (c). (d) A model for cell population-type selection. The horizontal axis (I_{AbrB}) is a parameter indicating the external activation rate of AbrB (see "Materials and Methods").

Therefore, it is not necessary to consider sporulation here. On the other hand, the inputs of the cell

state control system include external environmental conditions and auto-inducing signals that represent

cell density. There are various environmental conditions. For example, poor nutritional conditions may

be input to activate S, but we considered the formation of periodic growth colonies, and the initial

nutrient concentration was sufficiently high. Therefore, the nutritional conditions were constant and

sufficient at the time of colony formation. As an auto-inducing signal related to cell-type control, a

small peptide ComX is secreted by *B. subtilis* cells [56-58]. In the following, its concentration is

denoted as X, which is an input from C to the cell-type regulation system [56, 57, 59]. The input

mode of environmental pH is of significant interest at this stage. According to previously performed

studies, H or A is the candidate input point for external pH in the current model [60, 61].

Here, we considered the most minimal and smooth dynamic model of these four variables ("Materials

and Methods"). The curve of the set of equilibrium points can be divided into two types: with respect

to the increase and decrease in X, the variable that represents the cell state (for example, S) is

monotonic and non-monotonic (Fig. 5(b), (c)) [62]. In the latter case, the curve is an S-shaped curve

with two turning points. Specifically, there are two cases where cell-type control is either not hysteretic

(Fig. 5(b)) or hysteretic (Fig. 5(c)), considering the increase/decrease in the cell density signal X. The

latter hysteretic control is necessary for biofilm life cycle generation and concentric colony formation.

When there is hysteresis in the choice of cell state for increasing and decreasing X, a life cycle is

created for the cell population as follows. Migration phase: the cell density information X at the

growth front decreases as the cell population disperses in the motility state, and when it falls below a

certain threshold X_1, the cell population switches to the matrix-producing state. Growth phase: as the

cell population grows and matures in the matrix-producing state, X increases, and when it exceeds a

certain threshold $X_2 \left(> X_1 \right)$, the cell population switches to the motility state again.

AbrB is upregulated and SigH is suppressed as the environmental pH decreases in the weakly acidic

region of interest [60, 61]. These two are potential candidates for inputting environmental pH

information. It is not easy to determine which of these two elements, which suppress each other in the

regulatory network, is upstream for inputting external pH information. According to the detailed

mathematical analysis of the equilibrium state of this model, a situation in which hysteresis occurs

with a decrease in environmental pH and concentric colonies are formed can be realized only when

AbrB is the input point for pH (Fig. 5(a), (d); "Materials and Methods") [62] . Therefore, we proposed

a model of cell-type regulation using AbrB as the input point for environmental pH information (Fig.

5(a)). On the basis of this cell-type regulation model, the generation mechanism of the life cycle of the

cell population in the parameter range for hysteretic control could be explained (Fig. 5(c)), and the

dependence on environmental conditions could be understood.

Discussion

We have presented a possible mechanism of formation of *B. subtilis* concentric colonies that expand periodically and embody the life cycle of the cell population. By performing culturing experiments and their quantitative analysis, the dynamics of colony growth and its environmental pH dependence were clarified from a multilevel perspective—from the cell level to the macroscopic morphological level. We sought to understand patten formation under the assumption that the predominant cell type switches between motile cells and immotile cells, such as matrix producers. Based on the accumulation of molecular biology research to date, the outline of the gene regulatory network responsible for the regulation between these two cell types of *B. subtilis* has been established. We proposed a model of cell-type regulation by environmental conditions and cell density by combining concentric colony culture experiments under different environmental pH conditions. The cell density is considered to be measured by the cells themselves by the concentration of the secreted quorum-sensing factor ComX [56-58]. The mathematical properties of the minimal smooth model describing the dynamics also suggested the input location of environmental pH in the cell-type regulation system. The obtained model should be a general control mechanism that selects an appropriate cell population structure depending on the situation. Particularly, the S-shaped curve of the set of equilibrium points obtained by this mathematical model analysis makes it possible to understand the hysteretic control of the cell population that forms concentric colonies, and subsequently, the migration cycle generation mechanism.

In the cell population migration cycle, phase switches are dependent on cell density. We investigated the switching mechanism and found that the migration cycle was stably modulated via hysteretic control of cell type. Further, the effect of environmental conditions on colony patterns can be predicted. At present, however, environmental changes and cell type control were investigated via concentric circle colony formation experiments under environmental pH (and only neutral to acidic) conditions. In the future, we plan to elucidate how *B. subtilis* cells collectively cope with fluctuations in a large

range of pH, particularly via the study of the responses at high pH. Moreover, to predict the response

to changes in other environmental conditions, such as nutrient concentration, temperature, and oxygen

concentration, it is necessary to understand a wider gene regulatory network. Nevertheless, the

conditions for biofilm life cycle generation can be better organized as per the presence or absence of

hysteresis. Of note, this classification will probably be applicable to biofilm life cycle studies focusing

on other microbial species.

A morphological diagram was created for understanding the colony morphology of *B. subtilis* in the

1990s [36, 63], in which one of the five colonies was formed depending on the two parameters, nutrient

concentration and agar concentration, in the medium. These five colonies are called Eden cluster-like

[42-44], DLA-like [45-48, 64, 65], concentric ring-like, disk-like [36, 37], and DBM-like [38-41],

respectively. The concept of cell type was not established at the time, but it is now well-understood,

and the colony pattern can be explained as follows. The dominant cell type is immotile cells, such as

matrix producers, in Eden cluster-like and DLA-like colonies; disk-like and DBM-like colonies have

motile cells as the predominant cell type. Finally, in the concentric ring-like colony, the dominant cell

type switches between motile cells and immotile cells. Thus, among the five patterns, concentric

colonies had multiple dominant cell types, which created unsolved problems. The mechanism of

pattern formation was unknown, and it was not understood or reproduced by a mathematical model

[66]. Our proposed model for cell-type regulation has presented one possible mechanism for this

problem.

Furthermore, the concentric colonies presented uncertainties, even with respect to the formation

conditions, such as an extremely narrow agar concentration range [21] and the dependence on culture

temperature [19]. Apparently, parameters other than agar and nutrient concentrations were likely

involved in the formation of concentric colonies. In other words, introducing some kind of a stress

parameter is effective for concentric pattern formation. We obtained the conditions to stably form

concentric colonies by introducing environmental pH. By drawing the morphology diagram in a three-

parameter space, it is possible to understand the entire cell population morphology correctly and in detail. By incorporating our cell-type regulation model into a multilevel morphogenesis model, we expect to obtain a simulation model that accurately predicts the growth of multicellular systems with non-uniform cell state distribution.

Robust biofilms and other complex cell populations of microorganisms are social structures involved in the organization of diverse cell types. Therefore, understanding the mode of state regulation of each cell should be the first key to understanding and controlling the behavior of the cell population as a whole. Therefore, higher-order structures of cell populations are self-organized in a trans-hierarchical manner: from gene-level and cell-level control to microscopic structures, macroscopic morphology and function of cell populations, and interaction with the surrounding environment. Thus, the construction of multilevel models for controlling the state of cell populations will strongly advance the research on various multicellular societies.

Materials and Methods

Experimental methods

B. subtilis wild-type strain OG-01 (JCM32485) was used [46, 67]. A very small quantity of bacteria (endospores) was incubated overnight (10–16 h) in 5 mL of Luria–Bertani liquid growth medium (LB Broth Base (Lennox L Broth Base), Gibco BRL Life Technologies, cat. no. 12780-052) at 35°C with shaking at 140 rpm. For inoculation on solid agar media, this final pellet was diluted with the buffer to an OD_{600} of 0.5.

A solution of 86 mM NaCl and 29 mM K_2HPO_4 was prepared. As a nutrient, 20 g L^{-1} peptone (Bacto Peptone, BD Biosciences, cat. no. 211677) was added to the solution, and the pH was then adjusted to a designated value by adding 6 M HCl, and 0.7% agar (Bacto Agar, BD Biosciences, cat. no. 214010) was added to the above liquid nutrient medium. After autoclaving at 121°C for 15 min, 20 mL of the solution was poured into each plastic Petri dish. The dishes were maintained at room temperature

overnight and dried at 50°C for 90 min.

The pre-cultured bacterial suspension was inoculated using a sterilized needle at the center of the

prepared solid medium surface. The dishes were incubated in a humidified box at 35°C for the

designated time. Macroscopic colony-transmitted light images were obtained using a flat head scanner

(GT-X980, Epson). Cell or cell cluster level morphology was observed by phase-contrast microscopy

using an inverted microscope (IX83, Olympus) with a camera (DP80, Olympus).

Mathematical model of dynamic cell-type regulation.

To describe the dynamics of cell-type control (Fig. 5(a)), we consider the following minimal smooth

model:

$$\frac{dS}{dt} = \frac{c_S H}{a_S + b_S C} - d_S S$$

$$\frac{dH}{dt} = \frac{c_H}{a_A + b_A A} - d_H H$$

$$\frac{dA}{dt} = \frac{c_A}{a_A + b_A S} - d_A A \tag{1}$$

$$\frac{dC}{dt} = c_C X A - d_C C$$

324 where a_*, b_*, c_*, d_* $\left(* = S, H, A, C\right)$ are positive constants. There are two cases where cell-type

325 control is non-hysteretic (Fig. 5(b)) and hysteretic (Fig. 5(c)), regarding the increase/decrease in the

326 cell density signal X. We set a parameter indicating the rate at which AbrB is activated from outside

327 the model system of cell-type regulation (equation (1); Fig. 5(a)) as follows:

$$I_A = \frac{b_H}{a_H} \times \frac{c_A / a_A}{d_A} \tag{2}$$

329 Similarly, we define the rate at which Spo0A~P and SigH are activated from outside the model

330 system as

331
$$I_{SH} = \frac{b_A}{a_A} \times \frac{c_S / a_S}{d_S} \times \frac{c_H / a_H}{d_H}$$
(3)

Particularly, the external activation rate of AbrB $I_A = I_{AbrB}$ is the horizontal axis in Fig. 5(d). Thus,

the following can be proved [62]: There exists $I_{SH}^* > 0$ such that

1. If $I_{SH} \leq I_{SH}^*$, then the steady states are anhysteretic (Fig. 5(b)).

2. If $I_{SH} > I_{SH}^*$, then there exists $1 < I_{A,L} < I_{A,U}$ such that

(i) If $0 < I_A \leq I_{A,L}$, then the steady states are anhysteretic (Fig. 5(b)).

(ii) If $I_{A,L} < I_A < I_{A,U}$, then the steady states are hysteretic (Fig. 5(c)).

(iii) If $I_{A,U} \leq I_A$, then the steady states are anhysteretic (Fig. 5(b)).

Case 2 above (the external activation of Spo0A~P and SigH is larger than the threshold $I_{SH} > I_{SH}^*$)

shows the situation in Fig. 5(d), and it is also in agreement with the experimental results, suggesting

that $I_A = I_{AbrB}$ plays the role of a parameter corresponding to the environmental pH reduction. Note

the mathematical feature that at least AbrB must be actively involved to reproduce the pH dependence

of cell type control [62]. In the past literature, SigH or AbrB, especially SigH, was a candidate [60,

61], but at least SigH alone does not show the observed pH dependence of cell type control in the

mathematical model.

CRediT authorship contribution statement

Madoka Nakayama: Conceptualization, Methodology, Formal analysis, Investigation, Writing –

Review & Editing, Visualization, Funding acquisition. **Izumi Takagi:** Formal analysis, Writing –

Review & Editing, Funding acquisition. **Jun-ichi Wakita:** Formal analysis, Resources, Writing –

Review & Editing. **Wataru Shoji:** Formal analysis, Writing – Review & Editing, Supervision. **Sohei**

Tasaki: Conceptualization, Methodology, Software, Formal analysis, Investigation, Writing – Original

Draft, Writing – Review & Editing, Visualization, Supervision, Project administration, Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was supported by JSPS KAKENHI, Grant Number 23K03225 (M.N.), 23K03176 (I.T.), 19K03645 (S.T.), 23K03208 (S.T.), and MEXT KAKENHI, Grant Number 17H06327 (S.T.).

References

[1] P. Stoodley, K. Sauer, D.G. Davies, J.W. Costerton, Biofilms as Complex Differentiated Communities, Annu. Rev. Microbiol., 56 (2002) 187-209.

[2] R.M. Donlan, Biofilms: microbial life on surfaces, Emerg. Infect. Dis., 8 (2002) 881-890.

[3] L. Hall-Stoodley, J.W. Costerton, P. Stoodley, Bacterial biofilms: from the natural environment to infectious diseases, Nat. Rev. Microbiol., 2 (2004) 95-108.

[4] S.S. Branda, Å. Vik, L. Friedman, R. Kolter, Biofilms: the matrix revisited, Trends Microbiol., 13 (2005) 20-26.

[5] K. Kobayashi, M. Iwano, BslA(YuaB) forms a hydrophobic layer on the surface of *Bacillus subtilis* biofilms, Mol. Microbiol., 85 (2012) 51-66.

[6] H. Vlamakis, Y. Chai, P. Beauregard, R. Losick, R. Kolter, Sticking together: building a biofilm the *Bacillus subtilis* way, Nat. Rev. Microbiol., 11 (2013) 157-168.

[7] L. Hobley, C. Harkins, C.E. MacPhee, N.R. Stanley-Wall, Giving structure to the biofilm matrix: an overview of individual strategies and emerging common themes, FEMS Microbiol. Rev., 39 (2015) 649-669.

[8] H.-C. Flemming, J. Wingender, U. Szewzyk, P. Steinberg, S.A. Rice, S. Kjelleberg, Biofilms: an emergent form of bacterial life, Nature Reviews Microbiology, 14 (2016) 563-575.

[9] P. Gilbert, J. Das, I. Foley, Biofilm susceptibility to antimicrobials, Adv. Dent. Res., 11 (1997) 160-167.

[10] T.F. Mah, G.A. O'Toole, Mechanisms of biofilm resistance to antimicrobial agents, Trends
Microbiol., 9 (2001) 34-39.

[11] C.W. Hall, T.-F. Mah, Molecular mechanisms of biofilm-based antibiotic resistance and tolerance
in pathogenic bacteria, FEMS Microbiol. Rev., 41 (2017) 276-301.

[12] M. Kostakioti, M. Hadjifrangiskou, S.J. Hultgren, Bacterial biofilms: development, dispersal, and
therapeutic strategies in the dawn of the postantibiotic era, Cold Spring Harb. Perspect. Med., 3 (2013)
a010306-a010306.

[13] A. Algburi, N. Comito, D. Kashtanov, L.M.T. Dicks, M.L. Chikindas, Control of Biofilm
Formation: Antibiotics and Beyond, Appl. Environ. Microbiol., 83 (2017) e02508-02516.

[14] S. Galié, C. García-Gutiérrez, E.M. Miguélez, C.J. Villar, F. Lombó, Biofilms in the Food
Industry: Health Aspects and Control Methods, Front. Microbiol., 9 (2018).

[15] K.A. Bisset, The zonation phenomenon and structure of the swarm colony in *Proteus mirabilis*, J.
Med. Microbiol., 6 (1973) 429-433.

[16] O. Rauprich, M. Matsushita, C.J. Weijer, F. Siegert, S.E. Esipov, J.A. Shapiro, Periodic
phenomena in *Proteus mirabilis* swarm colony development, J. Bacteriol., 178 (1996) 6525-6538.

[17] T. Matsuyama, Y. Takagi, Y. Nakagawa, H. Itoh, J. Wakita, M. Matsushita, Dynamic Aspects of
the Structured Cell Population in a Swarming Colony of *Proteus mirabilis*, J. Bacteriol., 182 (2000)
385-393.

[18] Y. Yamazaki, T. Ikeda, H. Shimada, F. Hiramatsu, N. Kobayashi, J.-i. Wakita, H. Itoh, S. Kurosu,
M. Nakatsuchi, T. Matsuyama, M. Matsushita, Periodic growth of bacterial colonies, Physica D:
Nonlinear Phenomena, 205 (2005) 136-153.

[19] H. Fujikawa, Periodic growth of *Bacillus subtilis* colonies on agar plates, Physica A: Statistical
Mechanics and its Applications, 189 (1992) 15-21.

[20] H. Itoh, J.-i. Wakita, T. Matsuyama, M. Matsushita, Periodic pattern formation of bacterial
colonies, J. Phys. Soc. Japan, 68 (1999) 1436-1443.

[21] J. Wakita, H. Shimada, H. Itoh, T. Matsuyama, M. Matsushita, Periodic colony formation by
bacterial species *Bacillus subtilis*, J. Phys. Soc. Japan, 70 (2001) 911-919.

[22] H. Shimada, T. Ikeda, J.-i. Wakita, H. Itoh, S. Kurosu, F. Hiramatsu, M. Nakatsuchi, Y. Yamazaki,
T. Matsuyama, M. Matsushita, Dependence of local cell density on concentric ring colony formation
by bacterial species *Bacillus subtilis*, J. Phys. Soc. Japan, 73 (2004) 1082-1089.

[23] S. Tasaki, M. Nakayama, W. Shoji, Morphologies of *Bacillus subtilis* communities responding to
environmental variation, Dev. Growth Differ., 59 (2017) 369-378.

[24] D. López, H. Vlamakis, R. Kolter, Generation of multiple cell types in *Bacillus subtilis*, FEMS
Microbiol. Rev., 33 (2009) 152-163.

[25] D.B. Kearns, R. Losick, Swarming motility in undomesticated *Bacillus subtilis*, Mol. Microbiol.,

49 (2003) 581-590.

[26] D.B. Kearns, F. Chu, R. Rudner, R. Losick, Genes governing swarming in *Bacillus subtilis* and

evidence for a phase variation mechanism controlling surface motility, Mol. Microbiol., 52 (2004)

357-369.

[27] D.B. Kearns, A field guide to bacterial swarming motility, Nat. Rev. Microbiol., 8 (2010) 634-

644.

[28] M. Marvasi, P.T. Visscher, L. Casillas Martinez, Exopolymeric substances (EPS) from *Bacillus*

subtilis: polymers and genes encoding their synthesis, FEMS Microbiol. Lett., 313 (2010) 1-9.

[29] T.J. Smith, S.A. Blackman, S.J. Foster, Autolysins of *Bacillus subtilis*: multiple enzymes with

multiple functions, Microbiology, 146 (2000) 249-262.

[30] Y. Chai, T. Norman, R. Kolter, R. Losick, An epigenetic switch governing daughter cell separation

in *Bacillus subtilis*, Genes Dev., 24 (2010) 754-765.

[31] R. Chen, S.B. Guttenplan, K.M. Blair, D.B. Kearns, Role of the sigmaD-dependent autolysins in

Bacillus subtilis population heterogeneity, J. Bacteriol., 191 (2009) 5775-5784.

[32] R. Honda, J.-i. Wakita, M. Katori, Self-Elongation with Sequential Folding of a Filament of

Bacterial Cells, J. Phys. Soc. Japan, 84 (2015) 114002.

[33] M.M. Nakano, R. Magnuson, A. Myers, J. Curry, A.D. Grossman, P. Zuber, *srfA* is an operon

required for surfactin production, competence development, and efficient sporulation in *Bacillus*

subtilis, J. Bacteriol., 173 (1991) 1770-1778.

[34] M.M. Nakano, L.A. Xia, P. Zuber, Transcription initiation region of the *srfA* operon, which is

controlled by the *comP-comA* signal transduction system in *Bacillus subtilis*, J. Bacteriol., 173 (1991)

5487-5493.

[35] J. van Gestel, H. Vlamakis, R. Kolter, From cell differentiation to cell collectives: *Bacillus subtilis*

uses division of labor to migrate, PLoS Biol., 13 (2015) e1002141.

[36] J. Wakita, K. Komatsu, A. Nakahara, T. Matsuyama, M. Matsushita, Experimental investigation

on the validity of population dynamics approach to bacterial colony formation, J. Phys. Soc. Japan, 63

(1994) 1205-1211.

[37] J. Wakita, H. Kuninaka, T. Matsuyama, M. Matsushita, Size distribution of bacterial cells in

homogeneously spreading disk-like colonies by *Bacillus subtilis*, J. Phys. Soc. Japan, 79 (2010)

094002.

[38] D. Grier, E. Ben-Jacob, R. Clarke, L.M. Sander, Morphology and microstructure in

electrochemical deposition of zinc, Phys. Rev. Lett., 56 (1986) 1264-1267.

[39] Y. Sawada, A. Dougherty, J.P. Gollub, Dendritic and fractal patterns in electrolytic metal deposits,

Phys. Rev. Lett., 56 (1986) 1260-1263.

[40] E. Ben-Jacob, P. Garik, The formation of patterns in non-equilibrium growth, Nature, 343 (1990)

523-530.

[41] J. Wakita, I. Ràfols, H. Itoh, T. Matsuyama, M. Matsushita, Experimental investigation on the formation of dense-branching-morphology-like colonies in bacteria, J. Phys. Soc. Japan, 67 (1998) 3630-3636.

[42] M. Eden, A two-dimensional growth process, in: Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Volume 4: Contributions to Biology and Problems of Medicine, University of California Press, Berkeley, Calif., 1961, pp. 223-239.

[43] T. Vicsek, M. Cserző, V.K. Horváth, Self-affine growth of bacterial colonies, Physica A: Statistical Mechanics and its Applications, 167 (1990) 315-321.

[44] J. Wakita, H. Itoh, T. Matsuyama, M. Matsushita, Self-affinity for the growing interface of bacterial colonies, J. Phys. Soc. Japan, 66 (1997) 67-72.

[45] T.A. Witten, L.M. Sander, Diffusion-limited aggregation, a kinetic critical phenomenon, Phys. Rev. Lett., 47 (1981) 1400-1403.

[46] H. Fujikawa, M. Matsushita, Fractal growth of *Bacillus subtilis* on agar plates, J. Phys. Soc. Japan, 58 (1989) 3875-3878.

[47] M. Matsushita, H. Fujikawa, Diffusion-limited growth in bacterial colony formation, Physica A: Statistical Mechanics and its Applications, 168 (1990) 498-506.

[48] H. Fujikawa, M. Matushita, Bacterial fractal growth in the concentration field of nutrient, J. Phys. Soc. Japan, 60 (1991) 88-94.

[49] S. Tasaki, M. Nakayama, W. Shoji, Self-organization of bacterial communities against environmental pH variation: Controlled chemotactic motility arranges cell population structures in biofilms, PLoS One, 12 (2017) e0173195.

[50] S.A. Blackman, T.J. Smith, S.J. Foster, The role of autolysins during vegetative growth of *Bacillus subtilis* 168, Microbiology, 144 (1998) 73-82.

[51] W. Vollmer, B. Joris, P. Charlier, S. Foster, Bacterial peptidoglycan (murein) hydrolases, FEMS Microbiol. Rev., 32 (2008) 259-286.

[52] D.B. Kearns, R. Losick, Cell population heterogeneity during growth of *Bacillus subtilis*, Genes Dev., 19 (2005) 3083-3094.

[53] K. Kobayashi, Gradual activation of the response regulator DegU controls serial expression of genes for flagellum formation and biofilm formation in *Bacillus subtilis*, Mol. Microbiol., 66 (2007) 395-409.

[54] Y. Chai, F. Chu, R. Kolter, R. Losick, Bistability and biofilm formation in *Bacillus subtilis*, Mol. Microbiol., 67 (2008) 254-263.

[55] L.S. Cairns, L. Hobley, N.R. Stanley-Wall, Biofilm formation by *Bacillus subtilis*: new insights into regulatory strategies and assembly mechanisms, Mol. Microbiol., 93 (2014) 587-598.

[56] R. Magnuson, J. Solomon, A.D. Grossman, Biochemical and genetic characterization of a
competence pheromone from *B. subtilis*, Cell, 77 (1994) 207-216.

[57] M. Ansaldi, D. Marolt, T. Stebe, I. Mandic-Mulec, D. Dubnau, Specific activation of the *Bacillus*
quorum-sensing systems by isoprenylated pheromone variants, Mol. Microbiol., 44 (2002) 1561-1573.

[58] M. Okada, I. Sato, S.J. Cho, H. Iwata, T. Nishio, D. Dubnau, Y. Sakagami, Structure of the *Bacillus*
subtilis quorum-sensing peptide pheromone ComX, Nat. Chem. Biol., 1 (2005) 23-24.

[59] A.D. Grossman, R. Losick, Extracellular control of spore formation in *Bacillus subtilis*, Proc. Natl.
Acad. Sci. USA, 85 (1988) 4369-4373.

[60] W.M. Cosby, P. Zuber, Regulation of *Bacillus subtilis* σH (Spo0H) and AbrB in response to
changes in external pH, J. Bacteriol., 179 (1997) 6778-6787.

[61] J.C. Wilks, R.D. Kitko, S.H. Cleeton, G.E. Lee, C.S. Ugwu, B.D. Jones, S.S. BonDurant, J.L.
Slonczewski, Acid and base stress and transcriptomic responses in *Bacillus subtilis*, Appl. Environ.
Microbiol., 75 (2009) 981-990.

[62] S. Tasaki, M. Nakayama, I. Takagi, J. Wakita, W. Shoji, Necessary and sufficient condition for
hysteresis in the mathematical model of the cell type regulation of *Bacillus subtilis*, arXiv, (2020)
2009.06524.

[63] M. Ohgiwari, M. Matsushita, T. Matsuyama, Morphological changes in growth phenomena of
bacterial colony patterns, J. Phys. Soc. Japan, 61 (1992) 816-822.

[64] E. Ben-Jacob, H. Shmueli, O. Shochet, A. Tenenbaum, Adaptive self-organization during growth
of bacterial colonies, Physica A: Statistical Mechanics and its Applications, 187 (1992) 378-424.

[65] E. Ben-Jacob, O. Schochet, A. Tenenbaum, I. Cohen, A. Czirok, T. Vicsek, Generic modelling of
cooperative growth patterns in bacterial colonies, Nature, 368 (1994) 46-49.

[66] M. Mimura, H. Sakaguchi, M. Matsushita, Reaction–diffusion modelling of bacterial colony
patterns, Physica A: Statistical Mechanics and its Applications, 282 (2000) 283-303.

[67] T. Matsuyama, M. Matsushita, Comments on 'Classification and genetic characterization of
pattern-forming *Bacilli*' *Mol Microbiol* (1998) 27: 687–703, Mol. Microbiol., 31 (1999) 1278-1279.