Species- and strain-level assessment using *rrn* long-amplicons suggests donor's influence on gut microbial transference via fecal transplants in metabolic syndrome subjects

Alfonso Benítez-Páez[1,2*], Annick V. Hartstra[3], Max Nieuwdorp[3], Yolanda Sanz[1*]

1 Microbial Ecology, Nutrition & Health Research Unit. Institute of Agrochemistry and Food Technology, Spanish National Research Council (IATA-CSIC). 46980 Paterna-Valencia, Spain.

2 Host-Microbe Interactions in Metabolic Health Laboratory. Principe Felipe Research Center (CIPF). 46012 Valencia, Spain.

3 Department of Internal and Vascular Medicine, Amsterdam University Medical Centres. 1105 AZ Amsterdam, The Netherlands.

Running title: *Donors' influence on gut microbiota engraftment after FMT*

* To whom any correspondence should be addressed: ABP E-mail abenitez@cipf.es, YS E-mail yolsanz@iata.csic.es.

Abstract

Fecal microbiota transplantation (FMT) is currently used for treating *Clostridium difficile* infection and explored for other clinical applications in experimental trials. However, the effectiveness of this therapy could vary, and partly depend on the donor's bacterial species engraftment, whose evaluation is challenging because there are no cost-effective strategies for accurately tracking the microbe transference. In this regard, the precise identification of bacterial species inhabiting the human gut is essential to define their role in human health unambiguously. We used Nanopore-based device to sequence bacterial *rrn* operons (16S-ITS-23S) and to reveal species-level abundance changes in the human gut microbiota of a FMT trial. By assessing the donor and recipient microbiota before and after FMT, we further evaluated whether this molecular approach reveals strain-level genetic variation to demonstrate microbe transfer and engraftment. Strict control over sequencing data quality and major microbiota covariates was critical for accurately estimating the changes in gut microbial species abundance in the recipients after FMT. We detected strain-level variation via single-nucleotide variants (SNVs) at *rrn* regions in a species-specific manner. We showed that it was possible to explore successfully the donor-bacterial strain (e.g., *Parabacteroides merdae)* engraftment in recipients of the FMT by assessing the nucleotide frequencies at rrn-associated SNVs. Our findings indicate that the engraftment of donors' microbiota is to some extent correlated with the improvement of metabolic health in recipients and that parameters such as the baseline gut microbiota configuration, sex, and age of donors should be considered to ensure the success of FMT in humans.

Trial registration. The study was prospectively registered at the Dutch Trial registry - NTR4488 (https://www.trialregister.nl/trial/4488).

Keywords: Nanopore sequencing, MinION, gut microbiota, species engraftment, single-nucleotide variation, species-level resolution, *rrn* operon.

Introduction

Microbiome research advances rapidly and the study of microbial communities demands the implementation of sequencing strategies permitting the identification of microbial species and strains of complex ecosystems and allowing tracing of microbes engraftment in clinical trials evaluating microbiome replacement strategies. Among them, fecal microbiota transplantation (FMT) is widely recognized as an efficient therapy for *Clostridioides difficile* infection that outperforms antibiotic treatment [1]. Despite the potential risk for the transmission of infectious diseases by FMT, this approach has shown efficacy in ameliorating intestinal inflammation and metabolic disorders, such as insulin resistance [2-4]. The mode of FMT action is thought to depend on the engraftment of live microbes from donors to recipients, although some clinical trials leave the door open for hypotheses about alternative mechanisms [5]. The demonstration that the disease amelioration is linked to the ability of donor's microbes to thrive in the recipient gut is challenging but necessary to ascertain whether particular microbes play a therapeutic role in FMT trials. Metagenomics based on DNA shotgun sequencing is currently the only strategy that enables us to reach such a resolution level, but it is still very expensive, and few strain profiling results adopting such protocols have been provided in FMT studies [6-8]. Most of FMT microbiome surveys have been performed through the targeted amplification of universal gene markers, mainly including a few hypervariable regions (e.g., V3 and/or V4) of the bacterial 16S rRNA gene [9, 10]. This technique shows limitations since it only allows changes affecting the global microbiome structure to be captured and does not permit precise taxonomic identifications at the species level, mostly because of the limited resolution of this gene marker in discriminating closely related microbes. This issue is of particular relevance to accurately trace species engraftment in FMT or demonstrate microbe colonization in probiotic-based clinical trials. We previously developed a Nanopore-based amplicon sequencing method to improve the resolution of taxonomic identifications by studying the microbial genetic variability of nearly complete 16S rRNA genes [11]. This approach has been demonstrated to

perform well in a wide variety of microbiota studies [12-15]. More recently, we have also pioneered a

new methodology combining the sequencing of an extremely variable multilocus region with

sample multiplexing [16] to improve the species-level characterization of complex microbial

communities [12, 17]. The implementations mentioned above of nanopore-based technology and their

multiple applications to assess microbial communities allow us to gain knowledge in the

configuration of different analytical pipelines improving the processing of such data as updated

protocols are delivered [18-20].

We have made a great effort to compile a great deal of genetic information on *rrn* sequences from

thousands of bacterial species (*rrn_db*, freely available at https://github.com/alfbenpa/rrn_db) [16].

Nevertheless, the limited number of properly annotated DNA sequences available in public

repositories of bacteria from multiple ecosystems restricts the utilization of this approach in less

explored habitats [12, 21]. Notwithstanding, the utility of *rrn*-based analysis for surveying biological

diversity has been explored across kingdoms, showing promising results in the identification of

metazoans by metabarcoding approaches [22].

Despite the issues concerning unlinked 16S rRNA and 23S rRNA markers [23] that can be addressed

in certain bacterial groups and environmental samples via the study of individual regions, we think

the long-read amplicon sequencing of bacterial *rrn* regions could help fill the existing gap to

demonstrate the transfer and stable colonization of microbes under FMT strategies in a cost-

effective manner.

In this study, we aimed to evaluate the donor's influence on the transference of specific intestinal

microbes to FMT recipients using a unique-donor-to-multiple-recipients approach and amplicon

sequencing of the nearly complete *rrn* regions using Nanopore-based sequencing in a trial

evaluating the ability of FMT to alleviate metabolic syndrome. We obtained *de novo* information on

the *rrn* region from predominant bacterial species inhabiting the human gut, enriching the *rrn*

existing database with the annotations of dozens of microbial species and strains. We assessed to

what extent the FMT changed the gut microbiota composition of recipients at the species level and,

more importantly, tracked the strain-level variation to investigate microbiota engraftment. Overall,

our results demonstrated a specific microbial transference between donors and recipients, which

likely mediated health effects in metabolic syndrome patients, and that this transference partly

depends on the donor-recipient pair, their baseline gut microbiota configuration, and certain donor-

associated demographic variables.

Results

Assembly of human gut microbiota-derived rrn sequences

The evaluation of the gut microbiota in subjects with metabolic syndrome submitted to FMT aimed

at revealing microbiota engraftment; it required the comprehensive analysis of Nanopore

sequencing data to generate reliable genetic information supporting species- and strain-level

variation additionally to that compiled in our previous *rrn* repository [16]. During the first step of *rrn*

assembly (Figure 1), based on mapping forward reads against the nonredundant NCBI 16S database

and the selection of high-quality alignments, we detected the presence of 381 different species.

However, preliminary *rrn* assembly was initiated only for those species with at least 10X coverage

(250 in total), supporting high-quality consensus *rrn* sequences in downstream analyses. The

mapping of reverse reads against the preliminary *rrn* assemblies permitted us to confirm the

detection of 229 microbial species. Full *rrn* sequences assembled with reliable taxonomy

annotations were retrieved disregarding their individual coverage level (e.g. low, intermediate, or

high- Figure S1).

Species abundance shifts associated with FMT

Sample-derived forward and reverse reads were mapped to assess shifts in diversity and taxonomic

features. Among all reads, only 43% supported alignments with the highest quality (according to

identity and length) and were used in downstream analyses. After taxonomic assignment, we found

no large changes in any of the alpha diversity descriptors analyzed. Nevertheless, we found that

FMT increased the richness (Chao's index) of the microbiota in six out of the ten recipients (Figure

S3), which showed a gain of 31 species on average. In the remaining four recipients, there was a

reduction in richness resulting in an average loss of 16 species as a result of FMT. The beta

diversity evaluation based on the Bray-Curtis dissimilarity index indicated no major shifts in the

microbial structure of recipients as a consequence of FMT (PERMANOVA = 1.02, p = 0.403).

Nevertheless, the gut microbiota composition of the recipients was strongly influenced from greater

to lesser extents by the sequencing run (PERMANOVA = 3.69, p = 0.001), sex (PERMANOVA =

1.70, p = 0.032), and donor (PERMANOVA = 1.69, p = 0.004). The results of this multivariate

analysis are shown in Figure 2A. Globally, we observed that POST-FMT samples tended to be

mapped closer to those from donors. To further assess this hypothesis, we compared the distances

(Bray-Curtis weighted metrics) between the respective donors and the PRE-FMT microbiota or the

POST-FMT microbiota. As a result, we noted that donor microbiota-PRE-FMT pairs had a trend to

be more dissimilar than the donor microbiota-POST-FMT pairs, as indicated by a decrease in the

Bray-Curtis distance (PRE-FMT vs Donor median = 0.79, POST-FMT vs Donor median = 0.71,

average delta = 0.09, p = 0.075) (Figure 2B).

We performed an LMM analysis, which provided similar results to the beta diversity evaluation.

We found that sequencing batch, sex, and donor were the main covariates influencing the

microbiota data. Furthermore, we found that age and baseline BMI also explained the gut

microbiota variation among the subjects of this study to some extent. After including the variables

mentioned above as random effects in the model, a list of microbial species showing alterations as a

consequence of FMT was retrieved (Table 1), and some of them exhibiting an increased abundance

as a consequence of the FMT are plotted in Figure 2C. Ten different microbial species were found

to be differentially increased when paired samples obtained before and after FMT were compared,

and only three of them (e.g. *Bifidobacterium adolescentis*) seemed to show a decline because of

transplantation (Table 1).

Transferred and engrafted gut microbial species

To assess species engraftment, we next designed a proof of concept analysis to discriminate strain-

level variation and thus trace microbes engraftment more accurately. For this purpose, we selected

Parabacteroides merdae, a predominant species in the samples assessed that showed the greatest

increase as a result of FMT (Figure 3A), and *Faecalibacterium prausnitzii*, a highly abundant

species in the analyzed samples for which evident transfer between the donor-recipient pairs was

not observed (Figure 3B), to detect single-nucleotide variations (SNVs) indicative of strains

identity/transference. After a massive analysis of the total dataset merging the species-specific *rrn*

reads derived from all subjects (see methods), we identified three potential informative SNV sites in

the *P. merdae rrn*, whereas six were found in the *F. prausnitzii* counterpart (Figure 3C-D). In an

attempt to explore the natural existence of such genotypic variation inferred from our analysis in a

predictive manner, we looked for the presence of similar variants in the NCBI non-redundant

nucleotide database using the 100 nt surrounding region as a query for all *P. merdae* SNVs. The

inferred genetic variability was replicated at least for SNV-3527, where multiple matches presented

A and G nucleotides at this position (Figure S4). When we studied the nucleotide frequencies of

these particular SNVs across samples, a clear transfer pattern of the *P. merdae* genotype from donor

to recipient was detected, as indicated by the decreased genetic distance (Bray-Curtis index for

assessing the abundance of the four nucleotides in all SNVs at once, one-sided Wilcoxon Signed-

Rank test = 11, p = 0.051) between donor samples and POST-FMT compared to that of POST-FMT

and PRE-FMT pairs (Figure 3E). By contrast, the genetic distances retrieved after the comparison

of *F. prausnitzii* genotypes between POST-FMT or PRE-FMT samples and their donors did not

indicate the transfer of strains belonging to this species from donors to recipients (Figure 3F).

Comparative analysis of the delta Bray-Curtis distance (POST vs DONOR minus PRE vs DONOR

samples) between both species also suggests that only *P. merdae* was transferred from donors to

recipients, since the distance between POST-FMT and donor samples tended to decrease (Figure

3G, *p* = 0.056). The direct Sanger sequencing of *P. merdae* SNV-1254, SNV-2293, and SNV-3527

in the two recipients and their common donor supported the results of our long-read-based

assessment of FMT (Figure 3H). Globally, we observed the presence of a mixture of strains in some

of the samples analyzed, given the basecalling profile visualized in the electropherograms (e.g.,

SNV-3527 of Donor1 and 06-Pre samples). This pattern of strain coexistence was more evident in

the POST-FMT samples of both analyzed recipients (see SNV-2293 and SNV-3527 in Figure 3H).

The predominant haplotype observed in Donor1 (A1254-C2293-A3527) was transferred to the two

recipients explored and became dominant after FMT. Similar patterns of transfer were observed in

other donor-recipient pairs. Globally, our results demonstrated that the increased abundance of *P.*

merdae after the FMT was a direct consequence of donor strain transmission. This is likely the case

for other bacterial species that increased as a result of FMT.

Correlation between clinical variables and species abundance after FMT

Among the multiple clinical variables evaluated in these subjects with metabolic syndrome, FMT

tended to induced positive changes in the markers of glucose metabolism and blood pressure in the

recipients (Table 2). Fasting insulin (*p* = 0.030), fasting glucose (*p* = 0.074), and consequently

HOMA-IR (*p* = 0.005) tended to improved in the recipients after FMT. In line with these findings,

there was an apparent decrease in the glycosylated hemoglobin concentration after FMT (*p* =

0.060). On the other hand, systolic and diastolic blood pressures were lowered as a consequence of

donor FMT treatment, being reduced by 10% (*p* = 0.028) and 19% (*p* = 0.0008), respectively (Table

2). The concentrations of fecal SCFAs, such as fecal butyrate (*p* = 0.018) and acetate (*p* = 0.033),

were decreased after FMT, whereas that of propionate was not (*p* = 0.280). Regarding markers of

lipid metabolism, HDL cholesterol levels in plasma were reduced (*p* = 0.032). By computing rank-

based Kendall's *τ* (tau) parameter, we established associations between these clinical variables and

the alterations in the abundance of microbial species as a consequence of FMT (see Table 1).

Species such as *P. merdae*, which was increased after FMT, were negatively correlated with

diastolic blood pressure (τ = -0.44 and FDR = 0.062), whereas *A. shahii* was negatively correlated

with the abundance of fecal acetate and butyrate (τ = -0.50, -0.42 and FDR = 0.017, 0.042,

respectively). The above correlations were not detected for other closely related and abundant

species (e.g., *Parabacteroides johnsonii*, *Parabacteroides distasonis*, *Parabacteroides goldsteinii,*

Alistipes finegoldii, Alistipes onderdonkii, Alistipes putredinis or *Alisti*pes *indistinctus*).

Donors' influence on species engraftment and amelioration of metabolic dysfunction

After showing that microbiota engraftment took place between donor and recipients, and that

reversion of certain metabolic alterations may have been guided by particular microbial species

transferred from donors to recipients, we wanted to explore to what extent the donors and their

microbiota configurations can explain the FMT success. For such an aim, we ranked the metabolic

improvement individually (from 1 to 10 , from the lowest to the highest), including the reduction of

blood pressure, fasting glucose and insulin levels, glycosylated hemoglobin, HOMA-IR, weight

loss, and BMI of the ten subjects of the FMT. Then, the health score resulting from the sum of all

scores for every subject was estimated. This score was related to t the dissimilarity index (Bray-

Curtis) between the baseline microbiota configuration of donor-recipient pairs and the community

structural shift resulting from the FMT in every recipient, taking into account their donor pair

(Figure 4). Globally, we observed that metabolic health scores correlate to some extent with the

baseline dissimilarity between donors and recipients microbiotas and the community structural

changes after FMT. Consequently, the more dissimilarity in the baseline gut microbiota of donors

and recipients, the lower the benefits to revert the metabolic dysfunction in recipients and less

impact on their gut microbiota structure after FMT. The above observation applies to almost all

donor-recipient pairs, except for those including the Donor2. Although we observed a linear

relationship between the averaged dissimilarity between donor-recipient pairs and the respective

shift in the community structure after FMT (adjusted R-squared: 0.956, p = 0.013), a rank-based approach (Kendall's tau) showed those parameters were not well correlated with the metabolic health score ($\tau \geq$ -0.24, p $\geq$ 0.381). Therefore, additional traits yet to explore in donor-recipient pairs should help in the selection of subjects for succeeding in FMT. In line with this hypothesis, we finally evaluated the relationship between the metabolic health score and the recipients sex, showing that women exhibited a larger improvement in metabolic parameters than men (women [N = 5] = 49.6 ± 13.8 health score, men [N = 5] = 33.6 ± 11.3 health score, *t*-test = 6.62, p = 0.003). Moreover, we found a strong rank-based correlation between the age difference of donor-recipient pairs (donor age - recipient age) and the metabolic health score (Kendall's τ = -0.51, p = 0.046), suggesting that the youngest the donor (with respect to the recipient), the greater reversion of metabolic dysfunction.

Discussion

Here, we explored the *rrn* amplicon sequencing of the human gut microbiota through Nanopore sequencing and the *de novo* assembly of sequences of this multilocus hypervariable region to gain insights into the corresponding genetic markers and to demonstrate their potential use for deeply characterizing complex communities at the species and strain levels, permitting the assessment of microbe engraftment between donors and recipients in an FMT clinical trial. The utility of *rrn* operon sequencing via the MinION[TM] platform has already demonstrated a notable performance in resolving microbial communities at the species level, with the potential to define strain variability for particular species [24, 25].

We provided *de novo* reliable assemblies for more than two hundred *rrn* region sequences of human gut microbes. We were able to reconstruct a large proportion of *rrn* regions from approximately 40 new microbial species, and more than 150 new strains were absent in the first database release [16]. These numbers, limited mainly by sequencing coverage, could also be reduced by the DNA

extraction method used, in which a physical disruption step was omitted to maintain DNA integrity

and ensure the successful amplification of the 5 kbp amplicons. However, the homogeneous

processing of all samples resulted in comparable microbial profiles before and after FMT and did

not affect the species-specific evaluation conducted in this study.

By using a dataset consisting of approximately 400 K Nanopore reads derived from 24 samples,

employing the updated *rrn* database, and controlling the covariates influencing the microbiota, we

revealed changes in the human gut microbiota at the species and strain levels as a consequence of

an FMT intervention. This long amplicon-based approach enabled us to detect increases in species

such as *Parabacteroides merdae*, *Butyricimonas virosa*, *Prevotella copri*, and *Sutterella*

massiliensis as a result of FMT. In addition to age, sex, sequencing runs (recognized stochastic run-

to-run variation confounding the interpretation of microbiota data [26, 27]), and baseline BMI, we

found that the donor was a critical covariate contributing to the impact of FMT on the recipient

microbiota when using a unique donor for multiple recipients. These results are in agreement with

those previously published on the basis of short reads from the V4 hypervariable region of the 16S

rRNA bacterial gene explored in a larger cohort of subjects [3], for instance, the significant increase

of *P. copri* after FMT; however, our results outperform the taxonomic resolution achieved

previously, thus providing a more accurate gut microbiota survey. Interestingly, a recent FMT study

on the treatment of ulcerative colitis (UC) in which the microbiota was analyzed by sequencing

short reads also showed increases in *Sutterella* species as a consequence of the intervention [28].

Similarly, the abundance of *Butyricimonas* species was shown to be increased as a consequence of

an FMT intervention aimed at eradicating antibiotic-resistant bacteria [29]. These findings suggest that

some species may frequently show changes as a result of FMT in humans, regardless of the

condition of the recipient. Nevertheless, the presumable transfer of species between donors and

recipients and their engraftment could not be confirmed in those studies based on short-read

amplicon technology due to the lack of sufficient resolution. Additionally, potential replacement

with closely related species in the recipient could be impossibly monitored in conventional

microbiota surveys based on short reads. To shed light on the ability of the *rrn* sequencing approach

to assess SNVs that are likely associated with strain diversity, thus demonstrating donors' microbe

engraftment in recipients, we designed a proof of concept analysis selecting *P. merdae*, a species

that exhibited a remarkable increase in the recipients' gut after FMT, and *F. prausnitzii*, which

seemed to not be affected by FMT intervention. The combined information on different informative

SNVs from the *P. merdae rrn* sequence clearly suggests that the increase in this species observed in

recipients after undergoing FMT was likely because of strain transfer from the donors. This

inference is further supported by the existence of nucleotide variants inferred from Nanopore data

processing in either the samples evaluated in the present study or the genomic data present in public

repositories and derived from different sequencing methodologies.

The taxonomic resolution of our sequencing approach, proving the transference of specific bacterial

strains from donors to recipients, helped draw partial community structures at the species level and

determine compositional distances and their shifts as a consequence of the FMT. This strategy

helped to more firmly support a causal relationship between the changes in the gut microbiota and

the improvements in the recipient's clinical variables. In this regard, we attempted to demonstrate

that overall improvement of metabolic health (global score) was associated with baseline

dissimilarity between microbial communities of donors and recipients and with their microbiota

configuration shifts after FMT. We observed that the more similar is the gut microbiota of the

donor-recipient pairs at baseline, the better is the microbiota transference and the improvement in

metabolic parameters. However, this correspondence was not proven for all pairs; possibly,

additional traits of donors and recipients yet to be determined need to be considered for an optimal

selection of donors to ensure the clinical success of the FMT. One of those unknown additional

factors could be the sex of the recipient and donor. Indeed, we observed that women recipients

exhibited higher metabolic health score after FMT than men. Given that the sex of donors and

recipients was matched, further studies with a mixed-sex distribution would be needed to determine

i) if women's fecal material can be better engrafted regardless of the recipient sex, or ii) whether

women respond better to FMT irrespective of the sex origin of the FMT material. We have

previously unveiled differential responses of women and men gut microbiota to a particular dietary

intervention, suggesting that such disparate sex-associated microbial configurations and related

traits could plausibly explain the success of FMT [30]. Nonetheless, previous FMT interventions in

the context of IBS show no differential response between women and men [31]. Finally, an individual-

specific response to FMTs cannot be disregarded as in our trial the material from Donor1 exerted

the most positive impact on the metabolic health of their three different recipients. Consequently,

the role of the personal features of the donor, particularly those influencing the gut microbiota

composition and metabolic potential (e.g. diet, lifestyle) as well as other host covariates (e.g. age)

linked to successful FMTs should be further investigated to better guide donor selection and

identify microbial signatures improving metabolic health.

Conclusions

The molecular methods and data analyses presented here enabled reliable microbial surveys at the

species level and the inference of strain variations, taking into account the most abundant human

intestinal bacterial species. To the best of our knowledge, this is the first microbiota assessment of

fecal samples derived from donors and recipients involved in a FMT clinical trial to demonstrate

transference and engraftment of specific bacterial species by exploring the genetic diversity at the

strain level using amplicon sequencing. The genetic and taxonomy information retrieved from

advanced sequencing methods used here permitted us to evaluate the donors' influence on the gut

microbiota transference and clinical success of the FMT. The determinants of the success include a

high relatedness between the donor-recipient baseline microbiota configurations, the sex and the

age of the donor, which seem to influence the engraftment of intestinal microbes and the

amelioration of metabolic syndrome markers. Further studies should be warranted to confirm our

findings in larger intervention trials and demonstrate their predictive value.

Materials and Methods

Subjects, samples, and clinical data

Samples were obtained upon receiving informed consent from participants in a previous FMT

clinical trial carried out under the framework of the MyNewGut project (http://www.mynewgut.eu/)

between September 2014 and March 2017. The details of the study design are publicly available

elsewhere [3]. Briefly, a total of twenty-four fecal samples were analyzed in the present study.

Twenty samples were obtained from 10 recipients undergoing allogenic FMT, who provided one

sample before (PRE-FMT fecal samples) and one sample 4 weeks after (POST-FMT fecal samples)

the intervention. The samples of 4 donors were also analyzed, and the effect of this variable on the

recipients was considered in the data analysis. This subset of samples was mainly assessed to

explore differential species engraftment in multiple FMT recipients with common donors. The

sample size evaluated in this study is still representative according to parameters described

previously for this intervention arm (two-sided significance level of 0.05 and power of 80%,

minimum sample size = 10) [3]. Normality test was completed prior to execute parametric or non-

parametric tests to assess changes in the clinical variables explored.

DNA extraction, multilocus amplification, and sequencing

Microbial DNA was recovered from 100 mg feces samples by using the QIAamp® Fast DNA Stol

Mini kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions and omitting cell

disruption by mechanical methods (bead-beating) to preserve DNA with the high molecular weight

needed to amplify amplicons of 5 kbp in length. The *rrn* region comprising the nearly full-length

bacterial RNA ribosomal operon (16S, ITS, and 23S) was amplified as previously reported [16]. Dual-

barcoded purified PCR products were mixed in equimolar proportions before sequencing library

preparation. In total, three different libraries were prepared from ~1 µg of mixed amplicon DNA

(containing 7, 8, or 10 barcoded samples) using the SQK-LSK108 sequencing kit (Oxford

Nanopore Technologies, Oxford, UK) following the manufacturer's instructions to produce 1D

reads. Each library was individually loaded into FLO-MIN107 (R9.5) flow cells (Oxford Nanopore

Technologies, Oxford, UK), and sequencing was carried out in a portable MinION™ MkIb

sequencer (Oxford Nanopore Technologies, Oxford, UK), operated with *MINKNOW* v1.10.23

software (Oxford Nanopore Technologies, Oxford, UK). Flow cells were primed according to the

manufacturer's instructions, and an ~18 h run of 1D sequencing (outperforming the 2D approach for

this multiplex and amplicon setup, and improving de-multiplexing by using the barcode+primer

tags) was then executed for each library and flow cell.

Data preprocessing

Fast5 files were processed with the *Albacore* v2.1.3 basecaller and default configuration (minimum

q-score filter = 7), and *fasta* files were retrieved for downstream analyses. The barcode and primer

(forward or reverse) sequence information was used for demultiplexing into the DNA reads

generated from forward and reverse primers according to previous procedures [16]. A size filtering

step was configured to retain those reads of least 1,500 nt in length. Barcode and primer sequences

were then removed by trimming 50 nucleotides at the 5' end of forward and reverse reads. The

sequencing chemistry and Nanopore data-processing algorithms were the latest available by January

2018, when data was generated. A high-performance basecaller released after the end of this

analysis was tested on fast5 data produced getting no significant difference in sequences quality

(Figure S4).

Assembly of rrn regions

The study design and the assembly of *rrn* sequences are graphically explained in Figure 1. This

pipeline carries out what we refer to as the strand-wise and reference-assisted method for *rrn* region

reconstruction via the following steps:

•   Demultiplexing of Nanopore data using barcode and primer sequence information

365   generated two datasets per sample comprising DNA reads led from the 5´ end of *rrn*

366   amplicons (initiated from forward primer mapping at the 16S rRNA gene start site =

367   forward reads), and DNA reads beginning at the 3' end of *rrn* amplicons (initiated from

368   reverse primer mapping at the 23S rRNA gene end site = reverse reads).

369   •   Reads from all 24 samples were merged into forward and reverse read subsets.

370   •   Forward read binning into precise microbial species via competitive alignment against the

371   nonredundant 16S NCBI database (release January 2018). The *LAST* aligner [32] with a -s 2 -

372   q 1 -b 1 -Q 0 -a 1 -r 1 configuration was used for this purpose. The alignment score was the

373   main criterion for selecting top hits. In the case of multiple hits with the same top score, the

374   alignment was discarded for downstream processing. High-quality alignments were retained

375   according to sequence identity (sequence identity above the 33rd percentile $\geq 85\%$) and

376   length information (alignments ≥ 1500 nt).

377   •   A maximum of 500 forward reads per species (randomly shuffled), producing high-quality

378   alignments, were selected, and aligned through iterative refinement methods implemented

379   in *MAFFT* v7.310 with the default parameters [33], and the consensus sequence was then

380   obtained by using the *hmmbuild* and *hmmemit* algorithms implemented in *HMMER3* [34]. No

381   significant improvements in consensus *rrn* sequences were obtained using more than 500

382   sequences per species. Assemblers supporting nanopore data, temporary available at the

383   date of analysis, were tested to produce *de novo* consensus sequences but they failed to

384   produce outputs supporting the expected species-level diversity because of the coverage

385   bias across data.

386   •   Consensus sequences were annotated according to the NCBI taxonomy and used as a

387   reference to bin reverse reads in a similar manner as that used for forward reads. The

388   thresholds for the selection of high-quality alignments based on reverse reads and the

preliminary assembly of *rrn* regions were 85% sequence identity (upper 45th percentile)

and $\geq$ 3500 nt in length.

• Reverse reads were compared against the consensus sequences obtained from the forward

read assemblies and binned into the respective species according to NCBI taxonomic

annotation. A maximum of 500 reverse reads per species (randomly shuffled), producing

high-quality alignments, were selected, merged with a maximum of 500 forward reads

assigned reciprocally to the same species (randomly shuffled), and aligned through iterative

refinement methods implemented in *MAFFT*, after which a new consensus sequence was

obtained by using the *hmmbuild* and *hmmemit* algorithms implemented in *HMMER3*.

• Final consensus sequences were annotated according to the concordant NCBI taxonomy,

obtained from forward and reverse reads, and used as a reference to study the abundance

and prevalence of gut microbiota members at the species level.

• A BLAST-based search of the *rrn* assemblies against the nonredundant nucleotide and

reference 16S NCBI databases was performed to evaluate the representation of the

reconstructed and identified operons in public databases. The identification of *rrn*

assemblies was compared against the SILVA database [35] with SINA aligner [36]. Top hit

selection was based on the taxonomy score, TS = Log_{10}[alignment score*sequence

identity*alignment length].

Long-read mapping and variant calling

The *rrn* database [16] was updated with more than two hundred new *rrn* operon sequences assembled

in this study (labeled operONT) and re-annotated to different taxonomic levels according to the

NCBI taxonomy database. The mapping of sample reads was performed against *rrn_DBv2*

(https://github.com/alfbenpa/rrn_DBv2) by competitive alignment using the *LAST* aligner, as stated

above. Taxonomic assignment was based on the best hit retrieved by the calculation of TS (see

above), and reliable taxonomic assignments were filtered to those based on alignments with at least

85% sequence identity that were longer than 1500 nt. Species read counts were normalized using

the *compositions::clr* R v3.6 function, and a covariate-controlling linear mixed model-based

method was used to explore the differential abundance of taxonomic features between conditions as

stated in the paragraphs below. Species engraftment was evaluated by the selection of all sample

reads mapped to the *Parabacteroides merdae* and *Faecalibacterium prausnitzii* species. To detect

informative sites, based on single-nucleotide variants that were probably linked to strain variation,

we first proceeded to map all sample reads matching these species to the respective reference *rrn*

sequences using *LAST* aligner and the parameters set across this study (see *rrn* assembly). The *maf-*

convert algorithm (*LAST* aligner toolkit) was used to generate the respective *SAM* files. The

algorithms compiled within *SAMtools* v1.3.1 were used to index, order, and pileup reads as well as

to retrieve information regarding the nucleotide frequency per site and coverage (*vcf* files). Site

selection was based on *rrn* positions showing no nucleotide dominance (<70% frequency for a

single nucleotide), without reductions in the coverage (>75% of the relative coverage), thus

obtaining sites with a balanced representation of at least the two predominant alleles likely

representing strain variation across all samples. Second, the sample reads that were mapped to the

reference species were individually assessed to detect meaningful changes in intestinal bacterial

genotypes between the recipients PRE-FMT and POST-FMT samples. The changes in nucleotide

frequencies at all selected SNV sites (three for *P. merdae* and six for *F. prausnitzii*) were assessed

by calculating the Bray-Curtis dissimilarity index between pairs of POST-FMT and PRE-FMT

samples or POST-FMT and corresponding donor samples following statistical evaluation by using

the Wilcoxon signed rank test for paired samples.

Sanger sequencing and validation of Parabacteroides merdae SNVs

The reference *rrn* sequence from *P. merdae* was submitted to the Primer-BLAST web server

(https://www.ncbi.nlm.nih.gov/tools/primer-blast/) to retrieve species-specific primer pairs to

selectively amplify this *rrn* sequence and primers flanking SNV-1254, SNV-2293, and SNV-3527

for Sanger sequencing. Comparisons against the nonredundant NCBI database and *Parabacteroides*

merdae [taxid:46503] as a reference organism were fixed as checking parameters for primer

prediction. The *rrn* region from *P. merdae* was amplified via 28 PCR cycles including the following

stages: 95°C for 20 s, 61°C for 30 s, and 72°C for 150 s. Phusion High-Fidelity Taq Polymerase

(Thermo Scientific) and the pm623 (TGCCGTTGAAACTGGGTTACTTGA) and pm3785

(TTTCGCACAGCCATGTGTTTTGTT) primer pairs were used in the amplification reaction. The

PCR products were cleaned with an Illustra GFX PCR DNA and gel band purification kit (GE

Healthcare, Chicago, IL, USA) and sequenced by using Sanger technology in an ABI 3730XL

sequencer (STAB-VIDA. Caparica, Portugal) using the primers described above (pm623 and

pm3785) in addition to the pm1855 (ACCCCTTACGGAGTTTATCGTGGA) and pm2686

(TTCGCGTCTACTCACTCCGACTAT) primers. The sequencing electropherograms from the *ab1*

files were visualized with FinchTV v1.4.0 (Geospiza Inc.).

Diversity and taxonomic analyses at the species level

The prevalence and abundance of a total of 2,519 species contained in the database were evaluated,

and diversity analyses were completed taking into account the species with a > 0.01% relative

abundance on average (~250 species in total). Alpha diversity descriptors such as Chao's index,

Shannon's entropy, Simpson's reciprocal index, and dominance were obtained by using *QIIME*

v1.9.1 [37]. Similarly, *QIIME* was used to calculate the Bray-Curtis dissimilarity indexes among

samples and to perform multivariate exploratory (principal coordinate analysis - PCoA) and

statistical (PERMANOVA) analyses. A linear mixed model (LMM – *nlme::lme* R v3.6 function)

analysis was also conducted on clr-transformed data to detect differential features in the microbiota

before and after the intervention. The inherent variation due to individual features was set as a

random effect for each variable analyzed (fixed effect). The possible covariates of the clinical and

fecal microbiota that showed differences between the study groups (p-value ≤ 0.05) were selected.

Recognized covariates of microbiota, such as age, sex, baseline BMI, and sequencing batch, were

identified as significant in this study and included as random effects in the LMM. To identify

microbial species potentially linked to clinical variables that were altered as a consequence of FMT,

Kendall's τ (tau) parameter was estimated between variable pairs and corrected for multiple testing

using the false discovery rate (FDR) approach. Associations were selected according to an FDR p-

value ≤ 0.1. Graphics were generated in R v3.6 using the *ggplot2* package.

List of abbreviations

470   BMI, body mass index; FDR, false discovery rate; FMT, fecal microbiota transplantation; LMM,

471   linear mixed model; NCBI, National Center for Biotechnology Information; MAG, metagenome-

472   assembled genome; PCoA, principal coordinate analysis; PCR, polymerase chain reaction; RDP,

473   Ribosomal Database Project; rrn, bacterial ribosome RNA operon (16S-ITS-23S); SNP, single-

474   nucleotide polymorphism, SNV, single-nucleotide variation; UC, ulcerative colitis.

Declarations

Ethics approval and consent to participate

The study was prospectively registered at the Dutch Trial registry

(https://www.trialregister.nl/trial/4488) and was conducted according to the guidelines laid down in

the Declaration of Helsinki and the ethical standards of the responsible local committee on human

experimentation of the Amsterdam UMC (location AMC) [3]. The study was registered on August 1st,

2014. The first participant was enrolled on September 1st, 2014.

Availability of data and material

The Albacore-basecalled *fast5* files obtained from specific runs are publicly available in the

European Nucleotide Archive under accession number PRJEB33947. The updated *rrn* database

(*rrn_DBv2*) is publicly accessible at the GitHub repository https://github.com/alfbenpa/rrn_DBv2.

Competing interests

The authors have no conflicts of interest to declare.

Funding

This study was supported by the EU Project MyNewGut (No. 613979) of the European

Commission 7th Framework Programme and grant PID2020-119536RB-I00 from the Ministry of

Science and Innovation (Spain). This research study was also completed thanks to the CP19/00132

grant from Miguel Servet programe to ABP from the Institute of Health Carlos III (ISCIII) and its

co-funding from the European Social Fund (ESF/FSE).

Authors' contributions

ABP conceived and designed the study. ABP performed the experimental sequencing research and

data analysis, and AVH and MN performed the clinical research. ABP and YS directed the study.

ABP and YS wrote the manuscript. All authors reviewed and approved the final version of the

manuscript.

Acknowledgements

The authors thank the Bioinformatics and Biostatistics Unit from Principe Felipe Research Center

(CIPF) for providing access to the cluster, co-funded by European Regional Development Funds

(ERDF/FEDER) in Valencian Community 2014-2020.

References

1.   Cammarota G, Ianiro G, Gasbarrini A. Fecal microbiota transplantation for the treatment of
Clostridium difficile infection: a systematic review. J Clin Gastroenterol 2014; 48:693-702.

2.   Chu ND, Crothers JW, Nguyen LTT, Kearney SM, Smith MB, Kassam Z, et al. Dynamic
Colonization of Microbes and Their Functions after Fecal Microbiota Transplantation for Inflammatory
Bowel Disease. mBio 2021; 12:e0097521.

3.   Hartstra AV, Schuppel V, Imangaliyev S, Schrantee A, Prodan A, Collard D, et al. Infusion of donor
feces affects the gut-brain axis in humans with metabolic syndrome. Mol Metab 2020; 42:101076.

4.   Sokol H, Landman C, Seksik P, Berard L, Montil M, Nion-Larmurier I, et al. Fecal microbiota
transplantation to maintain remission in Crohn's disease: a pilot randomized controlled study. Microbiome
2020; 8:12.

5.   Ott SJ, Waetzig GH, Rehman A, Moltzau-Anderson J, Bharti R, Grasis JA, et al. Efficacy of Sterile
Fecal Filtrate Transfer for Treating Patients With Clostridium difficile Infection. Gastroenterology 2017;
152:799-811 e7.

6.   Aggarwala V, Mogno I, Li Z, Yang C, Britton GJ, Chen-Liaw A, et al. Precise quantification of
bacterial strains after fecal microbiota transplantation delineates long-term engraftment and explains
outcomes. Nat Microbiol 2021; 6:1309-18.

7.   Verma S, Dutta SK, Firnberg E, Phillips L, Vinayek R, Nair PP. Identification and engraftment of
new bacterial strains by shotgun metagenomic sequence analysis in patients with recurrent Clostridioides
difficile infection before and after fecal microbiota transplantation and in healthy human subjects. PLoS One
2021; 16:e0251590.

8.   Wilson BC, Vatanen T, Jayasinghe TN, Leong KSW, Derraik JGB, Albert BB, et al. Strain
engraftment competition and functional augmentation in a multi-donor fecal microbiota transplantation trial
for obesity. Microbiome 2021; 9:107.

9.   Rinott E, Youngster I, Yaskolka Meir A, Tsaban G, Zelicha H, Kaplan A, et al. Effects of Diet-
Modulated Autologous Fecal Microbiota Transplantation on Weight Regain. Gastroenterology 2021;
160:158-73 e10.

10.   Yu EW, Gao L, Stastka P, Cheney MC, Mahabamunuge J, Torres Soto M, et al. Fecal microbiota
transplantation for the improvement of metabolism in obesity: The FMT-TRIM double-blind placebo-
controlled pilot trial. PLoS Med 2020; 17:e1003051.

11.   Benitez-Paez A, Portune KJ, Sanz Y. Species-level resolution of 16S rRNA gene amplicons
sequenced through the MinION portable nanopore sequencer. Gigascience 2016; 5:4.

12.   Cusco A, Catozzi C, Vines J, Sanchez A, Francino O. Microbiota profiling with long amplicons
using Nanopore sequencing: full-length 16S rRNA gene and whole rrn operon. F1000Res 2018; 7:1755.

13.   Sakai J, Tarumoto N, Kodana M, Ashikawa S, Imai K, Kawamura T, et al. An identification protocol
for ESBL-producing Gram-negative bacteria bloodstream infections using a MinION nanopore sequencer. J
Med Microbiol 2019; 68:1219-26.

14.   Shin H, Lee E, Shin J, Ko SR, Oh HS, Ahn CY, et al. Elucidation of the bacterial communities
associated with the harmful microalgae Alexandrium tamarense and Cochlodinium polykrikoides using
nanopore sequencing. Sci Rep 2018; 8:5323.

15.   Shin J, Lee S, Go MJ, Lee SY, Kim SC, Lee CH, et al. Analysis of the mouse gut microbiome using
full-length 16S rRNA amplicon sequencing. Sci Rep 2016; 6:29681.

16.   Benitez-Paez A, Sanz Y. Multi-locus and long amplicon sequencing approach to study microbial
diversity at species level using the MinION portable nanopore sequencer. Gigascience 2017; 6:1-12.

17.   Kerkhof LJ, Dillon KP, Haggblom MM, McGuinness LR. Profiling bacterial communities by
MinION sequencing of ribosomal operons. Microbiome 2017; 5:116.

18.   Ciuffreda L, Rodriguez-Perez H, Flores C. Nanopore sequencing and its application to the study of
microbial communities. Comput Struct Biotechnol J 2021; 19:1497-511.

19.   Kerkhof LJ. Is Oxford Nanopore sequencing ready for analyzing complex microbiomes? FEMS
Microbiol Ecol 2021; 97.

20.   Pavlovic J, Cavalieri D, Mastromei G, Pangallo D, Perito B, Marvasi M. MinION technology for
microbiome sequencing applications for the conservation of cultural heritage. Microbiol Res 2021;
247:126727.

21.   Peker N, Garcia-Croes S, Dijkhuizen B, Wiersma HH, van Zanten E, Wisselink G, et al. A
Comparison of Three Different Bioinformatics Analyses of the 16S-23S rRNA Encoding Region for Bacterial
Identification. Front Microbiol 2019; 10:620.

22.   Krehenwinkel H, Pomerantz A, Henderson JB, Kennedy SR, Lim JY, Swamy V, et al. Nanopore
sequencing of long ribosomal DNA amplicons enables portable and simple biodiversity assessments with high
phylogenetic resolution across broad taxonomic scale. Gigascience 2019; 8.

23.   Brewer TE, Albertsen M, Edwards A, Kirkegaard RH, Rocha EPC, Fierer N. Unlinked rRNA genes
are widespread among bacteria and archaea. ISME J 2020; 14:597-608.

24.   Dowden RA, McGuinness LR, Wisniewski PJ, Campbell SC, Guers JJ, Oydanich M, et al. Host
genotype and exercise exhibit species-level selection for members of the gut bacterial communities in the
mouse digestive system. Sci Rep 2020; 10:8984.

25.   Ibironke O, McGuinness LR, Lu SE, Wang Y, Hussain S, Weisel CP, et al. Species-level evaluation
of the human respiratory microbiome. Gigascience 2020; 9.

26.   Gibbons SM, Duvallet C, Alm EJ. Correcting for batch effects in case-control microbiome studies.
PLoS Comput Biol 2018; 14:e1006102.

27.   Schloss PD, Gevers D, Westcott SL. Reducing the effects of PCR amplification and sequencing
artifacts on 16S rRNA-based studies. PLoS One 2011; 6:e27310.

28.   Paramsothy S, Nielsen S, Kamm MA, Deshpande NP, Faith JJ, Clemente JC, et al. Specific Bacteria
and Metabolites Associated With Response to Fecal Microbiota Transplantation in Patients With Ulcerative
Colitis. Gastroenterology 2019; 156:1440-54 e2.

29.   Bilinski J, Grzesiowski P, Sorensen N, Madry K, Muszynski J, Robak K, et al. Fecal Microbiota
Transplantation in Patients With Blood Disorders Inhibits Gut Colonization With Antibiotic-Resistant
Bacteria: Results of a Prospective, Single-Center Study. Clin Infect Dis 2017; 65:364-70.

30.   Benitez-Paez A, Hess AL, Krautbauer S, Liebisch G, Christensen L, Hjorth MF, et al. Sex, Food, and
the Gut Microbiota: Disparate Response to Caloric Restriction Diet with Fiber Supplementation in Women
and Men. Mol Nutr Food Res 2021; 65:e2000996.

31.   El-Salhy M, Casen C, Valeur J, Hausken T, Hatlebakk JG. Responses to faecal microbiota
transplantation in female and male patients with irritable bowel syndrome. World J Gastroenterol 2021;
27:2219-37.

32.   Kielbasa SM, Wan R, Sato K, Horton P, Frith MC. Adaptive seeds tame genomic sequence
comparison. Genome Res 2011; 21:487-93.

33.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in
performance and usability. Mol Biol Evol 2013; 30:772-80.

34.   Eddy SR. Accelerated Profile HMM Searches. PLoS Comput Biol 2011; 7:e1002195.

35.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA
gene database project: improved data processing and web-based tools. Nucleic Acids Res 2013; 41:D590-6.

36.   Pruesse E, Peplies J, Glockner FO. SINA: accurate high-throughput multiple sequence alignment of
ribosomal RNA genes. Bioinformatics 2012; 28:1823-9.

37.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME
allows analysis of high-throughput community sequencing data. Nat Methods 2010; 7:335-6.

Tables

Table 1. Changes in abundance and prevalence of gut microbial species after FMT.

Species	Abundance PRE-FMT (N = 10)[1]	Abundance POST-FMT (N = 10)[1]	LMM estimate[2]	Prevalence PRE-FMT (N = 10)	Prevalence POST-FMT (N = 10)	*p*-value
Parabacteroides merdae	0.58 ± 1.37	2.80 ± 0.79	2.07	100%	100%	0.002
Prevotella copri	-0.04 ± 0.50	0.90 ± 0.57	1.35	90%	100%	0.033
Sutterella massiliensis	0.44 ± 1.61	2.36 ± 1.53	1.64	90%	100%	0.003
Paraprevotella clara	-0.31 ± 1.14	1.18 ± 1.26	1.49	70%	100%	0.016
Alistipes shahii	1.36 ± 0.87	2.27 ± 0.66	1.05	90%	100%	0.005
Butyricimonas virosa	-0.38 ± 0.86	0.43 ± 0.92	1.01	80%	100%	0.013
Bacteroides eggerthii	-0.03 ± 0.89	0.50 ± 0.57	0.49	100%	100%	0.029
Clostridium saccharolyticum	-0.03 ± 0.84	-0.54 ± 0.76	-0.57	80%	100%	0.076
Bifidobacterium adolescenti	0.38 ± 0.54	-0.50 ± 0.91	-0.92	70%	70%	0.027
Phascolarctobacterium faecium	1.64 ± 1.96	0.51 ± 1.45	-1.47	70%	90%	0.058

600 1 Data expressed as the mean of clr-normalized counts ± standard deviation (sd). N, sample size.

601 2 LMM estimate resulting from the global pattern of abundance in POST-FMT samples with
602 respect to baseline information (PRE-FMT samples). Species altered as a consequence of FMT
603 ordered by the size effect (LMM estimate).

606   Table 2. Clinical variables altered after FMT.

Clinical outcome	PRE-FMT (N = 10)[1]	POST-FMT (N = 10)[1]	Statistics
women = 5 (50%)			
age (years) = 60.2 ± 6.4			
BMI (kg/m^2) = 32.9 ± 3.5			
Diastolic blood pressure	87.4 ± 8.2	72.0 ± 9.9	v = -16.9, $p < 0.001$
Systolic blood pressure	138.6 ± 15.3	124.5 ± 14.2	v = -14.2, $p = 0.028$
Fasting glucose (mmol/L)	5.49 ± 0.35	5.31 ± 0.50	v = -0.18, $p = 0.074$
Fasting insulin (mg/dL)	84.7 ± 30.9	63.6 ± 21.9	v = -19.0, $p = 0.030$
HOMA-IR	2.96 ± 0.99	2.16 ± 0.78	v = -0.93, $p = 0.005$
Hba1c (mmol/L)	36.5 ± 4.1	35.7 ± 3.6	v = -0.96, $p = 0.060$
Plasma HDL (mmol/L)	1.59 ± 0.29	1.40 ± 0.23	v = -0.191 $p = 0.032$
Fecal butyrate (µmol/g)	88.6 ± 44.2	60.3 ± 40.9	v = -31.6, $p = 0.018$
Fecal acetate (µmol/g)	434.0 ± 148.3	308.2 ± 121.1	v = -135.7, $p = 0.033$

607   1 Data expressed as the mean ± standard deviation (sd). N, sample size.

608   v = variation between groups analyzed by applying an LMM (PRE-FMT group as reference), HDL
609   = high-density lipoprotein, Hba1c = glycosylated hemoglobin.

Figure legends

Figure legends

Figure 1. Graphical description of the study. Data acquisition and processing steps, including

sample selection, amplicon sequencing, and the general pipeline for the *de novo* assembly of rrn

regions from human gut microbiota, are depicted.

Figure 2. Beta diversity of the microbial communities assessed by *rrn* sequencing. A - Scatter plot

compiling data from the multivariate analysis (principal coordinate analysis - PCoA) of the

microbiota from recipients and donors participating in FMT. The donor and recipient samples and

the sampling time points are defined according to the legend at the top. PCo; principal coordinate

(the two most informative coordinates are shown). Color legend corresponds with the distribution of

the different PRE-FMT and POST-FMT samples into the groups linked to the four donor samples

evaluated (see legend into the plot). B - A Genetic distance-based approach for evaluating

microbiota transfer between donor and recipient pairs. The microbial community structures of the

PRE-FMT and POST-FMT samples were compared with those of the donors through the

calculation of the Bray-Curtis dissimilarity index and are represented as boxplots. The Wilcoxon

signed-rank test for paired samples was used to compare differences in genetic distances. C –

Microbiota abundance shifts after FMT. Abundance data distribution for three species selected from

those described in Table 1 with meaningful increase after FMT is shown in boxplot fashion.

Figure 3. Single-nucleotide variation (SNV) analysis to detect species transfer and engraftment.

The species *Parabacteroides merdae* and *Faecalibacterium prausnitzii rrn*s were used to reveal

SNVs within each species and their abundance as a consequence of FMT (A and B panels,

respectively). Variation in one unit of clr-normalized counts is roughly equal to a doubling in raw

DNA read counts according to data distribution in both cases. C and D - The mapping, indexing,

and pileup of thousands of reads of *P. merdae* and *F. prausnitzii rrns* allowed us to identify

polymorphic sites exhibiting an even frequency for at least two nucleotides. These potential SNVs

are highlighted inside open circles, and the position number in the *rrn* sequence is also indicated.

The structural arrangement of *rrn* is illustrated on the x-axis. The black lines indicate the relative

frequency of the dominant allele, whereas the gray lines indicate the relative coverage per site

related to the average across the *rrn*. E and F - The distribution of the Bray-Curtis dissimilarity

index of the microbiota between POST-FMT and PRE-FMT samples and the corresponding donors,

using the combined nucleotide frequencies of SNVs detected in *P. merdae* and *F. prausnitzii*,

respectively. The Wilcoxon signed-rank test for paired samples was used to compare the genetic

distances of the microbiotas. G – Assessment of delta Bray-Curtis distances computed as POST-

FMT vs DONOR samples minus PRE-FMT vs DONOR samples for *Faecalibacterium prasunitzii*

and *Parabacteroides merdae* species. H - Electropherograms obtained from Sanger sequencing for

the strains of *P. merdae* SNV-3527, SNV-2293, and SNV-1254. The predominant alleles inferred

for every sample (based on the Q-score of base calling) supported the hypothesis that the strains

were transferred between donor and recipient pairs, as anticipated during the Nanopore-based

assessment.

Figure 4. Donors' influence on FMT success. A scatter plot showing the comparison of gut

microbiota between donor-recipient pairs at baseline (x-axis, Bray-Curtis dissimilarity index

computed) versus the shift in the microbiota community structure of every recipient before and after

the FMT (y-axis). The size of filled dots is determined by the sum of ranks linked to the

improvement of all metabolic parameters evaluated (blood pressure, fasting glucose and insulin,

glycosylated hemoglobin, HOMA-IR, weight loss, and BMI loss) (metabolic health score). Color

legend is associated with donor information. Asterisks and size represent the average values of all

recipients linked to every donor. Dashed grey line indicates linear correlation based on averaged

values (asterisks).

Figure S1. BLAST-based results showing the correct identification of assembled *rrn*s. The top hits

supporting the identification are shown for six different *rrn*s assembled with low, intermediate or

high coverage (number of reads).

Figure S2. Alpha diversity of the recipients' microbiota before and after FMT. The distributions of

the values obtained for four alpha diversity indicators, including Chao's index, Shannon's index, the

reciprocal Simpson's index, and the dominance index, are shown as boxplots. The results of the

Wilcoxon signed-rank test applied to establish differences between the two groups of samples

(POST-FMT and PRE-FMT) are also shown.

Figure S3. Searching for SNV genotypic variation in the NCBI nonredundant nucleotide database.

The nucleotide sequences spanning positions 1180 to 1280, 2250 to 2350, and 3470 to 3570 of the

P. merdae rrn operon were used as queries in searches against the nonredundant nucleotide

database on the BlastN web server (https://blast.ncbi.nlm.nih.gov/Blast.cgi). The results showed the

intrinsic variation attributable to SNV-3527 (as an example) present in genomic records from the

NCBI nonredundant nucleotide database including strains with genotype A3527 and others with

G3527, both of which were variants inferred from our Nanopore-based sequencing. Nucleotides

highlighted in yellow show the 5-mer surrounding the region of SNV-3527, revealed in detail by

Sanger sequencing, as shown in Figure 4H. The C2293 genotype was retrieved from a unique

matching sequence (CL06T03C08 strain), and A1254 was predominantly found in most of the *P.*

merdae strains evaluated (data not shown).

Figure S4. FastQC-based assessment of sequences produced by nanopore device. Results from the

main basecaller, *Albacore v2.1.3* and the more recently released high-performance basecaller of

nanopore data (*Guppy v5.0.11*) are shown for all three different sequencing runs generated in this

study.

A

PCo2 - 11.6% variability

PCo1 - 19.5% variability

Donor
- Donor1
- Donor2
- Donor4
- Donor6

☐ Recipients' PRE-FMT samples
■ Recipients' POST-FMT samples
● Donors' samples

B

p = 0.075

Bray-Curtis dissimilarity index

C

p = 0.002   *p* = 0.003   *p* = 0.033

clr-normalized counts

Parabacteroides merdae   *Sutterella massiliensis*   *Prevotella copri*

☐ **PRE-FMT**   ◼ **POST-FMT**