1 **Title: Principles of resilient coding for plant ecophysiologists**

Abstract

Plant ecophysiology is founded on a rich body of physical and chemical theory, but it is
challenging to connect theory with data in unambiguous, analytically rigorous, and
reproducible ways. Custom scripts written in computer programming languages (coding)
enable plant ecophysiologists to model plant processes and fit models to data reproducibly
using advanced statistical techniques. Since many ecophysiologists lack formal programming
education, we have yet to adopt a unified set of coding principles and standards that could
make coding easier to learn, use, and modify. We identify eight principles to help in plant
ecophysiologists without much programming experience to write resilient code: 1)
standardized nomenclature, 2) consistency in style, 3) increased modularity/extensibility for
easier editing and understanding, 4) code scalability for application to large datasets, 5)
documented contingencies for code maintenance, 6) documentation to facilitate user
understanding; 7) extensive tutorials, and 8) unit testing. We illustrate these principles using a
new R package, {photosynthesis}, which provides a set of analytical and simulation tools for
plant ecophysiology. Our goal with these principles is to advance scientific discovery in plant
ecophysiology by making it easier to use code for simulation and data analysis, reproduce
results, and rapidly incorporate new biological understanding and analytical tools.

Keywords: Gas exchange, hydraulics, photosynthesis, modeling, stomatal conductance,
curve fitting, software, R

Background

Computer coding is becoming an increasingly important skill in biological research (Sayres *et al.*, 2018), especially within plant ecophysiology. A disconnect in coding skill and a lack of formal computer science training can make it difficult for biologists to create or modify programs to incorporate new understanding of biological processes. In other words, sophisticated code (by trained programmers) is efficient, but difficult to modify by biologists for new uses. So why code at all? Coding allows for consistent, reproducible, transparent, and scalable analyses of scientific data, while at the same time minimizing human workhours compared to using pre-packaged software. However, most published ecophysiological analyses use spreadsheet-based methods rather than computer code, which comes with some limitations. For example, Sharkey *et al.* (2007) have an Excel spreadsheet-based method for fitting photosynthetic CO_2 response (A-C_i) curves (also see Bellasio *et al.*, 2016). A spreadsheet-based method can take several minutes per curve and involves a substantial amount of subjective decision-making (e.g. 'eye-balling' where transitions between CO_2- and RuBP- limited photosynthesis occur). Likewise, analysis of pressure-volume curves for hydraulic parameters is usually done via an Excel spreadsheet-based method (Sack *et al.*, 2003), which can be time-consuming, requires subjective decisions, and spreadsheets are usually not published with manuscripts, obscuring methodology. The total workload is time per spreadsheet multiplied by the number of curves, which can be inefficient in large studies. Cryptic changes in the spreadsheets can occur without a record of the change, potentially leading to compounding errors. Furthermore, spreadsheet tools often break, requiring a fresh, unaltered spreadsheet to be used for each CO_2 response curve. Another option, provided by Gu *et al.* (2010) (leafweb.org) provides an online service that analyses A-C_i curves, however in this case, the analysis is a black-box and could be misused by users lacking an understanding of the fitting process, and the data are stored on a government server which may cause some users discomfort.

Meanwhile, Duursma (2015) developed an R package, {plantecophys}, that can obtain similar outputs to the Sharkey et al. (2007) fitting tools in seconds, with far fewer subjective decisions that can easily be outlined in the code used in the fitting process, while providing a similar, but transparent approach as in Gu et al. (2010). Like the {plantecophys} package, analytical methods should be fully transparent and reproducible. As such, authors should publish their code, which is still not the norm in plant ecophysiology (but see Kumarathunge et al., 2019 for an example of published code). As a community, increased adoption and

dissemination of code will help the field perform more sophisticated analyses and model comparison (e.g. Walker *et al.* 2021). Coding may also streamline integration between theory and data analysis, especially for complex mathematical formulations that require computationally intensive numerical methods, a common situation in plant ecophysiology. Ideally, we would like a workflow in which we state our assumptions mathematically, derive empirical predictions, and test those predictions or estimate parameters with data. The process of translating a mathematical model of biology into code can also help novice and advanced coders deepen their understanding of models and their assumptions before confronting them with data. Open-source, research-grade computer algebra systems like SymPy (Meuer *et al.*, 2017) and numerical solvers aid mathematical derivation and are part of or can be readily integrated with programming languages that are widely used for data manipulation and analysis, such as R (R Core Team, 2021), Python (Python Software Foundation), or Julia (Bezanson *et al.* 2017).

Although coding can speed up large analyses, reduce errors, make analyses reproducible, and integrate theory with data, writing robust code that can be understood and reused by other scientists is not easy. First, one must learn one or more programming languages (e.g. R, Python, Matlab, Julia), which can involve steep learning curves. Second, even though coding one's own analysis can make it easier to catch errors associated with inappropriate use of black-box proprietary software, one must still understand the assumptions and limitations of statistical techniques and conceptual tools. Finally, code can be as unique as someone's handwriting, which can make it difficult even for an experienced programmer to make sense of a "transparent" analysis unless there is sufficient annotation within the code.

In this perspective, we propose seven principles of coding tailored to the specific needs of the plant ecophysiology research community. For example, guidance in other scientific fields often emphasize computational speed. However, given the typical scale of ecophysiological datasets (~MB, i.e. small-batch, artisanal datasets) and the computer power of personal computers (~GB of RAM, ~GHz of processing power), computational speed is usually not a major limitation. Instead, ecophysiologists often need to estimate parameters derived from complex biophysical/chemical models. Coding is important as the complex models required to fit different response curves involve many interacting equations, numerical solvers, and parameters that either need to be set or estimated. For example, there are seven different

models that can be used to fit temperature responses which ultimately require different
equations and fixed parameters (Arrhenius, 1915; Johnson *et al.* 1942; Medlyn *et al.*, 2002;
Kruse *et al.*, 2006; Heskel *et al.*, 2016; Liang *et al.*, 2018). In this domain, code flexibility and
modularity are usually more important than computational speed. Furthermore, flexibility and
modularity in code would enhance the sustainability of software after publication, which can
be an issue (Prlić & Procter, 2012). Here we demonstrate coding principles designed for plant
ecophysiology using a new R package called {photosynthesis}. We caution that this software
is a work-in-progress that does not yet completely adhere to all of the coding principles to
which we aspire, but will be refined in future releases. This perspective, written by trained
biologists not programmers, is intended to convey some of the lessons we have learned so
far to provide guidance for plant ecophysiologists who are thinking about or starting to code
their workflows, especially using R. We recognize that many other scientists in this field are
adept coders who have already honed their practices through experience. Hence, this
perspective is intended to guide for less experienced coders rather than a mandate for the
entire field. We hope our perspective spurs experienced coders to share "best practices" with
less-experienced peers and expand the principles below to other languages besides R. As
computational plant ecophysiology matures, we hope that this perspective will help move the
field toward more standardized and sustainable software practices like those in more
computationally-intensive subfields of biology like population genetics (Adrion *et al.* 2020).

Description
Principles of Coding
The overarching concept we propose is making code resilient by making it easier to use,
reproduce, and modify. Obviously not every possible discovery and need within a scientific
field can be predicted, but the code can be written to allow easy modification and
accommodation of the source code as the science progresses. Functional programming in R
and other languages provides a powerful tool for writing functions that take functions as
arguments and easily process newly written code into a standardized output without the need
for ever modifying the original function itself (Wickham, 2019). Such an approach helps to
write modular code that is easy to modify and understand, while minimizing
interdependencies between functions.

Freely available resources already exist for good coding practices in R packages and can be
applied to R scripts as well, primarily from the efforts of Hadley Wickham (Wickham, 2014,
2015, 2016, 2017, 2019; Wickham & Grolemund, 2016). As well, guides to best practices for
scientific computing exist (see Wilson et al. 2014 for a list of best practices). Here we propose
principles of coding for plant ecophysiology that, if implemented, could circumvent some of
the common coding issues encountered when modifying the code of others, reduce the
learning difficulty for nascent coders, and make software maintenance much easier:

1.   Standardized nomenclature for variables and functions
2.   Consistent style
3.   Modularity and extensibility
4.   Scalability
5.   Documented contingencies
6.   Documentation
7.   Extensive tutorials
8.   Unit testing

We think that adopting some or all of these principles will improve code reproducibility and
help advance scientific discovery, but our goal is not to rigidly prescribe how plant
ecophysiologists should do their work. First, we recognize that others will have different, well-
reasoned preferences and/or apply principles we have not covered here. Second, those who
find these principles useful may find implementing all of them time-consuming at first. We
strongly encourage incremental progress and not making perfection the enemy of the good.
Indeed, the {photosynthesis} package described below only partially implements our
principles, with much left to do in future development.

Principle 1: standardized nomenclature
Names vary wildly between functions with published code and data and even amongst
instruments within the same company (e.g. for net CO_2 assimilation, "*A*" is used in the Li-Cor
6800 and "PHOTO" is used in the Li-Cor 6400). Ideally, we need both standardized
nomenclature in the field (e.g. Reid *et al.* 2005) and standardized construction of variable and
function names to enhance readability and reduce the burden for learning how to use new
packages and functions or testing published code. For example, g is always in reference to
conductance, where a subscript term would then describe the physical pathway (e.g. s for

stomata, c for cuticle, or m for mesophyll) as well as the gas (e.g. c for CO_2, w for water
vapor). For example, g_{sw} would mean stomatal conductance to water vapor. Standardizing
nomenclature across both mathematical models and data files can also streamline theory-
data integration, but this also requires standard translation between mathematical and
computer notation, which is beyond our scope here.

For example, in {photosynthesis}, every function is named in a descriptive manner: e.g.
`fit_t_response` fits specified temperature responses model to data, while `fit_gs_model`
fits specified models of stomatal conductance. Variable names are also standardized: e.g.
"T_leaf" always means leaf temperature in Kelvin (K), "A_net" always means net CO_2
assimilation in µmol m^{-2} s^{-1}. In this regard, standard units should also be imposed in the
analysis (e.g. in R via the {units} package (Pebesma et al., 2016)), to remove any ambiguities
when interpreting the output. To allow for differences in variable names from the raw data
(e.g. from using different machines), the "varnames" list is used to translate input names (note
that this convention is adopted from {plantecophys} (Duursma, 2015)). We propose adopting
Wickham's (Wickham, 2019) style in that functions that *do* something have a verb name, e.g.
`fit_aci_response`, while functions that act as objects within other functions (e.g. stomatal
conductance models) should have a noun name, e.g. `gs_model`.

Principle 2: consistent style
Consistent coding style makes reading code easier - certain conventions, e.g. commenting
what the *next* line of code does, can make it easier to understand code documentation. Our
preference is for the 'tidy style', which applies to both data and code structure, and much else
(see the *The tidyverse Style Guide:* https://style.tidyverse.org/). For data, tidy style advocates
that each column is a variable, and each row is an observation, since R is particularly suited
for this style of data structure. Popular R packages like {dplyr} (Wickham *et al.*, 2020) and
{tidyr} (Wickham and Henry, 2020) facilitate tidy data and many other packages, like
{photosynthesis}, use them for consistent style. For code, computers do not care about style,
as long as it is correctly formatted, but for humans reading code, adherence to well-designed
style can be helpful, especially for beginners trying to learn from others. A benefit of tidy style
in particular is that R packages {styler} (Müller and Walthert, 2020), {lintr} (Hester *et al.* 2020),
and {formatR} (Xie, 2019) can automate conformity to style. Ideally, a consistent style would
be adopted across the field, however this may be too rigid. Style can be highly personal, and

many experienced coders likely have developed their own style, formal or informal, that works
for them. Our proposal is geared for beginning coders who are looking for guidance on an
established and easy-to-implement style. At the very least, a consistent style *within* a project
will make it easier to read, understand, and modify the code.

Principle 3: modularity and extensibility
Arguably, code written for plant ecophysiologists, whether formally trained in coding or not,
should be written in a modular manner, much like Lego bricks, where one component (e.g.
Arrhenius function) can be easily swapped with another (e.g. peaked Arrhenius function), or
extended (e.g. hypothetical mechanistic temperature response model). Note that this may
increase apparent complexity of software packages by creating more functions and make it
more difficult to work with at first. However, it will make adding, subtracting, or modifying code
modules easier for researchers who need to make on-the-fly changes to code as new
biological processes are discovered or old ones re-evaluated. To achieve modularity in the
structure of photosynthesis, we used principles of functional programming to develop a set of
key functions for processing data and running quality control checks: `fit_many`,
`analyze_sensitivity`, `compile_data`, and `print_graphs`. Both `fit_many` and
`analyze_sensitivity` can be run with any function within and outside of {photosynthesis}
to run multiple curve fits or sensitivity analyses on assumed input parameters. Meanwhile,
`compile_data` is used for processing the list outputs from `fit_many` into a form usable for
further analyses and export from R, and `print_graphs` is used to export all graphs from a
list as either .jpeg or compiled as a .pdf.

For curve fitting functions with multiple models (e.g. temperature responses, g_s models), we
use a basic function (e.g. `fit_t_response`), which contains fitting procedures for each of
the seven temperature response models in the package. Meanwhile, a `t_functions` file
contains all the temperature response functions. To extend the capabilities and add in a new
temperature response model, we simply need to add the new model to `t_functions`, and
the fitting procedure to `fit_t_response`. Currently, adding new functions requires
modifying the source code, but future versions should increase extensibility by allowing users
to supply any temperature-response function. This principle of function building increases the
extensibility of the code, while consistent style and standardized nomenclature provide the
rules for writing the extended components.

Modularity also applies to modelling. The {photosynthesis} functions `photo` and `photosynthesis` model C_3 photosynthesis using the Faquhar-von Caemmerer-Berry biochemical model (Farquhar *et al.* 1980). To account for temperature dependence, a user can specify leaf temperature, or they can provide additional inputs (e.g. air temperature, leaf size, wind speed, etc.) to model leaf temperature using energy balance in the R package {tealeaves} (Muir 2019). Both {photosynthesis} and {tealeaves} packages are modular in that they can work independently or be readily integrated (**Methods S1**). Ideally, future modeling packages would add modules to model environmental and plant parameters either on their own or integrated with these tools.

Principle 4: scalability

A major advantage in using code to analyze data is the ability to scale up an analysis to reduce time spent on repetitive tasks common in spreadsheet-based methods such as copy-and-paste, selecting data, choosing menu options, etc. Functions allow the same model to be fit across groups within a dataset using a consistent method. For this, our `fit_many` function and the principles of functional programming are how we achieve scalability within the package. Rather than writing functions for each type of model or curve, we have a single multiple fitting function, sensitivity analysis function, and printing function. R even has generic functions for scaling such as `apply` (base R language) and `map` ({purrr} package [Henry & Wickham, 2020]) which can be easily parallelized for speed (e.g. {parallel} and {furrr} [Vaughn & Dancho, 2018] packages). This makes it easy to scale a new function within the software to a large dataset.

Principle 5: documented contingencies

By documenting which functions are dependent on one another, it becomes easier to troubleshoot issues when modifying code and to pre-empt issues when adding or replacing a component. For example, `fit_aq_response` depends on `aq_response` - if we want to change from the non-rectangular hyperbola model to a rectangular hyperbolic model, then `fit_aq_response` needs to be modified in addition to `aq_response`. To document contingencies, we created a function, `check_dependencies`, which uses {pkgnet} (Burns et al., 2020) to generate an html report that automatically documents R package inter-dependencies and function inter-dependencies. This is particularly useful when adding,

subtracting, or modifying functions in the package, as it allows planning to minimize issues
that could break code.

Principle 6: documentation
Code annotations allow a new user to readily understand what a line of code is doing, how it
is doing it, and why the code is written in a particular way. By providing exhaustive line-by-line
annotation of a function, a new user can more rapidly understand the blueprint of the function.
This is especially useful for R scripts and code hosted on GitHub (unfortunately, comments
are erased from code upon submission to CRAN). For example, in `fit_t_response`, we
outline the need for running looped iterations for the starting values of nonlinear least squares
curve-fitting (Fig. 1). In the case of R packages hosted on CRAN, R documentation files
provide information on how to use a function, though as a terser set of instructions as per
CRAN policies (https://cran.r-project.org/doc/manuals/r-devel/R-exts.html).

```r
#Basically, use Arrhenius curve to feed Ea into Topt function start
#Try approach where you start Hd from 1 to 1000 to ensure model fit
#select minimum residual
model <- nlsLM(data = data,
               Par ~ Par25 * t_response_arrhenius(Ea,
                                                  Tleaf = Tleaf),
               start = start,
               lower = c(0, 0),
               upper = c(1e10, 10 * max(data$Par)),
               control = nls.control(maxiter = 100)
)

#Create empty dataframe to fill with 1000 curve fits
model_fm <- as.data.frame(cbind(rep(0, 1000),
                                rep(0, 1000),
                                rep(0, 1000),
                                rep(0, 1000),
                                rep(0, 1000),
                                rep(varnames$Par[[1]], 1000)))
#Assign column names
colnames(model_fm) <- c("Ea", "Hd", "kopt", "Topt", "residual",
                        "Parameter")
```

`#Make sure variable classes are appropriate`

Figure 1. Example of coding annotations to explain the given analytical approach.

Enough metadata and commenting should be provided for a new user to understand how to
use the written code (which can be an issue that affects widespread use of a program,
Mangul *et al.* 2019).

Principle 7: extensive tutorials

As with any tool, software will only be used if potential users can understand how it works.
Extensive tutorials, while providing function-by-function examples of how to use the software,
should also incorporate basic data-wrangling examples and explanations of why a given
approach to data analysis is used in the field. The benefits of this approach include: making
the code easier to adopt into your own analysis, making it easier for new coders to learn
enough of the language to use the package effectively, and help trainees learn the
appropriate theory behind the measurements and analytical approach. The net effect should
be to increase the inclusivity of the field by reducing barriers to success since not all
individuals will have equal access to workshops or experienced colleagues.

Principle 8: unit testing and benchmarking

For reproducibility, code should yield the same results when it is run by other users months or
years into the future. Unit testing, a common practice in software development that is still rare
in scientific code, evaluates whether various components, such as custom functions, perform
as expected. If all the components work as expected, it provides confidence that the whole
body of code does what it is supposed to. Most scientists informally test their functions as
they develop them, but formal unit testing involves writing scripts to test code and can be
rerun to periodically check whether code still works as expected. More dedicated efforts
automate testing and quantify code coverage, the fraction of code that is evaluated during
automated tests. There are many ways to implement unit testing, but the {testthat} package is
one option for R packages (Wickham, 2011) that {photosynthesis} uses for some (but not yet
all) of its source code. A related concept is benchmarking, by which we mean comparing
parameter estimates from the same data set using different software or later versions of the
same software. Benchmarking can help determine if parameter estimation is consistent
between software packages. For example, parameter estimates of photosynthetic CO_2
response parameters (Farquhar *et al.*, 1980) are very similar using comparable settings in
{photosynthesis} and {plantecophys} (Notes S1).

Examples of resilient coding in the {photosynthesis} package for R

We built a package containing analytical tools for plant ecophysiology (Stinziano *et al.*, 2020), embedding our coding principles into the package itself. The R package contains functions for fitting photosynthetic CO_2 (Farquhar *et al.*, 1980; von Caemmerer, 2000; Gu *et al.*, 2010; Duursma, 2015) and light response curves (Marshall & Biscoe, 1980), temperature responses of biological processes (Arrhenius, 1915; Medlyn *et al.*, 2002; Kruse *et al.*, 2006; Heskel *et al.*, 2016; Liang *et al.*, 2018), light respiration (Kok, 1956; Laisk, 1977; Yin et al., 2009, 2011; Walker & Ort, 2016), mesophyll conductance (Harley *et al.*, 1992), stomatal conductance models (Ball *et al.*, 1987; Leuning, 1995; Medlyn *et al.*, 2011), pressure-volume curves (Tyree & Hammel, 1972; Koide *et al.*, 2000; Sack *et al.*, 2003), hydraulic vulnerability curves (Pammenter & van der Willigen, 1998; Ogle *et al.*, 2009), and sensitivity analyses (Table 1; Table S2). It also contains functions for modeling C_3 photosynthesis using the Farquhar-von Caemmerer-Berry biochemical model (Farquhar *et al.*, 1980). The default kinetic parameters for gas exchange fitting procedures are taken from *Nicotiana tabacum* (Bernacchi *et al.*, 2001, 2002). The {photosynthesis} package is currently limited to C_3 photosynthesis, but future releases should expand its functionality to other photosynthetic pathways. A comprehensive illustration of how to use the package can be found in the vignette of the package (**Notes S2**, "photosynthesis-curve-fitting-sensitivity-analyses.rmd"). There are currently two vignettes available for the package that function as tutorials on CRAN (https://CRAN.R-project.org/package=photosynthesis). The first vignette (titled "photosynthesis-curve-fitting-sensitivity-analyses") demonstrates how to use curve fitting and sensitivity tools and the second (titled "introduction to the photosynthesis package") demonstrates how to simulate photosynthetic rate using the Farquhar-von Caemmerer-Berry C_3 biochemical model, define leaf and environmental parameters, replace default parameters, and solve for chloroplastic CO_2 concentrations.

Table 1. List of {photosynthesis} functions with applications and descriptions. The documentation for each function describes the estimated or simulated parameters, constants, and other calculated values. Documentation is updated to describe new functionalities as they are added.

Base Functions		
Applications	Function	Description
Gas Exchange	`fit_aci_response`	Fits A-C_i curves, provides parameters/graphs
Gas Exchange	`fit_aq_response`	Fits A-Q curves, provides parameters/graphs
Gas Exchange	`fit_g_mc_variableJ`	Fits g_{mc}, adds g_{mc} and dCcdA to dataframe for reliability checking
Gas Exchange	`fit_gs_model`	Fits the Ball *et al.* 1987, Leuning 1995, and Medlyn *et al.* 2011 models of stomatal conductance, provides parameters/graphs
Hydraulics	`fit_hydra_vuln_curve`	Fits the sigmoidal and Weibull models to hydraulic vulnerability data, provides parameters/graphs
Hydraulics	`fit_PV_curve`	Fits pressure volume curves, provides parameters/graphs
Gas Exchange	`fit_r_light`	Fits r_light according to the Kok (1956) method, Yin method (Yin *et al.* 2009, 2011), or Walker & Ort (2015) method.
Gas Exchange, Biochemistry	`fit_t_response`	Fits an Arrhenius (Arrhenius 1915), Heskel (Heskel *et al.* 2016), Kruse (Kruse *et al.* 2006), Medlyn (Medlyn *et al.* 2002), MMRT (Hobbs *et al.* 2013), and quadratic temperature response models, provides parameters/graphs
Modeling	`photo`	Simulates C_3 photosynthesis over a parameter set
Modeling	`make_parameters`	A set of functions (e.g. `make_enviropar`, `make_leafpar`) that generates the required inputs for `photo`
Meta-functions & Utilities		
Application	Function	Description
Software modification	`check_dependencies`	Generates HTML with package and function dependencies
All components	`compile_data`	Compiles the output from the `fit_many` function
All components	`fit_many`	Fits a function many times through a grouping variable
All components	`print_graphs`	Prints graphs from a list of graphs
All components	`sensitivity_analysis`	Allows up to 2-factor sensitivity analysis of any function

The package is specifically designed to accommodate new analytical tools and discoveries and be easily maintained by new users. Nonlinear curve fitting procedures use the nlsLM

function from {minpack.lm} (Elzhov *et al.*, 2016), which provides a more robust fitting

procedure for non-linear functions than the base R `nls` function. Graphical outputs are

provided using {ggplot2} (Wickham, 2016). Meta-functions were constructed with the tools

provided for generalizing functions and arguments in {rlang} (Henry & Wickham, 2019).

The principles of modularity and functional programming have been used to substantially

reduce code interdependencies within the software. For example, the `fitaci` function from

{plantecophys} has over 30 function dependencies (Fig. 2a). By applying our principles, we

were able to reduce this to just 4 function dependencies (Fig. 2b), by reengineering the fitting

procedure and eliminating redundant functions and code. Arguably, fewer dependencies

could indicate less modularity, even though each of the components is modular, but fewer

dependencies may reduce the number of bugs introduced by revisions in other components.

Figure 2. Dependencies of the *A-C$_i$* fitting functions in (a) {plantecophys} and (b)

{photosynthesis}.

Example Dataset

To demonstrate the fitting functions of the package, we use a combination of data collected

for the package and previously published data. A CO_2 by light response curve and CO_2 by

temperature response curve were collected in sunflower (*Helianthus annuum*) grown in a

rooftop greenhouse at the University of New Mexico (35.0843°N, 106.6198°W, 1587 m a.s.l.,

18.3 to 21.1/15.6 to 21.1 °C day/night temperature with daily irradiances of 600 to 1,200 µmol

m^{-2} s^{-1}). CO_2 response curves were measured at irradiances of 1,500, 150, 375, 125, 100, 75,

50, and 25 µmol m^{-2} s^{-1} at a T_{leaf} of 25 °C. CO_2 response curves were also measured at T_{leaf}

of 17.5, 20, 22.5, 25, 27.5, 30, 32.5, 35, 37.5, and 40 °C at an irradiance of 1,500 µmol m^{-2} s^{-}

^{1}. Data to demonstrate hydraulic vulnerability curve fitting methods were drawn from (Hudson

et al. 2018), while data for leaf pressure/volume analysis come from an unpublished dataset

collected at the University of New Mexico. Below we illustrate some of the functionality of the

package. This data is freely available in the package so potential users can test out the

functions and different analyses in the code. We refer potential users to the package vignette

for more worked examples (**Notes S2**, "photosynthesis-curve-fitting-sensitivity-

analyses.rmd").

Photosynthetic light response curve fitting

The `fit_aq_response` function returns a list containing the fitted light response model,

model parameters, and a graph showing the model fit to the data (Fig. 3a). This function

estimates the light-saturated net CO_2 assimilation rate, quantum yield of CO_2 assimilation, an

empirical an empirical curvature factor, and respiration (Marshall & Biscoe, 1980).

Figure 3. Gas exchange curve fitting outputs. a) Output from {`fit_aq_response`
showing the data (black points), the model fit (blue line), and the standard error on the
model fit (grey region). The light response at a [CO₂] of 100 µmol mol⁻¹ is shown. *A_net*:
net CO₂ assimilation. b) Graph from `fit_aci_response` showing modelled *A_net* (*A_mod*,
black line), CO₂-limited *A_net* (*A_c*, blue), RuBP regeneration-limited *A_net* (*A_j*, orange),
triose phosphate utilization-limited *A_net* (*A_p*), and the data (white dots). *A_net*: net CO₂
assimilation; C_i: intercellular CO₂ concentration. c) Output from `fit_t_response`
showing the Heskel temperature response of *J_max*. Data are black dots, model fit is the

 blue line, and the grey shaded region is the standard error on the model fit. J_{max}:
 maximum rate of electron transport; T_{leaf}: leaf temperature.

Photosynthetic CO_2 response curve-fitting

The `fit_aci_response` function returns a list containing the fitted parameters, a data

frame with the modelled data output, and a graph showing the model fit to the data (Fig. 3b).

It estimates the standard parameters of the Farquhar-von Caemmerer-Berry C_3 biochemical

model (Farquhar *et al.* 1980) and parameter standard errors to help evaluate results. As with

any nonlinear regression, failure of the solver to converge on a solution or very large standard

errors usually indicate problems fitting the model to the data and unreliable parameter

estimates.

Photosynthetic temperature response curve fitting

A series of temperature response functions can be fit using the package, with the outputs

including the fitted model, model parameters, and a graph (Fig. 3c). As with other functions,

details about parameters given in the package documentation.

Fitting g_m using the variable J method

The `fit_g_mc_variableJ` function using the method of Harley *et al*. (1992) using

chlorophyll fluorescence and gas exchange data to estimate g_{mc}. Both g_{mc} and $\delta C_c/\delta A$ are

calculated, where $\delta C_c/\delta A$ between 10 and 50 are deemed to be "reliable" (Harley *et al*.,

1992), and an average g_{mc} value is estimated based on the reliable values. This makes it

relatively easy to assess the reliability of g_{mc} estimates (Fig. 4).

Figure 4. Relationship between g_{mc} estimated through the variable J method and $\delta C_c/\delta A$ to test for reliability. The `fit_g_mc_variableJ` function was used on the CO_2 by light response data in sunflower. g_m: mesophyll conductance; C_c: chloroplastic CO_2 concentration; A: net CO_2 assimilation

Hydraulic vulnerability curve fitting

The `fit_hydra_vuln_curve` fits hydraulic vulnerability data using both a sigmoidal and Weibull function. Outputs include model fits, parameters and a graph (Fig. 5a).

Figure 5. a) Example output from `fit_hydra_vuln_curve` showing both model fits overlaid on the data (black dots). *PLC*: **percent loss of conductivity;** *Pe*: **air entry point;** *P50*: **water potential at 50%** *PLC*; *Pmax*: **hydraulic failure threshold. b, c) Example output from fit_pv_curve showing the b) water mass graph and c) the pressure-volume curve. Grey lines are fit to the linear regions of the data. Ψ: water potential; RWC: relative water content.**

Pressure-volume curves

The fit_pv_curve fits pressure-volume curves, returning parameters such as relative water
content and water potential at turgor loss points, relative capacitance at full turgor, and others.
Outputs include parameters and graphs (Fig. 5bc).

Sensitivity analyses

Both `analyze_sensitivity` and `compute_sensitivity` are used in combination for
sensitivity analyses. `analyze_sensitivity` allows up to two assumed parameters to be
varied in a fitting function, while `compute_sensitivity` runs two types of local sensitivity
calculations based on a user-defined reference value: parameter effect (Bauerle et al., 2014)
and control coefficient (Capaldo & Pandis, 1997). We can look at the impact of varying g_m and
Γ^* at 25 °C on fitted V_{cmax} (Fig. 6). We can see that g_m and Γ^* at 25 °C have an orthogonal
impact on V_{cmax}, with Γ^* having a stronger control than g_m on V_{cmax}.

Figure 6. Control coefficients of g_m and Γ^* at 25 °C calculated from
`analyze_sensitivity` and `compute_sensitivity`.

Moving forward - standardized practices and code editors

It is not easy to rewrite software, and we are not arguing as such. Rather, going forward as a community, we argue that we should adopt a set of coding principles and guidelines to create code as flexible as the biology we study. We present the R package, {photosynthesis}, as an example of these principles and guidelines. The consequences of this are not to be understated: it will be easier for new trainees and beginner coders to learn, understand, and write code for the community; and it will be easier to tailor existing code to our projects.

The drawback is that code may run more slowly, which may be a worthwhile tradeoff for some but not others. For example, computational speed may take precedence over flexibility for eddy flux covariance, genomics, and other "big data" applications. In ecophysiology, many datasets are often small enough that even complex analyses may only take 1 hr on one computer core of a multi-core system – as a community we can often afford slower-running code for greater flexibility and ease-of-understanding, especially as this could save days or weeks of coding to write a desired analysis. Our code should be as flexible as, and easier to understand, than the biology it describes.

However, providing code according to these standards is not sufficient - we also need code-competent editorial staff for journals who can properly review and test submitted code to ensure that it runs as intended. In some cases, code for a published dataset does not work even after comprehensive modification (Stinziano, pers. comm.). Standardized coding practices will help to reduce the burden on code editors by making it easier to read and understand code submissions.

Data Availability Statement

All data and code used in the manuscript are available at

https://github.com/cdmuir/photosynthesis.

Supporting Information

Methods S1 – Description of variables used in {photosynthesis}

Tables S2 – Table of other utility functions in {photosynthesis}.

Notes S1 – Benchmark comparison of `plantecophys::fitaci` and `photosynthesis::fit_aci_response`

Notes S2 – The {photosynthesis} R package tar.gz file.

Notes S3 – Examples tidy data file (hydraulic_vulnerability.csv).

Acknowledgements

We thank Tom Buckley, Markus Hauck, and three anonymous reviewers for feedback on earlier versions of the manuscript. This is publication #145 from the School of Life Sciences, University of Hawai'i at Mānoa.

References

Arrhenius S. **1915.** Quantitative laws in biological chemistry. Bell.

Baker M. 2017. Scientific computing: code alert. Nature **541**:563-565.

Ball JT, Woodrow IE, Berry JA. 1987. A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions, in Progress in Photosynthesis Research, Proceedings of the VII International Congress on Photosynthesis, vol. 4, edited by **I. Biggins**, pp. 221–224, Martinus Nijhoff, Dordrecht, Netherlands.

Bauerle WL, Daniels AB, Barnard DM. 2014. Carbon and water flux responses to physiology by environment interactions: a sensitivity analysis of variation in climate on photosynthetic and stomatal parameters. *Climate Dynamics* **42**:2539-2554.

Bellasio C, Beerling J, Griffiths H. 2016. An Excel tool for deriving key photosynthetic parameters from combined gas exchange and chlorophyll fluorescence: theory and practice. *Plant, Cell & Environment* **39**:1180-1197.

Bernacchi CJ, Singsaas EL, Pimentel C, Portis AR, Long SP. 2001. Improved temperature response functions for models of rubisco-limited photosynthesis. *Plant Cell & Environment* **24**:253-259.

Bernacchi CJ, Portis AR, Nakano H, von Caemmerer S, Long SP. 2002. Temperature response of mesophyll conductance. Implications for the determination of rubisco enzyme kinetics and for limitations to photosynthesis in vivo. *Plant Physiology* **130**:1992-1998.

Burns B, Lamb J, Qi J. 2020. pkgnet: Get Network Representation of an R Package. R package version 0.4.1. https://CRAN.R-project.org/package=pkgnet

Capaldo KP, Pandis SN. 1997. Dimethylsulfide chemistry in the remote marine atmosphere: evaluation and sensitivity analysis of available mechanisms. *Journal of Geophysical Research* **102**:23251-23267

Carey MA, Papin JA. 2018. Ten simple rules for biologists learning to program. *PLoS Computational Biology* **14**:e1005871.

Duursma RA. 2015. Plantecophys – an R package for analyzing and modeling leaf gas exchange data. *PLoS ONE* **10**:e0143346.

Elzhov TV, Mull KM, Spiess A-N, Bolker B. 2016. minpack.lm: R interface to the Levenberg-Marquardt nonlinear least-squares algorithm found in MINPACK, plus support for bounds. R package version 1.2-1. https://CRAN.R-project.org/package=minpack.lm

Farquhar GD, von Caemmerer S, Berry JA. 1980. A biochemical model of photosynthetic CO2 assimilation in leaves of C3 species. *Planta* **149**: 78-90.

Gu L, Pallardy SG, Tu K, Law BE, Wullschleger SD. 2010. Reliable estimation of biochemical parameters from C3 leaf photosynthesis-intercellular carbon dioxide response curves. *Plant Cell & Environment* **33**:1852-1874.

Harley PC, Loreto F, Di Marco G, Sharkey TD. 1992. Theoretical considerations when estimating mesophyll conductance to CO2 flux by analysis of the response of photosynthesis to CO2. *Plant Physiology* **98**:1429 - 1436.

Henry L, Wickham H. 2019. Rlang: functions for base types and core R and 'Tidyverse' features. R package version 0.4.2. https://CRAN.R-project.org/package=rlang.

Henry L, Wickham H. 2020. purrr: Functional Programming Tools. R package version 0.3.4. https://CRAN.R-project.org/package=purrr

Heskel MA, O'Sullivan OS, Reich PB, Tjoelker MG, Weerasinghe LK, Penillard A, Egerton JJG, Creek D, Bloomfield KJ, Xiang J, Sinca F, Stangl ZR, la Torre AM, Griffin KL, Huntingford C, Hurry V, Meir P, Turnbull MH, Atkin OK. 2016. Convergence in the temperature response of leaf respiration across biomes and plant functional types. *Proceedings of the National Academy of Sciences USA* **113**:3832-3837

Hester J, Angly F, Hyde R. 2020. lintr: A 'Linter' for R Code. R package version 2.0.1. https://CRAN.R-project.org/package=lintr

Hudson PJ, Limousin JM, Krofcheck DJ, Boutz AL, Pangle RE, Gehres N, McDowell NG, Pockman WT. 2018. Impacts of long-term precipitation manipulation on hydraulic architecture and xylem anatomy of piñon and juniper in Southwest USA. *Plant Cell & Environment* **41**: 421-435.

Johnson F, Eyring H, Williams R. 1942. The nature of enzyme inhibitions in bacterial luminescence: sulphanilamide, urethane, temperature, pressure. *Journal of Cell Comparative Physiology* **20**, 247– 268.

Koide RT, Robichaux RH, Morse SR, Smith CM. 2000. Plant water status, hydraulic resistance and capacitance. In: Plant Physiological Ecology: Field Methods and Instrumentation (eds RW Pearcy, JR Ehleringer, HA Mooney, PW Rundel), pp. 161-183. Kluwer, Dordrecht, the Netherlands

Kok B. 1956. On the inhibition of photosynthesis by intense light. *Biochimica et Biophysica Acta* **21**: 234–244

Laisk A. 1977. Kinetics of photosynthesis and photorespiration in C3 plants. Nauka, Moscow.

Leuning R. 1995. A critical appraisal of a coupled stomatal- photosynthesis model for C3 plants. *Plant Cell & Environment* **18**:339-357

Liang LL, Arcus VL, Heskel MA, O'Sullivan OS, Weerasinghe LK, Creek D, Egerton JJG,
Tjoelker MG, Atkin OK, Schipper LA. 2018. Macromolecular rate theory (MMRT) provides a
thermodynamics rationale to underpin the convergent temperature response in plant leaf
respiration. *Global Change Biology* **24**:1538-1547

Mangul S, Martin LS, Eskin E, Blekhman R. 2019. Improving the usability and archival
stability of bioinformatics software. *Genome Biology* **20**:47.

Marshall B, Biscoe P. 1980. A model for C3 leaves describing the dependence of net
photosynthesis on irradiance. *Journal of Experimental Botany* **31**:29-39

Medlyn BE, Dreyer E, Ellsworth D, Forstreuter M, Harley PC, Kirschbaum MUF, Le Roux
X, Montpied P, Strassemeyer J, Walcroft A, Wang K, Loustau D. 2002. Temperature
response of parameters of a biochemically based model of photosynthesis. II. A review of
experimental data. *Plant Cell & Environment* **25**:1167-1179

Medlyn BE, Duursma RA, Eamus D, Ellsworth DS, Prentice IC, Barton CVM, Crous KY,
Angelis PD, Freeman M, Wingate L. 2011. Reconciling the optimal and empirical
approaches to modelling stomatal conductance. *Global Change Biology* **17**:2134-2144

Muir CD. 2019. tealeaves: an R package for modelling leaf temperature using energy
budgets. *AoB PLANTS* **11**:plz054

Müller K, Walthert L. 2020. styler: Non-Invasive Pretty Printing of R Code. R package
version 1.3.2. https://CRAN.R-project.org/package=styler

Ogle K, Barber JJ, Willson C, Thompson B. 2009. Hierarchical statistical modeling of xylem
vulnerability to cavitation. *New Phytologist* **182**:541-554

Pammenter NW, Van der Willigen CV. 1998. A mathematical and statistical analysis of the
curves illustrating vulnerability of xylem to cavitation. *Tree Physiology* **18**:589-593

Pebesma R, Mailund T, Hiebert J. 2016. Measurement units in R. R Journal 8:486-494.

Prlić A, Proctor PB. 2012. Ten simple rules for the open development of scientific software.
PLoS Computational Biology **8**:e1002802.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Reid DEB, Silins U, Mendoza C, Lieffers VJ. 2005. A unified nomenclature for quantification
and description of water conducting properties of sapwood xylem based on Darcy's law. *Tree*
Physiology 25:993-1000.

Sack L, Cowan PD, Jaikumar N, Holbrook NM. 2003. The 'hydrology' of leaves: co-
ordination of structure and function in temperate woody species. *Plant, Cell & Environment*
26: 1343-1356.

Sayres MAW, Hauser C, Sierk M, Robic S, Rosenwald AG, Smith TM, Triplett EW, et al.
2018. Bioinformatics core competencies for undergraduate life sciences education. *PLoS*
ONE **13**:e0196878.

Tyree MT, Hammel HT. 1972. Measurement of turgor pressure and water relations of plants
by pressure bomb technique. *Journal of Experimental Botany* **23**:267

Vaughan D, Dancho M. 2018. furrr: Apply Mapping Functions in Parallel using Futures. R
package version 0.1.0. https://CRAN.R-project.org/package=furrr

von Caemmerer S. 2000. Biochemical models of leaf photosynthesis. CSIRO Publishing,
Collingwood.

Walker AP, Johnson AL, Rogers A, Anderson J, Bridges RA, Fisher RA, Lu D, Ricciuto
DM, Serbin SP, Ye M. 2021. Multi-hypothesis comparison of Farquhar and Collatz
photosynthesis models reveals the unexpected influence of empirical assumptions at leaf and
global scales. *Global Change Biology* **27:**804-822.

Walker BJ, Ort DR. 2015. Improved method for measuring the apparent CO_2
photocompensation point resolves the impact of multiple internal conductances to CO_2 to net
gas exchange. *Plant Cell & Environment* **38**:2462- 2474

Wickham H. 2019. Advanced R, 2nd Ed. Chapman & Hall.

Wickham H. 2017. Tidyverse Tidyweb. http://tidyverse.org/.

Wickham H. 2016. Tidyverse: easily install and load 'tidyverse' packages. https://CRAN.R-
project.org/package=tidyverse.

Wickham H. 2016. ggplot2: elegant graphics for data analysis. Springer-Verlag New York.

Wickham H. 2015. R Packages: Organize, Test, Document, and Share Your Code. O'Reilly
Media Inc. Sebastopol, CA, USA.

Wickham H. 2014. Tidy data. *Journal of Statistical Software* **59**:1-23.

Wickham H. 2011. testthat: Get Started with Testing. *The R Journal* **3**: 5-10.

Wickham H, Grolemund G. 2016. R for Data Science. O'Reilly Media Inc. Sebastopol, CA,
USA.

Wickham H, Henry L. 2020. tidyr: Tidy Messy Data. R package version 1.1.0.
https://CRAN.R-project.org/package=tidyr

Wickham H, François R, Henry L, Müller K. 2020. dplyr: A Grammar of Data Manipulation.
R package version 1.0.0. https://CRAN.R-project.org/package=dplyr

Wilson G, Aruliah DA, Brown CT, Hong NPC, Davis M, Guy RT, Haddock SHD, et al.
2014. Best practices for scientific computing. *PLoS Biology* **12**:e1001745.

Xie Y. 2019. formatR: format R code automatically. R package version 1.7. https://CRAN.R-
project.org/package=formatR.

Yin X, Struik PC, Romero P, Harbinson J, Evers JB, van der Putten PEL, Vos J. 2009.
Using combined measurements of gas exchange and chlorophyll fluorescence to estimate
parameters of a biochemical C3 photosynthesis model: a critical appraisal and a new
integrated approach applied to leaves in a wheat (Triticum aestivum) canopy. *Plant Cell &*
Environment 32:448-464

Yin X, Sun Z, Struik PC, Gu J. 2011. Evaluating a new method to estimate the rate of leaf
respiration in the light by analysis of combined gas exchange and chlorophyll fluorescence
measurements. *Journal of Experimental Botany* **62**: 3489–3499

Supplementary Information

Table S2. List of additional {photosynthesis} functions with applications and descriptions.

Functions		
Applications	Function	Description
Gas Exchange	`aq_response`	Contains the light response model used by `fit_aq_response`
Modeling	`bake-par`	Checks that `bake` has worked
Modeling	`bake`	Temperature scales input parameters
Modeling	`conductance`	Provides calculations for leaf conductances to CO_2
Modeling	`constants`	Checks formatting of physical constant inputs
Modeling	`enviro-par`	Checks formatting of environmental inputs
Modeling	`FvCB`	Contains the model of Farquhar et al. (1980) for modeling
Gas Exchange	`gs_models`	Contains all stomatal conductance models that can be fit with `fit_gs_model`
Gas Exchange	`j_calculations`	Contains the electron transport models for fitting A-C_i curves
Modeling	`leaf-par`	Ensures proper formatting of leaf parameter inputs
Modeling	`parameter_names`	Gets a vector of parameter names
Modeling	`photosynthesis`	Contains several functions necessary for `photo`
Gas Exchange	`t_functions`	Contains all temperature response functions that can be fit with `fit_t_response`
Modeling	`utils`	Contains unit conversion functions for modeling