Cevipabulin-tubulin complex reveals a novel agent binding site on α-tubulin and provides insights into microtubule dynamic instability

Jianhong Yang[1][*][#], Yamei Yu[1][*], Yong Li[1][*], Haoyu Ye[1][*], Wei Yan[1], Lu Niu[1], Yunhua Zheng[1], Zhoufeng Wang[2], Zhuang Yang[1], Heying Pei[1], Haoche Wei[1], Min Zhao[1], Jiaolin Wen[1], Linyu Yang[1], Liang Ouyang[1], Yuquan Wei[1], Qiang Chen[1], Weimin Li[2][#], Lijuan Chen[1][#]

[1]State Key Laboratory of Biotherapy and Cancer Center, West China Hospital of Sichuan University, Chengdu 610041, China

[2] Department of Respiratory Medicine, West China Hospital, Sichuan University, Chengdu, 610041, China

*These authors equally contribute to this work

#Correspondence should be addressed to L.C. (email: chenlijuan125@163.com) or W.L. (weimi003@scu.edu.cn) or J.Y. (yjh1988@scu.edu.cn) .

Abstract

Microtubule, composed of αβ-tubulin heterodimers, remains as one of the most popular anticancer targets for decades. To date, anti-microtubule drugs mainly target β-tubulin to inhibit microtubule dynamic instability (MDI) while agents binding to α-tubulin are less well characterized and also the molecular mechanism of MDI is far from being articulated. Cevipabulin, an oral microtubule-active antitumor clinical candidate, is widely accepted as a microtubule stabilizing agent (MSA) but binds to the microtubule -destabilization vinblastine site on β-tubulin and this unusual phenomenon has so far failed to be explained. Our X-ray crystallography study reveals that, in addition binding to the vinblastine site, cevipabulin also binds to a novel site on α-tubulin (named the seventh site) which located at the region spatially corresponding to the vinblastine site on β-tubulin. Interestingly, cevipabulin exhibits two unique site-dependent functions. Cevipabulin binding to the seventh site promotes tubulin degradation through interaction of the non-exchengeable GTP to reduce tubulin stability. Cevipabulin binding to the vinblastine site enhances longitudinal interactions but inhibits lateral interactions of tubulins, thus inducing tubulin protofilament polymerization (but not microtubule polymerization like MSAs), and then tangling into irregular tubulin aggregates. Importantly, the tubulin-cevipabulin structure is an intermediate between "bent" and "straight" tubulins and the involved bent-to-straight conformation change will be helpful to fully understand the molecular mechanism of tubulin assembly. Our findings confirm cevipabulin is not an MSA and shed light on the development of a new generation of anti-microtubule drugs targeting the novel site on α-tubulin and also

provide new insights into MDI.

Key words

Cevipabulin; Tubulin Inhibitor; Microtubule Dynamic Instability; Novel Binding Site;

Tubulin Degrader.

Introduction

Microtubules play key roles in many important cell events, especially cell division, and

thus remain as one of the most popular anticancer targets for decades [1, 2].

Microtubules are composed of αβ-tubulin heterodimers assembled into linear

protofilaments and their packaging demands both lateral and longitudinal interactions

between tubulins [3]. To date, various tubulin inhibitors have been reported to alter the

lateral and/or longitudinal interactions to promote microtubule assembly or disassembly,

including the clinical most popular anticancer drugs: vinca alkaloids, taxanes, eribulin

et al [4, 5]. These drugs all target β-tubulin, which has five different binding sites

(colchicine, vinblastine, paclitaxel, laulimalide and maytansine sites) [5]. By

overexpression of β-tubulin isoforms, especially βIII-tubulin, cancer cells are prone to

become resistant to these therapies [6]. So far, the pironetin site is the only one located

on α-tubulin [5, 7]. However, this site is too small and pironetin has six chiral centers

in its molecular structure, making it difficult to be synthetized. Since the crystal

structure of tubulin-pironetin was reported in 2016 [5, 7], no significant progress has

been made in the design of pironetin-binding-site inhibitors or even analogues of

pironetin.

Microtubule dynamic instability (MDI) is referred to the random switching between

microtubule regrowth and shrinkage, which accompanied by periodic cycles of "bent"

to "straight" conformation change in tubulin protofilament [8]. However, the detailed

molecular mechanism of bent-to-straight conformation transition is unclear. Structural

study of complexes of tubulin with the six known binding-site inhibitors allows in detail

description of how inhibitors bind to and change the conformation of tubulin to alter

MDI [3, 5, 9-12]. For example, inhibitors binding to paclitaxel or laulimalide site, the

only two microtubule stabilization agents (MSAs) sites [13], stabilize the M-loop on β-

tubulin to enhance lateral interactions to promote tubulin polymerization [9, 11],

revealing a structuring of the M-loop into a short helix during tubulin polymerization

[11]; Inhibitors binding to colchicine site, a widely known microtubule destabilization

agents (MDAs) site [12], bind to the intra-dimer interfaces to inhibit flipping in of T7

loop on β-tubulin to inhibit tubulin polymerization [12, 14], thus demonstrating a

flipping in and out of the T7 loop participating in MDI [14]. With the current existing

tubulin-inhibitors complexes, some of the local conformation changes of MDI are

easily observed. However, due to the lack of an intermediate structure between "bent"

and "straight" of tubulin, the most important "bent" to "straight" conformation change

has never been detailly described and the underlying molecular mechanism remains

elusive.

Cevipabulin (or TTI-237) is a synthetic tubulin inhibitor with *in vivo* anticancer

activity and has been used in clinical trials investigating the treatment of advanced

malignant solid tumors [15]. Competition experiment showed it competed with [3]H-

vinblastine but not [3]H-paclitaxel for binding to microtubules, indicating it binds to the

classic tubulin-depolymerization vinblastine site [16]. However, an *in vitro* tubulin

polymerization assay exhibited that cevipabulin did not inhibit tubulin polymerization

as vinblastine but promoted tubulin polymerization as paclitaxel [16]. These studies

concluded that cevipabulin seems displaying mixed properties between paclitaxel and

vinblastine. More recently, Kovalevich *et al.* identified two unusual characters of

cevipabulin. Cevipabulin could promote tubulin degradation and induce lots of tubulin

aggregates in cell cytoplasm which have never observed in reported known tubulin

inhibitors[17]. Despite extensive efforts, the unusual phenomenon and characters of

cevipabulin have so far failed to be explained and the underlying mechanism of action

on microtubule has not been clearly defined [16, 18-21]. Recently, Gonzalo *et al*.

synthetized an analogue of cevipabulin (named compound **2** in this paper) and got the

crystal structure of tubulin-compound **2** complex (PDB code: 5NJH) and prove

compound **2** binds to the vinblastine site of β -tubulin to enhance longitudinal

interactions and induced formation of tubulin bundles in cell, which seems like that

compound **2** binding to vinblastine site could really induce tubulin polymerization in a

paclitaxel-like manner[18]. However, they didn't observe compound **2** induced

microtubule through transmission electron microscope (TEM) *in vitro*, which makes

their conclusion not rigorous enough.

In this study, we solved the crystal structure of tubulin-cevipabulin complex and were

surprised to find that cevipabulin simultaneously binds to two spatially independent

sites: the vinblastine site and a new site on α-tubulin (called the seventh site). Detailed

mechanism revealed that cevipabulin binding to the two sites exhibited two different

and independent function: binding to the seventh site induces tubulin degradation and

binding to vinblastine site leads to tubulin protofilament polymerization and then

formation of irregular tubulin aggregates. Our study reveals that the increase in turbidity

caused by cevipabulin is not the consequence of microtubule polymerization but the

results of protofilament polymerization of tubulin, which well explains its paclitaxel

like phenomenon but undefined function previously. Structure-activity-relationship

demonstrate that trifluoropropanyl of cevipabulin plays a critical role in binding to the

seventh site. Our study reveals a novel binding site on α-tubulin related to tubulin

degradation effect and lays a foundation for the rational design of new generation of

anticancer drugs. Importantly, we define a novel tubulin inhibition mechanism:

enhancing longitudinal and inhibiting lateral interactions to induce formation of

irregular tubulin aggregates and the involved bent-to-straight conformation change

provides new insights into MDI.

Results

Cevipabulin induces tubulin degradation and formation of irregular tubulin aggregates

To elucidate the cellular effect of cevipabulin at an early time point, we carried out

label-free quantitative proteomic analysis on six-hour cevipabulin treated human

cervical adenocarcinoma cell line-HeLa. Cevipabulin significantly down-regulated the

protein level of α, β-tubulin and their isoforms with high selectivity (Fig.1a).

Immunoblotting study confirmed cevipabulin decreased tubulin proteins in HeLa,

human colon colorectal carcinoma cell line Hct116, human large cell lung carcinoma

cell line H460 and human B cell lymphoma cell SU-DHL-6 in a dose-dependent manner

(Fig.1b) and time dependent manner in HeLa cells (Fig. S1a), demonstrating that the

reduction of tubulin is a common biochemical consequence of cevipabulin treatment in

cancer cells. The quantitative PCR assay showed that cevipabulin had no effect on α-

and β-tubulin mRNA levels (Fig.S1b), indicating that the downregulation of tubulin

protein by cevipabulin is post-transcriptional. MG132, a proteasome inhibitor, could

completely block cevipabulin-induced tubulin degradation (Fig.S1c). All these proved

that cevipabulin promoted tubulin degradation in a proteasome dependent pathway.

Immunofluorescence staining of tubulin is commonly used to detect microtubule

morphology in cells treated with tubulin inhibitors [22]. Untreated cells presented

normal microtubule network in cells (Fig.1c). MSAs, such as paclitaxel, induced

excessive tubulin polymerization and presented bunches of microtubules in cells

(Fig.1d). MDAs, such as colchicine, inhibited tubulin polymerization and completely

destroyed microtubules (Fig.1e). Vinblastine, another MDA, inhibited tubulin

polymerization at low concentration (the same as colchicine, Fig. 1f), but induced the

formation of tubulin paracrystals in the cytoplasm at high concentration (Fig. 1g),

which was considered as packing of spiral protofilaments [3]. Interestingly, in cells

treated with cevipabulin we observed a large number of irregular tubulin aggregates

formation throughout the cytoplasm (Fig.1h), which was totally different from

traditional MSAs and MDAs.

148
Figure 1. Cevipabulin promotes α-and β-tubulin degradation and induces the formation of
irregular tubulin aggregates. (**a**) Label-free quantitative proteomic analysis of total proteins from
HeLa cells treated with 1 μM cevipabulin for 6 h. This graph presents fold-changes of 1825
quantified proteins between cevipabulin and DMSO treatment groups versus the p value (t test;
triplicate analysis). Three biological repetitions are performed. (**b**) Immunoblotting analysis of both
α and β-tubulin levels in HeLa, Hct116, H460 and SU-DHL-6 cells, which all are treated with
indicated concentrations of cevipabulin for 16 h. Results are representative of three independent
experiments. (**c-h**) Hela cells are treated with (**c**) DMSO, (**d**) paclitaxel (1 μM), (**e**) colchicine (1
μM), (**f**) vinblastine (1 μM), (**g**) vinblastine (10 μM) and (**h**) cevipabulin（3 μM）for 1 hour and
then subjected for immunofluorescence analysis with α-tubulin antibody to monitor morphology of
microtubule. Bar=10μm. Results are representative of three independent experiments. Cev:
cevipabulin.

Crystal structure of tubulin-cevipabulin reveals its simultaneously binding to the

vinblastine site and a novel site on α-tubulin

To analyze the binding details of cevipabulin (Fig. 2a) to tubulin, we soaked cevipabulin

into the crystals consisting of two tubulin heterodimers, one stathmin-like protein RB3

and one tubulin tyrosine ligase (T2R-TTL) [11]. The crystal structure of tubulin-

cevipabulin complex was determined to be 2.6 Å resolution (Table S1). The whole

structure was identical to previously reported [11], two tubulin heterodimers were

arranged in a head to tail manner (α1β1-α2β2) with the long helix RB3 comprising both

dimers and tubulin tyrosine ligase docking onto α1-tubulin (Fig. 2b). The Fo–Fc

difference electron density unambiguously revealed two cevipabulin molecules binding

to two different sites (Fig. 2c and 2d): one at the inter-dimer interfaces between the β1-

and α2-tubulin subunits (the vinblastine site) and the other one at the intra-dimer

interfaces between α2- and β2-tubulin subunits (Fig. 2b) and the later binding region is

a new binding site (here named as the seventh site).

The binding region of cevipabulin in the vinblastine site was formed by residues from

βH6, βH7, βT5 loop, αH10 and αT7 loop (Fig.2e). As presented in Figure 2f, the side

chain of βY224 made π-π stacking interactions with triazolopyrimidinyl group of

cevipabulin and the guanine nucleobase of GDP (Fig.2f). Seven hydrogen bonds (N1

atom to side chain of βY224; N3 atom to main-chain nitrogen of βY224 through a water;

N4 atom to main-chain nitrogen of βY224; 5- chlorine atom to both main-chain nitrogen

of βY224 and βT223; 2'- fluorine atom to site chain of βY224 and main-chain nitrogen

of βN206) between cevipabulin and β1-tubulin were observed. The -NH- group on the

cevipabulin side chain formed a salt bridge with βD211. Besides, cevipabulin also

exhibited four hydrogen bonds with α2-tubulin (oxygen atom on side chain to the side

chain of αN329; 2'- fluorine atom to the main-chain nitrogen of αN326; one fluorine

atom of trifluoropropanyl to both main and side chain of αN326) (Fig.2g).

The seventh site on α2-tubulin is formed by residues from αH1, αH6, αH7, αT5, βH10

and βT7 (Fig.2h). Similar to the vinblastine site, triazolopyrimidinyl of cevipabulin at

this site also made π-π stacking interactions with the side chain of αY224 and the

guanine nucleobase of GTP (Fig. 2i). There were eight hydrogen bonds (N1 atom to

side chain of αY224; N4 atom to main-chain nitrogen of αY224; 5- chlorine atom to

main-chain nitrogen of αT223; 2'- fluorine atom to site chain of αN206; 6'- fluorine

atom to site chain of αR221; One fluorine atom of trifluoropropanyl to side chain of

αN206; Another fluorine atom of trifluoropropanyl to both O2' AND O3' of GTP)

between cevipabulin and α2-tubulin and a salt bridge between the -NH- group of

cevipabulin side chain and αD211 (Fig. 2i). Notably, there is no hydrogen bond between

cevipabulin and β2-tubulin at this new site.

Figure 2. Crystal structure of tubulin-cevipabulin complex. (**a**) Chemical structure of
cevipabulin. (**b**) Overall structure of tubulin-cevipabulin complex. TTL is colored yellow, RB3 is
green, α-tubulin is dark and β-tubulin is grey. Cevipabulin on β1-tubuin and α2-tubulin are all shown
in spheres and colored yellow. (**c, d**) Electron densities of cevipabulins on (**c**) β1-tubulin or (**d**) α2-
tubulin. The Fo-Fc omit map is colored light blue and contoured at 3δ. (**e, f**) Close-up view of
cevipabulin binding to (**e**) β1-tubulin or (**f**) α2-tubulin. GDP or GTP is shown in magenta sticks.
Cevipabulin is shown in yellow sticks. Side chain of β1-Y224 or α2-Y224 is show in grey sticks.
(**g, h**) Interactions between (**g**) β1-tubulinn and vinblastine-site cevipabulin or (**h**) α2-tubulin and
vinblastine-site cevipabulin. Coloring is the same as in (**e**). Residues from tubulin that form
interactions with vinblastine-site cevipabulin are shown as sticks and labeled. Hydrogen bonds are
drawn with red dashed lines. (**i**) Interactions between α2-tubulin and the-seventh-site cevipabulin,
color is the same as in (**f**), residues from tubulin that form interactions with the-seventh-site
cevipabulin are shown as sticks and are labeled. Hydrogen bonds are drawn with red dashed lines.
Cev: cevipabulin.

Cevipabulin binding to vinblastine site induces the formation of irregular tubulin

aggregates while binding to the seventh site induces tubulin degradation

To address the functions of these two sites, we used vinblastine to block the vinblastine

site or single amino acid substitution (Y224G on α-tubulin) to block the seventh site. When vinblastine site was occupied, cevipabulin lost its ability to induce irregular tubulin aggregation, while retaining the tubulin-degradation effect (Fig. 3a and 3b). When the seventh site was mutant, cevipabulin lost the tubulin-degradation effect but persist inducing irregular tubulin aggregation (Fig. 3c and 3d). These data indicate that cevipabulin binds to the vinblastine site inducing irregular tubulin aggregation, while binds to the seventh site led to tubulin degradation.

To independently study the functions and structure activity relationship between these two sites, we employed two reported cevipabulin analogues (compounds **1 [17]** and **2 [18])** for further study **(**Fig. 3e), which only bound to the seventh site or the vinblastine site, respectively. Compared with cevipabulin, compound **1** lacks the N- substituted side chain. Further, the trifluoropropanyl in compound **1** was replaced by an azabicyclo to obtain compound **2**. We found compound **1** only induced tubulin degradation (Fig. 3f) and did not lead to irregular tubulin aggregation (Fig. 3g), in contrast, compound **2** only induced irregular tubulin aggregation but not tubulin degradation (Figs. 3f, 3g). Competition assay indicated that αY224G mutation, but neither vinblastine nor compound **2,** inhibited compound **1** induced tubulin degradation (Fig. 3h, S2a and S3b). Vinblastine, rather than compound **1** or αY224G mutation, suppressed compound **2** caused irregular tubulin aggregation (Fig. 3i, S2c and S2d). These results demonstrated that compound **1** only bond to the seventh site while compound **2** only bound to the vinblastine site, and also implied that trifluoropropanyl of cevipabulin played critical role in binding to the seventh site.

Figure 3. Cevipabulin binds to vinblastine site to induce formation of irregular tubulin aggregates while binds to the seventh site to induce tubulin degradation. (**a**) HeLa cells were treated with 10 μM vinblastine for 1 h and then further treated with 1 μM cevipabulin for 16 h. The α-tubulin protein level was detected by immunoblotting. Results are representative of three independent experiments. (**b**) HeLa cells treated with or without 10 μM vinblastine for 1 h before treated with 1 μM or 10 μM cevipabulin for another hour. Irregular tubulin aggregates were detected using immunofluorescence and the number of irregular tubulin aggregates was counted for randomly chosen 30 cells. ***p<0.00001. Results are representative of three independent experiments. (**c**) Vectors expressing either wild type or Y224G mutant GFP-tubulin were transfected to HeLa cells. After 24 hours, cells were treated with or without 1μM cevipabulin for 16 h. Then the protein level of GFP-α-tubulin was detected by immunoblotting. Results are representative of three independent experiments. (**d**) Vectors expressing either wild type or Y224G mutant GFP-α-tubulin were transfected to HeLa cells. After 24 hours, cells were treated with or without 1 μM cevipabulin for 1 h. Irregular tubulin aggregates were detected using immunofluorescence and the number of irregular tubulin aggregates was counted for randomly chosen 30 cells. ns: no significant difference. Results are representative of three independent experiments. (**e**) Chemical structure of cevipabulin derivatives. (**f**) Hela cells were treated with indicated compounds for 16 h. Then the protein level of α-tubulin was detected by immunoblotting. Results are representative of three independent experiments. (**g**) Hela cells were treated with indicated compounds for 1 h and irregular tubulin aggregates were detected using immunofluorescence and the number of irregular tubulin aggregates was counted for randomly chosen 30 cells. ns. no significant difference. ***p<0.0001 in comparison with the control. Results are representative of three independent experiments. (**h**) HeLa cells were treated with or without 30 μM compound **2** for 1 hour before treated with 10 μM compound **1** for

16 h and then the protein level of α-tubulin was detected by immunoblotting. Results are
representative of three independent experiments. (**i**) HeLa cells were treated with or without 30 μM
compound **1** for 1 hour before treated with 10 μM compound **2** for another hour. The irregular
tubulin aggregates were detected by immunofluorescence and the number of irregular tubulin
aggregates was counted for randomly chosen 30 cells. ns. no significant difference. Results are
representative of three independent experiments. Cev: cevipabulin. **1**: compound **1**; **2**: compound **2**;
Vin: vinblastine.
272

*Compound **2** promotes protofilaments polymerization by enhancing longitudinal and*

inhibiting lateral interactions of tubulins

Gonzalo *et al.* reported the crystal structure of tubulin-compound **2** complex (PDB code:

5NJH) and revealed compound **2** bound to the vinblastine site can to enhance

longitudinal interactions and induce the formation of tubulin bundles in cells. It seems

that the binding of compound **2** to vinblastine site could induce tubulin polymerization

in a paclitaxel-like manner [18]. Here we further and detailly investigated its effect on

tubulin in cells. Immunofluorescence study showed that compound **2** induced short

tubulin bundles (seems like microtubule bundles) formation at low concentrations (3

μM) for 1 hour treatment (Fig. 4b), which was in consistent with the published data

[18]. However, when the concentration of compound **2** was increased to 10 μM, some

of the tubulin bundles turned into tubulin aggregates (Fig. 4c). At higher concentration

of 30 μM, the whole cytoplasm was all filled with irregular tubulin aggregates, and no

tubulin bundles were observed (fig.4d), which was the same as cevipabulin treatment.

Interestingly, as cells treated with 3 μM compound **2** for longer time (4h or 8h), the

short tubulin bundles will also turn into irregular tubulin aggregates (Fig. 4e and 4f).

To further investigate this unusual characteristic, we treated purified tubulin with

compound **2** and then analyzed with TEM. As presented in Figure 4g, compound **2**

induced numbers of linear structure formation, which entangled each other to form a

bundle of tubulin or tubulin aggregates. The diameter of the thinnest linear structure

was about 6~7 nm (Fig. 4h), which was much smaller than that of microtubule (Fig.

s3a), but perfectly matched the diameter of tubulin protofilament. These results

indicated that compound **2** induced tubulin polymerizing into tubulin protofilament, but

not microtubule as previously suggested [18]. Thus, it is reasonable to assume that the

unusual irregular tubulin aggregates caused by compound **2** observed in cell cytoplasm

are the consequence of randomly stacking and aggregation of tubulin protofilaments.

Figure 4. Compound 2 induced tubulin protofilaments polymerization. (a-f) Hela Cells treated
with (**a**) 0, (**b**) 3, (**c**) 10 or (**d**) 30 μM compound **2** for 1 hour or treated with 3 μM compound **2** for

302 (**e**) 4 or (**f**) 8 h. Cells were then subjected for immunofluorescence analysis with α-tubulin

303 antibody to monitor the morphology of microtubule. Bar=10μm. Results are representative of

304 three independent experiments. (**g**, **h**) Purified tubulin (2mg/ml was incubated with 50 μM

305 compound **2** for 30min at room temperature before imaged with TEM. Results are representative

306 of three independent experiments. Both large tubulin protofilaments aggregation (**g**) and single

307 tubulin protofilament (**h**) were observed. Results are representative of three independent

308 experiments.

310 Crystal structures of tubulin-compound **2** (PDB code: 5NJH) and tubulin-cevipabulin

311 could be superimposed very well in whole (Fig S3b, with a root-mean-square deviation

312 (RMSD) of 0.45 Å over 1,930 Cα atoms) or in their binding region (Fig S3c). Hence,

313 we used the tubulin-compound **2** structure for structural mechanism analysis. We

314 superimposed the β1-tubulin subunit of tubulin-compound **2** to the one of the apo-

315 tubulin structure (PDB code: 4I55). In the inter-dimer interfaces, compound **2** led to a

316 6.7 Å shift of the αH10 helix of α2-tubulin towards β1-tubulin (Fig. S3d), and thus the

317 conformation of tubulin-compound **2** complex was arranged in a more "straight"

318 manner than that of apo tubulin complex (Fig. S3e), implying compound **2** enhanced

319 longitudinal interactions of tubulin dimers. We then superimposed tubulin-compound **2**

320 complex to a polymerized microtubule structure (PDB code: 6DPV). The individual

321 β1-tubulin or α2-tubulin in tubulin-compound **2** complex align better than those in apo

322 tubulin structure to the corresponding subunits in polymerized microtubule structure

323 (RMSD: 0.895 Å and 0.830 Å for β1-tubulin and α2-tubulin in tubulin-Compound **2**

324 complex, respectively; 1.392 Å and 1.261 Å for β1-tubulin and α2-tubulin in apo

325 tubulin, respectively), suggesting compound **2** caused both β1-tubulin or α2-tubulin to

326 take a more "polymerized" state. Focusing on the interface of β1-tubulin and α2-tubulin,

327 compound **2** binding caused significant movement of αT7, αH8 and αH10 to from a

depolymerized-to-polymerized state transformation (Fig. 5a and 5b). We noticed that

in tubulin-compound **2** complex, the T5 loop on α2-tubulin in the intra-dimer interface

showed an obvious shift from depolymerized to polymerized state (Fig.5c). As there is

no ligand binding to the intra-dimer interface in the tubulin-compound **2** complex, the

αT5 loop outward shift might be allosterically mediated by the inter-dimer interface

conformation change. We infer that the more compact intra-dimer interface induced by

compound **2** can make tubulin prone to form a straight dimer like that in the

polymerized microtubule (Fig.5d), and thus straighten tubulin protofilaments.

Therefore, we uncovered a continuous conformational change which could mimic the

bent-to-straight conformation change during tubulin polymerization. With compound **2**

binding as a small wedge, these straight protofilaments still has a 4.2° curvature at the

inter-dimer interface (Fig. 5e). Also, we could clearly observe a clash between αH10 in

polymerized microtubule structure and compound **2** (Fig. S3f), suggesting compound

2 binding obstructs the straight conformation of tubulins. This is in line with the fact

that compound **2** can not be incorporated into polymerized microtubule [18]. To analyze

the lateral interaction, we aligned tubulin-compound **2** complexes to two adjacent

protofilaments in polymerized microtubule structure based on β1-tubulin. The M-loop,

which is important for lateral interaction, exhibited a 5.4 Å shift between tubulin-

compound **2** complex and polymerized microtubule (Fig. S3g), suggesting the M-loop

is in a polymerization unfavored status. Thus, we revealed for the first time that

cevipabulin or compound **2** enhanced tubulin longitudinal interactions while inhibited

lateral interactions to induce excessive polymerization of tubulin protofilaments

 (Fig.5f). Our results also confirm that cevipabulin is not an MSA as previously reported.

Figure 5. Structural mechanism of cevipabulin and compound 2 induced tubulin protofilaments polymerization. (**a**) Overview of the aligned structures of apo tubulin (PDB code: 4I55) and a polymerized microtubule (PDB code: 6DPV). The crystal structures of apo tubulin (green) and the polymerized microtubule (violetpurple) are superimposed on α2-tubulin subunits. β1, α2 and β2-tubulin subunits of apo tubulin structure are shown while only α2-tubulin subunit the polymerized microtubule structure is shown. αH10, αH8, αT7 and αT5 are colored blue in apo tubulin structure while red in the polymerized microtubule structure. (**b**) Tubulin-compound **2** (salmon, PDB code: 5NJH) is aligned to the superimposed complexes in (**a**) based on α2-tubulin subunit. Close-up view of the β1-α2-tubulin inter-dimer interface reveals that αH10 and αH8 in tubulin- compound **2** have a significant movement from a "depolymerized" (apo tubulin) state to a "polymerized" (polymerized microtubule structure) state. (**c**) Close-up view of the α2-β2 intra-dimer interface. The αT5 loop in tubulin-compound **2** has an outward shift to match the αT5 loop in the polymerized microtubule structure. (**d**) View of the α2, β2-tubulin interaction of the aligned complexes in (**a**). The αT5 loops of apo tubulin and microtubule are highlighted in red and blue, respectively. The αT5 loop outward shift makes room for β2-tubulin to bind closer to α2-tubulin. (**e**) View of the β1, α2-tubulin interaction of the aligned complexes in (**b**). Compared to the polymerized microtubule structure (violetpurple), The β1, α2-tubulin inter-dimer interface exhibits a 13.5° bend angle in apo tubulin structure (green) while only 4.2° in tubulin-compound **2** complex (salmon). (**f**) The molecular mechanism of cevipabulin induced tubulin protofilaments polymerization. Cev: cevipabulin. Cpt 2: compound **2**.

Cevipabulin and compound 1 destabilize tubulin by interacting with the GTP on "nonexchangeable site" to promote tubulin degradation

We then investigated the tubulin degradation effect of cevipabulin binding to the seventh site. At the seventh site, cevipabulin bound to the intra-dimer interface and made multiple polar contacts with α2-tubulin. In particular, the trifluoropropanyl of cevipabulin formed two hydrogen bonds with non-exchangeable GTP (Fig. 2i), which plays a structural role and is important for the stability of tubulin dimers [23, 24]. This non-exchangeable GTP forms a number of hydrogen bonds with surrounding amino acid residues and a magnesian ion [23]. Single mutation abolishing hydrogen bond with this GTP could reduce the affinity of GTP and absence of the magnesian ion would reduce the protein stability [23, 25]. We speculated that the interaction between cevipabulin and the non-exchangeable GTP could decrease tubulin stability and subsequently promote tubulin degradation. Differential scanning fluorimetry (DSF) , a method monitoring protein unfolding, results showed that tubulin treated by compound **1** had an obvious lower Tm value (melting temperature) than DMSO treated, while compound **2** increased the Tm value and cevipabulin had no significant effect on the Tm value (Fig. 6a). Further the DSF results showed that compound **1** (only binds to the seventh site) reduces the stability of tubulin but compound **2** (only binds the vinblastine site) increases the stability of tubulin, and cevipabulin (binds to both sites) may balance these two effects and represent a neutralized output. TEM results also showed that compound **1** decreased tubulin stability as evidenced by lots of spherical tubulin aggregates (denatured or unfolding tubulin) observed upon compound **1** treatment (Fig.

6b). Using a thiol probe, tetraphenylethene maleimide (TPE-MI), which is non-

fluorescent until conjugated to a thiol [26], we further measured whether these

compounds promote unfolding of tubulin. TPE-MI alone did not increase fluorescence

of tubulin while addition of 4M guanidine hydrochloride (non-selective protein

denaturant) significantly increased fluorescence (Fig. 6c). Cevipabulin and compound

1 obviously increased tubulin fluorescence while vinblastine and compound **2** had no

such effects (Fig. 6c), demonstrating that cevipabulin or compound **1** could promote

unfolding of tubulin. In addition, PYR-41, an inhibitor of ubiquitin-activating enzyme

E1, totally blocked compound **1** induced tubulin degradation (Fig. 6d), suggesting

destabilized tubulin is removed by normal housekeeping ubiquitinylation. Therefore,

cevipabulin and compound **1** decrease tubulin stability by direct interaction with the

non-exchangeable GTP to subsequently promote its destabilization and degradation.

Figure 6. Cevipabulin or compound 2 decrease tubulin stability to promote tubulin destabilization and degradation. (**a**) Thermal unfolding curves of DMSO, cevipabulin (10 μM) , vinblastine (10 μM), compound **1** (10 μM) or compound **2** (10 μM) treated purified tubulin （2 μM） by a differential scanning fluorimetry (DSF) method. Plots of the fluorescence F350/F330 ratio and its first derivative are shown. The maximal values of the first derivatives are regarded as the melting temperature (Tm value). Results are representative of three independent experiments. (**b**) Purified tubulin (2mg/ml was incubated with 50 μM compound **1** for 30min at room temperature before imaged with TEM. Results are representative of three independent experiments. (**c**) Tubulin unfolding detected by TPE-MI. Tubulin (0.2 mg/ml) in PIPES buffer was mixed with 50 μM TPE-MI and the indicated compounds. Fluorescence (Excitation wavelength: 350nm; Emission wavelength: 470nm) were detected every half minute for 60 min. Results are representative of three independent experiments. (**d**) Hela cells were treated with or without PYR-41(20 μM) for 1 hour before treated with 10 μM compound **1** for 16 h. Protein level of α-tubulin were detected by immunoblotting. Results are representative of three independent experiments. Cev: cevipabulin; Vin: vinblastine. **1**: compound **1**；**2**: compound **2**.

Discussion

Our study identifies a novel binding site on α-tubulin, the seventh site. As this new site

is located near the non-exchangeable GTP site and this GTP is important for tubulin

stability [23-25], inhibitors such as cevipabulin and compound **1** binding to the seventh

site may reduce tubulin stability and promote tubulin degradation. This novel site on α-

tubulin is spatially corresponding to the vinblastine site on β-tubulin, which is also

bound by cevipabulin. Cevipabulin binding to the vinblastine site enhances the

longitudinal interaction within tubulin protofilaments to make them take a straighter

conformation while blocks tubulin lateral interaction, causing excessive tubulin

protofilaments polymerization, which randomly stack into irregular tubulin aggregates.

The binding pocket of cevipabulin to these two sites is very similar (formed by αH1,

αH6, αH7, αT5, βH10, βT7 for the seventh site and βH1, βH6, βH7, βT5, αH10, αT7

for the vinblastine site) and the binding modes of cevipabulin are also similar except

the trifluoropropanyl of cevipabulin adopts different conformations. Vinblastine-site

cevipabulin is mainly located on β1-tubulin and makes lots of hydrogen bond with β1-tubulin while its trifluoropropanyl is oriented towards α2-tubulin and makes four hydrogen bond interactions with α2-tubulin. The-seventh-site cevipabulin is totally located on α2-tubulin and makes lots of hydrogen bond with α2-tubulin and its trifluoropropanyl is also oriented towards α2-tubulin to establish hydrogen bonds with the non-exchangeable GTP. Of note, compound **2** lacking the trifluoropropanyl could not bind to the seventh site and showed no tubulin degradation effect, suggesting the trifluoropropanyl-GTP interaction is important for cevipabulin binding to the seventh site. We noticed that in the tubulin-compound **2** complex, although compound **2** bound only to the vinblastine site, the αT5 loop at the seventh site also had an outward shift like in the tubulin-cevipabulin complex. It seems to suggest that cevipabulin binds to the vinblastine site to allosterically affect the αT5 loop with an unknown mechanism, and then creates the pocket for cevipabulin binding to α2-tubulin. However, vinblastine binding showed no such allosteric effect on αT5 loop (Fig.S3h) and can not block cevipabulin binding to the seventh site, implying cevipabulin can bind to the seventh site and affect αT5 loop itself. As compound **2** has no degradation effect but has allosteric effect on αT5 loop, we can be sure that the αT5 loop shift does not contribute to the degradation effect. Although we confirmed that compound **1** binds only to the seventh site and not to the vinblastine site, we unfortunately did not obtain the crystal structure of tubulin-compound **1** complex (possible due to compound **1**'s lower affinity to the seventh site) which might provide other vital information of the seventh site.

We confirmed that cevipabulin is not an MSA as previous reported. The previously

observed turbidity increase [16] in *in vitro* tubulin polymerization assay is likely due to tubulin protofilaments polymerization rather than microtubule polymerization. Therefore, we believe that confirmation of MSA using only in vitro tubulin polymerization assay is not rigorous. In many CNS diseases, dysregulation of microtubule structure and dynamics is commonly observed in neurons [17, 27, 28] and stabilization of microtubules by MSA is a promising therapeutic strategy [27, 29]. However, the traditional MSAs binding to the paclitaxel site have relative large molecule weights and can not penetrate the blood–brain barrier [27]. Researchers then focus on developing brain-penetrant MSAs with smaller molecule weight that could be readily synthesized, such as cevipabulin and its derivatives for the treatment of CNS diseases [30-32]. Our study indicates cevipabulin and its derivatives do not stabilize microtubule instead inversely promoting its degradation or inducing excessive tubulin protofilaments formation. These new mechanisms should be considered when studying the cevipabulin and its derivatives on CNS diseases, or there are other undiscovered mechanisms supporting their effects on CNS diseases.

Microtubules assembly demands bent-to-straight conformation change of tubulin dimers and protofilaments [8]. This conformation change is widely accepted, but the detailed molecular mechanism remains elusive. Microtubules polymerization demands GTP-bound tubulin dimers and there are two opposite models to describe the connection between GTP and tubulin dimer conformation change. The allosteric model claims that GTP binds to tubulin dimers, causing a remote allosteric conformation change to generate straighter tubulin protofilaments, and then lateral interactions

establishing lateral interactions to form microtubules [33-35]. However, the lattice

model suggests that the GTP binding has no change on tubulin dimer conformation, but

the lattice assembly straightens tubulin dimer to incorporate into microtubules [33, 36].

Since most of the evidences in support of the two models are indirect, the debates can

not be resolved. Here, high-resolution crystal structures of tubulin-cevipabulin and

tubulin-compound **2** complexes directly revealed these compounds can cause a bent-

to-straight conformation change in tubulin protofilament: two adjacent tubulin dimers

get closer, transforming the inter-dimer-interface α-tubulin into a "polymerized" state,

including significant movement of αH10, αH8 and αT7 in the inter-dimer interface and

a succedent remotely allosteric mediated αT5 outward shift in the intra-dimer interface.

The αT5 outward shift then makes the intra-dimer-interface β-tubulin bind to the α-

tubulin closer to form a straight dimer. We think this continuous conformation change

could reflect the bent-to-straight conformation change of tubulin dimers in normal

physiological condition, and provide clearly insights into MDI. More importantly, with

these compounds binding as a small wedge, these straight protofilaments still have a

4.2° curvature at the inter-dimer-interface and establish no lateral interaction,

demonstrating the lateral interaction requires strict straight tubulin protofilaments and

the lateral interaction is the consequence rather than cause of straight conformation

formation. Therefore, our data supports the allosteric model.

Here, we reported a novel binding site on α-tubulin that possessed tubulin degradation

effect that was distinct from the traditional MDAs and MSAs. Using this specific site,

a new class of tubulin degraders can be designed as anticancer drug targeting α-tubulin.

We also presented a novel function of tubulin inhibitors-induced irregular tubulin

aggregation by enhancing longitudinal but blocking lateral interaction of tubulin, and

the involved conformation change will provide insights into MDI.

Materials and Methods

Reagents

Colchicine, vinblastine, paclitaxel, β,γ-Methyleneadenosine 5′-triphosphate disodium

salt (AMPPCP), Tetraphenylethene maleimide (TPE-MI), and DL-dithiothreitol (DTT)

were purchased from Sigma; Guanidine，hydrochloride, MG132 and PYR-41 were

obtained from Selleck; Cevipabulin was from MedChemExpress; Purified tubulin was

bought from Cytoskeleton, Inc.; Antibodies (α-tubulin antibody, β-tubulin antibody,

GAPDH antibody and gout anti mouse second antibody) were bought from Abcam.

Chemistry

All the chemical solvents and reagents used in this study were analytically pure without

further purification and commercially available. TLC was performed on 0.20 mm silica

gel 60 F_{254} plates (Qingdao Ocean Chemical Factory, Shandong, China). Visualization

of spots on TLC plates was done by UV light. NMR data were measured for ^{1}H at 400

MHz on a Bruker Avance 400 spectrometer (Bruker Company, Germany) using

tetramethylsilane (TMS) as an internal standard. Chemical shifts were quoted in parts

per million. High Resolution Mass Spectra (HRMS) were recorded on a Q-TOF Bruker

Daltonics model IMPACT II mass spectrometer (Micromass, Manchester, UK) in a

positive mode.

Scheme1: Reagents and conditions: a) diethyl malonate, NaH, CuI, dioxane, r.t.-reflux; b) 3-amino-1,2,4-triazole, tributylamine, 180 °C; c) POCl₃, reflux; d) amine, K₂CO₃, DMF, r.t.

General procedure for the synthesis of diethyl 2-(2,4,6-trifluorophenyl)malonate (**5**)

To a stirred solution of diethyl malonate (320 mg, 2.0 mmol) in 1,4-dioxane was added 60% sodium hydride (96 mg, 2.4 mmol) by portions at room temperature. Then cupper (I) bromide (380 mg, 2.0 mmol) and compound **4** (211mg, 1.0 mmol) was added. The reaction mixture was stirred at room temperature for 30 minutes and then refluxed for 8 hours under nitrogen protection. After completion of the reaction, the mixture was cooled to room temperature and hydrochloric acid (12 N, 50 mL) was added slowly. The organic phase was separated off and the aqueous phase was extracted with ethyl acetate (×2). The combined organic phase was concentrated in vacuo. The residue was purified by chromatograph on silica gel with petroleum ether and ethyl acetate as eluent to give compound **5** as a white solid. Yield: 62%. ^{1}H NMR (400 MHz, DMSO) δ 7.36 − 7.18 (m, 2H), 5.15 (s, 1H), 4.18 (q, J = 7.1 Hz, 4H), 1.23 − 1.14 (m, 6H). HRMS-ESI: calcd for $[C_{13}H_{13}F_3O_4+Na]^+$ 313.0664, found: 313.0663.

General procedure for the synthesis of 5,7-dichloro-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5-*a*]pyrimidine (**7**)

A mixture of 3-amino-1,2,4-triazole (84 mg, 1.0 mmol), compound **5** (290 mg, 1.0 mmol) and tributylamine (1.0 mL) was heated at 180 °C for 4 hours. After the reaction mixture was cooled to room temperature, the residue was diluted with dichloromethane,

washed with dilute hydrochloric acid and water and crystallized from diisopropyl ether to yield 116 mg of compound **6** (brown solid, 41% yield). Then phosphorus oxitrichloride (10 mL) was added to a 25 mL round-bottom flask filled with compound **6** (282 mg, 1.0 mmol), and refluxed for 4 hours. After completion of the reaction, the reaction mixture was cooled to room temperature and the solvent was distilled off. The residue was diluted with water and ether acetate. The organic phase was separated, washed with dilute sodium bicarbonate solution and brine, dried, concentrated in vacuo and purified by chromatograph on silica gel with petroleum ether and ethyl acetate as eluent to give compound **7** as a white solid. Yield: 66%, ^{1}H NMR (400 MHz, DMSO) δ 8.90 (s, 1H), 7.62-7.55 (m, 2H). HRMS-ESI: calcd for $[C_{11}H_3Cl_2F_3N+H]^+$ 318.9765, 320.9736, found: 318.9764, 320.9739; calcd for $[C_{11}H_3Cl_2F_3N+Na]^+$ 340.9585, 342.9555, found: 340.9576, 342.9565.

General procedure for the synthesis of **1-2**

Compounds **1** and **2** were prepared as described in Zhang et al[37]. Compound **7** (160 mg, 0.5 mmol), (S)-1,1,1-trifluoropropan-2-amine hydrochloride (75 mg, 0.5 mmol, for **1**), or (1R,3r,5S)-3-methoxy-8-azabicyclo[3.2.1]octane (71 mg, 0.5 mmol for **2**), and potassium carbonate (276mg, 2.0 mmol) was dissolved in DMF (5 mL) and stirred at room temperature for 4 hours. After completion of the reaction, water and ethyl acetate was added. The organic phase was separated, washed with brine, dried over anhydrous sodium sulfate, concentrated in vacuo and purified by chromatograph on silica gel with petroleum ether and ethyl acetate as eluent to give compounds **1 and 2** as white solid. Yield: 48%-63%.

(S)-5-chloro-6-(2,4,6-trifluorophenyl)-N-(1,1,1-trifluoropropan-2-yl)-[1,2,4]triazolo[1,5-a]pyrimidin-7-amine (**1**)

Yield: 48%, ^{1}H NMR (400 MHz, CDCl₃) δ 8.40 (s, 1H), 6.93-6.89 (m, 2H), 5.96 (d, J = 10.6 Hz, 1H), 4.75 (s, 1H), 1.43 (t, J = 10.0 Hz, 3H). HRMS-ESI: calcd for $[C_{14}H_8ClF_6N_5+H]^+$ 396.0451, found 396.0488; calcd for $[C_{14}H_8ClF_6N_5+Na]^+$ 418.0270, found 418.0263.

5-chloro-7-((1*R*,3*r*,5*S*)-3-methoxy-8-azabicyclo[3.2.1]octan-8-yl)-6-(2,4,6-

576 trifluorophenyl)-[1,2,4]triazolo[1,5-*a*]pyrimidine (**2**)

577   Yield: 63%, ^{1}H NMR (400 MHz, DMSO) δ 8.57 (s, 1H), 7.52-7.48 (m, 2H), 4.58

578 (s, 2H), 3.43 (t, *J* = 4.0 Hz, 1H), 3.17 (s, 3H), 2.01 (dt, *J* = 10.2, 5.1 Hz, 4H), 1.90 (d, *J*

579 = 14.6 Hz, 2H), 1.77 – 1.67 (m, 2H). HRMS-ESI: calcd for $[C_{19}H_{17}ClF_3N_5O+H]^+$

580 424.1152, found 424.1152; calcd for $[C_{19}H_{17}ClF_3N_5O+Na]^+$ 446.0971, found 446.0964.

581 *Cell culture*

582   HeLa, Hct116, H460 and SU-DHL-6 cells were all sourced from American Type

Culture Collection. H460 cells were cultured in RPMI 1640 medium and HeLa, Hct116

and SU-DHL-6 cells were cultured in Dulbecco's Modified Eagle's medium. Both

media were supplemented with 5%-10% fetal bovine serum and about 1% penicillin-

streptomycin. The culture temperature was set at 37℃, and cells were grown in a

humidified incubator with 5% CO_2. All cells have been authenticated by STR tests and

are free of mycoplasma.

Label free Quantitative Proteomics

HeLa cells were treated with or without 1μM cevipabulin for six hours and then all cells

were collected and lysed with radioimmunoprecipitation assay buffer (containing

proteinase inhibitor mixture) for 30min on ice. Then all samples were centrifuged at

10,000 g for 30 minutes to pellet cell debris. Supernatants were collected and stored at

-80℃ before analysis. We have done three biological repeats. Then the following label-

free quantitative proteomic analysis of these samples were carried out following the

procedure as described previously[38].

Immunoblotting

Cells were plated on six-well plates and cultured for 24hours before treated with

different compounds for different time. Total cells were harvested and washed by

phosphate buffer saline (PBS) before centrifuged at 1000 g for 3min. Then 1×loading

buffer (diluted from 6×loading buffer by radioimmunoprecipitation assay buffer l) was

added to the cell pellets and lysed for 10min. Samples were then incubated in boiling

water for 10 min and then stored at -20℃ before use. Equal volume of samples was

loaded to 10% SDS-PAGE for electrophoresis and then transferred to a polyvinylidene

difluoride (PVDF) membranes at 4℃ for 2 hours. Proteins on PVDF membranes were

incubated in blocking buffer (5% skim milk diluted in 1×PBST(PBS buffer with 0.1%

Tween-20)) for 1hours. Then the PVDF membranes were incubated with first

antibodies (diluted in blocking buffer) for 12hours and washed for three times with

PBST before incubated with second antibody (diluted in blocking buffer) for 45 min

and washed for three times with PBST again. At last, the PVDF membranes were

immersed in enhanced chemiluminescence reagents for 30 seconds subjected to image

with a chemiluminescence image analysis system (Tianneng, China).

Immunofluorescence

HeLa cells were grown on microscope cover glass in 24-well plates for 24 hours before

treated with different compounds for various time. Then the medium was removed and

cells were washed with prewarmed (37℃) PBS for 2 min before fixed with 50%

methanol/ 50% acetone for 3 min. The fixed cells were washed with PBS for 2 min

again before incubated with α-tubulin antibody (dilute in PBST containing 5% bovine

serum albumin) for 4 h at room temperature. Cells were then washed with PBST for

tree times (3×5 min) and followed by incubation in fluorescent second antibody (dilute

in PBST containing 5% bovine serum albumin) for 45min at room temperature. Three

times wash with PBST was performed again to remove unbounded second-antibody

before imaging using a fluorescence microscope (Zeiss, Germany)

Quantitative-PCR

HeLa and Hct116 cells were plated on six-well plates and culture for 24hours before

treated with cevipabulin for different time. Total mRNA of both HeLa and Hct116 cells

were extracted with TRIzol (Invitrogen, USA) agents following the manufacturer's

protocol and then qualified using a NanoDrop1000 spectrophotometer (Thermo Fisher

Scientific, USA. The cDNA synthesis was carried out using a high Capacity cDNA

Reverse Transcription Kit (Applied Biosystems, USA). Taq Universal SYBR Green

Supermix (BIO-RAD, USA) was employed for further Quantitative PCR analysis on a

CFX96 Real-time PCR System (BIO-RAD, USA). Relative mRNA level of both *α-*

tubulin and *β-tubulin* were normalized to that of GAPDH.

Single amino acid substitution on α-tubulin

The pIRESneo-EGFP-alpha Tubulin plasmid was obtained from Addgene (USA) and

mutation (Y224G) of α-Tubulin were performed using a Q5 Site-Directed Mutagenesis

kit (NEB #E0554S, USA). Hela cells were plated on six-well plates and incubated for

24 hours before transfected with these plasmids by Lipofectamine 2000. Then cells

were culture for another 24hours before treated with or without different compounds

for 16hours. Total protein was extracted and analyzed by immunoblotting to detect the

content of GTP-α-tubulin and GAPDH was employed as loading control.

Transmission electron microscopy

Purified porcine tubulin (2mg/ml) was diluted in PIPES buffer (80 mM PIPES, pH 6.9,

0.5 mM EGTA, 2 mM $MgCl_2$) supplemented with 1 mM GTP. Different compounds

were then incubated with tubulin at room temperature for 30 min. About 5μL of each

sample solution was added to a 230-mesh per inch, carbon films supported formvar.

Then the sample was stained with 2% (w/v) phosphotungstic acid for 60 seconds. A

Tecnai G2 F20 S-TWIN electron microscope (FEI, USA) was used for observation.

Differential Scanning Fluorimetry

Purified porcine tubulin (0.2mg/ml) was diluted in PIPES buffer supplemented with 1

mM GTP. Different compounds were added to tubulin solution and incubated for15min

at room temperature. Then capillaries were immersed into tubulin solutions to load the

samples for tests using the nanoDSF (Prometheus NT.48, NanoTemper, Germany). The

temperature range was set at 20-80°C and heating rate at 1°C/min. The fluorescence of

tryptophan fluorescence at 330nm (330F) and 350nm (350F) were detected and the

melting temperatures (Tm value) of tubulin were calculated as the maximum of the first

derivative of the F350/F330 fluorescence ratios.

TPE-MI as a thiol probe to detect unfolded protein

TPE-MI is a small molecule which is inherently non-fluorescent until covalently binds

to a thiol by its maleimide [26, 39]. This molecule could be used to monitor purified

protein unfolding *in vitro* [26]. Purified tubulin (0.2mg/ml) was diluted in PIPES buffer

supplemented with 1 mM GTP and then mixed with 50 μM TPE-MI and different

compounds. The samples were then immediately subjected to a microplate reader

(Biotek, USA) to detect the fluorescence (Excitation wavelength:350nm; Emission

wavelength: 470nm) every half minute for 60 min.

Structural Biology

Protein expression and purification were detailly described in our precious study [40]. Tubulin, RB3 and TTL (2:1.3:1.2 molar ratio) were mixed together, then 5 mM tyrosine, 10 mM DTT and 1 mM AMPPCP were added and then the mixture was concentrated to about 15 mg/ml at 4 ℃. The crystallization is conducted using a sitting-drop vapor-diffusion method under 20℃ and the crystallization buffer is optimized as: 6% PEG4000, 8% glycerol, 0.1 M MES (pH 6.7), 30 mM $CaCl_2$, and 30 mM $MgCl_2$. Seeding method was also used to obtain single crystals. Crystals appeared in about 2-days and in a rod like shape and the size reached maximum dimensions within one week. About 0.1 μL cevipabulin (diluted in DMSO with a concentration of 100 mM) was added to a drop containing tubulin crystal and incubated for 16 h at 20 ℃. The following data collection and structure determination were the same as previous description [40].

Statistical analysis

Data are presented as means. Statistical differences were determined using an unpaired Student's t test. p values are indicated in figure legend when necessary: **, $p < 0.001$; ***, $p < 0.0001$.

Acknowledgement

Project, Sichuan University (2020SCU12036); China Postdoctoral Science Foundation

Grants (2019T120855 and 2019M650248); Sichuan Science and Technology Program

Grants (2019YJ0088, 2019YFH0123 and 2019YFH0124). Post-doctoral Research

Project, West China Hospital, Sichuan University Grant 2018HXBH027.

Author contributions

J.Y. performed most of the cellular and biochemical experiments and wrote the draft.

J.Y., Y.Y., W.Y, L.N. and Q.C. performed the structural biology experiments. Y.L.

synthesized all chemical compounds. H.Y., Y.Z., Z.W., Z.Y., H.P., H.W., M.Z., J. W.

L.Y., and L.O., performed some of these biochemical experiments. W.Y., J.Y., W.L. and

L.C. conceived the idea and supervised the study. J.Y., Q.C., W.L. and L.C. revised the

manuscript.

All authors approved the final manuscript.

Conflict of interest

The authors declare no competing financial interests.

Data availability.

Atomic coordinates and structure factors of tubulin complexed with cevipabulin have

been deposited in the Protein Data Bank under accession code 7CLD. Further

information and requests for resources and reagents should be directed to and will be

fulfilled by Jianhong Yang (yjh1988@scu.edu.cn).

Reference

[1] Arnst KE, Banerjee S, Chen H, Deng S, Hwang DJ, Li W, et al. Current advances of tubulin inhibitors as dual acting small molecules for cancer therapy. Medicinal research reviews. 2019;39:1398-426.
[2] Haider K, Rahaman S, Yar MS, Kamal A. Tubulin inhibitors as novel anticancer agents: an

overview on patents (2013-2018). Expert opinion on therapeutic patents. 2019;29:623-41.

[3] Gigant B, Wang C, Ravelli RB, Roussi F, Steinmetz MO, Curmi PA, et al. Structural basis for the

regulation of tubulin by vinblastine. Nature. 2005;435:519-22.

[4] Doodhi H, Prota AE, Rodríguez-García R, Xiao H, Custar DW, Bargsten K, et al. Termination of

protofilament elongation by eribulin induces lattice defects that promote microtubule

catastrophes. Current Biology. 2016;26:1713-21.

[5] Yang J, Wang Y, Wang T, Jiang J, Botting CH, Liu H, et al. Pironetin reacts covalently with

cysteine-316 of α-tubulin to destabilize microtubule. Nature communications. 2016;7:1-9.

[6] Kavallaris M. Microtubules and resistance to tubulin-binding agents. Nature Reviews Cancer.

2010;10:194-204.

[7] Prota AE, Setter J, Waight AB, Bargsten K, Murga J, Diaz JF, et al. Pironetin binds covalently to

αCys316 and perturbs a major loop and helix of α-tubulin to inhibit microtubule formation. Journal

of Molecular Biology. 2016;428:2981-8.

[8] Manka SW, Moores CA. The role of tubulin–tubulin lattice contacts in the mechanism of

microtubule dynamic instability. Nature structural & molecular biology. 2018;25:607-15.

[9] Prota AE, Bargsten K, Northcote PT, Marsh M, Altmann KH, Miller JH, et al. Structural basis of

microtubule stabilization by laulimalide and peloruside A. Angewandte Chemie International

Edition. 2014;53:1621-5.

[10] Prota AE, Bargsten K, Diaz JF, Marsh M, Cuevas C, Liniger M, et al. A new tubulin-binding site

and pharmacophore for microtubule-destabilizing anticancer drugs. Proceedings of the National

Academy of Sciences. 2014;111:13817-21.

[11] Prota AE, Bargsten K, Zurwerra D, Field JJ, Díaz JF, Altmann K-H, et al. Molecular mechanism

of action of microtubule-stabilizing anticancer agents. Science. 2013;339:587-90.

[12] Ravelli RB, Gigant B, Curmi PA, Jourdain I, Lachkar S, Sobel A, et al. Insight into tubulin

regulation from a complex with colchicine and a stathmin-like domain. Nature. 2004;428:198-202.

[13] Steinmetz MO, Prota AE. Microtubule-targeting agents: strategies to hijack the cytoskeleton.

Trends in cell biology. 2018;28:776-92.

[14] Dorléans A, Gigant B, Ravelli RB, Mailliet P, Mikol V, Knossow M. Variations in the colchicine-

binding domain provide insight into the structural switch of tubulin. Proceedings of the National

Academy of Sciences. 2009;106:13775-9.

[15] Ayral-Kaloustian S, Zhang N, Beyer C. Cevipabulin (TTI-237): preclinical and clinical results for

a novel antimicrotubule agent. Methods and findings in experimental and clinical pharmacology.

2009;31:443.

[16] Beyer CF, Zhang N, Hernandez R, Vitale D, Lucas J, Nguyen T, et al. TTI-237: a novel

microtubule-active compound with in vivo antitumor activity. Cancer research. 2008;68:2292-300.

[17] Kovalevich J, Cornec A-S, Yao Y, James M, Crowe A, Lee VM-Y, et al. Characterization of brain-

penetrant pyrimidine-containing molecules with differential microtubule-stabilizing activities

developed as potential therapeutic agents for Alzheimer's disease and related tauopathies. Journal

of Pharmacology and Experimental Therapeutics. 2016;357:432-50.

[18] Sáez-Calvo G, Sharma A, de Asís Balaguer F, Barasoain I, Rodríguez-Salarichs J, Olieric N, et

al. Triazolopyrimidines are microtubule-stabilizing agents that bind the Vinca inhibitor site of

tubulin. Cell chemical biology. 2017;24:737-50. e6.

[19] Nasrin SR, Kabir AMR, Konagaya A, Ishihara T, Sada K, Kakugo A. Stabilization of microtubules

by cevipabulin. Biochemical and biophysical research communications. 2019;516:760-4.

[20] Nasrin SR, Ishihara T, Kabir AMR, Konagaya A, Sada K, Kakugo A. Comparison of microtubules
stabilized with the anticancer drugs cevipabulin and paclitaxel. Polymer Journal. 2020:1-8.
[21] Beyer CF, Zhang N, Hernandez R, Vitale D, Nguyen T, Ayral-Kaloustian S, et al. The
microtubule-active antitumor compound TTI-237 has both paclitaxel-like and vincristine-like
properties. Cancer chemotherapy and pharmacology. 2009;64:681-9.
[22] Stanton RA, Gernert KM, Nettles JH, Aneja R. Drugs that target dynamic microtubules: a new
molecular perspective. Medicinal research reviews. 2011;31:443-81.
[23] Menéndez M, Rivas G, Díaz JF, Andreu JM. Control of the structural stability of the tubulin
dimer by one high affinity bound magnesium ion at nucleotide N-site. Journal of Biological
Chemistry. 1998;273:167-76.
[24] Spiegelman BM, Penningroth SM, Kirschner MW. Turnover of tubulin and the N site GTP in
Chinese hamster ovary cells. Cell. 1977;12:587-600.
[25] Keays DA, Tian G, Poirier K, Huang G-J, Siebold C, Cleak J, et al. Mutations in α-tubulin cause
abnormal neuronal migration in mice and lissencephaly in humans. Cell. 2007;128:45-57.
[26] Chen MZ, Moily NS, Bridgford JL, Wood RJ, Radwan M, Smith TA, et al. A thiol probe for
measuring unfolded protein load and proteostasis in cells. Nature communications. 2017;8:1-11.
[27] Zhang B, Maiti A, Shively S, Lakhani F, McDonald-Jones G, Bruce J, et al. Microtubule-binding
drugs offset tau sequestration by stabilizing microtubules and reversing fast axonal transport
deficits in a tauopathy model. Proceedings of the National Academy of Sciences. 2005;102:227-
31.
[28] Zhang F, Su B, Wang C, Siedlak SL, Mondragon-Rodriguez S, Lee H-g, et al. Posttranslational
modifications of α-tubulin in alzheimer disease. Translational neurodegeneration. 2015;4:1-9.
[29] Zhang B, Carroll J, Trojanowski JQ, Yao Y, Iba M, Potuzak JS, et al. The microtubule-stabilizing
agent, epothilone D, reduces axonal dysfunction, neurotoxicity, cognitive deficits, and Alzheimer-
like pathology in an interventional study with aged tau transgenic mice. Journal of Neuroscience.
2012;32:3601-11.
[30] Ballatore C, Brunden KR, Trojanowski JQ, Lee VM-Y, Smith III AB. Non-naturally occurring
small molecule microtubule-stabilizing agents: A potential tactic for CNS-directed therapies. ACS
Publications; 2017.
[31] Lou K, Yao Y, Hoye AT, James MJ, Cornec A-S, Hyde E, et al. Brain-penetrant, orally
bioavailable microtubule-stabilizing small molecules are potential candidate therapeutics for
Alzheimer's disease and related tauopathies. Journal of medicinal chemistry. 2014;57:6116-27.
[32] Brunden KR, Trojanowski JQ, Smith III AB, Lee VM-Y, Ballatore C. Microtubule-stabilizing
agents as potential therapeutics for neurodegenerative disease. Bioorganic & medicinal chemistry.
2014;22:5040-9.
[33] Rice LM, Montabana EA, Agard DA. The lattice as allosteric effector: structural studies of $\alpha\beta$-
and γ-tubulin clarify the role of GTP in microtubule assembly. Proceedings of the National
Academy of Sciences. 2008;105:5378-83.
[34] Nogales E, Wang HW. Structural mechanisms underlying nucleotide-dependent self-
assembly of tubulin and its relatives. Current Opinion in Structural Biology. 2006;16:221-9.
[35] Wang H-W, Nogales E. Nucleotide-dependent bending flexibility of tubulin regulates
microtubule assembly. Nature. 2005;435:911-5.
[36] Buey RM, Díaz JF, Andreu JM. The nucleotide switch of tubulin and microtubule assembly: a
polymerization-driven structural change. Biochemistry. 2006;45:5933-8.

[37] Zhang N, Ayral-Kaloustian S, Nguyen T, Afragola J, Hernandez R, Lucas J, et al. Synthesis and
SAR of [1, 2, 4] triazolo [1, 5-a] pyrimidines, a class of anticancer agents with a unique mechanism
of tubulin inhibition. Journal of medicinal chemistry. 2007;50:319-27.
[38] Yang J, Li Y, Yan W, Li W, Qiu Q, Ye H, et al. Covalent modification of Cys-239 in β-tubulin by
small molecules as a strategy to promote tubulin heterodimer degradation. Journal of Biological
Chemistry. 2019;294:8161-70.
[39] Liu Y, Yu Y, Lam JW, Hong Y, Faisal M, Yuan WZ, et al. Simple Biosensor with High Selectivity
and Sensitivity: Thiol-Specific Biomolecular Probing and Intracellular Imaging by AIE Fluorogen on
a TLC Plate through a Thiol–Ene Click Mechanism. Chemistry–A European Journal. 2010;16:8433-
8.
[40] Niu L, Yang J, Yan W, Yu Y, Zheng Y, Ye H, et al. Reversible binding of the anticancer drug
KXO1 (tirbanibulin) to the colchicine-binding site of β-tubulin explains KXO1's low clinical toxicity.
Journal of Biological Chemistry. 2019;294:18099-108.

814
Figure S1: Cevipabulin promotes proteasome-dependent degradation of α-and β-tubulin. (**a**)
HeLa cells were treated with 1 μM cevipabulin for the indicated times and then the α and β-
tubulin levels were detected by immunoblotting. Results are representative of two independent
experiments. (**b**) HeLa and Hct116 cells were treated with indicated concentrations of cevipabulin
for 16 hours, and then mRNA levels of both *α-tubulin* and *β-tubulin* were measured by
quantitative-PCR. Data were shown as means ± SD of three independent experiments. (**c**) Cells
were treated with or without MG132 (20 μM) for one hour before treated with different
concentrations of cevipabulin for 16 hours. Protein levels of both α- and β-tubulin were detected
by immunoblotting. Results are representative of two independent experiments. Cev: cevipabulin.
824
825

Figure S2. Compound 1 binds to vinblastine site to induce formation of irregular tubulin aggregates and compound 2 binds to the seventh site to induce tubulin degradation. (**a**) HeLa cells were treated with or without 10 μM vinblastine for 1 hour before treated with 10 μM compound 1 for 16 h and then the protein level of α-tubulin was detected by immunoblotting. Results are representative of three independent experiments. (**b**) Vectors expressing either wild type or Y224G mutant GFP-tubulin were transfected to HeLa cells. After 24 hours, cells were treated with or without 10 μM compound **1** for 16 h. Then the protein level of GFP-α-tubulin was detected by immunoblotting. Results are representative of three independent experiments. (**c**) HeLa cells were treated with or without 10 μM vinblastine for 1hour before treated with 10 μM compound **2** for another hour, and irregular tubulin aggregates were detected using immunofluorescence and the number of irregular tubulin aggregates were counted for randomly chosen 30 cells. ***p<0.0001. Results are representative of three independent experiments. (**d**) Vectors expressing either wild type or Y224G mutant GFP-α-tubulin were transfected to HeLa cells. After 24 hours, cells were treated with or without 10μM compound **2** for 1 h. Irregular tubulin aggregates were detected using immunofluorescence and the number of irregular tubulin aggregates were counted for randomly chosen 30 cells. ns: no significant difference. Results are representative of three independent experiments. Vin: vinblastine. **1**: compound **1**；**2**: compound **2**.

Figure S3. **Structural mechanism of cevipabulin and compound 2 induced tubulin**

protofilaments polymerization. (**a**) TEM analysis of 50 μM paclitaxel treated purified tubulin.

(**b**) Overview of the aligned structures of tubulin-cevipabulin complex (dark) and tubulin-

compound **2** complex (salmon) (PDB code: 5NJH). (**c**) Close-up view of the cevipabulin and

compound **2** binding to inter-dimer interface in the aligned complexes in (**b**). (**d**) Close-up view of

the inter-dimer interface of the aligned structures of apo tubulin (green) and tubulin-compound **2**

complex (dark), which are superimposed on β1-tubulin subunit. Compound **2** is shown as yellow

stick. (**e**) Overview of the aligned structures in (**d**). (**f**) Close-up view of the inter-dimer interface

of the aligned structures of tubulin-compound **2** complex (salmon) and the polymerized

microtubule (violetpurple, PDB code: 6DPV), which are superimposed on β1-tubulin subunit.

Compound **2** is shown as yellow stick. (**g**) Two tubulin-compound **2** complexes aligned to two

adjacent protofilaments in polymerized microtubule structure on β1-tubulin respectively to

analyze the lateral interaction. M-loop in tubulin-compound **2** complex and the polymerized

microtubule structure are colored in red and purple respectively. (**h**) Tubulin-compound **2**

(salmon), tubulin-vinblastine (cyan, PDB code: 5J2T) and apo tubulin (green) were aligned on α2-

tubulin and the close-up view of T5 loops of α2-tubulin were shown. Cev: cevipabulin. cpd2:

compound **2**.

Table S1. Data collection and refinement statistics.

	Tubulin-cevipabulin
Data collection	
Space group	$P2_12_12_1$
Cell dimensions	
a, b, c (Å)	104.4 160.8 174.8
α, β, γ (°)	90.0 90.0 90.0
Resolution (Å)	50.0-2.60 (2.64-2.60) *
R_{pim}	3.1 (42.2)
$I/\sigma I$	23.6 (2.0)
Completeness (%)	100 (100)
Redundancy	13.4 (13)
Refinement	
Resolution (Å)	50.0-2.61
No. reflections	83938
R_{work}/ R_{free}	20.7/25.8
No. atoms	
Protein	17464
Ligand/ion	241
Water	294
B-factors	
Protein	44
Ligand/ion	56
Water	54
R.m.s deviations	
Bond lengths (Å)	0.008
Bond angles (°)	0.789

    *Highest resolution shell is shown in parenthesis.

a 1825 quantified proteins

β4B-Tub
α1B-Tub

β-Tub
α4A-Tub

α1C-Tub

p-Value

Fold Change

b

Cev	0	0.1	1	10		0	0.1	1	10		0	0.1	1	10		0	0.1	1	10 μM

Hela   Hct116   H460   SU-DHL-6

α-Tub

β-Tub

GAPDH

a

Trifluoropropanyl

Side Chain

Triazolopyrimidinyl

Difluorphenyl

b

TTL

CEV   CEV

β1-Tub   α2-Tub   β2-Tub

α1-Tub

RB3

c   **d**

e

α2H10

β1H6   Cev

β1T5

βY224

β1H7   α2H1

β1H1   GDP

h

β2H10

α2H6   Cev

α2T5

α2H7   αY224

α2H1   GTP   β2T7

f   β1-Tub

β1H7   β1H6   βD211

β1H6   βT223

βY224

βN206   Cev

GDP   β1T5

g   α2-Tub

α2H10   αK326   α2T7

αN329

Cev

i   α2-Tub

α2H7   αT223   αR221

α2H1   αY224   α2H6

αD211

GTP   αN206   Cev

α2T5

a

Ra

0.90
0.85
0.80
0.75

— Con Tm=61.1 ℃
— Vin Tm=61.5 ℃
— Cev Tm=61.3 ℃
— 1   Tm=56.9 ℃
— 2   Tm=63.2 ℃

First Derivae

20   30   40   50   60   70   80
Temperature (℃)

b

Con

50nm

c

RFU(AU)

— DMSO
— 4M GuHCl
— Vin 5μM
— 1 2 μM
— 1 5 μM
— 2 5μM
— Cev 2μM

5 10 15 20 25 30 35 40 45 50 55 60
Time(min)

d

PRY-41   -   -   +   +
1   -   +   -   +

α-Tub

GAPDH

a

b

β2-Tub α2-Tub β1-Tub α1-Tub

RB3

c

βH6 αH9 Cev αH10
βH7 Cpd2
βT5
GTP αT7
βH1 αH4

d

αH9 αH10
6.7Å Cpd7
α2-Tub β1-Tub βH6

e

β2-Tub α2-Tub β1-Tub α1-Tub

RB3

f

αH10
cpd2
α2-Tub β1-Tub βH6

g

H2 S3
H2-S3
5.2Å M-loop
α-Tub S2 α'-Tub

h

αT5