S494 O-glycosylation site on the SARS-COV-2 RBD Affects the Virus Affinity to ACE2 and its Infectivity; A Molecular Dynamics Study[†]

Shadi Rahnama,[‡] Maryam Azimzadeh Irani,[*,¶] Mehriar Amininasab,[§] and Mohammad Reza Ejtehadi[*,∥]

[‡]Institute for Nanoscience and Nanotechnology, Sharif University of Technology, Tehran 14588, Iran.

[¶]Faculty of Life Sciences and Biotechnology, Shahid Beheshti University, Tehran 1983963113, Iran.

[§]Department of Cell and Molecular Biology, School of Biology, College of Science, University of Tehran, Tehran, Iran.

[∥]Department of Physics, Sharif University of Technology, Tehran 14588, Iran.

E-mail: m_azimzadeh@sbu.ac.ir; ejtehadi@sharif.edu

Phone: +98 (0)21 29901; +98 (0)21 6616 4525. Fax: +98 (0)21 22431919; +98 (0)21 6602 2711

Abstract

SARS-COV-2 is a strain of Coronavirus family which caused the extensive pandemic of COVID-19, which is still going on. Several studies showed that the glycosylation of virus spike (S) protein and the Angiotensin-Converting Enzyme 2 (ACE2) receptor on the host

---

[†]O-glycosylation of the SARS-COV-2 RBD

cell is critical for the virus infectivity. Molecular Dynamics (MD) simulations were used to explore the role of a novel mutated O-glycosylation site (D494S) on the Receptor Binding Domain (RBD) of S protein. This site was suggested as a key mediator of virus-host interaction. We showed that the decoration of S494 with core and elongated O-glycans results in stabilized interactions on the direct RBD-ACE2 interface with more favorable binding free energies for longer oligosaccharides. Hence, the further drug design attempts should take this crucial factor into account, while suggesting any novel therapeutic candidate.

Introduction

Severe Acute Respiratory Syndrome Corona Virus 2 (SARS-COV-2) is a positive-sense single-stranded RNA virus which caused the pandemic of COVID-2019 that is still going on. In Wuhan, China, the virus is believed to transmit from bat to human[1] and underwent several inter and intra species passages. This large-scaled pandemic motivated several scientific groups to control the virus's spread and treat the infected patients by all means. The researchers are mainly focused on the phylogeny and origin of SARS-COV-2,[1] Structural assembly and the dynamics of virion-host cell proteins[2–4] and the experimental efforts to obtain a SARS-COV-2 neutralizing antibody.[5,6]

The SARS-COV-2 virion assembly and its infusion into the human cell are similar to the one known for SARS-COV.[2] The fusion protein known as the spike (S) protein interacts with ACE2 on the host cell via its Receptor Binding Domain (RBD)[3,7] (Figure 1).The RBD-ACE2 interaction takes place via four β strands on RBD ($\beta4$-$\beta7$) and two α helices on ACE2 binding interface (H1,H2)[3](Figure 1). Electron microscopy and X-ray crystal structures elucidated the active form of SARS-COV-2 S protein which assembles as a trimer,[2] and one RBD of one S protein complex ACE2,[3] respectively. The structures of S protein in the apo and ACE2 bound complexes show that RBD adopts a 3D fold similar to the RBD of SARS-COV;.[2] However, six novel mutations on SARS-COV-2 RBD could lead to different binding

affinity to ACE2.[1] Speculations regarding SARS-COV-2-ACE2 binding affinity suggested two different scenarios. One reported a dramatic increase (up to 20-fold) in the binding affinity of SARS-COV-2 to ACE2 compared to SARS-COV.[2] While, the other suggests a similar binding affinity for SARS-COV-2 and SARS-COV to ACE2.[3] Either way, the mutations in SARS-COV2 RBD certainly results in attenuated binding affinity of all designed inhibitors which successfully work on SARS-COV.[2] It was a challenging issue for anti-SARS-COV-2 drug design attempts since the pandemic has started.

Figure 1: The extracellular domain of human ACE2 and RBD of SARS-COV-2 S protein is shown with gray and purple, respectively (A and B). The RBD attached O-glycan is shown with green spheres (C). N-glycans in the proximity of RBD-ACE2 interface are shown with pink spheres (D). Two O-glycan interaction sites H1 and H2, are shown with orange (E).

In addition to the 3D structures of RBD-ACE2 complex, several recent Molecular Dynamics (MD) studies addressed the complex dynamics to design peptide inhibitors. That block the ACE2-RBD interaction and evaluate the currently targeted binding epitopes' accessibility for further improvements[8] .[4,9] Earlier MD studies explored the effect of pH[10] and temperature[11] on the dynamics of SARS-COV-ACE2 complex. To the best of our knowledge, there is currently no approved therapeutic inhibitor for SARS-COV-2.

A pioneering study showed that the observed mutations at the junction of S1 and S2 subdomains of the SARS-COV-2 S protein result in the emergence of a polybasic cleavage site adjacent to three O-glycosylation sites which are novel to SARS-COV-2 and were not observed in any other related virion.[1] It was proposed that O-glycosylation (addition of glycan building blocks to hydroxyl oxygen of Ser/Thr residues which is an enzymatic post transnational modification[12]) could lead to recognition of polybasic cleavage site by Furin enzyme and thus resulted in higher infectivity and broadens the host range of the virus.[1] Recent comprehensive experimental studies reported a couple of O-glycosylated sites on the SARS-COV-2 S protein as well.[13]The structure and dynamics of these O-glycans were also explored in a computational work.[14]

Six mutations have been reported on the RBD of SARS-COV-2 compared to the SARS-COV. One of these six occurring mutations is D494S on the RBD of S protein. Based on the proposed mechanism for the O-glycosylation of Furin cleavage site and the Evidences mentioned above,[1,13,14] we propose that Serine494 of SARS-COV-2 RBD could also become O-glycosylated even though it is not experimentally confirmed (Figure 1).

Our comprehensive review of literature on pathology and glycosylation pattern of the envelope proteins of all other coronaviruses and their close relatives showed strong evidence for the critical role of O-glycosylation in regulation and immune evasion of the viruses.[1,15–17] Also, former studies showed that the O-glycosylation of the membrane (M) protein of Mouse. Hepatitis Virus (MHV)15 triggers a lower interferon level than the N-glycosylated M protein.[15] 3A protein of the MHV, an accessory protein to the M protein is also known to

be O-glycosylated. Thus, it could be shielded from Peptide-N Glycosidase F (PNGase F) which cleaves most monosaccharide attached to protein. On the other hand, a complex non-template mechanism of the O- glycosylation could occur via several Polypeptide N-acetyl Galactoseamine Transferases (PPGalNAcTs) specific to various tissues was shown to be a common mechanism for the pathology of several high-evasion viruses such as Influenza.[18] In fact, the experimental findings suggest that the mucin proteins of human respiratory airways are heavily O- glycosylated with carbohydrate chains.[19–21] These dense O-glycan chains are known to trap invading microorganisms such as various strains of bacteria.[22]

Putting all the information mentioned above together, although O-glycosylation was not reported in the direct RBD-ACE2 interface of other coronaviruses, it is highly plausible that D494S mutation has an evolutionary origin and leads to O-glycosylation of RBD in the SARS-COV-2. Hence, the O- glycosylation could be the overlooked factor for the SARS-COV-2 increased binding affinity to ACE2, explaining the virus's high infectivity in the human host and exposing people with a higher level of blood sugar at higher risk.[23] Our observations may explain the reason for the higher vulnerability of diabetic patients to infection.

Herein, we explored this possibility by modeling the O-glycosylated RBD structure and dynamics in interaction with ACE2 (Figure 2). We show that O-glycosylation does indeed increase SARS-COV-2 RBD-ACE2 binding affinity. Also, we show that by increasing the O-glycan chain's length, the binding affinity will increase. Furthermore, we also considered the structural role of N-glycosylation (addition of glycan building blocks to the nitrogen atom of Asn located at the Asn-X-Ser/Thr motif, enzymatic post-translationalmodification) of both RBD and ACE2 in all of the simulations.

In summary, this work provides strong evidence that the O-glycosylation of SARS-COV2 RBD in the respiratory airways leads to stronger interactions between the virion and the host cell receptor. Hence, further drug design attempts should take this crucial factor into account, while suggesting any novel therapeutic candidate.

Methods

O- and N- glycosylated models of RBD-ACE2 complexes and Molecular Dynamics simulations

The crystal structure of SARS-COV-2 RBD in complex with the extracellular domain of ACE2 (PDB ID: 6M0J) was used as the starting structure. Zn^{2+} and Cl^- ions which were resolved in the crystal structure for their crucial role in stabilizing ACE2 structure were also preserved at S1 and S2 subunits of ACE2. All Aspargines located within N-glycosylation motifs from RBD (Asn343) and ACE2 (Asn53, Asn90, Asn103, Asn322, Asn432, and Asn546) were glycosylated. The GLYCAM online builder[24] was used to attach the oligosaccharide glycans to each site. This oligosaccharide model was selected as a primary model of glycosylation which occurs in normal human cells.[25] Figure 2 provides details on the structure of attached oligosaccharides. H++ algorithm[26] was used to estimate the pKa of ionizable residues of complex at physiological pH. Calculated pKa was used to assign the protonation states to residues, H34, H195, H345, H374, H378, H417, and H493 of ACE2 that were protonated and gain a positive charge. O-glycosylation of Serine494 of RBD was carried out by attaching three models of core and branched typical human O- glycans using GLYCAM online builder.[24] Models I, II, and III are consist of two, five, and six monosaccharide units, respectively (Figure 2). These O-glycan models were attached to RBD, while RBD-ACE2 complex was kept fully N-glycosylated (Figure 1). The RBD-ACE2 model without attached O-glycan will be referred to as model 0 in the rest of the text and figures. All systems were solvated using TIP3P water model[27] and were neutralized by attaining a buffered environment in 150 mM NaCl. The resulting water box approximately has a dimension of $121 \times 111 \times 149$ Å^3 and are typically consist of 190000 atoms each. All systems were parameterized with the CHARMM36m force field,[28,29] and NAMD2.12[30] was used to perform the MD simulations. Periodic boundary conditions under NPT ensemble was applied, while employing Langevin thermostat and Nose–Hoover Langevin piston method to keep systems'

temperature and pressure at 310 K and 1 bar, respectively. A cutoff of Å was assigned for computation of short range nonbonded van der Waals interactions. The particle mesh Ewald method[31] was used for long-range electrostatic interactions. R-RESPA multiple time-step schemes were used for the integration of motion equations.[30] Using this integrator, the Lennard-Jones interactions and bonded ones were updated every step and electrostatic interactions every two steps. The time steps for integration were set to 2fs. Final models of all complexes were minimized for 5000 steps to remove any steric clashes. Following minimization, the systems firstly equilibrated in NVT ensemble carried out for 0.5 ns. Subsequently, system equilibration was continued in NPT ensemble for an additional 0.5 ns. During both steps, the position of atoms in the complex (proteins, Zn^{+2} and Cl^{-1} ions) was restrained using a harmonic potential with spring constant k = 1 kcal/(molÅ^2). Production runs were then performed in three replicates of each system for 50 ns. During the production runs, all restraints on the protein were turned off, while dummy bonds between Zn^{+2} and Cl^{-1} ions and their adjunct atoms within radius of 3 Å were kept intact.

Binding free energy calculations

The binding free energy for all sets of ligand-receptor complexes was calculated with Molecular Mechanics Poison Boltzmann Surface Area (MMPBSA) using CaFE pipeline tool[32] on 100 frames from the last 10 ns of MD simulations and an ensemble-averaged overall replica of each model. Binding free energy can be decomposed into the contribution of different interactions and expressed as:[33]

$$\Delta G_{bind} = \Delta G_{gas} + \Delta G_{sol} \tag{1}$$

$$\Delta G_{gas} = \langle \Delta G_{mm} \rangle - T\Delta S \tag{2}$$

$$\Delta G_{sol} = \Delta G_{PB} + \Delta G_{SA} \tag{3}$$

Where $\langle \Delta G_{mm} \rangle$, -T$\Delta$S, ΔG_{PB} and ΔG_{SA} are the ensemble-averaged protein-ligand interactions, entropy contributions, electrostatic, and nonpolar solvation energies, respectively. The entropic part of the binding free energy was not taken into account as we only tried to show relative binding free energy of homologous ligands.[33,34] For electrostatic energy calculations, APBS method[35] was used. The dielectric constants for solvent and protein were set to 80.0 and 1.5 respectively, where the latter was selected due to the proposed fitting curve by Tingjun Hou et al.[36] Probe radius for calculation of SASA was set to 1.4 Å.

MD simulations analyses

Root Mean Squared Deviation (RMSDs), Root Mean Squared Fluctuation (RMSFs), and the distances between RBD and ACE2, H1 and H2 were calculated by tcl scripts taking advantage of VMD.[37] RMSD of each system was calculated by considering the backbone atoms (C_α, C, and N). RMSF and the distance among the geometric centers of domains were calculated for C_α atoms. All trajectories were fitted to the starting conformation to present the conformational changes during simulation time. All the plots were generated by Matplotlib[38] library of Python[39] and the figures by VMD[37] and PyMOL.[40]

Results and Discussion

Dynamics of the O- and N- glycosylated RBD-ACE2

It is worth mentioning again that as all the systems are fully N-glycosylated here, the N-glycans' global effect is explicit in the dynamics. However, the interpretation of results is mainly focused on O-glycosylation. RMSD plots of RBD-ACE2 complex simulations show that among three O-glycosylated systems, model III is the least flexible one, while models II and I come just after in order (Figure 3 A and B). Interestingly, Model 0 falls between models II and I (Figure 3). Thus, as it is expected, the length of attached O-glycan is

Figure 2: (A) The overlaid presentation of O-glycosylated RBD-ACE2 complex is shown with red sticks. The extracellular domain of human ACE2 and RBD of SARS-COV-2 S protein is shown with gray and purple, respectively. N-glycans in the proximity of RBD-ACE2 interface are shown with green sticks. O-glycan models I (B), II (C) and III (D) are shown with atom name sticks.

important for RBD-ACE2 interactions. The longer glycans lead to a more stabilized RBD-ACE2 interactions. It could play a significant role in the high evasion of SARS-COV-2.

RMSF plots are also consistent to these observations (Figures 4,5) and show that which show the most dramatic increase of fluctuations occurs in residues 430 to 450 of RBD in model I. While, models II and III and the 0 present similar fluctuation patterns (Figure 4). Two residues (GLY446 and Tyr449) within this region are known to interact with the ACE2 directly.[41] These results are interesting as we know which due to the experimental findings, the O-glycans of respiratory airways are often elongated oligosaccharides.[22]

Figure 3: (A) Average backbone RMSD plots for RBD-ACE2, calculated from all replicates of each system and are shown for model 0 (black), model I (lime), model II (orange) and model III (blue). (B) Probability Density Function (PDF) of RMSD sampled over the last 30 ns of the simulations (dashed line).

Figure 4: RMSF per residue of RBD calculated from all replicates of each system and are shown for model 0 (black), model I (lime), model II (orange) and model III (blue).

Detailed interactions

Detailed investigations show that decreased flexibility is more pronounced for RBD (Figure 6), while ACE2 average RMSD plots present a noticeable overlap among all systems (Figure

Figure 5: (A) RMSF per residue of ACE2 calculated from all replicates of each system and are shown for model 0 (black), model I (lime), model II (orange) and model III (blue). The highlighted regions indicates H1 and H2. (B) The relative difference in RMSD, D_{RMSF} is shown per residue for all the models (with the same color codes).

7.A-C).

Figure 6: (A) Average backbone RMSD plots for Receptor binding RBD calculated from all replicates of each system and are shown for model 0 (black), model I (lime), model II (orange) and model III (blue). (B) Probability Density Function (PDF) of RMSD sampled over the last 30 ns of the simulations (dashed line).

Figure 7: (A) Average backbone RMSD plots for ACE2 calculated from all replicates of each system and are shown for model 0 (black), model I (lime), model II (orange) and model III (blue). (A') Probability Density Function (PDF) of RMSD sampled over the last 30ns of the simulations (dashed line). (B) Average backbone RMSD plots for α-Helix1 calculated from all replicates of each system and are shown for model 0 (black), model I (lime), model II (orange) and model III (blue). (B') Probability Density Function (PDF) of RMSD sampled over the last 30ns of the simulations (dashed line).(C) Average backbone RMSD plots for α-Helix2 calculated from all replicates of each system and are shown for model 0 (black), model I (lime), model II (orange) and model III (blue). (C') Probability Density Function (PDF) of RMSD sampled over the last 30ns of the simulations (dashed line).

ACE2 extracellular domain is a sizable receptor with about 600 amino acids; hence, it is expected that the overall RMSD of receptor would not change by the O-glycosylation of

the RBD. As we aimed to study the alterations in ACE2-RBD interface, we calculated the RMSD of two α-Helices (Figures 7.B and7.C) which are known as RBD interacting sites within ACE2.[3] Average RMSD plots of H1 and H2 show that the first helix is the most affected by the O-glycan (Figure 7.B and 7.C). H1 is more flexible in all O-glycosylated models (Figure 7.B). H2 is less flexible in all the systems and is more persistent (Figure 7.C).

ACE2 RMSF plots support these observations, showing decreased fluctuations of H2 in the simulations (Figure 5.A). To highlight the differences, we have shown the relative differences compared to model 0 in Figure 5.B. The relative difference is presented as $D_{\mathrm{RMSF}} = \frac{RMSF_0 - RMSF_i}{RMSF_0 + RMSF_i}$ where $RMSF_i$ is per residue RMSF of each model. According to the definition more positive D_{RMSF} means lower relative RMSF.

The visualization of dynamics shows that O-glycan forms the most numerous polar contacts with H2 (Figure 8) In contrast, according to the crystal structures, H1 forms more contacts with RBD than H2.[3] One should note that the RBD is not O-glycosylated in the Xray crystal structures. In fact that we see more interactions between O-glycan and H2 in our simulations suggests that the O-glycan keeps the RBD-ACE2 together by not interfering in the main ligand-receptor interactions takes place at H1.

Biding affinity of O-glycosylated RBD to ACE2

The decreased fluctuations of O-glycosylated models II and III, does not necessarily prove stronger RBD-ACE2 interaction. However, calculating the distance between RBD-ACE2, RBD-H1 and RBD-H2 could provide hints on this interaction. Distributing RBD-ACE2 distance values shows a similar pattern (peaks at 49.5 Å) in models I and III simulations (Figure 9.A). While, in model 0 simulations, RBD and ACE2 are by less than 1 Å closer (Figure 9.A). The distance fluctuation between RBD and ACE2 in model II is more altering.

In addition to the ACE2-RBD, distances between RBD, H1, and H2 of ACE2 were also measured (Figures 9.B and 9.C). Plots show that O-glycosylation always leads to a

Figure 8: The extracellular domain of human ACE2 and RBD of the SARS-COV-2 S protein are shown with gray and purple respectively. The RBD attached O-glycan is shown with atom name sticks. N-glycans in the proximity of RBD-ACE2 interface are shown with orange spheres. Persistent polar interactions between H1 and H2 of ACE2 and the attached O-glycan are shown in A-C.

decrease between the RBD and the two helices (Figures 9.B and 9.C). This decrease is more significant in H2-RBD distance for all O-glycosylated models (Figure 9.C). While, H1-RBD distance distribution only presents a decrease in the most elongated attached O-glycan model (Figure 9.B). These observations suggest that RBD-ACE2 binding should be stronger at the direct interface upon O-glycosylation with oligosaccharides.

The binding free energy between the RBD and ACE2 also shows that the interaction is more favorable upon O-glycosylation in all the systems (Figure 10.A).

The binding free energies between RBD-ACE2, calculated with the MMPBSA method, show a monotonic decrease in ΔG by increasing the size of O-glycans (Figure 10.A) and staring from -159±6.34 for model 0 to -173±6.46, -177±5.93 and -187±3.50 for models I, II and III, respectively (Figure 10.B and 10.C).

Chosen value of solute dielectric constant has shown to have a dramatic effect on the

Figure 9: (A) PDF of RBD to ACE2 averaged distance distribution with a histogram and maximum likelihood gaussian distribution fit: calculated from all replicate of each system and are represented in black, lime, orange and blue for models 0-III respectively. (B) PDF of RBD to H1 averaged distance distribution with a histogram and maximum likelihood gaussian distribution fit: calculated from all simulations of each system and represented in black, lime, orange and blue for models 0-III respectively. (C) PDF of RBD to H2 averaged distance distribution with a histogram and maximum likelihood gaussian distribution fit: calculated from all simulations of each system and represented in black, lime, orange and blue for models 0-III respectively. Sampling were done from the last 30 ns of simulations.

prediction of binding free energy, while using the MMPBSA method[36,42,43] (see method section 2.2).

Regardless of the given value to the dielectric constant, the decrease in the O-glycosylated models binding free energy is always persistent and statistically significant (Figure 10.A).

Comparing the free energy (Equation 2, 3), non-bonded electrostatics and Van Der Waals interactions are always increasing by the size of glycans (Figure 10.B and 10.C).

Conclusions

S494 that only occurs in SARS-COV-2 and not in SARS-COV is located in direct RBD-ACE2 binding interface has the potential to be decorated by O-glycans. Previous experimental and computational studies emphasized the role of O-glycosylation in SARS-COV-2 high

Figure 10: Relative binding free energy components for RBD-ACE2 complex. (A) Overall binding free energy of models 0-III are shown in black, lime, orange and blue.(B) Molecular Mechanics (MM) part of free energy for all models. Colors are same as A. (C) Solvation part of free energy for all models. Colors are same as A

infectivity. The results of atomistic molecular dynamics simulations of SARS-COV-2 RBD in complex with ACE2 suggest that the O-glycosylation of S494 leads to the stronger interaction between RBD-ACE2 which can increase the virus infectivity. We tested three models of core and elongated O-glycans attached to RBD which lead to less flexible ACE-RBD dynamics and a decrease of distances between RBD and two major H1, H2 helices of ACE2 in the direct binding interface. Relative binding free energy of RBD-ACE2 is also more favorable in all O-glycosylated models. The increase of RBD binding affinity to ACE2 depends on the size of attached O-glycan. By increasing the size of O-glycan, the RBD-ACE2 binding affinity will increase. These findings add insightful information to the current status of SARS-COV-2-ACE2 glycosylation and its role in the virus's high evasion rate. This hypothesis is a suitable target for experimental validation, and if proven to be critical, for future therapeutic designs.

Acknowledgement

We are grateful to the HPC of SUT to provide the computational resources of this study. We are also thankful to H.Rajabi-Maham, H.Mobasheri, R.S.Haltiwanger and ,R.J.Woods for their very useful suggestions.

References

(1) Andersen, K. G.; Rambaut, A.; Lipkin, W. I.; Holmes, E. C.; Garry, R. F. The proximal origin of SARS-CoV-2. *Nature medicine* **2020**, *26*, 450–452.

(2) Wrapp, D.; Wang, N.; Corbett, K. S.; Goldsmith, J. A.; Hsieh, C.-L.; Abiona, O.; Graham, B. S.; McLellan, J. S. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* **2020**, *367*, 1260–1263.

(3) Lan, J.; Ge, J.; Yu, J.; Shan, S.; Zhou, H.; Fan, S.; Zhang, Q.; Shi, X.; Wang, Q.; Zhang, L.; Wang, X. Crystal structure of the 2019-nCoV spike receptor-binding domain bound with the ACE2 receptor. *bioRxiv* **2020**,

(4) Brielle, E. S.; Schneidman-Duhovny, D.; Linial, M. The SARS-CoV-2 exerts a distinctive strategy for interacting with the ACE2 human receptor. *bioRxiv* **2020**,

(5) Wang, C.; Li, W.; Drabek, D.; Okba, N. M.; van Haperen, R.; Osterhaus, A. D.; van Kuppeveld, F. J.; Haagmans, B. L.; Grosveld, F.; Bosch, B.-J. A human monoclonal antibody blocking SARS-CoV-2 infection. *Nature communications* **2020**, *11*, 1–6.

(6) Catalan-Dibene, J. Human antibodies can neutralize SARS-CoV-2. 2020.

(7) Yan, R.; Zhang, Y.; Li, Y.; Xia, L.; Guo, Y.; Zhou, Q. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science* **2020**, *367*, 1444–1448.

(8) Han, Y.; Král, P. Computational Design of ACE2-Based Peptide Inhibitors of SARS-CoV-2. *ACS nano* **2020**, *14*, 5143–5147.

(9) He, J.; Tao, H.; Yan, Y.; Huang, S.-Y.; Xiao, Y. Molecular mechanism of evolution and human infection with the novel coronavirus (2019-nCoV). *bioRxiv* **2020**,

(10) Tan, J.; Verschueren, K. H.; Anand, K.; Shen, J.; Yang, M.; Xu, Y.; Rao, Z.; Bigalke, J.; Heisen, B.; Mesters, J. R.; Chen, K.; Shen, X.; Jiang, H.; Hilgenfeld, R. pH-dependent Conformational Flexibility of the SARS-CoV Main Proteinase (Mpro) Dimer: Molecular Dynamics Simulations and Multiple X-ray Structure Analyses. *Journal of Molecular Biology* **2005**, *354*, 25 – 40.

(11) Rath, S. L.; Kumar, K. Investigation of the effect of temperature on the structure of SARS-Cov-2 Spike Protein by Molecular Dynamics Simulations. *bioRxiv* **2020**,

(12) Mariño, K.; Bones, J.; Kattla, J. J.; Rudd, P. M. A systematic approach to protein glycosylation analysis: a path through the maze. *Nature chemical biology* **2010**, *6*, 713–723.

(13) Shajahan, A.; Supekar, N. T.; Gleinich, A. S.; Azadi, P. Deducing the N-and O-glycosylation profile of the spike protein of novel coronavirus SARS-CoV-2. *bioRxiv* **2020**,

(14) Zhao, P. et al. Virus-Receptor Interactions of Glycosylated SARS-CoV-2 Spike and Human ACE2 Receptor. *bioRxiv* **2020**,

(15) Fung, T. S.; Liu, D. X. Post-translational modifications of coronavirus proteins: roles and function. *Future Virology* **2018**, *13*, 405–430.

(16) Parsons, L. M.; Bouwman, K. M.; Azurmendi, H.; De Vries, R. P.; Cipollo, J. F.; Verheije, M. H. Glycosylation of the viral attachment protein of avian coronavirus is essential for host cell and receptor binding. *Journal of Biological Chemistry* **2019**, *294*, 7797–7809.

(17) Uslupehlivan, M.; Şener, E. Glycoinformatics approach for identifying target positions to inhibit initial binding of SARS-CoV-2 S1 protein to the host cell. *bioRxiv* **2020**,

(18) Watanabe, Y.; Bowden, T. A.; Wilson, I. A.; Crispin, M. Exploitation of glycosylation in enveloped virus pathobiology. *Biochimica et Biophysica Acta (BBA)-General Subjects* **2019**,

(19) Hounsell, E. F.; Davies, M. J.; Renouf, D. V. O-linked protein glycosylation structure and function. *Glycoconjugate journal* **1996**, *13*, 19–26.

(20) Singh, K.; Tripathi, R. P. An Overview on Glyco-Macrocycles: Potential New Lead and their Future in Medicinal Chemistry. *Current Medicinal Chemistry* **2020**,

(21) Ridley, C.; Thornton, D. J. Mucins: the frontline defence of the lung. *Biochemical Society Transactions* **2018**, *46*, 1099–1106.

(22) Lamblin, G.; Lhermitte, M.; Klein, A.; Houdret, N.; Scharfman, A.; Ramphal, R.; Roussel, P. The carbohydrate diversity of human respiratory mucins: a protection of the underlying mucosa. *Am Rev Respir Dis* **1991**, *144*, S19–S24.

(23) Cristelo, C.; Azevedo, C.; Marques, J. M.; Nunes, R.; Sarmento, B. SARS-CoV-2 and Diabetes: New Challenges for the Disease. *Diabetes Research and Clinical Practice* **2020**, 108228.

(24) Kirschner, K. N.; Yongye, A. B.; Tschampel, S. M.; González-Outeiriño, J.; Daniels, C. R.; Foley, B. L.; Woods, R. J. GLYCAM06: a generalizable biomolecular force field. Carbohydrates. *Journal of computational chemistry* **2008**, *29*, 622–655.

(25) Mariño, K.; Bones, J.; Kattla, J. J.; Rudd, P. M. A systematic approach to protein glycosylation analysis: a path through the maze. *Nature chemical biology* **2010**, *6*, 713–723.

(26) Anandakrishnan, R.; Aguilar, B.; Onufriev, A. V. H++ 3.0: automating p K prediction and the preparation of biomolecular structures for atomistic molecular modeling and simulations. *Nucleic acids research* **2012**, *40*, W537–W541.

(27) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. *The Journal of chemical physics* **1983**, *79*, 926–935.

(28) Klauda, J. B.; Venable, R. M.; Freites, J. A.; O'Connor, J. W.; Tobias, D. J.; Mondragon-Ramirez, C.; Vorobyov, I.; MacKerell Jr, A. D.; Pastor, R. W. Update of the CHARMM all-atom additive force field for lipids: validation on six lipid types. *The journal of physical chemistry B* **2010**, *114*, 7830–7843.

(29) Huang, J.; Rauscher, S.; Nawrocki, G.; Ran, T.; Feig, M.; de Groot, B. L.; Grubmüller, H.; MacKerell, A. D. CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nature methods* **2017**, *14*, 71–73.

(30) Phillips, J. C.; Braun, R.; Wang, W.; Gumbart, J.; Tajkhorshid, E.; Villa, E.; Chipot, C.; Skeel, R. D.; Kale, L.; Schulten, K. Scalable molecular dynamics with NAMD. *Journal of computational chemistry* **2005**, *26*, 1781–1802.

(31) Darden, T.; York, D.; Pedersen, L. Particle mesh Ewald: An N log (N) method for Ewald sums in large systems. *The Journal of chemical physics* **1993**, *98*, 10089–10092.

(32) Liu, H.; Hou, T. CaFE: a tool for binding affinity prediction using end-point free energy methods. *Bioinformatics* **2016**, *32*, 2216–2218.

(33) Genheden, S.; Ryde, U. The MM/PBSA and MM/GBSA methods to estimate ligand-binding affinities. *Expert opinion on drug discovery* **2015**, *10*, 449–461.

(34) Wang, E.; Sun, H.; Wang, J.; Wang, Z.; Liu, H.; Zhang, J. Z.; Hou, T. End-point binding

free energy calculation with MM/PBSA and MM/GBSA: strategies and applications in drug design. *Chemical reviews* **2019**, *119*, 9478–9508.

(35) Baker, N. A.; Sept, D.; Joseph, S.; Holst, M. J.; McCammon, J. A. Electrostatics of nanosystems: application to microtubules and the ribosome. *Proceedings of the National Academy of Sciences* **2001**, *98*, 10037–10041.

(36) Hou, T.; Wang, J.; Li, Y.; Wang, W. Assessing the performance of the MM/PBSA and MM/GBSA methods. 1. The accuracy of binding free energy calculations based on molecular dynamics simulations. *Journal of chemical information and modeling* **2011**, *51*, 69–82.

(37) Humphrey, W.; Dalke, A.; Schulten, K. VMD – Visual Molecular Dynamics. *Journal of Molecular Graphics* **1996**, *14*, 33–38.

(38) Hunter, J. D. Matplotlib: A 2D graphics environment. *Computing in science & engineering* **2007**, *9*, 90–95.

(39) Van Rossum, G.; Drake, F. L. *Python 3 Reference Manual*; CreateSpace: Scotts Valley, CA, 2009.

(40) Schrödinger, LLC, The PyMOL Molecular Graphics System, Version 1.8. **2015**, There is no corresponding record for this reference.

(41) Lan, J.; Ge, J.; Yu, J.; Shan, S.; Zhou, H.; Fan, S.; Zhang, Q.; Shi, X.; Wang, Q.; Zhang, L., et al. Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature* **2020**, *581*, 215–220.

(42) Li, Y.; Cong, Y.; Feng, G.; Zhong, S.; Zhang, J. Z.; Sun, H.; Duan, L. The impact of interior dielectric constant and entropic change on HIV-1 complex binding free energy prediction. *Structural Dynamics* **2018**, *5*, 064101.

(43) Srivastava, H. K.; Sastry, G. N. Molecular dynamics investigation on a series of HIV protease inhibitors: assessing the performance of MM-PBSA and MM-GBSA approaches. *Journal of chemical information and modeling* **2012**, *52*, 3088–3098.