Glutamine deprivation triggers NAGK-dependent hexosamine salvage

Sydney L. Campbell[1,2], Clementina Mesaros[3], Luke Izzo[1,2], Hayley Affronti[1,2], Michael Noji[1,2],

Bethany E. Schaffer[4], Tiffany Tsang[1,2], Kathryn Sun[5], Sophie Trefely[1,2,6], Salisa Kruijning[1,2],

John Blenis[4], Ian A. Blair[3], Kathryn E. Wellen[1,2,7]

[1]Department of Cancer Biology, [2]Abramson Family Cancer Research Institute, [3]Department of

Systems Pharmacology and Translational Therapeutics, University of Pennsylvania,

Philadelphia, PA 19104; [4]Meyer Cancer Center and Department of Pharmacology, Weill Cornell

Medicine, New York, NY 10021; [5]Pancreatic Cancer Research Center, Perelman School of

Medicine, University of Pennsylvania, Philadelphia, PA 19104; [6]Center for Metabolic Disease

Research, Department of Microbiology and Immunology, Lewis Katz School of Medicine,

Temple University, Philadelphia, PA 19140. [7]Corresponding author.

Abstract

Tumors frequently exhibit aberrant glycosylation, which can impact cancer progression and

therapeutic responses. The hexosamine biosynthesis pathway (HBP) produces uridine

diphosphate N-acetylglucosamine (UDP-GlcNAc), a major substrate for glycosylation in the cell.

Prior studies have identified the HBP as a promising therapeutic target in pancreatic ductal

adenocarcinoma (PDA). The HBP requires both glucose and glutamine for its initiation. The

PDA tumor microenvironment is nutrient poor, however, prompting us to investigate how

nutrient limitation impacts hexosamine synthesis. Here, we identify that glutamine limitation in

PDA cells suppresses de novo hexosamine synthesis but results in increased free GlcNAc

abundance. GlcNAc salvage via N-acetylglucosamine kinase (NAGK) is engaged to feed UDP-

GlcNAc pools. *NAGK* expression is elevated in human PDA, and *NAGK* deletion from PDA cells

impairs tumor growth in mice. Together, these data identify an important role for NAGK-

dependent hexosamine salvage in supporting PDA tumor growth.

Introduction

Altered glycosylation is frequently observed in malignancies, impacting tumor growth as well

as immune and therapeutic responses (Akella et al., 2019; Mereiter et al., 2019; Munkley,

2019). Several types of glycosylation, including O-GlcNAcylation and N-linked glycosylation, are

dependent on the glycosyl donor uridine diphosphate N-acetylglucosamine (UDP-GlcNAc),

which is synthesized by the hexosamine biosynthesis pathway (HBP). The HBP branches off

from glycolysis with the transfer of glutamine's amido group to fructose-6-phosphate (F-6-P) to

generate glucosamine-6-phosphate (GlcN-6-P), mediated by the rate limiting enzyme

glutamine—fructose-6-phosphate transaminase (GFPT1/2). The pathway further requires

acetyl-CoA, ATP, and uridine triphosphate (UTP) to ultimately generate UDP-GlcNAc. O-

GlcNAcylation, the addition of a single N-acetylglucosamine (GlcNAc) moiety onto a serine or

threonine residue of intracellular proteins, is upregulated in multiple cancers (Akella, et al.,

2019). Targeting O-GlcNAcylation suppresses the growth of breast, prostate, and colon cancer

tumors (Caldwell et al., 2010; Ferrer et al., 2017; Gu et al., 2010; Guo et al., 2017; Lynch et al.,

2012). Similarly, highly branched N-glycan structures are sensitive to HBP flux and are

upregulated in malignant tissue, and targeting the relevant Golgi GlcNAc transferase enzymes

can limit tumor growth and metastasis in vivo (Granovsky et al., 2000; Li et al., 2008; Zhou et

al., 2011). Thus, improved understanding the regulation of the HBP in cancer could point

towards novel therapeutic strategies.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Pancreatic ductal adenocarcinoma (PDA) is a deadly disease with a 5-year survival rate of 9%

and a rising number of annual deaths (Rahib et al., 2014) (ACS Cancer Facts and Figures 2019,

NIH SEER report 2019). Mutations in *KRAS* occur in nearly all cases of human PDA and drive

extensive metabolic reprogramming in cancer cells. Enhanced flux into the HBP was identified

as a primary metabolic feature mediated by mutant KRAS in PDA cells (Ying et al., 2012).

Hypoxia, a salient characteristic of the tumor microenvironment (Lyssiotis and Kimmelman,

2017), was shown to further promote expression of glycolysis and HBP genes in pancreatic

cancer cells (Guillaumond et al., 2013). Notably, the glutamine analog 6-diazo-5-oxo-L-

norleucine (DON), which inhibits the HBP, suppressed PDA metastasis and sensitized PDA

tumors to anti-PD1 therapy (Sharma et al., 2020). DON has also been reported to sensitize PDA

cells to the chemotherapeutic gemcitabine in vitro (Chen et al., 2017). Additionally, a recently

developed inhibitor targeting the HBP enzyme phosphoacetylglucosamine mutase 3 (PGM3)

enhances gemcitabine-mediated reduction of xenograft tumor growth in vivo (Ricciardiello et al.,

2020). Thus, the HBP may represent a therapeutic target in PDA, although the regulation of

UDP-GlcNAc synthesis and the optimal strategies to target this pathway for therapeutic benefit

in PDA remain poorly understood.

An outstanding question is the impact of the tumor microenvironment on UDP-GlcNAc

synthesis. The HBP has been proposed as a nutrient-sensing pathway since its rate-limiting

step, mediated by GFPT1/2, requires both glutamine and the glycolytic intermediate fructose-6-

phosphate (Denzel and Antebi, 2015). In hematopoietic cells, glucose deprivation limits UDP-

GlcNAc levels and dramatically reduces levels of the N-glycoprotein IL3Rα at the plasma

membrane in a manner dependent on the HBP (Wellen et al., 2010). Similarly, O-GlcNAcylation

of certain nuclear-cytosolic proteins, including cancer-relevant proteins such as Myc and Snail,

has been demonstrated to be nutrient sensitive, impacting protein stability or function (Housley

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

et al., 2008; Park et al., 2010; Swamy et al., 2016). Yet, the PDA tumor microenvironment is

thought to be particularly nutrient poor, owing to its characteristic dense stroma (Halbrook and

Lyssiotis, 2017). This raises the question of how nutrient deprivation impacts the synthesis of

UDP-GlcNAc and its utilization for glycosylation. Understanding how PDA cells regulate these

processes under nutrient limitation could identify therapeutic vulnerabilities. In this study, we

investigated the impact of nutrient deprivation on the HBP and glycosylation in PDA cells,

identifying a key role for hexosamine salvage through the enzyme N-acetylglucosamine kinase

(NAGK) in PDA tumor growth.

Materials and methods

KEY RESOURCES TABLE				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
cell line (Homo sapiens)	MIA PaCa-2	ATCC	CRL-1420	
cell line (Homo sapiens)	PANC-1	ATCC	CRL-1469	
cell line (Homo sapiens)	HPAC	ATCC	CRL-2119	
cell line (Homo sapiens)	AsPC-1	ATCC	CRL-1682	
cell line (Homo sapiens)	BxPC-3	ATCC	CRL-1687	
cell line (Homo sapiens)	HCT 116	ATCC	CCL-247	

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

cell line (Homo sapiens)	SW480	ATCC	CCL-228	
Plasmid	lentiCRISPR v2	Addgene	52961	
Antibody	O-GlcNAc CTD1106	Cell Signaling	9875S	
Antibody	tubulin	Sigma	T6199	
Antibody	HSP60	Cell Signaling	12165S	
Antibody	NAGK	Atlas Antibodies	HPA035207	
Antibody	NAGK	Proteintech	15051-1-AP	
Antibody	AKT phosphoS473	Cell Signaling	4060	
Antibody	Viniculin	Sigma	V9264	
Chemical compound	[U-13C]-glucose	Cambridge Isotopes	CLM-1396-1	
Chemical compound	13C GlcNAc	Omicron Biochemicals	GLC-006	
Strain	NCr nude mice	Taconic	CrTac:NCr-Foxn1nu	
Sequenced-based reagent	*GFPT1* forward	This paper	RT-qPCR primers	CTCTGGCTTTGGTGGATA AA
Sequenced-based reagent	*GFPT1* reverse	This paper	RT-qPCR primers	GCAACCACTTGCTGAAGA
Sequenced-based reagent	*NAGK* forward	This paper	RT-qPCR primers	GTGCTCATATCTGGAACA GG
Sequenced-based reagent	*NAGK* reverse	This paper	RT-qPCR primers	ACCCTCATCACCCATCAT A
Sequenced-based reagent	*HPRT* forward	This paper	RT-qPCR primers	ATTATGCCGAGGATTTGG AA
Sequenced-based reagent	*HPRT* reverse	This paper	RT-qPCR primers	CCCATCTCCTTCATGACA TCT

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Sequenced-based reagent	*RPL19* forward	This paper	RT-qPCR primers	CAAGAAGGAGGAGATCATCAAG
Sequenced-based reagent	*RPL19* reverse	This paper	RT-qPCR primers	ATCACAGAGGCCAGTATGTA
Sequenced-based reagent	sgMGAT5 mouse forward	Doench et al., 2016	CRISPR deletion primers	CACCGGCTGTCATGACACCAGCGTA
Sequenced-based reagent	sgMGAT5 mouse reverse	Doench et al., 2016	CRISPR deletion primers	AAACTACGCTGGTGTCATGACAGCC
Sequenced-based reagent	sgNAGK#1 forward	Doench et al., 2016	CRISPR deletion primers	CACCGTTGACGTAGCCGATATCATG
Sequenced-based reagent	sgNAGK#1 reverse	Doench et al., 2016	CRISPR deletion primers	AAACCATGATATCGGCTACGTCAAC
Sequenced-based reagent	sgNAGK#2 forward	Doench et al., 2016	CRISPR deletion primers	CACCGTGCTTGGTGTGCGATCCAGT
Sequenced-based reagent	sgNAGK#2 reverse	Doench et al., 2016	CRISPR deletion primers	AAACACTGGATCGCACACCAAGCAC
Sequenced-based reagent	sgNAGK#3 forward	Doench et al., 2016	CRISPR deletion primers	CACCGCTCTACACCCCCATAGATCG
Sequenced-based reagent	sgNAGK#3 reverse	Doench et al., 2016	CRISPR deletion primers	AAACCGATCTATGGGGGTGTAGAGC
Sequenced-based reagent	siRNA non-targeting control	Santa Cruz Biotechnology	SC-37007	
Sequenced-based reagent	siGALE	Santa Cruz Biotechnology	SC-78950	
Sequenced-based reagent	siGNE	Santa Cruz Biotechnology	SC-60693	

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Sequenced-based reagent	siGALK2	Santa Cruz Biotechnology	SC-90002	

Cell culture

Cells were cultured in DMEM high glucose (Gibco, 11965084) with 10% calf serum (Gemini

GemCell U.S. Origin Super Calf Serum, 100-510), unless otherwise noted. Glucose- or

glutamine-restricted media was prepared using glucose, glutamine, and phenol red free DMEM

(Gibco, A1443001) supplemented with glucose (Sigma-Aldrich, G8769), glutamine (Gibco,

25030081), and dialyzed fetal bovine serum (Gemini, 100-108). For all glutamine restriction

experiments except S2.3 D, cells were plated 2-3x more densely for the nutrient restricted

condition samples to achieve similar confluency at the experiment endpoint. 1% oxygen levels

were achieved by culturing cells in a Whitley H35 Hypoxystation (Don Whitley Scientific). ATCC

names and numbers for the cell lines used in this study are: MIA PaCa-2 (ATCC# CRL-1420),

PANC-1 (ATCC# CRL-1469), HPAC (ATCC# CRL-2119), AsPC-1 (ATCC# CRL-1682), BxPC-3

(ATCC# CRL-1687), HCT 116 (ATCC# CCL-247), and SW480 (ATCC# CCL-228). All cells were

routinely tested for mycoplasma and authenticated by short tandem repeat (STR) profiling using

the GenePrint 10 System (Promega, B9510).

Generation of CRISPR cell lines

96 sgRNA sequences targeting *NAGK* or *Mgat5* from the Brunello and Brie libraries (Doench et al.,

97 2016) were cloned into the lentiCRISPRv2 vector (Sanjana, Shalem et al., 2014). Lentivirus was

98 produced in 293T cells according to standard protocol. Cells were then infected with the

99 CRISPR lentivirus and selected with puromycin. Cells were plated at very low density into 96

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

100   well plates to establish colonies generated from single cell clones. *Mgat5* gene disruption was

101   validated by qPCR and L-PHA binding. *NAGK* gene disruption was validated by qPCR, western

102   blot, and ^{13}C-GlcNAc tracing. Seven *NAGK* knockout clonal cell lines established from three

103   different sgRNAs, four in PANC-1 cells and three in MIA PaCa-2 cells, were chosen for use in

104   the study. Please see table at end of methods for primer sequences of guides used.

Western blotting

For protein extraction from cells, cells were kept on ice and washed three times with PBS, then

scraped into PBS and spun down at 200g for 5 minutes. The cell pellet was resuspended in 50-

100 µL RIPA buffer [1% NP-40, 0.5% deoxycholate, 0.1% SDS, 150 mM NaCl, 50 mM Tris plus

protease inhibitor cocktail (Sigma-Aldrich, P8340) and phosSTOP (Sigma-Aldrich,

04906845001)] and lysis was allowed to continue on ice for 10 minutes. Cells were sonicated

with a Fisherbrand Model 120 Sonic Dismembrator (Fisher Scientific, FB120A110) for three

pulses of 20 seconds each at 20% amplitude. Cell lysate was spun down at 15,000g for 10

minutes at 4^{o}C and supernatant was transferred to a new tube. For protein extraction from

tissue, the sample was resuspended in 500 µL RIPA buffer and homogenized using a

TissueLyser (Qiagen, 85210) twice for 30s at 20 Hz. Following incubation on ice for 10 minutes

the same procedure was followed as for cells. For both cells and tissue, lysate samples were

stored at -80^{o}C until analysis by immunoblot. All blots were developed using a LI-COR Odyssey

CLx system. Antibodies used in this study were: O-GlcNAc CTD110.6 (Cell Signaling 9875S),

tubulin (Sigma T6199), HSP60 (Cell Signaling 12165S), NAGK (Atlas Antibodies, HPA035207),

and PARP (Cell Signaling 9532).

For blots showing the mobility shift for NAGK in low glutamine, samples were prepared in lysis

buffer containing 50 mM Tris pH 8.0, 150 mM NaCl, 0.5% IGEPAL CA-630 (Sigma, I3021), 1

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

mM PMSF, 1.5 µM aprotinin, 84 µM leupeptin, 1 µM pepstatin A, -/+ 10 mM NaF and 20 mM

Na$_3$VO$_4$ as indicated. To visualize the NAGK mobility shift in response to low glutamine, 20 µg

total protein per sample was separated across 12.5 cm of 11% SDS-PAGE resolving space

under reducing conditions using the large electrophoresis systems available from C.B.S.

Scientific until approximately 3 cm of separation was obtained between the 25 and 37 kDa

protein standards (Bio-Rad; 1610375). Using electrophoresis, proteins were transferred (30 V,

4°C, overnight) to 0.45 uM pore size nitrocellulose membrane (Amersham, 10600002). The

primary antibodies used were NAGK (Proteintech, 15051-1-AP), AKT phosphoS473 (Cell

Signaling Technology, 4060), and Vinculin (Sigma, V9264). Membranes were developed using

the LI-COR Odyssey CLx system.

RT-qPCR

For RNA extraction from cells, cells were put on ice, washed with PBS, and scraped into PBS.

Samples were then spun down at 200g for 5 minutes and resuspended in 100 µL Trizol (Life

Technologies). For RNA extraction from tissue, samples were resuspended in 500 µL Trizol and

homogenized using a TissueLyser twice for 30s at 20 Hz. For both cells and tissue, RNA was

extracted following the Trizol manufacturer protocol. cDNA was prepared using high-capacity

RNA-to-cDNA master mix (Applied Biosystems, 4368814) according to kit instructions. cDNA

was diluted 1:20 and amplified with PowerUp SYBR Green Master Mix (Applied Biosystems,

A25778) using a ViiA-7 Real-Time PCR system. Fold change in expression was calculated by

the $\Delta\Delta C_t$ method using HPRT as a control. Please see table at end of chapter for primer

sequences.

Lectin binding assay

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Cells were put on ice, washed with PBS and then scraped into PBS. Samples were then spun

down at 200g for 5 minutes and resuspended in 3% BSA with fluorophore-conjugated lectin

added 1:1000 (Vector Labs FL-1111-2). Samples were covered and incubated on ice for 30

minutes at room temperature, then spun down and resuspended in PBS before analysis with an

Attune NxT Flow Cytometer (Thermo Fisher Scientific). Data was further analyzed using FlowJo

8.7.

Metabolite quantitation and labeling

For all metabolite quantitation experiments, each sample was collected from a 10 cm sub-

confluent plate of cells. To achieve similar confluency and protein content at the experiment

endpoint, cells were initially plated more densely for the nutrient deprived samples than for the

nutrient replete samples. For low glutamine experiments, PANC-1 cells were plated $3x10^5$ for 4

mM glutamine samples and $5.5x10^5$ for 0.05 mM samples. MIA PaCa-2 cells were plated $3x10^5$

for 4 mM samples and $1.2x10^6$ for 0.05 mM samples.

Samples were prepared according to Guo et al. (Guo et al., 2016). Briefly, cells were put on ice

and washed 3x with PBS. Then, 1 mL of ice cold 80% methanol was added to the plate, and

cells were scraped into solvent and transferred to a 1.5 mL tube. For quantitation experiments,

internal standard containing a mix of ^{13}C labeled metabolites was added at this time. Samples

were then sonicated and spun down, and the supernatants were dried down under nitrogen.

The dried samples were then resuspended in 100 μL of 5% sulfosalicylic acid and analyzed by

liquid chromatography-high resolution mass spectrometry as reported (Guo et al., 2016) with the

only modification that the LC was coupled to a Q Exactive-HF with a heated ESI source

operating in negative ion mode alternating full scan and MS/MS modes. The [M-H]$^-$ ion of each

analyte and its internal standard was quantified, with peak confirmation by MS/MS. GlcNAc

quantification was done on a triple quadropole instrument exactly as described (Guo et al.,

2016). Data analysis was conducted in Thermo XCalibur 3.0 Quan Browser and FluxFix (Trefely

et al., 2016). For quantitation experiments, samples were normalized first to peak integrations of

^{13}C-labeled internal standard components and then to protein content in the sample, measured

by BCA assay. Relative quantification was then calculated by normalizing to the control

condition in each experiment.

For glucose labeling experiments, cells were cultured in DMEM without glucose, glutamine, or

phenol red supplemented with 10 mM [U-^{13}C]-glucose (Cambridge Isotopes, CLM-1396-1), 4

mM glutamine, and 10% dialyzed fetal bovine serum. Cells were incubated for the indicated

time, and samples were prepared as above. For GlcNAc labeling experiments, cells were

cultured in DMEM without glucose, glutamine, or phenol red supplemented with 10 mM N-[1,2-

13C2]acetyl-D-glucosamine (^{13}C GlcNAc) (Omicron Biochemicals, GLC-006), 4 mM glutamine,

10 mM glucose, and 10% dialyzed fetal bovine serum. Cells were incubated for the indicated

time, and samples were prepared as above.

Soft agar colony formation assay

Cells were trypsinized and counted using a Bright-Line hemacytometer (Sigma, Z359629). The

bottom agar layer was prepared by adding Bacto Agar (BD Bioscience, 214050) to cell culture

media for a final concentration of 0.6%. 2 mL bottom agar was added to each well of a 6-well

tissue culture plate. Once bottom agar solidified, top layer agar was prepared by combining

trypsinized cells with the bottom agar mix for a final concentration of 0.3% Bacto Agar. 1 mL top

layer agar was added to each well with a bottom layer of agar. Cells were plated 2.5x10^4 per

well. 0.5 mL DMEM high glucose with 10% calf serum was added to cells every 7 days. Images

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

190   were taken after 3 weeks. Images were blinded and colonies per image were counted using

191   ImageJ (Schneider et al., 2012).

2D Proliferation assay

193   Cells were plated 3.5×10^4 per well of a 6-well plate. For each day that counts were recorded,

194   three wells were trypsinized and cells were counted twice using a hemocytometer (Sigma,

195   Z359629). The average of the two counts was recorded for each well, and the average count of

196   the three wells was used to graph the data. For proliferation assays in 0.05 mM glutamine,

197   trypan blue was used during cell counts.

Bioinformatics data analysis

The PDAC expression profiling dataset (GEO accession GSE16515, Pei et al., 2009) from NCBI

GEO Profile database (Edgar et al., 2002) was used to compare the expression level between

human normal and PDAC tumor samples. The dataset consists of 52 samples, in which 16

samples are matched tumor and normal tissues, and 20 samples are only tumor tissues. The

statistical analysis was conducted by one-way ANOVA, the level of significance was evaluated

by $p < 0.01$ and plotted in box-and-whisker diagram. Comparison of HBP gene expression

between tumor (TCGA PAAD dataset) and normal tissue (GTEx) was also conducted using

GEPIA2 (Tang et al., 2019).

Tumor growth in vivo

3×10^6 PANC-1 NAGK CRISPR cells were injected with 1:1 Matrigel (Corning, CB354248) into

the flanks of NCr nude mice and measured with calipers once per week for 22 weeks. At the

experiment endpoint (22 weeks or when tumor reached 20 mm in length), mice were euthanized

with CO_2 and cervical dislocation. Tumors were removed, weighed, cut into pieces for analysis,

and frozen. All animal experiments were approved by the University of Pennsylvania and the

Institutional Animal Care and Use Committee (IACUC).

Results

Tetra-antennary N-glycans and O-GlcNAcylation are minimally impacted by nutrient
limitation in pancreatic cancer cells

To examine the effects of nutrient deprivation on glycosylation, we cultured cells under

glucose or glutamine limitation and examined O-GlcNAc levels and cell surface

phytohemagglutinin-L (L-PHA) binding, a readout of N-acetylglucosaminyltransferase 5

(MGAT5)-mediated cell surface N-glycans (Fig S1A, B), which are highly sensitive to UDP-

GlcNAc availability (Lau et al., 2007). We focused on glucose and glutamine because of their

requirement to initiate the HBP (Fig. 1A). First, as a positive control, we examined HCT-116 and

SW480 colon cancer cells, previously documented to have glucose-responsive O-GlcNAcylation

(Park et al., 2010; Steenackers et al., 2016), which we also confirmed in HCT-116 cells (Fig.

1B). Indeed, L-PHA binding was suppressed by glucose restriction in SW480 cells and by

glutamine restriction in both colon cancer cell lines (Fig. 1C). Next, to test whether glycans were

sensitive to nutrient restriction in PDA cells, we examined L-PHA binding and O-GlcNAc levels

under nutrient deprivation conditions in a panel of human PDA cell lines, including PANC-1, MIA

PaCa-2, AsPC-1, and HPAC. Across these cell lines, no consistent changes in L-PHA binding

were observed under glucose or glutamine limitation (Fig. 1D, E; Fig. S1C). We also examined

L-PHA binding in PDA cells under oxygen- or serum-deprived conditions and observed minimal

changes (Figure S1D, E). O-GlcNAcylation was minimally altered by culture in low glutamine

and exhibited variable changes in response to glucose limitation (Fig. 1F), consistent with

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Figure 1: MGAT5-dependent N-glycans are minimally impacted by glucose or glutamine deprivation in PDA cells.
A) Overview of the hexosamine biosynthesis pathway (HBP). **B)** O-GlcNAc levels in HCT-116 cells in high and low nutrients; cells were incubated in indicated concentrations of glucose and glutamine for 48 hours. **C)** Phytohemagglutinin-L (LPHA) binding in colon cancer cells. Cells were incubated in the indicated concentrations of glucose (left) or glutamine (right) for 48 hours and then analyzed by flow cytometry. Graph shows mean fluorescence intensity (MFI) relative to control condition. Statistical significance was calculated by unpaired t-test. **D-E)** Phytohemagglutinin-L (LPHA) binding in pancreatic ductal adenocarcinoma (PDA) cells in low nutrients. Cells were incubated in the indicated concentrations of glucose (D) or glutamine (E) for 48 hours and then analyzed by flow cytometry. Statistical significance was calculated by one-way ANOVA. **F)** O-GlcNAc levels in PDA cells in high and low nutrients. Cells were incubated in the indicated concentrations of glucose or glutamine for 48 hours. **G)** Western blot for O-GlcNAc and PARP in PANC-1 cells cultured in the indicated concentrations of glutamine with or without Thiamet-G (TMG) treatment for the indicated time. For all bar graphs, mean +/- standard error of the mean (SEM) of three biological triplicates is represented. Panels B) - G) are representative of at least two independent experimental replicates. *, p ≤ 0.05; **, p ≤ 0.01; ***, p ≤ 0.001.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

stress-induced regulation of this modification (Taylor et al., 2008). Since glycosylation may be

maintained through either sustained ability to add the modifications or through changes in

turnover, we assayed active O-GlcNAcylation by inhibiting O-GlcNAcase with Thiamet G (TMG).

TMG treatment resulted in equivalently elevated O-GlcNAcylation levels in high and low

glutamine conditions (Fig. 1G; Fig. S1F), indicating that glutamine restriction does not limit the

capacity of cells to add the O-GlcNAc modification. Mia-PaCa-2 cells exhibited some cell death

in low glutamine, though this was not exacerbated by TMG treatment (Fig. S1F). Thus, under a

variety of nutrient stress conditions, neither L-PHA binding nor O-GlcNAcylation were

consistently suppressed in pancreatic cancer cell lines. Glutamine restriction in particular had

remarkably little impact on O-GlcNAcylation and L-PHA binding, raising the question of how

UDP-GlcNAc is generated during nutrient limitation.

De novo UDP-GlcNAc synthesis is suppressed upon glutamine limitation

We therefore next asked whether the abundance of HBP metabolites is impacted by nutrient

limitation. We measured HBP metabolites after glucose or glutamine restriction using HPLC-MS

(Guo et al., 2016). In low glutamine conditions, GlcN-6-P levels were potently decreased relative

to glutamine replete conditions in PANC-1 cells, while UDP-GlcNAc abundance was maintained

(Fig. 2A). In MIA PaCa-2 cells, UDP-GlcNAc abundance actually increased upon glutamine

restriction (Fig S2.1A). These data indicate that UDP-GlcNAc might be generated through

mechanisms other than de novo synthesis. Glycolytic intermediates were minimally impacted by

low glutamine conditions, and TCA cycle intermediates such as α-KG and malate decreased as

expected (Fig. 2A, Fig. S2.1A). In contrast to that in glutamine restriction, UDP-GlcNAc

abundance declined in 5 mM or 0.1 mM relative to 10 mM glucose conditions (Fig. S2.1B),

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Figure 2: *De novo* UDP-GlcNAc synthesis is supressed upon glutamine limitation.
A) Metabolite measurements in PANC-1 and MIA PaCa-2 cells after culture for 48 hours in 0.05 mM glutamine. Quantification is normalized to 4 mM glutamine condition. Statistical significance was calculated by unpaired t-test.Mean +/- SEM is represented. **B)** Overview of the GlcNAc salvage pathway feeding into the HBP. GlcNAc scavenged from O-GlcNAc removal or lysosomal breakdown of glycans can be phosphorylated by NAGK and used to regenerate UDP-GlcNAc. **C)** Overview of the incorporation of ^{13}C glucose into UDP-GlcNAc. Different parts of the molecule can be labeled from glucose-derived subunits, thus isotopologues up to M+16 can be derived from glucose. **D)** ^{13}C glucose tracing into F-6-P, GlcNAc-P, and UDP-GlcNAc in indicated glutamine concentrations. % labeled GlcN indicates sum of M+6 and M+8 isotopologues for GlcNAc-P and sum of M+6, M+8, M+11, and M+13 for UDP-GlcNAc, % labeled Ribose indicates sum of M+5, M+7, M+11, and M+13 for UDP-GlcNAc. All isotopologues are graphed in Figure S2. Statistical significance was calculated by unpaired t-test. Mean +/- SEM is represented. *, p ≤ 0.05; **, p ≤ 0.01; ***, p ≤ 0.001.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

suggesting that glutamine limitation specifically may trigger an adaptive response to sustain

UDP-GlcNAc pools.

We sought to understand how UDP-GlcNAc pools are sustained during glutamine restriction.

In addition to de novo synthesis of UDP-GlcNAc via the HBP, free GlcNAc in the cell can also

be phosphorylated via N-acetylglucosamine kinase (NAGK) to produce GlcNAc-P and then

regenerate UDP-GlcNAc (Fig. 2B). However, NAGK's roles in physiology and cancer biology

have been minimally studied. To investigate the possibility that UDP-GlcNAc is generated

through mechanisms other than its synthesis from glucose, we first designed a stable isotope

labeling strategy to quantify the fraction of the glucosamine ring that is synthesized *de novo* in

glutamine-replete versus -restricted conditions. Since multiple components of UDP-GlcNAc

[glucosamine ring, acetyl group, uridine (both the uracil nucleobase and the ribose ring)] can be

synthesized from glucose, UDP-GlcNAc isotopologues up to M+16 can be generated from

glucose (Moseley et al., 2011) (Fig. 2C). In order to measure the glucose carbon incorporated

into GlcNAc-P and UDP-GlcNAc via the HBP, all isotopologues containing a fully labeled

glucosamine ring are added together (% labeled GlcN indicates sum of M+6, M+8, M+11, and

M+13 for UDP-GlcNAc and sum of M+6 and M+8 for GlcNAc-P) (Fig. 2C).

After 48 hours of glutamine restriction, cells were incubated with fresh low glutamine medium

containing [U-^{13}C]-glucose to track the incorporation of glucose carbons into hexosamine

intermediates. Across multiple PDA cell lines, the fractional labeling of the glucosamine ring in

both GlcNAc-P and UDP-GlcNAc pools was markedly suppressed by glutamine restriction,

indicating decreased de novo synthesis in low glutamine conditions (Fig. 2D, Fig. S2.1C-E).

Notably, labeling into the ribose component of UDP-GlcNAc was also suppressed [% labeled

ribose indicates sum of isotopologues containing M+5 (i.e., M+5, M+7, M+11, and M+13); Fig.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

2D, Fig. S2.1C]. Consistently, incorporation of ^{13}C glucose into uridine triphosphate (UTP) was

suppressed upon glutamine restriction (Fig. S2.2A), even though UTP levels were maintained or

increased (Fig. 2A, Fig. S2.1A), suggesting a role for nucleoside salvage in maintaining

nucleotide pool in these conditions. This is consistent with previous reports demonstrating that

autophagy/ ribophagy is a source of nucleosides in amino acid deprived conditions (Guo et al.,

2011; Wyant et al., 2018). Indeed, silencing of either of the uridine salvage enzymes uridine

kinase 1 or 2 (UCK1/2) resulted in decreased UDP, UTP, and UDP-GlcNAc levels (Fig. S2.2B-

C), indicating that nucleoside salvage contributes to maintaining uridine phosphate and UDP-

GlcNAc pools.Thus, glutamine restriction suppresses the *de novo* synthesis of both GlcNAc-P

and UTP, both of which are required to produce UDP-GlcNAc.

We noted that GlcNAc-P and UDP-GlcNAc pools labeled from glucose with similar but not

identical kinetics. While this is potentially due to limitations in detection since GlcNAc-P is much

less abundant than UDP-GlcNAc, we considered whether GlcNAc-P-independent pathways

may also have minor contributions to glucose-dependent UDP-GlcNAc labeling. Although

pathways through which glucose can feed into UDP-GlcNAc's glucosamine ring independent of

the HBP have not been described in mammalian cells to our knowledge, we nevertheless tested

the two major metabolic branch points diverging from UDP-GlcNAc, which mediate UDP-

GalNAc and sialic acid synthesis. UDP-galactose-4-epimerase (GALE) interconverts UDP-

GlcNAc and UDP-GalNAc, and UDP-GlcNAc-2-epimerase/ManAc kinase (GNE) initiates sialic

acid biosynthesis. Silencing of neither GALE nor GNE reduced UDP-GlcNAc labeling from

glucose, however, indicating that these enzymes are unlikely to facilitate a bypass pathway (Fig.

S2.2D-E). Although a minor contribution from another unknown pathway cannot be ruled out,

the slight apparent differences in timing of GlcNAc-P and UDP-GlcNAc labeling most likely

311 reflect technical limitations. Regardless, the data clearly indicate that UDP-GlcNAc abundance

312 is maintained despite reduced *de novo* hexosamine synthesis from glucose (Fig. 2D).

313 **GlcNAc salvage feeds UDP-GlcNAc pools in pancreatic cancer cells**

314 As mentioned, UDP-GlcNAc can be generated via phosphorylation of free GlcNAc by NAGK

315 generating GlcNAc-6-P (Fig. 2B). When supplemented, GlcNAc is salvaged into the UDP-

316 GlcNAc pool (Ryczko et al., 2016; Wellen et al., 2010). Endogenous sources of GlcNAc may

317 include removal of O-GlcNAc protein modifications or breakdown of glycoconjugates and

318 extracellular matrix components. Notably, intracellular levels of GlcNAc increase upon glutamine

319 restriction (Fig. 3A). Yet, the significance of GlcNAc salvage to maintenance of UDP-GlcNAc

320 pools has been little studied, and the proportion of UDP-GlcNAc generated via the NAGK-

321 dependent salvage pathway is unknown.

322 *NAGK* mRNA expression increased in PDA cell lines in low glutamine conditions, and in some

323 cell lines also in low glucose (Fig. S2.3A, B). *GFPT1* expression was also induced in both low

324 glucose conditions, consistent with a prior report (Moloughney et al., 2016), and in low

325 glutamine conditions (Fig. S2.3A), even though *de novo* synthesis is suppressed when

326 glutamine is limited. Protein levels of NAGK did not increase in concordance with mRNA at

327 these time points, however, although a mobility shift potentially indicative of post-translational

328 modification was apparent when protein lysates were run on a gel using a large electrophoresis

329 system (see methods; Fig S2.3C-D). Removal of the phosphatase inhibitor Na_3VO_4 from the

330 sample buffer prevented the mobility shift, suggesting that NAGK may be phosphorylated on

331 one or more residues in low glutamine conditions (Fig. S2.3D). Taken together, these data

332 indicate that under low glutamine conditions, GlcNAc availability for salvage increases and the

333 salvage enzyme NAGK is subject to regulation.

334 This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic
Acid to Support Growth."

Figure 3: GlcNAc salvage feeds UDP-GlcNAc pools in pancreatic cancer cells.
A) Measurement of GlcNAc in PANC-1 and MIA PaCa-2 cells after incubation in the indicated concentrations of glutamine for 24 and 48 hours. Mean +/- SEM of four biological replicates is represented. Statistical significance was calculated by unpaired t-test. **B)** Measurement of 13C GlcNAc labeled on the acetyl group into GlcNAc-P and UDP-GlcNAc in NAGK knockout cells. Cells were incubated with 10 mM 13C GlcNAc for 6 hours. Mean +/- SEM of three biological replicates is represented. Statistical significance was calculated by unpaired t-test comparing the mean incorporation of the two CRISPR clones and the empty vector (EV) control. **C)** Labeling of F-6-P and GlcNAc-P from 13C glucose in PANC-1 (left) and MIA PaCa-2 (right) NAGK knockout cells. Statistical significance was calculated by unpaired t-test comparing the mean incorporation of the four CRISPR clones and the EV control. Mean +/- SEM of three biological replicates is represented. **D)** Percent of combined UDP-GlcNAc isotopologues containing a labeled glucosamine ring or a labeled ribose from UTP, calculated from S3.1 (E). Statistical significance was calculated by unpaired t-test comparing the mean incorporation of the four CRISPR clones and the EV control. Mean +/- SEM of three biological replicates is represented. **E)** Measurement of HBP metabolites in PANC-1 NAGK knockout cells cultured in the indicated concentrations of glutamine. Statistical significance was calculated by unpaired t-test comparing the mean incorporation of the four CRISPR clones and the EV control. Mean +/- SEM of four biological replicates is represented. *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.01$.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

These findings prompted us to investigate the role of NAGK in UDP-GlcNAc synthesis in PDA

cells. We functionally examined the role of NAGK in PDA cell lines by using CRISPR-Cas9 gene

editing to generate NAGK knockout (KO) PANC-1 and MiaPaCa-2 clonal cell lines (Fig. S3.1A-

B). N-[1,2-$^{13}C_2$]acetyl-D-glucosamine (^{13}C GlcNAc) was efficiently salvaged in control cells, and

this was suppressed by NAGK deletion, as evidenced by reduced fractional labeling of GlcNAc-

P and UDP-GlcNAc (Fig. 3B). Since we did not observe any residual protein expression, we

hypothesized that the N-acetylgalactosamine (GalNAc) salvage enzyme GalNAc kinase

(GALK2) might be responsible for the remaining GlcNAc salvage in the absence of NAGK.

Indeed, silencing of GALK2 further suppressed incorporation of ^{13}C GlcNAc into GlcNAc-P and

UDP-GlcNAc in the NAGK KO cells (Fig. S3.1C).

We hypothesized that knockout cells would conversely conduct increased de novo UDP-

GlcNAc synthesis. To test this, we incubated cells with [U-^{13}C]-glucose and examined

incorporation into GlcNAc-P and UDP-GlcNAc. Indeed, in the absence of NAGK, we observed

increased glucose-dependent fractional labeling of the glucosamine ring of UDP-GlcNAc and

GlcNAc-P, but not the ribose component of UDP-GlcNAc (Fig. 3C-D; Fig. S3.1D-E). This effect

was also observed with knockdown of NAGK by shRNA, though to a lesser extent (Fig. S3.2A-

B). Incorporation of glucose into F-6-P did not change (Fig. 3C) and the proportion of UDP-

GlcNAc containing an M+5 ribose ring was also unchanged in knockout cells (Fig. 3D), as

expected. Thus, when NAGK is deleted and GlcNAc salvage is suppressed, de novo

hexosamine synthesis increases.

We next assessed changes in the levels of hexosamine intermediates in control and NAGK

KO cell lines. In PANC-1 KO cells in 4 mM glutamine, GlcN-P increased significantly, consistent

with increased de novo synthesis in the absence of NAGK (Fig. 3E). GlcNAc-P was modestly

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

reduced in KO cells, though UDP-GlcNAc levels were maintained (Fig. 3E). In MIA PaCa-2

cells, GlcNAc-P was markedly suppressed in the absence of NAGK, though UDP-GlcNAc was

not (Fig. S3.2C). We also measured HBP metabolites under glutamine restriction, where we

expected NAGK would play a more significant role in UDP-GlcNAc generation. We were only

able to measure metabolites accurately in PANC-1 KO cells because MIA PaCa-2 NAGK KO

cells began to die quickly in low glutamine, which will be discussed further in the next section.

GlcN-P decreased in control and KO cells, consistent with reduced de novo hexosamine

synthesis (Fig. 3E). GlcNAc-P levels decreased in low glutamine in control cells, and decreased

further in cells lacking NAGK, consistent with contributions from both de novo synthesis and

salvage (Fig. 3E). Reciprocally, GlcNAc abundance was elevated upon glutamine limitation in

both control and NAGK KO cells (Fig. 3E). UDP-GlcNAc abundance was modestly reduced in

NAGK KO cells relative to controls under glutamine restriction, though levels were still

comparable to that in high glutamine (Fig. 3E), possibly reflecting changes in utilization. GALK2

silencing did not further suppress UDP-GlcNAc in NAGK KO cells (Fig. S3.2D), suggesting that

GALK2 may not have a major role in physiological GlcNAc salvage. Cumulatively, the data

demonstrate that GlcNAc is salvaged into UDP-GlcNAc pools in PDA cells in a manner

dependent at least in part on NAGK.

NAGK knockout limits tumor growth *in vivo*

To test the role of NAGK in cell proliferation, we first monitored growth of NAGK KO cells

compared to controls in 2D and 3D culture in 4 mM glutamine, finding minimal differences (Fig.

4A, Fig. S4A). We hypothesized that NAGK KO cell proliferation would be impaired in 0.05 mM

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Figure 4: NAGK expression is increased in human PDA tumors and NAGK knockout reduces tumor growth in vivo.
A) 2D proliferation assay by cell count in PANC-1 and MIA PaCa-2 NAGK knockout cells. Mean +/- SEM of three technical replicates is represented. Statistical significance was calculated using one-way ANOVA at each time point. **B)** Gene expression data for NAGK, GFPT1, PGM3, and UAP1 in human PDA tumors compared with matched normal tissue. Statistical analysis was conducted by one-way ANOVA, and level of significance was defined as p ≤ 0.01. **C)** Final tumor volume and **D)** final tumor weight of subcutaneous tumors generated from PANC-1 NAGK knockout cells in vivo. Cells were injected into the right flank of NCr nude mice and tumor volume was calculated from caliper measurements. Statistical significance was calculated by one-way ANOVA comparing each mean to the EV control mean. Mean +/- SEM of biological replicates is represented (n = 8 each group). *, p ≤ 0.05; **, p ≤ 0.01; ***, p ≤ 0.001.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

glutamine, where *de novo* UDP-GlcNAc synthesis is suppressed. Indeed, MIA PaCa-2 KO cells

died more quickly in 0.05 mM glutamine than did control cells (Fig. 4A). PANC-1 KO cells did

not show this effect (Fig. 4A), but we hypothesized that NAGK loss might have a stronger effect

in vivo where tumor growth can be constrained by nutrient availability.

To gain initial insight into whether NAGK is likely to play a functional role in PDA progression

in vivo, we queried publicly available datasets. From analysis of publicly available microarray

data (Pei et al., 2009) and gene expression data from the Cancer Genome Atlas (TCGA), we

indeed found *NAGK* expression to be increased in tumor tissue relative to adjacent normal

regions of the pancreas or to pancreas GTEx data (Fig. 4B, Fig. S4B). *GFPT1* expression was

also increased in tumor tissue (Fig. 4B, Fig. S4B), consistent with its regulation by mutant KRAS

(Ying et al., 2012). Two other HBP genes, *PGM3* and *UAP1*, did not show significantly

increased expression in PDA tumors in these datasets (Fig. 4B, Fig. S4B). We then studied the

role of NAGK in tumor growth *in vivo* by injecting NAGK CRISPR KO cells into the flank of NCr

nude mice. Final tumor volume and weight were markedly reduced in the absence of NAGK

(Fig. 4C-D). Of note, initial tumor growth was comparable between control and KO cells, but the

NAGK knockout tumors either stopped growing or shrank while control tumors continued to

grow larger (Fig. S4C). Interestingly, KO tumor samples showed increased L-PHA signal (Fig.

S4D), indicating that NAGK deficiency results in altered glycosylation within tumors. This could

possibly reflect either elevated de novo synthesis in the small tumors that form or differences in

cellular composition. For example, activated fibroblast marker α-smooth muscle actin (α-SMA)

was more abundant in the NAGK KO tumors (Fig S4D). Residual NAGK signal in whole tumors

also presumably reflects expression in other cell types (Fig. S4D), since NAGK was

undetectable in the clonal cell lines used for injections (Fig. S3.1B). Taken together, these data

are consistent with the notion that NAGK is dispensable when nutrients are abundant but

becomes more important as the tumors outgrow their original nutrient supply and become more

dependent on scavenging and recycling and indicate that NAGK-mediated hexosamine salvage

supports tumor growth *in vivo*.

DISCUSSION

In this study, we identify a key role for NAGK in salvaging GlcNAc for UDP-GlcNAc synthesis

in PDA cells. We show that glutamine deprivation suppresses *de novo* hexosamine

biosynthesis, which is reciprocally increased upon NAGK deletion. Glutamine deprivation also

results in increased availability of GlcNAc for salvage. *NAGK* expression is elevated in human

PDA tumors, and NAGK deficiency suppresses GlcNAc salvage in cells and tumor growth in

mice.

This work raises several key questions for future investigation. First, the sources of GlcNAc

salvaged by NAGK remain to be fully elucidated. GlcNAc may be derived from recycling of

GlcNAc following O-GlcNAc removal or breakdown of glycoconjugates. Additionally, GlcNAc

may be recovered from the environment. Nutrient scavenging via macropinocytosis is a key

feature of PDA (Commisso et al., 2013; Kamphorst et al., 2015). Macropinocytosis has mostly

been associated with scavenging of protein to recover amino acids, but lysosomal break down

of glycoproteins may also release sugars including GlcNAc. Additionally, ECM components,

including hyaluronic acid (HA), which is a polymer of GlcNAc and glucuronic acid disaccharide

units, may be additional sources of GlcNAc for salvage in the tumor microenvironment. Indeed,

in a manuscript co-submitted with this one, Kim, Halbrook, and colleagues identify HA as a

major source of scavenged GlcNAc (Kim et al., 2020). Our manuscript and the Kim, Halbrook et

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

al manuscript together indicate that NAGK may take on a heightened importance in the context

of high GlcNAc availability and nutrient deprivation, a situation that is likely to occur within the

tumor microenvironment.

Further, the key fates of UDP-GlcNAc that support tumor growth remain to be elucidated.

Sufficient UDP-GlcNAc is required for protein glycosylation to maintain homeostasis and

prevent ER stress, particularly in a rapidly dividing cell. Additionally, a wide range of cancers

exhibit elevated O-GlcNAc, which could contribute to driving pro-tumorigenic transcriptional and

signaling programs. In PDA specifically, the glycan CA19-9 is currently used as a biomarker for

disease progression and recent studies point to a functional role for CA19-9 in tumorigenesis

(Engle et al., 2019). UDP-GlcNAc is also required for HA synthesis, which is present in low

amounts in normal pancreas but increases in PanIN lesions and PDA (Provenzano et al., 2012).

PDA cells are capable of producing HA in vitro (Mahlbacher et al., 1992). Depletion of

fibroblasts in an autochthonous PDA mouse model results in a decrease in collagen I but not

HA in the tumor microenvironment, indicating that HA must be generated by another cell type,

possibly the tumor cells themselves (Özdemir et al., 2014). Previous studies demonstrated that

treatment of PDA with exogenous hyaluronidase can increase vascularization and improve drug

delivery to the tumor (Jacobetz et al., 2013; Provenzano et al., 2012), although a phase III

clinical trial reported no improvement in overall patient survival when combining pegylated

hyaluronidase with nab-paclitaxel plus gemcitabine (Cutsem et al., 2020). Recently, it was

shown that inhibiting the HBP by treatment with 6-diazo-5-oxo-l-norleucine (DON) depletes HA

and collagen in an orthotopic mouse model. DON treatment also increased CD8 T-cell

infiltration into the tumor, sensitizing the tumor to anti-PD1 therapy (Sharma et al., 2020). Thus,

targeting the HBP holds promise for improving the efficacy of other therapeutics. The findings of

the current study suggest that in addition to de novo hexosamine synthesis, targeting of

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

hexosamine salvage warrants further investigation in terms of potential for therapeutic

intervention. Of note, an inhibitor targeting PGM3, which converts GlcNAc-6-P to GlcNAc-1-P

and is thus required for both de novo UDP-GlcNAc synthesis and GlcNAc recycling, showed

efficacy in treating gemcitabine-resistant patient-derived xenograft PDA models (Ricciardiello et

al., 2020), as well as in breast cancer xenografts (Ricciardiello et al., 2018).

Finally, almost nothing is currently known about the role of NAGK and GlcNAc salvage in

normal physiology. Even in non-cancerous IL-3-dependent hematopoietic cells, a substantial

proportion of the UDP-GlcNAc pool remains unlabeled from ^{13}C-glucose (Wellen et al., 2010),

suggesting that salvage may contribute to UDP-GlcNAc pools in a variety of cell types.

However, while GFPT1 is required for embryonic development in mice, NAGK knockout mouse

embryos are viable (Dickinson et al., 2016). NAGK deficiency has not yet been characterized in

postnatal or adult mice. Perhaps GlcNAc salvage is dispensable when nutrients are available

and cells are not dividing, as in most healthy tissues. However, in a tumor, in which cells are

proliferating and nutrients are spread thin, NAGK and GlcNAc salvage may become more

important in feeding UDP-GlcNAc pools. Related questions include elucidating the mechanisms

regulating NAGK gene expression and putative post-translational modification, as well as

understanding the role of GALK2 in hexosamine salvage.

In sum, we report a key role for NAGK in feeding UDP-GlcNAc pools in PDA cells and in

supporting xenograft tumor growth. Further investigation will be needed to elucidate the

physiological functions of NAGK, as well as the mechanisms through which it supports tumor

growth and its potential role in modulating therapeutic responses.

Funding Sources: This work was supported by R01CA174761 and R01CA228339 to K.E.W.

This work was also funded in part under a grant with the Pennsylvania Department of Health to

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

K.E.W. and I.A.B. The Department specifically disclaims responsibility for any analyses,

interpretations, or conclusions. I.A.B. acknowledges support of NIH Grants P30ES013508 and

P30CA016520. J.B. acknowledges support of NIH Grant R01CA046595. S.L.C. received

support from T32CA115299 and F31CA217070, as well as from a Patel Family Scholar Award.

H.A. was supported by post-doctoral fellowship K00CA212455. T.T. is supported by the

National Cancer Institute through pre-doctoral fellowship F31CA243294 and acknowledges the

Blavatnik Family for a predoctoral fellowship. L.I. is supported by T32 GM-07229 and T32

CA115299. S.T. is supported by the American Diabetes Association through post-doctoral

fellowship 1-18-PDF-144. Funding sources were not involved in study design, data collection

and interpretation, or the decision to submit the work for publication.

Conflict of interest: I.A.B. is a founder of Proteoform Bio and a paid consultant for Calico,

Chimerix, PTC Therapeutics, Takeda Pharmaceuticals, and Vivo Capital.

References

Akella, N.M., Ciraku, L., and Reginato, M.J. (2019). Fueling the fire: Emerging role of the

hexosamine biosynthetic pathway in cancer. BMC Biol. *17*, 1–14.

Caldwell, S.A., Jackson, S.R., Shahriari, K.S., Lynch, T.P., Sethi, G., Walker, S., Vosseller, K.,

and Reginato, M.J. (2010). Nutrient sensor O-GlcNAc transferase regulates breast cancer

tumorigenesis through targeting of the oncogenic transcription factor FoxM1. Oncogene *29*,

2831–2842.

Chen, R., Lai, L.A., Sullivan, Y., Wong, M., Wang, L., Riddell, J., Jung, L., Pillarisetty, V.G.,

Brentnall, T.A., and Pan, S. (2017). Disrupting glutamine metabolic pathways to sensitize

gemcitabine-resistant pancreatic cancer. Sci. Rep. *7*, 1–14.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Commisso, C., Davidson, S.M., Soydaner-Azeloglu, R.G., Parker, S.J., Kamphorst, J.J., Hackett, S., Grabocka, E., Nofal, M., Drebin, J.A., Thompson, C.B., et al. (2013). Macropinocytosis of protein is an amino acid supply route in Ras-transformed cells. Nature *497*, 633–637.

Cutsem, E. Van, Tempero, M.A., Sigal, D., Oh, D., Fazio, N., Macarulla, T., Hitre, E., Hammel, P., Hendifar, A., Bates, S., et al. (2020). Randomized Phase III Trial of Pegvorhyaluronidase Alfa With Nab-Paclitaxel Plus Gemcitabine for Patients With Hyaluronan-High Metastatic Pancreatic Adenocarcinoma abstract. J. Clin. Oncol.

Denzel, M.S., and Antebi, A. (2015). Hexosamine pathway and (ER) protein quality control. Curr. Opin. Cell Biol. *33*, 14–18.

Dickinson, M.E., Flenniken, A.M., Ji, X., Teboul, L., Wong, M.D., White, J.K., Meehan, T.F., Weninger, W.J., Westerberg, H., Adissu, H., et al. (2016). High-throughput discovery of novel developmental phenotypes. Nature *537*, 508–514.

Doench, J.G., Fusi, N., Sullender, M., Hegde, M., Vaimberg, E.W., Donovan, K.F., Smith, I., Tothova, Z., Wilen, C., Orchard, R., et al. (2016). Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat. Biotechnol. *34*, 184–191.

Edgar, R., Domrachev, M., and Lash, A.E. (2002). Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Res. *30*, 207–210.

Engle, D.D., Tiriac, H., Rivera, K.D., Pommier, A., Oni, T.E., Alagesan, B., Lee, E.J., Yao, M.A., Lucito, M.S., Spielman, B., et al. (2019). The Glycan CA19-9 Promotes Pancreatitis and Pancreatic Cancer in Mice. *364*, 1156–1162.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Ferrer, C.M., Lu, T.Y., Bacigalupa, Z.A., Katsetos, C.D., Sinclair, D.A., and Reginato, M.J.

(2017). O-GlcNAcylation regulates breast cancer metastasis via SIRT1 modulation of FOXM1

pathway. Oncogene *36*, 559–569.

Granovsky, M., Fata, J., Pawling, J., Muller, W.J., Khokha, R., and Dennis, J.W. (2000).

Suppression of tumor growth and metastasis in Mgat5-deficient mice. Nat. Med. *6*, 306–312.

Gu, Y., Mi, W., Ge, Y., Liu, H., Fan, Q., Han, C., Yang, J., Han, F., Lu, X., and Yu, W. (2010).

GlcNAcylation plays an essential role in breast cancer metastasis. Cancer Res. *70*, 6344–6351.

Guillaumond, F., Leca, J., Olivares, O., Lavaut, M.-N.N., Vidal, N., Berthezène, P., Dusetti, N.J.,

Loncle, C., Calvo, E., Turrini, O., et al. (2013). Strengthened glycolysis under hypoxia supports

tumor symbiosis and hexosamine biosynthesis in pancreatic adenocarcinoma. Proc. Natl. Acad.

Sci. U. S. A. *110*, 3919–3924.

Guo, H., Zhang, B., Nairn, A. V., Nagy, T., Moremen, K.W., Buckhaults, P., and Pierce, M.

(2017). O-Linked N-Acetylglucosamine (O-GlcNAc) Expression Levels Epigenetically Regulate

Colon Cancer Tumorigenesis by Affecting the Cancer Stem Cell Compartment via Modulating

Expression of Transcriptional Factor MYBL1. J. Biol. Chem. *292*, 4123–4137.

Guo, J.Y., Chen, H.Y., Mathew, R., Fan, J., Strohecker, A.M., Karsli-Uzunbas, G., Kamphorst,

J.J., Chen, G., Lemons, J.M.S., Karantza, V., et al. (2011). Activated Ras requires autophagy to

maintain oxidative metabolism and tumorigenesis. Genes Dev. *25*, 460–470.

Guo, L., Worth, A.J., Mesaros, C., Snyder, N.W., Jerry, D., and Blair, I.A. (2016).

Diisopropylethylamine/hexafluoroisopropanol-mediated ion- pairing UHPLC-MS for phosphate

and carboxylate metabolite analysis: utility for studying cellular metabolism. Rapid Commun

Mass Spectrom. *30*, 1835–1845.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Halbrook, C.J., and Lyssiotis, C.A. (2017). Employing Metabolism to Improve the Diagnosis and

Treatment of Pancreatic Cancer. Cancer Cell *31*, 5–19.

Housley, M.P., Rodgers, J.T., Udeshi, N.D., Kelly, T.J., Shabanowitz, J., Hunt, D.F., Puigserver,

P., and Hart, G.W. (2008). O-GlcNAc regulates FoxO activation in response to glucose. J. Biol.

Chem. *283*, 16283–16292.

Jacobetz, M.A., Chan, D.S., Neesse, A., Bapiro, T.E., Cook, N., Frese, K.K., Feig, C.,

Nakagawa, T., Caldwell, M.E., Zecchini, H.I., et al. (2013). Hyaluronan impairs vascular function

and drug delivery in a mouse model of pancreatic cancer. Gut *62*, 112–120.

Kamphorst, J.J., Nofal, M., Commisso, C., Hackett, S.R., Lu, W., Grabocka, E., Vander Heiden,

M.G., Miller, G., Drebin, J.A., Bar-Sagi, D., et al. (2015). Human Pancreatic Cancer Tumors Are

Nutrient Poor and Tumor Cells Actively Scavenge Extracellular Protein. Cancer Res. *75*, 544–

553.

Kim, P.K., Halbrook, C.J., Kerk, S.A., Wisner, S., Kremer, D., Sajjakulnukit, P., Hou, S.W.,

Thurston, G., Anand, A., Yan, L., et al. (2020). Hyaluronic Acid Fuels Pancreatic Cancer

Growth. BioRxiv.

Lau, K.S., Partridge, E.A., Grigorian, A., Silvescu, C.I., Reinhold, V.N., Demetriou, M., and

Dennis, J.W. (2007). Complex N-Glycan Number and Degree of Branching Cooperate to

Regulate Cell Proliferation and Differentiation. Cell *129*, 123–134.

Li, D., Li, Y., Wu, X., Li, Q., Yu, J., Gen, J., and Zhang, X.-L. (2008). Knockdown of Mgat5

Inhibits Breast Cancer Cell Growth with Activation of CD4 + T Cells and Macrophages . J.

Immunol. *180*, 3158–3165.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Lynch, T.P., Ferrer, C.M., Jackson, S.R.E., Shahriari, K.S., Vosseller, K., and Reginato, M.J.

(2012). Critical role of O-linked β-N-acetylglucosamine transferase in prostate cancer invasion,

angiogenesis, and metastasis. J. Biol. Chem. *287*, 11070–11081.

Lyssiotis, C.A., and Kimmelman, A.C. (2017). Metabolic Interactions in the Tumor

Microenvironment. Trends Cell Biol. *27*, 863–875.

Mahlbacher, V., Sewing, A., Elsässer, H.P., and Kern, H.F. (1992). Hyaluronan is a secretory

product of human pancreatic adenocarcinoma cells. Eur. J. Cell Biol. *58*, 28—34.

Mereiter, S., Balmaña, M., Campos, D., Gomes, J., and Reis, C.A. (2019). Glycosylation in the

Era of Cancer-Targeted Therapy: Where Are We Heading? Cancer Cell *36*, 6–16.

Moloughney, J.G.G., Kim, P.K.K., Vega-cotto, N.M.M., Wu, C.-C., Zhang, S., Adlam, M., Lynch,

T., Chou, P.-C., Rabinowitz, J.D.D., Werlen, G., et al. (2016). mTORC2 Responds to Glutamine

Catabolite Levels to Modulate the Hexosamine Biosynthesis Enzyme GFAT1. Mol. Cell *63*,

811–826.

Moseley, H.N.B., Lane, A.N., Belshoff, A.C., Higashi, R.M., and Fan, T.W.M. (2011). A novel

deconvolution method for modeling UDP-N-acetyl-D-glucosamine biosynthetic pathways based

on 13C mass isotopologue profiles under non-steady-state conditions. BMC Biol. *9*.

Munkley, J. (2019). The glycosylation landscape of pancreatic cancer (Review). Oncol. Lett. *17*,

2569–2575.

Özdemir, B.C., Pentcheva-Hoang, T., Carstens, J.L., Zheng, X., Wu, C.C., Simpson, T.R.,

Laklai, H., Sugimoto, H., Kahlert, C., Novitskiy, S. V., et al. (2014). Depletion of carcinoma-

associated fibroblasts and fibrosis induces immunosuppression and accelerates pancreas

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

cancer with reduced survival. Cancer Cell *25*, 719–734.

Park, S.Y., Kim, H.S., Kim, N.H., Ji, S., Cha, S.Y., Kang, J.G., Ota, I., Shimada, K., Konishi, N.,

Nam, H.W., et al. (2010). Snail1 is stabilized by O-GlcNAc modification in hyperglycaemic

condition. EMBO J. *29*, 3787–3796.

Pei, H., Li, L., Fridley, B.L., Jenkins, G.D., Kalari, K.R., Lingle, W., Petersen, G., Lou, Z., and

Wang, L. (2009). FKBP51 Affects Cancer Cell Response to Chemotherapy by Negatively

Regulating Akt. Cancer Cell *16*, 259–266.

Provenzano, P.P., Cuevas, C., Chang, A.E., Goel, V.K., Von Hoff, D.D., and Hingorani, S.R.

(2012). Enzymatic Targeting of the Stroma Ablates Physical Barriers to Treatment of Pancreatic

Ductal Adenocarcinoma. Cancer Cell *21*, 418–429.

Rahib, L., Smith, B.D., Aizenberg, R., Rosenzweig, A.B., Fleshman, J.M., and Matrisian, L.M.

(2014). Projecting cancer incidence and deaths to 2030: The unexpected burden of thyroid,

liver, and pancreas cancers in the united states. Cancer Res. *74*, 2913–2921.

Ricciardiello, F., Votta, G., Palorini, R., Raccagni, I., Brunelli, L., Paiotta, A., Tinelli, F., D'Orazio,

G., Valtorta, S., De Gioia, L., et al. (2018). Inhibition of the Hexosamine Biosynthetic Pathway

by targeting PGM3 causes breast cancer growth arrest and apoptosis. Cell Death Dis. *9*.

Ricciardiello, F., Gang, Y., Palorini, R., Li, Q., Giampà, M., Zhao, F., You, L., La Ferla, B., De

Vitto, H., Guan, W., et al. (2020). Hexosamine pathway inhibition overcomes pancreatic cancer

resistance to gemcitabine through unfolded protein response and EGFR-Akt pathway

modulation. Oncogene 4103–4117.

Ryczko, M.C., Pawling, J., Chen, R., Abdel Rahman, A.M., Yau, K., Copeland, J.K., Zhang, C.,

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Surendra, A., Guttman, D.S., Figeys, D., et al. (2016). Metabolic Reprogramming by

Hexosamine Biosynthetic and Golgi N-Glycan Branching Pathways. Sci. Rep. *6*, 23043.

Sanjana, N.E., Shalem, O., and Zhang, F. (2014). Sanjana, Shalem et Zhang. Nat. Med. *11*,

783–784.

Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of

image analysis. Nat. Methods *9*, 671–675.

Sharma, N.S., Saluja, A., Banerjee, S., Sharma, N.S., Gupta, V.K., Garrido, V.T., Hadad, R.,

Durden, B.C., Kesh, K., Giri, B., et al. (2020). Targeting tumor-intrinsic hexosamine biosynthesis

sensitizes pancreatic cancer to anti-PD1 therapy Graphical abstract Find the latest version :

Targeting tumor-intrinsic hexosamine biosynthesis sensitizes pancreatic cancer to anti-PD1

therapy. *130*, 451–465.

Steenackers, A., Olivier-Van Stichelen, S., Baldini, S.F., Dehennaut, V., Toillon, R.A., Le

Bourhis, X., El Yazidi-Belkoura, I., and Lefebvre, T. (2016). Silencing the nucleocytoplasmic O-

GlcNAc transferase reduces proliferation, adhesion, and migration of cancer and fetal human

colon cell lines. Front. Endocrinol. (Lausanne). *7*, 1–11.

Swamy, M., Pathak, S., Grzes, K.M., Damerow, S., Sinclair, L. V, van Aalten, D.M.F., and

Cantrell, D.A. (2016). Glucose and glutamine fuel protein O-GlcNAcylation to control T cell self-

renewal and malignancy. Nat. Immunol. *17*, 1–11.

Tang, Z., Kang, B., Li, C., Chen, T., and Zhang, Z. (2019). GEPIA2: an enhanced web server for

large-scale expression profiling and interactive analysis. Nucleic Acids Res. *47*, W556–W560.

Taylor, R.P., Parker, G.J., Hazel, M.W., Soesanto, Y., Fuller, W., Yazzie, M.J., and McClain,

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

D.A. (2008). Glucose deprivation stimulates O-GlcNAc modification of proteins through up-
regulation of O-linked N-acetylglucosaminyltransferase. J. Biol. Chem. *283*, 6050–6057.

Trefely, S., Ashwell, P., and Snyder, N.W. (2016). FluxFix: Automatic isotopologue
normalization for metabolic tracer analysis. BMC Bioinformatics *17*, 1–8.

Wellen, K.E., Lu, C., Mancuso, A., Lemons, J.M.S.S., Ryczko, M., Dennis, J.W., Rabinowitz,
J.D., Coller, H.A., and Thompson, C.B. (2010). The hexosamine biosynthetic pathway couples
growth factor-induced glutamine uptake to glucose metabolism. Genes Dev. *24*, 2784–2799.

Wyant, G.A., Abu-Remaileh, M., Frenkel, E.M., Laqtom, N.N., Dharamdasani, V., Lewis, C.A.,
Chan, S.H., Heinze, I., Ori, A., and Sabatini, D.M. (2018). NUFIP1 is a ribosome receptor for
starvation-induced ribophagy. Science (80-.). *360*, 751–758.

Ying, H., Kimmelman, A.C., Lyssiotis, C.A., Hua, S., Chu, G.C., Fletcher-Sananikone, E.,
Locasale, J.W., Son, J., Zhang, H., Coloff, J.L., et al. (2012). Oncogenic kras maintains
pancreatic tumors through regulation of anabolic glucose metabolism. Cell *149*, 656–670.

Zhou, X., Chen, H., Wang, Q., Zhang, L., and Zhao, J. (2011). Knockdown of Mgat5 inhibits
CD133+ human pulmonary adenocarcinoma cell growth in vitro and in vivo. Clin. Invest. Med.
34, E155-62.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Supplemental Figure 1: LPHA binding detects MGAT5-dependent glycans.
A) Diagram of LPHA binding. LPHA recognizes specifically the β1-6 linkage established by MGAT5. **B)** LPHA binding on MGAT5 knockout cells isolated from a KPCY tumor (Li, Byrne et al., 2018), representative flow plot and quantification. Graph shows MFI relative to control cells. Statistical significance was calculated by unpaired t-test. **C)** Representative flow plots for LPHA binding graphed in Fig. 1D, E on PANC-1 cells in low glucose and low glutamine. **D-E)** LPHA binding in PDA cells in low nutrients. Cells were incubated in the indicated concentrations of serum (D) or oxygen (E) for 48 hours and then analyzed by flow cytometry. Graph shows MFI relative to control condition. **F)** Western blot for O-GlcNAc and PARP in MIA PaCa-2 cells cultured in the indicated concentrations of glutamine with or without TMG for the indicated time. Statistical significance was calculated by one-way ANOVA (D) and unpaired t-test (E). Panels B) - E) are representative of at least two independent experimental replicates. *, p ≤ 0.05; **, p ≤ 0.01; **, p ≤ 0.001.

646   This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Supplemental Figure S2.1: De novo hexosamine synthesis is suppressed in low glutamine conditions.
A) Metabolite measurements in MIA PaCa-2 cells after culture for 48 hours in 0.05 mM glutamine. Quantification is normalized to 4 mM glutamine condition. Statistical significance was calculated by unpaired t-test. Mean +/- SEM of four biological replicates is represented. **B)** Measurement of HBP metabolites in PANC-1 cells after culture for 48 hours in 5 mM or 0.1 mM glucose. Mean +/- SEM of four (5 mM) or five (0.1 mM) biological replicates is represented. Statistical significance was calculated by unpaired t-test comparing each low glucose condition to the 10 mM glucose control in each experiment. **C)** Measurement of 13C glucose incorporation into UDP-GlcNAc in high and low glutamine. Mean +/- SEM of three biological replicates is represented. **D)** Measurement of 13C glucose incorporation into GlcNAc-P in high and low glutamine. Mean +/- SEM of three biological replicates is represented. For (C-D), components of molecule labeled in each isotopologue are indicated, with dots representing the number of carbons; highlighted panels identify glucosamine ring labeling. **E)** Measurement of 13C glucose into fructose-6-phosphate in high and low glutamine. Mean +/- SEM of three biological replicates is represented. *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$.

648   This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Supplemental Figure S2.2: Uridine is salvaged in glutamine restricted conditions.
A) Measurement of incorporation of 13C glucose into uridine triphosphate (UTP) in high and low glutamine. Mean +/- SEM of three biological replicates is represented. **B)** Measurement of metabolites in PANC-1 cells after transfection with the indicated siRNAs and incubation in 4 mM (+) or 0.05 mM (-) glutamine for 48 hours. Mean +/- SEM of four biological replicates is represented. **C)** Gene expression of the indicated genes in PANC-1 cells after transfection with the indicated siRNAs and incubation in 4 mM (+) or 0.05 mM (-) glutamine for 48 hours. Mean +/- SEM of four biological replicates is represented. **D)** Measurement of incorporation of 13C glucose into UDP-GlcNAc in PANC-1 cells transfected with the indicated siRNAs and incubated in 4 mM or 0.05 mM glutamine for 48 hours. Mean +/- SEM of three biological replicates is represented. **E)** Gene expression of the indicated genes in PANC-1 cells after transfection with the indicated siRNAs and incubation in 4 mM (+) or 0.05 (-)mM glutamine for 48 hours. Mean +/- SEM of three biological replicates is represented. *, p ≤ 0.05; **, p ≤ 0.01; ***, p ≤ 0.001.

650   This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Supplemental Figure S2.3: NAGK protein expression does not increase in low glutamine despite increase in NAGK gene expression.
A) Gene expression of the indicated genes in PDA cell lines after incubation in 4 mM or 0.05 mM glutamine for 48 hours. Mean +/- SEM of three biological replicates is represented. Statistical significance was determined by unpaired t-test. **B)** Gene expression of the indicated genes in PDA cell lines after incubation in 10 mM or 0.1 mM glucose for 48 hours. Mean +/- SEM of three biological replicates is represented. Statistical significance was determined by unpaired t-test. **C)** Western blot for NAGK in low nutrient conditions. PANC-1 and MIA PaCa-2 cells were cultured in the indicated concentrations of glucose and glutamine for 48 hours. **D)** Western blot for NAGK in low nutrient conditions. PANC-1 cells were cultured in the indicated glutamine concentrations and lysates were prepared with or without phosphatase inhibitors. A band shift is observed for NAGK in lysates with Na_3VO_4.
*, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$.

652   This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Supplemental Figure S3.1: NAGK knockout cells show increased enrichment of 13C glucose into hexosamine intermediates.
A) Gene expression of NAGK in PANC-1 NAGK CRISPR cells. Mean +/- SEM of three biological replicates is represented. Statistical significance was determined by one-way ANOVA. **B)** Western blot for NAGK and HSP60 in PANC-1 (left) and MIA PaCa-2 (right) NAGK CRISPR knockout cells. **C)** Measurement of incorporation of 13C GlcNAc labeled on the acetyl group into UDP-GlcNAc in PANC-1 NAGK knockout cells. Cells were incubated with 10 mM 13C GlcNAc for 3 hours. Mean +/- SEM of three biological replicates is represented. Statistical significance was determined by unpaired t-test comparing siCon and siGALK2 for each clone. **D)** Measurement of incorporation of 13C glucose into GlcNAc-P in NAGK knockout cells. Mean +/- SEM of three biological replicates is represented. **E)** Measurement of incorporation of 13C glucose into UDP-GlcNAc in NAGK knockout cells. Mean +/- SEM of three biological replicates is represented. For (D-E), components of molecule labeled in each isotopologue are indicated, with dots representing the number of carbons; highlighted panels identify glucosamine ring labeling. *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.01$.

654   This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Supplementary Figure S3.2: HBP metabolites in NAGK KO cells.
A) Gene expression of NAGK in PANC-1 cells transfected with shRNA against NAGK. One biological sample is represented. **B)** Incorporation of 13C glucose into the glucosamine ring of UDP-GlcNAc in PANC-1 cells transfected with the indicated shRNAs targeting NAGK. Mean +/- SEM of three biological replicates is represented. **C)** Measurement of HBP metabolites in MIA PaCa-2 NAGK knockout cells cultured in 4 mM glutamine. Mean +/- SEM of four biological replicates is represented. Statistical significance was calculated by unpaired t-test comparing the mean of the three CRISPR clones and the empty vector control. **D)** Measurement of UDP-GlcNAc in PANC-1 NAGK knockout cells. Mean +/- SEM of four biological replicates is represented. Statistical significance was calculated by one-way ANOVA comparing each knockout to the EV control. *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$.

656   This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."

Supplementary Figure S4: NAGK knockout tumors show increased LPHA binding.
A) Soft agar colony formation in NAGK knockout and control cells. Mean +/- SEM of three biological replicates is represented. **B)** Gene expression data plotted using GEPIA2 comparing TCGA pancreatic cancer samples with TCGA normal pancreas samples and GTEx data. Differential analysis between tumor and normal tissues was analyzed by one-way ANOVA. *, $p < 0.05$. **C)** Tumor volume of NAGK knockout tumors at earlier time points in the experiment. Initial growth of NAGK knockout cells in vivo was comparable to control cells, but over time NAGK knockout tumors either stopped growing or regressed. **D)** Western blots for NAGK and α-SMA, and blot probed for LPHA binding, in whole tumor lysate samples. Blot for HSP60 as loading control.

This manuscript has been co-submitted with Kim et al. (2020), "Pancreatic Cancers Scavenge Hyaluronic Acid to Support Growth."