A spatial model of the plant circadian clock reveals design principles for coordinated timing under noisy environments

Short title: Coordinated timing through clock coupling

Mark Greenwood[1,2], Isao T. Tokuda[3*], James C.W. Locke[1,4*]

[1]Sainsbury Laboratory, University of Cambridge, Cambridge, UK

[2]Department of Biochemistry, University of Cambridge, Cambridge, UK

[3]Department of Mechanical Engineering, Ritsumeikan University, Kusatsu, Japan

[4]Microsoft Research, Cambridge, UK

[*]For correspondence: james.locke@slcu.cam.ac.uk or isao@fc.ritsumei.ac.jp

Abstract

Individual plant cells possess a genetic network, the circadian clock, that times internal processes to the day-night cycle. Mathematical models of the clock network have driven a mechanistic understanding of the clock in plants. However, these models are typically either 'whole plant' models that ignore tissue or cell type specific clock behavior, or 'phase only' models that do not include clock network components explicitly. It is increasingly clear that in order to reveal the design principles of the plant circadian clock, clock network models must address spatial differences. This is because complex spatial behaviours have been observed in tissues and cells in plants, including period and phase differences between cells and spatial waves of gene expression between organs. Here, we implement an up to date clock network model on a spatial template of the plant. In our model, the sensitivity to light inputs varies across the plant, and cells communicate their clock timing locally via the levels of core clock mRNA levels by cell-to-cell coupling. We found that differences in sensitivities to environmental input in the model can explain the experimentally observed differences in clock periods in different

organs, and we show using the model that a plausible coupling mechanism can generate the

experimentally observed waves in clock gene expression across the plant. We then examined

what features of the plant circadian system allow it to keep time under noisy light-dark (LD)

cycles. We found that differences in sensitivity to light can allow regional flexibility in phase even

under LD cycles, whilst local cell-to-cell coupling minimized variability in clock rhythms in

neighboring cells. Thus, local sensitivity to environmental inputs combined with cell-to-cell

coupling allows for flexible yet robust circadian timing under noisy environments.

Abbreviations: *CCA1*, *CIRCADIAN CLOCK ASSOCIATED 1*; CV, coefficient of variation; DD,

constant dark; *ELF4*, *EARLY FLOWERING 4*; LD, light-dark; *LHY*, *LATE ELONGATED*

HYPOCOTYL; LL, constant light; *LUC*, *LUCIFERASE*; *LUX*, *LUX ARRHYTHMO*; *PRR*,

PSEUDO-RESPONSE REGULATOR; *TOC1*, *TIMING OF CAB EXPRESSION 1*; YFP, yellow

fluorescent protein.

Introduction

The circadian clock is a 24 h genetic oscillator found in many organisms. The clock consists of a

circuit of interlocking feedback loops of mRNAs and proteins that generate daily oscillations in

circuit component levels. Signals from the environment align the timing of these oscillations to

the day-night cycle [1]. Once set, circadian clocks act as an internal timing signal, allowing

biological processes to anticipate the external environmental cycles. The clock modulates a

diverse range of processes in plants, including cell division, tissue growth, flowering time, and

scent emission [2–5]. Altogether this daily timing provides a significant fitness advantage to the

plant [6,7].

Individual plant cells possess a robust circadian clock [8]. However, substantial differences in

the period and phase of clock rhythms across the plant have been observed. Time lapse

imaging experiments with luciferase and fluorescent reporter genes have shown that rhythms in

core clock genes oscillate with different speeds in different organs under constant light (LL) [8–

13]. Further, experiments under a range of conditions have shown that differences in clock

speed and phase can be caused by organs having different sensitivity to environmental signals

[12–15]. Differences in the clock network between tissues may also contribute to generating

differences in rhythms across the plant, as although the clock genes are broadly expressed [13],

some are tissue enriched [16], and mutations can affect organs differently [11,15,17].

The observed differences in clock rhythms across the plant raises the question of how clocks in

different cells and tissues remain coordinated with each other. One mechanism would be for

cells to communicate their timing with their neighbors. High resolution experiments have

measured or inferred local coupling of clock rhythms between cells [8,11,14,16,18–20], and

local coupling can drive spatial waves of clock gene expression across the plant [8,14,18–20].

Longer-distance coupling between clocks is also possible. For example, molecular signals

communicate circadian temperature information from the shoot to the root [11,21], and light

information may be piped down the stem to entrain the root [22].

Mathematical modeling has played a crucial role in gaining a mechanistic understanding of plant

circadian clocks. Models of the network have increased in complexity over time in parallel with

the growing number of experiments [23–28]. Recently, these detailed molecular models have

been used to probe the differences between the shoot and root clock [13]. However, the

coupling of clocks between cells was not considered. For this, more computationally tractable

models of the clock network are necessary. Reduced models of the network have already been

constructed that capture many of the features of the single cell clock dynamics [29–32].

However, these models have not been applied to study spatial dynamics. Instead, 'phase only'

models that lack any genetic network information and only consider the phases of individual

cellular rhythms have been preferred [8,18–20]. Although these models allow the simulation of

general oscillatory behavior, owing to their simplicity they are unsuitable for investigating

molecular mechanisms of the clock.

Recently, a 'phase only' Kuramoto model [33] was used to propose a mechanism for whole-

plant coordination of clocks [8,14]. In order to match experimentally measured rhythms, the

model fixed clock periods to different speeds in each region of the plant. It assumed faster

rhythms in the cotyledons, hypocotyl, and root tip, and slower rhythms in the rest of the root, as

observed experimentally. With these periods fixed, cells were allowed to communicate clock

phase through local cell-to-cell coupling. With these assumptions, simulations of the model

generated waves of clock gene expression within and between organs, as observed

experimentally [8,14]. Thus, local cell-to-cell coupling could enable coordination between organs

in plants.

Multiple questions remain about how the plant clock coordinates rhythms that cannot be

addressed using a 'phase only' model. For example, how are the periods set differently in

different parts of the plant? What molecular mechanisms allow the coupling of clock rhythms

from cell-to-cell? How can the plant clock network 'filter' both internal and environmental noise

to robustly entrain to the environment? To begin to address these questions, in this work we

developed a spatial network model of the plant circadian system. We modified a previously

generated simplified network model and implemented it on a multicellular template of a plant. In

our model, the sensitivity to light varies across the plant, and cells communicate via the mRNA

levels of a core clock gene. Simulations capture the organ specific period differences and

spatial waves observed under LL, demonstrating a plausible mechanism of circadian

coordination. We then applied our model to examine how the system keeps time under noisy

light-dark (LD) cycles. We found that regional differences persist even under LD cycles, but cell-

to-cell minimized differences between neighbor cells. Thus, the combination of local sensitivity

to inputs and cell-to-cell coupling allows for coordinated timing in noisy environments.

Results

A locally coupled spatial model of the plant circadian clock network

We first implemented a reduced network model of the *Arabidopsis thaliana* circadian clock [31].

To decrease the complexity of the model, the authors grouped functionally similar genes into

single entities (Fig 1A). The compact network model incorporates known light inputs to the

network and qualitatively recapitulates clock dynamics under both light dark cycles and constant

light. At only 9 equations and 34 parameters the model is also computationally tractable for

spatial simulations. We modified the De Caluwe model to include a repression rather than

activation interaction between *PSEUDO-RESPONSE REGULATOR 9 (PRR9)*/*PRR7* and

CIRCADIAN CLOCK ASSOCIATED 1 (*CCA1*) / *LATE ELONGATED HYPOCOTYL* (*LHY*), as

this has recently been shown experimentally [34] (Methods).

We implemented the modified De Caluwe model on a simplified template of a seedling. The

template consisted of approximately 800 cells, classified into cotyledon, hypocotyl, root, and

root tip regions (Fig 1B and Methods). Although a number of studies have demonstrated local

cell-to-cell coupling between clocks in *A. thaliana* [8,11,14,16,18–20], the identity of the coupling

components are unclear. Initially, to model the coupling (Fig 1C), the level of *CCA1*/*LHY* in one

cell was assumed to be locally coupled to the level of the cell's neighbors. The coupling strength

was set to 2 (Methods). To simulate the variability observed in single cell clock rhythms [8] we

multiplied the level of each mRNA and protein by a time scaling parameter that was randomly

selected from a normal distribution (Methods). The range of this normal distribution was set

differently for each organ to qualitatively match the variability observed experimentally (S1 Fig).

130 **Different light sensitivities can explain organ differences in phase and period**

131 We next attempted to recapitulate in our model the differences in clock period and phase in

132 different organs that have been observed in experiments using a single cell CCA1-YFP reporter

133 [8] and a *GIGANTEA* luciferase reporter [14]. In these experiments, faster rhythms were

134 observed in the cotyledon, hypocotyl, and root tip, with slower rhythms in the rest of the root.

135 We first reanalyzed existing luciferase data [14] and confirmed that these relationships held for

136 several of the core clock genes in our model, *PSEUDO-RESPONSE REGULATOR 9* (*PRR9*),

137 *TIMING OF CAB EXPRESSION 1* (*TOC1*), and *EARLY FLOWERING 4* (*ELF4*) (Fig 2A).

138 Whereas in our previous 'phase only' model, we fixed the periods to be different in each part of

139 the plant, with our spatial network model, we could now investigate what causes the differences

140 in periods. Previously it has been hypothesized that different sensitivities to environmental

141 inputs alter periods across the plant [13,14]. For example, cells in the cotyledon may be fast

142 because they are more sensitive to light. Experimental tests of this, however, are confounded by

143 changes in metabolism and development caused by light [35]. To test this in our model, we

144 simulated differences in sensitivity by setting the intensity of light input in our model to differ

145 depending on the region, and examined whether this can generate the period differences

146 observed across the plant.

147

148 We entrained the cells in our simulations to LD cycles for 4 days before releasing them into LL

149 for a further 6 days and measured the periods, as carried out in previous experiments [14].

150 When assuming high sensitivity to light in the cotyledon (L = 1.35) and hypocotyl (L = 1.10), but

151 lower in the root (L = 0.90) and root tip (L = 1.03), all organs entrained to the LD cycles (S2 Fig).

152 Upon transfer to LL, we were able to generate different periods (Fig 2B) and phases (Fig 2D–F

153 and S3 Fig) in each organ, matching those observed experimentally. Thus, our results revealed

154 that different sensitivities to environmental inputs are sufficient to generate the experimentally

155 observed spatial differences in period and phase across the plant. This is due to higher light

sensitivity causing the clock to run faster in our simulations (S4 Fig), as expected for a diurnal

organism [36].

Local sharing of clock mRNA levels can drive spatial waves of clock gene expression

Previously we observed two waves of clock gene expression, one traveling up, and one down

the root, in *CCA1*, *PRR9* and *GI* reporters [8,14]. These waves could be explained in a phase

only model by local cell-to-cell coupling [8,14,19,20]. We next tested whether a plausible

mechanism for cell-to-cell coupling, sharing of mRNA between cells [37], can recapitulate the

experimental observations. To provide a benchmark, we first analyzed the seedlings carrying

transcriptional reporters for *TOC1*, *PRR9*, and *ELF4* [14] at the sub-tissue level (Methods). As in

previous studies [8,14,18–20], space-time plots revealed spatial waves of gene expression

within and between organs (Fig 3A and S5 Fig). The direction of the waves can be clearly

observed in plots of the final peaks of expression (Fig 3B and S5 Fig). For each gene, the wave

patterns appeared similar, with two waves of clock gene expression in the root (Fig 3C, S5 Fig,

and S1 Video).

We next analyzed simulations at the sub-tissue level (S6 Fig and Methods) to see if our model

captures these spatial dynamics. For each gene, the wave patterns appeared similar to

experiments, traveling from the fast oscillating regions that are more sensitive to light, into the

slower regions with lower sensitivity to light (Fig 3C, D, S5 Fig and S2 Video). These waves

required cell-to-cell coupling through the local sharing of clock mRNA, as we only observed

waves with coupling strengths, J, above approximately 1 (Fig 3E and S7 Fig). Similar simulation

results were obtained using different clock genes as the coupling component, suggesting that

any cell-to-cell sharing of clock components can explain the experimentally observed spatial

dynamics (S8 Fig). We also found that our simulation results were qualitatively similar when

keeping regional differences of light sensitivity and local cell-to-cell coupling, but assuming no

cell-to-cell variability within regions (S9 Fig). Additionally, we ran simulations assuming longer

distance and global coupling. Increasing the local coupling to be between 8 rather than 4

neighbor cells gave qualitatively similar results. However, with global coupling all cells adopted

the same phase at higher coupling strengths, regardless of position in the plant (S10 Fig).

Finally, by setting the light inputs in our locally coupled model to be equal in all regions of the

seedling, we were also able to simulate the loss of waves observed in the light sensing mutant

phyb-9 [14] (S11 Fig). Taken together, these results show that the assumptions of local cell-to-

cell coupling and differential light sensitivity between regions are the key aspects of our model

that allow a match to experimental data.

Local flexibility persists under idealized and noisy LD cycles

Our modeling suggests that different sensitivities to environmental inputs across the plant allows

the clock to be locally flexible under LL, by allowing it to adopt different periods and phases

across the plant even under the same environment. We simulated our model under LD cycles to

investigate whether local flexibility persisted with rhythmic input. We first simulated idealized LD

cycles, where lights are switched fully on during the daytime and off at night (Fig 4A and

Methods). The sensitivity to light during the daytime differed depending on the region, as with LL

simulations, and we first assumed no cell-to-cell coupling ($J = 0$). In the absence of cell-to-cell

coupling, close inspection of the final peaks of expression revealed small differences in the

timing between regions. *PRR5/TOC1* expression in cells of the cotyledon peaked at 132.70 ±

0.50 h (median ± interquartile range), followed by those in the hypocotyl (133.40 ± 0.40 h) and

then the root, with cells at the tip (133.50 ± 1.35 h) peaking before those in the middle region of

the root (134.80 ± 1.20 h; Fig 4B, left and S12 Fig). These phase differences were qualitatively

the same as observed under LL (Fig 2 and 3) but were discontinuous between regions. With the

addition of cell-to-cell coupling, the phase differences between regions were no longer

discontinuous and were instead staggered, causing the emergence of spatial waves (Fig 4B,

middle and right, and S12 Fig). However, phase differences between the regions persisted and

did not reduce in size with increasing strengths of coupling (S12 Fig). For example, there was a

difference between the cotyledon and root of 1.90 ± 0.18 h (median ± interquartile range)

without coupling ($J = 0$), 2.00 ± 0.20 h with coupling ($J = 2$), and 2.00 ± 0.18 h with stronger

coupling ($J = 4$). A similar spatial structure was observed in *CCA1*/*LHY* and *PRR9*/*PRR7*

expression (S12 Fig). However, *ELF4*/*LUX* appeared more synchronized when the peak

coincided with the dark transition, as darkness causes strong repression of *ELF4* expression.

We next simulated more realistic LD cycles containing fluctuations in the light intensity (Fig 4C).

These LD cycles were designed to approximate the differences in input that cells may

experience due to weather or a cell's microenvironment (Methods). Examination of the times of

the final peaks of expression reveals large variation in peak times within regions when cell-to-

cell coupling was absent ($J = 0$; Fig 4D, left). This made any difference in phase between

regions difficult to resolve. However, with increasing strengths of cell-to-cell coupling we

observed a decrease in variability. For example, there was a difference of 5.00 ± 1.05 h (median

± interquartile range) between the earliest and latest peaking cells without coupling ($J = 0$), but

3.10 ± 0.23 h with coupling ($J = 2$), and 3.20 ± 0.15 h with stronger coupling ($J = 4$). This

decrease in cell-to-cell variation revealed an underlying spatial structure (Fig 4D, middle and

right, and S13 Fig), comparable to that observed under idealized LD cycles (Fig 4B, middle and

right, and S12 Fig). Phase differences and spatial waves persisted with increasing strengths of

coupling (S13 Fig). This spatial structure was also observed in expression of *CCA1*/*LHY* and

PRR9/*PRR7*, but not in *ELF4*/LUX expression (S13 Fig). Together this shows that local flexibility

in phase, and spatial waves, persisted under LD cycles when local cell-to-cell coupling was

present.

Cell-to-cell coupling maintains global coordination under noisy light-dark cycles

Simulations under noisy LD cycles showed that whilst local flexibility of regions persisted, cell-

to-cell coupling decreased the phase variation within regions (Fig 4). We further investigated the

function of cell-to-cell coupling by testing whether coupling also affected the global

synchronization of rhythms, or is limited to a local effect on rhythms. To test this, we quantified

the synchrony between all cells within the template seedling using a score for synchronization,

R [38]. The synchronization index, R, is the ratio of the variance of the average signal to the

average variance of the individual cells. When cells are synchronized these variances are equal

to each other and R becomes 1, whereas when desynchronized R becomes 0. For each gene

pair, we observed a high R value under idealized LD cycles, with or without cell-to-cell coupling

(Fig 5, black lines). However, cells were less synchronous under noisy LD cycles (Fig 5, red

lines). Each gene showed a reduced level of synchrony in the absence of cell-to-cell coupling (J

= 0). *PRR9/PRR7* was the most affected by the noise, with a synchronization index of 0.97 ±

0.002 (mean ± standard deviation) under idealized LD and 0.61 ± 0.008 under noisy LD cycles

(Fig 5B). This is likely because *PRR9/PRR7* synthesis is most strongly affected by light (Table

1).

We observed an increase in global synchronization under noisy LD cycles as coupling strengths

increased (Fig 5). At higher strengths of coupling (J > 2), for *CCA1/LHY* and *PRR5/TOC1* the

synchronization index under noisy LD cycles approximately equaled that under idealized LD

cycles (Fig 5A and C). For *PRR9/PRR7* and *ELF4/LUX* expression, the synchronization index

increased with increasing strength of coupling, but did not reach the level observed under

idealized LD cycles for all strengths tested (Fig 5B and D). Oscillations of protein expression

also increased in synchrony with cell-to-cell coupling (S14 Fig). Thus, local cell-to-cell coupling

improves the global coordination of circadian timing under noisy light-dark cycles, whilst still

allowing differences in clock timing between regions.

Discussion

Here we develop a spatial network model for the plant clock and use it to examine the design principles of clock coordination in plants. This simple multicellular model, which assumes regional differences in sensitivity to light and cell-to-cell coupling through the local sharing of clock mRNA levels, successfully simulates the period differences and phase waves across the plant observed under LL in previous time lapse experiments. Our simulations predict that phase flexibility persists under LD cycles, with cell-to-cell coupling reducing cell-to-cell differences but still allowing regional flexibility. We therefore find that the plant circadian clock system can combine regional differences in environmental signaling with cell-to-cell coupling to enable robust, yet flexible, circadian timing under noisy environmental cycles.

Previous models of the plant circadian clock have proven important for understanding aspects of plant physiology, including starch metabolism and flowering [39,40]. However, it was recently shown that the clock controls physiology in a tissue-specific manner. For example, clocks in the epidermis regulate growth, whereas those in the vasculature regulate flowering [41]. In future, our spatial model could be extended to investigate circadian control of physiology at the cell and tissue level. An interesting example will be cell division, which is regulated by the clock [2], and occurs in meristematic regions of the plant [42]. It will be interesting to see if, and how, flexibility of the circadian system impacts tightly controlled phenotypes such as this.

Including noise in LD cycles represents a step towards understanding how the plant circadian system works under natural conditions. Our simulations under noisy LD cycles reveal that local cell-to-cell coupling allows a balance between ensuring cells are coordinated locally even under a noisy environment, whilst still allowing regional differences that would not be possible with a global coupling mechanism. However, this is still a simplification of the real environment. For example, we do not consider temperature, which is phase shifted between the air and soil [43].

Although technically challenging, experiments under more realistic conditions such as this will

be important for furthering our understanding. Promising advances in this direction include the

GLO-Roots system [44], and the robotics of Bordage et al. [13], which allow the root to be

imaged in the dark under LD cycles. As experiments progress, we predict that spatial models

such as ours will be crucial for understanding how the clock integrates such a range of

environmental signals across the plant.

The molecular details underlying the spatial differences, and coordination, of the plant circadian

system are still being deciphered. Recently, *ELF3*, *ELF4*, and *LUX* were found to be central to

the difference between the shoot and root, possibly due to their interaction with the light

signaling protein PHYB [15]. However, interpretation is hindered by the fact that components

causing differences could also act as a cell-to-cell mobile coupling signal. For example, ELF4

has also been proposed to move from the shoot to couple with clocks in the root [21]. We note

that our spatial molecular model has the potential to delineate these two effects. In future work it

will be important to vary the identity of the coupling signal, and the underlying genetic network,

to generate testable predictions.

The plant circadian system, with local inputs to cells that are coupled together, represents a

decentralized structure. This is in contrast to the centralized mammalian circadian system [45].

Here we found that this decentralized structure can afford plants flexibility, allowing regions to

adopt different phases under noisy LD cycles. It could be speculated that this has a

physiological effect on plants, causing, for example, a stagger in the timing of growth across the

plant. However, further work is required to understand how, or whether, this flexibility is an

advantage to plants.

Acknowledgments

MG and JL were supported by the Gatsby Charitable Foundation (grant number GAT3395/GLC)

and IT by the Japan Society for the Promotion of Science (grant numbers 17H06313,

18H02477, and 20K11875). We thank Dr. Katie Abley (University of Cambridge) for critical

reading of the manuscript.

Conflicts of Interests

318   The authors declare that they have no conflict of interest.

Methods

Single cell molecular model

As a model for the plant circadian clock, we exploit the compact model introduced by De

Caluwe et al. [31]. The original compact model consists of 9 ordinary differential equations.

Among them, 8 equations describe the temporal evolution of the mRNA and protein levels of the

core clock genes. The clock genes are grouped into four sets of lumped pairs labeled as: CL

(*CCA1* and *LHY*), P97 (*PRR9* and *PRR7*), P51 (*PRR5* and *TOC1*) and EL (*ELF4* and *LUX*). The

9-th equation is for the light-sensitive protein P (PIF3 and PIL1).

In our modeling, one modification to the original compact model was made. According to the

experiments reported in [46], the compact model assumed that the P97 variable is activated by

CL. However, more recent work [28,34], has shown that *LHY* acts as a repressor of all other

clock components, including *PRR9* and *PRR7*. We therefore replaced the activation term that

represents the connection from CL to P97 with a repression. The revised single cell model is

now described by the following differential equations:

$$335 \quad \frac{dc_{CL}^{(m)}}{dt} = \left(v_1 + v_{1L} L(t) c_P^{(p)} \right) \frac{1}{1 + (c_{CL}^{(p)}/K_0)^2 + (c_{P97}^{(p)}/K_1)^2 + (c_{P51}^{(p)}/K_2)^2} - \left(k_{1L} L(t) + k_{1D} D(t) \right) c_{CL}^{(m)},$$

$\quad \dfrac{dc_{CL}^{(p)}}{dt} = \left(p_1 + p_{1L}L(t)\right)c_{CL}^{(m)} - d_1 c_{CL}^{(p)},$

$\quad \dfrac{dc_{P97}^{(m)}}{dt} = \left(v_{2L}L(t)c_P^{(p)} + v_{2A}\right)\dfrac{1}{1+(c_{P51}^{(p)}/K_4)^2+(c_{EL}^{(p)}/K_5)^2+(c_{CL}^{(p)}/K_{5b})^2} - k_2 c_{P97}^{(m)},$

$\quad \dfrac{dc_{P97}^{(p)}}{dt} = p_2 c_{P97}^{(m)} - \left(d_{2D}D(t) + d_{2L}L(t)\right)c_{P97}^{(p)},$

$\quad \dfrac{dc_{P51}^{(m)}}{dt} = v_3 \dfrac{1}{1+(c_{CL}^{(p)}/K_6)^2+(c_{P51}^{(p)}/K_7)^2} - k_3 c_{P51}^{(m)},$

$\quad \dfrac{dc_{P51}^{(p)}}{dt} = p_3 c_{P51}^{(m)} - \left(d_{3D}D(t) + d_{3L}L(t)\right)c_{P51}^{(p)},$

$\quad \dfrac{dc_{EL}^{(m)}}{dt} = L(t)v_4 \dfrac{1}{1+(c_{CL}^{(p)}/K_8)^2+(c_{P51}^{(p)}/K_9)^2+(c_{EL}^{(p)}/K_{10})^2} - k_4 c_{EL}^{(m)},$

$\quad \dfrac{dc_{EL}^{(p)}}{dt} = p_4 c_{EL}^{(m)} - \left(d_{4D}D(t) + d_{4L}L(t)\right)c_{EL}^{(p)},$

$\quad \dfrac{dc_P^{(p)}}{dt} = 0.3\left(1 + c_P^{(p)}\right)D(t) - c_P^{(p)}L(t).$

$c_\alpha^{(m)}$ and $c_\alpha^{(p)}$ represent the concentration of the α-th mRNA and protein (or protein complex)

respectively, for α = CL, P97, P51, EL, and P. L(t) represents the input light signal ($L = 0$, lights

off; $L > 0$, lights on) and $D(t)$ denotes a corresponding darkness input signal ($D = 1$, lights off; D

$= 0$, lights on). The model contains 34 parameters, the values of which have been obtained

previously through automated optimization [31]. The parameter values for K_0, K_4, K_5, K_{5b} related

to the modified interaction from CL to P97 were optimized by minimizing the cost function

detailed in the following subsection. Sobol search and simulated annealing were combined to

optimize within the range $K_i \in [0.1,10]$ for $i = 0,4,5,5b$. The optimized parameter values together

with the original values determined by De Caluwe et al. [31] are listed in S1 Table. The free

running periods of the revised model are 25.5 h (LL; $L = 1$) and 26.25 h (DD; $D = 1$). Under LD

cycles, peak times (ZT) of the gene expressions were CL = 0.50 h, P97 = 8.25 h, P51 = 17.50 h,

and EL = 11.75 h.

Cost function

In the parameter optimization of the single cell model, the cost function was computed as

follows. First, the model was simulated under 12-h light–12-h dark conditions for a total of 60

days and then released into LL for 60 days, followed by constant dark (DD) for 60 days. In each

light condition, the first 55 days were discarded as a transient dynamic. To ensure detectable

rhythmicity under LL and DD conditions, all variables were required to have a minimum

value of 0.1, as well as a minimum difference of 10 % between their minimum and maximum

values. Any solution that did not meet these criteria was penalized with an arbitrarily large

score. Then, the free-running period was calculated using the chi-square periodogram [47] from

CL gene expression, at a significance level of 1 %. A score of 0 was given to a solution having a

free-running period between 24 and 25 h under LL and between 25 and 28 h under DD.

Solutions with free-running periods outside these ranges were allocated the scores of $(\tau_{LL} -$

$24.5)^2/(0.1 \cdot 24.5)^2$ and $(\tau_{DD} - 26.5)^2/(0.1 \cdot 26.5)^2$, where τ_{LL} and τ_{DD} represent free-running

periods under LL and DD, respectively. For simulations under 12-h light–12-h dark cycles,

solutions that were not entrained to the LD cycles were penalized with an arbitrarily large score.

Entrained solutions were given a score of 0 for each gene that attained peak expression within $\pm$

1 h of the expected ZT, which were as follows: CL = 1.5, P97 = 6, P51 = 12, and EL = 9.

Expression peaks lying outside these intervals were scored as $(ZT_{CL} - 1.5)^2/(0.1 \cdot 24)^2$,

$(ZT_{P97} - 6)^2/(0.1 \cdot 24)^2$, $(ZT_{P51} - 12)^2/(0.1 \cdot 24)^2$, $(ZT_{EL} - 9)^2/(0.1 \cdot 24)^2$, where ZT_α denotes

Zeitgeber time of the α-th gene's peak expression.

Spatial molecular model

To simulate the spatial dynamics, we implemented the revised model on a simplified template of

a plant. The template consisted of approximately 800 cells, classified into cotyledon, hypocotyl,

root, and root tip regions (Fig 1B). Each cell contained an implementation of the revised

compact model. To simulate growth of the seedlings, we added a row of cells to the root tip

every 24 h. During this growth, the root tip region of the template was kept fixed in size. To do

this, after the addition of new cells, the previously uppermost root tip cells became root cells

instead.

As the coupling agents have yet to be clearly identified by experimental studies, the individual

cells are assumed to be coupled through CL. Our model for coupled plant cells is described as

follows:

$$\tau_i \frac{dc_{CL,i}^{(m)}}{dt} = \left(v_1 + v_{1L}L(t)c_{P,i}^{(p)}\right) \frac{1}{1 + (c_{CL,i}^{(p)}/K_0)^2 + (c_{P97,i}^{(p)}/K_1)^2 + (c_{P51,i}^{(p)}/K_2)^2}$$

$$- \left(k_{1L}L(t) + k_{1D}D(t)\right)c_{CL,i}^{(m)} + J\left(\bar{c}_{CL,i}^{(m)} - c_{CL,i}^{(m)}\right),$$

$$\tau_i \frac{dc_{CL,i}^{(p)}}{dt} = \left(p_1 + p_{1L}L(t)\right)c_{CL,i}^{(m)} - d_1 c_{CL,i}^{(p)},$$

$$\tau_i \frac{dc_{P97,i}^{(m)}}{dt} = \left(v_{2L}L(t)c_{P,i}^{(p)} + v_{2A}\right) \frac{1}{1 + (c_{P51,i}^{(p)}/K_4)^2 + (c_{EL,i}^{(p)}/K_5)^2 + (c_{CL,i}^{(p)}/K_{5b})^2} - k_2 c_{P97,i}^{(m)},$$

$$\tau_i \frac{dc_{P97,i}^{(p)}}{dt} = p_2 c_{P97,i}^{(m)} - \left(d_{2D}D(t) + d_{2L}L(t)\right)c_{P97,i}^{(p)},$$

$$\tau_i \frac{dc_{P51,i}^{(m)}}{dt} = v_3 \frac{1}{1 + (c_{CL,i}^{(p)}/K_6)^2 + (c_{P51,i}^{(p)}/K_7)^2} - k_3 c_{P51,i}^{(m)},$$

$$\tau_i \frac{dc_{P51,i}^{(p)}}{dt} = p_3 c_{P51,i}^{(m)} - \left(d_{3D}D(t) + d_{3L}L(t)\right)c_{P51,i}^{(p)},$$

$$\tau_i \frac{dc_{EL,i}^{(m)}}{dt} = L(t)v_4 \frac{1}{1 + (c_{CL,i}^{(p)}/K_8)^2 + (c_{P51,i}^{(p)}/K_9)^2 + (c_{EL,i}^{(p)}/K_{10})^2} - k_4 c_{EL,i}^{(m)},$$

$$\tau_i \frac{dc_{EL,i}^{(p)}}{dt} = p_4 c_{EL,i}^{(m)} - \left(d_{4D}D(t) + d_{4L}L(t)\right)c_{EL,i}^{(p)},$$

$$\tau_i \frac{dc_{P,i}^{(p)}}{dt} = 0.3 \left(1 + c_{P,i}^{(p)} \right) D(t) - c_{P,i}^{(p)} L(t).$$

As in the revised single cell model, $c_\alpha^{(m)}$ and $c_\alpha^{(p)}$ represent the concentration of the α-th mRNA

and protein (or protein complex) respectively for α = CL, P97, P51, EL, and P, in the i-th cell

$(i = 1, 2, ..., N)$. In the first equation, the CL gene is locally and diffusively coupled to its

neighboring cells, where J is the coupling strength and $\bar{c}_{CL,i}^{(m)} = \frac{1}{4}\sum_{j \in N(i)} c_{CL,j}^{(m)}$ represents averaged

expression level over 4 neighboring cells (left, right, above, and below). In the case of coupling

between 8 neighboring cells (left, right, above, below, left above, left below, right above, and

right below), the averaged expression level becomes $\bar{c}_{CL,i}^{(m)} = \frac{1}{8}\sum_{j \in N(i)} c_{CL,j}^{(m)}$, or $\bar{c}_{CL,i}^{(m)} = \frac{1}{N}\sum_{j=1}^{N} c_{CL,j}^{(m)}$

for the case of globally coupled cells. The coupling strength was set to $J = 0, 0.001, 0.01, 0.1, 1, 2,$

or 4.

Differential light input to cells

We set the light strength, L, to vary depending on the position of the cell. In doing so, we found

that modest differences in L were sufficient to generate period differences between regions. To

approximate the period differences that we observed between regions in experiments (Fig 2A)

we set L = 1.35 for cotyledon cells, L = 1.10 for hypocotyl cells, L = 0.90 for root cells, and L =

1.03 for root tip cells. As the template of the seedling grew, we kept the root tip region of fixed

size (see 'Spatial molecular model'). This meant that the uppermost root tip cells become root

cells instead. After this transition, the light input decreased so that cells received a level

characteristic of root cells, L = 0.90. This caused these cells to slow.

To generate cell-to-cell variability in periods, the time scaling parameters τ_i were set according

to normal distributions as N(1, 0.059), N(1, 0.028), N(1, 0.073), N(1, 0.089) for the four regions,

as informed by the analysis of single cell data [8]. For these values of L and τ_i, the intrinsic

periods of the 4 regions ($J = 0$) in the absence of growth were 23.2 ± 1.4 h, 25.5 ± 0.7 h, 27.9

± 2.0 h, and 26.3 ± 2.3 h (mean ± standard deviation) respectively.

Characterizing clock periods from experiments

We analyzed experimental luciferase and confocal data to characterize the periods within a

seedling. We previously completed an organ-level analysis of period and phase for *PRR9*::*LUC*,

ELF4::*LUC*, and *TOC1*::*LUC* expression [14]. In this analysis, 315 µm diameter regions of

interest (ROI) were defined to represent the cotyledon, hypocotyl, root, and root tip regions of

the seedling. These ROI were used to generate time series, from which period estimates were

made across a number of individual seedlings. Here, we pooled estimates made from

PRR9::*LUC*, *ELF4*::*LUC*, and *TOC1*::*LUC* lines, and calculated the mean for each region (Fig

2A). Only periods from time series classified as rhythmic (as defined previously [14]) were used

in the calculation.

To characterize the variability of the periods, we measured the between-cell variation of clock

periods in each region of the plant (S1 Fig). To do this we analyzed a dataset of single cell time-

lapse microscopy, CCA1-YFP translational fusion data [8]. We defined cells as being within the

root tip region if they are less than 315 µm from the most distal cell of the root tip. Where there

are multiple imaging sections in the same organ, cells are pooled to give one region for the

cotyledon, hypocotyl, and root. Variability was estimated using the coefficient of variation (CV;

standard deviation / mean) of all cells within these regions. Only periods from time series

classified as rhythmic (as defined previously [8]) were used in the calculation.

Organ-level analysis

To assess our model we quantified the times of the peaks of expression. For experimental data,

we used the ROI positions defined previously [14]. For the model, we defined 5-by-3 (width-by-

height) pixel ROI on the seedling template, approximating the positions used in the experimental

analysis of (S6 Fig). The mean of these ROI at each time point were taken to give the simulated

time series. Peaks within 24–144 h after transfer to LL were identified using the "findpeaks"

MATLAB (MathWorks, UK) function, with the constraint that peaks must be more than 19 h

apart. For visual clarity, only peaks in which all organs completed the full cycle within the time

window were plotted. Peaks of simulated expression were plotted against previously published

experimental data for comparison (e.g. Fig 2D–F) [14]. Only peaks from experimental time

series classified as rhythmic (as defined previously [14]) were plotted.

Space-time plots

To visualize the spatial clock dynamics, we created space-time plots of gene expression (e.g.

Fig 3A). We made plots of *PRR9*::*LUC*, *ELF4*::*LUC*, and *TOC1*::*LUC* expression as described

previously [14]. Individual time series of the space-time plots were background subtracted and

small gaps caused by segmentation errors were interpolated. A third-order Butterworth filter was

then applied to reduce the high frequency noise. To create space-time plots of simulated clock

gene expression, we also extracted the expression from sections that are perpendicular to the

primary axis of the seedling (S6 Fig). We take the mean expression from 1-by-5 cell cross-

sections of the template. To aid interpretation of space-time plots, amplitude trends were

removed from both experimental and model space-time plots using part of the mFourFit

algorithm [25].

We additionally plotted the times of the final peaks of expression of the space-time plots (e.g.

Fig 3B). The final oscillation in which all organs completed the full cycle was used. We detected

peaks within a 24 h window of the expected time of these peaks. If more than one peak was

detected, we plotted the peak with the highest height.

Entrainment to noisy LD cycles

We modelled entrainment by simulating light cycles with a day length of 12 h. The level of light

during the day varied depending on the region of the template, matching L values under LL. We

termed these cycles without noise 'idealized LD' (Fig 4A). To simulate fluctuations in the light

cycle we utilized an algorithm developed previously [48], which approximates meteorological

data [49]. Fluctuations lasted a random interval of time drawn from an exponential distribution of

mean 2.4 h. The intensity of light during each fluctuation was drawn from a second uniform

distribution of the range 0 to L_{max}, with L_{max} matching the L values for the cell under LL. The

values are set according to the position of the cell on the template, as under LL. We termed this

condition 'noisy LD' (Fig 4D). We made the simplifying assumption that each cell is exposed to

an independent noisy LD cycle due to their unique positions in the environment. LD cycles were

input to the molecular model through the parameter L.

Synchrony analysis

We estimated the global synchrony in simulations from the variance of the individual cells (Fig 5

and S14 Fig). Firstly, the average signal of all N cells of the simulation $\{x_h(t): h = 1, 2, ..., N\}$,

was computed as $M(t) = \frac{1}{N}\sum_{h=0}^{N} x_h(t)$. Then the overall level of synchrony was measured by

the synchronization index [38],

493
$$R = \frac{\langle M^2 \rangle - \langle M \rangle^2}{\frac{1}{N}\sum_{h}^{N}\{\langle x_h^2 \rangle - \langle x_h \rangle^2\}},$$

where $\langle\rangle$ denotes the time average, which in our case was between 0 and 144 h. The

synchronization index measures the ratio of the variance of the average signal to the average

variance of the individual cells. The parameter quantifies the distribution of both the phases and

amplitudes of the individual cells. However, we normalized the individual traces to remove

amplitude variance and therefore measured only the phase differences. *R* ranges from 1 when

cells are synchronized, to 0 when cells are completely desynchronized.

References

1.   Webb AAR, Seki M, Satake A, Caldana C. Continuous dynamic adjustment of the plant

circadian oscillator. Nat Commun. 2019;10: 550. doi:10.1038/s41467-019-08398-5

2.   Fung-Uceda J, Lee K, Seo PJ, Polyn S, De Veylder L, Mas P. The Circadian Clock Sets

the Time of DNA Replication Licensing to Regulate Growth in Arabidopsis. Dev Cell.

2018;45: 101–113. doi:10.1016/j.devcel.2018.02.022

3.   Nozue K, Covington MF, Duek PD, Lorrain S, Fankhauser C, Harmer SL, et al. Rhythmic

growth explained by coincidence between internal and external cues. Nature. 2007;448:

358–361. doi:10.1038/nature05946

4.   Fenske MP, Nguyen LP, Horn EK, Riffell JA, Imaizumi T. Circadian clocks of both plants

and pollinators influence flower seeking behavior of the pollinator hawkmoth Manduca

sexta. Sci Rep. 2018;8: 2842. doi:10.1038/s41598-018-21251-x

5.   Greenwood M, Locke JC. The circadian clock coordinates plant development through

specificity at the tissue and cellular level. Curr Opin Plant Biol. 2020;53: 65–72.

doi:10.1016/j.pbi.2019.09.004

6.   Dodd AN, Salathia N, Hall A, Kevei E, Toth R, Nagy F, et al. Plant Circadian Clocks

Increase Photosynthesis, Growth, Survival, and Competitive Advantage. Science.

2005;309: 630–633. doi:10.1126/science.1115581

7.   Green RM, Tingay S, Wang Z-Y, Tobin EM. Circadian rhythms confer a higher level of

fitness to Arabidopsis plants. Plant Physiol. 2002;129: 576–84. doi:10.1104/pp.004374

8.   Gould PD, Domijan M, Greenwood M, Tokuda IT, Rees H, Kozma-Bognar L, et al.

Coordination of robust single cell rhythms in the Arabidopsis circadian clock via spatial

waves of gene expression. Elife. 2018;7: e31700. doi:10.7554/eLife.31700

9.   Thain SC, Murtas G, Lynn JR, McGrath RB, Millar AJ. The circadian clock that controls

gene expression in Arabidopsis is tissue specific. Plant Physiol. 2002;130: 102–110.

doi:10.1104/pp.005405

10.   Yakir E, Hassidim M, Melamed-Book N, Hilman D, Kron I, Green RM. Cell autonomous

and cell-type specific circadian rhythms in Arabidopsis. Plant J. 2011;68: 520–531.

doi:10.1111/j.1365-313X.2011.04707.x

11.   Takahashi N, Hirata Y, Aihara K, Mas P. A Hierarchical Multi-oscillator Network

Orchestrates the Arabidopsis Circadian System. Cell. 2015;163: 148–159.

doi:10.1016/j.cell.2015.08.062

12.   James AB, Monreal JA, Nimmo GA, Kelly CL, Herzyk P, Jenkins GI, et al. The circadian

clock in Arabidopsis roots is a simplified slave version of the clock in shoots. Science.

2008;322: 1832–5. doi:10.1126/science.1161403

13.   Bordage S, Sullivan S, Laird J, Millar AJ, Nimmo HG. Organ specificity in the plant

circadian system is explained by different light inputs to the shoot and root clocks. New

Phytol. 2016;212: 136–49. doi:10.1111/nph.14024

14.   Greenwood M, Domijan M, Gould PD, Hall AJW, Locke JCW. Coordinated circadian

timing through the integration of local inputs in Arabidopsis thaliana. PLoS Biol. 2019;17:

e3000407. doi:https://doi.org/10.1371/journal.pbio.3000407

15.   Nimmo HG, Laird J, Bindbeutel R, Nusinow DA. The evening complex is central to the

difference between the circadian clocks of Arabidopsis thaliana shoots and roots. Physiol

Plant. 2020; ppl.13108. doi:10.1111/ppl.13108

16.   Endo M, Shimizu H, Nohales MA, Araki T, Kay SA. Tissue-specific clocks in Arabidopsis

show asymmetric coupling. Nature. 2014;515: 419–422. doi:10.1038/nature13919

17.   Lee HG, Seo PJ. Dependence and independence of the root clock on the shoot clock in

Arabidopsis. Genes Genomics. 2018;40: 1063–1068. doi:10.1007/s13258-018-0710-4

18.   Wenden B, Toner DLK, Hodge SK, Grima R, Millar AJ. Spontaneous spatiotemporal

550 waves of gene expression from biological clocks in the leaf. Proc Natl Acad Sci U S A.

551 2012;109: 6757–6762. doi:10.1073/pnas.1118814109

552 19. Fukuda H, Nakamichi N, Hisatsune M, Murase H, Mizuno T. Synchronization of Plant

553 Circadian Oscillators with a Phase Delay Effect of the Vein Network. Phys Rev Lett.

554 2007;99: 098102. doi:10.1103/PhysRevLett.99.098102

555 20. Fukuda H, Ukai K, Oyama T. Self-arrangement of cellular circadian rhythms through

556 phase-resetting in plant roots. Phys Rev E. 2012;86: 041917.

557 doi:10.1103/PhysRevE.86.041917

558 21. Chen WW, Takahashi N, Hirata Y, Ronald J, Porco S, Davis SJ, et al. A mobile ELF4

559 delivers circadian temperature information from shoots to roots. Nat Plants. 2020;6: 416–

560 426. doi:10.1038/s41477-020-0634-2

561 22. Nimmo HG. Entrainment of Arabidopsis roots to the light:dark cycle by light piping. Plant

562 Cell Environ. 2018;41: 1742–1748. doi:10.1111/pce.13137

563 23. Locke JCW, Southern MM, Kozma-Bognár L, Hibberd V, Brown PE, Turner MS, et al.

564 Extension of a genetic network model by iterative experimentation and mathematical

565 analysis. Mol Syst Biol. 2005;1: 2005.0013. doi:10.1038/msb4100018

566 24. Locke JCW, Kozma-Bognár L, Gould PD, Fehér B, Kevei E, Nagy F, et al. Experimental

567 validation of a predicted feedback loop in the multi-oscillator clock of Arabidopsis

568 thaliana. Mol Syst Biol. 2006;2: 59. doi:10.1038/msb4100102

569 25. Edwards KD, Akman OE, Knox K, Lumsden PJ, Thomson AW, Brown PE, et al.

570 Quantitative analysis of regulatory flexibility under changing environmental conditions.

571 Mol Syst Biol. 2010;6: 424. doi:10.1038/MSB.2010.81

572 26. Pokhilko A, Hodge SK, Stratford K, Knox K, Edwards KD, Thomson AW, et al. Data

573 assimilation constrains new connections and components in a complex, eukaryotic

574 circadian clock model. Mol Syst Biol. 2010;6: 416. doi:10.1038/msb.2010.69

575 27. Pokhilko A, Fernández AP, Edwards KD, Southern MM, Halliday KJ, Millar AJ. The clock

gene circuit in Arabidopsis includes a repressilator with additional feedback loops. Mol

Syst Biol. 2012;8: 574. doi:10.1038/msb.2012.6

28.   Fogelmark K, Troein C. Rethinking transcriptional activation in the Arabidopsis circadian

clock. PLoS Comput Biol. 2014;10: e1003705. doi:10.1371/journal.pcbi.1003705

29.   Akman OE, Watterson S, Parton A, Binns N, Millar AJ, Ghazal P. Digital clocks: simple

Boolean models can quantitatively describe circadian systems. J R Soc Interface. 2012;9:

2365–2382. doi:10.1098/rsif.2012.0080

30.   Foo M, Somers DE, Kim P-J. Kernel Architecture of the Genetic Circuitry of the

Arabidopsis Circadian System. Jensen LJ, editor. PLoS Comput Biol. 2016;12:

e1004748. doi:10.1371/journal.pcbi.1004748

31.   De Caluwé J, Xiao Q, Hermans C, Verbruggen N, Leloup J-C, Gonze D. A Compact

Model for the Complex Plant Circadian Clock. Front Plant Sci. 2016;7: 74.

doi:10.3389/fpls.2016.00074

32.   Tokuda IT, Akman OE, Locke JCW. Reducing the complexity of mathematical models for

the plant circadian clock by distributed delays. J Theor Biol. 2019;463: 155–166.

doi:10.1016/J.JTBI.2018.12.014

33.   Kuramoto Y. Self-entrainment of a population of coupled non-linear oscillators.

International Symposium on Mathematical Problems in Theoretical Physics.

Berlin/Heidelberg: Springer-Verlag; 1975. pp. 420–422. doi:10.1007/bfb0013365

34.   Adams S, Manfield I, Stockley P, Carre IA, Dowidar N, Dunaway D. Revised Morning

Loops of the Arabidopsis Circadian Clock Based on Analyses of Direct Regulatory

Interactions. Foulkes NS, editor. PLoS One. 2015;10: e0143943.

doi:10.1371/journal.pone.0143943

35.   Taiz L, Zeiger E, Møller IM, Murphy A. Plant Physiology and Development. 6th ed.

Sunderland: Sinauer Associates; 2018.

36.   Aschoff J, Pohl H. Phase relations between a circadian rhythm and its zeitgeber within

602   the range of entrainment. Naturwissenschaften. 1978;65: 80–4. doi:10.1007/BF00440545

603 37. Maizel A, Markmann K, Timmermans M, Wachter A. To move or not to move: roles and

604   specificity of plant RNA mobility. Curr Opin Plant Biol. 2020;57: 52–60.

605   doi:10.1016/j.pbi.2020.05.005

606 38. Garcia-Ojalvo J, Elowitz MB, Strogatz SH. Modeling a synthetic multicellular clock:

607   repressilators coupled by quorum sensing. Proc Natl Acad Sci U S A. 2004;101: 10955–

608   60. doi:10.1073/pnas.0307095101

609 39. Seaton DD, Ebenhöh O, Millar AJ, Pokhilko A. Regulatory principles and experimental

610   approaches to the circadian control of starch turnover. J R Soc Interface. 2014;11:

611   20130979. doi:10.1098/rsif.2013.0979

612 40. Seaton DD, Smith RW, Song YH, MacGregor DR, Stewart K, Steel G, et al. Linked

613   circadian outputs control elongation growth and flowering in response to photoperiod and

614   temperature. Mol Syst Biol. 2015;11: 776. doi:10.15252/MSB.20145766

615 41. Shimizu H, Katayama K, Koto T, Torii K, Araki T, Endo M. Decentralized circadian clocks

616   process thermal and photoperiodic cues in specific tissues. Nat Plants. 2015;1: 15163.

617   doi:10.1038/nplants.2015.163

618 42. Meyerowitz EM. Genetic control of cell division patterns in developing plants. Cell.

619   1997;88: 299–308. doi:10.1016/S0092-8674(00)81868-1

620 43. Dawson GB, Fisher RG. Diurnal and seasonal ground temperature variations at Wairakei.

621   New Zeal J Geol Geophys. 1964;7: 144–154. doi:10.1080/00288306.1964.10420166

622 44. Rellán-Álvarez R, Lobet G, Lindner H, Pradier P-L, Sebastian J, Yee M-C, et al. GLO-

623   Roots: an imaging platform enabling multidimensional characterization of soil-grown root

624   systems. Elife. 2015;4. doi:10.7554/eLife.07597

625 45. Bell-Pedersen D, Cassone VM, Earnest DJ, Golden SS, Hardin PE, Thomas TL, et al.

626   Circadian rhythms from multiple oscillators: lessons from diverse organisms. Nat Rev

627   Genet. 2005;6: 544–556. doi:10.1038/nrg1633

628   46.   Para A, Farré EM, Imaizumi T, Pruneda-Paz JL, Harmon FG, Kay SA. PRR3 Is a

629   vascular regulator of TOC1 stability in the Arabidopsis circadian clock. Plant Cell.

630   2007;19: 3462–73. doi:10.1105/tpc.107.054775

631   47.   Sokolove PG, Bushell WN. The chi square periodogram: Its utility for analysis of circadian

632   rhythms. J Theor Biol. 1978;72: 131–160. doi:10.1016/0022-5193(78)90022-X

633   48.   Pittayakanchit W, Lu Z, Chew J, Rust MJ, Murugan A. Biophysical clocks face a trade-off

634   between internal and external noise resistance. Elife. 2018;7. doi:10.7554/eLife.37624

635   49.   Gu L, Fuentes JD, Garstang M, Silva JT Da, Heitz R, Sigler J, et al. Cloud modulation of

636   surface solar irradiance at a pasture site in Southern Brazil. Agric For Meteorol.

637   2001;106: 117–129. doi:10.1016/S0168-1923(00)00209-4

Figures

Fig 1. The structure of the spatial circadian clock model. (A) Summary of the modified

compact circadian clock model used for simulations. The original compact model [31] is

modified to include a repression interaction between CCA1/LHY and PRR5/TOC1. Light input is

received by all gene pairs, but affects their production and degradation rates to different

degrees. Yellow dashed arrows represent light input at the transcriptional level and solid yellow

arrows represent light input at both the transcriptional and post-transcription level. "T" arrows

represent repression. (B) The network was implemented within each cell on a simplified

template of a seedling, with cells classified as either cotyledon (blue), hypocotyl (turquoise), root

(yellow), or root tip cells (green). (C) The network in each individual cell is coupled to the

neighboring cells through the level of *CCA1/LHY*. *CCA1*, *CIRCADIAN CLOCK ASSOCIATED 1*;

ELF4, *EARLY FLOWERING 4*; *LHY*, *LATE ELONGATED HYPOCOTYL*; LL, constant light;

LUC, *LUCIFERASE*; *LUX*, *LUX ARRHYTHMO*; *PRR*, *PSEUDO-RESPONSE REGULATOR*;

TOC1, *TIMING OF CAB EXPRESSION 1*.

Fig 2. Regional differences in light input can generate the period structure observed

experimentally. (A) Period estimates of *PRR9*::*LUC*, *TOC1*::*LUC*, and *ELF4*::*LUC* for different

organs imaged under LL. Each data point represents a period estimate from the organ of a

single seedling. The horizontal black line shows the mean. (B) Period estimates of simulated

PRR9/PRR7 expression, measured from regions within the template seedling. The black line

indicates the mean and the red shaded area one standard deviation of 18 independent

simulations. (C) Periods of individual cells plotted on the template seedling. The color of the cell

represents the speed of the oscillation. By assuming higher light input to the cells in the

cotyledon, hypocotyl, and root tip, the model approximates the period differences observed

between regions in experiments, A. A noise parameter was adjusted for each region to simulate

the variability of periods observed in single cell experiments (S1 Fig). (D−F) Times of peaks of

simulated *PRR9*/*PRR7* and *PRR9*::*LUC* (D), simulated *PRR5*/*TOC1* and *TOC1*::*LUC* (E), or

simulated *ELF4*/*LUX* and *ELF4*::*LUC* (F), in different organs under LL. Simulations include

different levels of light input to regions and cell-to-cell coupling ($J = 2$). Plots represent the 25th

percentile, median, and the 75th percentile for the peak times of oscillations scored as rhythmic,

n = 9 simulations. Experimental data is an analysis of time-lapse movies carried out previously

[14]. *CCA1*, *CIRCADIAN CLOCK ASSOCIATED 1*; CV, coefficient of variation; *ELF4*, *EARLY*

FLOWERING 4; *LHY*, *LATE ELONGATED HYPOCOTYL*; LL, constant light; *LUC*,

LUCIFERASE; *LUX*, *LUX ARRHYTHMO*; *PRR*, *PSEUDO-RESPONSE REGULATOR*; *TOC1*,

TIMING OF CAB EXPRESSION 1.

Fig 3. Local sharing of mRNA can reproduce experimentally observed spatial waves of clock gene expression. (A) Representative intensity plot of *PRR9*::*LUC* expression across longitudinal sections of a single seedling under LL. (B) Times of the final peaks of the *PRR9*::*LUC* intensity plot simulated under LL. (C) Representative intensity plot of simulated *PRR9*/*PRR7* expression across longitudinal sections of a single seedling under LL. Simulations include different levels of light input to regions and cell-to-cell coupling (J = 2). (D) Times of the final peaks of the simulated *PRR9*/*PRR7* intensity plot simulated under LL. (E) Times of the final peak of simulated *PRR9*/*PRR7* intensity plots, each simulated under LL with increasing strength of coupling, *J*. Experimental data is an analysis of time-lapse movies carried out previously [14].

705   *PRR*, *PSEUDO RESPONSE REGULATOR*; LL, constant light; *LUC, LUCIFERASE. TOC1,*

706   *TIMING OF CAB EXPRESSION 1.*

Fig 4. Regional phase differences persist under idealized and noisy LD cycles. (A)

Schematic of idealized LD cycles, without fluctuations in light levels during the daytime. (B)

Times of the final peak of simulated *PRR5*/*TOC1* intensity plots, each simulated under idealized

LD with increasing strength of coupling, *J*. (C) Schematic of noisy LD cycles, with fluctuations in

light levels during the daytime. Each line represents an LD cycle for a single cell. (D) Times of

the final peak of simulated *PRR5*/*TOC1* intensity plots, each simulated under noisy LD with

increasing strength of coupling, *J*. LD, light-dark; *PRR*, *PSEUDO-RESPONSE REGULATOR*;

TOC1, *TIMING OF CAB EXPRESSION 1*.

Fig 5. Cell-to-cell coupling maintains global synchrony under noisy LD cycles. (A−D)

Quantification of phase coherence by the synchronization index, *R*, for simulated *CCA1/LHY*

(A), *PRR9/PRR7* (B), *PRR5/TOC1* (C) or *ELF4/LUX* (D) mRNA expression under idealized or

noisy LD cycles. Simulations included increasing strengths of coupling, *J*. Color legends are as

in A. Data points represent the mean ± standard deviation, *n* = 9 simulations. *CCA1*,

CIRCADIAN CLOCK ASSOCIATED 1; *ELF4*, *EARLY FLOWERING 4*; LD, light-dark; *LHY*,

LATE ELONGATED HYPOCOTYL; *LUX*, *LUX ARRHYTHMO*; *LUC*, *LUCIFERASE*; *PRR*,

PSEUDO-RESPONSE REGULATOR; *TOC1*, *TIMING OF CAB EXPRESSION 1*.

750 **S1 Fig. Between-cell variability of CCA1-YFP rhythms**. Between-cell variability of periods

751 within different organs. The periods were estimated from single cell time-lapse movies of CCA1-

752 YFP expression carried out previously [8]. *CCA1*, *CIRCADIAN CLOCK ASSOCIATED 1*; CV,

753 coefficient of variation; YFP, yellow fluorescent protein.

S2 Fig. Simulated expression of clock genes in different regions under LD cycles.

Expression of simulated *CCA1/LHY*, *PRR9/PRR7*, *PRR5/TOC1*, and *ELF4/LUX* from regions of

the seedling template 96–144 h after the beginning of LD cycles. The light input, *L*, was varied

depending on whether the cell was within the cotyledon (*L* = 1.35), hypocotyl (*L* = 1.10), root (*L*

= 0.90), or root tip (*L* = 1.03) region, and coupling was present between cells (*J* = 2). Data

represent the mean ± standard error, *n* = 9 simulations. *CCA1*, *CIRCADIAN CLOCK*

ASSOCIATED 1; *ELF4*, *EARLY FLOWERING 4*; LD, light-dark; *LHY*, *LATE ELONGATED*

763   *HYPOCOTYL; LUC, LUCIFERASE; LUX, LUX ARRHYTHMO; PRR, PSEUDO-RESPONSE*

764   *REGULATOR; TOC1, TIMING OF CAB EXPRESSION 1.*

S3 Fig. Simulated expression of multiple core clock genes under LL qualitatively match

luciferase reporter experiments. (A–C) Expression of simulated *PRR9*/*PRR7* and *PRR9*::*LUC*

(A), simulated *PRR5*/*TOC1* and *TOC1*::*LUC (B)*, or simulated *ELF4*/*LUX* and *ELF4*::*LUC* (C) in

different organs under LL. Simulations include different levels of light input to regions and cell-

to-cell coupling (*J* = 2). Data represent the mean ± standard error of organs scored as rhythmic.

n = 9 simulations. Experimental data is from time-lapse movies carried out previously [14].

792 *ELF4, EARLY FLOWERING 4*; LL, constant light; *LUC, LUCIFERASE; LUX, LUX*

793 *ARRHYTHMO*; *PRR, PSEUDO-RESPONSE REGULATOR*; *TOC1, TIMING OF CAB*

794 *EXPRESSION 1.*

817 **S4 Fig. Higher light intensities cause simulations to run faster.** The mean peak-to-peak

818 time of simulated *PRR9*/PRR7 expression with increasing intensities of light, *L*. A single

819 instance of the model was implemented at each *L*, without cell-to-cell coupling or variation in

820 gene expression. *L* values below 0.8 caused damping in the original and modified De Caluwe

821 implementation. *PRR*, *PSEUDO-RESPONSE REGULATOR*.

S5 Fig. Local sharing of mRNA can reproduce experimentally observed spatial waves of

clock gene expression for multiple core clock genes. (A–D) Representative intensity plot of

TOC1::LUC (A), simulated *PRR5/TOC1* (B)*, ELF4::LUC* (C) and simulated *ELF4/LUX* (D)

expression across longitudinal sections of a single seedling under LL. Simulations include

different levels of light input to regions and cell-to-cell coupling ($J = 2$). (E–H) Times of the final

peak of *TOC1*::*LUC* (E), simulated *PRR5*/*TOC1* (F)*, ELF4*::*LUC* (G) and simulated *ELF4*/*LUX*

(H) intensity plots. Experimental data is an analysis of time-lapse movies carried out previously

[14]. *ELF4*, *EARLY FLOWERING 4*; *LHY*, *LATE ELONGATED HYPOCOTYL*; LL, constant light;

LUX, *LUX ARRHYTHMO*; *PRR5*, *PSEUDO-RESPONSE REGULATOR 5; TOC1*, *TIMING OF*

CAB EXPRESSION 1.

A   **B**

S6 Fig. Spatial analysis of rhythms across the simplified template of a plant. (A) To

measure the period and phase of an organ, the expression was extracted from 3-by-3 pixel

regions of interest (yellow squares). (B) To produce space-time plots of gene expression (e.g.

Fig 3A), the mean expression of sections perpendicular across the central axis (yellow line) of

the template were taken. Luciferase space-time plots were created similarly, as described

previously [14] (Methods).

S7 Fig. Times of the final peaks of simulated expression under LL with increasing

strengths of coupling. (A–C) Times of the final peak of simulated *PRR9/PRR7* (A), simulated

PRR5/*TOC1* (B)*,* and simulated *ELF4*/*LUX* (C) intensity plots, each simulated under LL with

increasing strength of coupling, *J. ELF4*, *EARLY FLOWERING 4*; LL, constant light; *LUX, LUX*

ARRHYTHMO; *PRR*, *PSEUDO-RESPONSE REGULATOR; TOC1*, *TIMING OF CAB*

EXPRESSION 1.

S8 Fig. Times of the final peaks of simulated expression under LL with the sharing of

different clock gene mRNA between cells. (A–C) Times of the final peak of simulated

PRR9/PRR7 intensity plots, each simulated with the sharing of *CCA1*/*LHY*, *PRR9*/*PRR7*,

PRR5/*TOC1*, or *ELF4*/*LUX* mRNA between neighbor cells. Simulations were under LL with

increasing strength of coupling, *J. CCA1, CIRCADIAN CLOCK ASSOCIATED 1*; *ELF4, EARLY*

FLOWERING 4; *LHY, LATE ELONGATED HYPOCOTYL*; LL, constant light; *LUX, LUX*

ARRHYTHMO; PRR, PSEUDO-RESPONSE REGULATOR; TOC1, TIMING OF CAB

EXPRESSION 1.

S9 Fig. Times of the final peaks of simulated expression under LL without variation in

gene expression. (A–C) Times of the final peak of simulated *PRR9*/*PRR7* (A)*, PRR5*/*TOC1*

(B), or *ELF4*/*LUX* (C) intensity plots, each simulated with the sharing of *CCA1*/*LHY* mRNA

between 4 neighbor cells, but without cell-to-cell variation in gene expression. Simulations were

under LL with increasing strength of coupling, *J. CCA1, CIRCADIAN CLOCK ASSOCIATED 1*;

ELF4, EARLY FLOWERING 4; *LHY, LATE ELONGATED HYPOCOTYL*; LL, constant light;

LUX, LUX ARRHYTHMO; *PRR, PSEUDO-RESPONSE REGULATOR; TOC1, TIMING OF CAB*

EXPRESSION 1.

S10 Fig. Times of the final peaks of simulated expression under LL with different

coupling rules. (A–C) Times of the final peak of simulated *PRR9/PRR7* intensity plots, each

simulated under LL with the sharing of *CCA1*/*LHY* mRNA between the 4 nearest neighbor cells

(A), 8 nearest neighbor cells (B), or globally (all-to-all) (C). Simulations were run under LL with

increasing strength of coupling, *J. CCA1*, *CIRCADIAN CLOCK ASSOCIATED 1*; *LHY, LATE*

ELONGATED HYPOCOTYL; LL, constant light; *PRR, PSEUDO-RESPONSE REGULATOR*.

S11 Fig. Simulations with equal light input to regions predicts the loss of spatial waves

observed in the *phyb-9* mutant. (A) Representative intensity plot of *GI*::*LUC* expression

across longitudinal sections of a single seedling under constant red light. Imaging was

performed in the light sensing mutant *phyb-9* background. (B) Representative intensity plot of

simulated *PRR5*/*TOC1* expression across longitudinal sections of a single seedling under LL.

Simulations were performed with equal low light to all regions (L = 0.85) and cell-to-cell coupling

(J = 2). (C) Times of the final peaks of *phyb-9 TOC1*::*LUC* intensity plots. (D) Times of the final

peaks of simulated *PRR5*/*TOC1* intensity plots, with equal low light to all regions (L = 0.85) and

cell-to-cell coupling (J = 2). Experimental data is an analysis of time-lapse movies carried out

previously [14]. Intensity plots are limited to the lower hypocotyl and root due to rapid elongation

of the hypocotyl in the *phyb-9* mutant. *PRR, PSEUDO RESPONSE REGULATOR*; LL, constant

light; *LUC, LUCIFERASE. TOC1, TIMING OF CAB EXPRESSION 1.*

S12 Fig. Times of the final simulated peaks of expression under idealized LD cycles with

increasing strengths of coupling. (A–D) Times of the final peak of simulated *CCA1*/*LHY* (A),

PRR9/PRR7 (B), *PRR5*/*TOC1* (C)*,* and *ELF4*/*LUX* (D) intensity plots, each simulated under

idealized LD cycles with increasing strength of coupling, *J*. Cells were exposed to idealized LD

cycles during simulations. Grey background indicates nighttime. *ELF4* appears synchronized

when the peak coincides with the dark transition as darkness causes strong repression of *ELF4*

expression. *CCA1, CIRCADIAN CLOCK ASSOCIATED 1*; *ELF4, EARLY FLOWERING 4*; LD,

light-dark; *LHY, LATE ELONGATED HYPOCOTYL*; *LUX, LUX ARRHYTHMO*; *PRR, PSEUDO*

RESPONSE REGULATOR; *TOC1, TIMING OF CAB EXPRESSION 1.*

S13 Fig. Times of the final simulated peaks of expression under noisy LD cycles with

increasing strengths of coupling. (A–D) Times of the final peak of simulated *CCA1*/*LHY* (A),

PRR9/PRR7 (B), simulated *PRR5*/*TOC1* (C)*,* and simulated *ELF4*/*LUX* (D) intensity plots, each

simulated under idealized LD cycles with increasing strength of coupling, *J*. Cells were exposed

to idealized LD cycles during simulations. Grey background indicates nighttime. *ELF4* appears

synchronized when the peak coincides with the dark transition as darkness causes strong

987   repression of *ELF4* expression. *CCA1, CIRCADIAN CLOCK ASSOCIATED 1*; *ELF4, EARLY*

988   *FLOWERING 4*; LD, light-dark; *LHY, LATE ELONGATED HYPOCOTYL*; *LUX, LUX*

989   *ARRHYTHMO*; *PRR, PSEUDO RESPONSE REGULATOR*; *TOC1, TIMING OF CAB*

990   *EXPRESSION 1*.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

S14 Fig. Cell-to-cell coupling increases the synchrony of protein oscillations under realistic LD cycles. (A–D) Quantification of phase coherence by the synchronization index, R, for simulated *CCA1*/*LHY* (A), *PRR9*/*PRR7* (B), *PRR5*/*TOC1* (C) or *ELF4*/*LUX* (D) protein expression under idealized or noisy LD cycles. Simulations included increasing strengths of coupling, J. Color legends are as in A. Data points represent the mean ± standard deviation, n = 9 simulations. CCA1, CIRCADIAN CLOCK ASSOCIATED 1; ELF4, EARLY FLOWERING 4; LD, light-dark; LHY, LATE ELONGATED HYPOCOTYL; LUX, LUX ARRHYTHMO; PRR, PSEUDO RESPONSE REGULATOR; TOC1, TIMING OF CAB EXPRESSION 1.

S1 Table. Parameter values for the spatial clock model.

Parameter	Description	Original value [31]	Re-optimized value
v_1	*CCA1/LHY* synthesis	4.6 nM h^{-1}	4.58 nM h^{-1}
v_{1L}	*CCA1/LHY* light-induced synthesis	3.0 nM h^{-1}	3.0 nM h^{-1}
v_{2A}	*PRR9/PRR7* synthesis	1.3 nM h^{-1}	1.27 nM h^{-1}
v_{2L}	*PRR9/PRR7* light-induced synthesis	5.0 nM h^{-1}	5.0 nM h^{-1}
v_3	*PRR5/TOC1* synthesis	1.0 nM h^{-1}	1.0 nM h^{-1}
v_4	*ELF4/LUX* synthesis	1.5 nM h^{-1}	1.47 nM h^{-1}
k_{1L}	*CCA1/LHY* mRNA degradation (light)	0.53 h^{-1}	0.53 h^{-1}
k_{1D}	*CCA1/LHY* mRNA degradation (dark)	0.21 h^{-1}	0.21 h^{-1}
k_2	*PRR9/PRR7* mRNA degradation	0.35 h^{-1}	0.35 h^{-1}
k_3	*PRR5/TOC1* mRNA degradation	0.56 h^{-1}	0.56 h^{-1}
k_4	*ELF4/LUX* mRNA degradation	0.57 h^{-1}	0.57 h^{-1}
p_1	CCA1/LHY translation	0.76 h^{-1}	0.76 h^{-1}
p_{1L}	CCA1/LHY light-induced translation	0.42 h^{-1}	0.42 h^{-1}
p_2	PRR9/*PRR7* translation	1.0 h^{-1}	1.01 h^{-1}
p_3	PRR5/TOC1 translation	0.64 h^{-1}	0.64 h^{-1}
p_4	ELF4/LUX translation	1.0 h^{-1}	1.01 h^{-1}
d_1	CCA1/LHY degradation	0.68 h^{-1}	0.68 h^{-1}
d_{2D}	PRR9/*PRR7* degradation (dark)	0.51 h^{-1}	0.5 h^{-1}
d_{2L}	PRR9/*PRR7* degradation (light)	0.30 h^{-1}	0.29 h^{-1}
d_{3D}	PRR5/TOC1 degradation (dark)	0.48 h^{-1}	0.48 h^{-1}
d_{3L}	PRR5/TOC1 degradation (light)	0.78 h^{-1}	0.78 h^{-1}
d_{4D}	ELF4/LUX degradation (dark)	1.2 h^{-1}	1.21 h^{-1}
d_{4L}	ELF4/LUX degradation (light)	0.38 h^{-1}	0.38 h^{-1}
K_0	Inhibition of CCA1/LHY by CCA1/LHY	n/a	5.07 nM
K_1	Inhibition of CCA1/LHY by PRR9/PRR7	0.16 nM	0.16 nM
K_2	Inhibition of CCA1/LHY by PRR5/TOC1	1.2 nM	1.18 nM
K_4	Inhibition of PRR9/PRR7 by PRR5/TOC1	0.23 nM	0.40 nM
K_5	Inhibition of PRR9/PRR7 by ELF4/LUX	0.30	0.62 nM
K_{5b}	Inhibition of PRR9/PRR7 by CCA1/LHY	n/a	1.2 nM
K_6	Inhibition of PRR5/TOC1 by CCA1/LHY	0.46	0.46 nM
K_7	Inhibition of PRR5/TOC1 by PRR5/TOC1	2.0	2.0 nM
K_8	Inhibition of ELF4/LUX by CCA1/LHY	0.36	0.36 nM
K_9	Inhibition of ELF4/LUX by PRR5/TOC1	1.9	1.9 nM
K_{10}	Inhibition of ELF4/LUX by ELF4/LUX	1.9	1.9 nM

CCA1, CIRCADIAN CLOCK ASSOCIATED 1; *ELF4, EARLY FLOWERING 4*; *LHY, LATE*
ELONGATED HYPOCOTYL; *LUX, LUX ARRHYTHMO*; *PRR, PSEUDO-RESPONSE*
REGULATOR; *TOC1, TIMING OF CAB EXPRESSION 1.*

S1 Video. Time-lapse of simulated *PRR9*/*PRR7* expression under LL. Representative

simulation of simulated *PRR9*/*PRR7* expression from individual cells of the plant template 24–

144 h after transfer to LL. LL, constant light; *PRR, PSEUDO RESPONSE REGULATOR.*

1034

1035 **S2 Video. Time-lapse of *PRR9*::*LUC* expression under LL.** *PRR9*::*LUC* expression from a

1036 single representative seedling 24–144 h after transfer to LL. LL, constant light; *PRR9, PSEUDO*

1037 *RESPONSE REGULATOR9.*