Enzymatic synthesis of fatty acid amides using microbial lipids as acyl group-donors and their

biological activities

Hatim A. El-Baz [1], Ahmed M. Elazzazy [2,3], Tamer S. Saleh [4], Panagiotis Dritsas [5], Jazem A.

Mahyoub [6], Mohammed N. Baeshen [2], Hekmat R. Madian [7], Mohammed Alkhaled [2] and George

Aggelis [2,5*]

[1] Department of Clinical Biochemistry, College of Medicine, University of Jeddah, Jeddah

21589, Saudi Arabia.

[2] Department of Biology, College of Science, University of Jeddah, Jeddah 21589, Saudi Arabia

[3] Department of Chemistry of Natural and Microbial Products, Division of Pharmaceutical and

Drug Industries, National Research Centre, Dokki, Giza 12622, Egypt

[4] Department of Chemistry, College of Science, University of Jeddah, Jeddah 21589, Saudi

Arabia

[5] Unit of Microbiology, Division of Genetics, Cell and Developmental Biology, Department of

Biology, University of Patras, Patras 26504, Greece

[6] Department of Biological Sciences, Faculty of Science, King Abdulaziz University, Jeddah

21589, Saudi Arabia.

[7] Egyptian Petroleum Research Institute, Nasr City, Cairo 11727, Egypt

[*] Corresponding author:

Professor George Aggelis, email address: George.Aggelis@upatras.gr

Unit of Microbiology, Division of Genetics, Cell and Developmental Biology

Department of Biology, University of Patras

Patras 26504, Greece

Abstract

Fatty acid amides (FAAs) are of great interest due to their broad industrial applications. They can

be synthesized enzymatically with many advantages over chemical synthesis. In this study, the

fatty acid moieties of lipids of *Cunninghamella echinulata* ATHUM 4411, *Umbelopsis isabellina*

ATHUM 2935, *Nannochloropsis gaditana* CCAP 849/5, Olive oil and an eicosapentaenoic acid

(EPA) concentrate were converted into their fatty acid methyl esters and used in the FAA (i.e.

ethylene diamine amides) enzymatic synthesis, using lipases as biocatalysts. The FAA synthesis,

monitored using *in situ* NMR, FT-IR and thin-layer chromatography, was catalyzed efficiently

by the immobilized *Candida rugosa* lipase. The synthesized FAAs exhibited a significant

antimicrobial activity, especially those containing oleic acid in high proportions (i.e. derived

from Olive oil and *U. isabellina* oil), against several human pathogenic microorganisms,

insecticidal activity against yellow fever mosquito, especially those of *C. echinulata* containing

gamma linolenic acid, and anti-cancer properties against SKOV-3 ovarian cancer cell line,

especially those containing EPA in their structures (i.e. EPA concentrate and *N. gaditana* oil).

We conclude that FAAs can be efficiently synthesized using microbial oils of different fatty acid

composition and used in specific biological applications.

Keywords: Single cell oil; Fatty acid methyl esters; Fatty acid amide synthesis; Antimicrobial;

Insecticidal; Anti-cancer activity

Abbreviations: *ANOVA*, Analysis of variance; *ASW*, Artificial sea water; *CLSI*, Clinical and

Laboratory Standards Institute; *DHA*, Docosahexaenoic acid; *EPA*, Eicosapentaenoic acid; *FA*,

Fatty acid; *FAAs*, Fatty acid amides; *FAMEs*, Fatty acid methyl esters; *FT-IR*, Fourier-transform

infrared; *GLA*, Gamma linolenic acid; *LC50*, Median lethal concentration; *MBC*, Minimum

bactericidal concentration; *MHA*, Mueller Hinton II Agar; *MIC*, Minimum inhibitory

concentration; NMR, Nuclear magnetic resonance; *OPSR*, Open-pond simulating reactor; *PDA*,

Potato dextrose agar; *PUFAs*, Polyunsaturated fatty acids; *SCOs*, Single cell oils; *TLC*, Thin-

layer chromatography

Introduction

Fatty acid amides (FAAs) are organic compounds formed from a fatty acid (FA) and an amine,

such as ethanolamine or an amino acid. FAAs can be synthesized from alkanolamine and a fatty

acyl donor, such as a free FA or a FA alkyl ester, by chemical or enzymatic esterification or

transesterification methods [1, 2]. The enzymatic synthesis of FAAs can be performed using

lipases [3, 4], or aminoacylases [5]. FAAs are of considerable interest due to their wide-ranging

industrial applications in the production of lubricants, detergents, shampoo, cosmetics and

surfactant formulations [6, 7]. In addition, FAAs, demonstrating a potent antimicrobial activity

against Gram-positive and Gram-negative bacteria [8] and possessing beneficial anti-

inflammatory properties [9], provide an exciting opportunity to produce new medicines and

nutraceuticals with applications in the treatment of several human diseases and in human

nutrition [7, 10].

There are different sources of FAs, such as common plant oils and animal fats, which

can be used in amide synthesis. Alternatively, microbial lipids, so called Single Cell Oils

(SCOs), derived from microalgae and fungi, which do not compete with the food supply chain,

could be considered for this purpose. Microalgae and fungi are on the forefront of

biotechnological interest due to their ability to produce SCOs rich in polyunsaturated fatty acids

(PUFAs) of medical and nutritional interest [11-17]. The high PUFA content of the

aforementioned lipids offers an additional interest in their use as acyl group-donors in FAA

synthesis, since several reports demonstrate that PUFAs or compounds containing PUFA

moieties in their molecule exhibited interesting biological activities [18-20]. Among microalgae,

Nannochloropsis is a prominent genus that include species able to efficiently grow under non-

aseptic conditions and accumulate lipids rich in PUFAs, such as eicosapentaenoic acid (EPA)

[21, 22]. As for fungi, genera belonging to Mucoromycota (including *Mucor, Rhizopus,*

Umbelopsis, Lichtheimia, Cunninghamella and *Mortierella*) are well known for their ability to

synthesize PUFAs, especially gamma linolenic acid (GLA), which is of great pharmaceutical

interest due to its anticancer properties, while it has been used to improve premenstrual tension

and various skin diseases [13, 17]. Especially, *Cunninghamella echinulata* is an important GLA

producer [13, 19, 23], while *Umbelopsis isabellina* is regarded as a promising SCO producer,

being able to accumulate lipids in high percentages, though less rich in GLA [16, 24-26].

The aim of this study was to produce through enzymatic synthesis FAAs using as acyl

group-donors SCOs of different FA composition, such as those derived from the fungi

Umbelopsis isabellina (containing oleic acid in high percentage and GLA in low percentage) and

Cunninghamella echinulata (containing GLA in high percentages) and the microalga

Nannochloropsis gaditana (containing EPA in high percentages). The biological activity of the

above FAAs was tested against important human pathogens, the larvae of *Aedes aegypti* and the

SKOV-3 cancer cell line and compared with that of FAAs produced using as acyl group-donors

Olive oil (containing oleic acid in very high percentages) and an EPA concentrate (i.e. a fish oil

derivative containing EPA in very high percentages). We concluded that FAAs can be efficiently

produced using lipids of microbial origin and employed as bioactive compounds in various

biological applications depending on their FA composition.

Materials and methods

Biological materials and culture conditions

The fungal strains *Cunninghamella echinulata* ATHUM 4411 and *Umbelopsis isabellina*

ATHUM 2935 (culture collection of National and Kapodistrian University of Athens, Greece)

were maintained on potato dextrose agar (PDA) (Biolab Zrt, Budapest, Hungary) at 7 ± 1 ^{o}C.

The microalga *Nannochloropsis gaditana* CCAP 849/5 was maintained in 250-mL conical flasks

containing 50 mL of artificial sea water (ASW) at 25 ± 1 ^{o}C. All cultures were regularly sub-

cultured.

Cultures of *C. echinulata* and *U. isabellina* were performed in 250-mL Erlenmeyer

flasks containing 50 mL of a culture medium with the following composition (in g/L): glucose

(AppliChem, Darmstadt, Germany), 60.0; KH_2PO_4 (AppliChem), 12.0; Na_2HPO_4 (AppliChem),

12.0; $CaCl_2 \cdot 2H_2O$ (Carlo Erba, Rodano, Italy), 0.1; $CuSO_4 \cdot 5H_2O$ (BDH, Poole, England), 0.0001;

$Co(NO_3) \cdot 6H_2O$ (Merck, Darmstadt, Germany), 0.0001; $MnSO_4 \cdot 5H_2O$ (Fluka, Steinheim,

Germany), 0.0001; $ZnSO_4 \cdot 7H_2O$ (Merck), 0.001 and $FeCl_3 \cdot 6H_2O$ (BDH), 0.08. The medium was

limited in nitrogen with yeast extract (Conda, Madrid, Spain) at 3.0 g/L being the sole nitrogen

source. Yeast extract was also served as source of magnesium and ferrum according to Bellou et

al. [13]. The flasks were sterilized at 121 °C for 20 min and inoculated with 1 mL of spore

suspension containing 10^7 fungal spores produced on PDA cultures for 5 days at 28 °C.

Incubation took place in an orbital shaker (ZHICHENG ZHWY 211C, Shanghai, China) at

temperature 28 ± 1 °C and an agitation rate of 180 rpm. pH after sterilization was 6.5 ± 0.5 and

remained practically stable during cultivation.

A modified ASW described in Dourou et al. [22] was used as growth medium for *N.*

gaditana. Prior to sterilization pH of ASW was calibrated at 8.5 ± 0.5 through the addition of 2

M NaOH (Merck) solution. Microalgal cultures were performed in a laboratory-made glass

bioreactor of total volume 8.7 L and working volume 5.0 L, served as an open-pond simulating

reactor (OPSR) [22]. Initially, the reactor was washed with 70% ethanol and filled with 4.5 L

sterilized (at 121 °C for 20 min) ASW medium. The reactor was inoculated with 500 mL of a

fresh inoculum containing 10^5 cells/mL, and incubated at temperature 25 ± 1 °C under constant

illumination of 300 μmol m^{-2} s^{-1} supplied by linear fluorescent day light tubes T5, 8W, 65OOk,

G5. OPSR cultures were performed at temperature 25 ± 1 °C and pH 8.5 ± 0.5, which was

automatically controlled. Agitation was achieved through the use of a circulator, in the entryway

of which natural air was provided to the culture with a gas flow rate of 30 L/h. Illumination of

245 μmol m^{-2} s^{-1} was provided by 8 W fluorescent lamps, which were placed at a distance of 20

cm above the culture surface.

Cell mass harvesting

Fungal mycelia were harvested by filtration through Whatman No. 1 paper. Microalgal cell mass

was harvested by centrifugation at 24,000 g for 15 min at 4 °C (Heraeus, Biofuge Stratus,

Osterode, Germany). In both cases, the collected biomass was washed twice with distilled water,

dried at 80 °C until constant weight and gravimetrically determined.

Lipid extraction and purification

Microbial lipids were extracted in chloroform: methanol (2:1, v/v) (Sigma-Aldrich) following the

Folch et al. [27] method. The extracts were filtrated through Whatman No. 1 paper and washed

with a KCl (Sigma-Aldrich) 0.88 % (w/v) solution to remove non-lipid components.

Subsequently, the solvents were dried over anhydrous Na_2SO_4 (Sigma-Aldrich) and evaporated

under vacuum using a Rotavapor R-20 device (BUCHI, Flawil, Switzerland). The total cellular

lipids (L) were gravimetrically determined and expressed as a percentage on dry cell mass (L/x,

%).

Fatty acid methyl esters preparation and gas chromatography analysis

The FA moieties of lipids (i.e. approx. 100 mg of microbial oils or Olive oil or EPA concentrate

produced as above described) were converted into their fatty acid methyl esters (FAMEs) in a

two-stage reaction in accordance with the AFNOR [28] method in order to avoid trans-

isomerization. Briefly, in the first stage the FAs that are esterified with glycerol were converted

into FAMEs and the free FAs (if present) were converted into sodium soaps in a sodium

methoxide solution under reflux. Following, the resulting soaps were also converted into FAMEs

after adding an acetyl chloride solution in excess in the above mixture. The reaction was stopped

by adding water and the FAMEs were extracted in 6 mL hexane (Fluka). Finally, the organic

phase was removed under vacuum and the FAME preparation was stored in dark under an argon

atmosphere.

FAME mixtures were analysed in a gas chromatography device (Agilent 7890A device,

Agilent Technologies, Shanghai, China), equipped with a flame ionization detector (working at

280 °C) and a HP-88 (J&W Scientific) column (60 m × 0.32 mm). Carrier gas was helium at a

flow rate 1 mL/min and the analysis was run at 200 °C. Peaks of FAMEs were identified through

comparison to authentic standards.

Free fatty acid preparation

For glycerides cleavage, 1 g of lipids was saponified in 10 mL KOH 1N ethanol solution (95%)

under reflux for 1 h and 45 min. The mixture was acidified with 10 mL HCl 4 N solution and the

free FAs were extracted three-times with 5 mL hexane. The organic phase was washed with

distilled water until the washes were neutral and dried over anhydrous Na_2SO_4 (Sigma). Finally,

the organic phase was removed under vacuum and the FFA preparation was stored under an

Argon atmosphere.

Enzymatic synthesis of amides

The lipase-catalyzed synthesis of amides was carried out in 50-mL Erlenmeyer flasks in nearly

anhydrous media using 100 mg Novozym 435 lipase (i.e. immobilized *C. antarctica* lipase,

enzymatic activity □2 Units/mg) or 100 mg lipase from *C. rugosa* (immobilized, enzymatic

activity □2 Units/mg), both purchased from Sigma Aldrich Co., St. Louis, MO, as biocatalysts.

The reaction was carried out in an orbital shaker at 40 ± 1 ^{o}C, 90 rpm, in 25 mL acetone (Sigma

Aldrich Co.), with ethylene diamine (Acros Organics, Thermo Fisher Scientific, Waltham, MA)

and FAMEs or FFAs (produced as above) as substrates, used at different molar ratios. After

several preliminary experiments FAME preparations were selected as substrate. The reaction

lasted until the FAME substrate was exhausted, as evidenced by TLC (see below), usually after

18 h of incubation. The reaction mixture was then filtered, the solvent was removed from the

filtrate by evaporation under reduced pressure and the reaction residue is partitioned in

dichloromethane and distilled water (20 mL each). The organic layer, containing the synthesized

amide, was washed with saturated aqueous NaCl (Sigma Aldrich Co.) (10 mL), dried over

$MgSO_4$ (Sigma Aldrich Co.), gravity-filtered and the solvent removed under reduced pressure to

get the crude product.

Monitoring the evolution of the reaction and product characterization

Thin-layer chromatography and FT-IR

The reaction was monitored by thin-layer chromatography (TLC) performed on precoated Merck

60 GF254 silica gel plates (Merck, US) with a fluorescent indicator, and visualized under

ultraviolet irradiation at 254 and 360 nm. A mixture of n-hexane: ethyl acetate (1:4) was used as

eluent and the progress of the reaction monitored until the disappearance of FAME spot in the

reaction mixture.

FT-IR spectra for FAMEs and the FAA products were recorded on a Smart iTR, which

is an ultrahigh-performance, versatile attenuated total reflectance sampling accessory on the

Nicolet iS10 FT-IR spectrometer (Thermo Fisher Scientific). FT-IR spectra were used to confirm

the formation of FAAs from FAMEs, by detecting the formation of the amidic carbonyl group.

Quantitative determination of the reaction yields through *in situ* NMR monitoring

The percent conversion of FAMEs to amides was calculated during the reaction via *in situ* NMR

monitoring. In detail, the protons of methyl group of FAMEs, which are present at δ 3.56 ppm,

were assigned and the progress of the reaction of FAMEs with ethylene diamine was monitored

by ^{1}H NMR at regular intervals of 6 h. This was achieved by drawing a sample from the reaction

mixture using a 1mL syringe connected with a syringe filter of 0.22 µm pore size, followed by

evaporation of the solvent and dissolution of the residue in $CDCl_3$. The ^{1}H NMR was noted a

new singlet signal at δ 3.91 ppm which matched the $-CH_2-CH_2-$ of the amine used and grew

concurrently with a decline in the intensity of the methyl of ester group signals. The latter signals

disappeared after 24 h for ratio of FAME: amine 1:5 indicating 100% conversion. Therefore, we

succeed to calculate the % conversion via integrations of the peaks originated one from the

product (p) and the other from the reactant (r) according to the formula:

$$\% \text{ Conversion} = Ip / (Ir+Ip)$$

Where Ip is the integration of the signal of the product and Ir is the integration of the signal of

the reactant.

The physical properties and spectral data of the prepared diamides are described below.

Diamide of *Cunninghamella echinulata* methyl esters: Viscous liquid; black; IR (v_{max}, cm^{-1}):

3266 (NH), 2924, 2853 (CH$_2$), 1651 (CO amidic); Diamide of *Umbelopsis isabellina* methyl

esters: Liquid; dark brown; IR (v_{max}, cm^{-1}): 3276 (NH), 2924, 2854 (CH$_2$), 1663 (CO amidic);

Diamide of *Nannochloropsis gaditana* methyl esters: Viscous liquid; Yellow , IR (v_{max}, cm^{-1}):

3266 (NH), 2925, 2854 (CH$_2$), 1663 (CO amidic); Diamide of Olive oil methyl esters: Semi solid

material; off-white color; IR (v_{max}, cm^{-1}): 3407 (NH), 2922, 2852 (CH$_2$), 1638 (CO amidic);

Diamide of EPA concentrate methyl esters: Liquid material; orange color; IR (v_{max}, cm^{-1}): 3291

(NH), 2959, 2925 (CH$_2$), 1663 (CO amidic).

Antimicrobial evaluation of FAAs

The antimicrobial activity of the synthesized FAAs was tested *in vitro* using agar well diffusion

assay, MIC and MBC (see below) against human pathogens including the Gram-negative

Escherichia coli ATCC 25922, *Klebsiella pneumoniae* ATCC 700603, *Pseudomonas aeruginosa*

ATCC 15442, *Salmonella typhimurium* ATCC 14028 the Gram-positive bacteria, *Bacillus*

subtilis ATCC 6633*,* MRSA *Staphylococcus aureus* ATCC 4330, *Staphylococcus aureus* ATCC

25923 and the unicellular fungus *Candida albicans* ATCC 10221.

Agar well diffusion assay

Fresh bacterial cultures grown on nutrient agar for 20 h at 37 °C were suspended in a saline

solution (0.85%, w/v) to a turbidity of 0.5 Mac-Farland standards. Then 100 µl (10^6 CFU/mL)

of each bacterial suspension was swabbed onto Mueller Hinton II Agar (MHA) plates. 6 mm

diameter wells were punched on the MHA and inside the wells 100 μl of FAAs solution was

poured. The plates were preincubated in a refrigerator (at T=4 °C) for 1 h and then incubated

overnight at 37 °C, in order to allow the FAAs diffusion into the agar. The diameter of the

inhibition zones were measured in mm using Clinical and Laboratory Standards Institute (CLSI)

guidelines. Experiments were done in triplicate.

Evaluation of minimum inhibitory concentration (MIC) and minimum bactericidal concentration

(MBC) values for FAAs

MICs were determined according to the CLSI broth microdilution method [29]. One hundred

microliters of the Mueller Hinton broth medium were distributed into the wells of the micro titer

plates. A FAA solution (10 μL), serial diluted from stock solutions to achieve 200, 100, 50, 25,

12.5 and 6.25 µg/mL, was added to the well together with one hundred microliters of bacterial

suspension (6 x 10^6 CFU). The microwell plates were incubated at 37 °C for 24 h, then 5 μl of

a resazurin solution (6.75 mg/mL) was added to each well and the plates incubated at 37 °C for

another 4 h. Changes of color indicating cell viability were recorded. The bacterial growth was

measured using a Bio-Rad Microplate Reader at 600 nm. MIC was determined as the lowest

concentration of FAAs that inhibit visible growth of the tested microorganism. MBC was the

lowest FAA concentration resulting in microbial death. It was determined by sub-culturing cells

from wells that exhibited no color change to sterile MHA plates. All experiments were carried

out in triplicate.

Larval bioassay

Tests were performed on a field strain of *Aedes aegypti* raised from wild larvae, collected from

Jeddah, Saudi Arabia, and maintained in the laboratory under controlled conditions of 27 ± 1 ℃

and 70 ± 5% R.H., with a 14:10 (L:D) photoperiod. The standard World Health Organization

larval susceptibility test method was used. Treatments were carried out by exposing early 4th

instar larvae of *A. aegypti* to various concentrations of the tested compounds for 48 h, in groups

of glass beakers containing 100 mL of a FAA solution in tap water. Five replicates per FAA

concentration of 20 larvae each, and so for control trials, were set up. The larvae were given the

usual larval food during these experiments. Larval mortalities were recorded at 48 h post-

treatment. Log concentration-probability regression lines were drawn for the tested compounds.

Statistical parameters were calculated using the method of Finney [30].

Quantitative analysis of cell apoptosis by flow cytometry

The apoptotic activity of the SKOV-3 ovarian cancer cell line in response to tested compounds

was determined by Annexin FITC, as per the manufacturer's instructions (BD Biosciences,

USA). Briefly, the SKOV-3 cells were grown in a 25-mL flask at a density of 3×10^5 cells/well.

The induction of apoptosis was investigated in untreated and treated SKOV-3 cells with

curcuminoids at a concentration of 30 µM for 48h. After harvesting by trypsinization and

washing with PBS, the cells were stained with 5 µL Annexin FITC, incubated for 15 min in the

dark and then immediately analyzed using a FACS flow cytometer (BD FACSAria™ II - BD

Biosciences) using BD FACSDiva™ Software (BD Biosciences, USA).

Statistical analysis

The acquired data were analyzed using SPSS 9.0 and the results were given as mean ± SD of

three replicates. The mean comparison between the various assessed groups was performed using

one-way analysis of variance (ANOVA). Statistical significance was defined when $p < 0.05$.

Results and discussion

Lipid production and FA composition

Two oleaginous fungi, i.e. *C. echinulata* and *U. isabellina*, as well as the marine microalgae *N.*

gaditana, were selected for this study thanks to their ability to accumulate PUFA-containing

lipids.

294   *C. echinulata* has been recognized as a great GLA producer cultivated in sugar-based

media with high C/N ratio [31] while, *U. isabellina* is known for its capability to accumulate

lipids in high quantities [32-34]. In this study, *C. echinulata* produced significant quantities of

biomass and cellular lipids (i.e. 12.9 g/L of dry biomass containing 30.0% w/w lipids) while *U.*

isabellina accumulated 74% of lipids in its dry biomass, both cultivated in a mineral medium

with glucose as the sole source of carbon and energy (Table 1). GLA was found in considerable

concentration in *C. echinulata* lipids, representing 12.8% of total FAs (Table 2). However, the

major FA in these lipids was oleic (C18:1) (with a percentage of 44%), followed by palmitic

(C16:0) and linoleic (C18:2) acids. Concerning the FA profile of *U. isabellina*, C18:1 was the

dominant FA, found up to 54.4% in total lipids, while C16:0 and C18:2 were also found at

significant percentages. GLA percentages were low (i.e. 2.6%) in the lipids of *U. isabellina*.

Chatzifragkou *et al*. [35] reported slightly higher GLA percentages in the lipids of both strains

compared to the current study. It seems that several factors, such as the carbon source, affect

lipid FA composition. For instance, Fakas *et al*. [36], studying the effect of different carbon

sources on growth and lipid accumulation rates of *C. echinulata* and *U. isabellina*, reported that

xylose, in contrast to glucose, induced lipid accumulation and GLA biosynthesis.

In the last few decades, the interest in microalgae as PUFA producers is constantly

increasing. Biomass production and lipid accumulation of *N. gaditana* cultivated in ASW under

constant illumination, were satisfying (i.e. 313.9 mg/L and 22.7%, respectively) and in

accordance with data reported by Dourou et al. [22]. The predominant FA was EPA, found at a

percentage of 25% in the total FAs, followed by the monounsaturated palmitoleic (C16:1) and

C18:1 acids (Table 2). Dourou et al. [22] reported that the same strain cultivated on different

bioreactor configurations, was able to synthesize myristic acid (C14:1) too, which can be utilised

in a wide variety of applications in the cosmetics industry. The unsaturated FA content, though,

could potentially be increased by optimizing the growth conditions and/or by using genetically

modified strains.

Optimization of the FAA synthesis

Initially, the conditions of the amidation reaction (Fig. 1) were optimized by taking the Olive oil

FAMEs as a model substrate. The reaction, the progress of which was monitored by TLC, has

been done for 24 h at 40 ^{o}C with shaking at 90 rpm utilizing acetone as a solvent in the presence

of immobilized lipase as a catalyst. The % conversion, which was taken as a criterion for

determining the optimum conditions, was quantified via *in situ* NMR monitoring (Fig. 2).

Besides, FT-IR analysis (Fig. 3) gave a reliable evidence for amide formation due to the

appearance of a band at 1638 cm^{-1}, corresponding to the carbonyl of amide, in parallel with the

disappearance of the band at 1743 cm^{-1}, due to the consumption of the carbonyl group of

FAMEs. In addition, amide formation is confirmed by the appearance of a broadband at 3403

cm^{-1} due to NH, which in line with the amide structure 4 and rule out the formation of structure 3

due to the absence of the amino group NH$_2$ band (Fig. 1).

Two immobilized lipases, namely Novozym 435 and lipase from *C. rugosa* (Lipase CR)

were used as catalysts for the reaction of Olive oil FAMEs with ethylene diamine (Table 3). The

conversion yield was only 12% in the absence of a catalyst (entry 9, Table 3), confirming the

importance of lipases in FAA synthesis. Wang et al. [2, 3] proved that even in the absence of a

catalyst amidation can be performed but at high temperature and long reaction times, conditions

that create undesired product quality. Additionally, it was demonstrated that, under the

conditions of the present experimental work, the immobilized lipase Novozym 435 showed

lower conversion yield (entries 1-4, Table 3) than the Lipase CR (entries 5-8, Table 3).

Immobilized enzymes have the advantage over free enzymes to be easily recycled, providing

sustainability to the process, and for this reason several researchers proposed the employment of

immobilized lipases as catalysts for FAA synthesis [37-39]. In the current work the reusability of

the Lipase CR was checked for several reaction cycles for the synthesis of amide of Olive oil

FAMEs under the optimized reaction conditions. Specifically, the enzyme was removed after the

completion of the reaction by filtration, washed with ethanol solvent in a Soxhlet extraction

apparatus and the recovered enzyme was reused for three times under the same reaction

conditions. It was found that the regenerated enzyme performed the reaction efficiently without

loss of its catalytic activity. Similarly, according to Khare et al. [8], Novozym 435 could be

repeatedly used without any decrease of its catalytic activity, while Sharma et al. [40] have

demonstrated six repeated cycles of reusability of Chirazyme L-2 used to synthesize secondary

amide surfactant from N-methylethanol amine.

Previously, Wang et al. [2] showed that the molar ratio of vinyl stearate to ethanolamine

has affected the synthesis and purity of N-stearoyl ethanolamine produced. Thus, in the present

research different molar ratios of Olive oil FAMEs: ethylene diamine were tested (Table 3, Fig.

4). The amine rather than the FAME concentration in the reaction medium affected the

conversion yield of Olive oil FAMEs to amide. The maximum conversion yield (i.e. 100 %) was

attained with Lipase CR and a ratio of Olive oil FAMEs: ethylene diamine 1:5 (Table 3, entry 8).

These results are almost similar with those reported by Liu et al. [41] who, studying the effect of

the FA: diethanolamine ratio on the lipase-catalyzed amidation, showed that the maximum amide

yield was achieved at a low ratio 1:4.

The reaction conditions were further optimized using the immobilized Lipase CR as a

catalyst in different quantities and the reaction progress was monitored by TLC. It was found that

a 100% conversion yield was obtained in the shortest reaction time when 0.1 g of the Lipase CR

was employed (Table 4, entry 2), while a higher enzyme quantity was not necessary. Wang et al.

[3] reported that the yield of the amidation reaction increased when lipase concentration

increased from 10 to 20% of the total reactants.

Due to the nature of the reactants organic solvents were used by many researchers as a

suitable medium for the production of FAAs (see for instance [41, 42]). In the current paper

when acetone was used as a solvent the duration of the reaction was shorter than when *t*-butyl

alcohol and isoamyl alcohol were employed (entries 2, 4, 5, Table 4). Acetone is an

environmentally benign and low toxicity solvent, previously used as the best solvent for

polyunsaturated FAA synthesis [43].

Following optimization of the amidation reaction FAAs were produced using FAMEs of

SCOs from *C. echinulata*, *U. isabellina* and *N. gaditana* and of EPA concentrate oil (Table 5).

The conversion yields of the above reactions were excellent, reaching the values of 90-100%

(Table 5). The structures of the obtained FAAs were confirmed on the basis of their FT-IR

spectra in which appearance of the broadband due to NH group and of carbonyl group of amides

was observed (see for instance Fig. 5, 6 and original spectra in Fig. S1-S5). These results are in

agreement with Mudiyanselage et al. [44] who showed that microalgal lipids can be converted

into FAAs in a two-step reaction, including transesterification to form FAMEs followed by

amidation. The synthesis of amide in this work is efficient and its potential application on a large

scale will not interfere with the food supply chain, since SCOs are alternative sources to the

traditional sources of PUFAs.

Antimicrobial activity of amide compounds

FAAs derived from FAMEs of *C. echinulata*, *U. isabellina*, *N. gaditana* SCOs, Olive oil and

EPA concentrate were tested against various human pathogens for their antimicrobial efficacy by

the agar well diffusion method, which resulted in the formation of a variable diameter zone of

inhibition (Table 6). Except for MRSA *Staphylococcus aureus*, which is inhibited only by *N.*

gaditana-FAAs, Olive oil-FAAs and EPA-FAAs, all tested pathogens were significantly

inhibited by all FAAs produced in this work. *U. isabellina*-FAA was probably the most efficient

preparation against all pathogens, except for *MRSA Staphylococcus aureus*. On the contrary, *C.*

echinulata-FAAs seemed to be less efficient than *U. isabellina*-FAAs against all pathogens

(statistically significant at $p < 0.05$). *N. gaditana*-FAAs successfully inhibited all tested

organisms, except for *Bacillus subtilis* ATCC 6633 in the culture of which the inhibition zone

was only 9.00 ± 0.00 mm. The inhibition demonstrated by *N. gaditana*-FAAs against

Staphylococcus aureus ATCC 25923 and *Candida albicans* ATCC 10221 was similar to that

observed when *U. isabellina*-FAAs were employed. The Olive oil-FAAs showed a significant

inhibitory activity against all tested organisms, especially against *Pseudomonas aeruginosa*

ATCC 15442 and *Candida albicans* ATCC 10221 presenting an inhibition zone, 17.67 ± 0.57

mm and 18.07 ± 0.11 mm, respectively. Finally, EPA-FAAs showed a high antimicrobial

activity against all tested organisms specifically against *Staphylococcus aureus* (both strains) and

Pseudomonas aeruginosa ATCC 15442 (i.e. inhibition zone around 20 mm).

The results of MIC and MBC determined for selected pathogens (Table 7) were in line

with those obtained by the agar diffusion method. In detail, all tested pathogenic strains are

sensitive to the *U. isabellina*-FAAs, while *C. echinulata*-FAAs are less effective. For *N.*

gaditana-FAAs, the highest MIC observed was 200 µg/mL and this was against *Bacillus subtilis*

ATCC 6633, while the other pathogens tested were much more sensitive. All strains are sensitive

to FAAs derived from Olive oil, especially *Bacillus subtilis* ATCC 6633 and *Pseudomonas*

aeruginosa ATCC 15442 in which MIC was only 25 µg/mL. Besides, the FAAs derived from

EPA concentrate significantly affected the growth of all the tested bacteria particularly of

Bacillus subtilis ATCC 6633 and *Klebsiella pneumoniae* ATCC 700603. With few exceptions

(case of EPA-FAAs) the MBC was estimated to be 100-200 µg/mL.

The results reported in this paper are in agreement with previous reports in which

various FAAs have been used as potential antimicrobial agents [7]. Khare et al. [8] showed that

FAAs possess a strong antimicrobial activity towards Gram-positive (such as *Bacillus subtilis*

and *Staphylococcus aureus*) and Gram-negative (such as *Proteus vulgaris* and *Klebsiella*

pneumoniae) bacteria. Concerning the mechanism of action of FAAs Novak et al. [45] observed

that FAAs containing an epoxy group exhibit a broad spectrum of antimicrobial activity, which

is further enhanced by unsaturation. Later, Stevens and Hofmeyr [46] indicated a disturbance in

the FA constituents of the cell plasma membrane which interferes with the proper membrane

functions leading to the loss of the integrity of the plasma lemma. This suggestion was further

strengthened by Shao *et al.* [47] who worked on the mechanism of action of oleamide observed

that this compound caused alterations in the FA composition of the cell membranes. In this paper

FAAs prepared using FAMEs rich in oleic acid (i.e. Olive oil-FAAs and *U. isabellina*-FAAs)

were more effective against pathogens than those prepared from FAMEs contained oleic acid in

lower concentration (i.e. *C. echinulata*-FAAs). Unexpectedly, the presence of GLA in the lipids

of *C. echinulata* did not improve the antimicrobial activity of *C. echinulata*-FAAs compared to

U. isabellina-FAAs. *N. gaditana*-FAAs and EPA-FAAs, although containing oleic acid in very

low concentrations, are both effective against all tested bacteria, and therefore their activity may

be attributed to their high EPA content.

Insecticidal activity assay of amide compounds

The yellow fever mosquito (*Aedes aegypti*) spreads dangerous human arboviruses that include

dengue, Zika, and chikungunya. Therefore, control of yellow fever mosquitoes is a critical public

health priority [48]. Chemical insecticides are a leading method of control, but they are

expensive, contribute to the development of insecticidal resistance, pose risks to the environment

and cause safety concerns to humans and non-target animal species.

The susceptibility of *Aedes aegypti* larvae to FAAs under laboratory conditions was

tested using dipping methods. The larvicidal activity of a product is usually improved by

increasing its concentration and exposure time [49]. Many FA-derived products manifest toxicity

to different mosquito species larvae [50, 51], and have been proposed as alternatives to

conventional mosquito larvicides. Komalamisra et al. [52] considered larvicidal compounds

exerting LC50□<□50 mg/L active, 50 mg/L<□LC50□<□100 mg/L moderately active,

100 mg/L <□LC50□<□750 mg/L effective, and LC50□>□750 mg/L inactive. Kiran et al. [53]

considered compounds with LC50□<□100 mg/L as exhibiting a significant larvicidal effect. In

the current study, *C. echinulata*-FAAs showed a strong insecticidal activity against *A. aegypti*

larvae with LC50 0.3 mg/L, which could be probably attributed to the presence of GLA in

significant concentrations, while Olive oil-FAAs, EPA-FAAs and *N. gaditana*-FAAs exhibited

active insecticidal effect, demonstrating LC50 18.3, 20.5 and 34.3 mg/L, respectively. Contrary,

U. isabellina-FAAs were less active, presenting a LC50 equal to 132.1 mg/L (Table 8, Fig. 7).

Many bioactive substances, such as plant essential oils [54, 55], FAs [56] and cyanobacterial

extracts [57, 58], have been tested against *A. aegypti* larvae. According to our knowledge, this is

the first report demonstrating a larvicidal activity of FAAs, in contrast with numerous works

dealing with plant-based derivatives [59].

Cell apoptosis of ovarian cancer cell line induced by FAAs

The results show that all FAAs produced in this work can induce apoptosis of the SKOV-3

ovarian cancer cell line. Higher percentage of apoptosis was observed in the cells treated with *N.*

gaditana-FAAs followed by EPA-FAAs, Olive oil-FAAs and *C. echinulata*-FAAs (i.e. 61.7,

54.7, 52.7 and 50.4, respectively). The results are presented in Fig. 8.

Santos et al. [60] suggested that the antiproliferative activity is influenced by the

structural variation in the FAAs. Several studies have shown that intake of EPA, the main FA of

Nannochloropsis sp. may play a role in the prevention of the development of different type of

cancer [61]. Particularly, EPA and DHA have been investigated as potential dietary-based agents

for breast cancer prevention [62], and they have been shown to exhibit multiple anticancer

mechanisms of action, including the alteration of cell signaling [63], inhibition of cell

proliferation [64], inflammation [65], metastasis [64, 65], as well as induction of apoptosis [64,

66]. Studies with synthetic FAAs showed antiproliferative activity against several tumor cells

[67] and therefore these amide mediators may provide promising new agents, active against

inflammatory and cancer diseases [68]. It seems that variation in the FA moieties on groups

attached to the nitrogen atom may be responsible for differences in antiproliferative profiles [69].

Conclusions

FAMEs derived from SCOs of different origin and FA composition can be used in FAA

(diamide) enzymatic synthesis catalyzed by immobilized lipases, such as the *Candida rugosa*

lipase. The reaction of FAA synthesis can be completed under environmentally friendly

conditions in 24 h, while both the solvent (acetone) and the enzyme can be recycled. The

biological activities (antimicrobial, insecticidal activity, anti-cancer) of the synthesized FAAs are

related, partially at least, to their FA profile. Therefore, we conclude that oleaginous

microorganisms, able of synthesizing a wide range of FAs, can be considered in the near future

as source FAs suitable for producing FAAs of different biological activities.

Author agreement

Hatim A. El-Baz, Ahmed M. Elazzazy, Tamer S. Saleh, Panagiotis Dritsas, Jazem A. Mahyoub,

Mohammed N. Baeshen, Hekmat R. Madian, Mohammed Alkhaled and George Aggelis have all

agreed to submission.

Acknowledgments

This work was funded by the *University of Jeddah*, Saudi Arabia, under grant No. (UJ-06-18-

ICP). The authors, therefore, acknowledge with thanks the University technical and financial

support.

Compliance with Ethical Standards

Conflict of Interest

The authors declare that there are no conflicts of interest.

References

1. Plastina, P., Meijerink, J., Vincken, J. P., Gruppen, H., Witkamp, R., & Gabriele, B. (2009).

Selective synthesis of unsaturated N-acylethanolamines by lipase-catalyzed N-acylation of

ethanolamine with unsaturated fatty acids. *Letters in Organic Chemistry*, *6*(6), 444–447.

https://doi.org/10.2174/157017809789124885.

2. Wang, X., Wang, T., & Wang, X. (2012). An improved method for the synthesis of N-stearoyl

and N-palmitoylethanolamine. *Journal of the American Oil Chemists' Society,* 89, 1305–1313.

https://doi.org/10.1007/s11746-012-2017-y.

3. Wang, X., Wang, X., & Wang, T. (2012). Synthesis of oleoylethanolamide using lipase.

Journal of Agricultural and Food Chemistry, *60*(1), 451–457.

https://doi.org/10.1021/jf203629w.

4. Dettori, L., Jelsch, C., Guiavarc'h, Y., Delaunay, S., Framboisier, X., Chevalot, I., & Humeau,

C. (2018). Molecular rules for selectivity in lipase-catalysed acylation of lysine. *Process

Biochemistry*, *74*, 50–60. https://doi.org/10.1016/j.procbio.2018.07.021.

5. Bourkaib, M. C., Delaunay, S., Framboisier, X., Hôtel, L., Aigle, B., Humeau, C., Guiavarc'h,

Y., & Chevalot, I. (2020). N-acylation of L-amino acids in aqueous media: evaluation of the

catalytic performances of *Streptomyces ambofaciens* aminoacylases. *Enzyme and Microbial

Technology*, *137*, 109536. https://doi.org/10.1016/j.enzmictec.2020.109536.

6. D. Adlercreutz, P. Tufvesson, Karlsson, A., & Hatti-Kaul, R. (2010). Alkanolamide

biosurfactants: techno-economic evaluation of biocatalytic versus chemical production.

Industrial Biotechnology, *6*(4), 204–211. https://doi.org/10.1089/ind.2010.6.204.

7. Tanvir, R., Javeed, A., & Rehman, Y. (2018). Fatty acids and their amide derivatives from

endophytes: new therapeutic possibilities from a hidden source. *FEMS Microbiology Letters*,

365(12), fny114. https://doi.org/10.1093/femsle/fny114.

8. Khare, S. K., Kumar, A., & Kuo, T. M. (2009). Lipase-catalyzed production of a bioactive

fatty amide derivative of 7,10-dihydroxy-8(E)-octadecenoic acid. *Bioresource Technology*,

100(3), 1482–1485. https://doi.org/10.1016/j.biortech.2008.08.011.

9. Dang, H. T., Kang, G. J., Yoo, E. S., Hong, J., Choi, J. S., Kim, H. S., Chung, H. Y., & Jung,

J. H. (2011). Evaluation of endogenous fatty acid amides and their synthetic analogues as

potential anti-inflammatory leads. *Bioorganic & Medicinal Chemistry*, *19*(4), 1520–1527.

https://doi.org/10.1016/j.bmc.2010.12.046.

10. Meijerink, J., Balvers, M., & Witkamp, R. (2013). N-acyl amines of docosahexaenoic acid

and other n–3 polyunsatured fatty acids—from fishy endocannabinoids to potential leads. *British*

Journal of Pharmacology, 169(4), 772–783. https://doi.org/10.1111/bph.12030.

11. Arjuna, A. (2014). Production of polyunsaturated fatty acids by fungi: a review. *International*

Journal of Pharma and Bio Sciences, *5*(3), 931–954.

12. Bellou, S., Baeshen, M. N., Elazzazy, A. M., Aggeli, D., Sayegh, F., & Aggelis, G. (2014).

Microalgal lipids biochemistry and biotechnological perspectives. *Biotechnology Advances*,

32(8), 1476–1493. https://doi.org/10.1016/j.biotechadv.2014.10.003.

13. Bellou, S., Triantaphyllidou, I. E., Aggeli, D., Elazzazy, A. M., Baeshen, M. N., & Aggelis,

G. (2016). Microbial oils as food additives: recent approaches for improving microbial oil

production and its polyunsaturated fatty acid content. *Current Opinion in Biotechnology*, *37*, 24–

35. https://doi.org/10.1016/j.copbio.2015.09.005.

14. Athenaki, M., Gardeli, C., Diamantopoulou, P., Tchakouteu, S. S., Sarris, D., Philippoussis,

A., & Papanikolaou, S. (2018). Lipids from yeasts and fungi: physiology, production and

analytical considerations. *Journal of Applied Microbiology*, *124*(2), 336–367.

https://doi.org/10.1111/jam.13633.

15. Kavadia, A., Komaitis, M., Chevalot, I., Blanchard, F., Marc, I., & Aggelis, G. (2001). Lipid

and γ-linolenic acid accumulation in strains of zygomycetes growing on glucose. *Journal of the*

American Oil Chemists' Society, *78*(4), 341-346. https://doi.org/10.1007/s11746-001-0266-3.

16. Papanikolaou, S., & Aggelis, G. (2019). Sources of microbial oils with emphasis to

Mortierella (*Umbelopsis*) *isabellina* fungus. *World Journal of Microbiology and Biotechnology*,

35, 63. https://doi.org/10.1007/s11274-019-2631-z.

17. Kothri, M., Mavrommati, M., Elazzazy, A. M., Baeshen, M. N., Moussa, T. A. A., &

Aggelis, G. (2020). Microbial sources of polyunsaturated fatty acids (PUFAs) and the prospect

of organic residues and wastes as growth media for PUFA-producing microorganisms. *FEMS*

Microbiology Letters, *367*(5), fnaa028. https://doi.org/10.1093/femsle/fnaa028.

18. Mozaffarian, D., & Wu, J. H. Y. (2011). Omega-3 fatty acids and cardiovascular disease:

effects on risk factors, molecular pathways, and clinical events. *Journal of the American College*

of Cardiology, *58*(20), 2047–2067. https://doi.org/10.1016/j.jacc.2011.06.063.

19. Alakhras, R., Bellou, S., Fotaki, G., Stephanou, G., Demopoulos N. A., Papanikolaou, S., &

Aggelis, G. (2015). Fatty acid lithium salts from *Cunninghamella echinulata* have cytotoxic and

genotoxic effects on HL-60 human leukemia cells. *Engineering in Life Sciences*, *15*(2), 243–253.

https://doi.org/10.1002/elsc.201400208.

20. Sayegh, F., Elazzazy, A., Bellou, S., Moustogianni, A., Elkady, A. I., Baeshen, M. N., &

Aggelis, G. (2016). Production of polyunsaturated single cell oils possessing antimicrobial and

anticancer properties. *Annals of Microbiology*, *66*, 937– 948. https://doi.org/10.1007/s13213-

015-1176-0.

21. Bellou, S., & Aggelis, G. (2012). Biochemical activities in *Chlorella* sp. and

Nannochloropsis salina during lipid and sugar synthesis in a lab-scale open pond simulating

reactor. *Journal of Biotechnology*, *164*(2), 318–329.

https://doi.org/10.1016/j.jbiotec.2013.01.010.

22. Dourou, M., Tsolcha, O. N., Tekerlekopoulou, A. G., Bokas, D., & Aggelis, G. (2018). Fish

farm effluents are suitable growth media for *Nannochloropsis gaditana*, a polyunsaturated fatty

acid producing microalga. *Engineering in Life Sciences*, *18*(11), 851–860.

https://doi.org/10.1002/elsc.201800064.

23. Bellou, S., Moustogianni, A., Makri, A., & Aggelis, G. (2012). Lipids containing

polyunsaturated fatty acids synthesized by zygomycetes grown on glycerol. *Applied*

Biochemistry and Biotechnology, *166*, 146–158. https://doi.org/10.1007/s12010-011-9411-z.

24. Ruan, Z., Hollinshead, W., Isaguirre, C., Tang, Y. J., Liao, W., & Liu, Y. (2015). Effects of

inhibitory compounds in lignocellulosic hydrolysates on *Mortierella isabellina* growth and

carbon utilization. *Bioresource Technology*, *183*, 18–24.

https://doi.org/10.1016/j.biortech.2015.02.026.

25. Harde, S. M., Wang, Z., Horne, M., Zhu, J. Y., & Pan, X. (2016). Microbial lipid production

from SPORL-pretreated Douglas fir by *Mortierella isabellina*. *Fuel*, *175*, 64–74.

https://doi.org/10.1016/j.fuel.2016.02.023.

26. Gardeli, C., Athenaki, M., Xenopoulos, E., Mallouchos, A., Koutinas, A. A., Aggelis, G., &

Papanikolaou, S. (2017). Lipid production and characterization by *Mortierella* (*Umbelopsis*)

isabellina cultivated on lignocellulosic sugars. *Journal of Applied Microbiology*, *123*(6), 1461–

1477. https://doi.org/10.1111/jam.13587.

27. Folch, J., Lees, M., & Stanley, G.H.S. (1957). A simple method for the isolation and

purification of total lipids from animal tissues. *The Journal of Biological Chemistry*, *226*(1),

497–509.

28. AFNOR. (1984). Recueil de normes françaises des corps gras, graines oléagineuses, produits

dérivés. Paris: Association française de normalisation.

29. Wiegand, I., Hilpert, K., & Hancock, R. E. W. (2008). Agar and broth dilution methods to

determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nature*

Protocols, *3*, 163–175. https://doi.org/10.1038/nprot.2007.521.

30. Finney, D. J. (1971). Probit analysis, 3[rd] edition, Cambridge University Press, Cambridge,

UK.

31. Gema, H., Kavadia, A., Dimou, D., Tsagou, V., Komaitis, M., & Aggelis, G. (2002).

Production of γ-linolenic acid by *Cunninghamella echinulata* cultivated on glucose and orange

peels. *Applied Microbiology and Biotechnology*, *58*, 303-307. https://doi.org/10.1007/s00253-

001-0910-7.

32. Papanikolaou, S., Komaitis, M., & Aggelis, G. (2004). Single cell oil (SCO) production by

Mortierella isabellina grown on high sugar content media. *Bioresource Technology*, *95*(3), 287–

291. https://doi.org/10.1016/j.biortech.2004.02.016.

33. Subramaniam, R., Dufreche, S., Zappi, M., & Bajpai, R. (2010). Microbial lipids from

renewable resources: production and characterization. *Journal of Industrial Microbiology &*

Biotechnology, *37*(12), 1271–1287. https://doi.org/10.1007/s10295-010-0884-5.

34. Dourou, M., Aggeli, D., Papanikolaou, S., & Aggelis, G. (2018). Critical steps in carbon

metabolism affecting lipid accumulation and their regulation in oleaginous microorganisms.

Applied Microbiology and Biotechnology, *102*(6), 2509–2523. https://doi.org/10.1007/s00253-

018-8813-z.

35. Chatzifragkou, A., Fakas, S., Galiotou-Panayotou, M., Komaitis, M., Aggelis, G., &

Papanikolaou, S. (2010). Commercial sugars as substrates for lipid accumulation in

Cunninghamella echinulata and *Mortierella isabellina* fungi. *European Journal of Lipid Science*

and Technology, *112*(9), 1048–1057. https://doi.org/10.1002/ejlt.201000027.

36. Fakas, S., Papanikolaou, S., Batsos, A., Galiotou-Panayotou, M., Mallouchos, A., & Aggelis,

G. (2009). Evaluating renewable carbon sources as substrates for single cell oil production by

Cunninghamella echinulata and *Mortierella isabellina*. *Biomass and Bioenergy*, *33*(4), 573–580.

https://doi.org/10.1016/j.biombioe.2008.09.006.

37. Gotor-Fernández, V., Busto, E., & Gotor, V. (2006). *Candida antarctica* lipase B: an ideal

biocatalyst for the preparation of nitrogenated organic compounds. *Advanced Synthesis &*

Catalysis, *348*(7-8), 797–812. https://doi.org/10.1002/adsc.200606057.

38. Tufvesson, P., Annerling, A., Hatti-Kaul, R., & Adlercreutz, D. (2007). Solvent-free

enzymatic synthesis of fatty alkanolamides. *Biotechnology and Bioengineering*, *97*(3), 447–453.

https://doi.org/10.1002/bit.21258.

39. Al-Mulla, E. A. J., Yunus, W. M. Z. W., Ibrahim, N. A. B., & Rahman, M. Z. A. (2010).

Enzymatic synthesis of fatty amides from palm olein. *Journal of Oleo Science*, *59*(2), 59–64.

https://doi.org/10.5650/jos.59.59.

40. Sharma, J., Batovska, D., Kuwamori, Y., & Asano, Y. (2005). Enzymatic chemoselective

synthesis of secondary amide surfactant from N-methylethanol amine. *Journal of Bioscience and*

Bioengineering, *100*(6), 662–666. https://doi.org/10.1263/jbb.100.662.

41. Liu, K. J., Nag, A., & Shaw, J. F. (2001). Lipase-catalyzed synthesis of fatty acid

diethanolamides. *Journal of Agricultural and Food Chemistry*, *49*(12), 5761–5764.

https://doi.org/10.1021/jf0107858.

42. Levinson, W. E., Kuo, T. M., & Kurtzman, C. P. (2005). Lipase catalyzed production of

novel hydroxylated fatty amides in organic solvents. *Enzyme and Microbial Technology*, *37*(1),

126–130. https://doi.org/10.1016/j.enzmictec.2005.02.001.

43. Tremblay, H., St-Georges, C., Legault, M., Morin, C., Fortin, S., & Marsault, E. (2014). One-

pot synthesis of polyunsaturated fatty acid amides with anti-proliferative properties. *Bioorganic*

& Medicinal Chemistry Letters, *24*(24), 5635–5638. https://doi.org/10.1016/j.bmcl.2014.10.084.

44. Mudiyanselage, A. Y., Yao, H., Viamajala, S., Varanasi, S., & Yamamoto, K. (2015).

Efficient production of alkanolamides from microalgae. *Industrial & Engineering Chemistry*

Research, *54*(16), 4060–4065. https://doi.org/10.1021/ie503980g.

45. Novak, A. F., Solar, J. M., Mod, R. R., Magne, F. C., & Skau, E. L. (1969). Antimicrobial

activity of some N-substituted amides of long-chain fatty acids. *Applied Microbiology*, *18*(6),

1050-1056.

46. Stevens, S., & Hofmeyr, J. H. S. (1993). Effects of ethanol, octanoic and decanoic acids on

fermentation and the passive influx of protons through the plasma membrane of *Saccharomyces*

cerevisiae. *Applied Microbiology and Biotechnology*, *38*, 656–663.

https://doi.org/10.1007/BF00182806.

47. Shao, J., He, Y., Li, F., Zhang, H., Chen, A., Luo, S., & Gu, J. D. (2016). Growth inhibition

and possible mechanism of oleamide against the toxin-producing cyanobacterium *Microcystis*

aeruginosa NIES-843. *Ecotoxicology (London, England)*, *25*(1), 225–233.

https://doi.org/10.1007/s10646-015-1582-x.

48. Powell, J. R. (2018). Mosquito-borne human viral diseases: Why *Aedes aegypti*? *The*

American Journal of Tropical Medicine and Hygiene, *98*(6), 1563–1565.

https://doi.org/10.4269/ajtmh.17-0866.

49. Rodrigues, A. M., Sampaio, C. de G., de Souza, J. S. N., Campos, A. R., da Silva, A. B. R.,

de Morais, S. M., & Martins, V. E. P. (2019). Different susceptibilities of *Aedes aegypti* and

Aedes albopictus larvae to plant-derived products. *Revista da Sociedade Brasileira de Medicina*

Tropical, *52*, e20180197. https://doi.org/10.1590/0037-8682-0197-2018.

50. Kannathasan, K., Senthikumar, A., Venkatesalu, V., & Chandrasekaran, M. (2008).

Larvicidal activity of fatty acid methyl esters of *Vitex* species against *Culex quinquefasciatus*.

Parasitology Research, *103*(4), 999–1001. https://doi.org/10.1007/s00436-008-1078-1.

51. Perumalsamy, H., Jang, M. J., Kim, J. R., Kadarkarai, M., & Ahn, Y. J. (2015). Larvicidal

activity and possible mode of action of four flavonoids and two fatty acids identified in *Millettia*

pinnata seed toward three mosquito species. *Parasites & Vectors*, *8*, 237.

https://doi.org/10.1186/s13071-015-0848-8.

52. Komalamisra, N., Trongtokit, Y., Rongsriyam, Y., & Apiwathnasorn, C. (2005). Screening

for larvicidal activity in some Thai plants against four mosquito vector species. *The Southeast*

Asian Journal of Tropical Medicine and Public Health, *36*(6), 1412–1422.

53. Ravi Kiran, S., Bhavani, K., Sita Devi, P., Rajeswara Rao, B. R., & Janardhan Reddy, K.

(2006). Composition and larvicidal activity of leaves and stem essential oils of *Chloroxylon*

swietenia DC against *Aedes aegypti* and *Anopheles stephensi*. *Bioresource Technology*, *97*(18),

2481–2484. https://doi.org/10.1016/j.biortech.2005.10.003.

54. Magalhães, L. A., Lima, M., Marques, M. O., Facanali, R., Pinto, A. C., & Tadei, W. P.

(2010). Chemical composition and larvicidal activity against *Aedes aegypti* larvae of essential

oils from four *Guarea* species. *Molecules (Basel, Switzerland)*, *15*(8), 5734–5741.

https://doi.org/10.3390/molecules15085734.

55. Aguiar, R. W. S., dos Santos, S. F., da Silva Morgado, F., Ascencio, S. D., de Mendonça

Lopes, M., Fernandes Viana, K., Didonet, J., & Ribeiro, B. M. (2015). Insecticidal and repellent

activity of *Siparuna guianensis Aubl.* (Negramina) against *Aedes aegypti* and *Culex*

quinquefasciatus. *PLOS ONE*, *10*(2), e0116765. https://doi.org/10.1371/journal.pone.0116765.

56. Harada, K., Suomalainen, M., Uchida, H., Masui, H., Ohmura, K., Kiviranta, J.,

Niku☐Paavola, M. L., & Ikemoto, T. (2000). Insecticidal compounds against mosquito larvae

from *Oscillatoria agardhii* strain 27. *Environmental Toxicology*, *15*(2), 114–119.

https://doi.org/10.1002/(SICI)1522-7278(2000)15:2<114::AID-TOX7>3.0.CO;2-P.

57. Berry, J. P., Gantar, M., Perez, M. H., Berry, G., & Noriega, F. G. (2008). Cyanobacterial

toxins as allelochemicals with potential applications as algaecides, herbicides and insecticides.

Marine Drugs, *6*(2), 117–146. https://doi.org/10.3390/md20080007.

58. Mizerakis, P., Stathopoulou, P., Tsiamis, G., Baeshen, M. N., Mahyoub, J. A., Elazzazy, A.

M., Bellou, S., Sakoulogeorga, E., Triantaphyllidou, I. E., Mazioti, T., Katsoris, P., & Aggelis,

G. (2017). Bacterial diversity of the outflows of a Polichnitos (Lesvos, Greece) hot spring,

laboratory studies of a *Cyanobacterium* sp. strain and potential medical applications. *Annals of*

Microbiology, *67*, 643–654. https://doi.org/10.1007/s13213-017-1293-z.

59. Kim, S. I., & Ahn, Y. J. (2017). Larvicidal activity of lignans and alkaloid identified in

Zanthoxylum piperitum bark toward insecticide-susceptible and wild *Culex pipiens pallens* and

Aedes aegypti. *Parasites & Vectors*, *10*(1), 221. https://doi.org/10.1186/s13071-017-2154-0.

60. dos Santos, D. S., Piovesan, L. A., D' Oca, C. R., Hack, C. R., Treptow, T. G., Rodrigues, M.

O., Vendramini-Costa, D. B., Ruiz, A. L., de Carvalho, J. E., & D' Oca, M. G. (2015).

Antiproliferative activity of synthetic fatty acid amides from renewable resources. *Bioorganic &*

Medicinal Chemistry, *23*(2), 340–347. https://doi.org/10.1016/j.bmc.2014.11.019.

61. Wall, R., Ross, R. P., Fitzgerald, G. F., & Stanton, C. (2010). Fatty acids from fish: the anti-

inflammatory potential of long-chain omega-3 fatty acids. *Nutrition Reviews*, *68*(5), 280–289.

https://doi.org/10.1111/j.1753-4887.2010.00287.x.

62. Siddiqui, R. A., Harvey, K. A., Xu, Z., Bammerlin, E. M., Walker, C., & Altenburg, J. D.

(2011). Docosahexaenoic acid: a natural powerful adjuvant that improves efficacy for anticancer

treatment with no adverse effects. *BioFactors (Oxford, England)*, *37*(6), 399–412.

https://doi.org/10.1002/biof.181.

63. Allam-Ndoul, B., Guénard, F., Barbier, O., & Vohl, M. C. (2017). A study of the differential

effects of eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) on gene expression

profiles of stimulated thp-1 macrophages. *Nutrients*, *9*(5), 424.

https://doi.org/10.3390/nu9050424.

64. Blanckaert, V., Ulmann, L., Mimouni, V., Antol, J., Brancquart, L., & Chénais, B. (2010).

Docosahexaenoic acid intake decreases proliferation, increases apoptosis and decreases the

invasive potential of the human breast carcinoma cell line MDA-MB-231. *International Journal*

of Oncology, *36*(3), 737–742. https://doi.org/10.3892/ijo_00000549.

65. Chen, H. W., Chao, C. Y., Lin, L. L., Lu, C. Y., Liu, K. L., Lii, C. K., & Li, C. C. (2013).

Inhibition of matrix metalloproteinase-9 expression by docosahexaenoic acid mediated by heme

oxygenase 1 in 12-O-tetradecanoylphorbol-13-acetate-induced MCF-7 human breast cancer

cells. *Archives of Toxicology*, *87*(5), 857–869. https://doi.org/10.1007/s00204-012-1003-3.

66. Ravacci, G. R., Brentani, M. M., Tortelli, T. C., Torrinhas, R. S., Santos, J. R., Logullo, A.

F., & Waitzberg, D. L. (2015). Docosahexaenoic acid modulates a HER2-associated lipogenic

phenotype, induces apoptosis, and increases trastuzumab action in HER2-overexpressing breast

carcinoma cells. *BioMed Research International, 2015*, 838652.

https://doi.org/10.1155/2015/838652.

67. Jóźwiak, M., Filipowska, A., Fiorino, F., & Struga, M. (2020). Anticancer activities of fatty

acids and their heterocyclic derivatives. *European Journal of Pharmacology*, *871*, 172937.

https://doi.org/10.1016/j.ejphar.2020.172937.

68. Giordano, C., Plastina, P., Barone, I., Catalano, S., & Bonofiglio, D. (2020). n-3

polyunsaturated fatty acid amides: new avenues in the prevention and treatment of breast cancer,

International Journal of Molecular Sciences, *21*(7), 2279. https://doi.org/10.3390/ijms21072279.

69. Burstein, S., & Salmonsen, R. (2008). Acylamido analogs of endocannabinoids selectively

inhibit cancer cell proliferation. *Bioorganic & Medicinal Chemistry*, *16*(22), 9644–9651.

https://doi.org/10.1016/j.bmc.2008.10.015.

Table 1 Biomass yield (x, g or mg/L) and lipid content (L/x, %) of the microorganisms used in this study as source of lipids. The cultures were performed in triplicate

Microorganism	x	L/x (%)
Cunninghamella echinulata	12.9 ± 0.9 g/L	30.0 ± 1.5
Umbelopsis isabellina	13.2 ± 1.2 g/L	74.0 ± 0.8
Nannochloropsis gaditana	313.9 ± 0.4 mg/L	22.7 ± 0.1

Table 2 Fatty acid composition of the methyl ester mixtures used as acyl-donors in the amide synthesis. Analyses were performed in three independent samples

Source of lipid	Fatty acid composition (%, w/w)										
	C16:0	C16:1 n-7	C18:0	C18:1 n-9	C18:2 n-6	C18:3 n-6	C18:3 n-3	C18:4 n-3	C20:1 n-9	C20:5 n-3	Others
Cunninghamella echinulata	15.9 ± 0.7	2.0 ± 0.5	7.9 ± 0.6	44.0 ± 1.4	13.0 ± 1.2	12.8 ± 1.0	nd	nd	nd	nd	4.4 ± 1.2
Umbelopsis isabellina	22.1 ± 1.1	3.6 ± 0.4	2.8 ± 0.4	54.4 ± 4.1	11.7 ± 0.9	2.6 ± 0.3	nd	nd	nd	nd	2.8 ± 0.7
Nannochloropsis gaditana	18.0 ± 0.7	20.4 ± 0.5	0.7 ± 0.0	13.7 ± 0.4	5.7 ± 0.1	0.8 ± 0.2	0.8 ± 0.1	nd	7.2 ± 0.4	25.0 ± 0.3	7.7 ± 2.7
Olive oil	12.2 ± 1.2	2.4 ± 0.2	2.7 ± 0.3	74.1 ± 1.1	7.0 ± 0.2	nd	0.4 ± 0.0	nd	nd	nd	1.2 ± 0.3
EPA concentrate	0.5 ± 0.0	0.5 ± 0.0	3.3 ± 0.2	10.0 ± 1.8	1.0 ± 0.1	0.8 ± 0.0	2.3 ± 0.2	2.4 ± 0.1	4.4 ± 0.8	72.3 ± 1.4	4.0 ± 0.5

Table 3 Synthesis of Olive oil-FAAs utilizing different immobilized lipases and different molar

ratio of Olive oil FAMEs: ethylene diamine

Entry	Immobilized enzyme	FAME: Amine Ratio	Conversion (%)
1	Novozym 435	1:2	31.15 ± 1.25
2	Novozym 435	1:3	57.05 ± 2.95
3	Novozym 435	1:4	76.30 ± 3.30
4	Novozym 435	1:5	95.00 ± 4.15
5	Lipase CR	1:2	39.02 ± 2.95
6	Lipase CR	1:3	71.11 ± 4.55
7	Lipase CR	1:4	89.50 ± 3.45
8	Lipase CR	1:5	100.00 ± 4.83
9	No catalyst	1:5	12.57 ± 0.43

Table 4 Optimization of reaction conditions for Olive oil-FAAs synthesis utilizing Lipase CR

Entry	Lipase CR (wt, g)	Solvent	Time (h)	Conversion (%)
1	0.05	Acetone	30	82.09 ± 2.15
2	0.10	Acetone	24	100.04 ± 3.48
3	0.15	Acetone	24	99.13 ± 4.22
4	0.10	t-butyl alcohol	30	98.95 ± 4.78
5	0.10	Iso-amyl alcohol	30	100.20 ± 3.45

Table 5 Synthesis of FAAs by Lipase CR in acetone medium utilizing FAMEs of different origin

Entry	Source of FAMEs	Time (h)	Conversion (%)
1	*Cunninghamella echinulata*	24	90.23 ± 2.51
2	*Umbelopsis isabellina*	18	95.36 ± 5.75
3	*Nannochloropsis gaditana*	18	90.95 ± 1.50
4	EPA concentrate	18	100.05 ± 3.00

Table 6 The antimicrobial activity of FAAs against pathogenic strains. Data represent the mean of the diameter of the inhibition zones of three replicates ±SD. The concentration of FAAs used to determine the diameter of the inhibition zones was 4 mg/mL

	Source of FAAs				
	Cunninghamella echinulata	*Umbelopsis isabellina*	*Nannochloropsis gaditana*	Olive oil	EPA concentrate
	Inhibition zone (mm)				
Escherichia coli (ATCC 25922)	9.00 ± 1.00	17.00 ± 0.00	16.1 ± 1.01	15.67 ± 1.15	14.0 ± 0.00
Klebsiella pneumoniae (ATCC 700603)	10.33 ± 0.58	16.00 ± 0.00	14.3 ± 0.15	13.57 ± 0.81	12.57 ± 0.11
Salmonella typhimurium (ATCC 14028)	9.33 ± 0.58	19.00 ± 0.0	17.33 ± 0.57	14.67 ± 1.15	16.00 ± 0.00
Pseudomonas aeruginosa (ATCC 15442)	12.00 ± 1.00	20.00 ± 0.5	18.00 ± 0.00	17.67 ± 0.57	20.00 ± 1.00
Bacillus subtilis (ATCC 6633)	14.17 ± 0.29	17.00 ± 0.17	9.00 ± 0.00	12.17 ± 1.28	17.00 ± 1.00
MRSA *Staphylococcus aureus* (ATCC 4330)	0.00 ± 0.00	0.00 ± 0.00	15.50 ± 0.50	13.67 ± 1.15	19.00 ± 1.00
Staphylococcus aureus (ATCC 25923)	9.17 ± 0.29	22.00 ± 0.5	21.00 ± 0.86	15.00 ± 0.00	20.00 ± 2.00
Candida albicans (ATCC 10231)	16.00 ± 2.00	21.00 ± 1.8	19.00 ± 0.5	18.07 ± 0.11	18.07 ± 0.11

Table 7 Determination of minimum inhibitory concentration (MIC, µg/mL) and minimum bactericidal concentration (MBC, µg/mL)

of FAAs against pathogenic strains

Test organisms	Source of FAAs									
	Cunninghamella echinulata		*Umbelopsis isabellina*		*Nannochloropsis gaditana*		Olive oil		EPA concentrate oil	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
Klebsiella pneumoniae ATCC 700603	200.00 ± 0.00	200.00 ± 0.00	50.00 ± 0.00	200.00 ± 0.00	50.00 ± 0.00	200.00 ± 0.00	83.33 ± 28.87	200.00 ± 0.00	50.00 ± 0.00	100.00 ± 0.00
Pseudomonas aeruginosa ATCC 15442	100.00 ± 0.00	183.33 ± 28.87	50.00 ± 0.00	100.00 ± 0.00	50.00 ± 0.00	200.00 ± 0.00	25.00 ± 0.00	100.00 ± 0.00	100.00 ± 0.00	200.00 ± 0.00
Bacillus subtilis (ATCC 6633)	66.67 ± 28.87	200.00 ± 0.00	66.67 ± 28.87	100.00 ± 0.00	200.00 ± 0.00	200.00 ± 0.00	25.00 ± 0.00	100.00 ± 0.00	25.00 ± 0.00	50.00 ± 0.00
Staphylococcus aureus (ATCC 25923)	100.00 ± 0.00	200.00 ± 0.00	83.33 ± 28.87	100.00 ± 0.00	50.00 ± 0.00	200.00 ± 0.00	100.00 ± 0.00	100.00 ± 0.00	50.00 ± 0.00	100.00 ± 0.00

Table 8 Susceptibility of *Aedes aegypti* larvae to FAAs under laboratory conditions by using dipping methods. Data represent the mean of six replicates

Source of FAAs	LC50 (mg/L)	Lower limit	Upper limit	RR
Cunninghamella echinulata	0.294	0.258	0.332	1.00
Umbelopsis isabellina	38.837	34.29	43.588	132.10
Nannochloropsis gaditana	10.073	5.481	23.647	34.26
Olive oil	5.391	4.228	6.685	18.34
EPA concentrate	6.039	2.592	10.733	20.54

Figure Captions

Fig. 1 Amidation reaction of ethylene diamine and FAMEs catalyzed by lipase

Fig. 2 *In situ* NMR monitoring for % conversion of Olive oil FAMEs to amide

Fig. 3 FT-IR analysis of Olive oil FAMEs and its amide

Fig. 4 % conversion of Olive oil FAMEs to amide in different ratios of Olive oil FAMEs:

ethylene diamine

Fig. 5 FT-IR analysis of EPA concentrate FAMEs and its amide

Fig. 6 FT-IR analysis of *Umbelopsis isabellina* FAMEs and its amide

Fig. 7 The larval mortality effect of FAAs of *C echinulata*, *U isabellina*, *N gaditana*, Olive oil

and EPA concentrate at different concentrations against *Aedes aegypti* after continuous exposure

for 48 hours

Fig. 8 Effect of compounds on SKOV-3 cell apoptosis. Flow cytometry analysis of apoptosis in

SKOV-3 cells either untreated or treated with 10 µg/mL of every compounds for 48h. After the

treatment period, the cells were stained with Annexin FITC and subsequently analyzed by flow

cytometry

Abbreviations: *C. echinulata* – FAAs: *C.e.* – FAAs; *U. isabellina* – FAAs: *U.i.* – FAAs; *N.*

gaditana – FAAs: *N.g.* – FAAs; Olive oil – FAAs; O.o. – FAAs; EPA concentrate – FAAs: EPA

– FAAs

R: alkyl group

time = 24h

time = 18h

time = 12h

New signal due to 2
sym. methylene group

time = 6h

Methyl group of Olive oil FAME

time = 0h

FT-IR of FAA

2852
2922

3403 cm⁻¹ (NH)

1638 cm⁻¹(−C(=O)N=)

FT-IR of olive oil methyl ester

2853
2923

1743 cm⁻¹(RCOO)

FT-IR of FAA

3276 cm^{-1}(NH)

1663 cm^{-1}(−C(=O)N=)

FT-IR of *Umbelopsis isabellina* methyl ester

2853 cm^{-1}

2923 cm^{-1}

1743 cm^{-1}(RCOO)