Self-reassurance reduces neural and self-report reactivity to negative life events

Jeffrey J. Kim[1,2]*, Ross Cunnington[1], & James N. Kirby[1]

[1]Compassionate Mind Research Group, School of Psychology, The University of Queensland, Brisbane, Queensland, Australia
[2]The Centre of Advanced Imaging, The University of Queensland, Brisbane, Queensland, Australia

*Correspondence:
Mr. Jeffrey J. Kim
School of Psychology
The University of Queensland
St Lucia
Brisbane QLD 4072
Australia

Phone: (07) 3365 6230
Fax: (07) 3365 4466
Email: Jeffrey.kim@uqconnect.edu.au

Abstract

Background: Whilst research has shown how self-criticism may increase both neural and self-report markers of negative emotion, less well known is how self-reassurance - a compassionately-motivated cognitive self-relating style - may regulate negative emotion.

Method: Using fMRI, we invited participants to engage in self-criticism and self-reassurance toward written descriptions of negative life events (mistakes, setbacks, failures).

Results: Our results identify that neural markers of negative emotion and self-report markers of trial intensity during fMRI are suppressed under conditions of self-reassurance, relative to self-criticism.

Limitations: Future work to control for autobiographical memory during this fMRI task is needed, to explore how memory can contribute to self-reassurance and self-criticism.

Conclusions: Engagement in self-reassurance can reduce the 'sting' of negative life-events, both neural and self-report, which holds important implications for therapy.

Keywords: Compassion; Self-Criticism; fMRI; Reassurance; Emotion

Introduction

Adverse life events are inescapable, be it a disruption in a career, dissolution of a relationship, or even a world-wide pandemic. These factors are known to take a toll on both physical and mental health outcomes (Solís et al., 2015) which can increase the likelihood of mortality (Puterman et al., 2020). These disappointments (e.g., making mistakes), losses (e.g., of hoped love) and fears (e.g., of rejection) are all triggers to self-criticism (Halamová et al., 2018; Kim et al., 2020). Indeed, self-criticism is a common self-relating style people use to cope, often resulting in an individual taking the frustration and anger out on themselves, which compounds the experience of pain psychologically and neurophysiologically (Kim et al., 2020). Whilst research has shown

how self-criticism may increase both self-report (Cox et al., 2000; Gilbert et al., 2004)

and neural (Longe et al., 2010; Lutz et al., 2016) markers of negative emotion, less well

known is how self-reassurance - a compassionately-motivated cognitive self-relating

style - may regulate how the brain responds toward negative life events.

Here we conducted an fMRI experiment which examined two distinct self-relating

styles, self-criticism and self-reassurance (Kim et al., 2020; Petrocchi et al., 2019),

when participants imagined themselves responding to mistakes, setbacks or failures.

Importantly, we designed our experiment to deliberately tease apart neural markers of

negative emotion, first by manipulating an emotional – neutral contrast at the first level

of fMRI analysis, and explored how this activation may differ across self-criticism and

self-reassurance. To anticipate our findings, we identified that how the brain encodes

negative emotion differs under conditions of self-criticism versus self-reassurance,

specifically showing how self-reassurance can down-regulate neural markers of

negative emotion.

70   **Methods:**

Whilst the program of research within the present paper has been reported on

previously, this examined the (neuro)physiological correlates of a brief, two-week

compassion training paradigm (Kim et al., 2020). Here we focus on the novel whole

brain markers of criticism and reassurance which have not been reported. As our fMRI

method as reported in the previous paper is also the same imaging method used for the

present paper, we have reproduced this section for clarity under a CC BY open access

licence.

Participants:

40 participants (Mean age = 22 years, SD = .49, 27 female) took part in the

present study. A University's Ethics Sub-Committee approved the experimental

protocol, and this project complies with the provisions contained in the *National*

Statement of Ethical Conduct in Human Research and complies with the regulations

governing experimentation on humans. Participants provided informed and voluntary,

written and/or electronic consent.

fMRI Stimuli:

We created 60 written stimuli in total, consisting of a personal mistake, setback

or failure. 30 statements were of emotional valence whereas 30 were neutral (i.e., "I fail

to keep up with my commitments in life", and "I keep up with my commitments in life",

respectively). Our neutral stimuli were created to describe a non-emotive, non-intense

control to counterbalance the emotional stimuli set. For both emotional and neutral sets

we assessed two metrics, valence (1-5, where 1 = Very Unpleasant) and intensity (1-5,

where 1 = Not Intense). Our emotional statements (n = 30) were revealed to be

sufficiently unpleasant (M = 1.89) and intense (M = 3.54), with all neutral statements (n

= 30) described as less unpleasant (M = 3.80) and comparatively not intense (M =

2.34).

fMRI Design:

Within the scanner we examined participant's neural responses to the validated

(affective and neutral) written stimuli when engaged in self-criticism and self-

reassurance (Figure 1). Participants were given one practice block with 16 trials (8

emotional and 8 neutral) repeated for self-criticism and self-reassurance, and made

button press ratings on perceived trial intensity. After each trial within a block of either

self-criticism or self-reassurance, participants rated how intense their degree of self-

criticism or self-reassurance was to each statement (button-press on an MR-compatible

button box which ranged from 1-4, where 1 = not very intense, and 4 = very intense). A

typical trial consisted of stimuli presented for a 6 second duration, followed by a rating of

intensity for a 3 second duration, and an inter-trial-interval of .5 seconds. The first order

of instruction for a particular block, that is, self-reassurance verses self-criticism, was

counterbalanced for a total of 8 blocks. As our focal contrast, we manipulated the

emotionality of the statements within scan runs ("emotive" vs "neutral"), in a

counterbalanced order across participants. 30 statements were quasi-randomized

across participants and presented for a total of 30 trials per fMRI run (~6.5 min total

duration) over a total of 8 repeated fMRI runs. Participants were given 10 practice trials

of emotional and neutral stimuli, and rated stimuli on intensity.

fMRI Acquisition and Pre-Processing:

We collected our fMRI data on a Siemens Magnetom Prisma Fit 3 Tesla MRI

scanner utilizing a 64-channel head-coil. A gradient-echo, echo-planar imaging

sequence was used to acquire functional images, with the following sequence

parameters: 60 horizontal slices (2 x 2-mm in-plane voxel resolution and 2-mm slice

thickness plus 10% gap), repetition time (TR) 1000 ms; echo time (TE) 30 ms. Eight

identical fMRI runs of 292 images (6 minutes each) were acquired. A 3D high-

resolution, unified and denoised T1-weighted MP2RAGE image across the entire brain

was also acquired and used as anatomical reference for subsequent pre-processing in

SPM12 (TR = 4000 ms, TE = 2.93 ms, FA = 6°, 176 cube matrix, voxel size = 1-mm).

Functional imaging data were pre-processed and analyzed using SPM12, implemented

in MATLAB. Structural T1-scans were co-registered to the average of the spatially

realigned functional slices. Next, an inbuilt segmentation routine was applied to register

each structural T1-image to the standard MNI template in MNI space. These transform

parameters elicited from segmentation were subsequently applied to all realigned

images, resliced to a 2x2x2-mm resolution and smoothed with 6-mm full-width-at-half-

maximum (FWHM) isotropic Gaussian kernel.

fMRI First and Second-Level Analyses:

For first-level data analysis, block-related neural responses to stimuli were

modelled as 2 separate conditions (all combinations of emotional/neutral, self-

criticism/self-reassurance) and convolved with the canonical hemodynamic response

function (HRF). For group level analysis, whole-brain contrasts of self-criticism

(emotional-neutral) stimuli were reported at a cluster-level threshold of $p < .05$,

corrected for family-wise error, with clusters formed with a voxel-level height threshold

at $p < .001$, uncorrected, with a cluster extent threshold of K = 144. Whole-brain

contrasts of self-reassurance (emotional-neutral) stimuli were reported at a cluster-level

threshold of $p < .05$, corrected for family-wise error, with clusters formed with a voxel-

level height threshold at $p < .001$, uncorrected, with a cluster extent threshold of K =

144. Whole-brain repeated-measures contrasts of self-criticism (emotional-neutral) -

self-reassurance (emotional-neutral) stimuli were reported at a cluster-level threshold

of $p < .05$, corrected for family-wise error, with clusters formed with a voxel-level height

threshold at $p < .001$, uncorrected, with a cluster extent threshold of K = 110. Brain

regions shown to be significant had their anatomical labels identified with the Automated

Anatomical Labelling (AAL) toolbox implemented in SPM12.

Self-report markers of intensity during fMRI:

Analysis of participant's mean level of intensity ratings for reassuring trials (emotional stimuli: $M = 2.45$, $SD = 0.48$, neutral statements: $M = 2.63$, $SD = 0.64$) and critical trials (emotional stimuli: $M = 2.92$, $SD = 0.45$, neutral stimuli: $M = 2.07$, $SD = 0.52$) revealed intensity ratings were significantly higher for critical (emotional – neutral) but not for reassuring (emotional – neutral) trials ($t(38) = 7.300$, $p < 0.001$, and $t(38) = -1.372$, $p = 0.178$, *ns*, respectively).

Results

First, group-level one-sample t-tests of the whole brain contrasts of emotional - neutral stimuli were conducted overall. We then examined how this contrast differed across self-criticism and self-reassurance. For neural markers of negative emotion during self-criticism, we observed activation in the "salience" (midcingulo-insular), "default-mode" (medial frontoparietal), and the occipital network (Uddin et al., 2019). Whilst neural markers of negative emotion during self-reassurance recruited activation in regions such as the medial-prefrontal cortex (MPFC) and visual cortex, we observed no activation of the salience network as shown under self-criticism. Across both these contrasts, clusters were formed at a cluster-level threshold of $p < .05$, corrected for family-wise error, with clusters formed with a voxel-level height threshold at $p < .001$, uncorrected (cluster extent threshold K = 144).

We next conducted a repeated-measures contrast between self-criticism (emotional – neutral) minus self-reassurance (emotional – neutral), as a marker of neural markers of negative emotion which differs between these two self-relating styles. Here, we identified brain activation across bilateral hippocampus (with a cluster which also included left putamen and left insula), thalamus, ACC, and occipital lobe, revealing

neural markers of negative emotion are driven by self-criticism but not self-reassurance

(cluster-level threshold of $p < .05$, corrected for family-wise error, with clusters formed

with a voxel-level height threshold at $p < .001$, uncorrected, with a cluster extent

threshold of K = 110). A repeated-measures contrast between self-reassurance

(emotional – neutral) minus self-criticism (emotional – neutral) returned non-significant.

Our experimental design is shown in Figure 1, and Figure 2 depicts the whole-brain

results. Tables which present the thresholded output for each contrast are available

online (https://osf.io/9wzu4/?view_only=91484c009daa4b03b676cbfa70940a6f).

Figure 1. Task diagram for a typical trial. Participants were presented with 30
alternating trials of emotional or neutral statements which describe a mistake, setback
or failure. Across 8 scan runs of 6 minutes each, participants were asked to engage with
these statements from two different perspectives – four blocks of self-criticism, and four
blocks of self-reassurance (order counterbalanced across participants). Example
statements are presented inset.

188
189
Figure 2. Neural markers of negative emotion across self-criticism and self-
reassurance. **Negative Emotion during Self-Criticism: Left.** Sagittal image of MPFC
(Left Circle), ACC (Middle Circle), and Left Lingual Gyrus and Cerebellum (Right Circle).
Right. Axial image of Subcortical Regions (Top Circle) and Bilateral Visual Cortex
(Bottom Circle). **Negative Emotion during Self-Reassurance: Left.** Sagittal image of
MPFC (Left Circle) and Visual Cortex (Right Circle). **Right.** Axial image of Visual
Cortex. **Negative Emotion during Self-Criticism – Negative Emotion during Self-**
Reassurance: Top Left. Sagittal image of ACC. **Top Right.** Sagittal image of posterior
cingulate. **Bottom Left.** Axial image of left putamen. **Bottom Right.** Axial image of
Right Hippocampus. Coordinates reported in MNI-space. $N = 40$.
200

Discussion

Here we investigated neural markers of negative emotion when participants engaged in self-criticism and self-reassurance toward negative life events (i.e., mistakes, setbacks or failures). Across both self-criticism and self-reassurance, our fMRI study revealed common activation across diverse regions such as the visual cortex (associated with mental imagery), salience network (associated with processing pain and threat), and default-mode network (associated with self-referential thought) (Uddin et al., 2019). Brain activation overall was more extensive for self-critical than self-reassuring trials, even though both contrasts did activate similar regions such as the MPFC and visual cortex. Furthermore, self-reassurance did not activate regions such as the insula, anterior cingulate cortex and amygdala. In addition, self-report ratings of intensity for emotional stimuli were supressed for self-reassuring versus self-critical trials. Importantly, a contrast of negative emotion between self-criticism minus self-reassurance revealed brain activation in regions such as the anterior cingulate cortex, insula and hippocampus. Taken together, our data show that neural and self-report markers of negative emotion are supressed during self-reassurance compared with self-criticism, providing evidence for how cultivating a reassuring self-relating style can regulate neural markers of negative emotion.

Whilst recruitment of the insula and anterior cingulate cortex have previously been shown for self-criticism (Hooley et al., 2012; Lutz et al., 2016), it is important to remark on bilateral hippocampus activation within the current experiment, which may be an indicator of autobiographical memory recall (Aly, 2020; McCormick et al., 2020). Given our paradigm instructions were for participants to engage in self-critical thoughts from the stimuli presented, it is entirely possible that for reference participants engaged

in their own first-person accounts from situations in their own lives (Holland &

Kensinger, 2010). However, given we did not explicitly control for autobiographical

memory, future work is needed to explore how memory may contribute to neural

markers of self-criticism or self-reassurance. To position our results in the broader

literature on the neuroscience of empathy and compassion, we have shown that brain

regions for processing negative emotion toward others (Ashar et al., 2017; Kim,

Cunnington & Kirby, 2020; Zaki et al., 2016) were shown to not be recruited during

compassion to the self. Specifically, we have shown that neural markers of negative

emotion are suppressed during attempts to be compassionate and reassuring to one's

suffering. Our data suggest that engagement in self-reassurance is a way to reduce the

'sting' of negative life-events, both neural and self-report, which is a timely finding in our

current global environment.

References

Aly, M. (2020). Spotlight Brain Dynamics Underlying Memory for Lifetime Experiences.

Trends in Cognitive Sciences, 1–2. https://doi.org/10.1016/j.tics.2020.06.010

Ashar, Y. K., Andrews-Hanna, J. R., Dimidjian, S., Wager, T. D. (2017). Empathic Care

and Distress: Predictive Brain Markers and Dissociable Brain Systems. *Neuron*,

94(6), 1263-1273.e4. https://doi.org/10.1016/j.neuron.2017.05.014

Cox, B. J., Rector, N. A., Bagby, R. M., Swinson, R. P., Levitt, A. J., & Joffe, R. T.

(2000). Is self-criticism unique for depression? A comparison with social phobia.

Journal of Affective Disorders, *57*(1–3), 223–228. https://doi.org/10.1016/S0165-

0327(99)00043-9

Gilbert, P., Clarke, M., Hempel, S., Miles, J. N. V., & Irons, C. (2004). Criticizing and

reassuring oneself: An exploration of forms, styles and reasons in female students.

British Journal of Clinical Psychology, *43*(1), 31–50.

https://doi.org/10.1348/014466504772812959

Halamová, J., Kanovský, M., Gilbert, P., Troop, N. A., Zuroff, D. C., Hermanto, N.,

Petrocchi, N., Sommers-Spijkerman, M., Kirby, J. N., Shahar, B., Krieger, T.,

Matos, M., Asano, K., Yu, F. Y., Basran, J., & Kupeli, N. (2018). The Factor

Structure of the Forms of Self-Criticising/Attacking & Self-Reassuring Scale in

Thirteen Distinct Populations. *Journal of Psychopathology and Behavioral*

Assessment, *40*(4), 736–751. https://doi.org/10.1007/s10862-018-9686-2

Holland, A. C., & Kensinger, E. A. (2010). Emotion and autobiographical memory.

Physics of Life Reviews, *7*(1), 88–131. https://doi.org/10.1016/j.plrev.2010.01.006

Hooley, J. M., Siegle, G., & Gruber, S. A. (2012). Affective and Neural Reactivity to

Criticism in Individuals High and Low on Perceived Criticism. *PLoS ONE*, *7*(9).

https://doi.org/10.1371/journal.pone.0044412

Kim, J. J., Cunnington, R., & Kirby, J. N. (2020). The neurophysiological basis of

compassion: An fMRI meta-analysis of compassion and its related neural

processes. *Neuroscience and Biobehavioral Reviews*, *108*, 112–123.

https://doi.org/10.1016/j.neubiorev.2019.10.023

Kim, J. J., Henderson, T., Best, T., Cunnington, R., & Kirby, J. N. (2020). Neural and

Self-Report Markers of Reassurance: A Generalized Additive Modelling Approach.

Frontiers in Psychiatry, *11*. https://doi.org/10.3389/fpsyt.2020.566141

Kim, J., Parker, S., Doty, J., Cunnington, R., Gilbert, P., & Kirby, J. (2020).

Neurophysiological and behavioural markers of compassion. *Scientific Reports*,

10(1), 6789. https://doi.org/10.1038/s41598-020-63846-3

Longe, O., Maratos, F. A., Gilbert, P., Evans, G., Volker, F., Rockliff, H., & Rippon, G.

(2010). Having a word with yourself: neural correlates of self-criticism and self-

reassurance. *NeuroImage*, *49*(2), 1849–1856.

https://doi.org/10.1016/j.neuroimage.2009.09.019

Lutz, J., Bruhl, A. B., Doerig, N., Scheerer, H., Achermann, R., Weibel, A., Jancke, L., &

Herwig, U. (2016). Altered processing of self-related emotional stimuli in

mindfulness meditators. *NeuroImage*, *124*(Pt A), 958–967.

https://doi.org/10.1016/j.neuroimage.2015.09.057

McCormick, C., Barry, D. N., Jafarian, A., Barnes, G. R., & Maguire, E. A. (2020).

vmPFC Drives Hippocampal Processing during Autobiographical Memory Recall

Regardless of Remoteness. *Cerebral Cortex*, 1–16.

https://doi.org/10.1093/cercor/bhaa172

Petrocchi, N., Dentale, F., & Gilbert, P. (2019). Self-reassurance, not self-esteem,

serves as a buffer between self-criticism and depressive symptoms. *Psychology*

and Psychotherapy: Theory, Research and Practice, *92*(3), 394–406.

https://doi.org/10.1111/papt.12186

Puterman, E., Weiss, J., Hives, B. A., Gemmill, A., Karasek, D., Mendes, W. B., &

Rehkopf, D. H. (2020). Predicting mortality from 57 economic, behavioral, social,

and psychological factors. *Proceedings of the National Academy of Sciences of the*

United States of America, *117*(28), 16273–16282.

https://doi.org/10.1073/pnas.1918455117

Solís, C. B., Kelly-Irving, M., Fantin, R., Darnaudéry, M., Torrisani, J., Lang, T., &

Delpierre, C. (2015). Adverse childhood experiences and physiological wear-and-

tear in midlife: Findings from the 1958 British birth cohort. *Proceedings of the*

National Academy of Sciences of the United States of America, *112*(7), E738–

E746. https://doi.org/10.1073/pnas.1417325112

Uddin, L. Q., Yeo, B. T. T., & Spreng, R. N. (2019). Towards a Universal Taxonomy of

Macro ￼ scale Functional Human Brain Networks How Many Functional Brain

Networks Are A fundamental construct in neuroscience is the definition. *Brain*

Topography, *0123456789*. https://doi.org/10.1007/s10548-019-00744-6

Zaki, J., Wager, T. D., Singer, T., Keysers, C., & Gazzola, V. (2016). The Anatomy of

Suffering: Understanding the Relationship between Nociceptive and Empathic Pain.

Trends in Cognitive Sciences, *20*(4), 249–259.

https://doi.org/10.1016/j.tics.2016.02.003

Author Contributions: JJK, RC and JNK conceived and designed the experiment; JJK
acquired, analysed and interpreted the data; JJK and JNK wrote the manuscript. All
authors have given final approval for submission.

Acknowledgements: JJK thanks Fernanda L. Ribeiro for helpful feedback and
comments on a manuscript draft.

Role of the Funding Source: This research was supported by a University of
Queensland Research and Teaching fellowship awarded to JNK. JJK was supported by
an Australian Postgraduate Scholarship.

Competing Interests Statement: The authors declare no competing interests.