1RS.1BL molecular resolution provides novel contributions to wheat improvement

Zhengang Ru[1,†], Angela Juhasz[2,†], Danping Li[3,†], Pingchuan Deng[4,†], Jing Zhao[3,†], Lifeng

Gao[3,†], Kai Wang[5,†], Gabriel Keeble-Gagnere[6], Zujun Yang[7], Guangrong Li[7], Daowen Wang[8],

Utpal Bose[9], Michelle Colgrave[2,9], Chuizheng Kong[3], Guangyao Zhao[3], Xueyong Zhang[3], Xu

Liu[3], Guoqing Cui[3], Yuquan Wang[1], Zhipeng Niu[1], Liang Wu[4], Dangqun Cui[8]★, Jizeng Jia[8,3]★,

Rudi Appels[10,6]★ and Xiuying Kong[3]★

[1] School of Life Science and Technology, Henan Institute of Science and Technology,

Xinxiang, China

[2] School of Science, Edith Cowan University, Joondalup, Australia

[3] The National Key Facility for Crop Gene Resources and Genetic Improvement, Institute of

Crop Sciences, Chinese Academy of Agricultural Sciences, Beijing, China

[4] State Key Laboratory of Rice Biology, College of Agriculture and Biotechnology, Zhejiang

University, Hangzhou, China

[5] Novogene Bioinformatics Institute, Beijing, China

[6] Agriculture Victoria Research, AgriBio, Centre for AgriBioscience, Bundoora, VIC,

Australia

[7] Center for Informational Biology, School of Life Science and Technology, University of

Electronic Science and Technology of China, Chengdu, China

[8]College of Agronomy, Collaborative Innovation Center of Henan Grain Crops, Henan

Agricultural University, Longzi Lake Campus, Zhengzhou, China

[9] CSIRO Agriculture and Food, 306 Carmody Rd, St Lucia, Australia

[10] University of Melbourne, Parkville, Australia

[†]These authors contributed equally to this article.

★Correspondence: (e-mails dangquncui@163.com; jiajizeng@caas.cn;

rudi.appels@unimelb.edu.au; kongxiuying@caas.cn).

Keywords: wheat, 1RS.1BL translocation, high-resolution structure and gene complement,

gene families.

SUMMARY

Wheat-rye 1RS.1BL translocation has a significant impact on wheat yield and hence food production globally. However, the genomic basis of its contributions to wheat improvement is undetermined. Here, we generated a high-quality assembly of 1RS.1BL translocation comprising 748,715,293 bp with 4,996 predicted protein-coding genes. We found the size of 1RS is larger than 1BS with the active centromere domains shifted to the 1RS side instead of the 1BL side in Aikang58 (AK58). The gene alignment showed excellent synteny with 1BS from wheat and genes from 1RS were expressed well in wheat especially for 1RS where expression was higher than that of 1BS for the grain-20DPA stage associated with greater grain weight and negative flour quality attributes. A formin-like-domain protein FH14 (*TraesAK58CH1B01G010700*) was important in regulating cell division. Two PPR genes were most likely the genes for the multi fertility restoration locus Rf^{multi}. Our data not only provide the high-resolution structure and gene complement for the 1RS.1BL translocation, but also defined targets for enhancing grain yield, biotic and abiotic stress, and fertility restoration in wheat.

INTRODUCTION

The 1RS.1BL translocation chromosome was one of the earliest of so-called alien chromatin additions into wheat and is generally considered to be associated with the disease resistance (Zeller, 1973) and a step-change in yield achieved with the release of the Veery lines by CIMMYT (Rajaram et al., 1983). A survey by R Schlegel showed that approximately 30% of wheat cultivars released after the year 2000 carry the 1RS.1BL translocation (http://www.rye-gene-map.de/ rye-introgression/) (Schlegel and Korzun, 1997). The *Lr26*, *Sr31*, *Yr9*, *Sr50* rust resistance genes and the powdery mildew, *Mlg* locus have been identified on 1RS.1BL chromosomes as well as genetic factors affecting root biomass (Mago et al., 2002 and 2015; Ehdaie et al., 2003; Waines and Ehdaie, 2007; Sharma et al., 2011). The grain yield associated with 1RS.1BL has been shown to be disrupted by 1RS-1BS recombinants in the terminal region of 1RS as characterized in field trials of wheat accessions carrying these

recombinants in the same genetic background (Lukaszewski et al., 2000; Howell et al., 2014

and 2019). The 1RS.1BL chromosome has also been factored into maintaining male sterility

in hybrid wheat programs (Lukaszewski et al., 2017).

The 1RS rye chromosome segment in wheat varieties has at least 3 sources, namely, as

1RS.1BL in *Triticum aestivum* cv Salmon (Japan), as 1RS.1BL in *T. aestivum* varieties

originating from Germany and as 1RS.1AL in *T. aestivum* cv Amigo from the USA (Schlegel

and Korzun, 1997). The 1RS.1BL translocation has been widely introduced into wheat

cultivars globally for conferring a broad-range of resistance to races of powdery mildew and

rusts, environmental adaptability and yield performance (Zeller, 1973; Rajaram et al., 1983;

Schlegel and Korzun, 1997; Bartoš and Bareš, 1971; Sukumaran et al., 2015).

At a structure level, 1RS.1BL was one of the early chromosomes to have specific DNA

sequences assigned to it through the analysis of chromosomes using inbred lines of rye to

facilitate locating molecular markers and agronomic traits to 1RS as well as the entire rye

genome (Lawrence and Appels, 1986; Miedaner et al., 2012; Bauer et al., 2016). *In situ*

hybridization technology with labelled sequenced probes has allowed the convenient

microscopic visualization and identification of translocations involving 1RS (Appels et al.,

1978; McIntyre et al., 1990; Liu et al., 2017).

The favorable agronomic features have been a significant driver for sequencing the 1RS.1BL

chromosome in the China wheat cultivar, AK58. In the available rye genome data from an

inbred rye line 27,784 gene models (and segments) were sourced for assigning gene models to

the 1RS segment and these generally fell within gene models aligned from the wheat

reference genome sequence (Bauer et al., 2016; IWGSC et al., 2018). This study provides the

linear genome level structure of AK58-1RS.1BL utilizing a combination of Illumina and

PacBio sequencing with de novo NR Magic for the initial assembly followed by HiC

scaffolding and alignment to high density molecular genetic maps to generate the final

assembly. The genome structure identified new gene models, several multi-gene families

likely to be involved in yield attributes associated with 1RS.1BL and resolved gene families

involved in disease resistance and other agronomic traits.

RESULTS

Assembly of the 1RS.1BL genome of wheat cv AK58

The details of the assembly are provided in Figure S1 and Methods, and include the de novo

NR Magic software to carry out a primary alignment into contigs, followed by the HiC

process for scaffolding the contigs. Finally, we generated a high-quality chromosome-scale

assembly of 1RS.1BL translocation with a total length of 748,715,293 bp and predicted 4,996

genes using four different annotation pipelines and alignment with the IWGSC RefSeq v1.0

annotation. The alignment of our AK58_chr1RS.1BL_v6 to the reference Chinese Spring (CS)

1B is shown in Figure 1a. The AK58_chr1RS.1BL_v6 assembly was examined in detail in the

terminal 22 Mb region because, in general, this region of wheat genome assemblies can be

problematical with respect to contig orientation. The assembly shown was the best alignment to

available genetic mapping information for 1RS.1BL, as shown in Figure 1b (Mago et al., 2002;

Howell et al., 2014; Sharma et al., 2009), using the markers, gamma secalin (8.82 Mb) and

BE444266 (24.41 Mb). The IB267 (0.815 Mb) and iag95 (3.92 Mb) markers were located

following discussions with J Dubcovsky (pers. comm., see Figure 1b).

Alignment and *in situ* cross-referencing of the AK58-1RS.1BL genome

The *in situ* probes are generally repetitive and although the number of repeats was clearly

collapsed during the assembly process, all the regions aligned by *in situ* hybridization using

double labelling could be assigned positions in the new assembly (McIntyre et al., 1990; Liu

et al., 2017; Zhang et al., 2004). Importantly the macro-level structure of 1RS.1BL could be

validated in this way (Figure 1c). The repetitive array of Sc119.2 sequences at 117.6 Mb (31

copies of the core 45 bp repeat unit) on 1RS were under-represented in the assembly, relative

to the array at 0.4 Mb (662, 45 bp repeat units) based on comparing the *in situ* hybridization

signals which indicated qualitatively similar signals (Figure 1c). The amplification of

repetitive gene families in 1RS (Sc119.2, Sc200) in positions that were not in a syntenic order

has occurred against a background of a conserved syntenic order of high confidence (HC)

gene models (Figure 1a). The repetitive gamma-secalin and omega-secalin gene families were located in syntenic positions relative to 1BS of CS. In Figure 1d, a portion of a diversity analysis is shown using a 660K SNP-chip for SNPs that could be clearly scored in 36 1RS containing lines (identified using the gamma-secalin based PCR probes) (Figure S2 and Table S1). A subset of 9 wheat lines are shown for comparison to confirm that at the macro-level the 1RS is a large haplotype block, as described by Cheng et al. (Cheng et al., 2019). At a micro-level at least 6 groups, or haplotypes, of 1RS.1BL could be identified using the AK58-1RS as a reference, and these are accounted for by considering the different rye genome sources used in the intense breeding efforts in China combining 1RS.1BL containing wheat lines in crosses with triticales, rye and alternative sources of 1RS.1BL (Figure 1d, Figure S2 and Table S1). The 1RS in AK58 groups with only 3 other 1RS lines.

In our assembly, the size of 1RS is 275 Mb, 28% larger than 1BS (215 Mb) and is consistent with the overall genome size of rye (7-8 Gb) being approximately one-third larger than the diploid genome of barley or wheat progenitors. A comparison of the TE-complement between 1RS, 1AS, and 1DS in AK58 and 1BS in CS indicated that 12 TE subfamilies (Figure S3 and Table S2) were dominant in 1RS with a total length in excess of 41.9 Mb (15.27% of the 1RS length), compared to only 4.9 Mb in 1AS, 3.5 Mb in 1DS and 4.8 Mb in 1BS of CS. The 12 rye dominant TE included nine LTRs. There were five TE families, *LTR-Gypsy-RLG_famc9.2, LTR-RLX_famc7, LTR-RLX famc21, Unknown-XXX_famc9,* and *Unknown-XXX_famc81,* in which the length ratios of 1RS/1AS, 1RS/1BS and 1RS/1DS are range from 4 to 203 (Table S2). Although most of the rye dominant TEs were distributed in 1RS, including gene-flanking regions, some were more prominent in the centromere region (LTR-Gypsy-RLG_famc36, Figure S3; see also centromere section below).

Centromere structure at the 1RS-1BL boundary

The availability of a rye centromere sequence (pAWRC, Francki, 2001) that could be distinguished from the wheat centromere repeats, CRWs, allowed a more detailed analysis of the rye-wheat hybrid centromere region. A 800 bp region from pAWRC (AF245032) that had no similarity to CRWs was used to define a 9.9 Mb region on the 1RS side of the 1RS.1BL

centromere while CRW/CCS1 (AB048244.1, a 249 bp repetitive unit), and Tail 1 (AB016967)

from the wheat centromere were used to define the 1BL side of the 1RS.1BL chromosome

(Figure 2a) (Francki, 2001; Keeble- Gagnère et al., 2018). In total the region covered by these

centromere markers was 10.87 Mb within a region of 272.02 Mb to 296.20 Mb (24.18 Mb).

The centromere marker sequences are evident in the matrix analysis (Figure 2a) as large arrays

of repetitive sequences. At the junction between 1RS and 1BL, to form the 1RS.1BL

chromosome, there exists a sharp change-over from the blocks of repetitive sequences carrying

pAWRC (Figure 2a-a1, blue dashed line box) to the CRW markers sequences (Figure 2a-a1,

red dashed line boxes). The matrix defines the junction between 281.93 Mb and 281.99 Mb

and is between a RLG_Taes_Abia_B_3Brph7-445 element (coordinates 281,918,330 to

281,927,173 bp) and a LTR, Gypsy; consensus sequence (coordinates 281,926,633 to

281,935,207 bp) using TREP database (http://botserv2.uzh.ch/kelldata/trep-db/) to identify the

LTR elements. The junction can be most easily modeled as resulting from a recombination

event involving an Abia sequence located in Abia-like segments within CRW/Cereba elements

in the wheat centromere and an Abia element in the original 1R chromosome.

The ca 150 Mb region that is relatively poor in annotated gene models (coordinates 165 to

313 Mb, Figure 1a and Figure 2a, dashed line boxes) houses predicted genes that code for

peptides less than 50 amino acids and has a good coverage of hits from RNA-seq data

originating from a range of tissues that could not be clearly assigned to gene models.

Immunoprecipitation of CENH3 binding genome sequences from CS and AK58 nuclei

provided a class of sequence to further define the core 31 Mb centromere region more clearly.

The CS-CENH3 sequences differentiated the wheat centromere segment on 1BL from 1RS

when aligned across the AK58_chr1RS.1BL_v6 assembly (Figure 2a-a2, coordinates 282 to

292 Mb) and also identified two sections of non-centromere (wheat) DNA (blue solid line

boxes in Figure 2a-a5) even though the well-known centromere transposable elements,

Cereba and Quinta, exist in these regions (Figure 2a-a1 and Figure 2a-a4). The AK58-CENH3

sequences mainly identified the rye centromere segment 272 to 282 Mb on 1RS (Figure

2a-a3). The dot matrix of CS vs AK58 core centromere sequences (Figure 2a-a5) indicated a

region where the two genomes are structurally rearranged relative to each other.

The *in situ* localization of the centromere Abia (rye) and CRW (wheat) sequences as well as

the CENH3 protein using fluorescent antibodies (Figure 2b) indicated that the AK58 CENH3

sequences mainly co-located with the Bilby sequences (rye, Abia TE family) and confirmed

that the CENH3-ChIP protocols selected specific sub-populations centromere sequences.

Expression of 1RS genes in a wheat background

Transferring alien chromosome or chromosomal fragments from wheat relatives to wheat is

an efficient approach for wheat improvement, relying on the expression of the alien

chromosome genes in a wheat background. There are 1,480 high confidence genes annotated

in 1RS and 1,560 on 1BS in CS (tissue expression summarized in Figure 3a) and entries of

particular interest relate to the deployment of the 1RS.1BL chromosome in wheat breeding

with a focus on those affecting grain quality and yield gains. Our study using the RNAseq

data sets indicated that the 1RS gene expression is generally successful in wheat backgrounds

and that 1RS genes do not interfere with wheat gene expression in total. A graphical

representation of tissue-based expression groups is summarized in Figure 3a and 3b for AK58

and CS. At this broad level it is evident that 1RS has a greater percentage of genes expressed

specifically in the grain tissue relative to CS (Figure 3b). In Figure 3c, the tissue specific gene

expression patterns among the AK58 and CS chromosome 1 pairs are compared in more detail

and it is evident that only 1BS in AK58 (= 1RS), has higher levels than 1BS of CS at the

grain-20 DPA stage, where we found 6.44× higher levels of total gene expression in 1RS of

AK58 relative to 1BS in CS. This grain-20 DPA stage is the active grain filling stage and the

higher levels of gene expression may thus relate to the large grain size (45 g/1000 grain) and

higher yield (7.5 t/h) of AK58 compared to the small grain size (35 g/1000 grain) and lower

yield (3.0 t/h) of CS.

The hierarchical clustering of gene expression in Figure S4a within the AK58-1RS gene space

of the developing grain provides a greater resolution the view of the gene families during the

progressive differentiation of the grain tissue occurs as the spike matures and was consistent

with IWGSC (IWGSC et al., 2018) and Ramírez-González (Ramírez-González et al., 2018).

In Figure S4b, the early, prominent, expression of the MIKC-type MADS-box transcription

factor family was striking. Genes that were highly expressed in grain tissue were of interest

because historically the presence of 1RS in bread wheat was considered detrimental for

qualities relating to the performance of flour from these wheat lines in the standard processing

methodologies (Lukaszewski et al., 2000; Gobaa et al., 2007; Li et al., 2016). Transcription

factor gene families prominently expressed in roots (as well as leaves) such as Cys3His zinc

finger protein (C3H), MYB protein family (MYB) and WRKY signature containing

transcription factor (WRKY) (Figure S4b) were of interest because this class of gene was well

represented in the region that was studied by Howell et al. in which disruption of the genome

structure through recombination with wheat 1BS reduced yield (more detailed analysis below)

(Howell et al., 2019).

The pentatricopeptide-repeat (PPR) gene family at the *rf*/*Rf* ^{multi} loci of 1RS and 1BS

PPR domain carrying proteins form the one of largest gene families of land plants and are

involved in the regulation of RNA metabolism including RNA editing, stability, processing,

and splicing to translation (Lurin et al., 2004; Cheng et al., 2016). Most of restorer of fertility

(*Rf*) genes cloned in model organisms encode P-class PPR proteins. AK58 is a 1RS.1BL

variety and the 1RS.1BL translocation replaces *Rf* ^{multi} locus from chromosome 1BS of wheat

(15.28 to 59.84 Mb in CS) in multiple CMS systems (*Aegilops kotschyi, Ae. uniaristata* and

Ae. mutica) to generate male sterile wheats (Tables S3 and 4) (Lukaszewski et al., 2017;

Tsunewaki, 2015). The syntenic region in AK58-1RS is at 0.58 to 52.35 Mb in our assembly

(Tables S5 and 6). The overall gene numbers in the regions that satisfy the definition of the

PPR gene models are 21 in CS and 8 in AK58(Tables S3-5). A cluster of 11 PPR-family

genes in CS-*Rf* ^{multi} are a tandem array (at location 56.49 to 58.34 Mb) and a similar cluster

exists in AK58-*rf* ^{multi} (at 48.60 to 49.42 Mb) but is not in an exactly matching syntenic

location (Figure 4a, Tables S6 and 7). One gene model, *TraesCS1B01G072900*, from the CS

cluster has a homologous sequence *TraesAK58CH1B01G045250* at 49.41 Mb in AK58 and

this gene in AK58 is also syntenic to *TraesCS1B01G074600* in CS-*Rf* ^{multi}. The

TraesCS1B01G074600 gene is, in turn, syntenic to another AK58 gene

(*TraesAK58CH1B01G043900*) in the AK58-*rf* ^{multi} region, so we consider it useful to define

this as one closely related group represented by *TraesCS1B01G072900*. These relationships

reflect a complex micro-level relationship between the AK58- $rf^{\,multi}$ and CS-$Rf^{\,multi}$ regions

which results in the different expression patterns that are relevant to the male

fertility/restoration of fertility (Figure 4b). The three P-class proteins from the CS-$Rf^{\,multi}$

region, namely, *TraesCS1B01G072300*, *TraesCS1B01G072900* and *TraesCS1B01G074600,*

and their 1RS homologs, have significant mitochondrial targeting scores (Figure 4c, Tables S6

and 8) and between these genes only *TraesCS1B01G072300* shows the most striking

difference in transcription in FM tissue when compared to its homolog in AK58

(*TraesAK58CH1B01G045100*) (Figure 4b).

Taken together, the CS-$Rf^{\,multi}$/AK58-$rf^{\,multi}$ comparison suggests the *TraesCS1B01G072300*

and *TraesCS1B01G072900* are the most likely candidate genes of the multi fertility restoring

locus $Rf^{\,multi}$ in the context of the 1RS.1BL translocation replacing the $Rf^{\,multi}$ locus from

chromosome 1BS in multiple *Aegilops* CMS systems that generate male sterile wheats and

male fertility being subsequently restored by the 1BS-$Rf^{\,multi}$ locus (Lukaszewski et al., 2017;

Tsunewaki, 2015).

The grain storage protein gene families

Gamma and omega secalin sequences were identified and aligned to gamma and omega

gliadins from the CS chromosome 1B. While the sequence similarity between gamma gliadins

and gamma secalins was over 70%, omega gliadin and omega secalin sequences differed

significantly both in the signal peptide as well as in their repetitive regions. Nineteen gamma

secalin coding sequences were identified from which 18 were located in a single cluster at

1RS between positions 8,722,719 bp and 8,904,808 bp (Figure 5a). Using an extensive

proteome database of rye grain peptides (Figure S1 and Methods) (Bose et al., 2019), half of

these genes were verified at peptide level as bone-fide gamma secalin protein coding genes.

Eight sequences (7 gamma secalins, 1 purinin) represented complete sequences, two

sequences were partial sequences and ten sequences contained frameshifts or internal stop

codons. Similarly, a cluster of 18 omega secalin genes were identified, between positions

18,457,234 bp and 18,690,191 bp with 13 verified at the proteome level (Figure 5b). Between

the gamma secalin and omega secalin loci a single purinin gene was identified in a conserved

position compared to chromosome 1B.

Within the identified gamma secalins there were two sequences (Gamma secalin 5 and
Gamma secalin 16) with Tryp_alpha_amyl domain (PF00234) while the majority of gamma
secalins possess a single Gliadin Pfam domain (PF13016). In bread wheat, all the identified
gamma gliadins and low molecular weight (LMW) glutenins have PF13016 (Gliadin)
domains, and Tryp_alpha_amyl domains were only found in alpha gliadins among the gliadin
and glutenin types. The Pfam domain composition of available secalin sequences from
UniProt DB indicates the Gliadin domain (PF13016) and Tryp_alpha_amyl domain (PF00234)
are both characteristic of rye secalins, while PF00234 domains were only identified in the 75
k gamma secalins. The Tryp_alpha_amyl domain containing proteins represent a sulphur-rich
domain structure and are primarily characterized to have functions related to defense
mechanisms against pathogens or lipid transport (non-specific lipid transfer proteins). As both
in 1BS and 1RS the major storage proteins are located within regions enriched in disease
resistance proteins this might indicate a potential alternative function of these proteins and
their possible involvement in stress responses.

There were eleven nsLTP genes found on the 1RS chromosome arm, ten of which represented
the PR60 nsLTP sub-type (UniRef100_B2C4K0) clustered between 56.6 Mb and 57.2 Mb and
specifically expressed in grain tissue (Figure 5c); a single nsLTP (at 42.24 Mb) was highly
expressed in roots. On the long arm, 7 nsLTPs, 5 LTPs and 4 LTP-like sequences were
identified from which 3 short LTP-like sequences were grain specific. All the prolamin genes
present on 1RS were specifically/preferentially expressed in the grain, while the nsLTPs were
also expressed in other tissues such as under-spike internodes and young spikes.

The disease resistance gene families
The *Pm8* gene, orthologue of the *Pm3*, provided a good model for the RGA gene in the region
although the expression of the gene itself was not confined to any single tissue category
(Hurni et al., 2013). The identification of *Pm8* suggested that 1RS of AK58 is from diploid
rye Petkus. The RGA genes specifically expressed in different tissue were assessed against a

total of 2,871 gene models with the NB-ARC domain in their structure were identified within

the IWGSC RefSeq v1.0. In the AK58 1RS.1BL chromosome, disease resistance gene models

were identified (Table 1, Figure S1 and Methods). The NB-ARC domains are the most

common feature within disease resistance genes and are key components of apoptosomes that

are involved in recognizing the presence of a pathogen or DNA damage in the cell and

responding to the problem by localized cell death to protect the organism per se

(Crespo-Herrera et al., 2017; Li et al., 2016). The range of gene diversity is consistent with

the range of intrinsic and extrinsic stimuli to which wheat is exposed. Among the 22

1RS-specific gene models we identified three broad categories, dominated by a large group

(15) which comprised members of RGA families located on wheat 1B as well as other

chromosomes and included genes that were identified as having disease resistance-like

protein and tyrosine kinase domains (Table S9). Four gene models were part of a very large

family encoding proteins with lrr-serine/threonine kinase domains. Three gene models were

of particular interest in that they were absent from 1BS and thus represent novel resistance

genes introduced by 1RS (details in Table S9). Two of these gene models

TraesAK58CH1B01G008800 (at 4,676,665 bp) and *TraesAK58CH1B01G010100* (at

6,010,958 bp) are closely linked to, and on the proximal side of, the marker BE405749.1

which defines a region housing *Lr26* (Mago et al., 2002), as well as having the protein fold

c6j5tc associated with the disease resistance gene rpp13-like protein 4, determined using

Phyre2 (Kelley et al., 2015).

Characterization of the 1RS region disrupted by recombination between 1RS and 1BS

The yield-related region (YR) comprised the genome region in the terminal 14 Mb on the

1RS.1BL chromosome based on published molecular marker evidence in Mago et al. and

Howell et al. (Mago et al., 2002; Howell et al., 2014). The region exists within the terminal 22

Mb and housed 259 genes, from a total of 465, that could be characterized by the clusters of

co-expression representing gene networks potentially disrupted by recombination events

between 1RS and 1BS. The analysis in Figure 6 shows the co-expression matrix of the

contig-356, -445, -624 and -517 1RS genes (Table S10). The gene expression was calculated

using the Morpheus analysis tool (https://software.broadinstitute.org/morpheus/) and a 0.5

FPKM cut-off. The co-expression similarity matrix was calculated using Pearson correlation

and the clustering was carried out using only the co-expression values > absolute value 0.7

FPKM as being significant (highlighted red = positively correlated, blue = negatively

correlated). The contigs annotated as "Tissues" were identified based on the stage and

tissue-specific clustering in Figure 6. The boxed regions identified gene clusters that formed

qualitatively major networks based on shared patterns of expression. The 1RS genes in

clusters 1-7 shown in Figure 6 are combinations of models that are tissue specific in

expression and ones that are more generally expressed and overall, the analysis indicates that

the genes in the YR region are widely networked to genes involved in a broad range of

biological activity with a particular focus on root specific genes (Table S10). The clusters also

include genes potentially interacting with nitrogen metabolism related genes as well as genes

involved in stress responses and adaptation and thus a disruption in the activity profiles is

predicted to have wide ranging effects on a complex phenotype such yield. In particular we

identified   TraesAK58CH1B01G010700   (formin-like   domain   protein)   and

TraesAK58CH1B01G007500 (Cathepsin-L, homolog) genes as having the highest number of

co-expression partners. The rice homolog for the *TraesAK58CH1B01G010700* gene is

Os05t0104000-00, a formin-like-domain protein FH14, and defines a class of protein which

interacts with microtubules and microfilaments to regulate cell division. This gene also shows

co-expression with a nitrate reductase at chromosome 6B. The *TraesAK58CH1B01G010700*

gene showed a broad co-expression-based network including interactions with auxin response

factors on chromosomes 1A, 1B, 1D, 2B, 5A, 7A, 7B, 7D.

DISCUSSION

Our study provides the genome assembly for chromosome 1RS.1BL, a chromosome which

was one of the early successes of genetic engineering at a chromosome level that has had a

significant impact on wheat yield and hence food production globally (Schlegel and Korzun,

1997). The value of our assembly is demonstrated through the identification of 1RS-specific

genes within an overall gene space that showed excellent synteny with 1BS from wheat. In

contrast the non-gene space is shown to have regions of highly amplified, relatively short

(100 – 500 bp), units of DNA sequences in regions which are not syntenic to equivalent

regions in 1BS and are suggested to be possible sites for recombination type events between

1R and 1B. The detailed analysis of the centromere identified the junction between 1RS and

1BL and this was found to comprise part of an Abia transposable element (TE) on the 1RS

side and a Cerebra TE on the 1BS side suggesting that a recombination event between the

arrays of Abia TEs that characterize the rye centromere and the arrays of Cereba TEs in the 1B

centromere, which can also carry Abia-like sequences (http://botserv2.uzh.ch/kelldata/trep-db/),

could have generated the 1RS.1BL chromosome (Francki, 2001). Our analysis of the CENH3

locations indicated a shift in the location of the centromere assembly defined by CENH3 to the

rye side of the translocation and provided a clear indication of mobility in the location of the

point of attachment of micro-fibrils for mitosis.

Specific families of gene models characterized in detail in this study included the genes

predicted to be involved in the male sterile/male fertility restoration interactions, resistance

gene analogs and the gamma and omega secalin storage proteins. The substitution of 1BS

with 1RS in 1RS.1BL wheat lines results in the replacement of three PPR proteins that have

high mitochondria target scores. In the context of the CS-Rf^{multi}/AK58-rf^{multi} comparison

carried out, it is suggested that *TraesCS1B01G072300* and *TraesCS1B01G072900* are

representative of the most likely candidate genes of the multi fertility restoring Rf^{multi} locus

because they show the most striking differences in transcription to its AK58 homolog and

could thus be most significant in restoring male fertility the 1BS-Rf^{multi} locus in the multiple

CMS systems-male sterile wheats where 1RS.1BL translocation replaces the Rf^{multi} locus

from chromosome 1BS (Lukaszewski et al., 2017; Tsunewaki, 2015). Our analysis of the

resistance gene analogs identified three genes that were absent from 1BS and thus represent

new resistance genes for varieties carrying the 1RS.1BL translocation. Two of these genes,

TraesAK58CH1B01G008800 (at 4,676,665 bp) and *TraesAK58CH1B01T0G0100* (at

6,010,958 bp), locate in the genome region housing genetically defined disease resistance

genes and are thus candidate genes for entities such as *Lr26*.

The substitution of 1BS in 1RS.1BL wheat lines also results in the replacement of a major

source of wheat gliadin proteins on 1BS with secalin protein coding genes on the 1RS.1BL

chromosome. The analysis of the gamma and omega secalin storage protein coding regions and expression identified pseudogenes as well as two sequences (Gamma secalin 5 and Gamma secalin 16) with Tryp_alpha_amyl domain (PF00234). The Tryp_alpha_amyl domain containing proteins, represent a sulphur-rich domain structure are more broadly associated with defense mechanisms against pathogens, lipid transport and storage function and suggests an involvement in stress responses. In the context of unwanted processing attributes ("sticky" dough) associated with flour from 1RS.1BL containing wheat cultivars genes related to the production of xylose and arabinoxylose and the xylanase domain protein (TraesAK58CH1B01G088700) specifically expressed at 4 DPA in the grain could be related to the moderation of arabinoxylans that may be associated with increased water absorption resulting in the "sticky" dough defect (Gobaa et al., 2007; Henry et al., 1989; Lee et al., 1995).

The mapping of published DNA probes to the 1RS assembly indicated that a 9 Mb region (in the terminal 14 Mb, "YR" region) was disrupted by the 1RS/1BS recombinants selected by Lukaszewski (Lukaszewski et al., 2000). The genes in this region are widely networked, as expected, based on co-expression analyses and have a striking representation of root-specific genes that are good candidates for relating the changes in root phenotype to the disruption of the 9Mb region defined by Howell et al (Howell et al., 2014 and 2019). Our study highlighted the complex negative and positive interactions among the genes in this YR region and identified TraesAK58CH1B01G010700 (formin-like domain protein) and TraesAK58CH1B01G007500 (Cathepsin-L, homolog) genes as having the highest number of co-expression partners. The rice homolog for the *TraesAK58CH1B01G010700* gene is *Os05t0104000-00*, a formin-like-domain protein FH14, showed a broad co-expression-based network across 8 chromosomes consistent with a gene model that encodes a protein regulating cell division.

In brief, we generated a high quality 1RS.1BL translocation chromosome sequence which provided a basis for defining gene underpinning the agronomic attributes of the 1RS.1BL translocation chromosome in wheat improvement. The structure, gene complement of 1RS.1BL and candidate genes identified in this study provides the resource-of-choice for

refining the contribution of this chromosome to wheat genetic improvement.

EXPERIMENTAL PROCEDURES

Gene annotation

Protein-coding identification and gene prediction were carried out using a combination of

homology-based prediction, *de novo* prediction, and transcriptome-based prediction methods

(Figure S1). Five *ab initio* gene prediction programs, Augustus (v.2.5.5), Genscan (v.1.0),

GlimmerHMM (v.3.0.1), Geneid, and SNAP, were used to predict coding regions in the

repeat-masked genome. Proteins from ten plant genomes (*T. aestivum*, *T. turgidum dicoccides,*

T. urartu, *Ae. tauschii*, *Hordeum vulgare*, *Brachypodium distachyon*, *Oryza. sativa*, *Zea mays*,

Sorghum bicolor and *Setaria italic*) were downloaded from EnsemblPlants

(http://plants.ensembl.org/index.html). *Panicum virgatum* genome was downloaded from

Phytozome (https://phytozome.jgi.doe.gov/pz/portal.html). Protein sequences from these

genomes were aligned to the AK58 assembly using TblastN with an *E-value* cutoff of 1e-5.

The BLAST hits were conjoined using Solar software. GeneWise was used to predict the

exact gene structure of the corresponding genomic regions for each BLAST hit. Homology

predictions were split into two sets, which included a high-confidence homology set

(HCH-set) with predictions from genomes with CS wheat and a low confidence homology set

(LCH-set).

A collection of wheat FLcDNAs (16,807 sequences) were directly mapped to the AK58

genome and assembled by Program to Assemble Spliced Alignments (PASA). Gene models

created by PASA were denoted as the PASA-FLC-set (PASA full length cDNA set), this gene

set was used to train the *ab initio* gene prediction programs. RNA-seq data were mapped to

the assembly using Tophat (v.2.0.8). Cufflinks (v.2.1.1) was then used to assemble the

transcripts into gene models (Cufflinks-set). In addition, a total of 2,016 Gb RNA-seq data

from different organs (root, leaf, internode, flower and developing seed) were assembled by

Trinity, creating several pseudo-ESTs. These pseudo-ESTs were also mapped to the AK58

assembly and gene models were predicted using the PASA. This gene set was denoted as

PASA-T-set (PASA Trinity set).

Gene model evidence from the HCH-set, LCH-set, PASA-FLC-set, Cufflinks-set, PASA-T-set and *ab initio* programs were combined by EvidenceModeler (EVM) into a non-redundant set of gene structures. Weights for each type of evidence were set as follows: HCH-set > PASA-FLC-set > PASA-T-set > Cufflinks-set > LCH-set > Augustus > GeneID = SNAP = GlimmerHMM = Genscan. Gene model output by EVM with low confidence scores was filtered using: (1) coding region lengths of 150 bp, and (2) supported only by *ab initio* methods and with FPKM<1.

In an approach similar to that described for *Gossypium raimondii* genome studies (Yan et al., 2016), we further filtered gene models based on Cscore (Cscore is a peptide BLASTP score ratio mutual best hits BLASTP score), peptide coverage (coverage is highest percentage of peptide aligned to the best of homologues) and overlap of its CDS with TEs. The Cscore and peptide coverage were calculated as described in *G. raimondii* (Yan et al., 2016). Only transcripts with a Cscore ≥ 0.5 and peptide coverage ≥ 0.5 were retained. For gene models with more than 20% of their CDS sharing an overlap with TEs, we required that its Cscore must be at least 0.8 and that its peptide coverage must be at least 80%. Finally, we also filtered out gene models of which more than 30% of the peptides in length could be annotated as Pfam or Interprot TE domains.

Transcript analysis

Two transcriptome datasets derived from AK58 genome sequencing project were used for this analysis. (1) 42 diverse samples (each 3 biological replicates, total 126 libraries) were collected for AK58, covering anther development to the tetrad stages, floret/spikelet meristem, three stages of stem development, three stages for flag leaf, five stages of grain development, and 7 day seedling for the leaf and root sample under normal or six abiotic stresses conditions. (2) Seven tissues shared between AK58 and CS (each 3 biological replicates, total 42 libraries), covering 21 days seedling for the leaf and root, floret meristems stage, and four grain development stages (4/10/15/20 days after flowering).

To align RNA-seq sequences to the genome assembly, the BLAT software within Apollo was

used in two modes (Lee et al., 2013), one allowing only perfect matches and a second mode

using the Hi-sat-2 default settings (https://ccb.jhu.edu/software/hisat2/index.shtml) equating

to approximately 80% similarity over at least 80% of the sequence. Only unique gene models

were used to define the 1RS.1BL synteny.

Gene co-expression matrix of the contig-356, -445, -624 and -517 1RS genes were developed

using   the   Morpheus   matrix   visualization   and   analysis   tool

(https://software.broadinstitute.org/morpheus/). Mean values of gene expression data obtained

from the different tissue samples with a 0.5 FPKM cut-off were used to calculate Pearson

correlation coefficients. The obtained similarity matrix was used for hierarchical clustering

with complete linkage. Rows and columns represent the individual gene models present in the

analyses 22 Mb region (Table S10). Tissue specific expression patterns and location within the

analyzed region are annotated by different colors were obtained from the analysis presented in

Figure 6. Co-expression similarity values > absolute value 0.7 FPKM as being significant are

highlighted in red (positively correlated) and blue (negatively correlated).

Chromosome preparation

The protocol for root tip mitotic metaphase chromosomes of AK58 was largely referred to the

previously report (Han et al., 2006). Briefly, the roots of AK58 grew to 1.5-2.0 cm in length

were excised and treated with nitrous oxide gas for 2 h under pressure under 1 MPa. The

treated roots were fixed in ice-cold 90% acetic acid for 10 min. Subsequently, the root tips

were dissected and digested by 2% cellulose Onozuka R-10 (Yakult Pharmaceutical, Japan)

and 1% pectolyase Y23 (Yakult Pharmaceutical) solution for 45 min at 37°C. After digestion,

the root sections were broken in a 90% acetic acid. The cell suspension was dropped and

air-dried on glass slides for chromosome observation.

Fluorescence *in situ* hybridization

The method of sequential FISH and Non-denaturing FISH (ND-FISH) with different labeled

probes for karyotype analysis of AK58 was mainly performed according to the previously

published protocol (Fu et al., 2015). The probe Sec1 for rye specific secalin was labeled with

Texas Red-5-dUTP (Invitrogen) or Alexa Fluor 488-5- dUTP (Invitrogen) using nick

translation for FISH (Clarke et al., 1996). The oligonucleotide probes for ND-FISH with

centromeric specific probe CCS1, 18S-45SrDNA probe pTa71, and probe pSc119.2 were

referred to Tang et al. (Tang et al., 2014). The repeats probes pSc200 and 5SrDNA were from

the previously published information (Fu et al., 2015; Lang et al., 2019). The synthetic oligo

probes were 5' end-labeled with 6-carboxyfluorescein (FAM) for the green signal and

6-carboxytetramethylrhodamine (Tamra) for the red signal. The slides after FISH and

ND-FISH were mounted with Vectashield mounting medium containing 1.5 μg/mL 4, 6 -

diamidino -2- phenylindole (DAPI, Vector Laboratories, Burlingame, CA, USA). The FISH

images were captured with an Olympus BX-53 microscope equipped with a DP-80 CCD

camera.

660K SNP Analysis of the 1RS Lines and Non-1RS Lines

Wheat 660K SNP Array designed by the Institute of Crop Sciences of the Chinese Academy

of Agricultural Sciences and synthesized by Affymetrix® was applied to genotype 36

1RS.1BL lines and 9 non 1RS.1BL lines. All the SNPs were merged to FASTA file for 45

samples. TreeBeST (1.9.2) (http://treesoft.sourceforge.net/treebest.shtml#intro) nj was used to

build neighbor-joining phylogenetic tree with parameters: "-b 1000" and MEGA7 was used to

visualize phylogenetic trees (Kumar et al., 2016).

ChIP-seq

We used chromatin immunoprecipitation and sequencing technique to find the centromeric

DNA by CENH3 antibody, which was a rabbit polyclonal antiserum and was raised against the

peptide 'CARTKHPAVRKTK' (Li et al., 2013). ChIP was conducted using young leaves of

AK58 as previously described (Nagaki et al., 2003). The enriched DNA samples were

sequenced using Illumina Hiseq X-10 to generate 150 bp paired-end sequences. Reads were

filtered with TrimGalore (http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/)

and aligned to the AK58 genome sequence using Bowtie 2 (Langmead and Salzberg, 2012). We

only retained reads that determined the unique position, which MAPQ >= 30, for further

analysis. The distribution of ChIP-seq reads were calculated using the unique read number per 1

kb window. ChIP-Seq data precipitated from CS in our previous study (SAMN11655702) were

analyzed as parallel.

Centromeric sequence analysis

In order to detect the sequence variation, we aligned the sequence of 1B from CS to AK58

genome using NUCmer program (parameter: -c 700) in MUMmer4 package (Marcais et al.,

2018). We used mummerplot program in the same software to draw 245–315 Mb of 1B

chromosome from AK58. Consensus of *CRW*, *Quinta* and *Bilby* were aligned to AK58

genome and calculated the percentage per 50 kb to reveal the sequence composition.

Immuno-co-localization Analysis of CENH3 with *CRW* and *Bilby*

Root tips of young seedling were collected for Immuno-staining as previously described

(Zhao et al., 2019). *CRW* and *Bilby* were labeled with biotin-16-Dutp and

digoxigenin-11-dUTP via nick translation respectively. The hybridized probes were detected

by fluorescein-conjugated goat anti-biotin and anti-digoxigenin-rhodamine Fab fragments

coupled with TAMRA respectively. CENH3 was identified with anti-CENH3 antibody

detected by Goat anti-Rabbit IgG (H+L) conjugated Alexa Fluor 647. Images were taken by a

confocal (ZEISS LSM880) and processed using Adobe Photoshop CS.

PPR gene analysis

The genome sequence data and gene annotation of CS were downloaded from EnsemblPlants

(http://plants.ensembl.org/index.html). PPR sequences were annotated by Pfam databases

(http://pfam.xfam.org/, v.32.0), and IPR database (https://www.ebi.ac.uk/interpro/, v.77.0),

respectively. The classification of PPR was referred to Cheng et al. (Cheng et al., 2016). The

PPR gene collinearity between 1RS of AK58 and 1BS of CS was analyzed by best reciprocal

hit BLAST (*E-value* cutoff of 1e-5) (Gabriel and Kristen, 2007). The PPR gene's subcellular

localization was predicted using PSI (http://bis.zju.edu.cn/psi/) and validated by experiment.

In briefly, full-length PPR cDNA was amplified and inserted in front of a GFP-coding

sequence on the pUC-35S-EGFP vector. Tobacco (*Nicotiana benthamiana*) protoplasts were

prepared and transfected according to the method described by Shan et al (2014). The

protoplasts were subsequently stained with 10μm MitoTracker™ Orange CMTMRos

(Invitrogen, M7510) for 10 min, and then examined on using a confocal scanning microscope

system (ZEISS LSM880).

The grain storage protein gene analysis

Prolamin super-family genes identified in the wheat reference genome were used to manually

annotate the prolamin genes in the AK58 genome sequence (IWGSC et al., 2018; Juhász et al.,

2018). Translated sequences were checked for the presence of signal peptides and the

conserved cysteine pattern and Pfam domains as previously described (Juhász et al., 2018).

Obtained sequences were aligned with gliadins and secalins retrieved from the Uniprot

database to confirm the protein sub-types. Expression of genes was analyzed using the grain

specific transcriptome data set obtained from 4, 10, 15 and 20 DPA grain libraries. Protein

level expression of the translated secalins and nsLTPs were analyzed using the published data

(Bose et al., 2019). LC-MS-MS data originally generated from tryptic digests of rye flour

protein extracts were re-analyzed and protein identification was undertaken using

ProteinPilotTM 5.0 software (SCIEX) with the Paragon and ProGroup algorithms with

searches conducted against the Poaceae subset of the Uniprot database appended with the

identified 1RS gene models and a contaminant database (Common Repository of Adventitious

Proteins) (Shilov et al., 2007). Obtained fully tryptic peptides were mapped to the secalin and

nsLTP sequences in CLC Genomics Workbench v12 (Qiagen, Aarhus, Denmark) using 100%

sequence identity to confirm the expression at individual protein levels.

RGA gene analysis

To predict RGAs in AK58 genome, a new plant RGAs database was constructed using protein

sequences from the RGAdb (https://bitbucket.org/yaanlpc/rgaugury/src/master/) and candidate

RGAs predicted in *Ae. tauschii* genome (Li et al., 2016; Luo et al., 2017). A total of 61372

disease resistance related sequences were obtained. Protein sequences of all annotated genes

of AK58 were aligned to the new RGAs database using BLASTP with *E-value* cutoff of 1e-5.

Potential RGAs were selected based on rule 80:80:80 (query sequence coverage more than

80%, target sequence coverage more than 80 and identity more than 80%). Eight

RGAs-related domains and motifs including NB-ARC, NBS, LRR, TM, STTK, LysM, CC

and TIR were searched and identified by RGAugury pipeline (Li et al., 2016). RGA

candidates were predicted in *T. aestivum* (CS) genome using the same method.

ACKNOWLEDGES

564 The authors are grateful to Yaoguang Liu and his laboratory team from the College of Life

565 Sciences of the South China Agricultural University for suggestions in analyzing PPR gene

566 family. This research was supported by The National Key Research and Development

567 Program of China (2016YFD0101802 and 2016YFD0101602), Talent Program and

568 Innovation Program of CAAS.

AUTHOR CONTRIBUTIONS

570 ZR, DC, JJ, RA, and XK initiated the project and designed the study. AJ, DL, PD, ZJ, LF, KW,

571 GKG, ZY, GL, DW, UB, MC, CK, GZ, XZ, XL, GC, YW, ZN, and LW performed the

572 research. AJ, DL, PD, ZJ, LF, KW, GKG, CK, and GZ generated and analyzed the data. RA,

573 XK, JJ, AJ, DL, PD, ZJ and KW wrote the paper.

CONFLICT OF INTEREST

575 The authors have no conflicts of interest to declare.

DATA AVAILABILITY

The whole genome sequence data reported in this paper have been deposited in the Genome

Warehouse in National Genomics Data Center, Beijing Institute of Genomics (China National

Center for Bioinformation), Chinese Academy of Sciences, under accession number

GWHANRF00000000 that is publicly accessible at https://bigd.big.ac.cn/gwh. And the

Assembly and availability of 1RS.1BL genome sequence can be found in URGI,

https://urgi.versailles.inra.fr/download/iwgsc/AK58_1RS.1BL/.

SUPPORTING INFORMATION

584 **Figure S1.** Protein-coding gene prediction process overview.

REFERENCES

Appels, R., Driscoll, C. and Peacock, W. J. (1978). Heterochromatin and highly repeated

DNA sequences in rye (*Secale cereale*). *Chromosoma*, **70**, 67-89.

Bartoš, P. and Bareš, I. (1971). Leaf and stem rust resistance of hexaploid wheat cultivars

salzmünder bartweizen and weique. *Euphytica*, **20**, 435-440.

Bauer, E., Schmutzer, T., Barilar, I., Mascher, M. and Scholz, U. (2016). Towards a

whole-genome sequence for rye (*Secale cereale* L.). *Plant J.* **89**, 853.

Bose, U., Broadbent, J.A., Byrne, K., Hasan, S., Howitt, C.A., and Colgrave, M.L. (2019).

Optimisation of protein extraction for in-depth profiling of the cereal grain proteome. J.

Proteomics, **197**, 23-33.

Cheng, H., Liu, J., Wen, J., Nie, X. and Jiang, Y. (2019). Frequent intra- and inter-species

introgression shapes the landscape of genetic variation in bread wheat. *Genome Biol*. **20**,

136.

Cheng, S., Gutmann, B., Zhong, X., Ye, Y., Fisher, M.F., Bai, F., Castleden, I., Song, Y.,

Song, B., Huang, J., et al. (2016). Redefining the structural motifs that determine RNA

binding and RNA editing by pentatricopeptide repeat proteins in land plants. *Plant J*. **85**,

532-547.

Clarke, B. C., Mukai, Y. and Appels, R. (1996). The *Sec-1* locus on the short arm of

chromosome 1R of rye (*Secale cereale*). *Chromosoma*, **105**, 269-275.

Crespo-Herrera, L. A., Garkava-Gustavsson, L. and Ahman, I. (2017). A systematic

review of rye (*Secale cereale* L.) as a source of resistance to pathogens and pests in wheat

(*Triticum aestivum* L.). *Hereditas*, **154**, 14.

Ehdaie, B., Whitkus, R. W. and Waines, J. G. (2003). Root biomass, water-use efficiency,

and performance of wheat–rye translocations of chromosomes 1 and 2 in spring bread wheat

'Pavon'. *Crop Sci*. **43**, 710.

Francki, M. G. (2001). Identification of Bilby, a diverged centromeric Ty1-copia

retrotransposon family from cereal rye (*Secale cereale* L.). *Genome*, **44**, 266-274.

Fu, S., Chen, L., Wang, Y., Li, M., Yang, Z., Qiu, L., Yan, B., Ren, Z., and Tang, Z. (2015).

Oligonucleotide probes for ND-FISH analysis to identify rye and wheat chromosomes. *Sci*.

Rep. **5**, 10552.

Gabriel, M. H. and Kristen, L. (2007). Choosing BLAST options for better detection of

orthologs as reciprocal best hits. *Bioinformatics,* **24**, 319-324.

Gobaa, S., Bancel, E., Kleijer, G., Stamp, P. and Branlard, G. (2007). Effect of the

1BL.1RS translocation on the wheat endosperm, as revealed by proteomic analysis.

Proteomics, **7**, 4349-4357.

Han, F., Lamb, J. C. and Birchler, J. A. (2006). High frequency of centromere inactivation

resulting in stable dicentric chromosomes of maize. *Proc. Natl. Acad. Sci. USA*, **103**,

3238-3243.

Henry, R. J., Martin, D. J. and Stewart, B. G. (1989). Cell-wall polysaccharides of
rye-derived wheats, investigations of the biochemical causes of dough stickiness. *Food*
Chem. **34**, 309-316.

Howell, T., Hale, I., Jankuloski, L., Bonafede, M., Gilbert, M., and Dubcovsky, J. (2014).
Mapping a region within the 1RS.1BL translocation in common wheat affecting grain yield
and canopy water status. *Theor. Appl. Genet*. **127**, 2695-2709.

Howell, T., Moriconi, J.I., Zhao, X., Hegarty, J., Fahima, T., Santa-Maria, G.E., and
Dubcovsky, J. (2019). A wheat/rye polymorphism affects seminal root length and yield
across different irrigation regimes. *J. Exp. Bot*. **70**, 4027-4037.

Hurni, S., Brunner, S., Buchmann, G., Herren, G., Jordan, T., Krukowski, P., Wicker, T.,
Yahiaoui, N., Mago, R., Keller, B. (2013). Rye *Pm8* and wheat *Pm3* are orthologous genes
and show evolutionary conservation of resistance function against powdery mildew. *Plant J*.
76, 957-969.

International Wheat Genome Sequencing Consortium(IWGSC), et al. (2018). Shifting
the limits in wheat research and breeding using a fully annotated reference genome. *Science*,
361, eaar7191.

Juhász, A., Belova, T., Florides, C.G., Maulis, C., Fischer, I., Gell, G., Birinyi, Z., Ong, J.,
Keeble-Gagnère, G., Maharajan, A., et al. (2018). Genome mapping of seed-borne
allergens and immunoresponsive proteins in wheat. *Sci. Adv*. **4**, eaar8602.

Keeble-Gagnère, G., Rigault, P., Tibbits, J., Pasam, R., Hayden, M., Forrest, K., Frenkel,
Z., Korol, A., Huang, B.E., Cavanagh, C., et al. (2018). Optical and physical mapping
with local finishing enables megabase-scale resolution of agronomically important regions
in the wheat genome. *Genome Biol*. **19**, 112.

Keeble-Gagnère, G., Isdale, D., Suchecki, R.-ł., Kruger, A., Lomas, K., Carroll, D., Li, S.,
Whan, A., Hayden, M., and Tibbits, J. (2019). Integrating past, present and future wheat
research with Pretzel. bioRxiv, 517953.

Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. and Sternberg M. J. (2015). The
Phyre2 web portal for protein modeling, prediction and analysis. *Nat. Protoc*. **10,** 845-858.

Kucherov, G., Noé, L., Noé, L. and Roytberg, M. (2006). A unifying framework for seed
sensitivity and its application to subset seeds. *J. Bioinform. Comput. Biol*. **4**, 553-569.

Kumar, S., Stecher, G. and Tamura, K. (2016). MEGA7, molecular evolutionary genetics
analysis version 7.0 for bigger datasets. *Mol. Biol. Evol*. **33**, 1870-1874.

Langmead, B. and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods, **9**, 357-359.

Lang, T., Li, G., Wang, H., Yu, Z., Chen, Q., Yang, E., Fu, S., Tang, Z., and Yang, Z.
(2019). Physical location of tandem repeats in the wheat genome and application for
chromosome identification. *Planta*, **249**, 663-675.

Lawrence, G. J. and Appels, R. (1986). Mapping the nucleolus organizer region, seed
protein loci and isozyme loci on chromosome 1R in rye. *Theor. Appl. Genet*. **71**, 742-749.

Lee, J. H., Graybosch, R. A. and Peterson, C. J. (1995). Quality and biochemical effects of
a 1BL/1RS wheat-rye translocation in wheat. *Theor. Appl. Genet*. **90**, 105-112.

Lee, E., Helt, G. A., Reese, J. T., Munoz-Torres, M. C. and Lewis, S. E. (2013). Web
Apollo, A web-based genomic annotation editing platform. *Genome Biol*. **14**, R93.

Li, B., Choulet, F., Heng, Y., Hao, W., Paux, E., Liu, Z., Yue, W., Jin, W., Feuillet, C., and
Zhang, X. (2013). Wheat centromeric retrotransposons, the new ones take a major role in
centromeric structure. *Plant J*. **73**, 952-965.

Li, P., Quan, X., Jia, G., Xiao, J., Cloutier, S., and You, F.M. (2016). RGAugury, a pipeline
for genome-wide prediction of resistance gene analogs (RGAs) in plants. *BMC Genomics*,
17, 852.

Li, Z., Ren, T., Yan, B., Tan, F., Yang, M., and Ren, Z. (2016). A mutant with expression
deletion of gene *Sec-1* in a 1RS.1BL line and its effect on production quality of wheat. *PLoS*
ONE, **11**, e0146943.

Liu, C., Song, Q., Zhang, H., Yang, Z. and Hu, Y. (2017). Molecular cytogenetic
characterization and phenotypic evaluation of new wheat–rye lines derived from hexaploid
triticale 'Certa'×common wheat hybrids. *Plant Breeding*, **136**, 809-819.

Lukaszewski, A. J. (2000). Manipulation of the 1RS.1BL translocation in wheat by induced
homoeologous recombination. *Crop Sci*. **40**, 216-225.

Lukaszewski, A. J. (2017). Chromosomes 1BS and 1RS for control of male fertility in wheats and triticales with cytoplasms of *Aegilops kotschyi*, *Ae. mutica* and *Ae. uniaristata*. *Theor. Appl. Genet.* **130**, 2521-2526.

Lurin, C., Andrés, C., Aubourg, S., Bellaoui, M., Bitton, F., Bruyère, C., Caboche, M., Debast, C., Gualberto, J., Hoffmann, B., et al. (2004). Genome-wide analysis of Arabidopsis pentatricopeptide repeat proteins reveals their essential role in organelle biogenesis. *Plant Cell*, **16**, 2089-2103.

Luo, M.C., Gu, Y.Q., Puiu, D., Wang, H., Twardziok, S.O., Deal, K.R., Huo, N., Zhu, T., Wang, L., Wang, Y., et al. (2017). Genome sequence of the progenitor of the wheat D genome *Aegilops tauschii*. *Nature*, **551**, 498-502.

Mago, R., Spielmeyer, W., Lawrence, J., Lagudah, S., Ellis, G., and Pryor, A. (2002). Identification and mapping of molecular markers linked to rust resistance genes located on chromosome 1RS of rye using wheat-rye translocation lines. *Theor. Appl. Genet.* **104**, 1317-1324.

Mago, R., Miah, H., Lawrence, G.J., Wellings, C.R., Spielmeyer, W., Bariana, H.S., McIntosh, R.A., Pryor, A.J., and Ellis, J.G. (2005). High-resolution mapping and mutation analysis separate the rust resistance genes *Sr31*, *Lr26* and *Yr9* on the short arm of rye chromosome 1. *Theor. Appl. Genet.* **112**, 41-50.

Mago, R., Zhang, P., Vautrin, S., Šimková, H. and Dodds, P. N. (2015). The wheat *Sr50* gene reveals rich diversity at a cereal disease resistance locus. *Nat. Plants*, **1**, 15186.

Marçais, G., Delcher, A.L., Phillippy, A.M., Coston, R., Salzberg, S.L., and Zimin, A. (2018). MUMmer4, a fast and versatile genome alignment system. *PLoS Comput. Biol.* **14**, e1005944.

McIntyre, C. L., Pereira, S., Moran, L. B. and Appels, R. (1990). New *Secale cereale* (rye) DNA derivatives for the detection of rye chromosome segments in wheat. *Genome*, **33**, 635-640.

Miedaner, T., Hübner, M., Korzun, V., Schmiedchen, B. and Reif, J. C. (2012). Genetic architecture of complex agronomic traits examined in two testcross populations of rye (*Secale cereal* L.). *BMC Genomics*, **13**, 706.

Nagaki, K., Talbert, P.B., Zhong, C.X., Dawe, R.K., Henikoff, S., and Jiang, J. (2003).

Chromatin immunoprecipitation reveals that the 180-bp satellite repeat is the key functional

DNA element of *Arabidopsis thaliana* centromeres. *Genetics*, **163**, 1221-1225.

Rajaram S., Mann C. E., Ortiz-Ferrara G. and Mujeeb-Kazi A. (1983). Adaptation,

stability and high yield potential of certain 1B/1R CIMMYT wheats. *In, S. Sakamoto (Ed).*

Proc. 6th Int. Wheat Genet. Symp., Kyoto, Japan, 613-621.

Ramírez-González, R.H., Borrill, P., Lang, D., Harrington, S.A., Brinton, J., Venturini,

L., Davey, M., Jacobs, J., van Ex, F., Pasha, A., et al. (2018). The transcriptional

landscape of polyploid wheat. *Science*, **361**, eaar6089.

Schlegel, R. and Korzun, V. (1997). About the origin of 1RS.1BL wheat-rye chromosome

translocations from Germany. *Plant Breeding*, **116**, 537-540.

Shan, Q., Wang, Y., Li, J., and Gao, C. (2014). Genome editing in rice and wheat using the

CRISPR/Cas system. *Nat. Protoc*. **9**, 2395-2410.

Sharma, S., Bhat, P.R., Ehdaie, B., Close, T.J., Lukaszewski, A.J., and Waines, J.G.

(2009). Integrated genetic map and genetic analysis of a region associated with root traits on

the short arm of rye chromosome 1 in bread wheat. *Theor. Appl. Genet*. **119**, 783-793.

Sharma, S., Xu, S., Ehdaie, B., Hoops, A., Close, T.J., Lukaszewski, A.J., and Waines,

J.G. (2011). Dissection of QTL effects for root traits using a chromosome arm-specific

mapping population in bread wheat. *Theor. Appl. Genet*. **122**, 759-769.

Shilov, I.V., Seymour, S.L., Patel, A.A., Loboda, A., Tang, W.H., Keating, S.P., Hunter,

C.L., Nuwaysir, L.M., and Schaeffer, D.A. (2007). The Paragon Algorithm, a next

generation search engine that uses sequence temperature values and feature probabilities to

identify peptides from tandem mass spectra. Mol. Cell. *Proteomics*, **6**, 1638-1655.

Sukumaran, S., Dreisigacker, S., Lopes, M., Chavez, P. and Reynolds, M. P. (2015).

Genome-wide association study for grain yield and related traits in an elite spring wheat

population grown in temperate irrigated environments. *Theor. Appl. Genet*. **128**, 53-363.

Tang, Z., Yang, Z. and Fu, S. (2014). Oligonucleotides replacing the roles of repetitive

sequences pAs1, pSc119.2, pTa-535, pTa71, CCS1, and pAWRC.1 for FISH analysis. *J.*

Appl. Genet. **55**, 313-318.

Tsunewaki, K. (2015). Fine mapping of the first multi-fertility-restoring gene, Rf^{multi}, of

wheat for three *Aegilops* plasmons, using 1BS-1RS recombinant lines. *Theor. Appl. Genet.*

128, 723-732.

Waines, J. G. and Ehdaie, B. (2007). Domestication and crop physiology, roots of

green-revolution wheat. *Ann. Bot*. **100**, 991-998.

Yan, R., Liang, C., Meng, Z., Malik, W., Zhu, T., Zong, X., Guo, S., and Zhang, R. (2016).

Progress in genome sequencing will accelerate molecular breeding in cotton (*Gossypium*

spp.). *3 Biotech*, **6**, 217.

Zhang, P., Li, W., Fellers, J., Friebe, B. and Gill, B. S. (2004). BAC-FISH in wheat

identifies chromosome landmarks consisting of different types of transposable elements.

Chromosoma, **112**, 288-299.

Zhao, J., Hao, W., Tang, C., Yao, H., Li, B., Zheng, Q., Li, Z., and Zhang, X. (2019).

Plasticity in Triticeae centromere DNA sequences, a wheat × tall wheatgrass (decaploid)

model. *Plant J*. **100**, 314-327.

Zeller, F. J. (1973). 1B/1R wheat-rye chromosome substitutions and translocations. *Proc. 4th*

Int. *Wheat Genet. Symp., Missouri Agric. Exper. Station, Columbia, MO*, 209-221.

Table 1. RGAs prediction in AK58 1RS compared to CS 1AS, 1BS and 1DS, respectively.

Species	Chr	Have NB-ARC or TIR domain C: CC; N: NBS; L: LRR; T: TIR;				Has a TM domain STTK; TM; LRR; LysM; CC			Total
		CN	CNL	NBS	NL	RLK	RLP	TM-CC	
AK58	Chr1RS	8	39	4	13	74	4	7	149
	Chr1AS	6	20	1	8	39	2	1	77
CS	Chr1BS	10	30	14	18	41	4	1	118
	Chr1DS	5	28	4	10	37	5	2	91

 FIGURE LEGENDS

Figure 1. Structure of the AK58 1RS.1BL chromosome.

(a) The high confidence (HC) gene models from the IWGSC-Refseq v1.0/-Refseq v2.0

assembly for chromosome 1B of CS wheat have been aligned with their syntenic partner gene

models in AK58 1RS.1BL using open source Pretzel (IWGSC et al., 2018; Keeble-Gagnère et

al., 2019).

(b) Summary of available genetic and cytogenetic details for the terminal 22 Mb region of

1RS showing 1RS (red)-1BS (yellow) recombinants that disrupt the yield attributes associated

with the 1RS.1BL chromosome (Howell et al., 2014 and 2019).

(c) *In situ* hybridization of repetitive sequence probes typically used to identify rye

chromosomes to allow a broad level validation of the 1RS assembly.

(d) Distribution of SNPs along representative sections of the 1RS.1BL chromosomes based on

a 660K SNP-chip analysis (Figure S2 and Table S1). Nine wheat cultivars (left most clusters)

and 36 1RS containing cultivars (remaining columns in the Figure) where the dark horizontal

lines indicate a SNP in the respective position that is different in AK58 reference genome

(blue box). Absence of a dark line indicates the alternate allele is the same as that in AK58.

Figure 2. The 1RS.1BL centromere.

(a) Dot-plot analysis of genome sequence at 245 Mb-315 Mb and BLAST-based distribution

of ChIP-Seq data based on the CENH3-precipitation in AK58 and CS. (a1) Dot matrix of the

AK58 centromere region of 70 Mb using YASS with the dashed-line boxes indicating the very

large blocks of repetitive sequences (Kucherov et al., 2006). (a2) Blast of CENH3-antibody

ChIP-Seq reads from CS on the 1RS.1BL centromere region showed two sub-domains which

basically consist of the *CRW* and *Quinta* TEs. (a3) Blast of CENH3-antibody ChIP-Seq reads

from AK58, the blast domain appears on the 1RS side (red) instead of 1BL relative to the

wheat CRW (green) and Quinta (blue). (a4) Distribution of retrotransposons in the 1RS.1BL

centromere region of AK58 - the boxes in dashed lines and solid lines are discussed in the text.

(a5) Dot matrix of the 70 Mb AK58 centromere region vs the 20 Mb core centromere region

from CS. The solid blue line boxes define regions addressed in the text.

(b) Late prophase nuclei (same in each frame) show the *in situ* co-localization of the CENH3

antibody (white), the rye Bilby sequence (red) and the Cereba (CRW, green) sequences. The

Bilby sequence (red) detects 1RS centromeres from the other 40 wheat centromeres (green).

The green dots conjugating with the red dot is the 1BL centromere. The CENH3 signals were

mainly co-located with the Bilby signals and this directly supports the centromere shift in

observed in Figure 2 (a3) in the BLAST analysis of AK58-CENH3 ChIP-Seq reads.

Figure 3. Comparative analysis of gene expression in group 1 chromosomes between AK58

and CS.

(a) Expression pattern of genes in 1RS (blue) and 1BS (yellow) shown as % of total gene

number expressed in the spike, root, grain and leaf tissues. Tissue-specific genes are

highlighted in darker shades both for 1RS (blue) and 1BS (yellow).

(b) Comparison of tissue-specificity in AK58 and CS with the numbers showing information

for each line and tissue as AK58/CS.

(c) Comparison of tissue specific gene expression across the short and long arms of the group

1 chromosomes. The 1BS/1RS panel highlights a major difference between AK58 and CS in

the grain-DPA20 stage where a 6.44-fold higher expression of transcripts in AK58 vs CS at

20DPA was identified.

Figure 4. PPR gene comparisons between 1RS of AK58 and 1BS of CS.

(a) Gene collinearity analysis.

(b) PPR gene expression comparison for the genes indicated in either orange (CS origin) or

AK58 (blue origin) for the tissues, seedling leaf, seedling root, FM (flowering meristem) and

grain tissue (DPA = days post anthesis).

(c) Subcellular localization of *TraesCS1B01G072300* and *TraesCS1B01G072900*. Protoplasts

transformed   with   35S: GFP   (pUC-35S-GFP),   35S:TraesCS1B01G072300-GFP

(*TraesCS1B01G072300*)   and   35S:TraesCS1B01G072900-GFP   (*TraesCS1B01G072900*)

constructs were analyzed using fluorescence microscopy. DIC indicated bright field. The dye

Mito-Tracker Orange was used as a mitochondrial marker. (Scale bars: 20 μm.).

Figure 5. The grain storage protein gene analysis.

(a) Gamma secalin locus on chromosome arm 1RS.

(b) Omega secalin locus on chromosome arm 1RS.

(c) nsLTP locus on chromosome arm 1RS.The tracks underneath each of genome map

locations provide the level of expression at the respective stages of grain development and

evidence for the representation of the gene models as grain proteins in extensive rye proteome

studies.

Figure 6. Co-expression analysis of homologous genes expressed in the 22 Mb region of

AK58 1RS. Co-expression similarity values are represented in AK58 1RS calculated using

Pearson correlation followed by hierarchical clustering using the pre-computed similarity

matrix values. Co-expression values below -0.7 are labelled in blue and represent strong

negative co-expression. Co-expression above 0.7 represents strong expression similarity and

is highlighted in red. Tissue specificity and location of the genes within the analyzed contigs

in the 22 Mb region are highlighted with different colors. Matrix was generated using

https://software.broadinstitute.org/morpheus/.

Figure 1

Figure 2

 Figure 3

Figure 4

 Figure 5

 Figure 6