Title

Human skeletal muscle organoids model fetal myogenesis and sustain uncommitted PAX7 myogenic progenitors

Authors

Lampros Mavrommatis[1,2,15], Hyun-Woo Jeong[3], Urs Kindler[1], Gemma Gomez-Giro[4], Marie-Cécile Kienitz[5], Martin Stehling[6], Olympia E. Psathaki[2,7], Dagmar Zeuschner[8], M. Gabriele Bixel[9], Dong Han[2], Gabriela Morosan-Puopolo[1], Daniela Gerovska[10], Ji Hun Yang[11], Jeong Beom Kim[12], Marcos J. Araúzo-Bravo[10], Jens C. Schwamborn[4], Stephan A. Hahn[13], Ralf H. Adams[9,14], Hans R. Schöler[2], Matthias Vorgerd[15], Beate Brand-Saberi[1], Holm Zaehres[1,2]

Ruhr-University Bochum, Medical Faculty, Institute of Anatomy, Department of Anatomy and Molecular Embryology, 44801 Bochum, Germany

Max Planck Institute for Molecular Biomedicine, Department of Cell and Developmental Biology, 48149 Münster, Germany

Max Planck Institute for Molecular Biomedicine, Sequencing Core Facility, 48149 Münster, Germany

Luxembourg Centre for Systems Biomedicine, LCSB, Developmental and Cellular Biology, University of Luxembourg, 4367 Belvaux, Luxembourg

Ruhr-University Bochum, Medical Faculty, Department of Cellular Physiology, 44801 Bochum, Germany

Max Planck Institute for Molecular Biomedicine, Flow Cytometry Unit, 48149 Münster, Germany

Center for Cellular Nanoanalytics Osnabrück, CellNanOs, University of Osnabrück, 49076 Osnabrück, Germany

Max Planck Institute for Molecular Biomedicine, Electron Microscopy Unit, 48149 Münster, Germany

Max Planck Institute for Molecular Biomedicine, Department of Tissue Morphogenesis, 48149 Münster, Germany

Computational Biology and Systems Biomedicine, Biodonostia Health Research Institute, San Sebastián 20014, Spain 9 IKERBASQUE, Basque Foundation for Science, Bilbao 48013, Spain

School of Mechanical Engineering, Korea University, Seoul 02841, Republic of Korea; Next & Bio Inc., Seoul National University, Seoul 08826, Republic of Korea

School of Life Sciences, Ulsan National Institute of Science and Technology (UNIST), Ulsan 44919, Republic of Korea

Ruhr University Bochum, Medical Faculty, Department of Molecular GI Oncology, 44780, Bochum, Germany

Westphalian Wilhelms-University Münster, Medical Faculty, 48149 Münster, Germany

Department of Neurology with Heimer Institute for Muscle Research, University Hospital Bergmannsheil, 44789 Bochum, Germany

[*] Correspondence: Lampros Mavrommatis (lampros.mavrommatis@rub.de); Hyun-Woo Jeong (hyun-woo.jeong@mpi- muenster.mpg.de) for bioinformatics; Holm Zaehres (holm.zaehres@rub.de)

Abstract

In vitro culture systems that structurally model human myogenesis and promote PAX7$^+$ myogenic progenitor maturation have not been established. Here we report that human skeletal muscle organoids can be differentiated from induced pluripotent stem cell lines to contain paraxial mesoderm and neuromesodermal progenitors and develop into organized structures reassembling neural plate border and dermomyotome. Culture conditions instigate neural lineage arrest and promote fetal hypaxial myogenesis towards limb axial anatomical identity, with generation of sustainable uncommitted PAX7 myogenic progenitors and fibroadipogenic (PDGFRa+) progenitor populations equivalent to those from the second trimester of human gestation. Single cell comparison to human fetal and adult myogenic progenitors reveals distinct molecular signatures for non-dividing myogenic progenitors in activated (CD44^{High}/CD98$^+$/MYOD1$^+$) and dormant (PAX7^{High}/FBN1^{High}/SPRY1^{High}) states. Our approach provides a robust 3D in vitro developmental system for investigating muscle tissue morphogenesis and homeostasis.

Introduction

Novel skeletal muscle model systems are required to further elucidate the process of human myogenesis as well as investigate muscular disorders and potential gene, cell or drug therapies. Two-dimensional (2D) culture conditions guide pluripotent stem cell (PSC) differentiation towards skeletal muscle lineage using sequential growth factor applications and/or conditional PAX7 expression (Chal et al., 2015, Xi et al., 2017, Shelton et al., 2014, Borchin et al., 2013, Darabi et al., 2012). Further, surface marker expression can be utilized to isolate myogenic progenitors with in vivo repopulation potential (Magli et al., 2017, Hicks et al., 2018, Al Tanoury et al., 2020, Sun et al., 2022). While the few described three-dimensional (3D) differentiation approaches have provided cohorts of terminally differentiated myofibers, focus on potential interactions with the vasculature and nervous system has neglected assessment of the developmental identity or sustainability of myogenic progenitors (Faustino Martins et al., 2020, Maffioletti et al., 2018, Rao et al., 2018). Single cell technologies increasingly provide databases for deciphering myogenic trajectories and expression profiles of myogenic stem and progenitor cells (Barruet et al., 2020, Rubenstein et al., 2020, Xi et al., 2020), enabling full evaluation of the ability of PSC differentiation protocols to mimic human development. Translation to model muscular dystrophies and investigate potential interventions in vitro necessitates methods that provide expandable populations of muscle progenitors while promoting self-renewal and preserving a quiescent, non-dividing, state (Quarta et al., 2016, Montarras et al.,2005).

Most vertebrate skeletal muscle progenitors develop from the paraxial mesoderm via transient embryonic developmental structures (somites and dermomyotome) into the skeletal muscle system that spans the whole body. Here, we evaluate human skeletal muscle organoids as a novel system to structurally investigate myogenic differentiation from human induced pluripotent stem cells (iPSCs) in a 3D environment, mimicking pathways described for chicken and mouse (Buckingham and Rigby, 2014). We develop a comprehensive supplementation/reduction protocol to first drive differentiation towards paraxial mesoderm through application of the GSK3 inhibitor CHIR99021, BMP inhibitor LDN193189 and bFGF. Subsequent stimulation with WNT1A, SHH, FGF and HGF is designed to promote derivation of organized structures reassembling neural plate border and dermomyotome. We then aim to arrest neural lineage via FGF removal , while stimulating with HGF to selectively promote propagation of undifferentiated myogenic progenitors and consequent generation of fetal myofibers. Our goal is to provide PAX7$^+$ myogenic progenitors in a non-dividing quiescent state sustainable over weeks after differentiation induction. Single cell analysis will position cells along the quiescent-activation myogenic trajectory discriminating dormant (PAX7+, FBN1+, SPRY11+, CHODL1+) and activated (CD44+, CD98+, MYOD1+, VEGFA+) states. We thus seek to develop and validate a new skeletal muscle organoid system for investigating human myogenesis with translational potential for disease modeling and therapy development.

Results

Generation of human fetal skeletal muscle organoids

Organoid cultures from pluripotent stem cells often require pre-patterning before Matrigel embedding to promote structural development **(**Lancaster et al., 2013; Spence et al., 2011; Koehler et al., 2017). In our 3D approach, we provided cells with immediate matrix support upon embryoid body formation to preserve cell state transitions. Initial pre- embedding screening indicated high expression of pluripotent markers OCT4, SOX2 and NANOG, moderate expression of neural tube marker PAX6 and low expression of mesodermal markers BRACHYURY and MSGN1 **(Figures 1, 1-figure supplement 1A)**. Upon Matrigel embedding, stimulation with CHIR99021, LDN193189 and bFGF promoted paraxial mesoderm formation through derivation of BRACHYURY$^+$ and TBX6$^+$ cells **(Figures 1A,C, 1-figure supplement 1A)**. Immunostaining at Day 5 depicted presence of mesodermal (BRACHYURY$^+$), paraxial mesodermal (TBX6$^+$) and neuromesodermal (SOX2$^+$/ BRACHYURY$^+$) progenitors (Gouti et al., 2017; Henrique et al., 2015) **(Figure 1C).** From this stage, we attempted to mimic determination front formation and promote anterior somitic mesoderm (ASM) (Aulehla and Pourquie,2010; Shimozono et al., 2013) by maintaining constant CHIR99021 and LDN193189 levels, reducing bFGF levels to 50% and simultaneously introducing retinoic acid to the culture. Consequently, distinct PAX3$^+$ but SOX2$^-$ cells emerged on the organoid surface **(Figure 1D)**, followed by a significant downregulation of PSM markers HES7, TBX6 and MSGN1 **(Figures 1E, 1-figure supplement 1A)**. Concomitantly, we observed significant upregulation of ASM marker MEOX2 and neural crest marker TFAP2A but not of SOX2 or PAX6, excluding a shift towards neural tube formation **(Figures 1E,F, 1-figure supplement 1A)**. Up to this stage, inter-organoid sizes showed small variability **(Figure 1B)**. Dermomyotomal fate was promoted by Sonic Hedgehog (SHH) and WNT1A stimulation, while maintaining BMP inhibition avoided lateral mesoderm formation **(Figure 1A)**. Expression profiling at Day 11 depicted presence of neural tube/crest (SOX2, PAX6, TFAP2A, SOX10 upregulation) and mesodermal (UNCX, TBX18, PAX3) lineages and further downregulation of PSM markers **(Figures 1E,F, 1-figure supplement 1A)**. Notably, the dermomyotomal/neural crest marker PAX7 emerged together with markers that define the dorsomedial (EN1) or ventrolateral portion (SIM1) of the dermomyotome (Cheng et al., 2004) **(Figures 1D,F, 1-figure supplement 1A)**.

Consequently, for favoring myogenesis we stimulated organoid culture with FGF and HGF (Charge and Rudnicki, 2004) **(Figure 1A)**. Surprisingly, at Day 17, organoids constituted a mosaic of neural crest and myogenic progenitor cells. Cells with epithelial morphology were TFAP2A$^+$, SOX2$^+$, PAX3$^+$ and PAX7$^+$, indicating formation of neural plate border epithelium (Roellig et al., 2017) **(Figure 1G)**. In cell clusters with mesenchymal morphology, we detected specified migrating

PAX3$^+$/SOX10$^+$ neural crest progenitors and PAX3$^+$ /SOX2$^-$/ SOX10$^-$ cells of myogenic origin **(Figure 1G)**. At this stage, myogenic lineages appeared to be primarily represented by PAX3$^+$ (9.35±0.07%) rather than PAX7$^+$ cells, as the PAX7 expression pattern (15.3±0.1% PAX3$^+$/PAX7$^+$, 5.11±0.13% PAX7$^+$) predominantly overlapped with that of SOX2 and TFAP2A **(Figures 1G, 1-figure supplement 1B)**. Morphologically, neural plate boarder and dermomyotomal populations exhibited uneven distribution, and thereby subsequent neural crest and myogenic migration processes introduced organoid size variability **(Figure 1B,G)**. From Day 17 onwards, we omitted FGF stimulation to cease neural crest development and promote delamination/migration of PAX3$^+$/ PAX7$^+$/SOX10$^-$ progenitor cells (Murphy et al., 1994) **(Figure 2A)**. Strikingly, until day 23, we observed committed myogenic populations through detection of fast MyHC myofibers in proximity of PAX7$^+$ but SOX2$^-$/TFAP2A$^-$/SOX10$^-$ negative cells **(Figures 1H, 1-figure supplement 1C to E)**. Consistently, expression profiling indicated downregulation of neural tube/crest lineage markers and significant upregulation of muscle precursor migrating markers (Buckingham and Rigby, 2014), such as LBX1, CXCR4 and MEOX2 **(Figures 1E, 2A, 2-figure supplement 2A)**.

At 8 weeks, organoids showed profound changes in transcription profiling **(Figures 2, 2-figure supplement 2B).** Gene ontology enrichment analysis highlighted an ongoing developmental transition with muscle development, muscle contraction and myofibril assembly among the top upregulated gene ontology terms and neurogenesis and nervous system development among the top downregulated terms **(Figure 2B)**. In addition, we detected downregulation of key markers characterizing neural tube and neural crest development (Soldatov et al., 2019), such as PAK3, DLX5, B3GAT1, FOXB1, HES5 and OLIG3 **(Figures 2, 2-figure supplement 2C)**. Interestingly, we could spatially visualize this process using immunocytochemistry for SOX2, TFAP2A and SOX10 expressing cells that were restricted to the inner portion of the organoid, and probably not susceptible to culture stimulation at 5 weeks **(Figures 1H, 2C,D, 2-figure supplement 2D)**. This neural / myogenic lineage spatial orientation could be visualized even at Day 84 through presence of TUJ1$^+$ neurons confined to inner organoid areas and close to SOX2$^+$ epithelium, while FastMyHC$^+$ myofibers occupied exterior areas **(Figure 2E).** On the other hand, substantial proportions of migratory cells in proximity of FastMyHC$^+$, MF20$^+$ myofibers expressed PAX7 (40±0.3%) but not SOX2, TFAP2A, SOX10 or MYOD1 (3.55±0.32%) (**Figures 2F, 2-figure supplement 2D,E**). This behavior is further illustrated by the presence of MYOD1$^+$ cells confined to organoid periphery **(Figure 2D)**.

Culture progression led to significant increase in PAX7$^+$ and MYOD1$^+$ cells at 8 weeks (45.3±3,4%

and 49.8±0.62%, respectively) (**Figures 2F, 2-figure supplement 2E**), suggesting an interval in which organoid culture predominantly commits to myogenic lineage. Expression profiling at this stage depicted upregulation of markers characteristic for limb migrating myogenic cells, e.g. LBX1, PAX7, SIX1/4, EYA1/4, PITX2, MYF5, TCF4 and MSX1 **(Figures 2, 2-figure supplement 2A)**. Consequently, HOX gene cluster profiling at 4 weeks correlated HOX A9, B9, C9 upregulation with limb bud site, while upregulation of the HOX 10-13 cluster during 8 to16 weeks attributed later organoid development to a more distal limb axial anatomical identity (Shubin et al., 1997; Xu et al., 2011, Raines et al., 2015) **(Figure 2G)**.

Lineage representation and developmental identity for skeletal muscle organoids

Analysis of organoid culture at single cell resolution indicated predominant presence of skeletal muscle lineage (n= 3945 cells, 91.26% of total population), complemented with two smaller cell clusters of mesenchymal / 'fibroadipogenic' (n=165 cells, 3.82% of total population) , and neural (n=213 cells, 4.93% of total population) origin **(Figures 3A, 3-figure supplement 1A)**. Skeletal muscle lineages further separated into distinct sub-clusters: myogenic muscle progenitors in non-dividing (PAX7$^+$/PAX3$^-$) and mitotic states (PAX7$^+$/CDK1$^+$/KI67$^+$), myoblasts (MYOD1$^+$), myocytes (MYOG$^+$) and myotubes (MYH3$^+$) **(Figures 3A, 3-figure supplement 1A,B)**. Further, investigation of mesenchymal cluster identity suggested fibroadipogenic potential of a distinct cell population with PDGFRa among the top upregulated genes (Uezumi et al., 2010,2011; Xi et al., 2020) **(Figures 3A,F, 3-figure supplement 1A,B).** Consistently, we detected upregulation of fibrotic markers. Moreover, adipogenic potential was highlighted through upregulation of PREF-1 and EBF2 **(Figure 3F)**. Occasionally, we detected structures resembling adipocytes from 9 to10 weeks onwards, with adipogenic nature verified by PRDM16 and Oil red O positive staining **(Figures 3B,D, 3-figure supplement 1C,D)**. A SOX2$^+$/TUBB3$^+$ positive neural cluster potentially derived from the neural plate border epithelium at younger stages could be located towards organoid interior **(Figures 3C, 3-figure supplement 1E).**

Myogenesis progression based on *t*-SNE feature and Violin plots of key markers from each stage indicated gradual transitions from myogenic progenitor to myotube subclusters **(Figures 3E, 3-figure supplement 1F)**. MyHC expression profiling at single cell resolution demonstrated predominant presence of embryonic (MYH3$^+$) and perinatal (MYH8$^+$) myofibers, co-expressing β-Enolase and M-cadherin in vicinity of PAX7$^+$ cells **(Figure 3 G - J)**. Consistently, we detected MCAM expression on muscle subclusters **(Figure 3G)**, while bulk RNA seq at 16 weeks highlighted expression of additional MyHC isoforms, e.g. MYH2 (II$_a$), MYH4(II$_b$) and transcription factor NFIX (Moore et al., 1993; Barbieri et al., 1990; Alexander et al. 2016; Messina et al., 2010). At more mature stages, we detected adult MYH1 or slow MYH7 isoforms **(Figures 3, 3-figure supplement**

1G), presumably due to myofiber stimulation via spontaneous contraction. In agreement, differential expression comparison between 8 and 16 week organoids indicated less variance in transcription profiling **(Figures 3, 3-figure supplement 1H, 4A)**. Consistently, sarcomere organization, ion transport, response to stimulus and synapse structure maintenance were among the upregulated gene ontology terms and mitosis, cell cycle, DNA packaging and nuclear division among the top downregulated terms **(Figures 3, 3-figure supplement 2B-D)**. Notably, ongoing maturation did not affect pool of progenitor cells, as even at 14 weeks we could report significant increases in PAX7$^+$ (50,16$\pm$2.19%) and MYOD1$^+$ (58.53$\pm$0.92%) cells **(Figures 2F, 3, 3-figure supplement 2E,F).**

Functionality and maturation of organoid derived myofibers

Regarding localization and functionality of organoid derived myofibers, immunocytochemistry revealed positive staining for dystrophin and a continuous laminin sheath around individual muscle fibers. Ultrastructure and 2- photon microscopy analysis depicted well developed sarcomeres **(Figures 3, 3-figure supplement 3A-D)**. Moreover, patch clamp experiments, upon superfusion of cells with acetylcholine (ACh), indicated inward current generations that rapidly declined to a low steady state level **(Figure 3K, 3-figure supplement 3E)**. The I/V-curve of this current component showed almost linear behavior and reversal potential around 0 mV **(Figure 3L)**. These data are in line with previous studies that described currents with analogous properties as characteristic nAChR currents (Jahn et al., 2001; Shao et al., 1998). Application of a fluorescent biosensor for monitoring changes in cytosolic Ca^{2+} (Twitch2B) revealed nAChR efficiency in modulating intracellular [Ca^{2+}]. Summarized FRET recordings **(Figures 3M, 3-figure supplement 3F)**, following application of ACh (10 μM), illustrated a rapid increase in FRET ratio that gradually declined in the presence of ACh, probably reflecting desensitization of nAChRs. These results demonstrated that nAChR in skeletal muscle cells are functional in terms of inducing Ca^{2+} release from intracellular stores.

Using our organoid protocol, we successfully derived fetal muscle progenitors and electrophysiologically functional myofibers from hiPSC lines with wild type and Duchenne Muscular Dystrophy genetic backgrounds (**Figures 3, 3-figure supplement 3E,F**).

Identity and sustainability of organoid derived PAX7 myogenic progenitors

Skeletal muscle organoids remarkably foster sustainable propagation of PAX7$^+$ progenitors **(Figures 2F, 3A)**. Initial screening of PAX7$^+$ progenitors indicated expression of several satellite cell markers (Fukada et al., 2007), such as CD82, CAV1, FGFR1, FGFR4, EGFR, M-Cadherin,

NCAM, FZD7, CXCR4, ITGβ1, ITGA7, SCD2 and SCD4 **(Figures 4, 4-figure supplement 1A).** Further investigation verified NOTCH signaling activity (HES1, HEYL, HEY1 and NRARP) in the myogenic subcluster, while dormant myogenic progenitors exhibited high expression of SPRY1 and cell cycle inhibitors p57 (CDKN1C), p21 and PMP22 (Xi et al., 2020; Fukada et al., 2007; Bjornson et al., 2012; Shea et al., 2010) **(Figures 4, 4-figure supplement 1B, 2A,D)**. Moreover, proliferation assays performed at the same stage (14 weeks) demonstrated 4.78±0.28% edU$^+$ cells, while substantial proportions of PAX7$^+$ myogenic progenitors remained quiescent **(Figures 4, 4-figure supplement 1C,D)**. t-SNE clustering divided myogenic progenitors into three distinct groups with unique molecular signatures: "CD44^{high}", "FBN1^{high}" and "CDK1$^+$" clusters **(Figures 4A, 4-figure supplement 2A)**. The "CD44^{high}" cluster, further characterized by CD98 upregulation, adopted a molecular signature similar to activated satellite cell state (Porpiglia et al., 2017) **(Figures 4A,B, 4-figure supplement 2A,D).** In this state, myogenic progenitors expressed VEGFA **(4-figure supplement 2A,D)** as previously described for murine satellite cells (Verma et al., 2018)**.** Consistently, the CD44$^+$ / PAX7$^+$ 'activated' myogenic population located at sites more accessible to HGF signaling, e.g. exterior organoid areas and forming bulges **(Figure 4B)**. The "FBN1^{High}" sub-cluster was further characterized by "PAX7^{high} / SPRY1^{high}/CHODL^{high} / FBN1^{high}" expression **(Figures 4A, 4-figure supplement 2A,D).** PAX7^{high} cells within the dormant state co-expressed NOTCH3, JAG1 and LNFG, together with CHODL markers. Further, we verified presence of FBN1$^+$ microfibrils, which, compared to CD44$^+$ cells, occupied areas without direct access to activation signals **(Figure 4C)**. Lastly, the "CDK1$^+$" cluster was the only proliferative population, further expressing markers mediating satellite cell activation and proliferation, such as DEK and EZH2 (Cheung et al., 2012; Juan et al., 2011) **(Figures 4A,D, 4-figure supplement 1A, 2A,B).**

Pseudotemporal ordering of myogenic progenitors indicated that the "FBN1^{High}" subcluster was the main progenitor population residing in dormant state, which by downregulating PAX7, FBN1 and CHODL and upregulating CD44, MYOD1, CD98 (SLC3A2) and VEGF generated activated state **(Figures 4D, 4-figure supplement 2B)**. In activated state, myogenic progenitors that upregulate MYOD1 entered mitosis, proliferated and followed a trajectory leading to myogenic commitment **(Figures 4D, 4-figure supplement 2B)**. This commitment was further accompanied by PARD3, p38a MAPK and CD9 expression (Porpiglia et al., 2017; Troy et al., 2012) **(4-figure supplement 2E).** Conversely**,** myogenic progenitors that upregulated NOTCH3, SPRY1 and PAX7 followed a loop trajectory that reinstated dormant stem cell state **(4-figure supplement 2D,E)**. Interestingly, the "FBN1^{high}" cluster highly expressed extracellular matrix collagens, e.g. COL4A1, COL4A2, COL5A1, COL5A2, COL5A3 and COL15A1. Notably, such expression declined upon

commitment and differentiation (**4-figure supplement 2C,F**).

Organoid derived myogenic progenitor comparison to human fetal and adult progenitor/stem cells

To evaluate developmental identity of the myogenic cluster, we isolated ITGβ1$^+$ / CXCR4$^+$ organoid derived myogenic progenitors via FACS (Garcia et al., 2018) and compared to human fetal muscle progenitors, demonstrating high similarity (Pearson´s correlation co-efficiency, rho=0.9), with myogenic markers such as PAX7, MYF5, MYOG and MYOD1 at comparable expression levels **(4-figure supplement 3A-F).** Differential expression comparison verified expression of extracellular matrix collagens and proteins, such as COL4A1, COL5A1, COL6A1, COL15A1, FBN1 and CHODL, in myogenic progenitors similar to 17th week human fetal tissue (Pearson´s correlation co-efficiency, rho=0.97) **(Figure 4E)**. Further, to evaluate myogenic potency in vitro, isolated ITGβ1$^+$ / CXCR4$^+$ organoid derived myogenic progenitor cells were re-plated and allowed to differentiate under the same organoid culture protocol, which demonstrated capacity to generate FastMyHC+ and PAX7+ populations within 10 days **(4-figure supplement 3B,C**). Subsequently, comparison to available transcriptomic dataset of human adult satellite cells at single cell resolution, divided myogenic progenitors and adult satellite cells into four clusters with distinct molecular signatures **(Figure 4F)**. Interestingly, myogenic progenitors were enriched for extracellular matrix proteins, while satellite cells mainly upregulated genes repressing transcription/translation, such as ZFP36 and TSC22D1, or related to early activation response, such as MYF5, JUN and FOS **(4-figure supplement 4A,B).** In line, organoid derived myogenic progenitors exhibited higher NOTCH signaling activity in comparison to satellite cells, with NOTCH3 and RBPJ being among enriched markers **(4-figure supplement 4B**). In contrast, adult satellite cells exhibited PAX7^{low} / MYF5^{high} expression profiling, presumably due to tissue dissociation, thereby indicating a tendency for activation rather than preservation or reinstatement of quiescency (Machado et al., 2017; Seale et al., 2000) **(Figure 4G).** Pseudotime ordering showed two distinct clusters, with adult satellite cells downstream from non-dividing myogenic progenitors **(4-figure supplement 4C).** Consistently, downregulation of genes promoting quiescence like PAX7, NOTCH3 and RBP, and upregulation of activation genes like MYF5, JUN and FOS along the trajectory (**4-figure supplement 4D)** was a further indication that organoid derived myogenic progenitors resided in dormant non-dividing state and that our organoid platform promoted sustainability of myogenic progenitors.

Organoid derived myogenic progenitors exhibited expression profiling similar to that of Stage 4

myogenic progenitors, correlating to 17-18th week human fetal development, as recently reported in the human skeletal muscle atlas (Xi et al., 2020). SPRY1, PLAGL1, POU4F1, TCF4, PAX7, CD82 and CD44 markers, as well as ECM proteins, nuclear factor I family members (NFIA, NFIB, NFIC) and NOTCH signaling are specifically upregulated markers. Furthermore, in comparison to 2D culture protocols, our organoid approach preserved myogenic progenitor dormancy without activating MYOD1 and exhibited higher expression of ECM proteins and key markers characterizing Stage 4 fetal myogenic progenitors (**4-figure supplement 5**). ERBB3 and NGFR markers (Hicks et al., 2017) demarcate populations of earlier developmental stages and were not upregulated either in organoid derived or Stage 4 fetal myogenic progenitors (**4-figure supplement 5**). In addition, organoid myogenic progenitors and Stage 4 fetal myogenic progenitors both downregulated cycling genes like MKI67, TOP2A, EZH2 and PCNA (**4-figure supplement 5**).

In addition, we have performed a developmental score distribution of myogenic progenitors based on the difference between up-regulated satellite cell and embryonic markers from the human reference myogenic atlases (Xi et al., 2020) and adult satellite cell data (Rubenstein et al., 2020, Xi et al., 2020) in comparison to our organoid protocol (**Figure 4H**). Organoid-derived myogenic progenitors represent the late fetal stage of maturation partially overlapping with adult satellite cell scoring. We note the heterogeneity of adult satellite cell populations when performing developmental scoring in line with recent reports (Barruet et al., 2020). Finally, we transplanted CD82-positive progenitors from our organoids into the tibialis anterior muscle of immunodeficient mice to complement our study with an in vivo experiment (Alexander et al., 2016, Marg et al., 2019, Al Tanoury et al., 2020). Six week post transplantation we could verify clusters of human Lamin A/C positive cells in the transplanted but not in the control group (**Figure 4I,J**).

Discussion

Human skeletal muscle organoids offer a new cell culture system to study human myogenesis, in particular fetal myogenic progenitors. We demonstrate that modulation of Matrigel embedded embryonic bodies with WNT, BMP and FGF signaling at early stages leads to paraxial mesoderm formation **(Figure 1B)**. Further, under guided differentiation, we could promote concomitant development of neural and paraxial mesodermal lineages and derive mesodermal populations with somitic and dermomyotomal origin **(Figure 1C - F).** With WNT1A and SHH stimulation, neural lineage is directed towards dorsal neural tube / crest development which benchmarks the structural recapitulation of neural plate border epithelium **(Figure 1G)**. In vitro neural plate border can be utilized to assess transcriptomic networks and cell fate decision during human neural crest formation.

Delaminating from the dermomyotome, undifferentiated PAX3 progenitor cells reorientate and align in the cranio- caudal axis to form the underlying myotome or completely detach from the dermomyotome and migrate into the surrounding mesenchyme at the limb sites, where they propagate and become committed to skeletal muscle progenitors (Relaix et al., 2005). By stimulating organoid culture at the neural plate border/dermomyotomal stage with bFGF/HGF we could further visualize both migration of myogenic progenitors and migration/specification of neural crest populations (**Figures 1A,G, H, 2A).** Further, by omitting FGF during organoid development, we could detect a continuous upregulation of genes involved in the myogenic migration process, such as LBX1, PAX3, PAX7 and CXCR4, but not in genes characterizing neural tube or neural crest development, such as SOX10, TFAP2A, PAK3, DLX5, B3GAT1, FOXB1, HES5 and OLIG3. This indicates that organoid culture conditions and specifically stimulation with HGF favored skeletal muscle over neural lineage **(Figures 2C-E)**. Interestingly, we could show that by stimulating organoid culture with SF/HGF, an essential signal for detachment and subsequent migration, but not for specification of the cells at the lateral lip of the dermomyotome (Dietrich et al., 1999), we could preserve the PAX3+/PAX7+ myogenic migratory population in an undifferentiated and uncommitted state **(Figure 2D,E)**. Strikingly, expression profiling based on HOX gene cluster supported this notion, as over time the organoid culture adopted more distal than proximal limb axial anatomical identify **(Figure 2F)**.

Fetal development is characterized by PAX3+ to PAX7+ myogenic progenitor transition (Seale et al., 2000; Relaix et al., 2005), which we were able to demonstrate in our organoid culture. Our data further support organoid culture representing fetal stages as we could detect NFIX upregulation and verify presence of myofibers expressing fetal MyHC isoform as well as NCAM, M-Cad or MCAM in the proximity of PAX7 progenitors **(Figure 3G to J)**. Consequently, transcriptome comparison indicated high similarity to human fetal muscle tissue (17-week, Pearson correlation, rho=0.9) **(Figure 4E)**, as well as expression of several satellite cell markers **(4-figure supplement 1A).** This was further verified by comparison to a single cell dataset from the human skeletal muscle cell atlas **(Figures 4, 4-figure supplement 4)** Interestingly, single cell resolution showed adult satellite cell clusters with organoid derived myogenic progenitors **(Figure 4F)**. In addition, pseudotime ordering indicated that organoid derived myogenic progenitors reside developmentally upstream of activated satellite cells, upregulating markers associated with quiescence such as NOTCH signaling and extracellular matrix proteins **(4-figure supplement 2E,F)**. Preservation of myogenic progenitors in a non-dividing state without activating MYOD1 and self-renewal **(Figures 4D, 4-figure supplement 2E,F))** appeared to be responsible for the observed sustainable propagation of PAX7 progenitor cells even 14 weeks post differentiation **(Figures 2F, 3A).**

Patterning in our 3D protocol provide progenitors in a more mature late fetal state partially overlapping with adult satellite cell developmental scoring **(Figure 4H)**. The 'uncommitted' Pax7

progenitor status is demonstrated by our precursor populations bulkRNAseq and scRNAseq profiling **(Figure 4 A,E, F).** In this context, we could observe high expression of extracellular matrix proteins and upregulated NOTCH signaling in dormant non-dividing myogenic progenitors **(4-figure supplement 2A,E)**. This phenotype is similarly described for human fetal tissue myogenic progenitors (**Figure 4E,** Pearson correlation, rho=0.97**, 4-figure supplement 5)**. Studies evaluating engraftment potential of satellite cells and fetal muscle progenitors propose that muscle stem cells in a quiescent non- committed state exhibit enhanced engraftment potential (Hicks et al., 2017; Quarta et al., 2016; Montarras et al., 2005; Tierney et al., 2016). Our data demonstrate that upon activation and commitment dormant myogenic progenitors downregulate extracellular matrix proteins and upregulate expression of morphogens / receptors that make them susceptible to signals, like VEGFA that communicates with vasculature during tissue reconstitution, or CD9, CD44, CD98 participating in cellular activation (Porpiglia et al., 2017) **(2-figure supplement 1B,D)**. CD82+ populations from our organoids engraft in the tibialis anterior muscle of immunodeficient mice (**Figure 4I,J**). It would be of interest for future studies to investigate whether increased engraftment can be achieved in 3D protocols (Faustino Martins et al., 2020; Shahriyari et al., 2022; ours) versus 2D patterned progenitor cells and to which degree this is attributed to high expression of extracellular matrix proteins. In particular, high Fibrillin1 expression on dormant non-dividing myogenic progenitors could potentially contribute to avoidance of fibrosis by myogenic progenitors through regulation of TGF-β signaling (Cohn et al., 2007)*.*

Different phases of human fetal myogenesis have been modelled with 2D differentiation protocols (Shelton et al., 2014, Chal et al. 2015, Xi et al. 2017). Our 3D differentiation protocol does not go beyond these protocols when it comes to provide maturated physiologically responsive skeletal muscle cells, which we illustrate with the electrophysiological recording of organoid-derived cells of different origins (**Figures 3 K,L,M, 3-figure supplement 3E,F**). Structural distinctions like the posterior paraxial mesoderm at day 5, specified neural crest dermomytome at day 17, myogenic progenitor migration at day 23 and neural crest lineage arrest at day 35 (**Figures 1 C-H, 2 C**) cannot be similarily observed in 2D protocols. In addition, our 3D organoid protocol provides myogenic progenitors in dormant and activated states for at least 14 weeks in culture. We demonstrate that organoid culture sustains uncommitted MyoD1-negative, Pax7-positive myogenic progenitors as well as fibroadipogenic (PDGFRa+) progenitors, both resembling their fetal counterpart. This supply of muscle progenitors preserved in a quiescent state indicates translative potential of the approach. Future work will elucidate signaling transduction pathways during skeletal muscle organoid development to model and understand human myogenesis in more detail.

Materials and Methods

hiPSCs culture

Human induced pluripotent stem cell (hiPSC) lines, Cord Blood iPSC (CB CD34$^+$, passage 15 – 35), Duchenne Muscle Dystrophy iPSC patient line iPSCORE_65_1 (WiCell, cat.no. WB60393, passage 22-30) and DMD_iPS1 (passage 21-30) & BMD_iPS1 (passage 17-25) (Boston Children's Hospital Stem Cell Core Facility), LGMD2A iPSC and LGMD2A-isogenic iPSC (Dorn et al., 2015; Mavrommatis et al., 2020; Park et al., 2008; Panopoulos et al., 2017) were cultured in TESR-E8 (StemCell Technologies) on Matrigel GFR (Corning) coated 6 well plates. The study was performed after ethical approval from the ethics commission of the Ruhr-University Bochum, Medical Faculty (15-5401, 08/2015).

Human skeletal muscle organoid differentiation protocol

Prior differentiation, undifferentiated human PSCs, 60-70% confluent, were enzymatically detached and dissociated into single cells using TrypLE Select (ThermoFisher Scientific). Embryoid bodies formed via hanging drop approach with each droplet containing 3-4x10^4 human single PSCs in 20 µl were cultured hanging on TESR-E8 supplemented with Polyvinyl Alcohol (PVA) at 4mg/ml (SigmaAldrich) and rock inhibitor (Y-27632) at 10µM (StemCell Technologies) at the lid of Petri dishes. At the beginning of skeletal muscle organoid differentiation, embryoid bodies at the size of 250-300µm embedded into Matrigel and cultured in DMEM/F12 basal media (ThermoFisher Scientific) supplemented with Glutamine (ThermoFisher Scientific), Non Essential Amino Acids (ThermoFisher Scientific), 100x ITS-G (ThermoFisher Scientific), (Basal Media) 3µM CHIR99021 (SigmaAldrich) and 0.5µM LDN193189 (SigmaAldrich). At Day 3, human recombinant basic Fibroblast Growth Factor (bFGF) (Peprotech) at 10ng/µl final concentration added to the media. Subsequently, at Day 5 the concentration of bFGF reduced at 5ng/µl and the media further supplemented with 10nM Retinoic Acid (SigmaAldrich). The differentiation media, at Day 7, supplemented only with human recombinant Sonic hedgehog (hShh) (Peprotech) at 34ng/µl, human recombinant WNT1A (Peprotech) at 20ng/µl and 0.5µM LDN193189. At Day 11 the cytokine composition of the media changed to 10ng/µl of bFGF and human recombinant Hepatocyte Growth Factor (HGF) at 10ng/µl (Peprotech). From Day 15 onwards, the basal media supplemented with ITS-X (ThermoFisher Scientific) and human recombinant HGF at 10ng/µl. The first 3 days of the differentiation the media changed on a daily basis, from 3rd till 30th every second day, while from Day 30 onwards every third day. We evaluated the organoid approach with six hiPSCs lines with independent genetic background, with more than five independent derivations per line, especially for the control line (CB CD34$^+$) more than 20 derivation, obtaining always similar results. Per derivation and upon embryoid body matrigel embedding, cultures exhibited high reproducibility.

Upon migration and skeletal muscle formation organoids are occupying whole matrigel droplet and through generation of additional bulge reach size of 4-5 mM. Concomitantly, myogenic progenitors fall off the organoid and generate a sheath of myogenic organoids and muscle fibers at the surface of the culture plate. For all the lines, we could demonstrate functional myofibers and PAX7 positive myogenic populations. Myogenic populations from different lines exhibit high similarity, (Pearson correlation, rho = 0.94-0,95, Figure 4 - figure supplement 3G).

Immunocytochemistry

Cryosection Immunochemistry: Organoids from different stages were fixed on 4% paraformaldehyde overnight at 4^{O}C under shakings conditions, dehydrated (30% sucrose o/n incubation) and embedded in OCT freezing media. Cryosections were acquired on a Leica CM3050s Cryostat. For the immunostaining process, cryosections were rehydrated with PBS and followed by permeabilization once with 0.1% Tween-20 in PBS, (rinsed 3x with PBS), and then with 0.1% Triton-X in PBS (rinsed 3x with PBS). Subsequently, the sections blocked with 1% BSA / 10% NGS in PBS for 1hr at room temperature. Primary antibody incubations were performed from 1 to 2 days at 4^{O}C, where secondary antibody incubations for 2hr at room temperature.

EdU staining: At 12-week post differentiation, control organoids incubated with 2.5μM BrdU final concentration overnight. To detect EdU, the sections were processed with Click-iT EdU Alexa Fluor 488 cell proliferation kit (Invitrogen) following the manufacturer's instructions. The samples incubated with secondary antibodies after the click reaction for detecting EdU. *Primary Antibodies*: anti-Brachyury (R&DSystems, 1:250), anti-TBX6(Abcam, 1:200), anti-PAX3 (DHSB, 1:250), anti- PAX7(DHSB, 1:250), anti-SOX10 (R&DSystems,1:125), anti-KI67 (ThermoFisher Scientific, clone SolA15, 1:100), anti- TITIN (DHSB, 9D-10, 1:300), anti-MyHC (DHSB, MF20, 1:300), anti-MYOD1 (Santa Cruz Biotechnologies, clone 5.8A, 1:200), anti-PRDM16 (Abcam, ab106410, 1:200), anti-TFAP2A (DHSB, 3B5, 1:100), anti-Dystrophin (Novocastra/Leica Biosystems, clone DY4/6D3, 1:200), anti-Laminin (SigmaAldrich, 1:200), anti-FastMyHC (SigmaAldrich, clone MY-32, 1:300), anti-M-Cadherin (Cell Signaling Technology, 1:200) anti-SOX2 (ThermoFisher Scientific, clone Btjce, 1:100), anti-CD44 (eBioscience, clone IM7, 1:100), anti-Fibrillin1 (Invitrogen, clone 11C1.3, 1:100).

Secondary antibodies: Alexa Fluor® 647 AffiniPure Fab Fragment Goat Anti-Mouse IgM, μ Chain Specific (Jackson Immunoresearch Laboratories, 1:100), Rhodamine RedTM-X (RRX) AffiniPure Goat Anti-Mouse IgG, Fcγ Subclass 1 Specific (Jackson Immunoresearch Laboratories,1:100), Alexa Fluor® 488 AffiniPure Goat Anti-Mouse IgG, Fcγ subclass 2a specific (Jackson Immunoresearch Laboratories,1:100), Alexa Fluor 488, Goat anti-Rat IgG (H+L) Cross- Adsorbed Secondary Antibody, (ThermoFisher Scientific, 1:500), Alexa Fluor 488, Donkey anti-Mouse IgG

(H+L) Cross-Adsorbed Secondary Antibody, (ThermoFisher Scientific, 1:500), Alexa Fluor 647, Donkey anti-Goat IgG (H+L) Cross-Adsorbed Secondary Antibody, (ThermoFisher Scientific, 1:500), Alexa Fluor 488, Donkey anti-Goat IgG (H+L) Cross-Adsorbed Secondary Antibody, (ThermoFisher Scientific, 1:500), Alexa Fluor 568, Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, (ThermoFisher Scientific, 1:500). Images were acquired on a ZEISS LSM780 inverted confocal microscope.

Oil O Red Staining

For histological visualization of adipocytes within the organoids, Oil O Red Stain kit (Abcam, ab150678) applied on frozen sections derived from PFA fixated organoids following manufactures recommended protocol. Organoid sections upon staining with Oil O Red were visualized at an Olympus BX61 upright microscope.

Flow Cytometry

FACS intracellular staining: Organoids during the 2^{nd}, 5^{th}, 9^{th} and 14^{th} week of differentiation were dissociated into single cells by incubating them till dissociating at 37^{o}C within papain solution under shaking conditions. Then, the cells pelleted at 400xg for 5min and followed incubation with TryplE Select for 10min to ensure dissociation into singe cells. Further, the cells passed through 70µM (2^{nd} week) – 100µM (5^{th}, 9^{th}, $14t^{h}$ week) cell strainer to avoid aggregates. For both digesting steps 10% FBS/DMEM-F12 as digesting deactivation solution applied to the cells. Then, for intracellular flow cytometric staining analysis the Transcription Factor Buffer set (BD Pharmigen) was applied and the cells were analyzed using flow cytometer (BD Biosciences FACS ARIAII). Primary antibodies used in this study: anti-PAX7, anti- MYOD1, anti-Pax3 in total amount of 400µg per staining, while as secondaries the Rhodamine RedTM-X (RRX) AffiniPure Goat Anti-Mouse IgG, Fcγ Subclass 1 Specific (Jackson Immoresearch Laboratories), Alexa Fluor® 488 AffiniPure Goat Anti-Mouse IgG, Fcγ subclass 2a specific (Jackson Immoresearch Laboratories) in 1:50 dilution. As isotype controls, Mouse IgG1 kappa Isotype Control (Invitrogen, clone P3.6.2.8.1), Mouse IgG2a kappa Isotype Control, (Invitrogen, clone eBM2a) were used at 400µg total amount per staining.

FACS EdU assay: At 15-week post differentiation, organoids incubated overnight with 5µM EdU final concentration. Next day, organoids were dissociated into single cells by incubation at 37^{o}C within papain solution for 1-2hr, followed by incubation with TryplE Select for 10min to ensure single cell dissociation. Then, the dissociated cells passed through a 70µm cell culture strainer to remove any remaining aggregates. To detect EdU, the cells were processed with Click- iT EdU Alexa Fluor

488 Flow Cytometry Assay Kit (Invitrogen) according to manufacturer instructions and then analyzed using the 488 channel of a BD Biosciences FACSAria Fusion flow cytometer.

FACS isolation of ITGβ1$^+$ / CXCR4$^+$ myogenic cell population for RNA sequencing: Organoids from Duchenne and Control iPSC lines and during 15th - 16th week post differentiation dissociated into single cells during incubation with Papain solution till we could observe dissociation upon gentle shaking (1-2hr). To acquire singlets, we filter the cells with through 40μm cell strainer and upon washing with 1% BSA solution we proceed with surface antigen staining. For surface antigen staining 20 min incubation with the Alexa Fluor 488 anti-human CD29 (Biolegend, clone TS2/16), PE anti-human CD184[CXCR4] (Biolegend, clone 12G5) were applied together with the corresponding isotype controls: PE Mouse IgG2a, κ Isotype Ctrl antibody (Biolegend, clone MOPC-173), 488 Mouse IgG1, κ Isotype Ctrl antibody (Invitrogen, clone P3.6.2.8.1). For removing residual antibodies, the cells were washed twice with 1% BSA staining solution and processed by BD Biosciences FACSAria Fusion flow cytometer. Briefly before FACS sorting to discriminate between dead and alive cells DAPI was added to the samples and then DAPI$^-$ / CD29$^+$ / CXCR4$^+$ cells populations were collected into tubes containing RLT buffer supplemented with b-mercaptoethanol to avoid RNA degradation. FACS gating strategy is further depicted in Figure 5 – figure supplement 3A.

Bulk RNA sequencing

RNA extraction: Total RNA was extracted from single organoids or cultured cells by using the RNAeasy Micro Kit (Qiagen) according to the manufacturer's instructions. Subsequently, before library preparation, the RNA integrity was evaluated on an Agilent 2100 Bioanalyzer by using the RNA 6000 Pico kit (Agilent). *cDNA library preparation*: For 4-week and 8-week organoids cDNA library was prepared by using the whole transcriptome Illumina TruSeq Stranded Total RNA Library Prep Kit Gold (Illumina), followed by evaluation on an Agilent 2100 Bioanalyzer by using the DNA 1000 kit. The resulting mRNA library sequenced as 2 X 75 bp paired-end reads on a NextSeq 500 sequencer (Illumina). For 16-weeks and ITGβ1$^+$ / CXCR4$^+$ sorted cells cDNA library was prepared using the whole transcriptome Ovation Solo RNA seq Library Preparation Kit (TECAN, NuGEN), followed by evaluation on an Agilent 2100 Bioanalyzer by using the DNA 1000 Chip. The resulting mRNA library sequenced as 1 X 150 bp single reads on a HiSeq. 3000 sequencer (Illumina).

Bulk RNA seq bioinformatic analysis

Sequenced reads were aligned to the human reference genome (hg38) with TopHat2 (version 2.1.1), and the aligned reads were used to quantify mRNA expression by using HTSeq-count (version 0.11.2). DESeq2 (Love at al., 2014) was used to identify differentially expressed genes

across the samples. ITGβ1$^+$ / CXCR4$^+$ organoid derived myogenic cell populations were compared to already available transcriptomic dataset of human fetal muscle progenitors (GSM2328841-2) (Hicks et al., 2018).

scRNA sequencing

Sample and cDNA library preparation: Single cells acquired upon incubation for 1hr with solution containing papain and EDTA. Upon dissociation, we estimated the cell number and viability; then cells resuspended on solution containing 0.5% BSA in PBS to reach concentration of 390 cells per μl. The cDNA library was prepared using the Chromium Single Cell 3′ Reagent Kits (v3): Single Cell 3′ Library & Gel Bead Kit v3 (PN-1000075), Single Cell B Chip Kit (PN-1000073) and i7 Multiplex Kit (PN-120262) (10x Genomics) according to the manufacturer's instructions. Then, the cDNA library was run on an Illumina HiSeq 3000 as 150-bp paired-end reads.

Single cell RNA seq bioinformatic analysis

Sequencing data were processed with UMI-tools (version 1.0.0), aligned to the human reference genome (hg38) with STAR (version 2.7.1a), and quantified with Subread featureCounts (version 1.6.4). Data normalization and further analysis were performed using Seurat (version 3.1.3, Stuart et al., 2019). For initial quality control of the extracted gene- cell matrices, we filtered cells with parameters low threshold = 500, high threshold =6,000 for number of genes per cell (nFeature_RNA), high threshold = 5 for percentage of mitochondrial genes (percent.mito) and genes with parameter min.cells = 3. Filtered matrices were normalized by LogNormalize method with scale factor = 10,000. Variable genes were found with parameters of selection.method = "vst", nfeatures = 2000, trimmed for the genes related to cell cycle (KEGG cell cycle, hsa04110) and then used for principal component analysis. Statistically significant principal components were determined by JackStraw method and the first 5 principle components were used for non-linear dimensional reduction (tSNE and UMAP) and clustering analysis with resolution=0.2. Monocle3 (version 0.2.0, Cao et al., 2019) was used for pseudotime trajectory analysis. We imported data matrix of Seurat objects (assays[["RNA"]]@counts) to Monocle R package and then performed dimensionality reduction with PCA method with parameters max_components=2 and then cluster_cells, learn_graph and order_cells functions were performed subsequently. Organoid-derived myogenic progenitors were compared to already available transcriptomic dataset of adult satellite cells (GSE130646) (Rubenstein et al., 2020).

Developmental score was calculated as described in Xi et al. 2020. Briefly, we used the ''AddModuleScore'' function to calculate the embyonic and adult score using a list of differentially expressed genes (DEGs) between adult and embryonic myogenic progenitor clusters. DEGs were

selected from the Supplementary information in Xi et al. 2020 table mmc3. The developmental score was further calculated by subtracting embryonic from the adult score. Embryonic and fetal datasets are filtered using a low threshold = 500 and adult datasets are filtered using a low threshold = 250 for genes per cell. In addition, we used scaling the data ("S.Score," "G2M.Score," "Stress" and "total Count" to the vars.to.regress' argument) to regress out the effects of the cell cycle, dissociation-related stress as well as cell size/sequencing depth to all datasets. Myogenic subpopulation of SMO and adult satellite cell datasets are selected by "PAX7" expression. Embryonic and fetal (weeks 5 to 18) and adult satellite cell (years 7,11,34,42) scRNAseq data are from Xi et al., 2020 (GSE147457), adult satellite cell (year 25) scRNAseq data are from Rubenstein et al., 2020 (GSE130646).

qPCR expression analysis

By pooling 3 organoids per sample, total RNA was extracted using the RNAeasy Mini Plus Kit (Qiagen). For first strand cDNA synthesis, the High capacity RNA to cDNA Kit (Applied Biosystems) applied using as template 2µg of total RNA. For setting qPCR reactions, the GoTaq qPCR Master Mix (Promega) was used with input template 4ng cDNA per reaction while the reaction was detected on a CFX 96 Real-Time PCR detection system (BIO-RAD). The relative quantification (ΔCT) method applied for detecting changes in gene expression of pluripotent, neural tube, neural crest, posterior/anterior somitic; and dermomyotomal markers, between different time points along the differentiation. qPCR primers applied for each marker are listed in **Table S1.**

Transmission Electron Microscopy (TEM)

Skeletal muscle organoids were fixed for 4h at room temperature (RT) in 2.5% glutaraldehyde (SigmaAldrich) in 0.1M cacodylate buffer pH 7,4 (Sciences Services, Germany), subsequently washed in 0.1M cacodylate buffer pH 7,4, post- fixed for 2h at RT in 1% osmium tetroxide (Sciences Services, Germany) in 0.1M cacodylate buffer pH 7,4, dehydrated stepwise in a graded ethanol series and embedded in Epon 812 (Fluka, Buchs, Switzerland). Ultrathin sections (70 nm, ultramicrotome EM UC7, Leica, Wetzlar, Germany) were afterwards stained for 30 min in 1% aqueous uranyl acetate (Leica, Germany) and 20 min in 3% lead citrate (Leica, Germany). TEM images were acquired with a 200 kV TEM JEM 2100Plus (Jeol, Japan), transmission electron microscope.

Second harmonic generation (SHG) imaging using multi-photon microscopy

A TriM Scope II multi photon system from LaVision BioTec was used to visualize skeletal muscle fiber organization inside organoids and distinct sarcomeres. The microscope setup is a single beam instrument with an upright Olympus BX51 WI microscope stand equipped with highly sensitive non-

descanned detectors close to the objective lens. The TriM Scope II is fitted with a Coherent Scientific Chameleon Ultra II Ti:Sapphire laser (tuning range 680-1080 nm) and a Coherent Chameleon Compact OPO (automated wavelength extension from 1000 nm to 1600 nm). A 20x IR objective lens (Olympus XLUMPlanFl 20x/1.0W) with a working distance of 2.0 mm was used. Muscle fiber SHG signals were detected in forward direction using TiSa light at 850 nm, a 420/40 band pass filter and a blue-sensitive photomultiplier (Hamamatsu H67080-01). 3D-images were acquired and processed with LaVision BioTec ImSpector Software.

Electrophysiology

Current measurement: Membrane currents were measured at ambient temperature (22-24°C) using standard whole- cell patch clamp software ISO2 (MFK, Niedernhausen, Germany). Cells were voltage-clamped at a holding potential of –90 mV, i.e. negative to EnAChR, resulting in inward Na^+ currents. Every 10 s, voltage ramps (duration 500 ms) from -120 mV to +60 mV were applied to assess stability of the recording conditions and to generate I/V curves (membrane currents in response to depolarizing voltage ramps are shown as downward deflections). Signals were filtered (corner frequency, 1 KHz), digitally sampled at 1 KHz and stored on a computer equipped with the hardware/software package ISO2 for voltage control, data acquisition and data analysis. Rapid exposure to a solution containing acetylcholine was performed by means of a custom-made solenoid-operated flow system permitting a change of solution around an individual cell with a half time of about 100 ms. For measurements cells devoid of contact with neighboring cells were selected.

Fluorescence microscopy and imaging: To monitor changes in $[Ca2^+]i$, skeletal muscle cells were transiently transfected with pcDNA3[Twitch-2B] (Addgene, 49531) (0.25 µg per 35 mm culture-dish). Skeletal muscle cells were transfected using either poly-ethyleneimine (PEI) or Lipofectamine (Invitrogen) according to the manufacturer's instructions. Prior to experiments, cells were seeded on sterile, poly-L-lysine-coated glass cover slips and analyzed or 48 h after transfections. All experiments were performed using single cells at ambient temperature. Fluorescence was recorded buckusing an inverted microscope (Zeiss Axiovert 200, Carl Zeiss AG, Göttingen, Germany) equipped with a Zeiss oil immersion objective (100x/1.4), a Polychrome V illumination source and a photodiode-based dual emission photometry system suitable for CFP/YFP-FRET (FEI Munich GmbH, Germany). For FRET measurements, single cells were excited at 435 nm wavelength with light pulses of variable duration (20 ms to 50 ms; frequency: 5 Hz) to minimize photo- bleaching. Corresponding emitted fluorescence from CFP (F480 or FCFP) or from YFP (F535 or FYFP) was acquired simultaneously and FRET was defined as ratio FYFP/FCFP. Fluorescent signals were recorded and digitized using a commercial hardware/software package (EPC10 amplifier with an integrated D/A board and Patch-master

software, HEKA, HEKA Elektronik, Germany). The individual FRET traces were normalized to the initial ratio value before agonist application (FRET/FRET0).

Solutions and chemicals: For FRET measurements an extracellular solution of the following composition was used (mmol/L): NaCl 137; KCl 5.4; $CaCl_2$ 2; $MgCl_2$ 1.0; Hepes/NaOH 10.0, pH 7.4. For whole cell measurements of membrane currents an extracellular solution of the following composition was used (in mmol/L): NaCl 120, KCl 20, $CaCl_2$ 0.5, $MgCl_2$ 1.0, HEPES/NaOH 10.0 (pH 7.4). The pipette solution contained (in mmol/L): K-aspartate 100, KCl 40, NaCl 5.0, $MgCl_2$ 2.0, Na_2ATP 5.0, BAPTA 5.0, GTP 0.025, and HEPES/KOH 20.0, pH 7.4. Standard chemicals were from Merck. EGTA, HEPES, Na_2ATP, GTP and acetylcholine chloride, were from SigmaAldrich.

Transplantation experiments

Flow cytometry: The organoid culture was dissociated using TrypLe (Thermo Fisher Scientific, 12563011) and were filtered through 70 - 40 µm cell strainer. Single cell suspension was stained with PE anti-human CD82 Antibody (BioLegend, 342103) in 2%BSA and 2mM EDTA and sorted using FACS sorter (Beckman Coulter, MoFlo Astrios Zellsortierer). For gating, unstained single cells from the same organoid culture were used as a baseline control. Sorted cells were further processed for transplantation experiments.

Transplantation of skeletal muscle progenitors into murine tibialis anterior (TA) muscle: 12 week old organoids are dissociated and single cells were sorted using FACS (CD82+). Cell Transplantation was carried out as described before (Alexander et al., 2016, Marg et al., 2019, Al Tanoury et al., 2020). Briefly, 24h before transplantation, Cardiotoxin (10 ul CTX, 40ng/ml, Sigma-Aldrich, 217503-1MG) was injected into TA of 2-3 month old male HsdCpb:NMRI-Foxn1nu mice. Under anesthesia $1*10^5$ cells are injected into the TA muscle on one side. 6 weeks after transplantation the mice are killed and the TA were fixed and sliced using cryosections. Immunofluorescence analyses are carried out using recombinant Anti-Lamin A + Lamin C antibody and anti-Dystrophin antibody.

All animal experiments were approved by the local authorities (81-02.04.2020.A476) and performed in accordance with the guidelines for Ethical Conduct in the Care and Use of Animals.

Cryosection Immunochemistry: TA from mice were fixed on 4% paraformaldehyde overnight at 4°C and kept in increasing sucrose solutions (10%, 20%, 30%) at 4 °C until the tissue sinked down (usually 24h) and embedded in Tissue-Tek® O.C.T.™ Compound media. Cross-sections of TA with 10 - 20 µm thickness were acquired on CryoStar NX50 (Thermo Scientifc). Sections are stored on objectives at -20°C.

Cryosections were rehydrated with PBS and followed by an antigen retrieval, an AffiniPure Fab Fragment Goat Anti-Mouse IgG treatment to prevent unspecific bindings during the staining process and permeabilized once with 1% (vol/vol) Triton-X100 and 125 mM glycine in PBS, (rinsed 3x with PBS). Subsequently, the sections blocked with 5% BSA / 10% NGS in PBS for 1hr at room temperature. Primary antibody incubations were performed for 1 day at 4°C and secondary antibody incubation for 2hr at room temperature.

Primary Antibodies: anti-Dystrophin (Leica, NCLDYS1, 1:20), recombinant Anti-Lamin A + Lamin C antibody (Abcam, ab108595, 1:150) Secondary antibodies: Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 488 (1:1000), Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 568 (1:1000)

Image acquisition and processing. Images were acquired on a Zeiss Scan.Z1. Images were processed using Zen Lite Blue version 4.0.3.

Statistics

All statistical analysis was conducted using GraphPad Prism6 software. For qPCR analysis we performed one-way ANOVA with Tukey´s multiple comparisons test for each marker. For the FACS intracellular staining quantification, we performed one-way ANOVA with Sidak´s multiple comparisons test between the different time points. For Western blot quantification we performed Mann-Whitney test between two groups. Significance asterisks represent *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, ns: not significant.

Data and code availability

RNA sequencing datasets produced in this study are deposited in the Gene Expression Omnibus (GEO) under accession code GSE147514. Detailed scripts and parameters used for the study are available from the authors upon reasonable request.

To review GEO accession GSE147514:
Go to https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE147514

ACKNOWLEDGEMENTS

We are grateful to Drs. Karl Köhrer, Tobias Lautwein, Patrick Petzsch and Thorsten Wachtmeister, Genomics & Transcriptomics Laboratory, Heinrich-Heine-University Düsseldorf for performing single cell and bulk RNAseq experiments with their Illumina HiSeq platform and data provision. We are further grateful to Dr. Oliver Griesbeck for the pcDNA3[Twitch-2B] plasmid. We would like to thank Ingrid Gelker, Martina Sinn, Max Planck Institute Münster as well as Eva-Maria Konieczny, Rana Houmany, Boris Burr, Ruhr-University Bochum for their technical assistance. We thank Dr. Johnny Kim, Max Planck Institute Bad Nauheim for discussions and Dr. Elisabeth Stevens, *English Scientific*, Düsseldorf for scientific editing of the manuscript. Electron microscopy experiments were supported by the Deutsche Forschungsgemeinschaft SFB 944. We thank Drs. George Q. Daley and Thorsten Schlaeger, Boston Children's Hospital for providing the Duchenne Muscular Dystrophy patient-derived iPS cell lines DMD-iPS1 and DMD-iPS2 in the course of our study. Our study was supported by research grants from FoRUM F873- 16, Medical Faculty, Ruhr University Bochum, from Deutsche Gesellschaft für Muskelkranke e.V (DGM Foundation), Freiburg, the Georg E. und Marianne Kosing-Stiftung, Deutsches Stiftungszentrum, Essen and the Deutsche Duchenne Stiftung.

AUTHOR CONTRIBUTIONS

L.M. conceived and designed the study, designed and performed experiments, single cell bioinformatic analyses, analyzed data, and wrote manuscript, H.W.J., U.K., D.G., J.H.Y.. M.J.A.B. performed (sc)RNAseq experiments and analyzes, U.K., G.G.G., M.C.K., M.S., O.E.P., D.Z., M.G.B., D.H., G.M.P, S.H.A. performed experiments and analyzed data, J.B.K., J.C.S., R.H.A., H.R.S. provided study materials and supervised experiments, M.V. provided study materials, biopsies and supervised experiments, B.B.S. provided study materials, supervised study and edited manuscript, H.Z. supervised study, experimental design, provided study materials and wrote manuscript.

REFERENCES

Alexander MS, Rozkalne A, Colletta A, Spinazzola JM, Johnson S, Rahimov F, Meng H, Lawlor MW, Estrella E, Kunkel LM, Gussoni E. 2016. CD82 is a marker for prospective isolation of human muscle satellite cells and is linked to muscular dystrophies. *Cell Stem Cell* **19:**800–807. DOI: https://doi.org/10.1016/j.stem.2016.08.006

Al Tanoury Z, Rao J, Tassy O, Gobert B, Gapon S, Garnier JM, Wagner E, Hick A, Hall A, Gussoni E, & Pourquié O. 2020. Differentiation of the human PAX7-positive myogenic precursors/satellite cell lineage in vitro. Development **147:** dev187344. DOI: https://doi.org/10.1242/dev.187344

Aulehla A, Pourquie O. 2010. Signaling Gradients during Paraxial Mesoderm Development. *Cold Spring Harb Perspect Biol*. **2:** a000869. DOI: https://doi.org/10.1101/cshperspect.a000869

Baghdadi MB, Castel D, Machado L, Fukada SI, Birk DE, Relaix F, Tajbakhsh S, Mourikis P. 2018. Reciprocal signalling by Notch–Collagen V–CALCR retains muscle stem cells in their niche. Nature **557:**714–718. DOI: https://doi.org/10.1038/s41586-018-0144-9

Barbieri G, DeAngelis L, Feo S, Cossu G, Giallongo A. 1990. Differential expression of muscle-specific enolase in embryonic and fetal myogenic cells during mouse development *Differentiation* **45:**179-184. DOI: https://doi.org/10.1111/j.1432-0436.1990.tb00471.x

Barruet E, Garcia SM, Striedinger K, Wu J, Lee S, Byrnes L, Wong A, Xuefeng S, Tamaki S, Brack AS, Pomerantz JH. 2020. Functionally heterogeneous human satellite cells identified by single cell RNA sequencing. *Elife.* 9:e51576. DOI: https//doi.org/10.7554/eLife.51576.

Bjornson CR, Cheung TH, Liu L, Tripathi PV, Steeper KM, Rando TA. 2012. Notch signaling is necessary to maintain quiescence in adult muscle stem cells. *Stem cells* **30:**232–242. DOI: https://doi.org/10.1002/stem.773

Borchin B, Chen J, Barberi, T. 2013. Derivation and FACS-mediated purification of PAX3+/PAX7+ skeletal muscle precursors from human pluripotent stem cells. *Stem Cell Reports* **1:**620-631. DOI: https://doi.org/10.1016/j.stemcr.2013.10.007

Buckingham M, Rigby PWJ. 2014. Gene Regulatory Networks and Transcriptional Mechanism that Control Myogenesis. *Developmental Cell* **28:**225-238. DOI: https://doi.org/10.1016/j.devcel.2013.12.020

Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, Zhang F, Mundlos S, Christiansen L, Steemers FJ, Trapnell C, Shendure J. 2019. The single-cell transcriptional landscape of mammalian organogenesis. Nature, **566**: 496–502. DOI: https://doi.org/10.1038/s41586-019-0969-x

Chal J, Oginuma M, Al Tanoury Z, Gobert B, Sumara O, Hick A, Bousson F, Zidouni Y, Mursch C, Moncuquet P, Tassy O, Vincent S, Miyanari A, Bera A, Garnier JM, Guevara G, Hestin M, Kennedy L, Hayashi S, Drayton B, Cherrier T, Gayraud-Morel B, Gussoni E, Relaix F, Tajbakhsh S, Pourquié O. 2015. Differentiation of pluripotent stem cells to muscle fiber to model Duchenne muscular dystrophy. *Nat Biotechnol.* **33:**962–969. DOI: https://doi.org/10.1038/nbt.3297

Chargé SB, Rudnicki MA. 2004. Cellular and molecular regulation of muscle regeneration. Physiol. Rev. **84:** 209–238. DOI: https://doi.org/10.1152/physrev.00019.2003

Cheng L, Alvares LE, Ahmed MU, El-Hanfy AS, Dietrich S. 2004. The epaxial-hypaxial subdivision of the avian somite. *Dev. Biol.* **274:**348–369. DOI: https://doi.org/10.1016/j.ydbio.2004.07.020

Cheung TH, Quach NL, Charville GW, Liu L, Park L, Edalati A, Yoo B, Hoang P, Rando TA. 2012. Maintenance of muscle stem-cell quiescence by microRNA-489. Nature **482:** 524–528. DOI: https://doi.org/10.1038/nature10834

Cohn RD, van Erp C, Habashi JP, Soleimani AA, Klein EC, Lisi MT, Gamradt M, ap Rhys CM, Holm TM, Loeys BL, Ramirez F, Judge DP, Ward CW, Dietz HC. 2007. Angiotensin II type 1 receptor blockade attenuates TGF-beta-induced failure of muscle regeneration in multiple myopathic states. *Nat Med*, **13**:204–210. DOI: https://doi.org/10.1038/nm1536

Darabi R, Arpke RW, Irion S, Dimos JT, Grskovic M, Kyba M, Perlingeiro RC. 2012. Human ES- and iPS-derived myogenic progenitors restore DYSTROPHIN and improve contractility upon transplantation in dystrophic mice. Cell Stem Cell, **10:**610–619. DOI: https://doi.org/10.1016/j.stem.2012.02.015

Dietrich S, Abou-Rebyeh F, Brohmann H, Bladt F, Sonnenberg-Riethmacher E, Yamaai T, Lumsden A, Brand-Saberi B, Birchmeier C. 1999. The role of SF/HGF and c-Met in the development of skeletal muscle. Development **126**:1621- 1629.

Dorn I, Klich K, Arauzo-Bravo MJ, Radstaak M, Santourlidis S, Ghanjati F, Radke TF, Psathaki OE, Hargus G, Kramer J, Einhaus M, Kim JB, Kögler G, Wernet P, Schöler HR, Schlenke P, Zaehres H. 2015. Erythroid differentiation of human induced pluripotent stem cells is independent of donor cell type of origin. *Haematologica* **100:** 32-41. DOI: https://doi.org/10.3324/haematol.2014.108068

Faustino Martins JM, Fischer C, Urzi A, Vidal R, Kunz S, Ruffault PL, Kabuss L, Hube I, Gazzerro E, Birchmeier C, Spuler S, Sauer S, Gouti M. 2020. Self-Organizing 3D Human Trunk Neuromuscular Organoids. *Cell Stem Cell* **26:** 172-186. DOI: https://doi.org/10.1016/j.stem.2019.12.007

Fukada S, Uezumi A, Ikemoto M, Masuda S, Segawa M, Tanimura N, Yamamoto H, Miyagoe-Suzuki Y, Takeda S. 2007. Molecular signature of quiescent satellite cells in adult skeletal muscle. *Stem Cells* **25:** 2448-2459. DOI: https://doi.org/10.1634/stemcells.2007-0019

Garcia SM, Tamaki S, Lee S, Wong A, Jose A, Dreux J, Kouklis G, Sbitany H, Seth R, Knott PD, Heaton C, Ryan WR, Kim EA, Hansen SL, Hoffman WY, Pomerantz JH. 2018. High-Yield Purification, Preservation, and Serial Transplantation of Human Satellite Cells. *Stem Cell Reports.* **10**:1160-1174. DOI: https//doi.org/ 10.1016/j.stemcr.2018.01.022.

Gouti M, Delile J, Stamataki D, Wymeersch FJ, Huang Y, Kleinjung J, Wilson V, Briscoe J. 2017. A Gene Regulatory Network Balances Neural and Mesoderm Specification during Vertebrate Trunk Development. Dev. Cell **41**: 243–261. DOI: https://doi.org/10.1016/j.devcel.2017.04.002

Henrique D, Abranches E, Verrier L, Storey KG. 2015. Neuromesodermal progenitors and the making of the spinal cord. Development **142**:2864–2875. DOI: https://doi.org/10.1242/dev.119768

Hicks MR, Hiserodt J, Paras K, Fujiwara W, Eskin A, Jan M, Xi H, Young CS, Evseenko D, Nelson SF, Spencer MJ, Handel BV, Pyle AD. 2018. ERBB3 and NGFR mark a distinct skeletal muscle progenitor cell in human development and hPSCs. *Nat Cell Biol.* **20:**46–57. DOI: https://doi.org/10.1038/s41556-017-0010-2

Jahn K, Mohammadi B, Krampfl K, Abicht A, Lochmüller H, Buffer J. 2001. Deactivation and desensitization of mouse embryonic- and adult- type nicotinic receptor channels currents. *Neuroscience Letters* **307:** 89-92. DOI: https://doi.org/10.1016/s0304-3940(01)01929-2

Juan AH, Derfoul A, Feng X, Ryall JG, Dell'Orso S, Pasut A, Zare H, Simone JM, Rudnicki MA, Sartorelli V. 2011. Polycomb EZH2 controls self-renewal and safeguards the transcriptional identity of skeletal muscle stem cells. Genes & development **25:** 789–794. DOI: https://doi.org/10.1101/gad.2027911

Koehler KR, Nie J, Longworth-Mills E, Liu XP, Lee J, Holt JR, Hashino E. 2017. Generation of inner ear organoids containing functional hair cells from human pluripotent stem cells. *Nat Biotechnol.* **35:**583–589. DOI: https://doi.org/10.1038/nbt.3840

Kramerova I, Kudryashova E, Wu B, Germain S, Vandenborne K, Romain N, Haller RG, Verity MA, Spencer MJ. 2009. Mitochondrial abnormalities, energy deficit and oxidative stress are features of calpain 3 deficiency in skeletal muscle. Human Molecular Genetics **18:**3194–3205. DOI: https://doi.org/10.1093/hmg/ddp257

Lancaster MA, Renner M, Martin CA, Wenzel D, Bicknell LS, Hurles ME, Homfray T, Penninger JM, Jackson AP, Knoblich JA. 2013. Cerebral organoids model human brain development and microcephaly. *Nature* **501:**373– 379. DOI: https://doi.org/10.1038/nature12517

Le Douarin NM, Calloni GW, Dupin E. 2008. The stem cells of the neural crest. *Cell Cycle* **7:**1013-1019. DOI: https://doi.org/10.4161/cc.7.8.5641

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15:**550. DOI: https://doi.org/10.1186/s13059-014-0550-8

Machado L, Esteves de Lima J, Fabre O, Proux C, Legendre R, Szegedi A, Varet H, Ingerslev LR, Barrès R, Relaix F, Mourikis P. 2017. In Situ fixation redefines quiescence and early activation of skeletal muscle stem cells *Cell Rep.* **21:** 1982-1993. DOI: https://doi.org/10.1016/j.celrep.2017.10.080

Maffioletti SM, Sarcar S, Henderson ABH, Mannhardt I, Pinton L, Moyle LA, Steele-Stallard H, Cappellari O, Wells KE, Ferrari G, Mitchell JS, Tyzack GE, Kotiadis VN, Khedr M, Ragazzi M, Wang W, Duchen MR, Patani R, Zammit PS, Wells DJ, Eschenhagen T, Tedesco FS. 2018. Three-dimensional human iPSC-derived artificial skeletal muscles model muscular dystrophies and enable multilineage tissue engineering. *Cell Rep.* **23:**899-908. DOI: https://doi.org/10.1016/j.celrep.2018.03.091

Magli A, Incitti T, Kiley J, Swanson SA, Darabi R, Rinaldi F, Selvaraj S, Yamamoto A, Tolar J, Yuan C, Stewart R, Thomson JA, Perlingeiro R. 2017. PAX7 Targets, CD54, Integrin α9β1, and SDC2, Allow Isolation of Human ESC/iPSC- Derived Myogenic Progenitors. Cell Rep. **19**:2867–2877. DOI: https://doi.org/10.1016/j.celrep.2017.06.005

Marg A, Escobar H, Karaiskos N, Grunwald SA, Metzler E, Kieshauer J, Sauer S, Pasemann D, Malfatti E, Mompoint D, Quijano-Roy S, Boltengagen A, Schneider J, Schülke M, Kunz S, Carlier R, Birchmeier C, Amthor H, Spuler A, Kocks C, Rajewsky N, Spuler S. 2019 Human muscle-derived CLEC14A-positive cells regenerate muscle independent of PAX7. Nat Commun. **10**:5776. DOI: 10.1038/s41467-019-13650-z.

Mavrommatis L, Jeong HW, Gomez-Giro G, Stehling M, Kienitz MC, Psathaki OE, Bixel MG, Morosan-Puopolo G, Gerovska D, Araúzo-Bravo MJ, Schwamborn JC, Schöler HR, Adams RH, Vorgerd M, Brand-Saberi B, Zaehres H. 2020. Human skeletal muscle organoids model fetal myogenesis and sustain uncommitted PAX7 myogenic progenitors. bioRxiv 2020.09.14.295832 DOI: https://doi.org/10.1101/2020.09.14.295832

Messina G, Biressi S, Monteverde S, Magli A, Cassano M, Perani L, Roncaglia E, Tagliafico E, Starnes L, Campbell CE, Grossi M, Goldhamer DJ, Gronostajski RM, Cossu G. 2010. Nfix regulates fetal-specific transcription in developing skeletal muscle. *Cell* **140:**554-566. DOI: https://doi.org/10.1016/j.cell.2010.01.027

Montarras D, Morgan J, Collins C, Relaix F, Zaffran S, Cumano A, Partridge T, Buckingham M. 2005. Direct isolation of satellite cells for skeletal muscle regeneration. *Science* **309**:2064-2067. DOI: https://doi.org/10.1126/science.1114758

Moore R, Walsh FS. 1993. The cell adhesion molecule M-cadherin is specifically expressed in developing and regenerating, but not denervated skeletal muscle. *Development* **117:**1409-1420.

Mourikis P, Sambasivan R, Castel D, Rocheteau P, Bizzarro V, Tajbakhsh S. 2012. A critical requirement for notch signaling in maintenance of the quiescent skeletal muscle stem cell state. *Stem Cells* **30**: 243-252. DOI: https://doi.org/10.1002/stem.775

Murphy M, Reid K, Ford M, Furness JB, Bartlett PF. 1994. FGF2 regulates proliferation of neural crest cells, with subsequent neuronal differentiation regulated by LIF or related factors. Development **120:**3519-3528.

Panopoulos AD, D'Antonio M, Benaglio P, *et.al.* 2017. iPSCORE: A Resource of 222 iPSC Lines Enabling Functional Characterization of Genetic Variation across a Variety of Cell Types. *Stem Cell Reports* **8:**1086-1100. DOI: https://doi.org/10.1016/j.stemcr.2017.03.012

Park IH, Arora N, Huo H, Maherali N, Ahfeldt T, Shimamura A, Lensch MW, Cowan C, Hochedlinger K, Daley GQ. 2008. Disease-specific induced pluripotent stem cells. *Cell* **134:** 877-886. DOI: https://doi.org/10.1016/j.cell.2008.07.041

Porpiglia E, Samusik N, Ho ATV, Cosgrove BD, Mai T, Davis KL, Jager A, Nolan GP, Bendall SC, Fantl WJ, Blau HM. 2017. High-resolution myogenic lineage mapping by single-cell mass cytometry. *Nat Cell Biol.* **19**:558–567. DOI: https://doi.org/10.1038/ncb3507

Quarta M, Brett JO, DiMarco R, De Morree A, Boutet SC, Chacon R, Gibbons MC, Garcia VA, Su J, Shrager JB, Heilshorn S, Rando TA. 2016. An artificial niche preserves the quiescence of muscle stem cells and enhances their therapeutic efficacy. *Nat Biotechnol.* **34**:752-759. DOI: https://doi.org/10.1038/nbt.3576

Raines MA, Magella B, Adam M, Potter SS. 2015. Key pathways regulated by HoxA9,10,11/HoxD9,10,11 during limb development. BMC Developmental Biology **15**:28. DOI: https://doi.org/10.1186/s12861-015-0078-5

Rao L, Qian Y, Khodabukus A, Ribar T, Bursac N. 2018. Engineering human pluripotent stem cells into a functional skeletal muscle tissue. Nat Commun **9:**126. DOI: https://doi.org/10.1038/s41467-017-02636-4

Relaix F, Rocancourt D, Mansouri A, Buckingham M. 2005. A Pax3/Pax7-dependent population of skeletal muscle progenitor cells. *Nature* **435**: 948–953. DOI: https://doi.org/10.1038/nature03594

Roellig D, Tan-Cabugao J, Esaian S, Bronner ME. 2017. Dynamic transcriptional signature and cell fate analysis reveals plasticity of individual neural plate border cells. eLife **6:**e21620. DOI: https://doi.org/10.7554/eLife.21620

Rubenstein AB, Smith GR, Raue U, Begue G, Minchev K, Ruf-Zamojski F, Nair VD, Wang X, Zhou L, Zaslavsky E, Trappe TA, Trappe S, Sealfon SC. 2020. Single-cell transcriptional profiles in human skeletal muscle. *Sci Rep.* **10:**229. DOI: https://doi.org/10.1038/s41598-019-57110-6

Seale P, Sabourin LA, Girgis-Gabardo A, Mansouri A, Gruss P, Rudnick MA. 2000. Pax7 is required for the specification of myogenic satellite cells. *Cell* **102**:777-786. DOI: https://doi.org/10.1016/s0092-8674(00)00066-0

Shao Z, Mellor IR, Brierley JM, Harris J, Usherwood PN. 1998. Potentiation and inhibition of nicotinic acetylcholine receptors by spermine in the TE671 human muscle cell line. *The Journal of Pharmacology and Experimental Therapeutics* **286:**1269-1276.

Shahriyari M, Islam MR, Sakib SM, Rinn M, Rika A, Krüger D, Kaurani L, Gisa V, Winterhoff M, Anandakumar H, Shomroni O, Schmidt M, Salinas G, Unger A, Linke WA, Zschüntzsch J, Schmidt J, Bassel-Duby R, Olson EN, Fischer A, Zimmermann WH, Tiburcy M. Engineered skeletal muscle recapitulates human muscle development, regeneration and dystrophy. J Cachexia Sarcopenia Muscle. 2022 Dec;13(6):3106-3121. doi: 10.1002/jcsm.13094.

Shea KL, Xiang W, LaPorta VS, Licht JD, Keller C, Basson MA, Brack AS. 2010. Sprouty1 regulates reversible quiescence of a self-renewing adult muscle stem cell pool during regeneration. *Cell Stem Cell* **6:**117-129. DOI: https://doi.org/10.1016/j.stem.2009.12.015

Shelton M, Metz J, Liu J, Carpenedo RL, Demers SP, Stanford WL, Skerjanc IS. 2014. Derivation and expansion of PAX7-positive muscle progenitors from human and mouse embryonic stem cells. *Stem Cell Reports* **3:**516–529. DOI: https://doi.org/10.1016/j.stemcr.2014.07.001

Sherwood RI, Christensen JL, Conboy IM, Conboy MJ, Rando TA, Weissman IL, Wagers AJ. 2004. Isolation of adult mouse myogenic progenitors: functional heterogeneity of cells within and engrafting skeletal muscle. *Cell* **119:**543-554. DOI: https://doi.org/10.1016/j.cell.2004.10.021

Shimozono S, Iimura T, Kitaguchi T, Higashijima S, Miyawaki A. 2013. Visualization of an endogenous retinoic acid gradient across embryonic development. Nature **496:**363–366. DOI: https://doi.org/10.1038/nature12037

Shubin N, Tabin C, Carroll S. 1997. Fossils, genes and the evolution of animal limbs. *Nature* **388:**639–648. DOI: https://doi.org/10.1038/41710

Soldatov R, Kaucka M, Kastriti ME, Petersen J, Chontorotzea T, Englmaier L, Akkuratova N, Yang Y, Häring M, Dyachuk V, Bock C, Farlik M, Piacentino ML, Boismoreau F, Hilscher MM, Yokota C, Qian X, Nilsson M, Bronner ME, Croci L, Hsiao WY, Guertin DA, Brunet JF, Consalez GG, Ernfors P, Fried K, Kharchenko PV, Adameyko I. 2019. Spatiotemporal structure of cell fate decisions in murine neural crest. *Science* **364:**eaas9536. DOI: https://doi.org/10.1126/science.aas9536

Southard-Smith EM, Kos L, Pavan WJ. 1998. Sox10 mutation disrupts neural crest development in Dom Hirschsprung mouse model. Nat. Genet. **18:**60-64. DOI: https://doi.org/10.1038/ng0198-60

Spence JR, Mayhew CN, Rankin SA, Kuhar MF, Vallance JE, Tolle K, Hoskins EE, Kalinichenko VV, Wells SI, Zorn AM, Shroyer NF, Wells JM. 2011. Directed differentiation of human pluripotent stem cells into intestinal tissue in vitro. *Nature* **470:**105–109. DOI: https://doi.org/10.1038/nature09691

Stemmer M, Thumberger T, del Sol Keyer M, Wittbrodt J, Mateo JL. 2015. CCTop: an intuitive, flexible and reliable CRISPR/Cas9 target prediction tool. *PLoS One* **10:**e0124633. DOI: https://doi.org/10.1371/journal.pone.0124633

Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, Hao Y, Stoeckius M, Smibert P, Satija R. 2019. Comprehensive Integration of Single-Cell Data. *Cell* **177**:1888-1902. DOI: https://doi.org/10.1016/j.cell.2019.05.031

Sun C, Kannan S, Choi IY, Lim H, Zhang H, Chen GS, Zhang N, Park SH, Serra C, Iyer SR, Lloyd TE, Kwon C, Lovering RM, Lim SB, Andersen P, Wagner KR, Lee G. 2022. Human pluripotent stem cell-derived myogenic progenitors undergo maturation to quiescent satellite cells upon engraftment. Cell Stem Cell. **29**:610-619.e5. DOI: 10.1016/j.stem.2022.03.004.

Tierney MT, Gromova A, Sesillo FB, Sala D, Spenlé C, Orend G, Sacco A. 2016. Autonomous Extracellular Matrix Remodeling Controls a Progressive Adaptation in Muscle Stem Cell Regenerative Capacity during Development. *Cell Rep.* **14**:1940-1952. DOI: https://doi.org/10.1016/j.celrep.2016.01.072

Troy A, Cadwallader AB, Fedorov Y, Tyner K, Tanaka KK, Olwin BB. 2012. Coordination of satellite cell activation and self-renewal by Par-complex-dependent asymmetric activation of p38α/β MAPK. *Cell Stem Cell* **11:** 541-553. DOI: https://doi.org/10.1016/j.stem.2012.05.025

Uezumi A, Fukada S, Yamamoto N, Takeda S, Tsuchida K. 2010. Mesenchymal progenitors distinct from satellite cells contribute to ectopic fat cell formation in skeletal muscle. Nat. Cell Biol. **12:**143-152. DOI: https://doi.org/10.1038/ncb2014

Uezumi A, Ito T, Morikawa D, Shimizu N, Yoneda T, Segawa M, Yamaguchi M, Ogawa R, Matev MM, Miyagoe-Suzuki Y, Takeda S, Tsujikawa K, Tsuchida K, Yamamoto H, Fukada S. 2011. Fibrosis and adipogenesis originate from a common mesenchymal progenitor in skeletal muscle. *J Cell Sci.* 124:3654-3664. DOI: https://doi.org/10.1242/jcs.086629

Verma M, Asakura Y, Murakonda BSR, Pengo T, Latroche C, Chazaud B, McLoon LK, Asakura A. 2018. Muscle satellite cell cross-talk with a vascular niche maintains quiescence via VEGF and Notch signaling. *Cell Stem Cell* **23:**530-543. DOI: https://doi.org/10.1016/j.stem.2018.09.007

Xi H, Fujiwara W, Gonzalez K, Jan M, Liebscher S, Van Handel B, Schenke-Layland K, Pyle, AD. 2017. In Vivo Human Somitogenesis Guides Somite Development from hPSCs. *Cell Rep.* **18:**1573-1585. DOI: https://doi.org/10.1016/j.celrep.2017.01.040

Xi H, Langerman J, Sabri S, Chien P, Young CS, Younesi S, Hicks M, Gonzalez K, Fujiwara W, Marzi J, Liebscher S, Spencer M, Van Handel B, Evseenko D, Schenke-Layland K, Plath K, Pyle AD. 2020. A Human Skeletal Muscle Atlas Identifies the Trajectories of Stem and Progenitor Cells across Development and from Human Pluripotent Stem Cells. Cell Stem Cell, **27**:158–176. DOI: https://doi.org/10.1016/j.stem.2020.04.017

Xu B, Wellik DM. 2011. Axial Hox9 activity establishes the posterior field in the developing forelimb. *Proc Natl Acad Sci U S A*.**108**:4888-4891. DOI: https://doi.org/10.1073/pnas.1018161108

A

Embryoid Bodies

Matrigel Embedding **3D**

Posterior Paraxial Mesoderm Neuromesoderm

Anterior Paraxial Mesoderm Neural Tube induction

Anterior Paraxial Mesoderm Dermomyotome Trunk Neural Crest

Wnt Activation Bmp Inhibition

Wnt Activation Bmp Inhibition bFGF

Wnt Activation Bmp Inhibition bFGF, Retinoic acid

Sonic Hedgehog Bmp Inhibition Wnt1a

bFGF, HGF

HGF

Day   -1   0   2   5   7   11   17   >17

B

Skeletal Muscle Organoid Development

Size (uM)

Day 0   Day 3   Day 5   Day 7   Day 11   Day 17

C

Neuromesoderm / Posterior paraxial mesoderm Day 5

TBX6 SOX2 T DAPI

TBX6 SOX2 T

SOX2 T

D

Anterior paraxial mesoderm

PAX3 SOX2 DAPI   Day 7

PAX3 DAPI   PAX7   Day 11

E

Upregulated Genes

Downregulated Genes

6
3
0
-3
-6

D7   D11   D17   D23

PAX3
MEOX2
UNCX
TBX18
TBX6
HES7

F

Upregulated Genes

Downregulated Genes

10
5
0
-5

D7   D11   D17   D23

PAX7
EN1
SIM1
LBX1
TFAP2A
SOX10

G

Neural Plate Boarder - Specified Neural crest Dermomyotome - Day 17

PAX7 SOX2 DAPI

TFAP2A SOX2 DAPI

SOX10 PAX7 DAPI

PAX3 SOX10 DAPI

PAX3 SOX2 DAPI

PAX3 PAX7 DAPI

PAX3 SOX10

PAX3 SOX2

H

Myogenic Progenitor Migration Day 23

PAX7 SOX2 DAPI

SOX2 DAPI FastMyHC

SOX10 PAX7 DAPI

SOX2 TFAP2A DAPI

Figure 1. Skeletal muscle organoid protocol and correlation to fetal development.
(A) Brightfield images of myogenic development stages, with corresponding cytokines/growth factors protocol applications **(B)** Graph depicting organoid development in size (Day 0, *n*= 51; Day 3, *n*= 46; Day 5, *n*= 48; Day 7, *n*= 50; Day 11, *n*= 50; Day 17, *n*= 33; for each timepoint organoids from 3 independent derivations were measured) **(C)** Representative mesodermal T+, neuromesodermal T+, SOX2+ and paraxial mesodermal TBX6+ organoid expression (in live DAPI-cells) at Day 5 **(D)** Representative organoid PAX3+, mesodermal SOX2- and neural SOX2+ expression at Day 7; PAX3/PAX7 coexpression at Day 11 **(E)** Graph depicting qPCR values for anterior PAX3/UNCX/MEOX2/TBX18 and posterior TBX6/HES7 somitic mesodermal markers **(F)** Graph depicting qPCR values for epaxial and hypaxial dermomyotomal PAX7/EN1/SIM1/ LBX1 and neural crest TFAP2A/SOX10 markers **(G)** Representative organoid neural plate border epithelial PAX3+ / PAX7+ / SOX2+ / TFAP2A+, paraxial mesodermal PAX3+/SOX2- and delaminating specified neural crest progenitor PAX3+ / SOX10+ expression at Day 17 **(H)** Representative organoid myogenic FastMyHC+, PAX7+ and neural SOX2+/TFAP2A+/SOX10+ expression at Day 23. Dashed line represents location of embryoid body embedded into Matrigel. Statistics: Values at each timepoint represent difference in mean relative expression for each gene (D7= Day 5 - Day 7, D11= Day 7 - Day 11, D17= Day 11 - Day 17, D23= Day 17 - Day 23) as derived by performing ordinary one-way ANOVA and Tukey´s multiple comparison tests (E,F). Scale bars: 200µm (G), 100µm (A,D,H), 50µm

A

Fetal / Regenerating Myotubes

Delamination - Migration

HGF   HGF   HGF

Day 18-25   35-42   >60

B

Top Up-regulated Enriched GO Terms

Muscle structure development
Muscle contraction
Muscle system process
Anatomical structure development
Anatomical structure morphogenesis
Developmental process
System development
Animal organ development
Multicellular organism development
Extracellular matrix organization

-log10(Pvalue)

Top Down-regulated Enriched GO Terms

Nervous system development
Neurogenesis
Generation of neurons
Neuron differentiation
Neuron development
Plasma membrane bounded cell projection organization
Cell projection organization
Central nervous system development
Chemical synaptic transmission
Anterograde trans-synaptic signaling

-log10(Pvalue)

C

Migration of Myogenic Progenitors
Neural Plate Boarder and Neural Crest Lineage arrest
Day 35

FastMyHC SOX2 DAPI

PAX7 SOX2 DAPI

D

PAX7 SOX10   MYOD   PAX7 MF20

E

FastMyHC SOX2 DAPI   SOX2 TμJ1 DAPI
Day 98   Day 84

FastMyHC TuJ1 DAPI
Day 84

SOX2 TuJ1 DAPI

F

PAX7

% Cells

Weeks 2 5 8 14

MYOD1

% Cells

Weeks 5 8 14

G

Count

Value

Proximal
Limb bud
HOX 9,10

Distal
HOX 9,10
HOX 11,12,13

HOXA1
HOXB1
HOXD1
HOXA2
HOXB2
HOXA3
HOXB3
HOXD3
HOXA4
HOXB4
HOXC4
HOXD4
HOXA5
HOXB5
HOXC5
HOXA6
HOXB6
HOXC6
HOXA7
HOXB7
HOXB8
HOXC8
HOXD8
HOXA9
HOXB9
HOXC9
HOXD9
HOXA10
HOXC10
HOXD10
HOXA11
HOXC11
HOXD11
HOXC12
HOXD12
HOXA13
HOXB13
HOXC13
HOXD13

Org_16wk Org_16wk Org_8wk Org_8wk Org_4wk Org_4wk Org_4wk

Figure 2. Neural lineage development during skeletal muscle organoid progression.
(A) Stepwise brightfield images depicting delamination / migration of progenitor population during organoid culture progression and corresponding myofiber formation **(B)** Gene ontology enrichment analysis comparing 4 and 8 w organoids attributes muscle identity at 8w post differentiation and highlights muscle system development and neural lineage arrest among the top upregulated and downregulated gene ontology terms, respectively **(C-E)** Organoid overview at Day 35 indicates predominant expression of FastMyHC$^+$ and PAX7$^+$ myogenic populations, while SOX2$^+$ neural populations demarcate SOX2 neural plate border epithelium location as observed at earlier stages (Day 16) **(C)**; PAX7 cells are of myogenic origin (PAX7$^+$/SOX10$^-$), MF20$^+$ myotubes are in their proximity and MYOD1$^+$ cells appear at organoid periphery **(D)**; TUJ1$^+$ neurons are restricted to inner organoid areas and close to SOX2$^+$ epithelium, while FastMyHC$^+$ myofibers occupy exterior organoid areas**(E) (F)** Histographs based on FACS intracellular quantification depicting percentage of PAX7$^+$ or MYOD1$^+$ cells through differentiation protocol. For each replicate we pooled 10 organoids, n=10 Statistics: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001, ns: not significant **(G)** Heatmap of HOX gene cluster emphasizes organoid culture limb axial anatomical identity, by depicting transition from an initial limb bud (HOX 9-10) towards a more distal identity (HOX 11-13) at 8w and 16w post differentiation, respectively. Scale bars, 500μm **(C)**, 200μm **(A, D, E)**.

Figure 3. Skeletal muscle organoid characterization at single cell resolution. (A) t-SNE visualization of color-coded clustering (n= 4.323 cells) at 12w post differentiation highlights predominant presence of skeletal muscle lineage, represented by clusters corresponding to myogenic progenitors (n=1625 cells, 37% of total population) in non-dividing (n=1317 cells) and mitotic (n=308 cells) state, myoblasts (n=731 cells), myocytes (n=1147 cells) and myotubes (n=442). Additionally, mesenchymal and neural lineages are represented by 2 smaller clusters of fibroadipogenic (n=165 cells) and neural (n=213 cells) progenitors, respectively **(B)** Immunocytochemistry at Day 135 indicates derivation of PRDM16$^+$ adipocyte clusters distinct from PAX7$^+$ myogenic progenitors **(C)** Organoid overview at Day 98 depicts expression of PAX7$^+$ myogenic populations and highlights SOX2+ neural populations towards organoid interior **(D)** Positive areas with Oil O Red staining indicate derivation of adipocytes in organoid culture at Day 135 **(E)** Violin plots of key markers PAX7, MYOD1, MYOG, MYH3 from each stage as in 3A depict relative expression levels and emphasize gradual transition from myogenic progenitor to myotube subcluster **(F)** Dot plot showing expression of representative genes related to adipogenesis and fibrogenesis across the 7 main clusters. Circle area represents percentage of gene$^+$ cells in a cluster, color reflects average expression level (gray, low expression; blue, high expression) **(G)** Dot plot showing expression of representative genes related to fetal myogenesis across the 7 main clusters. Circle area represents percentage of gene$^+$ cells in a cluster, color reflects average expression level (gray, low expression; blue, high expression). **(H-J)** Representative organoid overview at Day 98 indicates predominant expression of Fast MyHC fetal myofibers **(H,J),** positive for M-Cadherin and in PAX7$^+$ cells proximity **(I,J) (K)** Representative recording (n=6) of acetylcholine (ACh)-induced changes in holding current in a single skeletal muscle cell. ACh (10 μM) was applied as indicated by the bar. Holding potential -90 mV. Downward deflections represent membrane currents in response to depolarizing voltage ramps (duration 500 ms) from -120 mV to +60 mV. Dashed line indicates zero current level. **(L)** I/V curve of nAChR currents obtained by subtraction of voltage ramp-induced changes of current in presence and absence of ACh (10-5 mol/L), corresponding lower-case letters b - a **(M)** Summarized FRET-recordings from skeletal muscle cells transfected with Twitch2B to monitor the increase in [Ca2+]i during ACh application (h) Scale bars, 1mm (**C,H**), 200μm **(B)**, 100μm **(I,J)**.

Figure 4. Myogenic progenitor identity and comparison to progenitors derived from fetal and adult muscle tissue. (A) t-SNE plot visualization of color-coded clustering indicates myogenic progenitor sub-cluster with distinct molecular signatures: "dormant" PAX7^{high}/ CHODL^{high}/FBN1^{high}, "activated" CD44^{high} / CD98$^+$ / MYOD1$^+$ and "mitotic" KI-67$^+$/CDK1$^+$/TOP2A **(B)** Organoid overviews at Day 98 illustrate CD44 and PAX7 expressing cells at developing sites, which are more accessible to HGF activation signal, specificity of CD44 on MYOD$^+$/PAX7$^+$ progenitor expressing cells (arrows) and absence from FastMyHC$^+$ positive myofibers is highlighted **(C)** FBN1$^+$ microfibrils are located towards organoid interior **(D)** Pseudotime ordering for myogenic progenitors and myoblast corresponding clusters highlights distinct developmental trajectories promoting myogenic commitment and self-renewal **(E)** Correlation coefficient plot for Log$_2$ fold change (Log$_2$ FC) values for isolated myogenic progenitors from human fetal tissue (17w) and FACS sorted CXCR4$^+$ / ITGβ1$^+$ organoid derived myogenic progenitors (16w). PAX7, COL1A1, COL3A1, COL4A1, COL5A1, COL15A1, FBN1 and CHODL and further extracellular matrix related genes are highlighted on the plot. Pearson's correlation coefficient, rho=0.9 for global expression comparison and rho=0.97 for selected genes **(F)** UMAP color-based clustering divides non-dividing myogenic progenitors and adult satellite cells into 4 clusters with distinct molecular signatures: satellite cells characterized by SPRY1^{high} /ZFP36^{high} /MYF5^{high} expression, co-clustered with dormant SPRY1^{high} / FBN1^{high} / CHODL^{high} / PAX7^{high}, activated CD44^{high} / CD98^{high} / MYOD1$^+$ and committed, NEB ^{high}/ MYH3^{high} / MYOD1^{high} organoid derived myogenic progenitors. Dots correspond to adult satellite cells from GSE130646 database, triangles correspond to organoid derived myogenic progenitors **(G)** Violin plots depicting relative expression levels of markers specific for quiescent PAX7 or activated MYF5 muscle stem cell state across adult satellite cells (GSE130646) and organoid derived myogenic progenitors subclusters **(H)** Ridge plots of developmental score distribution of myogenic progenitors across in vivo or in vitro stages based on the difference between up-regulated satellite cell and embryonic markers from human reference atlases for weeks (Wk) 5 to 18 embryonic and fetal stages, years (Yr) 7 to 42 adult satellite cells and SM (skeletal muscle) organoid **(I,J)** In vivo engraftment potential of human myogenic progenitors. 10e5 CD82 sorted human cells were injected into tibialis anterior muscle of nude mice (**I**). Control mice were not injected (**J**). Six weeks post transplantation, cryosections were stained with huLamin A/C (green), Dystrophin (red) and DAPI (blue). Human cells appear green and red in contrast to murine cells which only show a dystrophin positive red staining. Scale bars 200 μm

Figure 1 – figure supplement 1. Lineage representation and organoid culture progression at early stages of differentiation protocol. (A) qPCR analysis depicting relative expression of pluripotent (OCT4, SOX2, NANOG), mesodermal / posterior somitic (BRACHYURY,MESOGENIN,TBX6,HES7), anterior somitic (PAX3,UNCX,TBX18,MEOX2), dermomyotomal (PAX7, SIM1, EN1, LBX1) and neural tube / crest markers (SOX2, PAX6, TFAP2A, SOX10) **(B)** Gating strategies and FACS intracellular quantification of PAX3 and PAX7 markers at Day 10 and 16 **(C)** Tile scan overview at Day 23 showing neural crest SOX10+ populations close to neural plate border epithelium **(D)** and SOX10-/PAX3+ populations at more outer locations (E). For each replicate we pooled 6 organoids, n=6; gene expression normalized to housekeeping gene. Statistics: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001, ns: not significant. FACS plots, yellow: unstained population, red: isotype control, gray: PAX7+ or PAX3+ population. Scale bars, 500uM (C), 200uM (D, E).

A

Color Key
and Histogram

B

Org_8wk vs Org_4wk

Up: 5.919

Down: 5.005

log fold change

Mean of normalized counts

MEOX2
MEOX1
MYOD1
SIM2
TCF4
LBX1
SIM1
MSX1
EYA4
PITX2
SIX1
MYF5
SIX4
PAX7
EYA1
PAX3
CXCR4
EN1
FOXC1
FOXC2

#1 #2 #3   #1 #2   #1 #2
Org_4wk   Org_8wk   Org_16wk

C

Color Key
and Histogram

Neural Tube
Markers

Neural Crest
Markers

HES5
PAX6
FOXB1
OLIG3
TUBB3
SOX1
SOX10
TFAP2A
SOX9
DLX5
PAK3
B3GAT1

#1 #2   #1 #2   #1 #2 #3
Org_8wk   Org_16wk   Org_4wk

D

Delamination / Migration - Myogenic
Progenitors

TFAP2A SOX2 DAPI

Day 35

PAX7 MF20 DAPI

Day 35

E

5 Weeks

FSC-A   FSC-W   PAX7   MYOD1

SSC-A   FSC-H

9 Weeks

FSC-A   FSC-W   PAX7   MYOD1

SSC-A   FSC-H

FSC-A   FSC-W

Rhodamine Red-X

Figure 2 – figure supplement 1. Myogenic versus neural fate during organoid development.
(A) Heatmap of limb myogenic progenitor markers emphasizes upregulation through organoid protocol development (**B**) Differential expression comparison highlights upregulated (n=5.919) and downregulated (n=5.005) genes between 8 vs 4w organoids. (Padj< 0.001) **(C)** Heatmap of neural tube and neural crest markers during 4, 8 and 16 w post differentiation highlights neural lineage arrest from 4 w onwards **(D)** Organoid overview at Day 35 indicates presence of MF20$^+$ myofibers in proximity of PAX7$^+$ cells and delamination migration of mainly myogenic origin since SOX2 and TFAP2A expression is restricted towards inner organoid areas **(E)** Gating strategy applied to quantify MYOD1$^+$ (committed state) and PAX7$^+$ (progenitor state) populations at 5 and 8w post differentiation. Yellow histogram: unstained population, red: isotype control, gray: PAX7$^+$ or MYOD$^+$ population. Scale bars, 500uM

Figure 3 – figure supplement 1. Single cell RNA seq expression profiling and lineage representation in organoid culture at 12 weeks (A) Gene signatures of *t*-SNE described clusters based on relative expression levels of the 50 most significant markers for each of the 7 clusters **(B)** *t*-SNE plot of myogenic and fibroadipogenic markers **(C)** Brightfield image at Day 135 highlights presence of structures reassembling adipocytes **(D)** Immunocytochemistry at Day 135 emphasizes derivation of PRDM16$^+$ cells in FastMyHC$^+$ myofibers proximity **(E)** Organoid overview at Day 98 highlights muscle system development at outer and neural lineage representation at more inner sites **(F)** *t*-SNE plot of key markers PAX7, MYOD1, MYOG, MYH3 from each stage depict relative expression levels and emphasize gradual transition from myogenic progenitor to myotube subcluster **(G)** Heatmap of embryonic or fetal myogenic markers depicts a fetal environment during 8 to16 w post differentiation **(H)** Two-dimensional principal component analysis based on gene expression profiling between samples separates skeletal muscle organoid cluster from pluripotent and highlights greater difference in variance between 4 and 8-16w subgroups in comparison to 8 and 16w subgroup. w.p.d.: weeks post differentiation. Scale bars, 1mM **(D),** 200uM **(E, F)**.

A

Org_16wk vs Org_8wk

Up: 2856

Down: 2369

log fold change

Mean of normalized counts

B

Top Up-regulated Enriched GO Terms

Positive regulation of RNA metabolic process
Positive regulation of nucleic acid-templated transcription
Positive regulation of RNA biosynthetic process
Positive regulation of transcription, DNA templated
Developmental process
Positive regulation of nucleobase containing compound metabolic process
Regulation of ion transport
Positive regulation of gene expression
Positive regulation of transcription by RNA pol. II
Anatomical structure development

-log10(Pvalue)

C

Top Down-regulated Enriched GO Terms

Cellular component organization or biogenesis
Cellular component organization
Organelle organization
DNA packaging
Nucleosome assembly
Chromatin assembly
Cellular component biogenesis
Mitotic nuclear division
Mitotic cell cycle
Mitotic cell cycle process

-log10(Pvalue)

D

Up-regulated Enriched GO Terms

Muscle structure development
Ion transmembrane transport
Cation transport
Regulation of cation transmembrane transport
Muscle system process
Striated muscle cell differentiation
Cellular response to chemical stimulus
Sarcomere organization
Striated muscle contraction
Muscle organ development
Maintenance of synapse structure
Response to stimulus regulating muscle adaptation
Neuromuscular synaptic transmission
Regulation of ryanodine-sensitive calcium-release channel activity

-log10(Pvalue)

E

14 Weeks

SSC-A

Cells 74.2

FSC-A

Single Cells 63.8

FSC-W

PAX7

47.3   52.7

MYOD1

40.4   59.6

Rhodamine Red-X

F

PAX7 TITIN DAPI

Day 98

MYOD DAPI

Day 98

Figure 3 – figure supplement 2. Skeletal muscle organoid culture maturation and identity.
(A) Differential expression comparison highlights upregulated (n=2.856) and downregulated (n=2.369) genes between 16 and 8w organoids (Padj<0.001) **(B-C)** Gene ontology enrichment analysis at 16w depicts top statistically significant upregulated **(B)** and downregulated **(C)** gene ontology (GO) terms; **(D)** statistically significant upregulated GO terms further highlight skeletal muscle maturation **(E)** Gating strategy applied to quantify MYOD1$^+$ (committed state) and PAX7$^+$ (progenitor state) population at 14 w post differentiation **(F)** Organoid overview at Day 98 depicts high proportion of PAX7$^+$ /MYOD$^+$ cells towards outer areas. Yellow histogram: unstained population, Red histogram: Isotype Control, Gray Histogram: PAX7$^+$ or MYOD$^+$ population. Scale bars, 500uM **(F)**

Figure 3 – figure supplemental 3. Functional properties of organoid derived skeletal muscle myofibers. (A) Second Harmonic Generation imaging (SHG) from areas with dense fiber network reveals myofibril formation with distinct sarcomeres (z plane of stack right) **(B)** Electron microscopy depicts myofibrils with well-developed sarcomeres **(C,D)** Skeletal muscle organoid derived myofibers expressed LAMININ and DYSTROPHIN **(E)** Representative recording of acetylcholine (Ach)-induced changes in holding current in a single skeletal muscle cell from organoid derived skeletal muscle cells from iPSC lines with Duchenne (DMD_IPS1, DMD_IPSCORE_65_1) and Becker (BMD_IPS1) muscular dystrophy patients. ACh (10 µM) was applied as indicated by the bar. Holding potential -90 mV. Downward deflections represent membrane currents in response to depolarizing voltage ramps (duration 500 ms) from -120 mV to +60 mV. Dashed line indicates zero current level. **(F)** Summarized FRET-recordings from organoid derived skeletal muscle cells from iPSC lines with Duchenne (DMD_IPS1, DMD_IPSCORE_65_1), Becker (BMD_IPS1) transfected with Twitch2B to monitor the increase in [Ca2+]i during ACh application. Scale Bars, 100uM **(C,D)**, 20uM in **(A)**, 500nM **(B)**

Figure 4 – figure supplement 1. Subclustering of myogenic progenitors and NOTCH signaling. (A) Dot plot showing expression of representative genes related to satellite cell identity across the 7 main clusters. **(B)** Dot plot showing expression of representative genes related to NOTCH signaling are identified as regulators of satellite cell quiescence across the 7 main clusters **(C)** Organoid overview at Day 98 indicates PAX7 myogenic progenitors that are Ki-67-/EdU- **(D)** Gating strategy applied to quantify EdU+ proliferating cells in organoid culture at Day 98 (14 weeks); histograph depicting percentage of EdU+ cells; Dot plot, circle area represents percentage of gene+ cells in a cluster, color reflects average expression level (gray, low expression; blue, high expression). Scale Bars, 100uM in (C).

Figure 4 – figure supplement 2. Pseudotime ordering of myogenic progenitor reveals distinct states and cell fate decisions. (A) Gene signatures of t-SNE described clusters based on relative expression levels of the 50 most significant markers for each of the 3 clusters **(B)** Expression of selected genes along pseudotime. Group of genes selected for myogenic progenitors: PAX7, SPRY1, CHODL (dormant state), CD44, CD98, MYOD1 (activated state), TOP2A (mitotic state), and for myoblasts: MYOD1 **(C, D)** UMAP feature plots depicting relative expression of extracellular matrix proteins COL1A2, , COL5A2, COL5A3, COL6A2, FN1 **(C)** and selected transcriptional regulators that define dormant (PAX7, FBN1, CHODL, SPRY1), activated (CD44, CD98, MYOD1, VEGFA) and mitotic (TOP2A) state of myogenic progenitor and myoblast (MYOD1) clusters **(D)**. **(E,F)** UMAP feature plots depicting relative expression of genes regulating asymmetric divisions and self-renewal (PARD3, p38a/b, CD9, NOTCH3) **(E)** and extracellular matrix collagens (COL1A1, COL3A1, COL4A1, COL5A1, COL6A1, COL6A1, COL15A1; COL4A2, COL6A2 **(F)**

Figure 4 – figure supplement 3. Organoid derived myogenic progenitors and correlation to fetal muscle progenitors. (A) Gating strategy applied for FACS sorting CD29+ / CXCR4+ cells from 15 -16 w skeletal muscle organoids **(B,C)** Re-plating CD29+ / CXCR4+ cells and culturing for 14 days highlights fetal myogenic potential, illustrated by brightfield and immunocytochemistry images for Fast MyHC+, TITIN+ and PAX7+ populations **(D)** t-SNE feature plots for ITGβ1 and CXCR4 demarcate expression of FACS isolated calls into myogenic progenitor subcluster **(E)** Gating strategy from (A) together with unstained population (yellow), isotype control (red) and CD29+ / CXCR4+ (gray) population **(F)** Correlation coefficient plot for Log2 fold change (Log2 FC) values for isolated myogenic progenitors from human fetal tissue (17w) and FACS sorted CXCR4+ / ITGβ1+ organoid derived myogenic progenitors (16w). PAX7, MYF5, MYOD1, MYOG are highlighted on the plot **(G)** Correlation coefficient plot for Log2 fold change (Log2 FC) values for FACS sorted CXCR4+ / ITGβ1+ organoid derived myogenic progenitors (16 w) from CB CD34+, DMD_IPS1, BMD_iPS1 and iPSCORE_65_1 pluripotent lines. Scale bars, 200uM (B), 100uM (C).

Figure 4 – figure supplement 4. Organoid derived myogenic progenitors and correlation to adult human satellite cells. (A) Violin plots highlighting developmental status for organoid derived myogenic progenitors and satellite cells through relative expression of key signature markers ZFP36, CD44, FBN1 and NEB for each subcluster **(B)** Enhanced Volcano plot from comparing transcript levels between all adult satellite cells (GSE130646) and organoid derived myogenic progenitors. Log2 fold-change in normalized gene expression vs. - Log_{10} adjusted p-value is plotted. Differentially expressed genes: blue, adjusted p-value < 0.001; green, adjusted p-value > 0.001 and log2 fold-change > 1;,red, log2 fold-change > 1 and adjusted p value < 0.001; gray, no significance. **(C)** Pseudotime ordering for organoid derived myogenic progenitors and adult satellite cells together with expression of selected genes **(D)** Expression of genes selected for dormant myogenic progenitors (PAX7, NOTCH3, RBPJ) and activated satellite cells (MYF5, JUN, FOS).

Figure 4 – figure supplement 5. **Comparison between 2D in vitro myogenic differentiation protocols and in vivo staging.** Violin plots highlighting expression of key myogenesis markers: Comparison between myogenic progenitors derived in 2D differentiation protocols (HX protocol, Xi et al., 2017; JC protocol, Chal et al., 2015; MS protocol, Shelton et al., 2014) and in vivo myogenic progenitors (5 stages: 1,2: embryonic; 3,4: fetal; 5: postnatal). Data were acquired from Pyle´s LAB UCLA website, (https://aprilpylelab.com/datasets/in-vivo-and-in-vitro-multiple-protocols-smpcs-and-scs/) and Xi et al. (2020).

Gene	Forward Primer Sequence	Reverse Primer Sequence
OCT4	AGTGAGAGGCAACCTGGAGA	ACACTCGGACCACATCCTTC
SOX2	GCCCTGCAGTACAACTCCAT	GACTTGACCACCGAACCCAT
NANOG	CTGAGATGCCTCACACGGAG	GTGGGTTGTTTGCCTTTGGG
PAX6	GTGAGAGTGGACAGACATCCG	CTGTTCTGCATGCTGGCTCT
BRACHYURY	TTCATAGCGGTGACTGCTTATCA	CACCCCCATTGGGAGTACC
MESOGENIN	CAGGATGAGGACCTTGGCAG	GGATCTTGGTGAGAGGCTGG
TBX6	CCCTACTCGGCTGCATTTCT	GAGCCCACATCCAGATAGCC
HES7	GGAACCCGAAGCTGGAGAAA	CGGAAACCGGACAAGTAGCA
UNCX	GGAGAAGGCGTTCAACGAGA	GGAACCAGACCTGAACTCGG
TBX18	CCACCCCGTGTGTACATTCA	TGGCCTTGGTCATCCAGTTC
MEOX2	CTCTGCAAACCAACTGGCAC	AAGAGTTGGAGCACAGGACG
PAX3	AGACTGGCTCCATACGTCCT	CATGCCCGGGTTCTCTCTTT
PAX7	AACCACATCCGCCACAAGAT	CTCCTGGTAGCGGCAAAGAA
EN1	CAGGAACTCAGCCTCAACGA	ACTCGCTCTCGTCTTTGTCC
SIM1	GAGTGGTGTTCCCAGAAGGG	ATCCAGGGTCTGGAGCAGAT
LBX1	GCGGAGAAGTTACTCGCTGT	CCTTAAACGTCTTGCTGGCG
TFAP2A	CCAAGTCCAACAGCAATGCC	CGACCCGGAACTGAACAGAA
SOX10	CCATCCAGGCCCACTACAAG	GCTCTGGCCTGAGGGGT
RPS16 (Housekeeping)	GCTATCCGTCAGTCCATCTCCAA	CCTTCTTGGAAGCCTCATCCAC

Table S1. qPCR primer pairs applied to detect relative expression on key markers during skeletal muscle organoid development.