Mountaintop Removal Mining Threatens the Survival and Recovery of Imperiled Species

Michael J. Evans[1,2] Christian Thomas[3], & Jacob W. Malcom[1,2]

[1]Center for Conservation Innovation
Defenders of Wildlife
1130 17th St. NW
Washington, DC 20036

[2]Environmental Science and Policy Dept.
George Mason University
4400 University Dr
Fairfax, VA 22030

[3]SkyTruth
P.O. Box 3283
Shepherdstown, WV 25443

**Abstract** Environmental laws need sound data to protect species and ecosystems. In 1996, a proliferation of mountaintop removal coal mines in a region home to over 50 federally protected species was approved under the Endangered Species Act. Although this type of mining can degrade terrestrial and aquatic habitats, the available data and tools limited the ability to analyze spatially extensive, aggregate effects of such a program. We used two large, public datasets to evaluate the aggregate effects of mountaintop removal coal mining on water quality over 15 years across Kentucky, Tennessee, Virginia, and West Virginia. We combined an annual map of the extent of surface mines in this region from 1985 to 2015 generated from Landsat satellite imagery with public water quality data collected over the same time period from 4,260 monitoring stations within the same area. We used linear mixed models to estimate the relationship between the proportion of the area draining into a monitoring station that was mined and measures of water quality relevant to standards for sustaining aquatic life. Chronic and acute thresholds for aquatic life were exceeded thousands of times between 1985 and 2015 in streams that were important to the survival and recovery of species on the Endangered Species List. We found that mined area was positively related to increases in conductivity, manganese, sulfate, sulfur, total dissolved solids, total suspended solids, and zinc. The likelihood that chronic thresholds for copper, lead, and zinc were exceeded also increased with mined area. Finally, the proportion of a watershed that was mined was positively related to the likelihood that a waterway would be designated as impaired under the Clean Water Act. Our results demonstrate that mining has downstream effects that must be considered under environmental law. These findings and the public data used in our analyses are pertinent to an upcoming re-evaluation of the effects of current mine permitting regulations to the recovery and survival of federally protected species.

## Introduction

Laws and policies that regulate environmentally damaging activities often require the analysis of the direct, indirect, and cumulative environmental consequences of those actions. In practice the scale of actions and the complexity of the systems they affect can make it challenging to assess all these potential effects. Synergistic and large-scale impacts of human activities are responsible for the continued degradation of natural ecosystems despite substantial global conservation efforts (1,2). For environmental laws to effectively protect natural ecosystems, a complete picture of detrimental effects needs to be analyzed and accounted for during the permitting of potentially damaging actions. Without consideration of these 'downstream' effects, environmental laws and policies will fail in their mission to protect the natural ecosystems upon which biodiversity and human life depend (3). Thus, data and approaches that can accommodate potentially complicated dynamics at large scales are needed for decision makers to come to better choices and meet their legal requirements (4,5).

While many direct, acute impacts of environmentally regulated actions are easy to measure (e.g. habitat loss and fragmentation from forest clearing), the myriad indirect and aggregate effects of human activities may require additional data or new techniques to uncover. For example, surface coal mining directly impacts local ecosystems through land clearing and habitat removal (6) and is regulated in many countries. However, effects to the surrounding ecosystem extend beyond forest clearing and bedrock disruption, as the conversion from interior to exterior forests further increases the area of land impacted by surface mining 1.5-5 times through edge effects (7). Additionally, the compaction of soil and alterations to soil chemistry at former mines are barriers to the successful reforestation of native forests, thus shifting successional patterns and increasing levels of non-native vegetation (8,9). This degradation may

be further underestimated if the disturbance of individual mines is evaluated in isolation without considering the accumulative effects of present, or future mines.

Mountaintop removal mining with valley fill (MTMVF) can have particularly extensive negative effects on environmental quality at large, landscape scales. MTMVF is a coal extraction practice in which coal seams in rugged terrain are accessed by first clearing overlying forest and then using explosives to remove overlying soil and bedrock. The leftover rock is then deposited into headwater stream valleys (10,11). The deposition of soil and bedrock into stream valleys distinguishes MTMVF from other surface and underground coal mining. Because of these impacts to headwater streams, MTMVF affects both terrestrial and aquatic ecosystems, with consequences extending well beyond the immediate footprint of a mine. The conversion from forested land to bare earth decreases water quality, and streams linked to mine sites have higher rates of erosion and nutrient pollution and decreased habitat quality (8,9,11). The release of alkaline mine drainage from weathering of rock and site waste at mine sites elevates conductivity and concentrations of metallic ions that negatively impact aquatic biota (12), leading to decreases in aquatic biodiversity (10,13). Additionally, pollutants can be transferred through food webs to downstream terrestrial ecosystems (14).

The potential for MTMVF to damage ecosystems through these interrelated direct and indirect effects is exacerbated by the massive spatial scale and extent of the practice. In the United States, MTMVF has been particularly prominent across Central Appalachia – a region encompassing the Appalachian Mountains in Kentucky, Tennessee, Virginia, and West Virginia - such that MTMVF is the primary cause of land cover change in the region (15). Over 5,900 km$^2$ of Central Appalachian forest has been cleared for mining or mining related activities (16). This level of landscape modification could pose a substantial threat to biodiversity anywhere,

and in Central Appalachia the impacts of MTMVF are of particular concern because the region is a hot spot for endemic imperiled species (8). Over 50 species on the endangered species list are found in the region. Many of these species are aquatic (e.g., darters, salamanders, and crayfish), making them particularly vulnerable to the negative water quality impacts of MTMVF.

In the United States, two critical environmental laws protecting aquatic habitats and species are the Clean Water Act (CWA) and the Endangered Species Act (ESA). The CWA is the primary federal law protecting the integrity of surface waters. Among other regulations, the CWA requires states to develop water quality standards for waterways to meet different usage criteria, including suitability for aquatic life. If these standards are consistently not met a waterway may be designated as impaired and the state is required to develop limits for allowable loads of pollutants that restrict discharge into the waterway (17). While the CWA primarily regulates water quality, the ESA focuses on the status of species. The ESA requires federal agencies to ensure that their actions do not jeopardize the existence of species listed on the endangered species list (hereafter 'listed species') and do not adversely modify their designated critical habitat. Federal agencies are required to consult with the US Fish and Wildlife Service (hereafter "the Service") or the National Marine Fisheries Service, the agencies that implement the ESA, on the effects of proposed projects and regulations on listed species.

In 1996, the Service consulted on a set of regulations and procedures by which state regulatory authorities could issue mining permits under the Surface Mining Control and Reclamation Act (SMCRA). SMCRA governs the permitting and regulation of surface mines in the United States, requiring a state or federal regulatory authority to identify all threatened or endangered species at a proposed mine site and 'adjacent area', and provide written assurance that the permit will not jeopardize their existence or adversely modify critical habitat (18). In the

Service's judgement these procedures were enough to prevent the jeopardization of all listed species that could potentially be affected by mining activity nationwide. Environmentalists for years have questioned whether the regulations and procedures in SMCRA provide adequate protections to imperiled species from the indirect and accumulative effects of MTMVF in Central Appalachia for two reasons. First, the analysis of effects to listed species required by SMCRA are limited to mine sites and 'adjacent areas,' potentially ignoring degradation of downstream habitats. Second, the aggregate effects of existing and future mining activity were excluded entirely (18). Spatially extensive analyses were considered impractical at the time, but new data and technology now make landscape-scale assessments possible.

In this paper, we use large public datasets and remote sensing analyses to quantify the indirect and aggregate impacts of MTMVF in Central Appalachia. The purpose of this study was to assess the impacts of MTMVF to water quality through the lens of protections to federally protected species required under the ESA. We evaluate whether statistical relationships exist between mountaintop removal mining activities as determined from satellite imagery and downstream water quality as measured at thousands of monitoring stations across the region. If downstream water quality can be predicted from remotely sensed, landscape-scale data of mining impacts, then improved estimates of baseline conditions and predictions of future conditions for imperiled species can be made. To that end, we evaluated the relationships between mined area and:

1. observed values of different water quality measures;
2. the frequency that chronic thresholds for water quality were exceeded; and
3. the frequency of waterway impairment under the Clean Water Act.

We demonstrate that such relationships often exist and link them to the specific laws and policies that provide requirements and mechanisms to ameliorate them. We focus on variables that are known or believed to be important for aquatic species persistence in order to provide insights that can be used to directly improve the conservation prospects for federally protected species.

## Methods

*Mining data*

We obtained spatial data delineating the footprints of all large surface mines across Central Appalachia in each year from 1985 to 2015. These data were generated using Landsat satellite imagery in a previous analysis measuring trends in the extent of mining activities over time (16). In order to avoid commission errors in mine identification affecting subsequent analyses, we cross-referenced these footprints with each year of available data from the National Land Cover Dataset (19) and eliminated all footprints that overlapped with areas flagged as agriculture or development. We used these mine footprints to define our study area as a contiguous selection of all the US counties containing these mines (Fig. 1).

**Figure 1. Chronic exposure thresholds for aquatic life were exceeded thousands of times in Central Appalachian waterways between 1985 and 2015.** Map shows the locations of water quality monitoring stations within the study area encompassing parts of Kentucky, Tennessee, Virginia, and West Virginia. Colors indicate the mean number of different water quality measures for which chronic exposure thresholds were exceeded each time a sample was taken at a given location.

*Imperiled Species Data*

We compiled a list of ESA-listed aquatic species whose range overlapped the study area (Supplemental material) by using the ECOS Data Explorer (https://ecos.fws.gov/ecp/report/adhocDocumentation?catalogId=species&reportId=species).

Potentially important water quality measures were determined by examining federal documents pertaining to these species including listing decisions, recovery plans, and five-year reviews (available at https://ecos.fws.gov/ecp). We generated a list of streams that were important to the survival and recovery of listed aquatic species as those streams identified in these same documents as containing extant populations or as being necessary for recovery. For species with designated critical habitat, we included these waterways as well. We refer to this combined list as streams important to imperiled species.

*Water quality data*

We obtained measurements of water quality from the national water quality data portal (20) using the *dataRetrieval* package (21) for R (22). The national water quality data portal aggregates data from monitoring stations nationally, primarily from the U.S. Geological Survey and the Environmental Protection Agency, but also the National Park Service and other state agencies. We selected a set of water quality measures related to mining activity that can also affect the health of aquatic species (Table 1). Additionally, we collected flow rate and temperature data. Water quality data from 1985 through 2015 were collected - corresponding to the same period over which mining footprint data was available - from all monitoring stations within the counties comprising our study area.

**Table 1. Chronic and acute toxicity thresholds were exceeded many times in waterways important to listed aquatic species.** Table shows the number of times that any recorded value (e.g. 'Dissolved', 'Total', etc.) of each water quality measure exceeded standard thresholds for aquatic life, and the number of different monitoring stations at which these events occurred. These data only consider measures taken from 1,887 monitoring stations whose drainage basin contained a stream that was designated as important to species survival and recovery.

| *Measure* | *Chronic* | *Acute* | *Sites (%)* |
|---|---|---|---|
| Aluminum | 3,853 | 871 | 973 (51.6) |
| Arsenic | 39 | 39 | 831 (44.0) |
| Cadmium | 659 | 629 | 853 (45.2) |
| *Calcium | - | - | - |

| | | | |
|---|---|---|---|
| Conductivity | 2,246 | 2,244 | 226 (12.0) |
| Copper | 1,052 | 750 | 927 (49.1) |
| *Iron | - | - | - |
| Lead | 795 | 271 | 878 (46.5) |
| Manganese | 3,651 | 2,276 | 1,073 (56.9) |
| Mercury | 72 | 62 | 722 (38.3) |
| pH | 15,419 | 15,419 | 1,187 (62.9) |
| Selenium | 113 | 79 | 793 (42.0) |
| *Sulfate | - | - | - |
| *Turbidity | - | - | - |
| *Total dissolve solids | - | - | - |
| *Total suspended solids | - | - | - |
| Zinc | 451 | 451 | 997 (52.8) |

*No chronic or acute thresholds provided for measure*

Water quality data were provided in different units, and we standardized all measures of concentration to ug/L, temperature to degrees Celsius, conductivity to uS/cm, and turbidity to NTU. To account for potentially mis recorded data, we flagged as outliers any observations above the 99.9th percentile for a given measure. This threshold was determined empirically by plotting the number of observations falling outside successively larger quantiles and selecting the quantile at which observations appeared to plateau.

To identify acute and chronic toxicity thresholds for different water quality measures, we used the state water quality standards for aquatic life administered by Virginia under the Clean Water Act. These standards are approved by the Environmental Protection Agency and are used to determine waterway impairment requiring mitigation. The water quality standards thus represent an agreed upon set of thresholds necessary to maintain suitability of waterways for aquatic species. The Virginia standards were identical to those from Kentucky, Tennessee and West Virginia. For each measure we flagged any observation where recorded levels exceeded either the acute or chronic exposure thresholds. Finally, we also obtained the locations of all

waterways declared as 'impaired' under section 303(d) of the Clean Water Act occurring within our study area from the EPA's Environmental Dataset Gateway (23). Impaired waterway designations were made on biannual cycles beginning in 1991.

*Spatial analyses*

To assess the relationship between mining and water quality measures recorded at monitoring stations, we associated mines and monitoring locations based on hydrography. We used the watershed and flow modeling tools from the *pyshed* package (24) for Python to delineate the geographic areas that drained into each monitoring station. These analyses required a model of surface elevation, and we used a 30m digital elevation model provided by NASA (25) clipped to our study area. Once drainage basins were delineated for each monitoring station, we calculated the proportion of each basin covered by surface mines in each year from 1985 to 2015. We also created a many-to-one spatial join identifying which mines fell within the drainage basin of each monitoring station.

We used the annual Cropland Data Layers (26) and NLCD data to estimate the proportion of drainage basins that were covered by agriculture or impervious surface in each year for which mining footprint data was available (1985 – 2015). Cropland data were available annually beginning in 2009, and NLCD impervious surface data were available from 1992, 2001, 2004, 2006, 2008, 2011 and 2013. We interpolated between and extrapolated beyond these observed data points to estimate agricultural and impervious surface data for missing years between 1985 and 2015.

In analyses pertaining to 303(d) impaired waterways we consider mined area within watersheds containing the waterway. We used U.S. Geological Survey HUC12 hydrologic units (27) to represent watersheds. We then repeated the above interpolation and extrapolation

procedure to obtain the area mined and covered by agriculture or impervious surface within a contiguous selection of watersheds overlapping the mine footprint data set.

Unless otherwise indicated, all spatial analyses were performed using the Google Earth Engine python API (28).

*Statistical analyses*

We tallied the frequency with which water quality standards were exceeded in streams important to imperiled species by spatially joining the locations of monitoring stations to linear stream features with attributes indicating whether the stream was important to imperiled species.

In all analyses estimating the relationship between mined area and water quality measures, we attempted to account for attenuation in pollutant concentrations with increasing distance between monitoring stations and mines. We adjusted the area of each mine footprint within a given drainage basin in proportion to the square root of the distance from the mine to the corresponding monitoring station. We refer to this measure as adjusted mined area.

Our first objective was to quantify the relationship between values of each water quality measure and the proportion of drainage basins that were mined each year, while controlling for agriculture and impervious surface in the drainage basin. To do so, we created linear mixed effects models with normal error distributions and random intercepts per year nested within monitoring sites. These models were used to estimate the increase or decrease in mean water quality measures as a function of adjusted mined area, percent agriculture, and percent impervious surface within drainage basins.

Our second objective was to determine whether the proportion of a drainage basin that was mined each year affected the probability that pollutant levels would exceed thresholds deemed safe for aquatic life. We specified a generalized linear mixed model using a binomial

error distribution and logit link, with random intercepts per year nested within monitoring sites. These models were used to predict the probability that an observed value for a given measure would exceed chronic thresholds as a function of adjusted mined area, percent agriculture, and percent impervious surface within drainage basins.

Our last objective was to estimate the relationship between the proportion of a watershed that was mined and the probability that a waterway therein would be designated as impaired based on the standards for aquatic life under the Clean Water Act. We specified a generalized linear mixed model with a logit link and binomial error distribution with random intercepts per year nested within monitoring stations. These models were used to predict the probability that a watershed would contain an impaired waterway at the end of a given biannual evaluation cycle as a function of adjusted mined area, percent agriculture, and percent impervious. Because impaired waterways are only tallied during biannual cycles, we used the maximum percent agriculture and impervious surface within two-year cycles as predictor variables. To account for lagged and accumulative effects, we used the cumulative sum of percent mined area within drainage basins over time.

In all regression analyses we included only sites with at least 10 observations. We estimated model parameters in a Bayesian framework using the *rstanarm* (29) package for R. For each model we generated four MCMC chains and tested for convergence using the Rhat statistic. A significant relationship was determined between mined area and response variables if the 95% credible interval around the relevant parameter estimate did not overlap zero.

## Results

We obtained water quality data from 4,260 different water quality monitoring sites across our study area. Distances between stations and mines were exponentially distributed, ranging

from 0 to 343 km (x = 77.7 km, $\sigma^2$ = 4071 km). The number of observations at each site ranged

from 1 to 275. There were 569 sites with at least 10 observations and drainage basin areas greater

than 400 $km^2$ that were included in modeling analyses.

As of 2018, 55 listed aquatic species potentially occurred within the counties comprising

our study area. These included 39 mollusk, 12 fish, 3 crustacean, and 1 snail species. Of these

listed species, 16 had designated critical habitat. Additionally, for 50 of these species we were

able to identify specific streams that were important to the species survival and recovery in either

critical habitat designations, listing decisions, five-year reviews, or recovery plans. Of the 4,260

monitoring stations, 2,881 of these drained areas containing important streams. Chronic and

acute toxicity thresholds for aquatic life were exceeded thousands of times at these monitoring

stations (Table 1). The most frequently exceeded threshold was that for pH, followed by

manganese, aluminum, and conductivity (Table 1). 54 streams designated as critical habitat were

sampled directly by 209 monitoring stations. Water quality thresholds were exceeded at least

once at each of those monitoring stations a total of 5,592 times with a maximum of 272 at a

single station.

Linear mixed models indicated significant positive relationships between the proportion

of a drainage basin that was mined and measured levels of conductivity, manganese, sulfate,

sulfur, total dissolved solids, total suspended solids, and zinc (Fig. 2). No measures were

significantly negatively associated with adjusted mined area (Table 2).

**Figure 2. Increases in the proportion of drainage basins that were mined lead to increases in multiple measures of water quality that are detrimental to aquatic species.** Graphs show the change in water quality measures per change in mined area as estimated by linear mixed models. Dashed lines encompass a 95% credible interval around estimated relationships.

**Table 2. Significant positive relationships were estimated between mined area and nine measures of water quality.** Table shows the regression coefficients quantifying the relationship between adjusted mined area on measures, as estimated by linear mixed models. The 50th, 2.5th,

and 97.5th percentile of the posterior distribution, as well as a measure of MCMC chain convergence (Rhat) are included. Bold text indicates estimates with 95% credible intervals that did not overlap zero.

| Toxin | Measure | Median | 2.5% | 97.5% | Rhat |
|---|---|---|---|---|---|
| Aluminum | Total | 3.57E+03 | -1.01E+03 | 8.17E+03 | 1.008177 |
| Aluminum | Dissolved | 3.74E+03 | -9.82E+02 | 8.52E+03 | 1.006467 |
| Arsenic | Total | -5.03E+01 | -1.38E+02 | 3.52E+01 | 1.003121 |
| Arsenic | Dissolved | 3.39E-01 | -5.26E+00 | 5.70E+00 | 1.003891 |
| Cadmium | Total | -8.21E+02 | -2.08E+03 | 4.52E+02 | 1.008137 |
| Cadmium | Dissolved | -6.92E+00 | -4.39E+01 | 3.11E+01 | 0.999856 |
| Calcium | Total | 6.34E+04 | -4.87E+02 | 1.25E+05 | 1.008877 |
| Calcium | Dissolved | 8.04E+04 | -3.35E+04 | 1.92E+05 | 1.001664 |
| **Conductivity** | **Total** | **3.30E+06** | **1.96E+06** | **4.67E+06** | **1.000473** |
| Copper | Total | -1.20E+02 | -4.52E+02 | 2.03E+02 | 1.000769 |
| Copper | Dissolved | -1.26E+02 | -4.67E+02 | 2.04E+02 | 1.001732 |
| Iron | Total | -2.04E+03 | -2.23E+04 | 1.87E+04 | 0.999076 |
| Iron | Dissolved | -1.44E+04 | -3.48E+04 | 5.94E+03 | 0.999708 |
| Lead | Total | -5.12E+01 | -2.32E+02 | 1.28E+02 | 1.000115 |
| Lead | Dissolved | -5.32E+01 | -2.27E+02 | 1.16E+02 | 1.000174 |
| **Manganese** | **Total** | **1.24E+04** | **3.25E+02** | **2.40E+04** | **1.009628** |
| **Manganese** | **Dissolved** | **1.24E+04** | **3.25E+02** | **2.40E+04** | **1.009628** |
| Mercury | Total | -5.52E+00 | -3.42E+01 | 2.33E+01 | 1.003634 |
| Mercury | Dissolved | -5.77E-01 | -2.82E+00 | 1.76E+00 | 1.005743 |
| pH | Dissolved | 2.68E-01 | -3.37E-01 | 9.02E-01 | 1.000024 |
| Selenium | Total | -1.78E+02 | -4.01E+02 | 4.65E+01 | 1.000092 |
| Selenium | Dissolved | -1.76E+02 | -4.10E+02 | 5.36E+01 | 1.000355 |
| **Sulfate** | **Dissolved** | **6.08E+05** | **1.98E+05** | **1.01E+06** | **1.000307** |
| **Sulfur** | **Total** | **6.58E+05** | **3.58E+05** | **9.45E+05** | **1.000377** |
| **Total dissolved solids** | **Total** | **6.87E+05** | **2.25E+05** | **1.14E+06** | **1.000442** |
| Total dissolved solids | Dissolved | 7.74E+05 | -3.56E+04 | 1.55E+06 | 1.000669 |
| **Total suspended solids** | **Total** | **3.01E+05** | **1.96E+05** | **4.02E+05** | **1.002787** |
| Turbidity | Total | -7.06E+02 | -2.79E+03 | 1.33E+03 | 0.999759 |
| **Zinc** | **Total** | **6.73E+01** | **3.19E+01** | **1.03E+02** | **1.006411** |
| **Zinc** | **Dissolved** | **6.85E+01** | **3.32E+01** | **1.04E+02** | **1.005122** |

We also found significant positive relationships between the cumulative proportion of a watershed that was mined over time and the log odds that a stream in that watershed would be designated as impaired for aquatic life under the Clean Water Act (Table 3).

**Table 3. Significant positive relationships were estimated between mined area in a watershed and the probability that streams therein were designated as impaired.** Table shows the 50th, 2.5th, and 97.5th percentiles of posterior distributions of regression coefficients quantifying the relationship between and probability of impairment, as estimated by logistic

regression models with random effects. Measures of MCMC chain convergence (Rhat) are
included

| Max percentage | Median | 2.5% | 97.5% | Rhat |
|---|---|---|---|---|
| Mined | 23.04 | 20.34 | 25.88 | 1.00 |
| Impervious | 97,232.78 | 74,617.36 | 118,089.30 | 1.02 |
| Cultivated | -36,071.50 | -40,454.80 | -32,007.00 | 1.00 |

Measures of conductivity all exceeded chronic exposure thresholds, and we were unable

to model the probability of exceeding thresholds as a function of land cover predictors. Models

indicated that the probabilities that chronic exposure thresholds for copper, lead, and manganese

were exceeded were all positively related to the area mined within drainage basins (Fig. 3). The

probabilities for no water quality measures exhibited a significant negative relationship.

**Figure 3. Increases in the proportion of drainage basins that were mined increased the
probability that chronic exposure toxicity thresholds would be exceeded for three water
quality measures**. Graphs show the change in probability of exceedance per change in mined
area as estimated by linear mixed models. Dashed lines encompass the 95% credible interval
around estimated relationships.

## Discussion

The scale of the human enterprise is large and growing (3). To minimize the

environmental impacts of human activities, analytical approaches that can address the full scale

and complexity of their effects are required. Many laws and regulations are theoretically capable

of handling such large-scale effects, but too often the data and science needed to fully understand

the effects of our actions are limited (4). This is arguably the case with mountaintop mining with

valley fill (MTMVF) in Central Appalachia, where dated and spatially limited analyses led to the

blanket approval of thousands of MTMVF mines decades ago. Here we combine two classes of

large, public datasets to demonstrate that surface mining is associated with degraded water

quality at landscape scales in ways that affect the survival and recovery of federally protected

species. We found consistent evidence linking changes in mined area with increases in

concentrations of toxics, conductivity, and dissolved and suspended solids. Far from being innocuous side effects, the measures with the strongest relationships to mining were among those that directly affect the survival of aquatic species (12). These findings demand regulatory action under federal environmental laws including the Endangered Species Act and Clean Water Act, because these activities are concentrated in an area with high numbers of imperiled species.

The positive relationships between the extent of mined areas and degradation in water quality that we identified were not surprising. A large body of previous ecological and hydrologic research has shown that surface mining can negatively impact water quality and reduce the suitability of streams for aquatic species at local and regional scales (6,10,11,13,14). Consistent with this research, we found substantial increases in stream conductivity, and the concentrations of manganese, sulfate, sulfur, zinc, and dissolved and suspended solids as a result of increases in the proportion of upstream areas that were mined. Our results build on previous work by illustrating that MTMVF not only degrades water quality immediately proximate to mines, but at a landscape scale - establishing a direct relationship between the aggregate area mined on the landscape and degradation in water quality and habitat suitability for aquatic species. In the context of environmental regulations governing the permitting and operation of MTMVF mines, these direct and aggregate effects illustrate the importance of considering a spatially extensive set of conditions when evaluating the environmental baseline and potential impacts to protected ecosystems and species.

Accounting for the aggregate effects of mine permitting at landscape scale is particularly important in the context of conserving species threatened with or on the brink of extinction, such as those on the endangered species list. We found that degradation in water quality progressed to potentially lethal levels for aquatic life - documenting thousands of instances in which chronic

and acute water quality thresholds for aquatic life were exceeded between 1985 and 2015. Our analyses demonstrate a positive relationship between the probability that chronic thresholds were exceeded and the amount of upstream area that was mined. Measured levels of dissolved copper, lead, and manganese were more likely to exceed safe thresholds for aquatic life as the proportion of upstream area that was mined increased. Together, these dynamics likely contributed to the emergent outcome we observed that a waterway was more likely to fail to meet water quality standards and be declared impaired under the Clean Water Act as the proportion of its watershed that was mined increased.

Mining activities thus directly impacted listed species, based on the observation that water quality thresholds were exceeded thousands of times at monitoring stations on streams designated as critical habitat and stations draining streams identified by the US Fish and Wildlife Service as important to listed species survival and recovery. For instance, monitoring station 211WVOWR-KE-000-004.50 collected water from a section of the Elk river designated as critical habitat for the diamond darter (*Crystallaria cincotta*). Out of 48 occasions, chronic and acute thresholds for aluminum were exceeded 33 and 8 times, respectively; manganese 14 and 5 times; copper 2 and 2 times; and cadmium once. The repeated exceedance of acute toxicity thresholds for aquatic life within designated critical habitat could constitute an adverse modification of critical habitat – an outcome federal agencies must legally avoid under the ESA. The regular and repeated exceedance of water quality thresholds for aquatic life in both critical habitat, and waterways important to the survival and recovery of listed species would seem to be in conflict with the purpose of the ESA; protecting and recovering imperiled species.

These results are highly relevant at a time when government agencies are revisiting surface mining authorizations. The Office of Surface Mining Regulation and Enforcement has

agreed to reinitiate consultation on the effects of SMCRA on imperiled species under section 7

of the ESA (30). As part of this process, the Service is required to critically evaluate both the

environmental baseline of conditions for threatened and endangered species in all areas where

surface mining is covered by SMCRA, and the effects of continued permitting of surface mines

by state and federal regulatory authorities (31). Because they are required to use best available

scientific and commercial data in their analyses (32), the Service will need to use the public

satellite and water quality data provided by other government agencies used here. These data

address one of the fundamental limitations of the original biological opinion by enabling a

quantitative analysis of the aggregate effects of the permitting program. This means that the

environmental baseline and the effects of possible future mines authorized under SMCRA can be

based on a solid, data-informed approach that is scientifically and legally defensible. As our

results demonstrate, these data strongly indicate that the continued operation and approval of

MTMVF mines in Central Appalachia impair the survival and recovery of listed species.

The re-initiation of consultation on SMCRA is relevant not only to the permitting of new

mines, but to regulations surrounding the reclamation of abandoned mines. The use of coal for

energy, and hence its production, has been declining in the United States and is expected to

continue to decline (33). While this means a decrease in new permits, it also means an increase

in mine abandonment and reclamation. In conjunction with re-initiation of the SMCRA

consultation, this economic trend creates an opportunity for regulatory agencies to require best

practices for mine reclamation that protect and recover imperiled species. The destructive nature

of MTMVF is a major barrier to the successful reestablishment of native Appalachian forests

(15), and reclamation efforts have historically emphasized restoring vegetation to mined sites

without prioritizing native species (34,35). In the context of imperiled species conservation, even

efforts that successfully re-establish native communities often exclude rare species (36).

Recently however, reclamation efforts have shifted to focus on reestablishing native forests on

mined lands. The Forestry Reclamation Approach provides management practices which allow

operators to meet the reclamation requirements of SMCRA, while also restoring native forests

(37). New requirements should meet these standards to ensure native ecosystems are restored on

reclaimed mines.

Of course, water quality degradation and waterway impairment are not solely attributable

to surface mining, and more than half of US waterways do not meet CWA water quality

standards. In addition to point source pollution, many land use factors can contribute to the

degradation of aquatic conditions, including the prevalence of impervious surfaces and

agriculture within watersheds (38,39). These pathways were reflected in our results, which show

positive relationships between impervious surface and agricultural extent within watersheds and

the likelihood that an encompassed waterway would be impaired. In recent years, however,

energy has been the largest driver of land use change in the United States (40), and policies that

minimize the negative environmental impacts of energy exploration, extraction, and production

thus have the potential to make a significant improvements to the conservation prospects of

biodiversity.

Our results demonstrate that even after accounting for additional sources of water quality

degradation, surface mining contributed to further increases in the concentration of toxics that

can impair aquatic biota in Central Appalachia. These findings indicate that, in situ, the growth,

continued operation, and legacy effects of MTMVF in the region likely directly limit the

prospects for survival and recovery of over 50 federally protected species. Thus, the surface mine

permitting program implemented under SMCRA and approved under the ESA is directly

counterproductive to the Act's purpose – protecting and restoring imperiled species. Fortunately, there are policy opportunities to ameliorate these effects. Designating critical habitat for the 39 listed aquatic species that occur in Central Appalachia that lack such designated habitat would add legal protections for hundreds of kilometers of waterways subject to degradation from mining. The re-initiated ESA consultation on SMCRA can use the same publicly available water quality and remote sensing data to analyze the effects of mine operation beyond the adjacent area, and the aggregate effects of a permitting program. While approved practices for assessing and permitting the operation of an individual mine may successfully mitigate impacts to immediately surrounding ecosystems, without a landscape scale assessment it is impossible to assess the net impact of a programmatic permitting program.

## Acknowledgements

We thank Kat Diersen and Robert Dreher for review and revisions of the manuscript. Kathryn Kay and Chelsea Proctor assisted with water quality data cleaning and preparation and reviewed federal documents to collect endangered species data. Finally, we thank the team at the Center for Conservation Innovation for providing continuous suggestions and input on the goals and methods of the analysis.

## References

1. Di Marco M, Venter O, Possingham HP, Watson JEM. Changes in human footprint drive changes in species extinction risk. Nat Commun. 2018 Dec 5;9(1):4621.
2. Dempsey J, Martin TG, Sumaila UR. Subsidizing extinction? Conserv Lett. 2020 Jan 24;(13):e12705.
3. IPBES. Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Diaz S, Settele J, Brondizio ES, Ngo HT, Gueze M, Agard J, et al., editors. Vol. 45. 2019. 56 p.
4. Stephenson PJ, Bowles-Newark N, Stanwell-Smith D, Diagana M, Höft R, Abarchi H, et al. Unblocking the flow of biodiversity data for decision-making in Africa. Biol Conserv. 2017 Sep 1;213:335–40.
5. Evans MJ, Malcom JW, Li YW. Novel data show expert wildlife agencies are important to endangered species protection. Nat Commun. 2019 Dec 1;10(1):1–9.
6. Palmer MA, Bernhardt ES, Schlesinger WH, Eshleman KN, Foufoula-Georgiou E,

Hendryx MS, et al. Mountaintop mining consequences. Science (80- ). 2010;327(5962):148–9.

7. Wickham JD, Riitters KH, Wade TG, Coan M, Homer C. The effect of Appalachian mountaintop mining on interior forest. Landsc Ecol. 2007;22(2):179–87.

8. Maigret TA, Cox JJ, Yang J. Persistent geophysical effects of mining threaten ridgetop biota of Appalachian forests. Front Ecol Environ. 2019;17(2):85–91.

9. Wickham J, Wood PB, Nicholson MC, Jenkins W, Druckenbrod D, Suter GW, et al. The overlooked terrestrial impacts of mountaintop mining. Bioscience. 2013;63(5):335–48.

10. Bernhardt ES, Lutz BD, King RS, Fay JP, Carter CE, Helton AM, et al. How many mountains can we mine? Assessing the regional degradation of central appalachian rivers by surface coal mining. Environ Sci Technol. 2012;46(15):8115–22.

11. Ross MRV, McGlynn BL, Bernhardt ES. Deep Impact: Effects of mountaintop mining on surface topography, bedrock structure, and downstream waters. Environ Sci Technol. 2016 Feb 16;50(4):2064–74.

12. Hartman KJ, Kaller MD, Howell JW, Sweka JA. How much do valley fills influence headwater streams? Hydrobiologia. 2005;532(1):91–102.

13. Pond GJ, Passmore ME, Borsuk FA, Reynolds L, Rose CJ. Downstream effects of mountaintop coal mining: Comparing biological conditions using family- and genus-level macroinvertebrate bioassessment tools. J North Am Benthol Soc. 2008;27(3):717–37.

14. Naslund LC, Gerson JR, Brooks AC, Walters DM, Bernhardt ES. Contaminant subsidies to riparian food webs in Appalachian streams impacted by mountaintop removal coal mining. Environ Sci Technol. 2020;54:3951–9.

15. Townsend PA, Helmers DP, Kingdon CC, McNeil BE, de Beurs KM, Eshleman KN. Changes in the extent of surface mining and reclamation in the Central Appalachians detected using a 1976-2006 Landsat time series. Remote Sens Environ. 2009;113(1):62–72.

16. Pericak AA, Thomas CJ, Kroodsma DA, Wasson MF, Ross MRV, Clinton NE, et al. Mapping the yearly extent of surface coal mining in central appalachia using landsat and google earth engine. PLoS One. 2018;13(7):1–15.

17. U.S. Congress. Clean Water Act of 1977. United States of America; 1977.

18. U.S. Fish and Wildlife Service. Formal Section 7 Biological Opinion and Conference Report on Surface Coal Mining and Reclamation Operations Under the Surface Mining Control and Reclamation Act of 1977. Vol. 26. 1996.

19. Fry JA, Xian G, Jin S, Dewitz JA, Homer CG, Yang L, et al. Completion of the 2006 national land cover database for the conterminous united states. Photogramm Eng Remote Sensing. 2011;77(9):858–64.

20. Read EK, Carr L, De Cicco L, Dugan HA, Hanson PC, Hart JA, et al. Water quality data for national-scale aquatic research: The Water Quality Portal. Water Resour Res. 2017;53:1735–45.

21. De Cicco LA, Lorenz D, Hirsch R, Watkins W. dataRetrieval: R packages for discovering and retrieving water data available from U.S. federal hydrologic web services. Reston, VA: U.S. Geological Survey; 2018.

22. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2018.

23. U.S. Environmental Protection Agency. Environmental Dataset Gateway [Internet]. [cited 2017 Jan 5]. Available from: https://edg.epa.gov/metadata/catalog/main/home.page

24.    Bartos M. Pysheds. 2020.

25.    Farr TG, Rosen PA, Caro E, Crippen R, Duren R, Hensley S, et al. The Shuttle Radar Topography Mission. Rev Geophys. 2007;45(2):RG2004.

26.    Boryan C, Yang Z, Mueller R, Craig M. Monitoring US agriculture: the US Department of Agriculture, National Agricultural Statistics Service, Cropland Data Layer Program. Geocarto Int. 2011 Aug;26(5):341–58.

27.    U.S. Geological Survey. Watershed boundary dataset for HUC12 [Internet]. 2017 [cited 2019 Nov 12]. Available from: http://datagateway.nrcs.usda.gov

28.    Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R. Google Earth Engine: Planetary-scale geospatial analysis for everyone. Remote Sens Environ. 2017 Dec 1;202:18–27.

29.    Goodrich B, Gabry J, Ali I, Brilleman S. rstanarm: Bayesian applied regression modeling via Stan. 2018.

30.    D.D.C. National Parks Conservation Association v. S.M.R. Jewell. Vol. 62 F. 2014. p. 7.

31.    U.S. Fish and Wildlife Service and National Marine Fisheries Service. Consultation Handbook: Procedures for Conducting Consultation and Conference Activities Under Section 7 of the Endangered Species Act. 1998;(March):xxii-9-8 +appendices.

32.    U.S. Congress. The Endangered Species Act Amendments of 1978. Federal Register, 1531 1978.

33.    Energy Information Administration. Annual Coal Report 2018. Washington, DC; 2019.

34.    Oliphant AJ, Wynne RH, Zipper CE, Ford WM, Donovan PF, Li J. Autumn olive (Elaeagnus umbellata) presence and proliferation on former surface coal mines in Eastern USA. Biol Invasions. 2017;19(1):179–95.

35.    Zipper CE, Burger JA, Skousen JG, Angel PN, Barton CD, Davis V, et al. Restoring forests and associated ecosystem services on Appalachian coal surface mines. Environ Manage. 2011;47(5):751–65.

36.    Holl KD. Long-term vegetation recovery on reclaimed coal surface mines in the eastern USA. J Appl Ecol. 2002;39(6):960–70.

37.    Adams MB, Angel P, Barton C, Burger J, Davis J, French M, et al. The Forestry Reclamation Approach: Guide to Successful Reforestation of Mined Lands. Gen Tech Rep NRS-169. 2017;(May):119.

38.    Brabec E, Schulte S, Richards PL. Impervious Surfaces and Water Quality: A Review of Current Literature and Its Implications for Watershed Planning. J Plan Lit. 2002 May 6;16(4):499–514.

39.    Verhoeven JTA, Arheimer B, Yin C, Hefting MM. Regional and global concerns over wetlands and water quality. Trends Ecol Evol. 2006;21(2):96–103.

40.    Trainor AM, McDonald RI, Fargione J. Energy sprawl is the largest driver of land use change in United States. PLoS One. 2016;11(9):1–16.

Figure 1

Figure 2

Figure 3