Functional assessment of the "two-hit" model for neurodevelopmental defects in *Drosophila* and *X. laevis*

Short title: Assessment of "two-hit" model for developmental defects

Lucilla Pizzo[1]*, Micaela Lasser[2]*, Tanzeen Yusuff[1], Matthew Jensen[1], Phoebe Ingraham[1], Emily Huber[1], Mayanglambam Dhruba Singh[1], Connor Monahan[2], Janani Iyer[1], Inshya Desai[1], Siddharth Karthikeyan[1], Dagny J. Gould[1], Sneha Yennawar[1], Alexis T Weiner[1], Vijay Kumar Pounraja[1], Arjun Krishnan[4,5], Melissa Rolls[1], Laura Anne Lowery[6] and Santhosh Girirajan[1,3#]

1. Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University Park, PA USA
2. Department of Biology, Boston College, Chestnut Hill, MA USA
3. Department of Anthropology, The Pennsylvania State University, University Park, PA USA
4. Department of Computational Mathematics, Science and Engineering, Michigan State, University, East Lansing, MI, USA
5. Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI, USA
6. Department of Medicine, Boston University Medical Center, Boston, MA, USA

*contributed equally to the work

#Corresponding author:
Santhosh Girirajan, MBBS, PhD
205A Life Sciences Building
Pennsylvania State University
University Park, PA 16802
E-mail: sxg47@psu.edu
Phone: 814-865-0674

Keywords: phenotypic variability, modifier, CNV

Abstract

We previously identified a deletion on chromosome 16p12.1 that is mostly inherited and

associated with multiple neurodevelopmental outcomes, where severely affected probands

carried an excess of rare pathogenic variants compared to mildly affected carrier parents. We

hypothesized that the 16p12.1 deletion sensitizes the genome for disease, while "second-hits" in

the genetic background modulate the phenotypic trajectory. To test this model, we examined

how neurodevelopmental defects conferred by knockdown of individual 16p12.1 homologs are

modulated by simultaneous knockdown of homologs of "second-hit" genes in *Drosophila*

melanogaster and *Xenopus laevis*. We observed that knockdown of 16p12.1 homologs affect

multiple phenotypic domains, leading to delayed developmental timing, seizure susceptibility,

brain alterations, abnormal dendrite and axonal morphology, and cellular proliferation defects.

Compared to genes within the 16p11.2 deletion, which has higher *de novo* occurrence, 16p12.1

homologs were less likely to interact with each other in *Drosophila* models or a human brain-

specific interaction network, suggesting that interactions with "second-hit" genes may confer

higher impact towards neurodevelopmental phenotypes. Assessment of 212 pairwise interactions

in *Drosophila* between 16p12.1 homologs and 76 homologs of patient-specific "second-hit"

genes (such as *ARID1B* and *CACNA1A*), genes within neurodevelopmental pathways (such as

PTEN and *UBE3A*), and transcriptomic targets (such as *DSCAM* and *TRRAP*) identified genetic

interactions in 63% of the tested pairs. In 11 out of 15 families, homologs of patient-specific

"second-hits" enhanced or suppressed the phenotypic effects of one or many 16p12.1 homologs.

In fact, homologs of *SETD5* synergistically interacted with homologs of *MOSMO* in both

Drosophila and *X. laevis*, leading to modified cellular and brain phenotypes, as well as axon

outgrowth defects that were not observed with knockdown of either individual homolog. Our

results suggest that several 16p12.1 genes sensitize the genome towards neurodevelopmental

defects, and complex interactions with "second-hit" genes determine the ultimate phenotypic

manifestation.

Author Summary

Copy-number variants, or deletions and duplications in the genome, are associated with multiple

neurodevelopmental disorders. The developmental delay-associated 16p12.1 deletion is mostly

inherited, and severely affected children carry an excess of "second-hits" variants compared to

mildly affected carrier parents, suggesting that additional variants modulate the clinical

manifestation. We studied this "two-hit" model using *Drosophila* and *Xenopus laevis*, and

systematically tested how homologs of "second-hit" genes modulate neurodevelopmental defects

observed for 16p12.1 homologs. We observed that 16p12.1 homologs independently led to

multiple neurodevelopmental features and weakly interacted with each other, suggesting that

interactions with "second-hit" homologs potentially have a higher impact towards

neurodevelopmental defects than interactions between 16p12.1 homologs. We tested 212

pairwise interactions of 16p12.1 homologs with "second-hit" homologs and genes within

conserved neurodevelopmental pathways, and observed modulation of neurodevelopmental

defects caused by 16p12.1 homologs in 11 out of 15 families, and 16/32 of these changes could

be attributed to genetic interactions. Interestingly, we observed that *SETD5* homologs interacted

with homologs of *MOSMO*, which conferred additional neuronal phenotypes not observed with

knockdown of individual homologs. We propose that the 16p12.1 deletion sensitizes the genome

to multiple neurodevelopmental defects, and complex interactions with "second-hit" genes

determine the clinical trajectory of the disorder.

Introduction

Rare recurrent copy-number variants (CNVs) account for about 15% of individuals with neurodevelopmental disorders, such as autism, intellectual disability, and schizophrenia (1, 2). While certain CNVs were initially associated with specific neuropsychiatric diagnoses, such as the 16p11.2 deletion and autism (3, 4), 3q29 deletion and schizophrenia (5), and 15q13.3 deletion and epilepsy (6), variable expressivity of phenotypes has been the norm rather than the exception for these CNVs (7). A notable example of this is the 520-kbp deletion encompassing seven genes on chromosome 16p12.1, which is associated with multiple neuropsychiatric disorders, including intellectual disability/developmental delay (ID/DD), schizophrenia, and epilepsy (8, 9). Furthermore, a large-scale study on a control population reported cognitive defects in seemingly unaffected individuals with the 16p12.1 deletion (10), suggesting that the deletion is sufficient to cause neuropsychiatric features. In contrast to other pathogenic CNVs that occur mostly *de novo*, the 16p12.1 deletion is inherited in more than 95% of individuals from a mildly affected or unaffected carrier parent (8, 9, 11). In fact, affected children with the deletion were more likely to carry another large CNV or deleterious mutation elsewhere in the genome ("second-hit") compared to their carrier parents (8, 9), providing evidence that additional rare variants modulate the effect of the deletion. These results suggest that the 16p12.1 deletion confers significant risk for disease and sensitizes the genome for a range of neuropsychiatric outcomes, while additional rare variants in the genetic background determine the ultimate phenotypic trajectory.

The extensive phenotypic variability and lack of chromosomal events, such as translocations and atypical deletions, have made causal gene discovery for variably-expressive CNVs such as the 16p12.1 deletion challenging. In particular, the developmental and neuronal

phenotypes associated with each individual 16p12.1 gene and the interaction models that explain

how "second-hit" genes modulate the associated phenotypes have not been assessed. Therefore, a

systematic evaluation of developmental, neuronal, and cellular defects caused by reduced

expression of individual 16p12.1 genes, as well as their interactions with each other and with

"second-hit" genes from patients with the deletion, would allow us to understand the functional

basis of the variable phenotypes associated with the deletion. *Drosophila melanogaster* and

Xenopus laevis serve as excellent models for systematic evaluation of developmental and tissue-

specific effects of multiple genes and their genetic interactions, as they are amenable for rapid

genetic manipulation and high-throughput evaluation. In fact, *Drosophila* have been classically

used to study the roles of genes and genetic interactions towards developmental and neurological

phenotypes (12-14). For example, Grossman and colleagues overexpressed human transgenes

from chromosome 21 in flies and identified synergistic interactions between *DSCAM* and

COL6A2, which potentially contribute to the heart defects observed in individuals with Down

syndrome (15). Furthermore, functional assays using *X. laevis* have uncovered developmental

defects, behaviors, and molecular mechanisms for several homologs of genes associated with

neurodevelopmental disorders, such as *NLGN1* (16), *CACNA1C* (17), *GRIK2* (18), and *PTEN*

(19).

Using *Drosophila* and *X. laevis* models, we recently found that multiple genes within the

variably expressive 16p11.2 and 3q29 deletion regions individually contribute to

neurodevelopmental defects (20, 21), suggesting that no single gene could be solely causative for

the wide range of defects observed with deletion of an entire region. Moreover, we identified

complex genetic interactions within conserved biological pathways among homologs of genes

affected by these CNVs. For example, fly and *X. laevis* homologs of *NCBP2* enhanced the

neuronal and cellular phenotypes of each of the other homologs of 3q29 genes (21), while fly

homologs of 16p11.2 genes interacted in cellular proliferation pathways in an epistatic manner to

enhance or suppress phenotypes of individual homologs (20). In fact, several aspects of the

interactions observed in our studies were also functionally or thematically validated in vertebrate

model systems, providing further evidence for the utility of these models to study complex

genetic interactions (22, 23). While our previous work showed pervasive interactions of

homologs within regions associated with neurodevelopmental disease, the deletions within these

regions occur primarily *de novo* (11), indicating a strong phenotypic impact associated with these

CNVs. In contrast, the 16p12.1 deletion is mostly inherited and more frequently co-occurs with

"second-hit" variants in affected individuals than other pathogenic CNVs (11), suggesting that

interactions involving "second-hit" genes may confer a higher impact towards the variable

neurodevelopmental phenotypes compared to those caused by interactions between genes within

the CNV region.

Here, using *Drosophila melanogaster* and *X. laevis* as two complementary model systems

of development, we present the first systematic assessment of conserved genes within the

16p12.1 deletion towards developmental, neuronal, and cellular phenotypes in functional models.

We found that knockdown of each individual 16p12.1 homolog affects multiple phenotypic

domains of neurodevelopment, leading to developmental delay and seizure susceptibility, brain

size alterations, neuronal morphology abnormalities, and cellular proliferation defects. These

defects were modulated by simultaneous knockdown of homologs of genes in established

neurodevelopmental pathways and transcriptome targets, as well as homologs of genes that

carried "second-hits" in affected children with the deletion, through genetic interactions and

"additive" effects. Our results suggest a model where reduced expression of each individual gene

within 16p12.1 is sufficient to sensitize the genome towards distinct neurodevelopmental defects,

which are then modulated by complex interactions with "second-hit" genes.

Results

Multiple homologs of 16p12.1 genes contribute to *Drosophila* and *X. laevis* development

We identified four conserved fly homologs out of the seven human protein coding 16p12.1 genes

using reciprocal BLAST and orthology prediction tools (**S1 Table**) (24). Using RNA

interference (RNAi) and the *UAS-GAL4* system (25), we reduced the expression of the four fly

homologs in a tissue-specific manner, and studied their individual contributions towards

developmental, neuronal, and cellular defects (**Fig 1**). A complete list of the fly lines used in this

study and full genotypes for all experiments are provided in **S1 File**. We authenticated the RNAi

lines by confirming 40-60% expression of the four homologs using RT-qPCR (**S1 Fig**). We note

that the genes are represented with fly gene names along with human counterparts at first

mention in the text, and as fly genes with allele names in the figures.

We first assessed the global role of 16p12.1 homologs during development by decreasing

their expression ubiquitously using the *da-GAL4* driver, and detected larval lethality with

knockdown of *Sin* (*POLR3E*) and larval and pupal lethality with knockdown of *UQCR-C2*

(*UQCRC2*) (**Fig 2A, S2 Fig**). Wing-specific *bx^{MS1096}-GAL4* mediated knockdown led to severe

phenotypes for *Sin* and severe defects and lethality for *UQCR-C2* fly models, recapitulating the

observations made with ubiquitous knockdown (**Fig 2A, S2 Fig**) and suggesting a role for these

homologs in signaling pathways required for early development (26-28). Next, we evaluated

whether decreased expression of the homologs leads to neuronal phenotypes frequently observed

in animal models of neurodevelopmental disease, including altered lifespan, susceptibility to

seizures, delayed developmental timing, changes in brain size, and dendritic arbor defects (29-

34). We observed early lethality in adult flies with nervous system-specific *Elav-GAL4*-mediated

knockdown of *Sin* and *CG14182* (*MOSMO*) (**Fig 2B**), while extended lifespan was observed

with knockdown of *UQCR-C2*, as previously reported for *Hsp26*, *Hsp27* (35), and *SOD* (36).

While altered mitochondrial activity has been shown to increase lifespan in *Drosophila* (37, 38),

further studies are necessary to understand the mechanism underlying this phenotype observed

with knockdown of *UQCR-C2*. *UQCR-C2* knockdown in the nervous system also led to

significantly greater recovery time when subjected to mechanical stress during bang sensitivity

assays, suggesting a higher susceptibility to developing seizures (32) (**S2 Fig**). Furthermore,

evaluation of developmental transitions revealed delayed pupariation and larval lethality with

knockdown of *Sin*, indicating a possible role for this gene in developmental timing, as well as

partial larval lethality for *CG14182* (**Fig 2C**). We also analyzed neuronal morphology in

Drosophila class IV sensory neurons using the *ppk-Gal4* driver (31, 39, 40), and identified

reduced complexity of dendritic arbors during development for *CG14182* (**Fig 2D**).

Measurements of total area of the developing third instar larval brain led to reduced brain sizes

with pan-neuronal knockdown of *CG14182* and *Sin* (**Fig 2E, S3 Fig**), which corresponded with a

decreased number of cells in the brain lobe stained with anti-phosphorylated-Histone 3 (pH3), a

marker for proliferating cells (**Fig 2F**). Interestingly, *Sin* knockdown also led to a reduction in

the number of apoptotic cells, as indicated by staining with anti-Death-caspase-1 (Dcp-1) (**S3**

Fig), likely reflecting its role in both proliferation and apoptotic processes (41, 42).

We then performed RNA-sequencing of fly heads with pan-neuronal knockdown of the

16p12.1 homologs. Gene Ontology (GO) enrichment analysis of differentially expressed genes

identified enrichments for multiple cellular, developmental, and neuronal processes (**S4 Fig, S2**

File). We found that each 16p12.1 homolog disrupted unique sets of genes and biological

functions, as 1,386/1,870 (74%) differentially expressed genes and 28/52 (53.8%) enriched GO

biological process terms were uniquely disrupted by one homolog (**S4 Fig**). Notably, we also

observed this trend among the human homologs of differentially-expressed genes, with 654/994

(65.8%) uniquely differentially expressed genes and 353/428 (82.5%) GO terms uniquely

disrupted by the 16p12.1 homologs, suggesting that they may act within independent pathways

(**S4 Fig**). For example, knockdown of *CG14182* altered the expression of fly homologs of human

genes involved in synapse assembly and transmission (*NLGN1*, *CEL*) as well as histone

methyltransferase binding (*NOP56, CBX1*). Similarly, human homologs of genes differentially

expressed with knockdown of *Sin* were involved in neuronal projection, neurotransmitter release

(such as *CHRNA7*, *KCNAB2*, and multiple solute carrier transport family genes, including

SLC6A1), and GABA pathways (such as *ADCY2*, *ADCY4*, and *ADCY7*), as well as in the

development of several non-neuronal organ systems, including cardiac, kidney, lung, and

muscle, further indicating the importance of *Sin* towards global development.

Next, we examined developmental phenotypes associated with decreased dosage of

homologs of 16p12.1 genes in *X. laevis*, a complementary vertebrate model system (**Fig 1** and

2G). We injected homolog-specific morpholinos at two- or four-cell stage embryos to reduce the

expression of each homolog to approximately 50% (partial knockdown), and further reduced

expression with higher morpholino concentrations (stronger knockdown) to increase our

sensitivity to detect more specific phenotypes (**S1 Fig**, see **Methods**). Reduced expression of

mosmo and *polr3e* led to severe craniofacial defects in stage 42 tadpoles, as measured by specific

facial landmarks, while milder defects were observed for *cdr2* (**Fig 2H, S5 Fig**). This suggests a

role for these homologs in key developmental processes involved in craniofacial morphogenesis,

such as neural crest cell formation and migration (43-48), and could potentially explain the

craniofacial changes observed in more than 50% of individuals with the 16p12.1 deletion (9). We

next examined axon outgrowth phenotypes in neural tube explants from stage 20-22 injected *X*.

laevis embryos, and found that stronger knockdown of *mosmo* (20 ng morpholino) led to a

significant reduction in axon length (**Fig 2I, S5 Fig**), suggesting a potential role for the homolog

in cytoskeletal signaling processes involved in axon outgrowth (49). Furthermore, stronger

knockdown of *mosmo* (20 ng morpholino) and *polr3e* (20 ng morpholino) resulted in decreased

forebrain and midbrain area (**Fig 2J, S5 Fig**), in concordance with the brain size defects we

observed in *Drosophila* models. Interestingly, partial knockdown of *mosmo* (12 ng morpholino)

also led to a severe reduction in forebrain and midbrain area (**S5 Fig**). Western blot analysis for

whole embryo lysates using anti-pH3 antibody as a marker for cellular proliferation showed

decreased proliferation with knockdown of *polr3e*, while knockdown of *mosmo* did not lead to

any significant changes (**S6 Fig**). Overall, these results suggest that homologs of 16p12.1 genes

individually contribute to multiple developmental defects and affect distinct developmental,

neuronal, and cellular processes in *Drosophila* and *X. laevis*.

Weak genetic interactions and combined independent effects of 16p12.1 homologs mediate

neurodevelopmental defects

Our previous studies identified several potential models for how genes within CNVs

combinatorially influence neurodevelopmental phenotypes (20, 21, 50). As multiple homologs of

16p12.1 genes contribute towards developmental, neuronal, and cellular phenotypes, we assessed

the effect of simultaneous knockdown of pairs of 16p12.1 fly homologs. The *Drosophila* fly eye

has been widely used to identify genetic interactions that disrupt ommatidial organization during

development (51), and modifier genes for homologs of several human diseases, including

Spinocerebellar ataxia type 1 (52), Huntington's disease (52), and Fragile X syndrome (53), have

been studied in flies. We assessed whether eye-specific *GMR-GAL4* mediated knockdown of

individual homologs led to phenotypes, and evaluated the severity of eye roughness using

Flynotyper, a tool that quantifies the levels of ommatidial disorderliness in the adult fly eye (54).

We observed that knockdown of *Sin* led to a subtle disruption of ommatidial organization

compared with controls, while no such phenotypes were observed with knockdown of other

homologs (**S7 Fig**). Further reduction in expression of the 16p12.1 homologs using *GMR-GAL4*

and overexpression of *Dicer2* led to more severe eye phenotypes for *CG14182* and *Sin* (**S7 Fig**).

As *GMR-GAL4*-mediated knockdown of the 16p12.1 homologs only exhibited modest eye

phenotypes, we used *Flynotyper* scores as a sensitive quantitative trait with a wide dynamic

range to assess for combinatorial effects of 16p12.1 homologs. We therefore generated *GMR-*

GAL4 eye-specific recombinant lines for each homolog and crossed them with multiple RNAi

lines for other 16p12.1 homologs (see **Methods**, **S1 File**), to test a total of 30 two-hit crosses for

12 pairwise knockdowns (**Fig 1** and **3A, S8 Fig**).

We performed two independent analyses to interpret the combinatorial effect of 16p12.1

homologs on eye phenotypes. *First*, we assessed whether the phenotypic severity observed with

knockdown of a 16p12.1 homolog is enhanced or suppressed with simultaneous knockdown of a

second homolog. We observed significant changes in eye severity for four pairwise knockdowns

compared with single hit recombinant lines crossed with controls, which were further validated

with multiple RNAi lines (**S3 File**). For example, we observed that simultaneous knockdown of

UQCR-C2 with *Sin* or *CG14182* led to an increase in eye phenotype compared to knockdown of

UQCR-C2 crossed with control (**Fig 3B** and **3C**). Similarly, decreased expression of *Sin* led to an

enhancement of the *CG14182* eye phenotype. *Second*, to quantitatively assess whether the

change in phenotypic severity due to pairwise knockdowns could be attributed to genetic

interactions, we applied a "multiplicative" model to the *Flynotyper* scores. The multiplicative

model estimates the expected combined effect (i.e., no interaction) of two gene mutations as the

product of the phenotypes observed with individual gene mutations (**Fig 3D, S9 Fig**), and

identifies any deviation of the observed phenotypes from the expected values as positive

(ameliorating the phenotype) or negative/synergistic (aggravating the phenotype) interactions.

This strategy has been widely applied to identify fitness-based genetic interactions (55, 56), and

more recently, to assess for interactions contributing to non-fitness-related quantitative

phenotypes, such as cell count, nuclear area (57), and protein folding in the endoplasmic

reticulum (58). After applying the multiplicative model to our pairwise interaction data, we

identified five pairwise combinations of 16p12.1 homologs that were validated using multiple

RNAi lines (**S2 Table**). Two pairwise combinations corresponded with no interactions, while the

remaining three were positive genetic interactions, with an observed phenotype milder than

expected (**S3 File**). Only one out of the four pairwise knockdowns that resulted in significant

changes in eye severity compared with single hit recombinant lines crossed with control

corresponded with a genetic interaction, while the rest were not validated across multiple fly

lines tested (**S3 File**). To contextualize these observations, we compared the strength of genetic

interactions among the 16p12.1 homologs to those of homologs of genes affected by the autism-

associated 16p11.2 deletion, a region with reported pervasive genetic interactions (20). We

quantified the magnitude of genetic interactions using "interaction scores", defined as the $\log_2$

ratio between the observed and expected phenotypic values from the multiplicative model (see

Methods), and found significantly lower interaction scores for the 16p12.1 homologs compared

to the 16p11.2 homologs (**S4 File**).

We further investigated the effects of the combined knockdown of homologs of *MOSMO*

and *POLR3E*, genes that individually contributed to multiple defects in both fly and *X. laevis*

models, towards *X. laevis* development (**Fig 1** and **3E**). Pairwise interactions in *X. laevis* models

were tested using partial knockdown of the homologs to avoid potential lethality with stronger

knockdown. Partial pairwise knockdown of *polr3e* (10 ng morpholino) and *mosmo* (12 ng

morpholino) showed significantly reduced forebrain and midbrain area when compared to

knockdown of *polr3e* alone (**Fig 3F**), but not when compared to knockdown of *mosmo* alone.

Similarly, we assessed whether *mosmo* and *polr3e* interact to modulate cellular proliferation

processes during *X. laevis* development and did not observe any changes in anti-pH3 signals with

combined knockdown of *polr3e* and *mosmo* compared with knockdown of *polr3e* alone (**Fig 3G,**

S6 Fig). Overall, our analysis in *Drosophila* and *X. laevis* suggest that 16p12.1 homologs

contribute towards neurodevelopmental phenotypes through both weak genetic interactions and

combined independent effects.

Homologs of 16p12.1 genes interact with genes in conserved neurodevelopmental pathways

We recently identified genetic interactions between fly homologs of CNV genes and conserved

genes in neurodevelopmental pathways, providing functional evidence that phenotypes of CNV

genes are modulated by key neurodevelopmental genes (20, 21). As our functional analyses

showed that knockdown of each 16p12.1 homolog resulted in multiple neuronal and

developmental phenotypes, we hypothesized that genes involved in conserved

neurodevelopmental pathways could modulate phenotypes due to knockdown of 16p12.1

homologs through genetic interactions. We therefore performed 255 crosses to test 116 pairwise

gene combinations between eye-specific recombinant lines for each of the four 16p12.1

homologs and 13 homologs of known neurodevelopmental genes and 39 homologs of

transcriptional targets (**Fig 4A, S10 Fig, S2 Table, S3 Table**). As validation, we used multiple

RNAi, mutant or overexpression lines when available (**Fig 1**). Details of the number of

homologs, fly lines, and crosses used for all interaction experiments are provided in **S2 Table**.

First, we screened for 55 combinations between homologs of 16p12.1 genes and 13 homologs of

human genes in established developmental pathways, such as synapse function

(*Prosap*/*SHANK3*), cell division (*Pten/PTEN*), and chromatin modulation (*kis/CHD8*), as well as

genes functionally related to 16p12.1 homologs (54, 59-61). Using *Flynotyper* to quantify adult

eye defects and the multiplicative model to identify genetic interactions, we identified

interactions specific to an individual 16p12.1 homolog or those involving multiple homologs

(**S11-S16 Fig, S3 File**). For example, *CG10465* (*KCTD13*) negatively interacted with *UQCR-C2*

and *CG14182*, leading to significantly more severe phenotypes than expected using the

multiplicative model (**Fig 4B**). Similarly, simultaneous knockdown of *Sin* with *kis* led to an

exaggerated eye phenotype, suggesting negative interactions between the genes (**Fig 4B**).

Overall, we confirmed 22 interactions out of the 55 pairwise combinations (40%) tested,

including both positive (12/55) and negative (10/55) effects (**Fig 4C-D, S17 Fig, S2 Table**).

Next, to identify interactors of 16p12.1 homologs towards developmental functions and pathways

(**S10 Fig, S3 Table)**, we screened for interactions of the homologs with 25 dysregulated fly

genes selected from our transcriptome studies as well as 14 genes within enriched Gene

Ontology categories, such as nervous system development and function (*Dscam1*, *Asp*, *mGluR*,

NaCP60E), protein folding (*Hsp23*, *Hsp26*, *Hsp70Ab*), and muscle contraction (*Actn*, *ck*). We

identified interactions for 42 out of 61 tested pairs (68.8%, **S2 Table**), validated using additional

lines when available (**S3 File**). For example, knockdown of *Gat* (*SLC6A1*), *Dscam4* (*DSCAM)*,

Nipped-A (*TRRAP*), and *aurB* (*AURKB*) each modified the eye phenotype due to knockdown of

Sin through positive or negative genetic interactions (**Fig 4C, S13 Fig, S16 Fig, S3 File**).

Furthermore, the protein-folding gene *Hsp26* (*CRYAA)* was differentially expressed with

knockdown of *Cen*, and its knockdown enhanced the phenotype of *Cen* through a negative

interaction (**Fig 4C, S12 Fig, S3 File**). Overall, we identified 64 pairwise interactions between

the 16p12.1 homologs and genes from established neuronal functions and transcriptome targets

(**Fig 4C-D, S2 Table**), suggesting that phenotypes of 16p12.1 homologs can be modulated by

genes within multiple neurodevelopmental pathways through genetic interactions.

Homologs of patient-specific "second-hit" genes modulate phenotypes of 16p12.1 homologs

We recently found that an increased burden of rare variants (or "second-hits") outside of disease-

associated CNVs, such as 16p11.2 deletion, 15q13.3 deletion, and 16p12.1 deletion, contributed

to variability of cognitive and developmental phenotypes among affected children with these

CNVs (8, 9, 11). In fact, we found that severely affected children with the 16p12.1 deletion had

additional loss-of-function or severe missense variants within functionally intolerant genes

compared to their mildly affected carrier parents (8, 9, 11). We hypothesized that homologs of

genes carrying patient-specific "second-hits" modulate the effects of individual 16p12.1

homologs not only "additively" but also through genetic interactions (**S10 Fig, S3 Table**). To

test this, we performed 227 crosses to study 96 pairwise interactions between eye-specific

recombinant lines for each of the four 16p12.1 homologs and 46 RNAi or mutant lines for 24

homologs of patient-specific "second-hit" genes identified in 15 families with the 16p12.1

deletion (**Fig 1, S18 Fig, S2-S4 Tables**) (9). Out of the 96 combinations tested, we identified 32

pairwise knockdowns that modulated the phenotype of a 16p12.1 homolog, confirmed with

additional lines when available, for 11 out of 15 families carrying "second-hit" genes (**Fig 5A-C,**

S2-S4 Tables). In fact, the phenotypic effects of 16 out of 32 combinations were attributed to

genetic interactions (**Fig 4D, S3 File**). Interestingly, we observed that different "second-hit"

homologs showed distinct patterns of interactions with homologs of 16p12.1 genes (**Fig 5B, S3**

File). For example, the affected child in family GL_11 carried "second-hit" pathogenic

mutations in *NRXN1* and *CEP135* (**Fig 5A**). Knockdown of the fly homolog *Nrx-1* enhanced the

eye phenotype caused by knockdown of *Sin* and *UQCR-C2*, while simultaneous knockdown of

Cep135 with *UQCR-C2* led to lower phenotypic score compared to knockdown of *UQCR-C2*

alone (**Fig 5B**). While the two-hit phenotypes were not significantly different from the expected

combined effects of the individual genes in these cases (**S3 File**), we observed genetic

interactions between *Nrx-1* and *CG14182*, and *Cep135* and *Sin* or *Cen*, suggesting potential

functional connections between these genes (**Fig 5D**). Interestingly, for 11/96 combinations

tested with the multiplicative model, we found that the phenotype of the pairwise knockdown

was more severe compared to the phenotype observed for the knockdown of an individual

16p12.1 homolog, but significantly less severe than the expected effects, suggesting potential

buffering against deleterious combined independent effects of the genes (62, 63) (**Fig 4C, Fig**

5D, S3 File). In another example, the affected child in family GL_01 carried inherited "second-

hit" variants in *LAMC3* and *DMD,* as well as a *de novo* loss-of-function mutation in the

intellectual disability-associated and chromatin regulator gene *SETD5* (64) (**S18 Fig**).

Knockdown of *Lanb2*, homolog of *LAMC3*, enhanced the phenotype caused by knockdown of

UQCR-C2, although they positively interacted towards a milder phenotype than expected (**S3**

File). Furthermore, *upSET*, homolog of *SETD5*, led to enhancements of the phenotypes caused

by knockdown of *Sin* and *CG14182* (**S3 File**). Interestingly, while the phenotype caused by

simultaneous knockdown of *Sin* and *upSET* was not different from expected using the

multiplicative model, *upSET* synergistically interacted with *CG14182*, leading to an enhanced

eye phenotype with pairwise knockdown (**Fig 6A**). To assess the cellular changes affected by

this interaction during development, we tested for alterations in apoptosis and proliferation in the

third instar larval eye discs, and found that simultaneous knockdown of *CG14182* and *upSET* led

to an increased number of cells undergoing proliferation and apoptosis compared to knockdown

of *CG14182* alone (**Fig 6B**). Interestingly, we also identified interactions between *CG14182* and

other chromatin modifier genes, including *Nipped-A*, a transcriptional target of *Sin*, and *Osa*,

homolog of the "second-hit" gene *ARID1B*, identified in family GL_13 (**S16 Fig, S3 File**). These

interactions also modulated cellular proliferation and apoptosis processes in the developing eye

discs observed with knockdown of *CG14182* (**S19 Fig**).

We further evaluated whether interactions between the fly homologs of *POLR3E* and

MOSMO with *SETD5* were also conserved during vertebrate development, and studied brain and

axon outgrowth phenotypes of homologs of these genes in *X. laevis* (**Fig 1**). We observed that

simultaneous knockdown of *polr3e* and *setd5* led to smaller forebrain and midbrain areas

compared with *polr3e* knockdown alone (**S20 Fig**). Similarly, simultaneous knockdown of

mosmo and *setd5* led to a further reduction in midbrain area than that observed with knockdown

of *mosmo* alone (**Fig 6C**). Furthermore, analysis of axon outgrowth in developing *X. laevis*

embryos showed that simultaneous knockdown of *mosmo* and *setd5* led to significantly reduced

axon length compared to the individual knockdowns of either *mosmo* or *setd5*, while no changes

were observed with simultaneous knockdown of *polr3e* and *setd5* (**Fig 6D, S20 Fig**). In fact, the

axon outgrowth defect observed with simultaneous knockdown of *mosmo* and *setd5* was not

observed with partial knockdown of either individual homolog. This result suggests a genetic

interaction between *mosmo* and *setd5* during vertebrate nervous system development. Overall,

our results show that interactions with "second-hit" genes can modulate neurodevelopmental and

cellular phenotypes associated with homologs of 16p12.1 genes.

Discussion

We previously described multiple models for how genes within CNVs contribute towards

neurodevelopmental phenotypes (20, 21, 50). Here, we analyzed neurodevelopmental defects and

cellular and molecular mechanisms due to individual and pairwise knockdown of conserved

16p12.1 homologs in *Drosophila* and *X. laevis*, and evaluated how these defects are modulated

by homologs of "second-hit" genes. Our results provide multiple hypotheses for how genes

within the deletion contribute to neurodevelopmental phenotypes. *First*, in line with our previous

findings for homologs of genes within CNV regions (20, 21), our results show that no single

homolog within the 16p12.1 region is solely responsible for the observed neurodevelopmental

phenotypes. In fact, we observed a global developmental role for multiple 16p12.1 homologs, as

well as specific roles of each homolog towards craniofacial and brain development (**S5 Table**).

This was further confirmed by interactions of 16p12.1 homologs with genes in conserved

neurodevelopmental pathways. Our findings are in accordance with the core biological functions

described for some of these genes. *For example*, *POLR3E* encodes a cofactor of the RNA

polymerase III, which is involved in the transcription of small RNA, 5S ribosomal RNA and

tRNA (65), while *MOSMO* is a negative regulator of the hedgehog signaling pathway (66).

Second, knockdown of individual homologs sensitized both model organisms towards specific

phenotypes. For example, knockdown of homologs of *MOSMO* led to neuronal morphology

defects and knockdown of homologs of *POLR3E* led to brain size phenotypes that correlated

with cellular proliferation defects in both model systems, while knockdown of *UQCR-C2* led to

seizure susceptibility in flies. *Third*, we found that the 16p12.1 homologs were less likely to

interact with each other (3 interactions out of 12 pairs tested) compared to their interactions with

downstream transcriptome targets (42 interactions out of 61 pairs tested, Fisher's exact test,

p=0.0077). These results suggest reduced functional overlap among the 16p12.1 homologs, an

observation supported by the distinct sets of biological functions enriched among the

differentially expressed genes obtained with knockdown of each individual homolog (**S4 Fig**).

Beyond the four conserved homologs evaluated in this study, little functional information is

available on the other genes in the region, including *VWA3A* and *PDZD9* as well as non-protein

coding genes. Results from mouse models of *EEF2K*, which encodes a kinase associated with

protein synthesis elongation, have postulated associations of this gene with synaptic plasticity

(67), learning and memory (68), atherosclerosis-mediated cardiovascular disease (69), and

depression (70). Although *EEF2K* could function in concert with the tested 16p12.1 genes to

contribute towards neurodevelopmental features, it showed low connectivity (29[th] percentile) to

other 16p12.1 genes in a human brain-specific interaction network compared to the pairwise

connectivity of all genes in the network (71, 72) (**S5 File**). *VWA3A* showed even lower

connectivity (5[th] percentile) to other 16p12.1 genes compared to all gene pairs. Further functional

analyses that includes all protein-coding and non-coding genes are necessary for a

comprehensive understanding of the consequences of the entire deleted region, as these genes

may also contribute towards the pathogenicity of 16p12.1 deletion.

We recently showed that additional variants or "second-hits" modulate the manifestation

of developmental and cognitive phenotypes associated with disease-causing variants, including

intelligence quotient and head circumference phenotypes (8, 9, 11). Using the 16p12.1 deletion

as a paradigm for a complex disorder, we examined how homologs of genes carrying "second-

hit" variants modulate the phenotypes caused by decreased expression of individual CNV

homologs. For example, homologs of *ARID1B*, *CEP135* and *CACNA1A* suppressed the eye

phenotypes and interacted with one or more 16p12.1 homologs. Furthermore, we identified a

negative interaction between homologs of *MOSMO* and *SETD5,* which led to novel

neurodevelopmental phenotypes in both *Drosophila* and *X. laevis* compared with knockdown of

either individual homolog. Interestingly, mouse embryonic stem cells lacking *Setd5* exhibited

dysregulation of genes involved in hedgehog signaling (73), a key pathway recently associated

with *MOSMO* function (66). Moreover, we observed that *MOSMO* and *SETD5* are highly

connected to each other in a human brain-specific interaction network compared to all genes in

the genome (top 84th percentile compared to all genetic interactions with *MOSMO*), suggesting

that the human genes may also be functionally related (71, 72). We further observed interactions

between *CG14182* and other genes with chromatin regulating function, such as *Nipped-A*

(*TRRAP*) and *osa* (*ARID1B*) (**S16 Fig**). Based on these observations, we propose that while

genes carrying "second-hit" variants in combination with the deletion may "additively"

contribute towards more severe phenotypes, they may also interact towards developmental

phenotypes, conferring high impact towards variable defects associated with the 16p12.1

deletion (**S21 Fig**). The ultimate nature of the interactions will depend on the role of the

individual CNV genes towards specific phenotypes, the identity of genes carrying "second-hits",

as well as the molecular complexity associated with each phenotypic domain (50).

The high inheritance rate of the 16p12.1 deletion (8, 9) suggests that while it confers risk

for several phenotypes, the CNV can be transmitted through multiple generations until additional

variants accumulate and cumulatively surpass the threshold for severe disease (7). In contrast,

other CNVs associated with neurodevelopmental disease, such as the autism-associated 16p11.2

deletion and the 17p11.2 deletion that causes Smith-Magenis syndrome, occur mostly *de novo*

and are less likely to co-occur with another "second-hit", suggesting a higher pathogenicity on

their own (11). For example, the 16p11.2 deletion occurs *de novo* in approximately 60% of the

cases, and only 8% of the affected children carry another rare large CNV, in contrast to 25% of

severely affected children with 16p12.1 deletion that carry a "second-hit" large CNV (11). When

we compared experimental results from 16p12.1 homologs with those from fly homologs of

16p11.2 genes (20), we found evidence that the varying pathogenicity of the CNVs could be

explained by differential connectivity and combinatorial effects of genes within each region (**Fig**

7A) (74). For example, we previously found that 24 out of the 52 tested pairwise knockdowns of

16p11.2 homologs led to enhancement or suppression of phenotypes, significantly modifying the

effect of the individual genes (20). In contrast, only four out of twelve tested combinations

between 16p12.1 homologs led to a slight change in phenotypic severity, which in aggregate

showed lower phenotypic scores than those observed for pairwise knockdown of 16p11.2

homologs (**Fig 7B** and **S22 Fig**). In fact, using a multiplicative model, we found that the

magnitude of interactions between homologs of 16p11.2 genes was stronger than that observed

between 16p12.1 homologs (**Fig 7C**). Moreover, transcriptome analyses showed a higher overlap

of differentially expressed genes among 16p11.2 homologs compared to 16p12.1 homologs,

further suggesting a higher functional relatedness among the 16p11.2 genes (**Fig 7D**). We

similarly compared the connectivity of genes within both CNV regions in a human brain-specific

interaction network (71, 72), and found that 16p11.2 genes were more strongly connected to each

other than were 16p12.1 genes, and were also more strongly connected to each other than with

other genes in the genome (**Fig 7E and 7F, S22 Fig**). Furthermore, the connectivity of 16p12.1

genes was lower than 99.6% of connectivity values of simulated sets of six 16p11.2 genes (0.4th

percentile). In fact, the connectivity values of genes within 16p12.1 and 16p11.2 were in the

lower (~6[th]) and higher (~79[th]) percentiles of connectivity values from simulated sets of

contiguous genes in the genome, respectively (**S23 Fig**). Interestingly, genes connecting pairs of

16p11.2 genes were enriched for genes intolerant to functional variation (**Fig 7E and 7G**), such

as *ASH1L*, a histone methyltransferase activator and autism candidate gene (75), and *CAMK2B,* a

protein kinase gene causative for intellectual disability (76). In contrast, connector genes unique

to 16p12.1 genes were not associated with neurodevelopmental disease or enriched for genes

intolerant to variation (**S5 File**). This suggests that the 16p11.2 deletion disrupts a tight network

of key genes in the brain, including other neurodevelopmental genes and genes with disease

relevance (77, 78). Overall, we propose that 16p12.1 genes contribute towards multiple

neurodevelopmental phenotypes through weak genetic interactions and "additive" effects with

"second-hit" genes and exhibit less functional connectivity compared with 16p11.2 genes,

leading to a high transmissibility of the deletion and allowing for higher impact "second-hit"

variants to modulate neurodevelopmental phenotypes.

Our study provides the first systematic analysis of individual and pair-wise contributions

of 16p12.1 homologs towards neurodevelopmental phenotypes and associated cellular and

molecular mechanisms, and identifies a key role of genetic interactions with "second-hit"

homologs towards variability of phenotypes. Our work does not intend to recapitulate human

disease, but rather highlights the basic cellular roles of individual conserved genes and their

interactions towards neurodevelopmental phenotypes. As such, these findings should be further

examined in higher-order model systems, including mouse and human cellular models. Our

functional analyses suggest a model where 16p12.1 genes sensitize an individual towards defects

in different domains of neurodevelopment, but the ultimate phenotypic manifestation may

depend on complex interactions with "second-hits" in the genetic background.

Materials and Methods

Ethics statement

All *X. laevis* experiments were approved by the Boston College Institutional Animal Care and

Use Committee (Protocol #2016–012), and were performed according to national regulatory

standards.

Drosophila stocks and genetics

Using *Ensembl* database (79), NCBI Protein-Protein BLAST tool (80), and DRSC Integrative

Ortholog Prediction Tool (DIOPT) (24), we identified four homologs out of the seven genes

within the 16p12.1 deletion region in *Drosophila melanogaster* (**S1 Table**). No fly homologs

were present for three genes, including *VWA3A*, *PDZD9* and *EEF2K*. Similar strategies were

used to identify fly homologs of conserved neurodevelopmental genes and genes carrying

"second-hits" in children with the 16p12.1 deletion. Fly Atlas Anatomy microarray expression

data from FlyBase confirmed the expression of each 16p12.1 homolog in the nervous system

during *Drosophila* development (**S1 Table**) (81), and expression data from Xenbase (82)

confirmed the expression of the homologs in *X. laevis* brain.

  Multiple RNAi lines were used to test neurodevelopmental defects of 16p12.1 homologs

(S1 File). RNAi, mutant, or overexpression lines for fly homologs were obtained from the

Vienna Stock Resource Center (VDRC), Bloomington *Drosophila* Stock Center (BDSC) (NIH

P40OD018537), or Kyoto Stock Center (S1 File). The following lines used were generated in

various research labs: *Drice$^{17–1}$* and *Drice$^{17–2}$* from Bergmann lab (83), *GluRIIB $^{Overexp\ EGFP}$* from

Sigrist lab (84), *Hsp26$^{Overexp\ Hsp26}$* from Benzer lab (35), and *Hsp70Ab$^{Overexp\ Hsp70-9.1}$* and

Hsp70Ab$^{Overexp\ Hsp70-4.3}$ from Robertson lab (85). Tissue-specific knockdown of homologs of

16p12.1 genes was achieved using the *UAS-GAL4* system (25), with specific *GAL4* lines

including *w^{1118};dCad-GFP, GMR-GAL4/CyO* (Zhi-Chun Lai, Penn State University),

w^{1118};GMR-GAL4; UAS-Dicer2 (Claire Thomas, Penn State University), *w^{1118},mcd8-GFP*, *Elav-*

GAL4/Fm7c;; UAS-Dicer2 (Scott Selleck, Penn State University), *w^{1118},Elav-GAL4* (Mike

Groteweil, VCU), *w^{1118};;Elav-GAL4,UAS-Dicer2* (Scott Selleck, Penn State University),

w^{1118};da- GAL4 (Scott Selleck, Penn State University), *w^{1118},bx^{MS1096}-GAL4;; UAS-Dicer2* (Zhi-

Chun Lai, Penn State University), and *UAS-Dicer2; ppk-GAL4, UAS-mCD8-GFP* (Melissa

Rolls, Penn State University). A list of full genotypes for all fly lines and crosses tested in this

study is provided in **S1 File**. Fly crosses were reared on a cornmeal-sucrose-dextrose-yeast

medium at room temperature (RT), 25°C or 30°C. For all experiments, RNAi lines were

compared to a control with the same genetic background to account for background-specific

effects (**S1 File**). Three different controls were used: *w^{1118}* from VDRC (GD stock # 60000), in

which inverted repeats are inserted by P-element insertion; *y,w^{1118}* from VDRC (KK stock #

60100), where inverted repeats are inserted by site-specific recombination; and *{y[1] v[1];*

*P{y[+t7.7]=CaryP}*attP2 from BDSC (TRiP stock # 36303).

RT-quantitative PCR for *Drosophila* RNAi knockdown of 16p12.1 homologs

Decreased expression of homologs of 16p12.1 genes in the nervous system was confirmed using

reverse transcription quantitative PCR (RT-qPCR) for individual *Drosophila* RNAi lines.

Decreased expression of the genes was achieved using *Elav-GAL4;;UAS-Dicer2* lines, reared at

25°C. As nervous system-specific knockdown of *Sin* with *Elav-GAL4;;UAS-Dicer2* caused

developmental lethality in all three RNAi lines tested (*Sin^{GD7027}, Sin^{KK101936}, Sin^{HMC03807}*), we

confirmed knockdown of *Sin* using *Elav-GAL4* without overexpression of *Dicer2* and reared at

RT. We note that all experiments with nervous system-specific knockdown of *Sin* were

performed under these conditions. Briefly, three biological replicates, each containing 35-40

female heads, were collected after being separated by repeated freezing in liquid nitrogen and

vortex cycles. Total RNA was extracted from *Drosophila* heads using TRIzol (Invitrogen,

Carlsbad, CA, USA), and cDNA was generated using qScript cDNA synthesis kit (Quantabio,

Beverly, MA, USA). Quantitative RT-PCR was performed in an Applied Biosystems Fast 7500

system using SYBR Green PCR master mix (Quantabio), with *rp49* as the reference gene.

Primers were designed using NCBI Primer-BLAST (86), with primer pairs separated by an

intron in the corresponding genomic DNA, if possible. **S6 Table** details the primers used to

quantify the level of expression of 16p12.1 homologs. The delta-delta Ct method was used to

calculate the percentage of expression compared to the control (87), and statistical significance

compared to the control was determined using t-tests.

Eye and wing imaging

Eye-specific knockdown of the 16p12.1 homologs was achieved using *GMR-GAL4* driver at

30°C. Female progeny were collected on day 2-3 and imaged using an Olympus BX53

compound microscope with LMPLan N 20X air objective and a DP73 c-mount camera at 0.5X

magnification, with a z-step size of 12.1μm (Olympus Corporation, Tokyo, Japan). Individual

image slices were captured using the CellSens Dimension software (Olympus Corporation,

Tokyo, Japan), and were stacked into their maximum projection using Zerene Stacker software

(Zerene Systems, Richland, WA, USA). Wing phenotypes were assessed in day 2-5 female

progeny from *bx^{MS1096}-GAL4* lines crossed to the RNAi lines at 25°C. Adult wings were imaged

using a Zeiss Discovery V20 stereoscope (Zeiss, Thornwood, NY, USA) and a ProgRes Speed

XT Core 3 camera (Jenoptik AG, Jena, Germany) with a 40X objective. Adult wing images were

captured using ProgRes CapturePro v.2.8.8 software. We characterized qualitative phenotypes

for between 10-20 wing images, including curly, wrinkled, shriveled, dusky or vein defects, and

10-30 eye images were assessed for rough, glazed, eye size, and necrotic patches defects.

Quantitative assessment of rough adult eye phenotypes was performed using a software called

Flynotyper (20, 54), which calculates a phenotypic score for each eye image by integrating the

distances and angles between neighboring ommatidia. The phenotypic scores generated by

Flynotyper were compared between RNAi lines and their respective controls using one-tailed

Mann-Whitney tests, with Benjamini-Hochberg correction for multiple tests.

Lifespan Measurement

Lifespan assessment of homologs of 16p12.1 genes was performed as previously reported (88).

Briefly, fly crosses were set up at 25°C with *Elav-GAL4;;UAS-Dicer2* for each of the fly

homologs, or *Elav-GAL4* at RT for *Sin^{GD7027}*. In all cases, the newly emerged progeny were

collected every day for five consecutive days, and the birth date was recorded. F1 flies were

allowed to mate for 24 hours, and were separated under CO_2 into at least four vials, each

containing 20 females. Vials were transferred every 2-3 days, and the age and number of living

flies were registered. One-way repeated measures ANOVA with post-hoc pairwise t-tests were

performed to identify changes in lifespan for the individual 16p12.1 homologs.

Bang-sensitive assay

Sensitivity to mechanical stress was assessed in females with decreased expression of 16p12.1

homologs in the nervous system, using *Elav-GAL4;;UAS-Dicer2* and reared at 25°C. *Sin* was

excluded from the analysis, as adult flies with *Elav-GAL4* knockdown of the gene exhibited

severe motor defects. Ten female flies from the progeny were collected on day 0-1 for ten

consecutive days, and experiments were performed on day 2-3. Flies were individually separated

under CO_2 24 hours before the experiments, collected in culture vials containing food, and

transferred to another empty culture vial the day of the experiment. Identification of bang-

sensitive phenotypes was performed as previously reported (89). Each vial was vortexed at

maximum speed (Fischer Scientific) for 15 seconds, and the time for each fly to recover from

paralysis was registered. Differences in bang-sensitivity compared with controls were identified

using two-tailed Mann-Whitney tests.

Assessment of delay in developmental timing

Pupariation time was assessed in third instar larvae obtained from crosses between RNAi lines

and w^{1118}*;;Elav-GAL4,UAS-Dicer2* or w^{1118}*,Elav-GAL4* flies. Developmentally-synced larvae

were obtained from apple juice plates with yeast paste, and were reared for 24 hours. Thirty

newly emerged first instar larvae were transferred to culture vials, for a total of four to ten vials

per RNAi line tested. The number of larvae transitioning to pupae were counted every 24 hours.

Significant differences in pupariation timing compared with the control across the duration of the

experiment were identified with one-way repeated measures ANOVA and post-hoc pairwise t-

tests.

Dendritic arborization experiments

Class IV sensory neuron-specific knockdown was achieved by crossing the RNAi lines to *UAS-*

Dicer2; ppk-GAL4 driver at 25°C in apple juice plates. First instar larvae were collected and

transferred to cornmeal-based food plates for 48 hours. Z-stack images of the dorsal side of third

instar larvae were obtained using a Zeiss LSM 800 (Zeiss, Thornwood, NY, USA) confocal

microscope. To perform Sholl analyses, we assessed the number of intersections of dendrite

branches with four concentric circles starting from the cell body and separated by 25 μm. The

total number of intersections was normalized to the width of the larval hemi-segment, and

significant changes compared with control were assessed using two-tailed Mann-Whitney tests.

Measurement of larval brain area

Larval brain area was assessed in third instar larvae obtained from crosses between the RNAi

lines with *Elav-GAL4*. Crosses were set up in apple plates containing yeast paste to control for

size effects generated by food availability. Fifteen first instar larvae were transferred to culture

vials containing a fixed volume (8-10 mL) of cornmeal-based food. Brains were dissected from

third instar larva in PBS (13mM NaCl, 0.7mM Na_2HPO_4, and 0.3mM NaH_2PO_4), fixed in 4%

paraformaldehyde in PBS for 20 minutes, washed three times in PBS, and mounted in Prolong

Gold antifade reagent with DAPI (Thermo Fisher Scientific, P36930). Z-stacks of *Drosophila*

brains were acquired every 10μm with a 10X air objective with 1.2X magnification using an

Olympus Fluoview FV1000 laser scanning confocal microscope (Olympus America, Lake

Success, NY). The area of the maximum projection of the Z-stack was measured using Fiji

software (90). Differences in brain area were assessed using two-tailed Mann-Whitney tests.

RNA sequencing and differential expression analysis in *Drosophila melanogaster*

RNA sequencing was performed for three biological replicates of RNA isolated from 35-40

Drosophila heads with *Elav-GAL4* mediated nervous system-specific knockdown of 16p12.1

homologs as well as controls with matching drivers and rearing temperatures. cDNA libraries

were prepared with TruSeq Stranded mRNA LT Sample Prep Kit (Illumina, San Diego, CA).

Single-end 100bp sequencing of the cDNA libraries was performed using Illumina HiSeq 2000

at the Pennsylvania State University Genomics Core Facility, at an average coverage of 35.1

million reads/sample. Quality control was performed using Trimmomatic (91), and raw

sequencing data was aligned to the fly reference genome and transcriptome build 6.08 using

TopHat2 v.2.1.1 (92). Total read counts per gene were calculated using HTSeq-Count v.0.6.1

(93). Differences in gene expression were identified using a generalized linear model method in

edgeR v.3.20.1 (94), with genes showing a $\log_2$-fold change >1 or < -1 and with a Benjamini-

Hochberg corrected FDR<0.05 defined as differentially expressed. Human homologs of

differentially-expressed genes in flies were identified using DIOPT v7.0. Biological pathways

and processes affected by downregulation of homologs of 16p12.1 genes, defined as significant

enrichments of Gene Ontology (GO) terms (p<0.05, Fisher's exact test with Benjamini-

Hochberg multiple testing correction), were identified using PantherDB (95).

Pairwise knockdowns in the fly eye

To study genetic interactions in the fly eye, we generated recombinant stock lines for each

16p12.1 homolog by crossing RNAi lines with eye-specific *GMR-GAL4*, as detailed in **S1 File**.

Various factors including presence of balancers, chromosomal insertion of the shRNA, lethality

with *Elav-GAL4*, and severity of eye phenotypes with *GMR-GAL4* were considered to select

RNAi lines for generating recombinant lines. For example, for *CG14182*, we used the GD RNAi

line *CG14182^{GD2738}*, which showed milder eye phenotypes, in order to test a wider range of

potential interactions. We assessed genetic interactions between homologs of 16p12.1 genes with

each other as well as with homologs of "second-hits" identified in children with the 16p12.1

deletion, conserved neurodevelopmental genes, and select transcriptional targets. A total of 24

homologs of genes carrying "second-hits" were selected as disease-associated genes carrying

rare (ExAC frequency ≤1%) copy-number variants, loss-of-function (frameshift, stopgain or

splicing) mutations, or *de novo* or likely-pathogenic (Phred-like CADD ≥25) missense mutations

previously identified from exome sequencing and SNP microarrays in 15 affected children with

the 16p12.1 deletion and their family members (9, 96, 97). We also selected seven conserved

genes strongly associated with neurodevelopmental disorders (20, 54) and six genes with

previously described functional associations with individual 16p12.1 genes, such as

mitochondrial genes for *UQCRC2* (59) and *Myc* for *POLR3E* and *CDR2* (60, 61). We also tested

interactions of the 16p12.1 homologs with 25 differentially-expressed genes (or "transcriptome

targets") and 14 genes selected from enriched Gene Ontology groups identified from RNA

sequencing experiments (**S3 File**). Overall, we tested 212 pairwise gene interactions including 96

interactions with homologs of "second-hit" genes, 55 with neurodevelopmental genes, and 61

with transcriptome targets, using multiple RNAi, mutant or overexpression lines per gene when

available (**S2 Table**).

GMR-GAL4 recombinant lines for the homologs of 16p12.1 genes were crossed with

RNAi or mutant lines for the interacting genes to achieve simultaneous knockdown of two genes

in the eye. We also tested overexpression lines for specific genes that are functionally related to

16p12.1 homologs, including *Myc*, *Hsp23*, and *Hsp26*. Our previous assessment showed no

changes in phenotypic scores for recombinant lines crossed with *UAS-GFP* compared to crosses

with controls, demonstrating that the lines have adequate *GAL4* to bind to two independent *UAS-*

RNAi constructs (20).

To evaluate how simultaneous knockdown of interacting genes modulated the phenotype

of 16p12.1 homologs, *Flynotyper* scores from flies with double knockdowns were compared to

the scores from flies obtained from crosses between 16p12.1 recombinant lines and controls

carrying the same genetic background as the interacting gene. Significant enhancements or

suppressions of phenotypes of 16p12.1 homologs were identified using two-tailed Mann-

Whitney tests and Benjamini-Hochberg multiple testing correction (**S3 File**).

Analysis of genetic interactions using the multiplicative model

The nature of genetic interactions between pairs of gene knockdowns was determined using a

multiplicative model, which has been widely used for evaluating genetic interactions for

quantitative phenotypes in yeast, *Drosophila*, and *E. coli* models (55-58, 98). This model tests

whether the observed phenotypic effect of simultaneously knocking down two genes is different

from the expected product of effects due to knockdown of individual genes. We first normalized

Flynotyper scores for each individual line and pairwise knockdown to the *Flynotyper* scores

from the background-specific controls to obtain normalized phenotypic scores as "*percentage of*

control." We then calculated the expected effects as the product of the averages of normalized

scores for the two individual gene knockdowns, and compared the expected scores to the

normalized scores from pairwise knockdowns using a two-tailed one-sample Wilcoxon signed

ranked test with Benjamini-Hochberg correction. The distributions of the expected and observed

values for all individual pairs of genes and pairwise knockdowns are shown in **Figure S9**.

Pairwise knockdowns where the observed effects were significantly higher than expected were

categorized as negative, aggravating or synergistic genetic interactions, while those with

observed values significantly lower than expected were considered as positive or alleviating

interactions. Pairwise knockdowns where the observed effects were not significantly different

from the expected effects were considered as not interacting. The magnitude of a genetic

interaction was measured using "*interaction scores*", calculated as the $\log_2$ ratio between the

observed and expected values (**S4 File**). Positive interaction scores indicated negative or

aggravating genetic interactions, while negative interaction scores indicated positive or

alleviating genetic interactions. An interaction was considered to be validated when the observed

trend was reproduced by multiple fly lines when available. Interactions assessed with only one

fly line were considered as "potential negative", "potential positive" or "potential no interaction"

(**S3 File**). To compare interactions of 16p12.1 homologs to 16p11.2 homologs, we obtained

Flynotyper phenotypic scores from single and pairwise *GMR-GAL4*-mediated knockdown of

16p11.2 homologs, previously described in Iyer *et al.* (20).

Immunohistochemistry of the developing brain and eye discs in *Drosophila melanogaster*

Third instar larvae brain or eye discs were dissected in PBS and fixed in 4% paraformaldehyde in

PBT (0.3% Triton X-100 in PBS), followed by three washes with PBT. Preparations were

blocked for one hour in blocking buffer (5% FBS or 1% BSA in 0.3% PBT), followed by

incubation overnight at 4°C with the primary antibody. We assessed for markers of proliferation

using mouse anti-pH3 (S10) (1:100; 9706, Cell Signaling Technology, Danvers, MA, USA) and

apoptosis using rabbit anti-Dcp-1 (Asp216) (1:100, 9578, Cell Signaling). Secondary antibody

incubation was performed using Alexa fluor 647 goat anti-mouse (1:100, Invitrogen, Carlsbad,

CA, USA) and Alexa fluor 568 goat anti-rabbit (1:100, Invitrogen) for 2 hours at 25°C, followed

by three washes with PBT. Tissues were mounted in Prolong Gold antifade reagent with DAPI

(Thermo Fisher Scientific, P36930) prior to imaging. Z-stacks of brain lobe or eye discs were

acquired every 4μm with a 40X air objective with 1.2X magnification using an Olympus

Fluoview FV1000 laser scanning confocal microscope (Olympus America, Lake Success, NY).

Image analysis was performed using Fiji (90). The number of cells undergoing proliferation or

apoptosis were quantified throughout the brain lobe, or posterior to the morphogenetic furrow in

the developing eye discs. The total number of Dcp-1 positive cells in larval brain and eye discs,

as well as pH3 cells in the eye discs, were manually counted from the maximum projections. The

total number of pH3 positive cells in the larval brain were quantified using the MaxEntropy

automated thresholding algorithm per slice, followed by counting the number of particles larger

than 1.5 μm. Differences in the number of positive pH3 or Dcp-1 cells were compared with

appropriate controls using two-tailed Mann-Whitney tests.

***Xenopus laevis* embryos**

Eggs collected from female *X. laevis* frogs were fertilized *in vitro*, dejellied, and cultured

following standard methods (**16, 99**). Embryos received injections of exogenous mRNAs or

antisense oligonucleotide strategies at the two- or four-cell stage, using four total injections

performed in 0.1X MMR media containing 5% Ficoll. Embryos were staged according to

Nieuwkoop and Faber (**100**).

***X. laevis* gene knockdown and rescue**

Morpholinos (MOs) were targeted to early splice sites of *X. laevis mosmo*, *polr3e*, *uqcrc2*, *cdr2*,

and *setd5*, or standard control MO (**S7 Table**), purchased from Gene Tools (Philomath, OR). In

knockdown experiments, all MOs were injected at either the 2-cell or 4-cell stage, with embryos

receiving injections two or four times total. *mosmo* and control MOs were injected at 12

ng/embryo for partial and 20 ng/embryo for stronger knockdown; *polr3e* and control MOs were

injected at 10 ng/embryo for partial and 20 ng/embryo for stronger; *uqcrc2* and control MOs

were injected at 35 ng/embryo for partial and 50 ng/embryo for stronger; *cdr2* and control MOs

were injected at 10 ng/embryo for partial and 20ng for stronger knockdown; and *setd5* and

control MOs were injected at 10 ng/embryo for partial knockdown. All double knockdown

experiments were performed with partial knockdown to avoid potential lethality.

Splice site MOs were validated using RT-PCR. Total RNA was extracted using Trizol

reagent, followed by chloroform extraction and ethanol precipitation from 2-day old embryos

injected with increasing concentrations of MO targeted to each 16p12.1 homolog, respectively.

cDNA was synthesized using SuperScript II Reverse Transcriptase (Invitrogen, Carlsbad, CA,

USA). PCR was performed in a Mastercycler using HotStarTaq DNA Polymerase (Qiagen,

Germantown, MD, USA) following manufacturer instructions. PCR was performed using

primers with sequences detailed in **S6 Table**. RT-PCR was performed in triplicate, and band

intensity was measured using the densitometry function in ImageJ (101) and normalized to the

uninjected control mean relative to the housekeeping control *odc1*. Phenotypes were rescued

with exogenous mRNAs co-injected with their corresponding MO strategies. *X. laevis* ORF for

mosmo was purchased from the European *Xenopus* Resource Center (EXRC, Portsmouth, UK)

and gateway-cloned into pCSF107mT-GATEWAY-3'GFP destination vector. Constructs used

were *mosmo*-GFP, and GFP in pCS2+. In rescue experiments, MOs of the same amount used as

for the knockdown of each homolog were co-injected along with mRNA (1000pg/embryo for

mosmo-GFP) in the same injection solution.

Quantifying craniofacial shape and size of *X. laevis* embryos

The protocol for quantifying craniofacial shape and size was adapted from Kennedy and

Dickinson (102). Embryos at stage 42 were fixed overnight in 4% paraformaldehyde in PBS. A

razor blade was used to make a cut bisecting the gut to isolate the head. Isolated heads were

mounted in small holes in a clay-lined dish containing PBS. Frontal and lateral view images were

taken using a Zeiss AxioCam MRc attached to a Zeiss SteREO Discovery.V8 light microscope

(Zeiss, Thornwood, NY, USA). ImageJ software (101) was used to perform craniofacial

measurements, including: 1) facial width, which is the distance between the eyes; 2) face height,

which is the distance between the top of the eyes and the top of the cement gland at the midline;

3) dorsal mouth angle, which is the angle created by drawing lines from the center of one eye, to

the dorsal midline of the mouth, to the center of the opposite eye; and 4) midface area, which is

the area measured from the top of the eyes to the cement gland encircling the edges of both eyes.

For all facial measurements, two-tailed student's t-tests were performed between knockdown

embryos and control MO-injected embryos with the same amount of morpholino.

Neural tube explants, imaging, and analysis

Embryos were injected with either control MO or 16p12.1 homolog-specific MO at the 2-4 cell

stage, and culturing of *Xenopus* embryonic neural tube explants from stage 20-22 embryos were

performed as previously described (99). For axon outgrowth analysis, phase contrast images of

axons were collected on a Zeiss Axio Observer inverted motorized microscope with a Zeiss

20X/0.5 Plan Apo phase objective (Zeiss, Thornwood, NY, USA). Raw images were analyzed by

manually tracing the length of individual axons using the NeuronJ plug-in in ImageJ (103). All

experiments were performed on multiple independent occasions to ensure reproducibility. Axon

outgrowth data were normalized to controls from the same experiment to account for day-to-day

fluctuations. Statistical differences were identified between knockdown embryos and control

MO-injected embryos with same amounts of morpholino using two-tailed student's t-tests.

Immunostaining for brain morphology, imaging, and analysis

For brain morphology analysis, half embryo KDs were performed at the two-cell stage. *X. laevis*

embryos were unilaterally injected two times with either control MO or 16p12.1 homolog-

specific MO and a GFP mRNA construct (300pg/embryo). The other blastomere was left

uninjected. Embryos were raised in 0.1X MMR through neurulation, and then sorted based on

left/right fluorescence. Stage 47 embryos were fixed in 4% paraformaldehyde diluted in PBS for

one hour, rinsed in PBS, and gutted to reduce autofluorescence. Embryos were processed for

immunoreactivity by incubating in 3% bovine serum albumin and 1% Triton-X 100 in PBS for

two hours, and then incubated in anti-acetylated tubulin (1:700, T7451SigmaAldrich, St. Louis

MO, USA) and goat anti-mouse Alexa Fluor 488 conjugate secondary antibody (1:1000,

Invitrogen, Carlsbad, CA, USA). Embryos were rinsed in 1% Tween-20 in PBS and imaged in

PBS. Removal of the skin dorsal to the brain was performed if the brain was not clearly visible

due to pigment.

Images were taken at 3.2X magnification using a Zeiss AxioCam MRc attached to a

Zeiss SteREO Discovery.V8 light microscope (Zeiss, Thornwood, NY, USA). Images were

processed in ImageJ (101). The areas of the forebrain and midbrain were determined from raw

images using the polygon area function in ImageJ. Brain sizes were quantified by taking the ratio

of forebrain and midbrain areas between the injected side versus the uninjected side for each

sample. All experiments were performed on at least three independent occasions to ensure

reproducibility, and data shown represent findings from multiple replicates. Statistical

differences were identified between knockdown embryos and control MO injected embryos with

the same amount of morpholino using two-tailed student's t-tests.

Western blot for cell proliferation

Embryos at stage 20 to 22 were lysed in buffer (50mM Tris pH 7.5, 1% NP40, 150mM NaCl,

1mM PMSF, 0.5 mM EDTA), supplemented with cOmplete™ Mini EDTA-free Protease

Inhibitor Cocktail (Sigma-Aldrich) and PhosSTOP Phosphatase Inhibitor Cocktail (Sigma-

Aldrich). Blotting was carried out using rabbit polyclonal antibody to Phospho-Histone H3

(Ser10) (1:500, PA5-17869, Invitrogen), with mouse anti-beta actin (1:2500, ab8224, Abcam,

Cambridge, MA, USA) as a loading control. Bands were detected by chemiluminescence using

Amersham ECL Western blot reagent (GE Healthcare Bio-Sciences, Pittsburgh, PA). Band

intensities were quantified by densitometry in ImageJ and normalized to the control mean

relative to β-actin.

Connectivity of 16p12.1 and 16p11.2 deletion genes in a human brain-specific interaction

network

We examined the connectivity of human 16p12.1 and 16p11.2 deletion genes in the context of a

human brain-specific gene interaction network that was previously built using a Bayesian

classifier trained on gene co-expression datasets (71, 72). As the classifier assigned weighted

probabilities for interactions between all possible pairs of genes in the genome, we first built a

network that only contained the top 0.5% of all pairwise interactions (predicted weights >2.0).

Within this network, we identified the shortest paths between 25/27 genes in the 16p11.2

deletion or 6/7 genes in the 16p12.1 region that were present in the network and all other protein-

coding genes in the genome, using the inverse of the probabilities as weights for each interaction.

For each shortest path, we calculated the overall length as a measure of connectivity between the

two genes, and also identified the connector genes located within the shortest path (**S5 File**). All

network analyses were performed using the NetworkX v.2.4 Python package (104). We

compared the average connectivity of the six 16p12.1 genes to the 25 16p11.2 genes, as well as

the predicted pathogenicity of connector genes for 16p11.2 and 16p12.1 interactions, using two-

tailed Mann-Whitney tests. To account for a larger number of genes in the 16p11.2 deletion, we

also compared the median connectivity of 16p12.1 genes to the distribution of median

connectivity values for 1000 simulations of six random genes from the 16p11.2 region. We

further compared the median connectivity of 16p12.1 genes to the connectivity of 1000 random

sets of six genes in the network. Finally, to compare these values to random sets of contiguous

genes in the genome, we calculated the median connectivity among 1000 randomly-selected

contiguous blocks of six and 25 genes, excluding blocks containing genes within canonical CNV

regions (11). For each comparison with simulated gene sets, we calculated one-tailed z-scores

and percentile values for the observed median connectivity values (**S6 File**).

Statistical analyses

All statistical analyses of functional data were performed using R v.3.4.2 (R Foundation for

Statistical Computing, Vienna, Austria). Details for each statistical test, including sample size, p-

values with and without multiple testing correction, confidence intervals, test statistics, and

ANOVA degrees of freedom, are provided in **S6 File**.

Data access

Gene expression data for the *Drosophila* RNAi knockdown of homologs of 16p12.1 genes and
controls have been submitted to the NCBI Gene Expression Omnibus (GEO;
https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE151330, and the raw RNA
Sequencing files are deposited in the SRA (Sequence Read Archive) with BioProject database
(https://www.ncbi.nlm.nih.gov/bioproject/) accession number PRJNA635495. Source code for
the RNA-Sequencing and network analysis is available on the Girirajan lab GitHub page at
https://github.com/girirajanlab/16p12_fly_project.

Acknowledgments

This work was supported by NIH R01GM121907 and resources from the Huck Institutes of the
Life Sciences to S.G, Fulbright Commission Uruguay-ANII to L.P, NIH T32-GM102057 to
M.J., and NIH R01GM085115 to M.M.R.

Disclosure declaration

Authors disclose no conflict of interest.

References

1.   Girirajan S, Campbell CD, Eichler EE. Human copy number variation and complex genetic disease. Annu Rev Genet. 2011;45:203-26.

2.   Wilfert AB, Sulovari A, Turner TN, Coe BP, Eichler EE. Recurrent de novo mutations in neurodevelopmental disorders: properties and clinical implications. Genome Med. 2017;9(1):101.

3.   Weiss LA, Shen Y, Korn JM, Arking DE, Miller DT, Fossdal R, et al. Association between microdeletion and microduplication at 16p11.2 and autism. N Engl J Med. 2008;358(7):667-75.

4.   Zufferey F, Sherr EH, Beckmann ND, Hanson E, Maillard AM, Hippolyte L, et al. A 600 kb deletion syndrome at 16p11. 2 leads to energy imbalance and neuropsychiatric disorders. Journal of medical genetics. 2012;49(10):660-8.

5.   Mulle JG. The 3q29 deletion confers >40-fold increase in risk for schizophrenia. Mol Psychiatry. 2015;20(9):1028-9.

6.   Helbig I, Mefford HC, Sharp AJ, Guipponi M, Fichera M, Franke A, et al. 15q13.3 microdeletions increase risk of idiopathic generalized epilepsy. Nat Genet. 2009;41(2):160-2.

7.   Girirajan S, Eichler EE. Phenotypic variability and genetic susceptibility to genomic disorders. Human molecular genetics. 2010;19(R2):R176-R87.

8.   Girirajan S, Rosenfeld JA, Cooper GM, Antonacci F, Siswara P, Itsara A, et al. A recurrent 16p12. 1 microdeletion supports a two-hit model for severe developmental delay. Nature genetics. 2010;42(3):203-9.

9.   Pizzo L, Jensen M, Polyak A, Rosenfeld JA, Mannik K, Krishnan A, et al. Rare variants in the genetic background modulate cognitive and developmental phenotypes in individuals carrying disease-associated variants. Genet Med. 2018.

10.   Stefansson H, Meyer-Lindenberg A, Steinberg S, Magnusdottir B, Morgen K, Arnarsdottir S, et al. CNVs conferring risk of autism or schizophrenia affect cognition in controls. Nature. 2014;505(7483):361-6.

11.   Girirajan S, Rosenfeld JA, Coe BP, Parikh S, Friedman N, Goldstein A, et al. Phenotypic heterogeneity of genomic disorders and rare copy-number variants. New England Journal of Medicine. 2012;367(14):1321-31.

12.   Gatto CL, Pereira D, Broadie K. GABAergic circuit dysfunction in the Drosophila Fragile X syndrome model. Neurobiol Dis. 2014;65:142-59.

13.   Jumbo-Lucioni PP, Parkinson WM, Kopke DL, Broadie K. Coordinated movement, neuromuscular synaptogenesis and trans-synaptic signaling defects in Drosophila galactosemia models. Hum Mol Genet. 2016;25(17):3699-714.

14.   Sears JC, Broadie K. Fragile X Mental Retardation Protein Regulates Activity-Dependent Membrane Trafficking and Trans-Synaptic Signaling Mediating Synaptic Remodeling. Front Mol Neurosci. 2017;10:440.

15.   Grossman TR, Gamliel A, Wessells RJ, Taghli-Lamallem O, Jepsen K, Ocorr K, et al. Over-expression of DSCAM and COL6A2 cooperatively generates congenital heart defects. PLoS Genet. 2011;7(11):e1002344.

16.   Chen SX, Tari PK, She K, Haas K. Neurexin-neuroligin cell adhesion complexes contribute to synaptotropic dendritogenesis via growth stabilization mechanisms in vivo. Neuron. 2010;67(6):967-83.

17.   Lewis BB, Wester MR, Miller LE, Nagarkar MD, Johnson MB, Saha MS. Cloning and characterization of voltage-gated calcium channel alpha1 subunits in Xenopus laevis during development. Dev Dyn. 2009;238(11):2891-902.

18.   Ishimaru H, Kamboj R, Ambrosini A, Henley JM, Soloviev MM, Sudan H, et al. A unitary non-NMDA receptor short subunit from Xenopus: DNA cloning and expression. Receptors Channels. 1996;4(1):31-49.

19.   Ueno S, Kono R, Iwao Y. PTEN is required for the normal progression of gastrulation by repressing cell proliferation after MBT in Xenopus embryos. Dev Biol. 2006;297(1):274-83.

20.   Iyer J, Singh MD, Jensen M, Patel P, Pizzo L, Huber E, et al. Pervasive genetic interactions
modulate neurodevelopmental defects of the autism-associated 16p11.2 deletion in Drosophila
melanogaster. Nat Commun. 2018;9(1):2548.
21.   Singh MD, Jensen M, Lasser M, Huber E, Yusuff T, Pizzo L, et al. NCBP2 modulates
neurodevelopmental defects of the 3q29 deletion in Drosophila and Xenopus laevis models. PLoS Genet.
2020;16(2):e1008590.
22.   McCammon JM, Blaker-Lee A, Chen X, Sive H. The 16p11.2 homologs fam57ba and doc2a
generate certain brain and body phenotypes. Hum Mol Genet. 2017;26(19):3699-712.
23.   Qiu Y, Arbogast T, Lorenzo SM, Li H, Tang SC, Richardson E, et al. Oligogenic Effects of
16p11.2 Copy-Number Variation on Craniofacial Development. Cell Rep. 2019;28(13):3320-8 e4.
24.   Hu Y, Flockhart I, Vinayagam A, Bergwitz C, Berger B, Perrimon N, et al. An integrative
approach to ortholog prediction for disease-focused and other functional studies. BMC Bioinformatics.
2011;12:357.
25.   Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates and generating
dominant phenotypes. Development. 1993;118(2):401-15.
26.   Strigini M, Cohen SM. A Hedgehog activity gradient contributes to AP axial patterning of the
Drosophila wing. Development. 1997;124(22):4697-705.
27.   Yan SJ, Gu Y, Li WX, Fleming RJ. Multiple signaling pathways and a selector protein
sequentially regulate Drosophila wing development. Development. 2004;131(2):285-98.
28.   Yusuff T, Jensen M, Yennawar S, Pizzo L, Karthikeyan S, Gould DJ, et al. Drosophila models of
pathogenic copy-number variant genes show global and non-neuronal defects during development. PLoS
Genet. 2020;16(6):e1008792.
29.   Callan MA, Cabernard C, Heck J, Luois S, Doe CQ, Zarnescu DC. Fragile X protein controls
neural stem cell proliferation in the Drosophila brain. Hum Mol Genet. 2010;19(15):3068-79.
30.   Kwon CH, Luikart BW, Powell CM, Zhou J, Matheny SA, Zhang W, et al. Pten regulates
neuronal arborization and social interaction in mice. Neuron. 2006;50(3):377-88.
31.   Lee A, Li W, Xu K, Bogert BA, Su K, Gao FB. Control of dendritic development by the
Drosophila fragile X-related gene involves the small GTPase Rac1. Development. 2003;130(22):5543-52.
32.   Parker L, Padilla M, Du Y, Dong K, Tanouye MA. Drosophila as a model for epilepsy: bss is a
gain-of-function mutation in the para sodium channel gene that leads to seizures. Genetics.
2011;187(2):523-34.
33.   Rujano MA, Sanchez-Pulido L, Pennetier C, le Dez G, Basto R. The microcephaly protein Asp
regulates neuroepithelium morphogenesis by controlling the spatial distribution of myosin II. Nat Cell
Biol. 2013;15(11):1294-306.
34.   Stuss DP, Boyd JD, Levin DB, Delaney KR. MeCP2 mutation results in compartment-specific
reductions in dendritic branching and spine density in layer 5 motor cortical neurons of YFP-H mice.
PLoS One. 2012;7(3):e31896.
35.   Wang HD, Kazemi-Esfarjani P, Benzer S. Multiple-stress analysis for isolation of Drosophila
longevity genes. Proc Natl Acad Sci U S A. 2004;101(34):12610-5.
36.   Orr WC, Sohal RS. Extension of life-span by overexpression of superoxide dismutase and
catalase in Drosophila melanogaster. Science. 1994;263(5150):1128-30.
37.   Copeland JM, Cho J, Lo T, Jr., Hur JH, Bahadorani S, Arabyan T, et al. Extension of Drosophila
life span by RNAi of the mitochondrial respiratory chain. Curr Biol. 2009;19(19):1591-8.
38.   Mahmoudi S, Xu L, Brunet A. Turning back time with emerging rejuvenation strategies. Nat Cell
Biol. 2019;21(1):32-43.
39.   Matthews BJ, Kim ME, Flanagan JJ, Hattori D, Clemens JC, Zipursky SL, et al. Dendrite self-
avoidance is controlled by Dscam. Cell. 2007;129(3):593-604.
40.   Soba P, Zhu S, Emoto K, Younger S, Yang SJ, Yu HH, et al. Drosophila sensory neurons require
Dscam for dendritic self-avoidance and proper dendritic field organization. Neuron. 2007;54(3):403-16.
41.   King KL, Cidlowski JA. Cell cycle and apoptosis: common pathways to life and death. J Cell
Biochem. 1995;58(2):175-80.

42.   King KL, Cidlowski JA. Cell cycle regulation and apoptosis. Annu Rev Physiol. 1998;60:601-17.
43.   Hunt P, Gulisano M, Cook M, Sham MH, Faiella A, Wilkinson D, et al. A distinct Hox code for
the branchial region of the vertebrate head. Nature. 1991;353(6347):861-4.
44.   Hunt P, Whiting J, Muchamore I, Marshall H, Krumlauf R. Homeobox genes and models for
patterning the hindbrain and branchial arches. Dev Suppl. 1991;1:187-96.
45.   Lasser M, Pratt B, Monahan C, Kim SW, Lowery LA. The Many Faces of Xenopus: Xenopus
laevis as a Model System to Study Wolf-Hirschhorn Syndrome. Front Physiol. 2019;10:817.
46.   Le Lievre CS, Le Douarin NM. Mesenchymal derivatives of the neural crest: analysis of
chimaeric quail and chick embryos. J Embryol Exp Morphol. 1975;34(1):125-54.
47.   Lumsden A, Sprawson N, Graham A. Segmental origin and migration of neural crest cells in the
hindbrain region of the chick embryo. Development. 1991;113(4):1281-91.
48.   Mills A, Bearce E, Cella R, Kim SW, Selig M, Lee S, et al. Wolf-Hirschhorn Syndrome-
Associated Genes Are Enriched in Motile Neural Crest Cells and Affect Craniofacial Development in
Xenopus laevis. Front Physiol. 2019;10:431.
49.   Goldberg JL. How does an axon grow? Genes Dev. 2003;17(8):941-58.
50.   Jensen M, Girirajan S. An interaction-based model for neuropsychiatric features of copy-number
variants. PLoS Genet. 2019;15(1):e1007879.
51.   Thaker HM, Kankel DR. Mosaic analysis gives an estimate of the extent of genomic involvement
in the development of the visual system in Drosophila melanogaster. Genetics. 1992;131(4):883-94.
52.   Branco J, Al-Ramahi I, Ukani L, Perez AM, Fernandez-Funez P, Rincon-Limas D, et al.
Comparative analysis of genetic modifiers in Drosophila points to common and distinct mechanisms of
pathogenesis among polyglutamine diseases. Hum Mol Genet. 2008;17(3):376-90.
53.   Cziko AM, McCann CT, Howlett IC, Barbee SA, Duncan RP, Luedemann R, et al. Genetic
modifiers of dFMR1 encode RNA granule components in Drosophila. Genetics. 2009;182(4):1051-60.
54.   Iyer J, Wang Q, Le T, Pizzo L, Gronke S, Ambegaokar SS, et al. Quantitative Assessment of Eye
Phenotypes for Functional Genetic Studies Using Drosophila melanogaster. G3 (Bethesda).
2016;6(5):1427-37.
55.   Baryshnikova A, Costanzo M, Kim Y, Ding H, Koh J, Toufighi K, et al. Quantitative analysis of
fitness and genetic interactions in yeast on a genome scale. Nat Methods. 2010;7(12):1017-24.
56.   Dixon SJ, Costanzo M, Baryshnikova A, Andrews B, Boone C. Systematic mapping of genetic
interaction networks. Annu Rev Genet. 2009;43:601-25.
57.   Horn T, Sandmann T, Fischer B, Axelsson E, Huber W, Boutros M. Mapping of signaling
networks through synthetic genetic interaction analysis by RNAi. Nat Methods. 2011;8(4):341-6.
58.   Jonikas MC, Collins SR, Denic V, Oh E, Quan EM, Schmid V, et al. Comprehensive
characterization of genes required for protein folding in the endoplasmic reticulum. Science.
2009;323(5922):1693-7.
59.   Duncan AM, Ozawa T, Suzuki H, Rozen R. Assignment of the gene for the core protein II
(UQCRC2) subunit of the mitochondrial cytochrome bc1 complex to human chromosome 16p12.
Genomics. 1993;18(2):455-6.
60.   Gomez-Roman N, Grandori C, Eisenman RN, White RJ. Direct activation of RNA polymerase III
transcription by c-Myc. Nature. 2003;421(6920):290-4.
61.   O'Donovan KJ, Diedler J, Couture GC, Fak JJ, Darnell RB. The onconeural antigen cdr2 is a
novel APC/C target that acts in mitosis to regulate c-myc target genes in mammalian tumor cells. PLoS
One. 2010;5(4):e10045.
62.   Hartman JLt, Garvik B, Hartwell L. Principles for the buffering of genetic variation. Science.
2001;291(5506):1001-4.
63.   Queitsch C, Carlson KD, Girirajan S. Lessons from model organisms: phenotypic robustness and
missing heritability in complex disease. PLoS Genet. 2012;8(11):e1003041.
64.   Grozeva D, Carss K, Spasic-Boskovic O, Parker MJ, Archer H, Firth HV, et al. De novo loss-of-
function mutations in SETD5, encoding a methyltransferase in a 3p25 microdeletion syndrome critical
region, cause intellectual disability. Am J Hum Genet. 2014;94(4):618-24.

65.   Hu P, Wu S, Sun Y, Yuan CC, Kobayashi R, Myers MP, et al. Characterization of human RNA
polymerase III identifies orthologues for Saccharomyces cerevisiae RNA polymerase III subunits. Mol
Cell Biol. 2002;22(22):8044-55.
66.   Pusapati GV, Kong JH, Patel BB, Krishnan A, Sagner A, Kinnebrew M, et al. CRISPR Screens
Uncover Genes that Regulate Target Cell Sensitivity to the Morphogen Sonic Hedgehog. Dev Cell.
2018;44(2):271.
67.   McCamphill PK, Farah CA, Anadolu MN, Hoque S, Sossin WS. Bidirectional regulation of eEF2
phosphorylation controls synaptic plasticity by decoding neuronal activity patterns. J Neurosci.
2015;35(10):4403-17.
68.   Gildish I, Manor D, David O, Sharma V, Williams D, Agarwala U, et al. Impaired associative
taste learning and abnormal brain activation in kinase-defective eEF2K mice. Learn Mem.
2012;19(3):116-25.
69.   Zhang P, Riazy M, Gold M, Tsai SH, McNagny K, Proud C, et al. Impairing eukaryotic
elongation factor 2 kinase activity decreases atherosclerotic plaque formation. Can J Cardiol.
2014;30(12):1684-8.
70.   Monteggia LM, Gideons E, Kavalali ET. The role of eukaryotic elongation factor 2 kinase in
rapid antidepressant action of ketamine. Biol Psychiatry. 2013;73(12):1199-203.
71.   Greene CS, Krishnan A, Wong AK, Ricciotti E, Zelaya RA, Himmelstein DS, et al.
Understanding multicellular function and disease with human tissue-specific networks. Nat Genet.
2015;47(6):569-76.
72.   Krishnan A, Zhang R, Yao V, Theesfeld CL, Wong AK, Tadych A, et al. Genome-wide
prediction and functional characterization of the genetic basis of autism spectrum disorder. Nat Neurosci.
2016;19(11):1454-62.
73.   Osipovich AB, Gangula R, Vianna PG, Magnuson MA. Setd5 is essential for mammalian
development and the co-transcriptional regulation of histone acetylation. Development.
2016;143(24):4595-607.
74.   Andrews T, Honti F, Pfundt R, de Leeuw N, Hehir-Kwa J, Vulto-van Silfhout A, et al. The
clustering of functionally related genes contributes to CNV-mediated disease. Genome Res.
2015;25(6):802-13.
75.   Iossifov I, Levy D, Allen J, Ye K, Ronemus M, Lee YH, et al. Low load for disruptive mutations
in autism genes and their biased transmission. Proc Natl Acad Sci U S A. 2015;112(41):E5600-7.
76.   Kury S, van Woerden GM, Besnard T, Proietti Onori M, Latypova X, Towne MC, et al. De Novo
Mutations in Protein Kinase Genes CAMK2A and CAMK2B Cause Intellectual Disability. Am J Hum
Genet. 2017;101(5):768-88.
77.   Krumm N, Turner TN, Baker C, Vives L, Mohajeri K, Witherspoon K, et al. Excess of rare,
inherited truncating mutations in autism. Nature genetics. 2015;47(6):582-8.
78.   Petrovski S, Wang Q, Heinzen EL, Allen AS, Goldstein DB. Genic intolerance to functional
variation and the interpretation of personal genomes. PLoS Genet. 2013;9(8):e1003709.
79.   Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, et al. Ensembl 2020.
Nucleic Acids Res. 2020;48(D1):D682-D8.
80.   Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and
PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res.
1997;25(17):3389-402.
81.   Chintapalli VR, Wang J, Dow JA. Using FlyAtlas to identify better Drosophila melanogaster
models of human disease. Nat Genet. 2007;39(6):715-20.
82.   Bowes JB, Snyder KA, Segerdell E, Jarabek CJ, Azam K, Zorn AM, et al. Xenbase: gene
expression and improved integration. Nucleic Acids Res. 2010;38(Database issue):D607-12.
83.   Xu D, Wang Y, Willecke R, Chen Z, Ding T, Bergmann A. The effector caspases drICE and dcp-
1 have partially overlapping functions in the apoptotic pathway in Drosophila. Cell Death Differ.
2006;13(10):1697-706.

84.   Schmid A, Hallermann S, Kittel RJ, Khorramshahi O, Frolich AM, Quentin C, et al. Activity-
dependent site-specific changes of glutamate receptor composition in vivo. Nat Neurosci.
2008;11(6):659-66.
85.   Xiao C, Mileva-Seitz V, Seroude L, Robertson RM. Targeting HSP70 to motoneurons protects
locomotor activity from hyperthermia in Drosophila. Dev Neurobiol. 2007;67(4):438-55.
86.   Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. Primer-BLAST: a tool to
design target-specific primers for polymerase chain reaction. BMC Bioinformatics. 2012;13:134.
87.   Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative
PCR and the 2(-Delta Delta C(T)) Method. Methods. 2001;25(4):402-8.
88.   Sun Y, Yolitz J, Wang C, Spangler E, Zhan M, Zou S. Aging studies in Drosophila melanogaster.
Methods Mol Biol. 2013;1048:77-93.
89.   Ganetzky B, Wu CF. Indirect Suppression Involving Behavioral Mutants with Altered Nerve
Excitability in DROSOPHILA MELANOGASTER. Genetics. 1982;100(4):597-614.
90.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.
91.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20.
92.   Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment
of transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol. 2013;14(4):R36.
93.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput
sequencing data. Bioinformatics. 2015;31(2):166-9.
94.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 2010;26(1):139-40.
95.   Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, et al. PANTHER: a
library of protein families and subfamilies indexed by function. Genome Res. 2003;13(9):2129-41.
96.   Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for
estimating the relative pathogenicity of human genetic variants. Nat Genet. 2014;46(3):310-5.
97.   Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, et al. Analysis of protein-
coding genetic variation in 60,706 humans. Nature. 2016;536(7616):285-91.
98.   Elena SF, Lenski RE. Test of synergistic interactions among deleterious mutations in bacteria.
Nature. 1997;390(6658):395-8.
99.   Lowery LA, Faris AE, Stout A, Van Vactor D. Neural Explant Cultures from Xenopus laevis. J
Vis Exp. 2012(68):e4232.
100.   Nieuwkoop PD FJ. Normal table of Xenopus laevis (Daudin) : a systematical and chronological
survey of the development from the fertilized egg till the end of metamorphosis: Garland Pub., 1994;
1994.
101.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nat
Methods. 2012;9(7):671-5.
102.   Kennedy AE, Dickinson AJ. Quantitative analysis of orofacial development and median clefts in
Xenopus laevis. Anat Rec (Hoboken). 2014;297(5):834-55.
103.   Popko J, Fernandes A, Brites D, Lanier LM. Automated analysis of NeuronJ tracing data.
Cytometry A. 2009;75(4):371-6.
104.   Hagberg AA SD, Swart PJ, editor Exploring network structure, dynamics, and function using
NetworkX.
7th Python in Science Conference SciPy 2008; 2008.
105.   D'Angelo D, Lebon S, Chen Q, Martin-Brevet S, Snyder LG, Hippolyte L, et al. Defining the
Effect of the 16p11.2 Duplication on Cognition, Behavior, and Medical Comorbidities. JAMA Psychiatry.
2016;73(1):20-30.
106.   Niarchou M, Chawner S, Doherty JL, Maillard AM, Jacquemont S, Chung WK, et al. Psychiatric
disorders in children with 16p11.2 deletion and duplication. Transl Psychiatry. 2019;9(1):8.
1157

 Figure Legends

Fig. 1. Strategy to evaluate the individual contributions of homologs of 16p12.1 genes and

their interactions with "second-hit" genes towards neurodevelopmental phenotypes. (A)

Ideogram of human chromosome 16 indicating the deleted region on UCSC genome build

GRCh37, hg19 (chr16:21,948,445-22,430,408) (also known as 16p12.2 deletion). Seven protein

coding genes are located within the 16p12.1 deletion region, including *UQCRC2*, *PDZD9*,

MOSMO, *VWA3A*, *EEF2K*, *POLR3E,* and *CDR2*. Four out of the seven genes are conserved in

both *Drosophila melanogaster* and *Xenopus laevis*. **(B)** We performed global and functional

domain-specific phenotypic assessment using RNAi lines and tissue-specific knockdown in

Drosophila, and morpholino-mediated whole embryo knockdown in *X. laevis*, to identify

individual contributions of 16p12.1 homologs towards different developmental and neuronal

features. We next evaluated the effect of pairwise knockdown of 16p12.1 homologs towards eye

phenotypes in *Drosophila*, and brain size and cellular proliferation defects in *X. laevis*. We

characterized 212 interactions between the 16p12.1 homologs and homologs of patient-specific

"second-hit" genes identified in children with the deletion, genes within conserved

neurodevelopmental pathways, and differentially-expressed genes identified from RNA-seq

analysis. We found that homologs of "second-hit" genes participate in complex genetic

interactions with 16p12.1 homologs to modulate neurodevelopmental and cellular phenotypes.

1176

Fig. 2. Multiple homologs of 16p12.1 genes contribute to neurodevelopmental defects in

***Drosophila melanogaster* and *X. laevis*.**

(A) Schematic showing multiple phenotypes affected by tissue-specific knockdown of individual

16p12.1 homologs in *Drosophila melanogaster*. Ubiquitous knockdown was achieved with *da-*

GAL4, eye-specific knockdown with *GMR-GAL4*, wing-specific knockdown with *bx^{MS1096}-*

GAL4, and nervous system-specific with *ppk-GAL4* or *Elav-GAL4*. See **S2A-C Figure** for details

on phenotypes observed for individual fly lines. (**B**) Nervous-system mediated knockdown using

Elav-GAL4 with overexpression of *Dicer2* at 25°C led to reduced lifespan with knockdown of

CG14182$^{GD2738\_2}$ (n=100, one-way repeat measures ANOVA with post-hoc pairwise t-test, days

6-61, $p<0.05$) and increased lifespan with knockdown of *UQCR-C2^{GD11238}* (n=120, days 51-81,

$p<0.05$). *Elav-GAL4* mediated knockdown of *Sin^{GD7027}* at RT without overexpression of *Dicer2*

led to reduced lifespan of adult flies (n=160, day 1-6, $p<0.05$). Data represented show mean ±

standard deviation of 4-8 independent groups of 20 flies for each line tested. (**C**) Nervous-system

mediated knockdown led to delayed pupariation time and larval lethality for *Sin^{GD7027}* (n=180,

one-way repeat measures ANOVA with post-hoc pairwise t-test, days 6-18, $p<0.05$) and partial

larval lethality for *CG14182$^{GD2738\_2}$* (n=120, days 7-10, $p<0.05$). Data represented show mean ±

standard deviation of 4-9 independent groups of 30 larvae for each line tested. (**D**) Knockdown

of 16p12.1 homologs in sensory class IV dendritic arborization neurons using *ppk-GAL4* with

overexpression of *Dicer2* showed reduced complexity of dendritic arbors (measured as sum of

intersections normalized to width) for *CG14182^{GD2738}* (n=12, two-tailed Mann-Whitney, *p=5.35

$\times 10^{-5}$). Scale bar represents 25 μm. (**E**) Third instar larvae with nervous system-specific

knockdown of 16p12.1 homologs showed reduced brain area for *CG14182$^{GD2738\_2}$* (n=15, two-

tailed Mann-Whitney, *p=0.047) and *Sin^{GD7027}* (n=17, *p= 0.001). (**F**) Developing third instar

larvae with knockdown of *CG14182$^{GD2738\_2}$* (n=15, two-tailed Mann-Whitney, *p=0.026) and

Sin^{GD7027} (n=10, *p= 9.74×10^{-4}) showed reduced number of phosphorylated Histone-3 (pH3)

positive cells in the brain lobe (green). Scale bar represents 50 μm. All control data for

Drosophila represents phenotypes observed for the GD VDRC control (Control^{GD}) crossed with

the indicated tissue-specific *GAL4* driver. (**G**) Schematic showing the phenotypes observed with

knockdown of 16p12.1 homologs in *X. laevis*. (**H**) Representative images of tadpoles injected

with control morpholino, indicating facial landmarks for face width (yellow) and orofacial area

(red), and tadpoles with knockdown of *polr3e* and *mosmo*. Knockdown of *cdr2* (n=54, two-tailed

student's t-test, *p=7.75 $\times 10^{-4}$), *polr3e* (n=37, *p=1.97 $\times 10^{-13}$) and *mosmo* (n=50, *p=1.36 $\times 10^{-}$

^{11}) led to decreased face width, while knockdown of *polr3e* (*p=3.29 $\times 10^{-16}$) and *mosmo*

(*p=1.47 $\times 10^{-8}$) led to decreased orofacial area. All measures were normalized to their respective

control injected with the same morpholino amount. Scale bar represents 500 μm. (**I**) Strong

knockdown of *mosmo* led to decreased axon length in neural tube explants (n=566, two-tailed

student's t-test, *p=7.40 $\times 10^{-12}$). All measures were normalized to their respective control

injected with the same morpholino amount. Representative schematic for axon length

measurements is shown on the left. (**J**) Representative images show forebrain (red on control

image) and midbrain (blue) areas of the side injected with morpholino (right, red asterisk), which

were normalized to the uninjected side (left). Strong knockdown of *mosmo* (n=67, two-tailed

student's t-test, *p<3.07$\times 10^{-13}$) and *polr3e* (n=48, *p<7.39$\times 10^{-4}$) led to decreased midbrain and

forebrain area of *X. laevis* tadpoles (stained with tubulin). Scale bar represents 500 μm. In all

cases, *X. laevis* data represents strong knockdown of the 16p12.1 homologs, except for *cdr2*,

which showed lethality and is represented with partial knockdown. All control data for *X. laevis*

represents controls injected with the highest amount of morpholino (50 ng, see **S5 Fig**). Boxplots

represent all data points with median, 25th and 75th percentiles, and red dotted lines indicate the

control median. Statistical details, including sample size, confidence intervals and p-values, are

provided in **S6 File**. A list of full genotypes for fly crosses used in these experiments is provided

in **S1 File**.

Fig. 3. Homologs of 16p12.1 genes contribute towards neurodevelopmental defects through

genetic interactions and combined independent effects.

(**A**) We generated eye-specific *GMR-GAL4* recombinant lines for the four 16p12.1 homologs to

test a total of twelve pairwise interactions for modulation of eye defects. (**B**) Representative

brightfield images of *Drosophila* adult eyes for recombinant lines of 16p12.1 homologs crossed

with RNAi lines for the other homologs, which show enhancement (Enh.) or suppression (Supp.)

of the phenotypes observed with crosses with control. Scale bar represents 100 μm. (**C**)

Simultaneous knockdown of $UQCR\text{-}C2^{GD11238}$ with $CG14182^{GD2738}$ (n=18, two-tailed Mann-

Whitney with Benjamini-Hochberg correction, *p=0.002) or Sin^{GD7027} (n=19, *p=0.023) led to a

significant enhancement in the eye phenotype (measured using *Flynotyper* scores) compared to

knockdown of $UQCR\text{-}C2^{GD11238}$ alone. Similarly, simultaneous knockdown of $CG14182^{GD2738}$

with Sin^{GD7027} (n=19, *p=0.021) enhanced the eye phenotype observed for $CG14182^{GD2738}$ alone.

Simultaneous knockdown of Cen^{GD9689} with $UQCR\text{-}C2^{GD11238}$ (n=20, *p=0.023) led to a milder

suppression of the eye phenotype compared to knockdown of Cen^{GD9689} alone. Double

knockdowns were compared to the recombinant lines of the 16p12.1 homologs crossed with

wild-type controls of the second 16p12.1 homolog. Note that only experiments with Control^{GD}

are represented here; see **S8 Fig** for results from other lines with KK and BL controls. (**D**) We

applied a multiplicative model to identify the nature of genetic interactions for the pairwise

knockdowns tested. The expected phenotype from simultaneous knockdown of homolog A and

homolog B, or when the combined effect indicates no genetic interaction (in blue), was

calculated as the product of the normalized phenotypic scores (i.e. percentage of control)

observed from knockdown of individual genes. Positive or alleviating genetic interactions were

identified for combinations where the observed phenotype was significantly milder than

expected (in green), while negative or aggravating interactions were identified when the

combined phenotypes were significantly more severe than expected (in red). One-sample

Wilcoxon signed rank tests with Benjamini-Hochberg correction for multiple testing were used

to identify significant interactions. (**E**) We generated double knockdowns of 16p12.1 homologs

in *X. laevis* models by co-injecting embryos with morpholinos of two homologs. All double

knockdown experiments were performed with partial knockdown of the genes, to avoid potential

lethality with stronger knockdown. (**F**) Representative images of tadpoles stained with anti-

tubulin show forebrain (red on control image) and midbrain (blue) areas of the side injected with

morpholino (right, red asterisk), which were normalized to the uninjected side (left).

Simultaneous knockdown of *polr3e* and *mosmo* led to decreased forebrain (n=36, two-tailed

student's t-test, *p=1.10×10^{-9}) and midbrain area (*p=1.98×10^{-7}), which showed no differences

compared to partial knockdown of *mosmo* alone. Control data represents control injected with

highest amount of morpholino (22ng). Scale bar represents 500 μm. (**G**) Representative western

blots show bands for phosphorylated histone-3 (pH3) and β-actin for the uninjected control,

knockdown of *polr3e*, knockdown of *mosmo*, and pairwise knockdown of *polr3e* and *mosmo*

(full western blots are shown in **S6 Fig**). Bar plot shows intensity of pH3 band normalized to β-

actin. Simultaneous knockdown of *polr3e* and *mosmo* does not lead to changes in the

proliferation defects observed with knockdown with *polr3e* alone. Boxplots represent all data

points with median, 25th and 75th percentiles, and red dotted lines indicate the control median.

Statistical details, including sample size, confidence intervals and p-values, are provided in **S6**

File. A list of full genotypes for fly crosses used in these experiments is provided in **S1 File**.

1271

Fig. 4. Homologs of 16p12.1 genes show complex interactions with conserved

neurodevelopmental genes and homologs of patient-specific "second-hit" genes.

(**A**) We evaluated how homologs of genes outside of the CNV region (*Gene B*), including genes

carrying "second-hit" variants in children with the 16p12.1 deletion, genes within conserved

neurodevelopmental pathways, and transcriptome targets, affect the phenotypes observed for

homologs of 16p12.1 genes. We crossed eye-specific recombinant lines for each homolog with a

total of 124 RNAi, mutant or overexpression lines for 76 interacting genes to test a total of 212

pairwise gene combinations. (**B**) Representative brightfield images of *Drosophila* adult eyes for

recombinant lines of 16p12.1 homologs crossed with background-specific controls (Control^{KK},,

also represented as C^{KK}) or RNAi lines for *kis* and *CG10465*, are shown as examples of genetic

interactions between the 16p12.1 homologs and homologs of neurodevelopmental genes. Bar

plots show normalized phenotypes (median ± interquartile range) for 16p12.1 recombinant lines

crossed with background-specific control or with RNAi lines for interacting genes. *Sin^{GD7027}*

negatively interacted with *kis^{KK100890}* and led to a more severe phenotype (two-tail one-sample

Wilcoxon signed rank test with Benjamini-Hochberg correction, n=11, *p=0.012, in red) than

expected (in blue) under a multiplicative model. Similarly, *CG10465^{KK105756}* negatively

interacted with *UQCR-C2^{GD11238}* (n=10, *p=0.024) and *CG14182^{GD2738}* (n=10, *p=0.015),

leading to more severe eye phenotypes than expected. Phenotypes are represented as percentage

of average, i.e. normalized to *Flynotyper* scores from control flies carrying the same genetic

background as the interacting gene. Scale bar represents 100 μm. (**C**) Heatmaps show interaction

scores calculated as the $\log_2$ ratio between the average of observed and expected phenotypic

scores. Positive scores represent negative aggravating genetic interactions (in red), while

negative scores represent positive alleviating interactions (in green). Grey boxes indicate

pairwise crosses that were not tested or were not validated by multiple lines. A complete list of

interaction scores is provided in **S4 File**. (**D**) Scatter plots depict interactions tested for 16p12.1

homologs. The plots show the average phenotypic score of the interacting gene on the x-axis,

and the average observed phenotypic score for the pairwise knockdown on the y-axis. The blue

line represents the expected phenotypic score of the pairwise knockdown calculated for each

16p12.1 homolog (value of first hit crossed with control, such as *UQCR-C2^{GD11238}* X Control^{BL}),

and all possible phenotypic scores (ranging from 0 to 60) of the interacting genes are represented

on the x-axis. All positive and negative (validated or potential) interactions are represented in

green and red, respectively, and fly lines of genes with no significant interactions are shown in

grey. Only lines from the BDSC stock center are represented here; **S17 Fig.** shows scatter plots

representing VDRC stock lines.

1306

Fig. 5. Homologs of patient-specific "second-hits" modulate phenotypes of 16p12.1

homologs.

(**A**) Representative pedigrees of families with 16p12.1 deletion (affected child in black, carrier

parent in grey) that were selected to study the effect of homologs (represented within

parenthesis) of genes carrying "second-hits" towards phenotypes of homologs of 16p12.1 genes.

(**B**) Observed phenotypic changes of 16p12.1 homologs by patient-specific "second-hit"

homologs. Plots show the changes in *Flynotyper* scores (mean ± s.d.) for *GMR-GAL4* control

(grey) or recombinant lines of 16p12.1 homologs crossed with either background-specific control

line (left) or with "second-hit" homologs (right). We note that represented changes in *Flynotyper*

scores for *Cen^{GD9689}/Nrx-1$^{GD14451\_2}$*, *UQCR-C2^{GD11238}/Zasp52^{HMJ22168}*, and

Sin^{GD7027}/Zasp52^{HMJ22168} were not validated with multiple RNAi lines for the "second-hit"

homolog. *Flynotyper* values for all the tested pairwise knockdowns are shown in **S11-S14 Fig**

and validated enhancements and suppressions (using Mann-Whitney tests) are shown in **S15-S16**

Fig. (**C**) Representative brightfield adult eye images for pairwise knockdowns that enhanced

(Enh.) or suppressed (Supp.) phenotypes of 16p12.1 homologs are shown. Scale bar represents

100 μm. (**D**) Examples of genetic interactions identified between the 16p12.1 homologs and

homologs of patient-specific "second-hit" genes using the multiplicative model. Bar plots show

normalized phenotypes (median ± interquartile range) for the 16p12.1 recombinant lines crossed

with background-specific controls (Control^{GD} or Control^{BL}, also represented as C^{GD}, C^{BL},

respectively) or with RNAi lines for *Nrx-1$^{GD14451\_2}$, Cep135$^{GD6121\_2}$, Zasp52^{HMJ22168}, shot^{HMJ23381},*

Rpn2^{HMS0533}, Dhc98^{DMB03402}. Sin^{GD7027} negatively interacted with *Cep135$^{GD6121\_2}$* and led to a

more aggravating phenotype (two-tail one-sample Wilcoxon signed rank test with Benjamini-

Hochberg correction, n=10, *p=0.012, in red) than expected (in blue) under a multiplicative

model, while other examples of pairwise knockdowns with homologs "second-hit" genes shown

here led to positive genetic interactions (*p<0.05, in green). Details of number of homologs, fly

lines and crosses, as well as a list of full genotypes for all interaction experiments are provided in

S1 File. Statistical details, including sample size, confidence intervals, and p-values, are

provided in **S6 File**.

1335

Fig. 6. Homolog of *SETD5* synergistically interacts with homolog of *MOSMO* to modify

neurodevelopmental defects.

(**A**) Pedigree of a family with 16p12.1 deletion, with the proband also carrying a *de novo*

pathogenic mutation in *SETD5*. Representative brightfield adult eye images for control and

GMR-GAL4 knockdown of *CG14182^{GD2738}, upSET^{HMC03177}*, and *CG14182^{GD2738}/upSET^{HMC03177}*

are shown. Data show a negative genetic interaction with simultaneous knockdown of

$CG14182^{GD2738}$ and $upSET^{HMC03177}$. Bar plots show normalized phenotypes (median ±

interquartile range) for recombinant lines of $CG14182^{GD2738}$ and Sin^{GD7027} crossed with

background-specific control (Control^{BL}, also represented as C^{BL}) or $upSET^{HMC03177}$. An

aggravating phenotype is observed with $CG14182^{GD2738}$/$upSET^{HMC03177}$ (two-tailed one-sample

Wilcoxon signed rank test with Benjamini-Hochberg correction, n=9, *p=0.018, in red) than

expected (in blue). (**B**) Representative confocal images of third instar larval eye discs stained

with anti-phosphorylated histone-3 (pH3, green) or anti-Dcp-1 (red), markers of cellular

proliferation and apoptosis, respectively. Positive pH3 or Dcp-1 cells were quantified posterior to

the morphogenetic furrow, indicated by white boxes in left panels. Double knockdown of

$CG14182^{GD2738}$/$upSET^{HMC03177}$ led to increased pH3 (n=17, two-tailed Mann-Whitney, *p=

0.046) and Dcp-1 (n=19, *p=0.006) positive cells compared to knockdown of $CG14182^{GD2738}$

alone. The double knockdown also led to increased Dcp-1 positive cells compared to knockdown

of $upSET^{HMC03177}$ alone (*p= 2.19×10^{5}). Scale bar represents 50 μm. (**C**) Representative images

of tadpoles stained with anti-tubulin show forebrain (red on control image) and midbrain (blue)

areas of the side injected with morpholino (right, red asterisk), which were normalized to the

uninjected side (left). Partial knockdown of *mosmo* with *setd5* led to a reduction in the midbrain

area compared to the knockdown of *mosmo* alone (n=16, two-tailed student's t-test, *p= 0.047).

Control data represents control injected with highest amount of morpholino (22ng). Scale bar

represents 500 μm (**D**) Normalized axon length of *X. laevis* tadpoles with simultaneous

knockdown of *mosmo* and *setd5* led to a significant reduction in axon length that was not

observed with partial knockdown of *mosmo* (n=438, two-tailed student's t-test, *p= 3.34×10^{-6})

or *setd5* (*p=1.86×10^{-9}). All measures were normalized to their respective controls injected with

the same morpholino amount (See **S20 Fig**). Control data represents controls injected with

highest amount of morpholino (22ng). All double knockdown experiments were performed with

partial knockdown of the genes, to avoid potential lethality with stronger knockdown. Boxplots

represent all data points with median, 25th and 75th percentiles, and red dotted lines indicate the

control median. A list of full genotypes for fly crosses used in these experiments is provided in

S1 File. Statistical details, including sample size, confidence intervals and p-values, are provided

in **S6 File**.

1371

Fig. 7. Functional relatedness of genes within disease-associated CNV regions correspond

with higher pathogenicity.

(**A**) Bar plot shows frequency of reported *de novo* occurrence of the 16p12.1 deletion (9, 11)

compared to the autism-associated 16p11.2 deletion (11, 105, 106). Schematic shows a model for

higher functional connectivity of genes within the 16p11.2 region compared to the 16p12.1

region. Only genes with *Drosophila* homologs are represented. (**B**) Phenotypic scores of

individual 16p11.2 homologs (grey) are significantly enhanced or suppressed by a second

16p11.2 homolog (orange). In contrast, little variation in phenotypic scores is observed for

16p12.1 homologs (grey) with simultaneous knockdown of another homolog (green). The

interacting homologs are labeled as follows: A: *Pp4-19C* (*PPP4C*), B: *CG17841* (*FAM57B*), C:

coro (*CORO1A*), D: *Ald1* (*ALDOA*), E: *Rph* (*DOC2A*), F: *Tao* (*TAOK2*), G: *Asph* (*ASPHD1*), H:

klp68D (*KIF22*), I: *Pa1* (*PAGR1*), J: *Pis* (*CDIPT*), K: *CG10465* (*KCTD13*), L: *CG15309*

(*YPEL3*), M: *Doc3*, (*TBX6*), N: *rl* (*MAPK3*), W: *CG14182* (*MOSMO*), X: *Cen* (*CDR2*), Y: *Sin*

(*POLR3E*), Z: *UQCR-C2* (*UQCRC2*). (**C**) Pairwise knockdown of homologs of 16p11.2 genes

(n=27) show a larger magnitude of interactions compared with those among 16p12.1 homologs

(n=5, two-tailed Mann-Whitney test, *p=0.011). Interaction values of zero (blue shade) represent

no interactions, while values above or below zero represent negative (in red) and positive (in

green) interactions, respectively. (**D**) Pairs of 16p11.2 homologs exhibit a higher proportion of

shared differentially-expressed genes compared to pairs of 16p12.1 homologs (n=30 for 16p11.2,

n=12 for 16p12.1, two-tailed Mann-Whitney test, *p= 0.031). (**E**) Network diagram shows

connections between human 16p11.2 or 16p12.1 genes within a brain-specific interaction

network. 16p12.1 genes are indicated in green, 16p11.2 genes in orange, connector genes in grey,

and connector genes that are intolerant to functional variation (RVIS $\leq 20^{th}$ percentile) in dark

green. *C16orf92* and *C16orf54* for 16p11.2 and *PDZD9* for 16p12.1 were not present in the brain

network and were therefore excluded from the network analysis. (**F**) Genes within the 16p11.2

region show higher average pairwise connectivity in a human brain-specific network, measured

as the inverse of the shortest paths between two genes, compared to 16p12.1 genes (n=25 for

16p11.2, n=6 for 16p12.1, two-tailed Mann-Whitney, *p=0.036, see **S5 File**). (**G**) 16p11.2

connector genes have lower RVIS percentile scores compared to 16p12.1 connector genes

(n=166 for 16p11.2, n=33 for 16p12.1, two-tailed Mann-Whitney, *p=0.017, see **S5 File**).

Functionally-intolerant genes are represented in dark green. Boxplots represent all data points

with median, 25th and 75th percentiles. Statistical details are provided in **S6 File**.

1404

Supporting Information Legends

S1 Fig. Expression levels of 16p12.1 homologs in *Drosophila* and *X. laevis*.

(**A**) *Drosophila* homologs of 16p12.1 genes were knocked down using nervous system-specific *Elav-GAL4* driver with overexpression of *Dicer2* at 25°C. RT-qPCR confirmed 40-60% knockdown of the 16p12.1 homologs (two-tailed student's t-test, *p<0.05). As knockdown of *Sin* caused embryonic lethality in these conditions, all experiments in the nervous system and RT-qPCR were performed without overexpression of *Dicer2* and reared at room temperature (RT). $Sin^{KK101936}$ and $Sin^{HMC03807}$ were also embryonic lethal without *Dicer2*. Sin^{GD7027_2} did not show knockdown of the homolog, and the RNAi line was therefore not used for further experiments. All experiments were performed in comparison to appropriate background-specific controls. Only one control is shown per gene. GD VDRC Control is shown in all cases for simplification (Control^{GD}). A list of full genotypes for fly crosses used in these experiments is provided in **S1 File**. (**B**) Normalized band intensity of RT-PCR of *X. laevis* tadpoles injected with different morpholino dosages of the 16p12.1 homologs compared to the uninjected control. Different morpholino sequences were used for the L and S alleles for *uqcrc2* and *mosmo*, while unique sequences were used for both L and S alleles for *cdr2* or *setd5*. As the S allele has not been annotated for *polr3e*, only the L allele was targeted. Colored bars represent the dosages of morpholinos used, with grey bars indicating amounts for "partial knockdown" (approximately 50% of expression) and black bars indicating amounts for "stronger knockdown". 10ng of morpholino was used for partial and stronger KD experiments for *mosmo* S allele, as increasing concentrations did not lead to differences in knockdown of the allele. Bar plots represent mean +/- SD, and red dotted lines indicate 50% expression. Statistical details are provided in **S6 File**.

S2 Fig. Global neurodevelopmental defects with knockdown of 16p12.1 *Drosophila*

homologs.

(**A**) Summary of phenotypes observed with tissue-specific knockdown of each of the 16p12.1

homologs. (**B**) Ubiquitous (*Da-GAL4*) and nervous system-specific (*Elav-GAL4*) knockdown of

multiple RNAi lines for each 16p12.1 homolog showed a range of lethality and developmental

defects. (**C**) Representative brightfield images of adult wings with knockdown of 16p12.1

homologs using the wing-specific driver bx^{MS1096}-*GAL4*. Severe phenotypes were observed for

UQCR-C2 and *Sin*, with some RNAi lines, including *UQCR-C2$^{GD11238\_2}$*, *UQCR-C2^{KK108812}* and

Sin^{GD7027}, showing lethality. Scale bar represents 500μm. (**D**) Bang sensitivity assay for adult

flies with nervous system-specific knockdown of the 16p12.1 homologs showed increased

recovery time for *UQCR-C2^{GD11238}* (n=95, two-tailed Mann-Whitney test, *p=0.003). *Sin^{GD7027}*

adult flies exhibited severe motor defects and could not be tested for the phenotype. Boxplots

represent all data points with median, 25th and 75th percentiles. Statistical details are provided in

S6 File. A list of full genotypes for fly crosses used in these experiments is provided in **S1 File**.

1442

S3 Fig. *Sin^{GD7027}* and *CG14182$^{GD2738\_2}$* lead to decreased brain area and decreased number

of proliferating cells in the brain.

(**A**) Representative confocal brightfield images of nervous system-specific knockdown of

16p12.1 homologs show decreased total brain area for *Sin^{GD7027}* and *CG14182$^{GD2738\_2}$*. Scale bars

represent 100μm. (**B**) *Elav-GAL4* mediated knockdown led to decreased number of

phosphorylated histone-3 positive cells (pH3, green) in the brain lobe (DAPI, blue) with

knockdown of *Sin^{GD7027}* (n=10, two-tailed Mann-Whitney, *p= 9.74×10^{-4}) and *CG14182$^{GD2738\_2}$*

(n=15, *p=0.026), indicating decreased proliferation with knockdown of the homologs.

Knockdown of Sin^{GD7027} led to decreased number of Dcp-1 positive cells (*p= 2.78×10^{-4}, red) in

the brain lobe. Scale bar represents 50μm. Boxplots represent all data points with median, 25th

and 75th percentiles. Statistical details, including sample size, confidence intervals and p-values,

are provided in **S6 File**. A list of full genotypes for fly crosses used in these experiments is

provided in **S1 File**.

1456

S4 Fig. Enriched GO terms observed with knockdown of 16p12.1 fly homologs in the

nervous system.

(**A**) Clusters of enriched Gene Ontology (GO) Biological Process terms for differentially

expressed fly genes observed with nervous system-specific knockdown of 16p12.1 homologs

(left) and their human homologs (right). While some clusters of terms overlap among 16p12.1

homologs, genes dysregulated with knockdown of individual homologs exhibit unique

enrichments for GO terms, suggesting their independent action towards neuronal development.

Venn diagrams show overlaps of enriched GO Complete Biological Processes terms for (**B**)

differentially-expressed fly genes observed with knockdown of individual 16p12.1 homologs, or

(**C**) human homologs of the fly genes. We also observed that most of the (**D**) fly homologs or (**E**)

human counterparts of the differentially-expressed genes were unique to each 16p12.1 homolog.

A list of differentially expressed genes with knockdown of 16p12.1 homologs, as well as a list of

all enriched GO terms for these gene sets, is detailed in **S2 File**. Venn diagrams were constructed

using Venny 2.1 software (https://bioinfogp.cnb.csic.es/ tools/venny).

1471

S5 Fig. Decreased dosage of 16p12.1 homologs leads to multiple neurodevelopmental

phenotypes in *X. laevis*.

(A) Representative images of tadpoles injected with control morpholino or morpholinos for

16p12.1 homologs, indicating facial landmarks for face width (yellow), height (blue), angle

(green), and orofacial (red) and eye (orange) area. Boxplots showing face height, width, angle,

and orofacial and eye area of each knockdown compared to its own control. Knockdown of

mosmo (n=50, two-tailed student's t-test, *p=0.010) and *cdr2* (n=54, *p=3.68×10^{-6}) led to

increased face height. Knockdown of *cdr2* (*p=7.75×10^{-4}), *polr3e* (n=37, *p=1.97×10^{-13}) and

mosmo (*p=1.36×10^{-11}) led to decreased face width, while knockdown of *cdr2* (*p=1.03×10^{-8}),

polr3e (*p=2.73×10^{-4}) and *mosmo* (*p=3.50×10^{-7}) led to decreased face angle. Knockdown of

polr3e (*p=3.29×10^{-16}) and *mosmo* (*p=1.47×10^{-8}) led to decreased orofacial area, and

knockdown of *polr3e* (*p=1.01×10^{-18}), *mosmo* (*p=7.23×10^{-10}) and *cdr2* (*p=0.009) led to

decreased eye area. Data represents strong knockdown of the 16p12.1 homologs, except for *cdr2*,

which showed lethality and is represented with partial knockdown. All measures were

normalized to their respective control injected with the same morpholino amount. Scale bars

represent 500μm. **(B)** Boxplots showing axon length of each knockdown compared to its own

control. Strong knockdown of *mosmo* led to decreased axon length in neural tube explants

(n=548, two-tailed student's t-test, *p=7.40×10^{-12}), which was rescued by co-injection with

overexpressed (OE) mRNA of the gene (n=566, *p=4.06×10^{-5}). All measures were normalized

to their respective control injected with the same morpholino amount. **(C)** Representative images

stained with anti-tubulin show forebrain (red on control image) and midbrain (blue) areas of the

side injected with morpholino (right, red asterisk), which were normalized to the uninjected side

(left). Partial knockdown of *mosmo* led to decreased forebrain (n=47, two tailed student's t-test,

*p=1.18×10^{-9}) and midbrain (*p=1.45×10^{-7}) area. Graphs represent contralateral ratio of brain

area compared to uninjected side of the embryo. Scale bars represent 500μm. All boxplots

represent all data points with median, 25th and 75th percentiles. In each case, measurements for

each knockdown were compared to controls injected with equal amounts of morpholino.

Statistical details, including sample size, confidence intervals and p-values, are provided in **S6**

File.

1501

S6 Fig. Whole western blot for phosphorylated histone-3 in *X. laevis* embryos with

knockdown of *polr3e*, *mosmo* and *setd5*.

(**A**) Three replicate western blot experiments were performed. The intensity of bands at 17 kDa,

corresponding with pH3 (top, indicated with arrow), were normalized to the β-actin loading

control (bottom). (**B**) Partial knockdown of *polr3e* shows reduced band intensity with anti-pH3

antibody compared to β-actin loading control. Bar plot represents mean ± SD.

1508

S7 Fig. Knockdown of *Sin* and *CG14182* lead to disruption of the fly eye in a dosage

sensitive manner.

(**A**) Representative brightfield adult eye images and *Flynotyper* phenotypic scores of eye-specific

knockdown of 16p12.1 homologs with *GMR-GAL4* and no overexpression of *Dicer2* at 30°C. A

mild eye phenotype was observed with knockdown of *Sin*, replicated across multiple RNAi lines

(two-tailed Mann-Whitney with Benjamini-Hochberg correction, *p<0.05). (**B**) Representative

images and *Flynotyper* scores of eye-specific knockdown of 16p12.1 homologs with *GMR-GAL4*

and overexpression of *Dicer2* at 30°C. Severe eye phenotypes were observed for all tested RNAi

lines of *Sin* (*p< 1.13×10^{-4}) and *CG14182* (*p< 4.70×10^{-4}). Scale bar represents 100 μm.

Boxplots represent all data points with median, 25th and 75th percentiles, and red dotted lines

indicate the control median. Statistical details, including sample size, confidence intervals and p-

values, are provided in **S6 File**. A list of full genotypes for fly crosses used in these experiments

is provided in **S1 File**.

1522

S8 Fig. Pairwise knockdown of homologs of 16p12.1 genes lead to moderate changes in eye

severity compared to individual knockdown of genes.

(**A-D**) Pairwise knockdown of homologs of 16p12.1 genes led to subtle changes in phenotypic

scores, with only four significant interactions (compared to control lines) validated by multiple

RNAi lines. *Sin* and *CG14182* enhanced the eye phenotype of *UQCR-C2*^{GD11238}, while *Sin*

enhanced *CG14182*^{GD2738} eye phenotype and *UQCR-C2* suppressed *Cen*^{GD9689} eye phenotype

(two-tailed Mann-Whitney with Benjamini-Hochberg correction, *p<0.05). Boxplots represent

all data points with median, 25th and 75th percentiles. Red dotted lines indicate the median of

recombinant lines crossed with control. Statistical details, including sample size, confidence

intervals and p-values, are provided in **S6 File**. A list of full genotypes for fly crosses used in

these experiments is provided in **S1 File**.

1534

S9 Fig. Range of observed and expected phenotypic scores using a multiplicative model.

Histograms representing the distribution of observed (in grey) and expected (in blue) phenotypic

scores of *GMR-GAL4*-mediated pairwise knockdowns of 16p12.1 homologs and interacting

genes (top panel) and *GMR-GAL4*-mediated single knockdowns of potential interacting genes

tested (bottom panel). The distribution shows an overlap of the observed and expected

phenotypic scores values.

1541

S10 Fig. Expected genetic interactions for each category of interacting gene analyzed in this

study. The diagram shows representative gene networks of each of the 16p12.1 homologs

analyzed. *UQCR-C2* is shown as an example, where the gene is closely connected to

transcriptome targets and functionally related genes, while neurodevelopmental genes and

patient-specific "second-hit" genes distribute more randomly between gene categories that can

lead to positive and negative genetic interactions (top bubble) or combined independent effects

(bottom bubble). These hypotheses aligned with our results, as we identified genetic interactions

for 42/61 pairwise combinations (68.8%) with direct transcriptome targets, compared to 22/55

(40%) interactions identified with functionally related and neurodevelopmental genes (p=0.0027,

Fisher's exact test). Moreover, we observed that homologs of genes carrying "second-hits" in

severely affected children with the 16p12.1 deletion interacted with 16p12.1 homologs in 37/96

(38.5%) of the pairs, although the proportion of genetic interactions was not as high compared to

that identified with functionally related genes or genes in neurodevelopmental pathways and

transcriptome targets (64/101, 63%, Fisher's exact test, p=0.019).

1556

S11 Fig. Phenotypic scores of the tested pairwise knockdowns of *UQCR-C2*[GD11238] with

homologs of "second-hit" or neurodevelopmental genes in *Drosophila* eye.

Flynotyper phenotypic scores of *UQCR-C2*[GD11238] crossed with RNAi, mutant or overexpression

lines of (**A**) neurodevelopmental genes or genes functionally related with *UQCR-C2* function, (**B**

and C) homologs of "second-hits" identified in children with 16p12.1 deletion, and (**D**)

transcriptome targets and functionally related groups identified in RNA-sequencing of *UQCR-C2*

knockdown model. Boxplots represent all data points with median, 25th and 75th percentiles.

Red dotted lines indicate the median of recombinant lines crossed with control. A list of full

genotypes and statistics, including sample size, confidence intervals and p-values, for these

experiments are provided in **S1 File** and **S6 File**.

1567

S12 Fig. Phenotypic scores of the tested pairwise knockdowns of *Cen*^{GD9689} with homologs of

"second-hit" or neurodevelopmental genes in *Drosophila* eye.

Flynotyper phenotypic scores of *Cen*^{GD9689} crossed with RNAi, mutant or overexpression lines of

(**A**) neurodevelopmental genes or genes functionally related with *Cen* function, (**B and C**)

homologs of "second-hits" identified in children with 16p12.1 deletion, and (**D**) transcriptome

targets and functionally related groups identified in RNA-sequencing of *Cen* knockdown model.

Boxplots represent all data points with median, 25th and 75th percentiles. Red dotted lines

indicate the median of recombinant lines crossed with control. A list of full genotypes and

statistics, including sample size, confidence intervals, and p-values, for these experiments are

provided in **S1 File** and **S6 File**.

1578

S13 Fig. Phenotypic scores of the tested pairwise knockdowns of *Sin*^{GD7027} with homologs of

"second-hit" or neurodevelopmental genes in *Drosophila* eye.

Flynotyper phenotypic scores of *Sin*^{GD7027} crossed with RNAi, mutant or overexpression lines of

(**A**) neurodevelopmental genes or genes functionally related with *Sin* function, (**B and C**)

homologs of "second-hits" identified in children with 16p12.1 deletion, and (**D and E**)

transcriptome targets and functionally related groups identified in RNA-sequencing of *Sin*

knockdown model. Boxplots represent all data points with median, 25th and 75th percentiles.

Red dotted lines indicate the median of recombinant lines crossed with control. A list of full

genotypes and statistics, including sample size, confidence intervals and p-values, for these

experiments are provided in **S1 File** and **S6 File**.

1589

S14 Fig. Phenotypic scores of the tested pairwise knockdowns of *CG14182*^{GD2738} with

homologs of "second-hit" or neurodevelopmental genes in *Drosophila* eye.

Flynotyper phenotypic scores of *CG14182*^{GD2738} crossed with RNAi, mutant or overexpression

lines of (**A**) neurodevelopmental genes or genes functionally related with *CG14182* function, (**B**

and C) homologs of "second-hits" identified in children with 16p12.1 deletion, and (**D**)

transcriptome targets and functionally related groups identified in RNA-sequencing of *CG14182*

knockdown model. Boxplots represent all data points with median, 25th and 75th percentiles.

Red dotted lines indicate the median of recombinant lines crossed with control. A list of full

genotypes and statistics, including sample size, confidence intervals and p-values, for these

experiments are provided in **S1 File** and **S6 File**.

1600

S15 Fig. Multiple homologs of patient-specific "second-hits" and neurodevelopmental genes

modulate phenotypes caused by knockdown of *UQCR-C2*^{GD11238} and *Cen*^{GD9689}.

Representative brightfield adult eye images and phenotypic scores of eyes from RNAi, mutant or

overexpression lines of neurodevelopmental genes, homologs of genes with "second-hits" in

children with 16p12.1 deletion, and transcriptome targets that significantly enhanced (Enh.) or

suppressed (Supp.) the phenotypes of recombinant lines of *UQCR-C2*^{GD11238} (**A**) or *Cen*^{GD9689} (**B**)

(*p<0.05, two-tailed Mann-Whitney tests with Benjamini-Hochberg correction). Scale bar

represents 100 μm. Boxplots represent all data points with median, 25th and 75th percentiles.

Red dotted lines indicate the median of recombinant lines crossed with control. A list of full

genotypes and statistics, including sample size, confidence intervals and p-values, for these

experiments are provided in **S1 File** and **S6 File**.

1612

S16 Fig. Multiple homologs of patient-specific "second-hits" and neurodevelopmental genes

modulate phenotypes caused by knockdown of *CG14182^{GD2738}* and *Sin^{GD7027}*.

Representative brightfield adult eye images and phenotypic scores of eyes from RNAi, mutant or

overexpression lines of neurodevelopmental genes, homologs of genes with "second-hits" in

children with 16p12.1 deletion, and transcriptome targets that significantly enhanced (Enh.) or

suppressed (Supp.) the phenotypes of recombinant lines of *CG14182^{GD2738}* (**A**) or *Sin^{GD7027}* (**B**)

(*p<0.05, two-tailed Mann-Whitney tests with Benjamini-Hochberg correction). Scale bar

represents 100 μm. Boxplots represent all data points with median, 25th and 75th percentiles.

Red dotted lines indicate the median of recombinant lines crossed with control. A list of full

genotypes and statistics, including sample size, confidence intervals and p-values, for these

experiments are provided in **S1 File** and **S6 File**.

1624

S17 Fig. Genetic interactions between 16p12.1 homologs and neurodevelopmental genes,

transcriptome targets and homologs of patient-specific "second-hit" genes. Scatter plots

depict the interactions tested for *GMR-GAL4* recombinant lines for *UQCR-C2^{GD11238}* (**A**),

Cen^{GD9689} (**B**), *CG14182^{GD2738}* (**C**), and *Sin^{GD7027}* (**D**). The plots show the average phenotypic

score of the interacting gene on the x-axis using VDRC GD (top) or KK (bottom) fly lines and

the average observed phenotypic score for the double knockdown on the y-axis. Blue line

represents the expected phenotypic score of the pairwise knockdown calculated as the product of

the first hit phenotype (*Flynotyper* score of first hit crossed with control, such as *UQCR-*

C2^{GD11238} X Control^{GD}) for each theoretical phenotypic value of interacting gene (ranging from 0

to 60) represented on x-axis. All positive and negative (validated or potential) interactions are

represented in green and red, respectively, and fly lines of genes with no significant interactions

are shown in grey.

1637

S18 Fig. "Second-hit" genes identified in probands with 16p12.1 deletion.

Pedigrees of 15 families with the 16p12.1 deletion, highlighting 23 genes with rare secondary

likely-pathogenic mutations (CNVs and SNVs) that were identified in severely affected children

with the 16p12.1 deletion. These 23 genes carrying "second-hit" mutations were selected for

Drosophila experiments to test how their decreased expression affects the neurodevelopmental

phenotypes observed for 16p12.1 homologs. Family members who carry either the 16p12.1

deletion or individual genes with "second-hits" are indicated in the pedigrees. Phenotypes

observed for affected children and other family members are indicated below each pedigree.

1646

S19 Fig. *Osa* and *Nipped-A* interact with *CG14182^{GD2738}* through cellular processes during

eye development.

(**A**) Representative brightfield adult eye images show that simultaneous knockdown of

CG14182^{GD2738} with *osa^{JF01207}* leads to a suppressed (Supp.) eye phenotype, while simultaneous

knockdown of *Nipped-A^{HMS00167}* leads to synergistic enhancement (Enh.) in eye phenotype

compared to single knockdown of *CG14182^{GD2738}*. Scale bar represents 100 μm. (**B**)

Representative confocal images of third instar larval eye discs stained with DAPI (blue) and anti-

phosphorylated histone-3 (pH3, green) or anti-Dcp-1 (red), markers of cellular proliferation and

apoptosis, respectively. Positive pH3 or Dcp-1 cells were quantified posterior to the

morphogenetic furrow (indicated by white boxes). Simultaneous knockdown of $CG14182^{GD2738}$

with $osa^{JF01207}$ led to an increase in the number of pH3 positive cells (n=20, two-tailed Mann-

Whitney, *p= 2.33×10^{-6}) compared to the single knockdown of $CG14182^{GD2738}$. (**C**)

Simultaneous knockdown of $CG14182^{GD2738}$ with $Nipped-A^{HMS00167}$ led to a significant reduction

in the number of Dcp-1 positive cells compared to single knockdown of $CG14182^{GD2738}$ (n=15,

in red, two-tailed, Mann-Whitney, *p=2.54×10^{-5}). Scale bars represent 50 μm. Boxplots

represent all data points with median, 25th and 75th percentiles. Statistical details, including

sample size, confidence intervals and p-values, are provided in **S6 File**. A list of full genotypes

for fly crosses used in these experiments is provided in **S1 File**.

1665

S20 Fig. *setd5* **modifies phenotypes observed with knockdown of** *polr3e* **and** *mosmo* **in** *X.*

laevis.

(**A**) Representative images stained with anti-tubulin show forebrain (red on control image) and

midbrain (blue) areas of the side injected with morpholino (right, red asterisk), normalized to the

uninjected side (left). Simultaneous knockdown of *polr3e* and *setd5* in *X. laevis* led to decreased

forebrain (n=28, two-tailed student's t-test, *p=6.01×10^{-7}) and midbrain area (*p=1.67×10^{-7})

compared to knockdown of *polr3e* alone, which were not different to the partial knockdown of

setd5 alone (p>0.05). Scale bar represents 500 μm. (**B**) Normalized axon length of *X. laevis*

tadpoles with simultaneous knockdown of *mosmo* and *setd5* showed decreased axon length

different from the control injected with 22ng of morpholino (n=438, two-tailed, student's t-test,

*p=2.95×10^{-7}) and from individual knockdown of *setd5* (*p=1.86×10^{-9}) or *mosmo* (*p= 3.34×10^{-}

^{6}), showing a synergistic effect of decreased dosage of the homologs towards neuronal

phenotypes. **(C)** Normalized axon length of *X. laevis* tadpoles with simultaneous knockdown of

polr3e and *setd5* showed no change in axon length (two-tailed student's t-test, p>0.05). In each

case, the individual knockdown was normalized and compared to the control injected with the

same amount of morpholino. Boxplots represent all data points with median, 25th and 75th

percentiles. Statistical details, including sample size, confidence intervals and p-values, are

provided in **S6 File**.

1684

S21 Fig. Interactions between 16p12.1 homologs and "second-hits" are of higher magnitude

than those between 16p12.1 homologs. (**A**) Absolute values of scores of interactions identified

between 16p12.1 homologs are lower compared to those identified between the 16p12.1

homologs and "second-hit" homologs (n=3 for 16p12.1, n=37 for interactions with "second-

hits", two-tailed Mann-Whitney, *p=0.0032). (**B**) Comparison of absolute values of interaction

scores between 16p12.1 homologs to those observed between the homologs and homologs of

"second-hits" split by proband. Only one RNAi line per interacting gene is shown.

1692

S22 Fig. 16p11.2 genes exhibit higher phenotypic scores than 16p12.1 genes and are more

connected to each other than to other genes in the genome. (**A**) Pairwise eye-specific

knockdown of 16p11.2 homologs (*rl*, *CG10465* and *Pp4-19C* crossed with other 16p11.2

homologs) lead to more severe phenotypic scores compared to pairwise knockdown of 16p12.1

homologs (n=36 for 16p11.2, n=12 for 16p12.1, one-tailed Mann-Whitney, *p=3.51 $\times 10^{-5}$). Grey

circles represent pairwise knockdowns, while 16p11.2 and 16p12.1 single-homolog knockdowns

are represented in orange and green, respectively. (**B**) Analysis of a human brain-specific

network shows higher average pairwise connectivity, measured as the inverse of the shortest path

between two genes, between pairs of 16p11.2 genes compared to the connectivity of 16p11.2

genes to the rest of the genome (n=25 two-tailed Mann-Whitney, *p= 6.64×10^{-3}). This trend was

not observed for 16p12.1 genes (n=6, p>0.05). Boxplots represent all data points with median,

25th and 75th percentiles. Statistical details are provided in **S6 File**.

1705

S23 Fig. 16p12.1 genes are less connected than 16p11.2 genes in a human brain-specific

interaction network (**A**) Cumulative frequency plot shows low connectivity of 16p12.1 genes

(observed value shown as red dotted line) in a human brain-specific interaction network when

compared with 1000 permutations of six 16p11.2 genes (blue, 0.4^{th} percentile, p=0.117, one-

tailed z-score test) and random set of genes in the genome (green, 9.2^{nd} percentile, p=0.143, one-

tailed z-score test). Cumulative frequency plot of 1000 simulations of bins of 25 contiguous

genes (green) (**B**) or six contiguous genes (blue) (**C**), compared with the median connectivity

values for 16p11.2 genes (green dotted line, 78.6^{th} percentile, one-tailed z-score test, p=0.246)

and 16p12.1 genes (blue dotted line, 5.7^{th} percentile, p=0.130).

1715

1716 **S1 Table. *Drosophila* and *X. laevis* homologs of 16p12.1 genes.**

1717 DIOPT (24) and reciprocal BLAST (80) searches were used to identify fly homologs of 16p12.1

1718 genes. The expression of homologs in the larval central nervous system during development was

1719 assessed using FlyAtlas Anatomy microarray expression data from FlyBase (81).

1720

S2 Table. Number of experiments and genetic interactions identified in this study.

This table lists the number of crosses, RNAi/mutant/overexpression lines, tested pairwise

combination of genes, and genes that significantly enhanced or suppressed the phenotype of

16p12.1 homologs (**S2A Table**). The number of confirmed and potential negative and positive

genetic interactions towards eye phenotypes using the multiplicative model is also provided (**S2B**

Table).

1727

1728   **S3 Table. Hypotheses, experimental design, and conclusions obtained for experiments**

1729   **performed in our study.**

1730   Explanation of hypotheses, experimental design, results, and conclusions that can be drawn or

1731   hypotheses that can be further tested for each experiment performed in this study is provided.

1732

S4 Table. "Second-hit" variants identified in families with 16p12.1 deletion and tested for

interactions with 16p12.1 homologs.

Genes carrying "second-hits" were previously identified (9) through exome sequencing and SNP

microarrays in 15 children with the 16p12.1 deletion, and selected as disease-associated genes

carrying rare (ExAC frequency $\leq$1%) copy-number variants, loss-of-function (frameshift,

stopgain or splicing) mutations, or *de novo* or likely-pathogenic (Phred-like CADD $\geq$25)

missense mutations.

1740

1741   **S5 Table. Phenotypes observed for individual 16p12.1 homologs in *Drosophila* and *X. laevis***

1742   **models.**

1743   Developmental and neuronal phenotypes observed with individual knockdown of 16p12.1

1744   homologs in *Drosophila* and *X. laevis*.

1745

1746 **S6 Table. Sequences of oligonucleotides used for confirming gene knockdown in *Drosophila***

1747 **and *X. laevis.***

1748

1749 **S7 Table. *X. laevis* morpholino sequences used in this study.**

1750

S1 File. Stock list and genotypes of *Drosophila* lines used in the study (Excel file).

This file details genotypes of tissue-specific drivers, and stock centers, stock numbers and

genotypes of RNAi used to knock down the four 16p12.1 homologs, as well as RNAi, mutant, or

overexpression lines for homologs of "second-hit" genes and genes within conserved

neurodevelopmental pathways. Genotypes of recombinant lines of 16p12.1 homologs crossed

with interacting genes are also detailed, as well as the individual controls used for each

experiment. Details of the number of homologs, fly lines, and crosses used for all interaction

experiments are also provided. BDSC: Bloomington Drosophila Stock Center, VDRC: Vienna

Drosophila Stock Center.

1760

S2 File. Differentially expressed genes and Gene Ontology enrichments for knockdown of

16p12.1 homologs (Excel file).

List of differentially expressed genes, defined as log-fold change >1 or <-1 and false discovery

rate (FDR) <0.05 (Benjamini-Hochberg correction), following RNA sequencing of *Drosophila*

fly heads with *Elav-GAL4*-mediated knockdown of the four 16p12.1 homologs. Human

homologs of differentially-expressed fly genes were identified using DIOPT (24). Enriched Gene

Ontology (GO) and PantherDB terms are also provided for differentially-expressed fly genes and

their human homologs.

1769

1770 **S3 File. Summary of genetic interactions identified with 16p12.1 homologs in *Drosophila***

1771 **(Excel file).**

1772 This file details the genes that modulated the phenotypes of the 16p12.1 homologs, as well as the

1773 genetic interactions identified in this study.

1774

1775 **S4 File. Interaction scores and observed and expected values for all tested genetic**

1776 **interactions of 16p12.1 homologs in *Drosophila* (Excel file).**

1777

1778 **S5 File. Connectors of 16p12.1 and 16p11.2 genes identified in a human brain network**

1779 **(Excel file).**

1780

1781 **S6 File. Statistical analysis of experimental data (Excel file).**

1782 This file provides details of all statistical information, including sample size, test statistics, p-

1783 values, and Benjamini-Hochberg FDR corrections for all data shown in the main and

1784 supplemental figures. Details for ANOVA tests include factors, degrees of freedom, tests

1785 statistics, and post-hot pariwise t-tests with Benjamini-Hochberg FDR corrections.

1786

A Pairwise interactions in *Drosophila*

GMR-GAL4, UAS-RNAi 1 ✕ UAS-RNAi 2

UQCR-C2 *CG14182* *Sin* *Cen*

B

UQCR-C2^GD11238/Control^GD Cen^GD9689/Control^GD Sin^GD7027/Control^GD CG14182^GD2738/Control^GD

UQCR-C2^GD11238/Sin^GD7027 (Enh.) UQCR-C2^GD11238/CG14182^GD2738 (Enh.) Cen^GD9689/UQCR-C2^GD11238 (Supp.) CG14182^GD2738/Sin^GD7027 (Enh.)

C

Phenotypic score

- - - control avg.
— exp

UQCR-C2^GD11238/Control^GD
UQCR-C2^GD11238/Sin^GD7027
UQCR-C2^GD11238/CG14182^GD2738
Cen^GD9689/Control^GD
Cen^GD9689/UQCR-C2^GD11238
CG14182^GD2738/Control^GD
CG14182^GD2738/Sin^GD7027

D Phenotype (percentage of control)

Mild   Severe

Control	100%
Homolog A	120%
Homolog B	140%
No interaction	Expected (120% x140% = 168%)
Positive interaction	130%
Negative interaction	200%

E Pairwise interactions in *X. laevis*

Homolog 1 Homolog 2 Homolog 1+2

F

Control * injected / uninjected
polr3e *
mosmo *
polr3e/mosmo *

Contralateral ratio of average brain area

Forebrain
Midbrain

Control polr3e mosmo polr3e/mosmo

* ns

G

pH3
β-actin

Control polr3e mosmo polr3e/mosmo

Normalized band intensity
pH3/β-actin

Control mosmo polr3e polr3e/mosmo

Interactions of 16p12.1 homologs with patient-specific "second-hits"

A Patient-specific "second-hit" interaction in flies

GL_01

16p12.1 del — ○

16p12.1 del
SETD5 (upSET)

Control^{BL}

CG14182^{GD2738}/Control^{BL}

upSET^{HMC03177}

CG14182^{GD2738}/*upSET^{HMC03177}* Enh.

Phenotype (percentage of control)

ns   *

Control^{BL}
Sin^{GD7027}/C^{BL}
upSET^{HMC03177}
Sin^{GD7027}/upSET^{HMC03177}
Exp. (Sin/upSET)
CG14182^{GD2738}/C^{BL}
Exp. (CG14182/upSET)
CG14182^{GD2738}/upSET^{HMC03177}

B Cell proliferation/apoptosis

Control^{BL} *CG14182^{GD2738}*/Control^{BL} *upSET*   *CG14182^{GD2738}*/*upSET^{HMC03177}*

pH3

Dcp-1

Number of positive cells

pH3   *   ns

Dcp-1   *   *

Control^{BL}
CG14182^{GD2738}/C^{BL}
upSET
CG14182/upSET

C Forebrain/midbrain area

Control   uninjected   injected *

mosmo   *

setd5   *

mosmo/setd5   *

Contralateral ratio of average brain area

ns
ns
Forebrain

*
ns
Midbrain

Control   mosmo   setd5   mosmo/setd5

D Axon length

axon

Normalized axon length

*
*

Control   mosmo   setd5   mosmo/setd5