# Syringeal vocal folds do not have a voice
# in zebra finch vocal development

Alyssa Maxwell, Iris Adam, Pernille S. Larsen, Peter G. Sørensen, Coen P.H. Elemans[*]

**Affiliation**: Department of Biology, University of Southern Denmark, 5230 Odense, Denmark

*Corresponding author: coen@biology.sdu.dk

**Abstract**

Vocal behaviour can be dramatically changed by both neural circuit development and postnatal maturation of the body. During song learning in songbirds, both the song system and syringeal muscles are functionally changing, but it is unknown if maturation of sound generators within the syrinx contributes to vocal development. Here we densely sample the respiratory pressure control space of the zebra finch syrinx *in vitro*. We show that the syrinx produces sound very efficiently and that key acoustic parameters, minimal fundamental frequency, entropy and source level, do not change over development in both sexes. Thus, our data suggests that the observed acoustic changes in vocal development must be attributed to changes in the motor control pathway, from song system circuitry to muscle force, and not by material property changes in the avian analog of the vocal folds. We propose that in songbirds, muscle use and training driven by the sexually dimorphic song system are the crucial drivers that lead to sexual dimorphism of the syringeal skeleton and musculature. The size and properties of the instrument are thus not changing, while its player is.

*Keywords:* Vocal learning, neuromechanics, motor control, voice production

**Introduction**

Many vertebrates change their vocal output over postnatal development[1–4]. Especially in species capable of vocal imitation learning, progressive changes in vocal output are attributed to changes in the underlying brain circuitry. However, the maturation of the vocal organ and tract have profound effect on voice during puberty in humans[5], and can explain transitions from juvenile to adult calls in marmosets[6]. Thus, the postnatal development of the body can drive changes in vocal behavior, and therefore needs to be included when studying vocal development.

The zebra finch is a widespread animal model to study the mechanisms underlying vocal imitation learning and production[7,8]. In male zebra finches, sound production changes dramatically over the 100-day period of from hatching to adulthood[2,9,10], developing from variable subsong to highly stereotyped crystallized song. These changes, including the ones in acoustic structure, are typically attributed to changes in neural circuitry, even though the vocal organ, the syrinx, is also undergoing significant anatomical and functional changes during this time[11–14]. Coinciding with the transition from the sensory to the sensorimotor phase of learning around 20 DPH, syringeal muscles in male zebra finches increase in mass and cross-sectional area[11]. Additionally, the contractile properties of male syringeal muscles increase in speed[12], which affects the transformation from neural commands to muscle force[14]. Thus, over development functional changes occur in the syrinx that change the transformation from neural signals into force which thereby change motor control of the vocal organ. However, it remains unknown if the organ itself also exhibits changes that will affect the complex transformation from muscle forces to sound.

Analogous to mammals, birds produce sound when expiratory airflow from the bronchi induces self-sustained vibrations of vocal fold-like structures within the syrinx[15–18] (Fig. 1a) following the myoelastic-aerodynamic (MEAD) theory for voice production[19]. Songbirds have two bilateral sets of independently controlled paired folds or labia, one in each bronchus or hemi-syrinx, with the lateral labium on the lateral side, and medial vibratory mass (MVM) on the medial side. The MVM is a tissue continuum in which the thicker part is called the medial labium (ML)[20–22] (Fig. 1b). These labia consist of multiple layers of tissue in most species investigated[22,23]. According to MEAD, acoustic features, such as fundamental frequency ($f_o$), or boundary conditions for vibration, such as the phonation threshold pressure, are largely set by the positioning of and driving pressures[19] on the vocal folds, as well as their resonance properties and shape. Vocal fold resonance properties in turn are determined by vocal fold length and ultrastructure, such as collagen and elastin fiber composition, the occurrence of

tissue layering and layer orientation[24]. During vocal development in humans[25–27], and

marmosets[6], changes in size, (ultra)structure and mechanical properties of the labia have been

linked to dramatic changes in vocal output. However, it is unknown whether material properties

or morphology of songbird labial change over vocal development and if so whether these

changes contribute to the changing vocal output.

Tension and positioning of labia are modulated directly by two independent motor

systems in songbirds: the respiratory and the syringeal motor system. The respiratory system

generates the pressure and flow driving vibration of the avian vocal folds. In zebra finches, a

sound pulse is produced when these tissues collide within the vibratory cycle[15]. Additionally,

the syrinx is suspended in a pressurized air sac, the interclavicular air sac (ICAS), that is

essential for sound production in birds[15,28,29]. The pressure differential between bronchial

pressure ($p_b$) and ICAS pressure ($p_{icas}$) results in a net pressure, the transmural pressure ($p_t$),

that applies a net force on the MVM. This force has been shown to correlate most closely to

changes in $f_o$ during manipulation of $p_{icas}$ *in vivo*[29] and *ex vivo*[15], but we do not know how these

pressures interact to drive acoustic parameters.

The syringeal motor system of songbirds consists of up to 8 pairs of intrinsic and extrinsic

syringeal muscles[21,30,31] (Fig. 1a), which control the positioning of individual bones within the

syringeal skeleton as well as the torque on them. Consequently, syrinx muscles thereby - either

directly or indirectly – modulate the adduction level and tension of the labia, which are

suspended between the bones[32]. The control space of syringeal muscles is vast due to its

multidimensionality and only limitedly understood[15,31,33]. Taken together, both, changes in the

motor control spaces or the sound production properties of the syrinx within this space could

contribute to vocal development, but this is currently unknown.

Here we test if the acoustic output of the zebra finch syrinx sound generators changes

over vocal development. We focused on the syringeal dynamics controlled by respiratory

pressures and densely sampled the control space of bronchial versus air sac pressure.

**Materials and Methods**

***In vitro* <u>sound production</u>**

*Animals use and care*

Zebra finches (*Taeniopygia guttata,* order Passeriformes) were kept and bred in group
aviaries at the University of Southern Denmark, Odense, Denmark on a 12:12 hour light:dark
photoperiod and given water and food *ad libitum*. All experiments were conducted in
accordance with the Danish law concerning animal experiments and protocols were approved
by the Danish Animal Experiments Inspectorate (Copenhagen, Denmark). Adult birds were
provided with nesting material, *ad libitum.* Nest boxes were monitored daily, so that birds could
be accurately aged. Breeding began in December 2017 and ended November 2018.

We studied vocal output of the syrinx *in vitro* in four age groups: 25 DPH (N = 5 males,
N = 5 females), 50 DPH (N = 4 males, N = 4 females), 75 DPH (N = 6 males, N = 4 females),
and 100 DPH (N = 5 males, N = 6 females). Sex was determined by dissection postmortem (25
DPH) or plumage (all other ages).

*Syrinx mounting procedure*

Animals were euthanized by Isoflurane (Baxter, Lillerød, Denmark) overdose. The
syrinxes were dissected out while being regularly flushed and then submerged in a bath of
oxygenated Ringer's solution (recipe cf. [34]) on ice. Each syrinx was photographed with a Leica
DC425 camera mounted on a stereomicroscope (M165-FC, Leica Microsystems, Switzerland)
prior to being placed inside the experimental chamber. Each bronchus was placed over
polyethylene tubing (outer diameter 0.97 mm x inner diameter 0.58 mm, Instech Salomon, PA,
USA) and secured with 10-0 nylon suture (AROSurgical, Newport Beach, CA, USA). The
trachea was placed into a 1 mL pipette which was cut to the length of 15 mm (outer diameter
2.4 mm, inner diameter 1.4 mm). Some adipose tissue was left on the trachea to ensure an
airtight connection. To control the pressure inside the chamber it was made airtight with a glass
lid, which also allowed visualization of the syrinx through the stereomicroscope.

*Experimental setup*

We designed an experimental setup that allows studying the mechanical behavior of the
syrinx *in vitro* or *ex vivo* (described in detail in[15]). The bronchial pressure ($p_b$) and pressure in
the experimental chamber (i.e. interclavicular air sac pressure, $p_{icas}$) can be controlled
independently by two dual-valve differential pressure PID controllers (model PCD, 0-10 kPa,

Alicat Scientific, AZ, USA). Bronchial flow through the syrinx was measured with a MEMS flow sensor (PMF2102V, Posifa Microsystems, San Jose, USA) about 7 cm upstream from the bronchial connection. Sound was recorded with a ½ inch pressure microphone-pre-amplifier assembly (model 46AD, G.R.A.S, Denmark), amplified and high-pass filtered (0.2 Hz, 3-pole Butterworth filter, model 12AQ, G.R.A.S., Denmark). The sound recording sensitivity was tested before each experiment using a 1 kHz tone (sound calibrator model 42AB, G.R.A.S., Denmark). The microphone was placed 15 cm from the tracheal connector outlet and positioned at a 45° angle to avoid the air jet from the tracheal outlet. The sound, pressure and flow signals were low-pass filtered at 10 kHz (model EF120, Thor Labs) and digitized at 50 kHz (USB 6259, 16 bit, National Instruments, Austin, Texas).

*Experimental protocol*

To explore the acoustic output of the avian syrinx (Fig. 1a), we subjected the syrinxes to $p_b$ ramps from 0-3 kPa (over atmospheric pressure) at 1 kPa/s with $p_{icas}$ at a constant pressure. In one run, we applied 13 $p_{icas}$ settings at equidistant 0.23 kPa intervals between 0 and 3 kPa in a randomized order with 2 s pause ($p_b$ and $p_{icas} = 0$ kPa) between ramps (Fig. 1c). We avoided high flow regimes ($p_t > 2$ kPa) to decrease desiccation and avoid potential damage to the syrinx. When water in the bronchi or trachea prevented normal sound production in part of the run, we repeated the entire run. We subjected left and right hemi-syrinx separately in randomized order to this paradigm by allowing air flow only through one side and kept a one-minute rest between runs. The total duration for the experiment was maximally 20 minutes per individual.

*Data Analysis*

Sound was filtered with a 0.1 kHz 3rd order Butterworth high-pass filter, pressure and flow signals were filtered with a 2 kHz 3rd order Butterworth low-pass filter, all with zero-phase shift implementation (filtfilt function, Matlab). Sound, pressure and flow signals were binned in 2 ms duration bins, with a sliding window of 1 ms, and we calculated several parameters per bin. We calculated the RMS values for all signals. For the sound signal, we additionally extracted the fundamental frequency ($f_o$), source level (SL) and Wiener entropy (WE).

Fundamental frequency was extracted using the yin-algorithm[35] by manually optimizing minimal aperiodicity (0.05 – 0.2) and power (0.25 – 5.0 mPa). Source level at 1 m distance (SL) of the emitted sound was defined as:

$$SL = 20\log_{10}\frac{P}{P_\text{o}} + 20\log_{10}r, \qquad (1)$$

where $p$ is the RMS of sound pressure (Pa), $p_\text{o}$ is the reference pressure in air of 20 μPa,

and r the distance (m) from the tracheal outlet to the microphone. To calculate WE, we first

computed the amplitude spectrum Px of each bin using the periodogram method (nfft = 2048,

overlap = 1024 samples). WE was defined as:

$$WE = \log_{10}\left(\frac{geomean(Px)}{mean(Px)}\right). \qquad (2)$$

The transmural pressure over the MVM was defined as $p_\text{t} = p_\text{b} - p_\text{icas}$. Phonation threshold

pressures (PTP) were identified for $p_\text{icas}$, $p_\text{b}$, and $p_\text{t}$ as the first pressure value for each $p_\text{b}$ ramp

where sound power crossed the set threshold. The minimal $f_\text{o}$ was defined as the average $f_\text{o}$ of

the first value for each $p_b$ ramp where sound power crossed the set threshold. Minimal SL was

defined as the lowest SL value of all $p_b$ ramps per side.

By combining pressure, flow and sound signals we were able to calculate the mechanical

efficiency (ME) as the ratio (in dB) of acoustical power ($P_\text{acoustic}$) over aerodynamic power

($P_\text{aerodynamic}$):

$$ME = 10\log_{10}\left(\frac{P_\text{acoustic}}{P_\text{aerodynamic}}\right), \qquad (3)$$

The acoustic power (W) was assumed to radiate over half a sphere: $P_\text{acoustic} = \frac{4\pi r^2 p^2}{\rho v}$, with

air density $\rho = 1.2$ kg m$^{-3}$, speed of sound $v = 344$ m s$^{-1}$, and $p$ the RMS sound pressure in Pa.

The aerodynamic power $P_{aerodynamic} = p_\text{b}V$ where $p_b$ is the bronchial pressure (Pa) and $V$ is the

flow rate (m$^3$ s$^{-1}$.) All data analysis was done in Matlab [36] and R version 3.5.1[37].

*MVM dimensions*

After the experimental protocol, each syrinx was pinned down in the *situ* position and

kept in 4% (w/v) paraformaldehyde for 24 hours at 4°C. It was then placed in PBS and stored

at 4°C. To quantify the projected size of the MVM of each hemi-syrinx, we carefully cut the

syrinx in half and made pictures of the medial side of the *medial tympaniform membranes*

(MTMs) (Fig. 8d,e), and the lateral view of each hemi-syrinx using a Leica DC425 camera

mounted on a stereomicroscope (M165-FC, Leica Microsystems, Switzerland) in Leica

Application Suite (LAS Ver. 4.7.0). We measured the distances between physiological

landmarks in the MTM following[20,32]: (i) the distance between the *medio-ventral cartilage*

(MVC) and the *lateral dorsal cartilage* (LDC), and (ii) the distance between the LDC and the

*medio-dorsal cartilage* (MDC). Distance measurements between cartilages were taken edge-

to-edge in the middle of the cartilage. Additionally, we measured the area of the LDC which is

fully suspended in the MVM. Length and area measurements were taken using ImageJ (Ver.
2.0.0-rc-69/1.52p;[38]).

**<u>In vivo sound production</u>**

To measure the lowest fundamental frequency produced by an unactuated syrinx *in vivo*,
we performed unilateral tracheosyringeal nerve lesions on the right side of the syrinx of
juvenile and adult males.

*Sound recordings*

Song was recorded in custom-built, sound attenuated recording boxes (60x95x57 cm)
and vocalizations were recorded continuously using a omnidirectional microphone (Behringer
ECM8000) mounted 12 cm above the cage connected to an amplitude triggered recording
software (Sound Analysis Pro;[39]). Sound was digitized at a sampling rate of 44.1kHz with 16-
bit resolution (Roland octa capture, amplification 40 dB). Adult birds were recorded right
before and 3 weeks after denervation. Juveniles were recorded at 50 and 100 DPH.

*Tracheosyringeal nerve lesions*

Unilateral denervations of the tracheosyringeal nerve were performed as previously
described[40]. In brief, birds were anesthetized with isoflurane (induction 3%, maintenance 1.5 -
2%) and a 5 mm section of the right tracheosyringeal nerve was removed from the trachea
through a small incision of the skin. After that the skin was re-sutured and the birds were
allowed to recover in a heated cage. Adults were subsequently transferred to holding cages.
Juveniles received surgery at 30 DPH and were returned to their parents after surgery.

*Song analysis*

Adult birds: For each bird (N = 5, age >120 DPH, days post-surgery: 20±2), the most
common motif was determined by two experienced observers (IA, CPHE) in recordings
acquired before denervation. Subsequently, the same motif was identified in recordings
acquired 3 weeks after denervation. A custom written GUI (Matlab) was used to extract motifs
from post denervation sound recordings. 200 motifs per bird were randomly selected for
analysis. WAV files were high pass filtered at 0.3 kHz using a 3$^{rd}$ order Butterworth filter.
Syllables were assigned to be produced by the left or right hemi-syrinx based on whether their
spectral content stayed the same or changed after denervation, respectively following earlier
results by[41,42].

Juvenile birds: Song of juvenile males (N = 6) was analyzed on recordings from 50 DPH
(DPH 52.8 ± 1, days post-surgery: 22.3 ± 0.8). At this age, the syllable structure is not fully
stereotyped, but motifs usually can be distinguished. We thus measured the lowest fundamental
frequency of 50 motifs per bird irrespective of syllable identity. As the juvenile birds were
denervated while singing subsong, we couldn't distinguish which syllable was produced on
which side. Under the assumption that the lowest fundamental is produced by the denervated
side, we classified the measured syllables as right produced.

In all birds, frequencies were measured on spectrograms of post denervation recordings
(FFT size: 1024, overlap: 75%, Hanning window) in SASlab (Avisoft, Germany). In all
juvenile birds the frequency of the lowest harmonic (i.e $f_o$) was measured. In all other cases, to
increase the $f_o$ resolution, we measured the 3 - 5th harmonic and $f_o$ was calculated as:

232
$$f_o = {f_n}/{n}. \qquad (4)$$

where $n$ is the $n^{th}$ harmonic. If several different syllables where analyzed for the adult
birds the syllable with the lowest mean fundamental frequency (per side) was selected for
statistical analysis.

**Statistics**

To asses which pressure ($p_b$, $p_{icas}$, $p_t$) drives $f_o$ and SL in adults, we used a delta Bayesian
information criterion (ΔBIC) approach. In a first step, we fit a linear model to the $f_o$-pressure
and SL-pressure relationship. We then computed the BIC values for these models and
subtracted the BIC of the null model with no pressure variable ($f_o = 1$; SL = 1) to calculate the
ΔBIC score. The model with the lowest ΔBIC score represents the model with the best fit.

To compare the $f_o$ - $p_t$ and SL – $p_b$ relationship over ages and between side and sex, we
fit individual linear models to the data and extracted the slopes. For $f_o$ we split the data in two
regions: region S1 ($p_t = 0 – 0.75$ kPa) and region S2 ($p_t = 0.75 – 2$ kPa).

To test whether age, sex or side of the syrinx had a significant effect on our response
variables, we fit linear mixed effect models (LMMs) to our data using the lmer function of the
lmerTest package[43] with maximum likelihood optimization. Sex, side and age were fit as fixed
effects and animal was included as a random effect to correct for dependence. Model equations
can be found in Table S1, S2 and S3. To determine which effects significantly contribute to
the model, we performed model selection using a Likelihood Ratio Test (LRT) with the Chi
squared distribution using the lme4 package[44]. The difference between the absolute difference
in minimal $f_o$ between sexes and the difference in minimal $f_o$ *in vivo* was assessed with Welch's

t-test. The difference between S1 and S2 slopes for $f_o$ were assessed using paired t-tests for

males and females.

All reported values in the text are mean ± SD. All error bars in figures are S.D. The

outputs of al LMMs are reported in Supplementary Table S1, S2, and S3. We chose to present

data in figures spilt by sex even when not significant. Statistical significance was accepted at

$p < 0.05$ for all statistical tests.

**Results**

     We developed an experimental paradigm to quantify sound production as a function of driving parameters, because predicting the behavior of this complicated nonlinear system from anatomy or isolated mechanical property tests alone will only present correlates or rough estimates. Because we still have limited knowledge on the biomechanical effect of all syringeal muscles, we focused on syringeal dynamics controlled by respiratory pressures.

*Acoustic output of the adult syrinx*

     To quantify the functional acoustic output of the adult male and female zebra finch syrinx, we densely sampled the bronchial ($p_b$) and interclavicular ($p_{icas}$) pressure control space in isolated left and right hemi-syrinxes *in vitro*. For all zebra finch syrinxes tested, sound was produced in a pressure space enclosed by a minimal $p_b$ and $p_{icas}$ and exclusively in the lower half of the $p_b$, $p_{icas}$ space (Fig. 1d). Here a differential or transmural pressure ($p_t$) exerts force on the MVM that is positive when directed outwards from bronchus to surrounding air sac. We quantified the phonation threshold pressures (PTP) for the three pressures (Fig. 1g). The $PTP_b$ was not significantly different for sex (LMM, $p = 0.418$) or side (LMM, $p = 0.395$) and was $1.01 \pm 0.28$ kPa (range: $0.64 - 1.68$ kPa, N = 11). The $PTP_{icas}$ was also not significant different for sex (LMM, $p = 0.638$) or side (LMM, $p = 0.223$) and was $0.45 \pm 0.27$ kPa (range: $0.22$-$1.07$ kPa, N = 11). The $PTP_t$ was also not significantly different for sexes (LMM, $p = 0.057$) or side (LMM, $p = 0.395$) and was $0.15 \pm 0.19$ kPa (range: 4e-5 – $0.75$ kPa, N = 11).

     Next, we quantified the acoustic output of the adult zebra finch hemi-syrinx within the $p_b$-$p_{icas}$ control space for three parameters: fundamental frequency ($f_o$), source level (SL) and Wiener entropy (WE). In all animals, fundamental frequency was gradually modulated depending on different combinations of $p_b$ and $p_{icas}$ (Fig. 1d-f). Frequency jumps were not observed. Of all pressures, $p_t$ described the $f_o$ data best ($\Delta$BIC = -528, see Methods). The continuous, smooth increase of $f_o$ with $p_t$ can be clearly seen in Fig. 1e,f. In males, the right typically produced higher frequencies than the left as a function of $p_t$ (Fig. 1e). However, the minimal $f_o$ produced was not significantly different and was $511 \pm 102$ Hz (range: $352 - 613$ Hz, N = 5) and $513 \pm 123$ Hz (range: $312 - 625$ Hz, N = 5) for left and right hemi-syrinx respectively (Fig. 1h). To describe $f_o$ as a function of $p_t$, we split the data in to two regions (Fig. 1e,f); region S1 ($p_t = 0 - 0.75$ kPa) and region S2 ($p_t = 0.75 - 2$ kPa) and fit linear models to the data. The slope of S1 was $189 \pm 76$ Hz/kPa (range: $67 - 453$ Hz/kPa, N = 5), and was significantly higher than the S2 slope of $55 \pm 58$ Hz/kPa (range: $25 - 310$ Hz/kPa, N = 5) by $134 \pm 18$ Hz/kPa (Fig. 1j) (paired t-test, t = 5.3061, df = 9, $p$ = 5e-4). In females, the minimal

$f_o$ produced was 529 ± 84 Hz (range: 435 – 678 Hz, N = 6) for the left and 568 ± 57 Hz (range:

518 – 650 Hz, N = 6) for the right hemi-syrinx (Fig. 1f). The slope of the S1 region was 181 ±

108 Hz/kPa (range: 67 – 453 Hz/kPa, N = 6) and was also significantly higher than the S2 slope

of 55 ± 58 Hz/Pa (range: -42 – 167 Hz/kPa, N = 6) by 126 ± 51 Hz/kPa (Fig. 1j) (paired t-test,

t = 3.31, df = 10, $p$ = 0.0079). Comparing the sexes, the difference in minimal $f_o$ between left

and right seemed smaller in males than in females (Fig. 1i), but this effect was not significant

(Welch's t-test, t = -1.08, df = 6.42, p = 0.32). The slopes of both S1 and S2 were not significant

different between sex and side and were 185 ± 92 Hz/kPa (range: 25 – 453 Hz/kPa, N = 11)

and 55 ± 58 Hz/kPa (range: -42 - 167, N = 11), for S1 and S2, respectively (Fig. 1j,

Supplementary Table S1).

Source level at 1 m distance was best described by $p_b$ ($\Delta$BIC = -78,996) and increased

linearly with pressure in both sexes (Fig. 2a-c). The minimum SL (Fig. 2d) was not significantly

different for sex (LMM, $p$ = 0.149) or side (LMM, $p$ = 0.225) and was 45 ± 4 dB re 20µPa @

1m (range: 37 - 51 dB re 20µPa @ 1m, N = 11). The slope of the SL-$p_b$ relationship (Fig. 2e)

did not differ significantly between males and females (LMM, $p$ = 0.704), but was significantly

higher on the right side (5 ± 2 dB/kPa, N = 11) than the left side (4 ± 1 dB/kPa, N = 11) (LMM,

$p$ = 0.016).

WE did not show any obvious relation to $p_b$, $p_{icas}$, or $p_t$, and therefore we considered the

mean of the entire control space (Fig. 3a). The mean WE did not differ significantly between

sexes (LMM, $p$ = 0.138) or sides (LMM, $p$ = 0.661) and was -1.8 ± 0.12 dB (range: -1.6 to -2.1

dB, N = 11) (Fig. 3b).

Our setup allowed us to calculate the mechanical efficiency (ME) of sound production,

which estimates how much of the power in the air flow is transformed into sound (see Methods,

Fig. 4). We found that ME did not vary systematically with $p_b$, $p_{icas}$, or $p_t$ and therefore

considered the mean of the entire control space (Fig. 4b). ME was not significantly different

between sexes (LMM, $p$ = 0.118), but was significantly lower (LMM, $p$ = 0.027) on the right

side (-36 ± 1.8 dB, range: -39 to -33, N = 10) compared to the left (-35 ± 2.2 dB, range; -38 to

-31 dB, N = 10) (Fig. 4c).

*Acoustic control space does not change over vocal development*

Next, we tested if the output of the isolated zebra finch syrinx changed over vocal

development from 25 to 100 DPH (Fig. 5). Like in the adult syrinx, sound was produced

exclusively in the lower half of the syringeal $p_b$, $p_{icas}$ space over vocal development. The PTP$_b$

was not significantly affected by age (LMM, $p$ = 0.257), side (LMM, $p$ = 0.268), or sex (LMM,

$p = 0.558$) and was $0.9 \pm 0.24$ kPa (range: $0.53 - 1.68$ kPa, N = 32). $PTP_{icas}$ demonstrated a

small but significant increase with age (LMM, $p = 0.002$), but no significant difference between

sexes (LMM, $p = 0.762$) or sides (LMM, $p = 0.274$), and went from $0.21 \pm 0.8$ kPa at 25 DPH

(range: $0.006 - 0.25$ kPa, N = 8) to $0.37 \pm 0.25$ kPa at 100 DPH (range: $0.21 - 1.1$ kPa, N =

11).

Consistently, $PTP_t$ also demonstrated a small but significant decrease with age from 0.17

$\pm 0.08$ kPa at 25 DPH (range: $0.06 - 0.31$ kPa, N = 8) to $0.03 \pm 0.06$ kPa at 100 DPH (range:

4e-5– 0.16 kPa, N = 11), and marginally, but significantly lower in males ($0.0846 \pm 0.0981$

kPa) compared to females ($0.1047 \pm 0.0875$).

We examined changes in the syringeal $p_b$-$p_{icas}$ control space for three acoustic

parameters: fundamental frequency ($f_o$), source level (SL) and Wiener entropy (WE). Over

development, $f_o$ was also gradually modulated within the $p_b$-$p_{icas}$ space and did not exhibit

frequency jumps (Fig. 5). In males and females, the minimal $f_o$ produced by each hemi-syrinx

*in vitro* did not change significantly from 25 DPH to 100 DPH (Fig. 5b), but the left hemi-

syrinx produced a significantly lower minimal $f_o$ than the right (males: L: $487 \pm 106$ Hz, N =

18, R: $519 \pm 104$ Hz, N = 18, females: L: $480 \pm 83$ Hz, N = 15, R: $590 \pm 69$ Hz N = 15) (LMM,

$p < 0.001$). Interestingly, in 5 out of 18 males the left-right difference was reversed.

The slope of the $f_o$-$p_t$ relationship in region S1 did not significantly change with age

(LMM, $p = 0.651$), sex (LMM, $p = 0.630$), or side (LMM, $p = 0.270$) and was $91 \pm 109$ Hz/kPa

(range: $-204 - 250$ Hz/kPa, N = 32). The slope in region S2 was significantly different between

ages (LMM, $p = 0.0273$), but not for sex (LMM, $p = 0.464$), or side (LMM, $p = 0.474$), and

was $88 \pm 43$ Hz/kPa (range: $31 - 146$ Hz/kPa, N = 8), $62 \pm 28$ Hz/kPa (range: $18 - 105$ Hz/kPa,

N = 7), $70 \pm 46$ Hz/kPa (range: $5 - 149$ Hz/kPa, N = 7), and $65 \pm 50$ Hz/kPa (range: $-43 - 135$,

N = 11) for 25, 50, 75, and 100 DPH respectively.

Source level increased with $p_b$ for all animals (Fig. 6a,e), but the minimal source level

did not change significantly with age (LMM, $p = 0.680$), sex (LMM, $p = 0.374$), or side (LMM,

$p = 0.271$) and was $44 \pm 4$ dB re20μPa @ 1m (range: $36 - 51$ dB, N = 32) (Fig. 6b-c). The SL-

$p_b$ slope also did not change with age (LMM, $p = 0.83$), or sex (LMM, $p = 0.874$), but was

significantly less steep (LMM, $p = 0.037$) on the left ($4.1 \pm 1.0$ dB/kPa, range: $1.9 - 6.2$ dB/kPa,

N = 32) versus right hemi-syrinx ($4.7 \pm 1.6$ dB/kPa, range: $0.9 - 7.4$ dB/kPa, N = 32) by $0.6 \pm$

0.3 dB (N = 32).(Fig. 6d). Mean WE did not change significantly for any of the parameters

tested (LMM, age: $p = 0.070$, sex: $p = 0.979$, side: $p = 0.084$), and was $-1.8 \pm 0.1$ dB (range: -

2.1 to -1.6 dB, N = 32; Fig. 7a-b, e-f).

Next, we calculated the mechanical efficiency (ME) to test if the ability of the syrinx to
convert mechanical power to sound changed over song development (Fig. 7c,g). ME did not
significantly change with sex (LMM, $p$ = 0.091) or side (LMM, $p$ = 0.101), but demonstrated
a small but significant decrease (LMM, $p$ = 0.046) from -33 ± 1.7 dB (range: -37 to -31 dB, N
= 8) at 25 DPH to -35 ± 1.8 dB (range: -39 to -33 dB, N = 10) in adults (Fig. 7d, h).
Taken together, we observed that the acoustic output of the syrinx control space did not
change considerably over development in males and females *in vitro*. To test whether these *in*
*vitro* results are representative for the performance of the syrinx during *in vivo* sound
production, we measured the acoustic output over development *in vivo*. Because both SL and
WE are affected by vocal tract filtering properties, we focused on fundamental frequency. We
measured the lowest fundamental frequency produced by males with unilateral syrinx
denervation at 50 and 100 DPH (see Methods). Consistent with the *in vitro* data, the minimal
$f_o$ did not change significantly (Welch's t-test, t = 0.49, df = 6.60, p = 0.64) with age (Right
hemi-syrinx: 555 ± 70 Hz (N = 6) and 524 ± 98 Hz (N = 5) for 50 and 100 DPH respectively.
Left adult hemi-syrinx: 516 ± 54 Hz (N = 5) and also did not differ between the left (intact)
and right (denervated) hemi-syrinx. Furthermore, the *in vivo* values were consistent with the *in*
*vitro* data (Fig. 5b), strongly suggesting that the output of the syrinx *in vitro* is representative
for the *in vivo* situation.
380
*Morphological changes over development*
We lastly investigated if the medial vibratory mass (MVM) dimensions change over
development. We focused on non-invasive photographic techniques because detailed
histological or material property testing was beyond the scope of the present study. The tension
in the MVM can be increased by the shortening of two syringeal muscles that insert on the
*medio-ventral cartilage* (MVC) and *medio-dorsal cartilage* (MDC)[21] (Fig. 8a-c). A third
cartilage, the *lateral dorsal cartilage* (LDC), is embedded within the MVM. A thickening of
the tissue between the MVC and LDC is called the medial labium. The two syringeal muscles
change the tension of the MVM by changing the distance between 1) MVC and LDC and 2)
LDC and MDC (Fig. 8b,c). The MVC - LDC length did not change with sex (LMM, $p$ = 0.72)
or age (LMM, $p$ = 0.446), but was significantly (LMM, $p$ = 0.002) longer on the left (1.13 ±
0.10 mm; range: 965 – 1297 μm, N = 29) compared to the right (1.05 ± 0.11 mm, range: 798 –
1363 μm, N = 29) (Fig. 8f). The LDC - MDC length also did not change with sex (LMM, $p$ =
0.982) or age (LMM, $p$ = 0.191), but it was significantly (LMM, $p$ = 0.149) longer on the left

(0.71 ± 0.16 mm, range: 387 – 1107 μm, N = 29) compared to right (0.65 ± 0.13 mm, range:

365 – 858 μm, N = 29) (Fig. 8g). The area of the LDC was smaller (LMM, $p < 0.001$) on the

left (4.1e4 ± 2.2e4 μm$^2$, range: 1978 – 9.9e4 μm$^2$, N = 29) compared to the right (6.2e4 ± 3.0e4

μm$^2$, range: 11867 – 1.1e5 μm$^2$, N = 29) hemi-syrinx and increased over development (LMM,

$p = 0.048$; Fig. 8h). Lastly, we approximated overall growth of the syrinx as the distance

between both ends of the B4 cartilage in the bronchus, and found no significant change for side

(LMM, $p = 0.14$), sex (LMM, $p = 0.17$), or age (LMM, $p = 0.8$) and was 1.85 ± 0.23 mm (range:

1280 – 2196 μm, N = 29; Fig. 8i).

**Discussion**

As songbirds learn to sing, both their song system and vocal organ are undergoing

postnatal changes that lead to dramatic changes in vocal behavior. Here we show that the

acoustic output of the sound generators within the zebra finch syrinx does not change

considerably over vocal development. This strongly suggests that the observed acoustic

changes during vocal development are caused by changes in the motor control pathway, from

song system circuitry to muscle force, and not by changes in the avian analog of the vocal

folds. In other words, the size and properties of the instrument are not changing, but its player

is.

We show that both sides of the syrinx, or hemi-syrinxes, produce sound in a well-

defined consistent pressure control space. The pressure space is bound by phonation threshold

pressures (PTP) for $p_b$, $p_{icas}$ and $p_t$, which are the same for both sexes and sides. The PTP$_b$

values in adults of 1.01 ± 0.28 kPa reported here are consistent with earlier reported PTP$_b$

values of 1.2 kPa in situ[45] and *in vivo*[31]. The possibility to control the two respiratory pressures

($p_b$ and $p_{icas}$) independently is very different from laryngeal systems where no air sacs directly

apply force on the vocal folds. To what extent birds can actively and independently control the

magnitude of $p_t$ *in vivo* during song remains unknown. Certainly, positive pressurization of $p_{icas}$

is an essential condition to achieve syringeal vibration in all investigated species *in vivo*, (e.g.,

zebra finches[29] and *in vitro*[15,46,47], and dynamic $p_t$ changes up to 1 kPa occur during

vocalizations in ringdoves[34,48–50].

Our data show that source level and fundamental frequency are modulated by

respiratory pressures. Source level was driven by bronchial pressure, consistent with laryngeal

voiced sound production[19]. Fundamental frequency ($f_o$) was set predominantly by $p_t$, consistent

with the idea that $p_t$ acts as a force on the MVM that increases tension[51,52], which has also been

observed in collapsible tubes[54]. Interestingly, especially close to the phonation onset $f_o$

modulation is steep (200 Hz/kPa) and a dynamic $p_t$ difference of 1 kPa could in zebra finches

lead to $f_o$ modulation of comparable magnitude to full stimulation of the ventral syringeal

muscle[15]. In contrast, Wiener Entropy (WE) was not modulated systematically in the pressure

space. However, over the course of song learning WE has been shown to change over multiple

time scales, from within a day to weeks and months[2,3,55] and increased entropy variance is

linked to better learning success[3,55]. These data suggest that WE is under control of the motor

systems of (i) the syrinx, by modulating collision force and thereby spectral slope of the sound

source[56], and (ii) the upper vocal tract, by changing resonance properties and thereby frequency

content of the radiated sound[57,58].

Mechanical efficiency may have been an important selecting factor during the evolution

of the syrinx as a vocal organ[59] and our data confirms that the mechanical efficiency of voiced

sound production is indeed high in the avian syrinx compared to the mammalian larynx. Our

data shows that about 0.05% (-33 dB) of the flow energy is converted into sound, while in

mammals, laryngeal ME is only about $1.10^{-3}$ to 0.01% (-50 to -40 dB) in tigers (Titze et al.,

2010) and $3.10^{-5}$ to $1.10^{-3}$ % (-65 to -50 dB) in marmosets[6]). ME in the rooster has been reported

to be as high as 1.6% *in vivo*[61]. The ME of the zebra finch *in vivo* is likely to be higher than we

measured *in vitro*, because our preparation does not include a vocal tract. The reported SL of

the zebra finch is about 70-80 dB re 20µPa @ 1m[62], while the SL of our isolated syrinx was

20-30 dB SPL lower. A conservative estimate of 10 dB SL increase (i.e. RMS is 3 times higher)

while maintaining the same aerodynamic power, results in a 10-fold increase in radiated

acoustic power and a ME of 0.5% (-23 dB). This is already comparable to the very high

efficiency reported by Brackenbury[61]. Furthermore, we observed slight differences of ME over

development, but these effects were not as pronounced as the changes observed in marmosets,

where the adult larynx produces louder and more efficient vocalisations than the infant larynx[6].

Thus, the ME of the zebra finch syrinx is very high already at 25 DPH and remains unaffected

by vocal development.

Our data shows that acoustic output of the syrinx is modulated smoothly within the

pressure boundaries: both SL and $f_o$ increased continuously with bronchial and air sac

pressures. Thus when driven by physiologically relevant pressures, this nonlinear dynamic

system exhibits only one bifurcation from steady (no vibration and no sound) to flow-induced

self-sustained vibration, without additional bifurcations, such as period doublings or jumps to

deterministic chaos[45,63–66]. Earlier work already showed that in a subset of the pressure control

space *in situ*, additional bifurcations did not occur[45]. Moreover, the supposed bifurcations

during song *in vivo*, e.g. frequency jumps, were shown to not be actual bifurcations[45], but more

likely millisecond scale modulation driven by superfast syringeal muscle[67]. Here we also show

that in the independently controlled and densely sampled pressure control space of the zebra

finch syrinx *in vitro*, additional bifurcations do not occur. Thus, driven by physiologically

relevant pressures, the acoustic output of the syrinx is continuous, which simplifies sensory-

motor control[45].

The stable acoustic output of the sound generators over development suggests that its

mechanical properties are not changing[22,23] and acoustic output[41,68,69] are common in songbirds.

In all investigated species to date, except for the Bengalese finch[15,69], the left hemi-syrinx

produces lower frequencies.

The $f_o$ of sound is determined by positioning of and driving pressures on the vocal folds,

as well as their resonance properties[20]. Small changes in material properties can drive changes

in vocalizations over development in marmosets[6]. Because we show that minimum frequency

does not significantly change over development *in vitro* and *in vivo*, our data strongly suggest

that the effective resonance properties of the syringeal sound generators when driven by

physiological realistic pressures do not change over vocal development. Next to resonance

properties, the change in MVM stiffness as response to shortening is also likely changing very

little over development. Although we didn't measure directly how syringeal muscle force

modulates MVM resonance properties over development, the $f_o$-$p_t$ relationship (S1) can be seen

as a proxy of the effect of MVM lengthening by e.g. VS contraction, as $p_t$ exerts a force on the

MVM that causes changes in strain[32]. Taken together, the MVM vibration frequencies and its

response to external force do not change, which strongly suggest that its material properties

also do not change over development.

We propose that syringeal sound production on the one hand, and its control on the

other, are shaped differentially by genetic and environmental factors. First, we propose that the

development and maintenance of the sound generators is predominantly under genetic control.

Second, we propose that the syringeal muscles and skeletal properties are heavily influenced

by use, which can ultimately lead to a sexually dimorph syrinx.

Our data suggest that MVM mechanical properties are set and maintained by genetic

factors. First, a salient feature in our dataset is that the adult left hemi-syrinx produces a lower

minimal $f_o$ compared to the right in both sexes, corroborating earlier findings[20]. In most

songbirds both hemi-syrinxes contribute to the vocal repertoire[70,71] and left-right differences in

labial morphology[22,23] and acoustic output[41,68,69] are common in songbirds. In all investigated

species to date, except for the Bengalese finch[15,69], the left hemi-syrinx produces lower

frequencies. Here we show for the first time that differences between left and right sound

generators, size and acoustic output, are already established at 25 DPH. Second, we show that

neither the morphology, nor the acoustic output of the bilateral sound generators changes over

development. Thus, the MVM properties have matured already at 25 DPH. This latter feature

makes the songbird system a wonderful model system to study vocal development compared

to mammalian systems where larynx maturation strongly contributes to vocal output[6,25–27].

Third, MVM properties remain constant even though they are colliding billions of times over

their lifetime and there must be a large variability in use between individuals and sexes. In

human vocal folds, strong collisions can lead to several types of lesions that severely affect

vocal output[72]. Thus, ultrastructural repair of vibratory tissues must lead to maintenance of

their properties, but what cellular and molecular mechanisms underlie this maintenance in birds

remains unknown. Taken together, these observations support the ideas that the sound

generators of the syrinx are matured at the onset of vocal learning, their composition remains

stable and is species specific, which suggests that their composition and dynamical behavior is

set by genetic factors and not affected by use. The zebra finch syrinx is thus a stable instrument

which's postnatal development does not add further complexity to song learning

In stark contrast, syringeal muscles are changing functionally over postnatal

development during song learning. Until 25 DPH the syrinx of male and female zebra finches

is not clearly distinguishable. However, after 25 DPH, the syringeal muscles start to exhibit

sexually dimorphic features, such as increased muscle mass[11] and speed in males[12], which leads

to a different action potential to force transform[14]. Furthermore, muscle force exerted on bones

is known to redirect bone deposition and thus can lead to significant bone remodelling[73]. Taken

together, we propose that in songbirds syringeal muscle activity and resulting forces acting on

the syrinx are crucially important drivers to the sexual dimorphism of bones and cartilages of

the syringeal skeleton. This implies that most anatomical changes are caused by muscle use

and training driven by the sexually dimorphic songbird brain (i.e., song system). As such the

instrument, i.e. the vocal folds, are not changing, but the player is.

**References**

1.  Takahashi, D. Y. *et al.* The developmental dynamics of marmoset monkey vocal production. *Science.* **349**, 734–738 (2015).

2.  Tchernichovski, O., Mitra, P. P., Lints, T. & Nottebohm, F. Dynamics of the vocal imitation process: How a zebra finch learns its song. *Science.* **291**, 2564–2569 (2001).

3.  Kollmorgen, S., Hahnloser, R. H. R. & Mante, V. Nearest neighbours reveal fast and slow components of motor learning. *Nature* **577**, 526–530 (2020).

4.  Goldstein, M. H., King, A. P. & West, M. J. Social interaction shapes babbling: Testing parallels between birdsong and speech. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 8030–8035 (2003).

5.  Oller, D. K. *et al.* Functional flexibility of infant vocalization and the emergence of language. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 6318–6323 (2013).

6.  Zhang, Y. S., Takahashi, D. Y., Liao, D. A., Ghazanfar, A. A. & Elemans, C. P. H. Vocal state change through laryngeal development. *Nat. Commun.* **10**, 1–12 (2019).

7.  Fee, M. S. & Scharff, C. The songbird as a model for the generation and learning of complex sequential behaviors. *ILAR J.* **51**, 352–377 (2010).

8.  Brainard, M. S. & Doupe, A. J. Translating birdsong: Songbirds as a model for basic and applied medical research. *Annu. Rev. Neurosci.* **36**, 489–517 (2013).

9.  Liu, W. C., Wada, K. & Nottebohm, F. Variable food begging calls are harbingers of vocal learning. *PLoS One* **4**, e5929 (2009).

10. Darshan, R., Wood, W. E., Peters, S., Leblois, A. & Hansel, D. A canonical neural mechanism for behavioral variability. *Nat. Commun.* **8**, 15415 (2017).

11. Godsave, S. F., Lohmann, R., Vloet, R. P. M. & Gahr, M. Androgen receptors in the embryonic zebra finch hindbrain suggest a function for maternal androgens in perihatching survival. *J. Comp. Neurol.* **453**, 57–70 (2002).

12. Mead, A. F. *et al.* Fundamental constraints in synchronous muscle limit superfast motor control in vertebrates. *Elife* **6**, e29425 (2017).

13. Adam, I. & Elemans, C. P. H. Sensory and Motor Systems Vocal Motor Performance in Birdsong Requires Brain-Body Interaction. *eNeuro* **6**, (2019).

14. Adam, I. & Elemans, C. P. H. Increasing muscle speed drives changes in the neuromuscular transform of motor commands during postnatal development in songbirds. *J. Neurosci.* **40**, 6722–6731 (2020).

15. Elemans, C. P. H. *et al.* Universal mechanisms of sound production and control in birds and mammals. *Nat. Commun.* **6**, 1–13 (2015).

16.    Jiang, W. *et al.* High-fidelity continuum modeling predicts avian voiced sound
production. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 4718–4723 (2020).

17.    Goller, F. & Larsen, O. N. A new mechanism of sound generation in songbirds. *Proc.*
*Natl. Acad. Sci.* **94**, 14787–14791 (1997).

18.    Mindlin, G. B. & Laje, R. *The Physics of Birdsong*. Springer Science & Business Media
(2006).

19.    Titze, I. R. *Principles of voice production*. (National Center for Voice and Speech,
2000).

20.    Fee, M. S. Measurement of the linear and nonlinear mechanical properties of the oscine
syrinx: Implications for function. *J. Comp. Physiol. A Neuroethol. Sensory, Neural,*
*Behav. Physiol.* **188**, 829–839 (2002).

21.    Düring, D. N. *et al.* The songbird syrinx morphome: A three-dimensional, high-
resolution, interactive morphological map of the zebra finch vocal organ. *BMC Biol.* **11**,
1 (2013).

22.    Riede, T. & Goller, F. Functional morphology of the sound-generating labia in the syrinx
of two songbird species. *J. Anat.* **216**, 23–36 (2010).

23.    Riede, T. & Goller, F. Morphological basis for the evolution of acoustic diversity in
oscine songbirds. *Proc. R. Soc. B Biol. Sci.* **281**, 20132306–20132306 (2014).

24.    Švec, J. G., Horáček, J., Šram, F. & Veselý, J. Resonance properties of the vocal folds:
In vivo laryngoscopic investigation of the externally excited laryngeal vibrations. *J.*
*Acoust. Soc. Am.* **108**, 1397–1407 (2000).

25.    Sato, K., Hirano, M. & Nakashima, T. Age-related changes of collagenous fibers in the
human vocal fold mucosa. *Ann. Otol. Rhinol. Laryngol.* **111**, 15–20 (2002).

26.    Hammond, T. H., Gray, S. D., Butler, J., Zhou, R. & Hammond, E. Age- and gender-
related elastin distribution changes in human vocal folds. *Otolaryngol. - Head Neck*
*Surg.* **119**, 314–322 (1998).

27.    Hammond, T. H. & Gray, S. D. A study of age- and gender-related elastin distribution
changes in human vocal folds. *Otolaryngol. - Head Neck Surg.* **117**, 100 (1997).

28.    Gaunt, A. S., Stein, R. C. & Gaunt, S. L. L. L. Pressure and air flow during distress calls
of the starling, Sturnus vulgaris (Aves;Passeriformes). *J. Exp. Zool.* **183**, 241–261
(1973).

29.    Amador, A. & Margoliash, D. A Mechanism for Frequency Modulation in Songbirds
Shared with Humans. *J. Neurosci.* **33**, 11136–11144 (2013).

30.    Setterwall, C. G. Syrinx hos polymyoda passeres. *Gleerupska Univ.* 1–105 (1901).

31.   Riede, T. & Goller, F. Peripheral mechanisms for vocal production in birds - differences
and similarities to human speech and singing. *Brain Lang.* **115**, 69–80 (2010).

32.   Düring, D. N., Knörlein, B. J. & Elemans, C. P. H. H. In situ vocal fold properties and
pitch prediction by dynamic actuation of the songbird syrinx. *Sci. Rep.* **7**, 11296 (2017).

33.   Srivastava, K. H., Elemans, C. P. H. & Sober, S. J. Multifunctional and Context-
Dependent Control of Vocal Acoustics by Individual Muscles. *J. Neurosci.* **35**, 14183–
14194 (2015).

34.   Elemans, C. P. H. *et al.* Syringeal muscles fit the trill in ring doves (Streptopelia risoria
L.). *J. Exp. Biol.* **209**, 965–977 (2006).

35.   de Cheveigné, A. & Kawahara, H. YIN, a fundamental frequency estimator for speech
and music. *J. Acoust. Soc. Am.* **111**, 1917–1930 (2002).

36.   The Mathworks Inc. MATLAB - MathWorks. *www.mathworks.com/products/matlab*
(2016).

37.   R Development Core Team. A Language and Environment for Statistical Computing. *R
Found. Stat. Comput.* (2018).

38.   Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis. *Nat.
Methods* **9**, 676–682 (2012).

39.   Tchernichovski, O., Nottebohm, F., Ho, C. E., Pesaran, B. & Mitra, P. P. A procedure
for an automated measurement of song similarity. *Anim. Behav.* (2000).

40.   Roy, A. & Mooney, R. Auditory plasticity in a basal ganglia-forebrain pathway during
decrystallization of adult birdsong. *J. Neurosci.* **27**, 6374–6387 (2007).

41.   Williams, H., Crane, L. A., Hale, T. K., Esposito, M. A. & Nottebohm, F. Right-side
dominance for song control in the zebra finch. *J. Neurobiol.* **23**, 1006–1020 (1992).

42.   Riede, T., Fisher, J. H. & Goller, F. Sexual dimorphism of the zebra finch syrinx
indicates adaptation for high fundamental frequencies in males. *PLoS One* **5**, e11368
(2010).

43.   Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in
Linear Mixed Effects Models . *J. Stat. Softw.* (2017).

44.   Bates, D., Maechler, M., Bolker, B. & Walker, S. lme4: linear mixed-effects models
using S4 classes. R package version 1.1-6. *R* (2014).

45.   Elemans, C. P. H., Laje, R., Mindlin, G. B. & Goller, F. Smooth operator: Avoidance of
subharmonic bifurcations through mechanical mechanisms simplifies song motor
control in adult zebra finches. *J. Neurosci.* **30**, 13246–13253 (2010).

46.   Rüppell, W. Physiologie und Akustik der Vogelstimme. *J. für Ornithol.* **81**, 433–542

(1933).

47.    Abs, M. On the Bioacoustics of the Breaking of the Birds Voice. *Zool. JAHRBUCHER-*

628           *ABTEILUNG FUR Allg. Zool. UND Physiol. DER TIERE* **84**, 289–382 (1980).

629    48.    Gaunt, A., Gaunt, S. L. & Casey, R. Syringeal Mechanics Reassessed: Evidence from

630           Streptopelia. *Auk* **99**, 474–494 (1982).

631    49.    Beckers, G. J. L., Nelson, B. S. & Suthers, R. A. Vocal-tract filtering by lingual

632           articulation in a parrot. *Curr. Biol.* **14**, 1592–1597 (2004).

633    50.    Elemans, C. P. H., Zaccarelli, R. & Herzel, H. Biomechanics and control of vocalization

634           in a non-songbird. *J. R. Soc. Interface* **5**, 691–703 (2008).

635    51.    Elemans, C. P. H. H., Muller, M., Larsen, O. N. & Van Leeuwen, J. L. Amplitude and

636           frequency modulation control of sound production in a mechanical model of the avian

637           syrinx. *J. Exp. Biol.* **212**, 1212–1224 (2009).

638    52.    Elemans, C. P. H., Heeck, K. & Muller, M. Spectrogram analysis of animal sound

639           production. *Bioacoustics* **18**, 183–212 (2008).

640    53.    Bertramt, C. D., Pedley, T. J. J., Bertram, C. D. D. & Pedley, T. J. J. A mathematical

641           model of unsteady collapsible tube behaviour. *J. Biomech.* **15**, 39–50 (1982).

642    54.    Bertram, C. D. Flow phenomena in floppy tubes. *Contemp. Phys.* **45**, 45–60 (2004).

643    55.    Derégnaucourt, S., Mitra, P. P., Fehér, O., Pytte, C. & Tchernichovski, O. How sleep

644           affects the developmental learning of bird song. *Nature* **433**, 710–716 (2005).

645    56.    Hunter, E. J., Titze, I. R. & Alipour, F. A three-dimensional model of vocal fold

646           abduction/adduction. *J. Acoust. Soc. Am.* **115**, 1747–1759 (2004).

647    57.    Ohms, V. R., Beckers, G. J. L., ten Cate, C. & Suthers, R. A. Vocal tract articulation

648           revisited: the case of the monk parakeet. *J. Exp. Biol.* **215**, 85–92 (2012).

649    58.    Riede, T., Suthers, R. A., Fletcher, N. H. & Blevins, W. E. Songbirds tune their vocal

650           tract to the fundamental frequency of their song. *Proc. Natl. Acad. Sci.* **103**, 5543–5548

651           (2006).

652    59.    Riede, T., Thomson, S. L., Titze, I. R. & Goller, F. The evolution of the syrinx: An

653           acoustic theory. *PLoS Biol.* **17**, 1–22 (2019).

654    60.    Ingo R. Titze, W. Tecumseh Fitch, Eric J. Hunter, Fariborz Alipour, Douglas

655           Montequin, Douglas L. Armstrong, JoAnn McGee, E. J. W. Vocal power and pressure-

656           flow relationships in excised tiger larynges. *J. Exp. Biol.* **213**, 3866–3873 (2010).

657    61.    Brackenbury, J. H. Physiological energetics of cock-crow. *Nature* **270**, 433–435 (1977).

658    62.    Brumm, H. & Ritschard, M. Song amplitude affects territorial aggression of male

659           receivers in chaffinches. *Behav. Ecol.* **22**, 310–316 (2011).

63.  Wilden, I., Tembrock, G., Herzel, H. & Peters, G. Subharmonics, biphonation, and
deterministic chaos in mammal vocalization. *Bioacoustics* **9**, 171–196 (1998).

64.  Laje, R., Gardner, T. J. & Mindlin, G. B. Neuromuscular control of vocalizations in
birdsong: A model. *Phys. Rev. E - Stat. Physics, Plasmas, Fluids, Relat. Interdiscip.*
*Top.* **65**, 051921 (2002).

65.  Gardner, T., Cecchi, G., Magnasco, M., Laje, R. & Mindlin, G. B. Simple motor gestures
for birdsongs. *Phys. Rev. Lett.* **87**, 208101 (2001).

66.  Amador, A., Perl, Y. S., Mindlin, G. B. & Margoliash, D. Elemental gesture dynamics
are encoded by song premotor cortical neurons. *Nature* **495**, 59–64 (2013).

67.  Elemans, C. P. H., Mead, A. F., Rome, L. C. & Goller, F. Superfast vocal muscles
control song production in songbirds. *PLoS One* **3**, 6–11 (2008).

68.  Nottebohm, F. & Nottebohm, M. E. Left hypoglossal dominance in the control of canary
and white-crowned sparrow song. *J. Comp. Physiol.* **108**, 171–192 (1976).

69.  Secora, K. R. *et al.* Syringeal specialization of frequency control during song production
in the bengalese finch (lonchura striata domestica). *PLoS One* **7**, e34135 (2012).

70.  Suthers, R. A. How birds sing and why it matters. in *Nature's Music: The Science of*
*Birdsong* 272–295 (2004).

71.  Suthers, R. A. & Zollinger, S. A. From brain to song: the vocal organ and vocal tract.
*Neuorscience of Birdsong* 78–98 (2008).

72.  Titze, I. R. & Hunter, E. J. Comparison of vocal vibration-dose measures for potential-
damage risk criteria. *J. Speech, Lang. Hear. Res.* **58**, 1425–1439 (2015).

73.  Robling, A. G., Castillo, A. B. & Turner, C. H. Biomechanical and molecular regulation
of bone remodeling. *Annu. Rev. Biomed. Eng.* **8**, 455–498 (2006).

**Acknowledgments**

The authors wish to thank Torben Christensen and Sonja Jakobsen for technical support, Emil B. Hansen for help with breeding, and John Jackson for assistance with statistical analysis.

**Author Contributions**

A.M., I.A. & C.E. conceived the study. A.M., I.A. & C.E. developed technical methods. I.A. & C.E. provided reagents and materials. A.M. & P.L. performed *in vitro* experiments and collected all data. I.A., P.S. & C.E. performed *in vivo* experiments and collected all data. A.M., I.A. & C.E. analysed the data. A.M. prepared all figures and tables. A.M., I.A. & C.E. wrote the manuscript. All authors read and approved the final version of the manuscript.

**Additional Information:** The authors declare that they have no competing interests.

**Funding**

This work was supported by the Danish Research Council (Grant DFF 5051-00195), the Carlsberg Foundation (Grant CF17-0949) to I.A. and Novo Nordisk Foundation (Grant NNF17OC0028928) to C.P.H.E

**Data Availability**

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

 **Figures and legends**

 Figure 1

**Figure 1. The adult zebra finch syrinx produces sound in a well-defined pressure control space with higher frequencies on the right. (a)** Ventral view of the male syrinx. One of the two sound generators are indicated by a dashed circle. **(b)** Schematic view on the sound generator's lateral vibratory mass. **(c)** Schematic sagittal cross-section through the syrinx of the left bronchus, indicating the movement of pressures around and through the syrinx. Example raw data of sound production induced by bronchial and interclavicular air sac pressures **(d)** Example of fundamental frequencies produced in the pressure control space for the right hemi-syrinx of an adult male zebra finch (100 DPH). Phonation threshold pressures (PTP) are indicated for: $PTP_{icas}$ (grey dot), and $PTP_b$ (red dot). Transmural pressure ($p_t$) is indicated by white dashed lines for $p_t = 0$, 1 and 3 kPa. **(ef)** Frequency increases with $p_t$ for both the left (dark red, dark blue) and right hemi-syrinx (light red, light blue) in males and **(c)** females, respectively. Black lines indicate linear regressions for region S1 ($0 < p_t < 0.75$) and S2 ($p_t > 0.75$). The phonation threshold pressure (PTP) values for individual $p_{icas}$ ramps are indicated by black circles. **(g)** PTP for $p_b$, $p_t$, and $p_{icas}$ (kPa), for the left and right hemi-syrinx. **(h)** Minimum frequency produced by the right hemi-syrinx was higher than the left hemi-syrinx (p = 0.09, see Table S1) for most individuals. **(i)** The absolute difference of minimal $f_o$ was not significantly different between males and females. **(j)** Slopes for the left and right hemi-syrinx did not differ significantly between sides but were significantly steeper in S1 than in S2. For statistics see Supplementary Table S1. *, p<0.05; **, p<0.01; ***, p<0.001.

Figure 2

**Figure 2. Source level increases with bronchial pressure in the adult zebra finch hemi-**
**syrinx. (a)** Example of source level (SL) produced in the pressure control space for the left
hemi-syrinx of an adult male zebra finch (100 DPH). Source level increases with $p_b$ for both
the left (dark red, dark blue) and right hemi-syrinx (light red, light blue) in **(b)** males and **(c)**
females, respectively. Black dashed lines indicate linear regressions (Left: long dash; Right:
short dash). **(d)** Minimum SL does not differ significantly between the left and right hemi-
syrinx or between males and females. **(e)** The SL increase with bronchial pressure was
significantly steeper for the right hemi-syrinx than the left hemi-syrinx. For statistics see
Supplementary Table S1. *, p<0.05.

Figure 3

**Figure 3. Wiener entropy is not affected by pressure in the adult zebra finch hemi-**
**syrinx**. **(a)** Wiener entropy (WE) produced in the pressure control space for the left hemi-
syrinx of an adult male zebra finch (100 DPH). **(b)** Mean WE of the left and right hemi-
syrinx was not significantly different between male and female zebra finches or for left and
right hemi-syrinxes. For statistics see Supplementary Table S1.

Figure 4. Mechanical efficiency is not affected by respiratory pressure, but lower on the right side of adult zebra finch. (a) Example of one $p_b$-ramp (from top to bottom) received sound pressure (Pa), bronchial and interclavicular air sac pressures (kPa), acoustic and aerodynamic power (W), and mechanical efficiency (ME) (dB). (b) Mechanical efficiency in the pressure control space for the left hemi-syrinx of an adult male zebra finch (100 DPH). (c, d) Mechanical efficiency was significantly higher for the left hemi-syrinx than the right hemi-syrinx in both sexes. For statistics see Supplementary Table S1. *, $p < 0.05$.

Figure 5

**Figure 5. Minimal frequency does not change over zebra finch song development**. **(a)**
Fundamental frequency in the pb picas pressure control space for the left and right hemi-syrinx in
males over vocal development $f_o$ values are averaged over all individuals within a group (insets
S.D.). **(b)** Minimum $f_o$ does not change with age in males and females. **(c)** Minimum frequency was
significantly different between the left and right hemi-syrinx, regardless of age or sex. **(d)** Slope S1
for the relationship between $f_o$ and $p_t$ was not significantly different for side. **(e)** Pressure control
spaces *in vitro* for the left and right hemi-syrinx in females. For statistics see Supplementary Table
S2. ***, p<0.001.

Figure 6

**Figure 6. Minimum source level (SL) does not change over zebra finch song development**. **(a)**

Mean SL pressure control spaces *in vitro* for the left and right hemi-syrinx in males (inset S.D.).

Minimum SL produced for **(b)** males and **(c)** females does not change over development, nor was it

significantly different between sides. **(d)** Slope for the relationship between SL and $p_b$ was

significantly steeper on the right compared to the left hemi-syrinx. **(e)** Mean SL pressure control

spaces *in vitro* for the left and right hemi-syrinx in females. For statistics see Supplementary Table

S2. *, $p < 0.05$.

**Figure 7. Wiener entropy and mechanical efficiency do not change over zebra finch song**
**development**. **(a)** Wiener entropy (WE) for the left and right hemi-syrinx in males age 25 and 100
DPH. **(b)** WE did not change for males over development. **(c)** Mechanical efficiency (ME) for the
left and right hemi-syrinx in males age 25 and 100 DPH. **(d)** The mean ME produced for males
significantly differed over development. **(e)** Wiener entropy (WE) for the left and right hemi-syrinx
in females age 25 and 100 DPH. **(f) In females,** WE did not significantly change over development.
**(g)** Mechanical efficiency (ME) for the left and right hemi-syrinx in females age 25 and 100 DPH.
**(h)** The mean ME of female syrinxes decreased significantly over development. For statistics see
Supplementary Table S2. *, $p < 0.05$.

Figure 8

789
790

**Figure 8. Morphological changes of the medial labia over zebra finch song development**. **(a)**

Schematic of lateral vibratory mass. **(b)** Physiological landmarks in the ML, the *medio-ventral*

*cartilage* (MVC), the *medio-dorsal cartilage* (MDC) and the *lateral dorsal cartilage* (LDC) and **(c)**

the distance between landmarks. **(d,e)** Example images of MTMs of left/right and male/female hemi-

syrinx. **(f)** MVC-LDC length was significantly different between sides, but not over development.

**(g)** LDC-MDC length significantly differed between sides, but not over development. **(h)** LDC area

for the four investigated age groups was significantly different and differed significantly between

the two sides. **(i)** Bronchial width did not significantly differ over development or side.  For statistics

see Supplementary Table S3. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.