The adaptive value of tandem communication in ants: insights from an agent-based model

Natascha Goy[1], Simone M. Glaser[1] & Christoph Grüter[1,2]

[1]Institute of Organismic and Molecular Evolutionary Biology, Johannes-Gutenberg University of Mainz, Mainz, Germany.

[2]Present address: School of Biological Sciences, University of Bristol, 24 Tyndall Avenue, BS8 1TQ Bristol, UK

Correspondence: natascha.goy@posteo.de

Keywords: communication, tandem running, agent-based model, *Temnothorax nylanderi*

Abstract

Social animals often share information about the location of resources, such as a food source or a new nest-site. One well-studied communication strategy in ants is tandem running, whereby a leader guides a recruit to a resource. Tandem running is considered an example of animal teaching because a leader adjusts her behaviour and invests time to help another ant to learn the location of a resource more efficiently. Tandem running also has costs, such as waiting inside the nest for a leader and a reduced walking speed. Whether and when these costs outweigh the benefits of tandem running is not well understood. We developed an agent-based simulation model to investigate the conditions that favour communication by tandem running during foraging. We predicted that the spatio-temporal distribution of food sources, colony size and the ratio of scouts and recruits affect colony foraging success. Our results suggest that communication is favoured when food sources are hard to find, of variable quality and long lasting. These results mirror the findings of simulations of honeybee communication. Scouts locate food sources faster than tandem followers in some environments, suggesting that tandem running may fulfil the criteria of teaching only in some situations. Furthermore, tandem running was only beneficial above a critical colony size threshold. Taken together, our model suggests that there is a considerable parameter range that favours colonies that do not use communication, which could explain why many social insects with small colony sizes forage solitarily.

Introduction

Finding food is critical for survival and reproduction, but also energy- and time-consuming. Foraging for food can be done independently or by using information provided by other organisms (Sumpter 2010; Hoppitt and Laland 2013). In social insects, such as ants, social bees or social wasps, new food sources are usually discovered by scouts that explore the environment

on their own (Hölldobler and Wilson 1990; Seeley 1995). After finding a profitable food source,

they return to their nest and often communicate their discovery to nestmates. The communicated

information depends on the species, but often includes the location of the resource, e.g. by

means of laying a pheromone trail (Hölldobler and Wilson 1990; Jarau and Hrncir 2009;

Czaczkes et al. 2015b). Recruitment communication allows colonies to exploit profitable

feeding sites fast, e.g. before competitors have discovered and consumed the food source. Once

the foragers have learned the location of the food source, they can use their route memory to

return to the feeding site (e.g. von Frisch 1967; Collett et al. 2013).

In social insects, foraging strategies should not only take into account short-term

individual success, but also how they affect colony foraging success. Thus, the value of

communication should ultimately be studied at the colony level. So far, most theoretical and

empirical studies that explored the value of communication for colony foraging success have

focused on honeybees (but see also e.g. Sumpter and Pratt 2003; Dechaume-Moncharmont et

al. 2005; Czaczkes et al. 2015a). These studies suggest that the value of communicating the

location of food sources by means of waggle dances depends on how food sources are

distributed (Sherman and Visscher 2002; Dornhaus and Chittka 2004; Dornhaus et al. 2006;

Beekman and Lew 2008; Donaldson-Matasci and Dornhaus 2012; Schürch and Grüter 2014;

I'Anson Price et al. 2019; reviewed in I'Anson Price and Grüter 2015). For example, Beekman

& Lew (2008) found that the value of the "dance language" (the spatial information provided

by the waggle dance) depends on the size and distance of the food patches. When patches were

large and close to the hive, colonies that did not use dance communication and instead followed

an individual foraging strategy were more successful. Dornhaus et al. (2006) concluded that

dance communication does not help colonies collect more energy if there are many food sources

that vary little in quality. Their models suggest that communication is beneficial if high-quality

food sources are available, but are hard to find and that dance communication could be

detrimental if food sources are easy to find (see also Dechaume-Moncharmont et al. 2005). In

the latter case, foragers should search for new food sources through scouting (independent

search) and return to known high-quality food sources using route memory (Schürch and Grüter

2014).

There is a well-known, but not yet fully understood link between colony size and the

method of recruitment in ants (Beckers et al. 1989; Planqué et al. 2010; Dornhaus et al. 2012).

While large colony size is associated with pheromone-based mass-recruitment, species with

smaller colony sizes often forage solitarily or they use a recruitment method called tandem

running (Beckers et al. 1989). In tandem running, an experienced ant (tandem leader) guides an

inexperienced nestmate (tandem follower) to a new nest-site or a rewarding food source

(Hingston 1929; Wilson 1959; Möglich et al. 1974; Franks and Richardson 2006; Pratt 2008;

Kaur et al. 2017; reviewed in Franklin 2014). It has been argued that tandem followers locate

resources quicker than scouts that search for resources by individual exploration and trial-and-

error learning (Franks and Richardson 2006). Additionally, ants that are recruited by a tandem

leader might find food sources of higher quality because foragers are more likely to perform

tandem runs after finding a better food source (Shaffer et al. 2013). On the other hand, tandem

running also has disadvantages. During a tandem run, both ants walk with reduced speed

(Franks and Richardson 2006; Kaur et al. 2017) and a substantial proportion of tandem runs fail

(e.g. Wilson 1959; Pratt 2008; Glaser and Grüter 2018; Grüter et al. 2018). Furthermore,

recruits experience time and opportunity costs as they wait inside their nest for a leader, rather

than search in the environment for food sources by themselves. These disadvantages could

explain why some ant species do not seem to use tandem communication when foraging, even

though tandem runs are used during colony migrations (Hölldobler 1984; Traniello and

Hölldobler 1984; Fresneau 1985; Maschwitz et al. 1986). More generally, a sizeable group of

ant species do not seem to use any form of communication during foraging (e.g. Beckers et al.

1989; Lanan 2014; Reeves and Moreau 2019). This raises the question whether, when and how

a communication method that is relatively slow and small-scale, like tandem running, improves

colony foraging success and whether the ecological circumstances that favour tandem running

match those that favour honeybee dance communication.

We developed an agent-based simulation model to investigate the importance of

recruitment communication in the form of tandem running for the foraging success of virtual

ant colonies. We compared colonies that could perform tandem runs with colonies that

consisted only of scouts, *i.e.* without tandem running in an environment that varied in the

number, quality, distance and longevity of food sources. Additionally, we tested whether colony

size affects the importance of communication for colony foraging success. Finally, we explored

the role of forager ratio (relative numbers of scouts and recruits) and tested if recruits indeed

locate food sources faster than scouts. Based on studies that simulated honeybee foraging, we

predicted that tandem running is beneficial when high quality food sources are hard to find

(Dornhaus et al. 2006; Beekman and Lew 2008), but is detrimental to colony success when food

sources are short-lived (Schürch and Grüter 2014). We also predicted that larger colonies

benefit more from tandem running.

The agent-based simulation model

An agent-based simulation model (ABM) was developed using the software Netlogo 6.1.1

(Wilensky 1999, Wilensky and Rand 2015) (the NetLogo file can be found in the online

material). The model simulates the foragers of an artificial ant colony in an environment

consisting of their nest and food sources. Some of the basic parameters, like the range of colony

sizes, walking speeds or energy collected by foragers were derived from the ant species

Temnothorax nylanderi (Glaser and Grüter 2018).

 Purpose

 The aim of our model was to explore the adaptive value of tandem running in ants by measuring

 the colony foraging success (as gained energy) and the time required by foragers to find a food

 source. We compared colonies that could perform tandem runs with colonies that consisted only

 of scouts, *i.e.* without tandem running. This latter situation is found in many ant species with

 small colony sizes, such as *Diacamma* or *Neoponera* (Hölldobler 1984, Traniello and

 Hölldobler 1984, Fresneau 1985, R. Kaur, pers. communication). In both situations, foragers

 could also use route memory (or private information) to return to food sources they visited in

 the past. We assessed the effects of communication depending on food source distribution

 (number, distance), their quality and stability as well as colony size and the scout-recruit ratio.

 Entities, state variables and scale

 Netlogo operates with patches that can be used to measure distances and ticks for time steps. In

 our model, 1 tick is equivalent to 1 second and 1 patch to 1 cm. The agents operate in a two-

 dimensional square grid of 140×140 patches (arena) with a nest and either 2 or 10 food sources

 (FS). This simulated environments with few or many food sources. The nest is located in the

 center (x=0, y=0), with a radius of 10 patches. The food patches were at a distance of either 40

 (default) or 20 patches from the outer edge of the nest, simulating natural conditions as *T.*

 nylanderi mostly forages within 50 cm from their nest (Heinze et al. 1996). Each food source

 had a size of 1 patch, which could represent a dead insect or a drop of honey dew, and could

 either be of high or low quality (FS_{high} and FS_{low}), simulating a sugar solution of either 1 molar

 or 0.1 molar concentration.

 Since all agents are foragers, our default colony size of 100 would correspond to a

 natural colony consisting of ~300-400 workers, assuming that foragers account for about 20-

 30% of a *Temnothorax* colony (e.g. Shaffer et al. 2013). Simulated colonies consisted of varying

ratios of scouts that search for resources independently and recruits that waited in the nest until

they are recruited to a food source. The default scout-recruit ratio was 1:4 (*i.e.* 20 scouts + 80

recruits in the default situation), similar to what has been observed in honeybees where scouts

represent about 5-35 % of the colony (von Frisch 1967; Seeley 1995). In colonies without

tandem running, all foragers were scouts. In the default configuration, scouts and recruits can

both assume any of the following seven states: (*1*) idle inside the nest, (*2*) feeding at food

sources, (*3*) returning to the nest with food, (*4*) unloading food, (*5*) searching for a follower

inside the nest, (*6*) leading a tandem run to the food source or (*7*) returning alone to the food

source (*i.e.* use private information). Additionally, scouts search for food sources independently,

while recruits wait inside the nest for a tandem leader. Recruits can then follow tandem runs to

a food source (Fig. 1).

Fig. 1. State diagram for the agent-based model for colonies with tandem runs. All foragers

start inside the nest. Green boxes represent agent states that are possible for both scouts and

recruits, blue boxes are states that are only possible for scouts and orange boxes represent
states that are only possible for recruits.

Colonies gain nest energy (NE) when agents return to the nest and "unload" the energy gained at food sources. We estimated the energy content of a crop load of a *T. nylanderi* forager the following way: we measured foragers (N = 21, from 3 colonies) and estimated that full foragers carry ~0.15 mg of sugar solution. Given the energetic value of sucrose, we calculated that a forager feeding at a 1 molar sucrose solution collects ~0.75 Joule per foraging trip, whereas a forager feeding at a 0.1 molar solution would obtain ~0.075 J.

Table 1: Overview of the model parameters and the used values

Variables	Description	Default values	Other values tested
FS distance	Distance from nest centre to food source	40	20
FS number	Number of food sources	2 or 10	
Colony size	Number of agents (foragers) in a colony	100	20-200
Scout-recruit ratio	The ratio of scouts and recruits in a colony	1:4 (r = 0.2) or (all scouts)	1:9 to 10:0 (r = 0.1 to 1.0)
FS_{High}	Energy gained from the high-quality food source	0.75 J	
FS_{Low}	Energy gained from the low-quality food source	0.075 J	
t_{max}	Duration of a simulation (1 tick ~ 1 second)	5400 ticks	21600 ticks
$v_{outside}$	Walking velocity of ants outside the nest	0.8 patch/ticks	
v_{nest}	Walking velocity of ants inside the nest	0.1 patch/ticks	
v_{tandem}	Walking velocity of Tandem leader and Tandem follower towards the respective food source	0.4 patch/ticks	
M_{cost}	Metabolic or energy cost of walking outside	$2.446 \times e^{-7}$ J/tick	$2.446 \times e^{-6}$, $2.446 \times e^{-8}$
t_{scouts}	Time a scout searches food before returning to nest	600-900 ticks	
$t_{nest-stay}$	Time a returned forager stays inside the nest	60 ticks	
$t_{tandemstarter}$	Time an active recruiter searches for a recruit inside the nest	120 ticks	
$t_{feeding}$	Feeding time of drinking agents	120 ticks	
$p_{break-up}$	Probability that tandems break up	0/tick	0.002/tick,

			0.005/tick
$p_{recruitment}$	Probability to recruit when "satisfied"	50%	

Process overview

The default simulation duration t_{max} was 5400 ticks (corresponding to 90 minutes), but we also

tested a duration of 21600 ticks (corresponding to 6 h). Time and distance in the model were

connected via the walking speeds (0.8 patches/tick outside the nest = $v_{outside}$, 0.4 patches/tick in

a tandem run= v_{tandem}), which were chosen to be similar to the walking velocity (in cm/sec) of

T. nylanderi ants (Glaser and Grüter 2018).

When the model was initialized (t=0), the nest and either 2 or 10 food sources and the

agents were created. In the situation without tandem running, only scouts were simulated. All

agents started in the centre of the nest. Scouts immediately started to perform a random walk to

search for food sources with the speed of $v_{outside}$, whereas recruits patrolled inside the nest with

speed v_{nest} (0.1 patches/tick) and waited to be recruited by another agent. All agents started with

an energy of zero. When leaving the nest, this energy decreases every tick by a metabolic cost

M_{cost} (see Table 1). M_{cost} was chosen so that the metabolic costs that accumulate during an

average foraging trip correspond to ~0.1% of the value of energy obtained during a typical

foraging trip (Fewell 1988). We estimated this by running several simulations and measuring

foraging trip duration of our agents. We also ran simulations with metabolic rates that were 10-

times higher or 10-times lower than our default value but found that this did not affect the

general patterns (Fig. S1).

When an agent finds a food source, it becomes a feeding agent and feeds for a duration

of 120 ticks. It gains either 0.75 J or 0.075 J, depending on whether the food source is of high

or low quality. If scouts do not find a food source within a certain time period (t_{scouts}), they

return to the nest. If they are at greater distances from the nest, unsuccessful scouts return

quicker (600 ticks). Unsuccessful scouts that are closer to the nest (35 patches from the center)

return if 900 ticks have passed. This was done to match observations that *T. nylanderi* scouts

often return to their nest if they had been searching unsuccessfully for several minutes (S.M.G.,

personal observation). After their return, unsuccessful scouts wait idle inside the nest for 60

ticks ($t_{nest-stay}$), before resuming to scout. At the end of the feeding time, agents return to the nest

either as "satisfied" or "unsatisfied" foragers. Foragers that found a high-quality food source

were always satisfied, whereas agents feeding at a low-quality food source had only a 10%

probability to become satisfied. After unloading for the duration of $t_{nest-stay}$, "satisfied" agents

become prospective tandem leaders with a 50% probability ($p_{recruitment}$), whereas unsatisfied

agents would not recruit. This leads to a recruitment probability of 5-50% per trip, which is

similar to what has been found in both *T. nylanderi* and *Pachycondyla harpax* (Glaser and

Grüter 2018; Grüter et al. 2018). Satisfied agents return to the same food source they had visited

before, either in a tandem run or alone. In other words, they use "route memory" to revisit a

high-quality food source, but were unlikely to return to a low-quality food source (10%

probability). Unsatisfied agents would not recruit and either wait inside the nest for a tandem

leader (recruits) or they search for a new food source (scouts). Fig. S2 is a screenshot of a

simulation showing the arrangement of the nest, food sources and some of the agent states.

Prospective tandem leaders stay inside the nest and search for a potential recruit for the

duration of 120 ticks ($t_{tandemstarter}$). A tandem run starts when a leader encounters a recruit on the

same patch. By default, tandem runs do not break up but we also tested situations with a break-

up probability of 0.002 and 0.005 per tick, which corresponds to tandem success rates of ~75 %

and ~50% for the default distance (calculated based on an average tandem run duration of 127

ticks for the default food source distance). Lost tandem followers first perform a random walk

for 180 ticks ($t_{search-time}$) and – if they do not find a food source – have an equal probability to

become either a scout or to return to the nest as an unsatisfied forager.

218   In the default settings, food sources were *ad libitum*, *i.e* they did not disappear during

219 the simulations. Since this may not always be the case, we also simulated food sources that

220 disappeared after they were visited by 10 agents to create a more dynamic foraging environment.

221 If a food source disappears before ants return to it (either alone or in a tandem), agents reaching

222 the old food source location search randomly for 180 ticks ($t_{search-time}$), then they become

223 unsatisfied foragers and return to the nest. If the food source vanishes during feeding, the agent

224 becomes an unsatisfied forager. Food sources that have disappeared are replaced by an identical

225 food source at the same position after 600 ticks have passed, which means that it has to be

226 discovered again by scouts. For each simulation run, new inexperienced agents were created as

227 described above.

229 *Tested factors*

230 • Spatio-temporal distribution and variability of food sources: we tested the effects of food

231  source number (2 or 10), variability (only high-quality or variable quality. In the latter

232  situation, food sources alternated in quality, *i.e.* FS 1 high-quality, FS2 low-quality, FS3,

233  high-quality etc.), distance, foraging (simulation) duration and food source longevity (stable

234  or short-lived).

235 • Scout-recruit ratio: next to our default ratio of 1:4, we tested several other ratios, including

236  the extreme case with only scouts.

237 • Colony size: in addition to simulating a colony size of 100 agents, we tested a range of other

238  colony sizes (Table 1).

239 • Food discovery time of scouts and recruits: For each simulation run, we quantified the time

240  scouts needed to discover their first food source. In recruits, we measured both their waiting

241  time inside the nest and the duration of the tandem run. These durations were averaged per

242  forager type and per simulation run. Agents that did not discover a food source during an

entire simulation were given the maximum value of 5400 ticks. Only conditions with

successful tandem runs were considered.

The total nest energy NE (total J of all individual collection trips minus the total J of the

metabolic costs) were measured for each simulation run.

Sensitivity of outcomes

Due to the stochasticity of simulations we performed 30 simulation runs for each tested

combination of parameters. To assess how sensitive our model is to changes in the default

parameters, we varied many of them and explored their effect, as mentioned above (see Table

1). We also simulated environments with only low-quality food sources. A pure scouting

strategy was always better under these circumstances (Fig. S3). This is because tandem runs

are very rare when all food sources are of low quality and recruits spend most of their time

inside the nest.

Statistical analyses

All statistical analyses were performed using the software R 3.6.3 (www.r-project.org). Since

different treatments occasionally had unequal variance (heteroscedasticity) or contained zeros

and in order to provide a consistent statistical approach we used non-parametric statistical tests

throughout. It should be noted, however, that when we compared parametric and non-

parametric methods (Anova's), they yielded very similar results. We used Mann-Whitney U

tests to compare two independent samples and Wilcoxon signed-rank tests for paired data. In

addition to the p-values, the R software provides the test statistic value W, which is a linear

transformation of the usual rank sum statistic U. When three groups were compared, we used

Kruskal-Wallis tests and Dunn tests with sequential Bonferroni corrections for post-hoc pair-

wise comparisons ("FSA" package, Ogle et al. 2020) (Sokal and Rohlf 1995).

Results

Distribution of food sources

We first tested if the number of food sources and their distance from the nest affect the value of communication. When colonies had access to few food sources, they were more successful with tandem recruitment (scout-recruit ratio of 1:4) than colonies consisting only of scouts, irrespective of whether food sources were of high-quality (Fig. 2) (Mann-Whitney U Test, W = 215, p = 0.0004) or of variable quality (W = 307, p = 0.034). In a rich environment, with 10 food sources, colonies collected overall more energy (Fig. 2). Tandem communication was beneficial when food source quality was variable (W = 112, p < 0.0001), whereas colonies consisting only of scouts performed better when all 10 food sources were of high quality (W = 773, p < 0.0001). This general pattern did not change when food sources were closer to the nest (20 patches instead of 40 patches) (2 food sources, high-quality: p < 0.0001; variable-quality: W = 210, p = 0.0003; 10 food sources, high-quality: W = 827, p < 0.0001; variable-quality: W = 54, p < 0.0001), but colonies gained overall more energy when all food sources were close to the nest (Fig. 2). Fig. 2e and 2f illustrate the temporal development of nest energy during exemplary simulation runs that correspond to the conditions shown in Fig. 2a and 2b with high-quality food sources.

Fig. 2. Nest energy with 2 or 10 food sources. In (a) and (b) food sources were at a distance of 40 patches, whereas in (c) and (d) food sources were at a distance of 20 patches. *p<0.05, **p<0.001, ***p<0.0001. In (e) and (f), nest energy is plotted over time for conditions as shown in (a) and (b) when all food sources were of high quality (5 simulation runs per treatment for visualisation of the trajectory).

Foraging duration and food source longevity

When we increased the foraging duration (*i.e.* the simulation duration) from 5400 to 21600 ticks, we found a similar pattern. Tandem running was highly beneficial when there were few food sources (high-quality: W = 0, p < 0.0001; variable-quality: W = 0, p < 0.0001). Tandem runs were also beneficial when there were many food sources of variable quality (W = 56, p < 0.0001). In the case of many high-quality food sources, pure scout colonies performed better

(W = 900, p < 0.0001). It is noteworthy that colonies with tandem communication were almost

as successful in an environment with 2 food sources as in an environment with 10 food sources

(Fig. 3a,b).

Fig. 3. Nest energy with 2 or 10 food sources. In (a) and (b), food sources simulations were 4-times
longer (21600 ticks instead of 5400). In (c) and (d), simulations lasted 5400 ticks and food sources
disappeared if they were visited by 10 ants. A new food source appeared after a delay.

So far, we assumed that food sources offered food during the entire simulation. Next,

we tested the effects of short-lived food sources. If food sources were unstable (sometimes

called non-renewable), a scouting strategy was more successful, irrespective of the number of

food sources and their variability (Fig. 3c,d) (2 food sources, high-quality: W = 897, p-value <

0.0001; variable-quality: W = 896, p-value < 0.0001; 10 food sources, high-quality: W = 900,

p-value < 0.0001; variable-quality: W = 900, p-value < 0.0001). Differences were particularly

pronounced when colonies were offered many food sources. Scouting remained the better

strategy when we increased the foraging duration to 21600 ticks (e.g. 2 food sources, high-

quality: W = 900, p-value < 0.0001; variable-quality: W = 900, p-value < 0.0001).

Tandem success rate

Tandems do occasionally break up and we tested how this affects the energy collected by

colonies. We compared colonies with 100% (default), ~75% and ~50% successful tandem runs

and colonies with only scouts in an environment with few food sources, *i.e.* under conditions

where tandem runs are beneficial (Fig. 2a). Our simulations show that a reduction in tandem

success rate has a negative impact on the energy intake that is collected by colonies (Fig. 4). If

only about 50% of the tandem runs are successful, colonies without any tandem running collect

more energy in an environment with few, stable food sources (Fig. 4) (high-quality, 50%

success rate vs. no tandems: $W = 246$, $p = 0.002$, variable-quality, 50% success rate vs. no

tandems: $W = 257$, $p = 0.004$).

Fig. 4. Nest energy with 2 food sources of high (a) and mixed (b) quality in relation to the tandem success rate. Adjacent treatment groups were compared, as indicated by asterisks or "n.s.". No tandems = only scouts. Default settings were used for the other parameters.

However, tandem runs with a high rate of failure (50%) are not always a disadvantage

compared to having no communication. When colonies can forage for longer (simulations of

21600 ticks), colonies that perform tandem runs with a ~50% break-up rate are more successful

than colonies consisting of only scouts (Fig. 5) (high-quality, 50% success rate vs. no tandems:

$W = 866$ $p < 0.0001$, variable-quality, 50% success rate vs. no tandems: $W = 689$, $p = 0.0003$),

highlighting the benefits of imperfect communication over longer time periods.

Fig. 5. Nest energy with 2 food sources of high or variable quality and a long foraging duration. Colonies
were scouting or could recruit with tandem runs that had a ~50% failure rate. Default settings were used
for the other parameters.

Colony size and scout-recruit ratio

We tested various colony sizes ranging from 20 to 200 agents in an environment with few,
variable food sources, *i.e.* an environment that favours tandem running under default conditions
(see Fig. 2a).

Fig. 6. The relationship between colony size and nest energy (a) and nest energy per agent (b) in
colonies with and without tandem runs in environments with two food sources of variable quality. Three

354 scout-recruit ratios were simulated, r = 0.2 and r = 0.4 and colonies consisting only of scouts, r = 1.0.
355 Grey area indicates confidence intervals. Significance tests refer to comparisons among ratios,
356 separately for each colony size. P-values for total nest energy (a) or energy per agent (b) are identical.
357 Default settings were used for the other parameters.
358

Colony size had a strong effect on the total collected energy that was collected (Fig. 6).

If colonies were very small (20 foragers), they were least successful if they performed tandem

runs and had a default scout-recruit ratio of 0.2 (Table 2). There was no difference in foraging

success when colony size ranged from 30 to 50 foragers. However, colonies with tandem

recruitment were more successful if they had at least 60 agents (Table 2). The most successful

colonies contained 40% scouts, suggesting that the scout-recruit ratio has a considerable impact

on colony success. Fig. 6b shows the nest energy collected per agent (nest energy/colony size).

In colonies with only scouts, individual agents collected a relatively constant amount of energy

irrespective of colony size (Spearman rank correlation: rho = 0.1, p = 0.09). In colonies with

tandem running, on the other hand, individual agents collected more energy on average as

colony size increased from 20 to 100 agents (r = 0.2, rho = 0.34, p < 0.0001; r = 0.4, rho = 0.35,

p < 0.0001).

Table 2: Effect of colony size on nest energy. Three conditions were tested: in two conditions, colonies
performed tandem runs and had a scout-recruit ratio of 0.2 or 0.4. In the third condition, colonies
consisted only of scouts (1.0). Pair-wise comparisons were performed if the overall p < 0.05 and p-
values were corrected using sequential Bonferroni.
376

Colony size	Kruskal-Wallis Test		p-value of pair-wise comparisons		
	χ^2	p-value	0.2 vs. 0.4	0.2 vs. 1.0	0.4 vs. 1.0
20	12.7	0.002	0.018	0.002	0.42
30	2.33	0.31	NA	NA	NA
40	3.92	0.14	NA	NA	NA
50	1.81	0.41	NA	NA	NA
60	8.98	0.01	0.077	0.4	0.01
70	22.85	<0.0001	0.0007	0.35	<0.0001
80	20.6	<0.0001	0.13	0.007	<0.0001
90	15.86	0.0004	0.09	0.048	0.0002
100	19.17	<0.0001	0.053	0.027	<0.0001
200	46.59	<0.0001	0.006	0.0001	<0.0001

To explore this further, we simulated different scout-recruit ratios and different colony

sizes to test how the balance between scouts and recruits affects colony foraging success.

Simulations suggest that the optimal proportion of scouts is ~40% for the simulated

environment, irrespective of colony size (Fig. 7). Interestingly, deviations from the optimal

ratio have a larger negative impact in larger colonies (see "pointiness" of curves in Fig. 7). For

example, there is no difference in success when colonies with 50 agents contain 40% or 80%

of scouts (W = 119; p = 0.54). When colony size is 200, however, colonies with 80% scouts

collect 31.5% less energy than colonies with 40% scouts (W = 199, p = 0.0001).

Fig. 7. The effect of the scout-recruit ratio with three different colony sizes. The line shows the best fit
line based on local polynomial regression using the LOESS method (locally estimated scatterplot
smoothing). The smallest ratio was 0.1. A ratio of 1.0 refers to colonies containing only scouts.
391
392

Discovery times

Unsurprisingly, foragers needed more time to find their first food source in an environment

with few food sources compared to when there were many food sources (Fig. 8). Recruits

needed less time in an environment with few, high-quality food sources compared to scouts

(Wilcoxon-signed rank test: W = 143, p-value < 0.0001), whereas there was no difference when

food sources were variable in quality (Fig. 8a) (W = 348, p-value = 0.13). However, in an

environment with many food sources, scouts did comparatively better and needed a similar

amount of time to locate their first food source when food sources were all high-quality ($W = 327$, p-value = 0.07). With many, variable food sources, scouts were significantly faster than recruits (Fig. 8b) ($W = 720$, p-value < 0.0001).

Fig. 8. Time until agents located their first food source in environments with few (a) or many (b) food sources of constant or variable quality. For recruits, the food discovery time consisted of the time waiting inside the nest and the tandem run duration. Default settings were used for the other parameters.

Discussion

Our simulations show that the spatio-temporal distribution of food sources greatly affects whether colonies with communication are more successful than colonies that employ a scouting strategy. Tandem running was beneficial when colonies were in an environment with few food sources (+ 57-83% nest energy) and when food sources were of variable quality (Fig. 2a,b). Colonies without communication were more successful (~15%) in a rich environment that offered only high-quality food sources. This is in line with studies that simulated honeybee foraging and found that communicating food source locations by waggle dancing is most beneficial if food sources are hard to find and of variable quality (Dornhaus et al. 2006; Beekman and Lew 2008; Schürch and Grüter 2014; I'Anson Price et al. 2019). Under such

circumstances, the probability that scouts find high-quality food sources on their own is low

and communicating the location of a relatively small number of high-quality patches becomes

advantageous. As food source variability decreases and the number of high-quality food sources

increases, scouts become more successful. Even though colonies with communication also

collect more energy in such an environment, the benefits of communication no longer offset the

costs of recruits waiting for information inside the nest. This highlights that communication

often has considerable time and opportunity costs (Seeley 1983; Seeley and Visscher 1988;

Dechaume-Moncharmont et al. 2005; Schürch and Grüter 2014; I'Anson Price et al. 2019).

It has been hypothesised that recruitment communication is particularly beneficial in an

ephemeral environment (Sherman and Visscher 2002; Dornhaus and Chittka 2004; Grüter and

Ratnieks 2011), *i.e.* when food sources last only for short time-periods and, thus, need to be

exploited quickly. Counterintuitively, a simulation model of honeybee foraging has found that

communication was less beneficial if food sources were shorter-lived (Schürch and Grüter

2014). Our simulations support their findings by showing that tandem running was a very

successful strategy in a stable environment with relatively long foraging durations (*i.e.* with

longer simulations) and few, variable food sources (Fig. 3a). A long-lasting food source could

be a large insect (Lanan 2014), floral nectars or a group of honeydew secreting insects (Carroll

and Janzen 1973; Quinet and Pasteels 1996; Völkl et al. 1999; Mailleux et al. 2003; Lanan

2014). A very different pattern was observed when resources were shorter-lived: colonies

without communication were always more successful, irrespective of the foraging (simulation)

duration (Fig. 3c,d). The most likely explanation is that colonies with communication pay time

costs without being able to take advantage of the benefits of communication over longer time

periods (see also Schürch and Grüter 2014). Our model differs from theirs in that our food

sources only disappeared if they were exploited, rather than with a constant probability. A food

source that disappears after it has been exploited could be a droplet of honeydew that fell on

vegetation. Honeydew droplets on leaf surfaces represent an important food source for the

tandem recruiting *Temnothorax curvispinosus* (Lynch et al. 1988).

Tandem runs occasionally break-up and success rates of ~50% to 90% are not

uncommon (Wilson 1959; Pratt 2008; Kaur et al. 2017; Glaser and Grüter 2018; Grüter et al.

2018). We simulated different success rates and found that colonies with more successful

tandem runs collected more energy (Fig 4). If the success rate was about 50%, colonies

consisting only of scouts collected more energy in an environment with few food sources, *i.e.*

a virtual environment that normally favours tandem running. When foraging durations were

longer, on the other hand, colonies with tandem runs gained the upper hand over scouting

colonies even though half of all tandem runs failed (Fig. 5). Under these circumstances, even a

relatively low number of successful recruitment events can be very important because the

discovered high-quality food sources can be exploited for longer time periods by successful

recruits. Additionally, tandem recruitment can lead to an exponential increase of ants at a feeder

even if a leader recruits <1 follower per trip. With exponential growth, the impact of

communication will increase over time (Fig. 2e).

We found that colony size had a considerable effect on the value of tandem

communication (Fig. 6). This contrasts with models of honeybee communication, where colony

size did not greatly affect the benefits of communication (Dornhaus et al. 2006; Schürch and

Grüter 2014), but is consistent with an empirical study on honeybee colony foraging success

(Donaldson-Matasci et al. 2013) and a mathematical model of ant communication (Planqué et

al. 2010). If colonies contained 60 or more foragers, tandem communication was usually

beneficial. However, a pure scouting strategy was equally or more successful when colonies

had 20 to 50 foragers, even in environments with few and variable food sources, *i.e.* a virtual

environment that normally favours tandem running. This number of foragers could be expected

in ant colonies with ~80-250 workers (assuming that foragers make up 20-30% of the worker

population, e.g. Shaffer et al. 2013), which is also the typical colony size of many ant species

that use tandem running and species with solitary foraging (Beckers et al. 1989). Our simulation

results could explain why some species, e.g. in the genera *Diacamma* or *Neoponera*, do not

perform tandem runs during foraging even though they use this recruitment method during

migrations (Hölldobler 1984; Traniello and Hölldobler 1984; Maschwitz et a. 1986). Whether

colonies employ tandem running might depend on the food sources they collect (e.g. small or

large items) and whether they are risk-averse or risk-prone because tandem recruitment was

often associated with a more unpredictable outcome in our simulations (greater variation in nest

energy gain among simulations of a particular situation, see Fig. 2). A better understanding of

the natural history of these species and similar species that do perform tandem runs (e.g.

Neoponera vs. *Pachycondyla*) is needed to understand why some species use communication,

while others forage solitarily.

Colony foraging performance depended on the proportions of scouts and recruits (Fig.

7). In our simulations with few food sources, colonies were most successful if scouts

represented about 40% of the forager population, but this is likely to depend on the number and

variability of food sources (see Fig. 2). Interestingly, having the right scout-recruit ratio is more

important in larger colonies than in smaller ones, possibly because the foraging success of

smaller colonies depends more on chance events, such as the discovery of a high-quality food

source by a single scout. This suggests that larger colonies would benefit from having the ability

to assess their current environment and adjust their use of communication accordingly. Whether

this is indeed common is not well known, but it has recently been reported that honeybees are

able to assess the value of communication and reduce their reliance on waggle dances if dance

information is not beneficial in the current environment (I'Anson Price et al. 2019).

In the simulations, we measured the time recruits and scouts need to locate their first

food source in environments with many or few food sources. We found that the food discovery

time depends strongly on the environment. Recruits were faster in environments with few high-quality food sources, whereas scouts found a food source sooner in an environment with many, variable food sources. Our measurements also included the time that recruits wait inside the nest to find a tandem leader. Franks and Richardson (2006) found that tandem followers found a food source faster in their experiment with one food source, which, in combination with their other findings, indicated that tandem running fulfils the criteria for animal teaching set out by Caro and Hauser (1992; namely, a teacher [*i*] modifies its behaviour in the presence of a naive observer, [*ii*] at some cost to the leader [*iii*] so that the observer can learn more quickly or efficiently). Our simulations suggest that this is the case only in certain environments, namely those with few, high-quality resources. In other situations, scouts are likely to learn food source locations quicker and tandem running might no longer fulfil the criteria for animal teaching (namely that a follower acquires knowledge or learns a skill more rapidly or efficiently than it might otherwise do, or that it would not learn at all, see Caro and Hauser 1992).

Taken together, our simulations show that the value of tandem communication is highly dependent on the environment and the size and composition of the colony. Future studies should explore whether and how foragers can assess their foraging environment and modify their communication behaviour (see also Grüter and Czaczkes 2019). It would also be desirable to test the conclusions from our simulations empirically, but so far it has been challenging to stop ants from performing tandem runs without affecting their behaviour.

Acknowledgments

We would like to thank the AG Foitzik for discussion and feedback on the model. S.M.G. was funded by the German Research Foundation (DFG: GR 4986/1-1).

References

Al Toufailia HM, Grüter C, Ratnieks FLW (2013) Persistence to unrewarding feeding locations by honeybee foragers (*Apis mellifera*): the effects of experience, resource profitability and season. Ethology 119:1096–1106

Beckers R, Goss S, Deneubourg J-L, Pasteels JM (1989) Colony size, communication and ant foraging strategy. Psyche 96:239–256

Beekman M, Lew JB (2008) Foraging in honeybees - when does it pay to dance? Behavioral Ecology 19:255–262

Caro TM, Hauser MD (1992) Is there teaching in nonhuman animals? Quarterly Review of Biology 67:151–174

Carroll CR, Janzen DH (1973) Ecology of foraging by ants. Annual Review of Ecology and Systematics 4:231–257

Collett M, Chittka L, Collett TS (2013) Spatial memory in insect navigation. Current Biology 23:R789–R800

Czaczkes TJ, Czaczkes B, Iglhaut C, Heinze J (2015a) Composite collective decision-making. Proceedings of the Royal Society of London B: Biological Sciences 282:20142723

Czaczkes TJ, Grüter C, Ratnieks FLW (2015b) Trail pheromones: an integrative view of their role in colony organization. Annual Review of Entomology 60:581–599

Dechaume-Moncharmont FX, Dornhaus A, Houston AI, McNamara JM, Collins EJ, Franks NR (2005) The hidden cost of information in collective foraging. Proceedings of the Royal Society of London Series B-Biological Sciences 272:1689–1695

Donaldson-Matasci MC, Dornhaus A (2012) How habitat affects the benefits of communication in collectively foraging honey bees. Behavioral Ecology and Sociobiology 66:583–592

Donaldson-Matasci MC, DeGrandi-Hoffman G, Dornhaus A (2013) Bigger is better: honeybee colonies as distributed information-gathering systems. Animal Behaviour 85:585–592

Dornhaus A, Chittka L (2004) Why do honey bees dance? Behavioral Ecology and Sociobiology 55:395–401

Dornhaus A, Klügl F, Oechslein C, Puppe F, Chittka L (2006) Benefits of recruitment in honey bees: effects of ecology and colony size in an individual-based model. Behavioral Ecology 17:336–344

Dornhaus A, Powell S, Bengston S (2012) Group size and its effects on collective organization. Annual Review of Entomology 57:123–141

Franklin EL (2014) The journey of tandem running: the twists, turns and what we have learned. Insectes Sociaux 61:1–8

Franks NR, Richardson T (2006) Teaching in tandem-running ants. Nature 439:153

Fresneau D (1985) Individual foraging and path fidelity in a ponerine ant. Insectes Sociaux 32:109–116

Galef BG, Giraldeau LA (2001) Social influences on foraging in vertebrates: causal mechanisms and
adaptive functions. Animal Behaviour 61:3–15

Glaser S, Grüter C (2018) Ants (*Temnothorax nylanderi*) adjust tandem running when food source
distance exposes them to greater risks. Behavioral Ecology and Sociobiology 72:40

Grüter C, Czaczkes TJ (2019) Communication in social insects and how it is shaped by individual
experience. Animal Behaviour 151:207-215

Grüter C, Ratnieks FLW (2011) Honeybee foragers increase the use of waggle dance information
when private information becomes unrewarding. Animal Behaviour 81:949–954

Grüter C, Wüst M, Cipriano AP, Nascimento FS (2018) Tandem recruitment and foraging in the
ponerine ant *Pachycondyla harpax* (Fabricius). Neotropical Entomology 47:742-749

Heinze J, Foitzik S, Hippert A, Holldobler B (1996) Apparent Dear-enemy Phenomenon and
Environment-based Recognition Cues in the Ant *Leptothorax nylanderi*. Ethology 102:510–
522

Hingston RWG (1929) Instinct and intelligence. Macmillan, New York

Hölldobler B (1984) Communication during foraging and nest-relocation in the African stink ant,
Paltothyreus tarsatus Fabr. (Hymenoptera, Formicidae, Ponerinae). Zeitschrift fur
Tierpsychologie 65:40–52

Hölldobler B, Wilson EO (1990) The Ants. The Belknap Press of Harward University, Cambridge,
Massachusetts

Hoppitt W, Laland KN (2013) Social learning: An Introduction to Mechanisms, Methods, and Models.
Princeton University Press, Princeton

I'Anson Price R, Dulex N, Vial N, Vincent C, Grüter C (2019) Honeybees forage more successfully
without the "dance language" in challenging environments. Science Advances 5:eaat0450

I'Anson Price R, Grüter C (2015) Why, when and where did honey bee dance communication evolve?
Frontiers in Ecology and Evolution 3:1–7

Jarau S, Hrncir M (2009) Food Exploitation by Social Insects: Ecological, Behavioral, and Theoretical
Approaches. CRC Press, Taylor & Francis Group, Boca Raton, Florida

Kaur R, Joseph J, Anoop K, Sumana A (2017) Characterization of recruitment through tandem
running in an Indian queenless ant *Diacamma indicum*. Royal Society Open Science 4:160476

Lanan M (2014) Spatiotemporal resource distribution and foraging strategies of ants (Hymenoptera:
Formicidae). Myrmecological News 20:53

Mailleux AC, Deneubourg J-L, Detrain C (2003) Regulation of ants' foraging to resource productivity.
Proceedings of the Royal Society of London Series B-Biological Sciences 270:1609–1616

Maschwitz U, Jessen K, Knecht S (1986) Tandem Recruitment and Trail Laying in the Ponerine Ant
Diacamma rugosum: Signal Analysis. Ethology 71:30–41

Möglich M, Maschwitz U, Hölldobler B (1974) Tandem calling: a new kind of signal in ant
communication. Science 186:1046–1047

Ogle DH, Wheeler P, Dinno A (2020) FSA: Fisheries Stock Analysis

Planqué R, van den Berg JB, Franks NR (2010) Recruitment strategies and colony size in ants. PLoS
ONE 5:e11664

Pratt SC (2008) Efficiency and regulation of recruitment during colony emigration by the ant
Temnothorax curvispinosus. Behavioral Ecology and Sociobiology 62:1369-1376

Quinet Y, Pasteels JM (1996) Spatial specialization of the foragers and foraging strategy in *Lasius*
fuliginosus (Latreille) (Hymenoptera, Formicidae). Insectes Sociaux 43:333-346

Reeves DD, Moreau CS (2019) The evolution of foraging behavior in ants (Hymenoptera:
Formicidae). 14

Richardson TO, Sleeman PA, McNamara JM, Houston AI, Franks NR (2007) Teaching with
evaluation in ants. Current Biology 17:1520–1526

Rieucau G, Giraldeau LA (2011) Exploring the costs and benefits of social information use: an
appraisal of current experimental evidence. Philosophical Transactions of the Royal Society of
London Series B-Biological Sciences 366:949–957

Schürch R, Grüter C (2014) Dancing bees improve colony foraging success as long-term benefits
outweigh short-term costs. PLoS ONE 9:e104660

Seeley TD (1995) The wisdom of the hive: The social physiology of honey bee colonies. Harward
University Press, Cambridge, Massachusetts

Seeley TD (1983) Division of labor between scouts and recruits in honeybee foraging. Behavioral
Ecology and Sociobiology 12:253–259

Seeley TD, Visscher PK (1988) Assessing the benefits of cooperation in honeybee foraging: search
costs, forage quality, and competitive ability. Behavioral Ecology and Sociobiology 22:229–
237

Sherman G, Visscher PK (2002) Honeybee colonies achieve fitness through dancing. Nature 419:920–
922

Sokal RR, Rohlf FJ (1995) Biometry, Third edition. W. H. Freeman and company, New York

Sumpter DJT, Pratt SC (2003) A modelling framework for understanding social insect foraging.
Behavioral Ecology and Sociobiology 53:131–144

Sumpter DJT (2010) Collective Animal Behavior. Princeton University Press, Princeton

Traniello JF, Hölldobler B (1984) Chemical communication during tandem running in *Pachycondyla*
obscuricornis (Hymenoptera: Formicidae). Journal of Chemical Ecology 10:783–794

Völkl W, Woodring J, Fischer M, Lorenz MW, Hoffmann KH (1999) Ant-aphid mutualisms: the
impact of honeydew production and honeydew sugar composition on ant preferences.
Oecologia 118:483–491

von Frisch K (1967) The dance language and orientation of bees. Harvard University Press,
Cambridge, Massachusetts

Wilson EO (1959) Communication by tandem running in the ant genus *Cardiocondyla*. Psyche 66:29–
34

Fig. S1

Fig. S1. The effects of 10-times higher and 10-times lower metabolic costs on nest energy

(see Table 1). Two (a) and ten (b) high-quality food sources were offered, default values were

used for all other parameters. The default conditions match those shown in Fig. 2a and 2b.

Mann-Whitney U tests, **p<0.001, ***p<0.0001.

647 **Fig. S2**

649 **Fig. S2.** NetLogo interface showing some of the different agent types in different colours. In

650 this situation, two food sources (FS 1 and FS 2) were offered.

662 **Fig. S3**

664 **Fig. S3**. Nest energy of colonies with or without tandem runs when all food sources are of low

665 quality. Two food source distances were simulated, 40 patches (a) or 20 patches (b). Default

666 values were used for all other parameters. Mann-Whitney U tests, ***p<0.0001.