Single-component, self-assembling, protein nanoparticles presenting the receptor binding domain and stabilized spike as SARS-CoV-2 vaccine candidates

Linling He[1], Xiaohe Lin[1], Ying Wang[4,5], Ciril Abraham[4], Cindy Sou[1], Timothy Ngo[1], Yi Zhang[4,5], Ian A. Wilson[1,3], and Jiang Zhu[1,2*]

[1]Department of Integrative Structural and Computational Biology, [2]Department of Immunology and Microbiology, [3]Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla, California 92037, USA

[4]Fels Institute for Cancer Research and Molecular Biology, and [5]Department of Microbiology and Immunology, Temple University, Philadelphia, Pennsylvania 19140, USA.

[*] Corresponding author (to whom correspondence should be addressed)
JZ: Phone (858) 784-8157; Email: jiang@scripps.edu

KEYWORDS
Coronavirus disease 2019 (COVID-19); heptad repeat 2 (HR2); self-assembling protein nanoparticle (SApNP); severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2); spike (S) protein; vaccine.

ABSTRACT (150 words)

Vaccination against SARS-CoV-2 provides an effective tool to combat the COIVD-19 pandemic. Here, we combined antigen optimization and nanoparticle display to develop vaccine candidates for SARS-CoV-2. We first displayed the receptor-binding domain (RBD) on three self-assembling protein nanoparticle (SApNP) platforms using the SpyTag/SpyCatcher system. We then identified heptad repeat 2 (HR2) in S2 as the cause of spike metastability, designed an HR2-deleted glycine-capped spike (S2GΔHR2), and displayed S2GΔHR2 on SApNPs. An antibody column specific for the RBD enabled tag-free vaccine purification. In mice, the 24-meric RBD-ferritin SApNP elicited a more potent neutralizing antibody (NAb) response than the RBD alone and the spike with two stabilizing proline mutations in S2 (S2P). S2GΔHR2 elicited two-fold-higher NAb titers than S2P, while S2GΔHR2 SApNPs derived from multilayered E2p and I3-01v9 60-mers elicited up to 10-fold higher NAb titers. The S2GΔHR2-presenting I3-01v9 SApNP also induced critically needed T-cell immunity, thereby providing a promising vaccine candidate.

ONE-SENTENCE SUMMARY (125 characters)

The SARS-CoV-2 receptor binding domain and S2GΔHR2 spike elicited potent immune responses when displayed on protein nanoparticles as vaccine candidates.

INTRODUCTION

Three coronaviruses (CoVs) have caused widespread outbreaks in humans, including severe acute respiratory syndrome CoV-1 (SARS-CoV-1), Middle East respiratory syndrome CoV (MERS-CoV), and SARS-CoV-2, which is the causative agent of COVID-19 (*1-3*) and has resulted in more than 1.9 million deaths worldwide (*4*). Enormous efforts are being undertaken to develop effective therapeutics and prophylactics for SARS-CoV-2. Small molecules that can block the host receptor, angiotensin-converting enzyme 2 (ACE2), and the transmembrane protease serine 2 (TMPRSS2) (*5*), which is required to process the spike protein, are being considered as treatments in addition to other interventions (*6*). While the immunology underlying COVID-19 is still being intensively studied (*6-8*), various vaccine candidates are now in clinical development (*9-12*). Inactivated virus vaccines have exhibited robust neutralizing antibody (NAb) responses in animals (*13, 14*), whereas viral vector vaccines based on human adenovirus (Ad5 and Ad26) and chimpanzee ChAdOx1 have been evaluated in nonhuman primates (NHPs) and human trials (*15-18*). Both DNA (*19-21*) and mRNA (*22, 23*) vaccines have been rapidly developed, with moderate NAb titers observed for the mRNA vaccine in medium and high dose groups (*22*). A recombinant spike protein adjuvanted with lipid nanoparticles (NPs), NVX-CoV2373, elicited high NAb titers in a human trial that were on average four-fold greater than in convalescent patients (*24, 25*). Efficacy was recently reported for a vector vaccine (AZD1222: 70.4%) (*26*) and two mRNA vaccines (mRNA-1273: 94.1% and BNT162b2: 95%) (*27, 28*). In December 2020, the U.S. Food and Drug Administration (FDA) issued the emergency use authorization (EUA) for the two mRNA vaccines.

The SARS-CoV-2 spike protein is a trimer of S1-S2 heterodimers. The S1 subunit contains a receptor-binding domain (RBD) that binds to ACE2 on host cells to initiate infection.

The S2 subunit consists of a fusion peptide (FP) and heptad repeat regions 1 and 2 (HR1 and

HR2). Upon endocytosis of the virion, the S1 subunit is cleaved off to facilitate FP insertion into

the host cell membrane, while the remaining S2 refolds to bring HR1 and HR2 together to fuse

the viral and host cell membranes (*29*). The spike protein harbors all NAb epitopes and is the

main target for vaccine development against SARS-associated CoVs (*30*). Convalescent plasma

(CP) has been used to treat COVID-19 patients with severe conditions (*31*), highlighting the

importance of NAbs in protection (*32*). Due to moderate sequence conservation of the RBDs

(~73%), some previously identified NAbs targeting the SARS-CoV-1 RBD have been shown to

bind and cross-neutralize SARS-CoV-2 (*33, 34*). Using single-cell technologies and the SARS-

CoV-2 RBD or spike as bait, potent NAbs have now been isolated from COVID-19 patients (*35-*

41). Camelid-derived single-chain NAbs have also been obtained by panning naïve or immune

llama single-chain antibody (VHH) libraries (*42, 43*). Structures of the SARS-CoV-2 spike and

RBD in unliganded (*44, 45*), ACE2-bound (*46-48*), and antibody-bound (*49-51*) states

determined by x-ray crystallography and cryo-electron microscopy (cryo-EM) have paved the

way for rational vaccine design. Cryo-EM and cryo-electron tomography (ET) have revealed the

inherent spike metastability and the co-existence of pre/post-fusion spikes on virions (*52*). A

double-proline mutation (*52*) has been used in most soluble spike (S2P) constructs and all but

inactivated vaccines, although a HexaPro version with greater yield and stability is now available

(*53*). Cryo-ET has also uncovered a dynamic, triple-hinged HR2 stalk that facilitates viral entry

and immune evasion (*54-56*).

In this study, we designed and optimized SARS-CoV-2 antigens for multivalent display

on self-assembling protein nanoparticles (SApNPs) (*57-59*), including a ferritin (FR) 24-mer and

two 60-mers (E2p and I3-01) containing an inner layer of locking domains (LD) and a cluster of

T-cell epitopes (*60*). To facilitate tag-free vaccine purification, we developed an immunoaffinity

column based on antibody CR3022 that binds to both SARS-CoV-1/2 RBDs (*34, 50*). We first

designed a scaffolded RBD trimer construct to mimic the "RBD-up" spike conformation. The

SARS-CoV-1/2 RBDs were attached to SApNPs using the SpyTag/SpyCatcher system (*61*),

providing a robust strategy for developing RBD-based nanoparticle vaccines. We then probed

the spike metastability by comparing two uncleaved spike antigens, S2P (K986P/V987P) and

S2G (K986G/V987G). The SARS-CoV-2 S2G spike exhibited abnormal behavior, suggesting

that an unidentified facet of the spike can promote conformational change and block antibody

access to the RBD. An HR2-deleted spike, S2GΔHR2, produced high-purity trimers, suggesting

that the HR2 stalk may be a trigger of spike metastability consistent with recent findings (*54-56*).

We next displayed S2GΔHR2 on three SApNPs by gene fusion, resulting in spike SApNPs with

high yield, purity, and antigenicity. In mouse immunization, the S2P spike protein elicited the

lowest level of NAb response. In contrast, the scaffolded RBD trimer registered two-to-three-

fold higher NAb titers, with another five-fold increase in NAb titer achieved by multivalent

display on FR. S2GΔHR2 elicited up to seven-fold higher NAb titers, while the large,

multilayered S2GΔHR2 E2p and I3-01 SApNPs induced up-to-10-fold higher NAb titers than

S2P. Further analysis indicated that the S2GΔHR2-presenting I3-01v9 SApNP can elicit a strong

Th1 response as well as other types of T-cell responses needed for protective cellular immunity.

Our study thus identifies the HR2 stalk as a major source of spike metastability, validates an

HR2-deleted spike design, and provides a set of RBD- and spike-based virus-like particles

(VLPs) as effective protein vaccine candidates against SARS-CoV-2.

RESULTS

Rational design of scaffolded RBD trimers and RBD-presenting SApNPs

RBD binding to the ACE2 receptor initiates the membrane fusion process (*5*). The crystal

structure of SARS-CoV-2 RBD/ACE2 complex revealed the atomic details of receptor

recognition (*62*). The SARS-CoV-2 RBD has been used as bait to isolate monoclonal antibodies

(mAbs) from patient samples (*35-41*). For SARS-CoV-1 and MERS-CoV, RBD-based vaccines

have induced potent NAbs that effectively block viral entry (*30*). Therefore, the RBD represents

a major target for humoral responses following infection and can be used to develop epitope-

focused vaccines.

We first hypothesized that RBD attached to a trimeric scaffold could mimic the "RBD-

up" spike conformation and elicit NAbs that block ACE2 binding. To test this possibility, we

designed a fusion construct containing SARS-CoV-1/2 RBD, a short 5-aa G_4S linker (with a 2-aa

restriction site), and a trimeric viral capsid protein, SHP (PDB: 1TD0) (**Fig. 1A**). Structural

modeling showed that the three tethered RBDs form a triangle of 92 Å (measured at L492),

which is 14 and 18 Å wider than the SARS-CoV-1 "two-RBD-up" spike (PDB: 6CRX, measured

at L478) (*63*) and the MERS-CoV "all-RBD-up" spike (PDB: 5X59, measured for L506) (*64*),

respectively, allowing NAb access to each RBD. We then developed an immunoaffinity

chromatography (IAC) column to facilitate tag-free purification. Previously, NAb-derived IAC

columns have been used to purify HIV-1 Env trimers/NPs (*58, 59, 65, 66*), hepatitis C virus

(HCV) E2 cores/NPs (*57*), and Ebola virus (EBOV) GP trimers/NPs (*60*). Tian et al. reported

that a SARS-CoV-1 NAb, CR3022, can bind SARS-CoV-2 RBD (*34*). The SARS-CoV-2

RBD/CR3022 structure revealed a conserved cryptic epitope that is shared by the two SARS-

CoVs, suggesting that transient breathing motions of the spike protein enabled CR3022 binding

to the RBD (*50*). Here, we examined the utility of CR3022 in IAC columns. The SARS-CoV-1/2

RBD-5GS-1TD0 constructs were transiently expressed in 100-ml ExpiCHO cells and purified on

a CR3022 column prior to size-exclusion chromatography (SEC) using a Superdex 200 10/300

GL column. While the SARS-CoV-1 RBD construct showed both aggregate (~8.6 ml) and trimer

(~12.7 ml) peaks in the SEC profile, the SARS-CoV-2 RBD construct produced a single, pure

trimer peak at ~12.8 ml (**Fig. 1B**). In sodium dodecyl sulfate-polyacrylamide gel electrophoresis

(SDS-PAGE), a monomer band of ~37 kD and a trimer band of ~100 kD were observed under

reducing and non-reducing conditions, respectively (**fig. S1A**). Antigenicity was assessed for the

two scaffolded RBD trimers in enzyme-linked immunosorbent assay (ELISA) after CR3022/SEC

purification (**Fig. 1C**). RBD-specific NAbs targeting SARS-CoV-1 (CR3022 (*67*), m396 (*68*),

80R (*69*), and S230 (*70*)) and SARS-CoV-2 (B38 (*38*), CB6 (*37*), S309 from a SARS survivor

(*33*), and P2B-2F6 (*36*)), were tested in ELISA. Overall, similar half maximal effective

concentration (EC_{50}) values were observed for the two RBD trimers binding to their respective

NAbs (**Fig. 1C**). The SARS-CoV-1 RBD trimer showed greater affinity for CR3022 than its

SARS-CoV-2 counterpart with a 1.3-fold difference in EC_{50} values, consistent with previous

findings (*34, 50*). Of the SARS-CoV-2 NAbs, B38 yielded a similar EC_{50} value to CR3022,

which can bind but not neutralize SARS-CoV-2. Antibody binding kinetics was measured by

biolayer interferometry (BLI) (**Fig. 1D** and **fig. 1B**). Overall, all tested mAbs exhibited a fast on-

rate, but with visible differences in their off-rates. B38 showed a faster off-rate than other SARS-

CoV-2 NAbs, while CR3022, the mAb used to purify SARS-CoV-1/2 RBD proteins, exhibited a

comparable kinetic profile.

We then hypothesized that the SpyTag/SpyCatcher (termed SPY) system could be used to

conjugate RBD to SApNPs to create multivalent RBD vaccines capable of eliciting a more

potent NAb response (**Fig. 1E**). The 13-aa SpyTag spontaneously reacts with the SpyCatcher

protein to form an irreversible isopeptide bond (*61*). The SPY system has been successfully used

to attach antigens to VLPs (*71*). Here, SpyTag was fused to the C terminus of RBD, while

SpyCatcher was fused to the N terminus of an SApNP subunit, both with a 5-aa G_4S linker. This

design was first tested for the 24-meric ferritin (FR). Here, we compared two production

strategies, co-expression of RBD-5GS-SpyTag and SpyCatcher-5GS-FR versus supernatant mix

after separate expression, both followed by purification on a CR3022 column. Protein obtained

from transient transfection in 50-ml ExpiCHO cells was analyzed by SEC on a Superose 6

10/300 GL column (**Fig. 1F**). Both production strategies produced a peak (12 ml) corresponding

to SApNPs. While the SARS-CoV-2 construct outperformed its SARS-CoV-1 counterpart in

particle yield (0.6-1.0 mg versus 0.3-0.5 mg after CR3022/SEC), the supernatant mix appeared to

be superior to co-expression for yield in both cases. The results thus suggest that both strategies

can be used to produce RBD SApNPs in Chinese hamster ovary (CHO) cells in an industrial

setting, as this mammalian expression system has been widely used to manufacture therapeutic

glycoproteins under good manufacturing practice (GMP) conditions (*72, 73*). Antigenicity was

assessed for SEC-purified RBD-5GS-SPY-5GS-FR SApNPs. In ELISA, RBD SApNPs showed

slightly improved mAb binding compared to the RBD trimers, as indicated by EC_{50} values (**Fig.**

1G). In BLI, a more pronounced effect of multivalent display on antigenicity was observed,

showing notably increased binding signals and plateaued dissociation (**Fig. 1H** and **fig. 1C**).

Structural integrity of various RBD SApNPs was analyzed by negative stain EM (nsEM) (**Figs.**

1I and **1J**). For SARS-CoV-1, a genetically fused RBD-10GS-FR construct produced very few,

albeit highly pure, SApNPs (**Fig. 1I**, left). In contrast, the RBD-5GS-SPY-5GS-FR construct

produced a high yield of SApNPs with visible surface decorations (**Fig. 1I**, right). For SARS-

CoV-2, the purified RBD-5GS-SPY-5GS-FR SApNPs, irrespective of the production strategy,

showed morphology of well-formed particles (**Fig. 1J**). Previously, we genetically fused a small

LD protein and a T-cell epitope to each subunit of the E2p and I3-01v9 60-mers, resulting in

"multilayered" SApNPs (*60*). PADRE, a 13-aa pan-DR epitope that activates CD4$^+$ T cells (*74*),

was used to promote B cell development toward NAbs. Here, two of the best designs, E2p-LD4-

PADRE (or E2p-L4P) and I3-01v9-LD7-PADRE (or I3-01v9-L7P), were tested for their ability

to display SARS-CoV-1/2 RBDs. Following the strategy established for FR, SARS-CoV-1/2

RBDs were attached to the I3-01v9-L7P SApNP using the SPY system (**Fig. 1K**). Despite the

modest yield in ExpiCHO cells (**Fig. 1L**), large and pure particles were observed in the EM

images (**Fig. 1M**). However, some impurities were noted for the RBD-presenting E2p-L4P

SApNPs (**fig. S1D**). Nonetheless, we illustrated the utility of the SPY system for rapid

development of SARS-CoV-1/2 RBD vaccines based on three different SApNP platforms.

Recently, RBDs were displayed on various SApNPs using the SPY system as MERS-

CoV and SARS-CoV-2 vaccine candidates (*75, 76*). Walls et al. also reported a SARS-CoV-2

RBD vaccine candidate based on the two-component NP platform (*77*), which requires a

"connector" component to facilitate NP assembly (*78*), likely resulting in suboptimal stability.

Additionally, all these vaccine candidates require a complex production process involving two

expression systems, separate purification steps, and an *in vitro* assembly step before final

purification. In contrast, our SPY-linked RBD SApNPs are single-component by nature and can

be produced in CHO cells with a simple purification scheme, offering unique advantages in

stability and manufacturability.

Rational design of prefusion spike through minimizing metastability

In addition to the RBD, the SARS-CoV-1/2 spikes contain other NAb epitopes (*30*), which are

all presented in a trimeric context (**Fig. 2A**). A double-proline mutation (2P) between HR1 and

the central helix (CH) has been used to stabilize the MERS-CoV (*79*) and SARS-CoV-1 spikes

(*63*). A similar 2P mutation (K986P/V987P) was introduced into the SARS-CoV-2 spike (termed

S2P), which has been used to isolate and characterize NAbs (*33, 35, 40, 42-45, 49*) and is the

antigen in almost all vaccine candidates in clinical development (*11, 12*). However, a recent

cryo-EM study revealed an unexpected packing of S1 in the S2P spike, positioned ~12 Å

outwards, compared to the full-length native spike, as well as a more ordered FP proximal region

(FPPR) in S2 (*52*). New designs have been generated to control the spike conformation (*80*) or to

further stabilize it with more prolines (HexaPro) (*53*). Recent cryo-EM and cryo-ET studies

revealed that the SARS-CoV-2 spikes could adopt diverse orientations on native virions due to

the highly flexible HR2 stalk (*54-56*). Previously, we identified an HR1 bend as the cause of

HIV-1 Env metastability (*58, 81*) and examined the role of an equivalent HR1 bend and the HR2

stalk in EBOV GP metastability (*60*). This understanding of metastability proved critical for

designing stable trimers and trimer-presenting SApNP vaccines for both viruses (*58-60, 81*). It is

therefore imperative to explore the cause(s) of spike metastability to facilitate rational vaccine

design for SARS-CoV-2.

We first created uncleaved spike ectodomain (S_{ECTO}) constructs for SARS-CoV-1/2, both

containing the 2P mutation (K968P/V969P and K986P/V987P, respectively), a 5-aa G_4S linker, a

trimerization motif (PDB: 1TD0), and a C-terminal His_6 tag. The two constructs were transiently

expressed in 50-ml ExpiCHO cells followed by purification on either a Nickel column or a

CR3022 column. The $S2P_{ECTO}$-5GS-1TD0-His_6 protein was characterized by SEC on a Superose

6 10/300 GL column (**Fig. 2B**, panels 1 and 2). After the Nickel column, both $S2P_{ECTO}$

constructs showed a trimer peak (~12 ml) with the left and right shoulders indicative of

aggregate and dimer/monomer species, respectively. CR3022 purification resulted in a consistent

trimer peak, as well as reduction in dimer/monomer species. We then compared a pair of S_{ECTO}

constructs for SARS-CoV-1/2, both containing a double glycine (2G) mutation, K968G/V969G

and K986G/V987G, respectively. The 2G mutation had little effect on the SARS-CoV-1 spike

but produced abnormal SEC profiles and showed no yield for the SARS-CoV-2 spike after

purification by Nickel and CR3022 columns, respectively (**Fig. 2B**, panels 3 and 4). Lastly, we

tested a pair of S2G constructs without the HR2 stalk (E1150-Q1208), termed S2GΔHR2.

Deletion of the HR2 stalk restored the SARS-CoV-2 trimer peak and reduced aggregates for both

SARS-CoV-1/2, as shown by the SEC profiles upon CR3022 purification (**Fig. 2B**, panel 5).

Since the triple-hinged HR2 stalk can generate diverse spike orientations on native virions (*54-*

56), and the fusion core is formed by HR1 and HR2, we hypothesized that HR2 may be a key

determinant of SARS-CoV-2 spike metastability (**Fig. 2C**, left). It is plausible that the

interactions between HR1 and HR2 of two neighboring spikes may facilitate the pre-to-post-

fusion transition in addition to ACE2 binding and S1 dissociation. Given the extensive sequence

difference in HR1 (9 amino acids in total) compared to SARS-CoV-1 (**Fig. 2C**, right), we sought

to examine the contribution of HR1 to SARS-CoV-2 spike metastability with two HR1-swapped

(HR1$_S$) spike constructs. Interestingly, while HR1 swapping proved ineffective, deletion of the

HR2 stalk once again restored the trimer peak, suggesting a more important role for HR2 (**fig.**

S2, **A** to **C**). Therefore, S2GΔHR2 appeared to be a general spike design for SARS-CoV-1/2 and

perhaps other CoVs. Four separate production runs of SARS-CoV-2 S2GΔHR2-5GS-1TD0 in

300-ml ExpiCHO cells resulted in nearly identical SEC profiles with trimer yields of 0.8-1.0 mg

(**Fig. 2D**, left). Consistently, blue native polyacrylamide gel electrophoresis (BN-PAGE) showed

high trimer purity across SEC fractions (**Fig. 2D**, right). Antigenicity was assessed for

CR3022/SEC-purified SARS-CoV-2 S2P$_{ECTO}$ and S2GΔHR2 spike proteins. In ELISA, the

S2GΔHR2 spike showed slightly higher affinity for the five representative mAbs than did the

S2P$_{ECTO}$ spike (**Fig. 2E**). When tested against three newly identified human NAbs, C105 (*49*)

and CC12.1/CC12.3 (*41*), the two spikes yielded similar EC$_{50}$ values (**fig. S2D**). In BLI, the

S2GΔHR2 spike showed higher binding signals than the S2P$_{ECTO}$ spike at the highest

concentration, while exhibiting similar binding kinetics (**Fig. 2F**). The use of NAb P2B-2F6 (*36*)

for spike purification resulted in higher trimer yield with similar purity to the CR3022 column

across SEC fractions (**fig. S2E**). Thermostability was assessed by differential scanning

calorimetry (DSC) for CR3022/SEC-purified SARS-CoV-2 S2P$_{ECTO}$ and S2GΔHR2 spikes (**fig.**

S2F). Upon the 2P-to-2G substitution and HR2 deletion, a small increase in thermal denaturation

midpoint (T$_m$) was observed (46.8 vs. 47.6 °C), with notably higher onset temperature (T$_{on}$), 35.2

vs. 40.0 °C, and narrower half width of the peak $(\Delta T_{1/2})$, 4.7 vs. 3.9 °C. Altogether, we

demonstrated that the HR2 stalk is a major source of spike metastability and S2GΔHR2 presents

an alternative spike design to S2P, although more stabilizing mutations may be required to

achieve greater thermostability.

Rational design of single-component self-assembling spike nanoparticles

Although it was possible to conjugate trimeric SARS-CoV-2 spikes to an SApNP using the SPY

system (*82*), the random, irreversible linking will likely result in irregular display with

unoccupied but spatially occluded anchoring sites on the surface. The SPY system is perhaps

more suitable for small individual antigens (e.g. the RBD). Using the gene fusion approach, we

previously designed single-component SApNPs displaying stabilized HIV-1 Env trimers (*58, 59*)

and optimized HCV E2 cores (*57*). Recently, we reengineered E2p and I3-01v9 60-mers to

incorporate an inner layer of LDs to stabilize the non-covalently formed NP shell, in addition to

a cluster of T-cell epitopes (*60*). This multilayered design may be essential to the stability of

resulting SApNP vaccines when they are used to display large, complex viral antigens such as

the SARS-CoV-2 spike.

Native SARS-CoV-2 virions present both pre- and post-fusion spikes (*52, 54, 55*) (**Fig.**

3A, top), while our vaccine strategy aims to develop single-component SApNPs that each

present 8 or 20 stable prefusion S2GΔHR2 spikes to the immune system (**Fig. 3A**, bottom). As

demonstrated in our previous studies (*58, 59*), different linker lengths may be needed to connect

a trimer to the SApNP surface, as the spacing between the N termini of NP subunits around each

three-fold axis varies: FR (20 Å), E2p (9 Å), and I3-01v9 (50.5 Å). Based on this consideration,

we displayed the S2GΔHR2 spike on FR with a 5-aa G_4S linker, on E2p with a 5-aa G_4S linker,

and on I3-01v9 with a 10-aa $(G_4S)_2$ linker, resulting in SApNPs with diameters of 47.9, 55.9, and

59.3 nm, respectively (**Fig. 3B**). The multilayered E2p-L4P and I3-01v9-L7P (*60*), which were

validated for presenting RBDs (**Fig. 1**, **K** to **L**; **fig. S1D**), were used here as NP carriers of the

S2GΔHR2 spike. Together, three S2GΔHR2 SApNP constructs were transiently expressed in

400-ml ExpiCHO cells, followed by CR3022 purification and SEC on a Superose 6 10/300 GL

column (**Fig. 3C**). Three production runs for each of the three constructs generated highly

consistent SEC profiles, despite the variation of low molecular weight (m.w.) impurities

observed for the FR and E2p-L4P. After CR3022 and SEC purification, we obtained 0.3-0.4, 0.5-

1.0, and 0.8-1.2 mg protein for S2GΔHR2-5GS-FR, S2GΔHR2-5GS-E2p-L4P, and S2GΔHR2-

10GS-I3-01v9-L7P, respectively. Overall, the I3-01v9-derived S2GΔHR2 SApNP appeared to

perform best in terms of particle yield, purity, and stability. The structural integrity was analyzed

by nsEM, which showed well-formed particles of 45-65 nm with recognizable protrusions on the

surface (**Fig. 3D**). The varying shape of these profusions may correspond to S2GΔHR2 spikes

with "open" RBDs, in contrast to an array of "closed" HIV-1 and EBOV trimers on the SApNP

surface (*58-60*). Such open conformation may facilitate induction of a strong RBD-specific NAb

response *in vivo*. Antigenicity of S2GΔHR2 SApNPs was assessed using the same panel of

mAbs. In ELISA, three SApNPs showed slightly improved binding to some, but not all, mAbs

compared to the individual spike (**Fig. 3E**). In BLI, we observed a clear correlation between peak

mAb binding signal and antigen valency, with E2p/I3-01v9 > FR > spike (**Fig. 3F**). Multivalent

display on the two 60-mers substantially increased mAb binding compared to the FR 24-mer. In

previous studies, we observed a similar correlation for HIV-1 Env trimer and HCV E2 core

versus their SApNPs (*57, 58*). In summary, these VLP-size SApNPs with 8 or 20 spikes on the

surface provide promising vaccine candidates for *in vivo* evaluation.

SARS-CoV-1/2 vaccine-induced binding antibody response

Selected SARS-CoV-1/2 RBD- and spike-based vaccine constructs were assessed in BALB/c

mice (**Fig. 4A**). We adopted a similar immunization protocol to be consistent with our previous

studies on HIV-1, HCV, and EBOV SApNPs (*57, 58, 60*). Briefly, groups of five mice were

immunized four times at three-week intervals. All antigens (50 μg per dose) were formulated

with AddaVax, an oil-in-water emulsion adjuvant, except for I3-01v9, which was mixed with

aluminum phosphate (AP) (*83*). We first analyzed the binding antibody (bAb) responses, as

measured by EC_{50} titers, in the two SARS-CoV-2 RBD vaccine groups (**Fig. 4B** and **fig. S4**).

The SPY-linked RBD SApNP (RBD-5GS-SPY-5GS-FR) elicited significantly higher bAb titers

than the scaffolded RBD trimer (RBD-5GS-1TD0) at w2 and w5, irrespective of the coating

antigen, and yielded a significant *P* value at w8 when the RBD was coated. Compared to the

$S2P_{ECTO}$ spike ($S2P_{ECTO}$-5GS-1TD0), the RBD SApNP elicited significantly higher bAb titers

against the RBD at w2, w5, and w8 (**Fig. 4B**, right), demonstrating a strong epitope-focusing

effect. Mouse sera bound the SARS-CoV-1 spike with lower EC_{50} titers than the SARS-CoV-2

spike but with similar patterns (**fig. S4A**). We then analyzed the bAb responses induced by

SARS-CoV-2 spikes S2P$_{ECTO}$-5GS-1TD0 and S2GΔHR2-5GS-1TD0, as well as three SApNPs

each displaying 8 or 20 S2GΔHR2 spikes (**Fig. 4C** and **fig. S5**). The S2GΔHR2 spike showed

two-three-fold higher average EC$_{50}$ titers than the S2P$_{ECTO}$ spike irrespective of the coating

antigen (of note, to facilitate a fair comparison, mouse sera from these two groups were tested

against their respective spike antigens). Three SApNPs exhibited different temporal patterns

depending on the coating antigen. In terms of spike-specific response, the I3-01v9 group

registered a steady increase in average EC$_{50}$ titer over time, showing the highest bAb titers at w2

and w8 and significantly outperforming the S2P$_{ECTO}$ group at all time points. The I3-01v9 group

also yielded higher EC$_{50}$ titers than the S2GΔHR2 group throughout, although not with

significant *P* values. The FR SApNP group exhibited a similar temporal pattern with lower EC$_{50}$

titers, which were still significantly higher than the S2P$_{ECTO}$ group. Among the three SApNPs,

E2p exhibited the lowest average EC$_{50}$ titer at w2 and reached the highest at w5, which then

decreased slightly at w8. In terms of RBD-specific response, the five groups showed a clear

ranking based on their average EC$_{50}$ titers, which remained consistent across time points. At w2,

I3-01v9 showed an average EC$_{50}$ titer of 175, whereas all other spike-based vaccines induced

little RBD-specific bAb response. At w5 and w8, S2GΔHR2 elicited higher bAb titers (on

average by two-fold) than S2P$_{ECTO}$, while all three SApNPs outperformed the individual

S2GΔHR2 spike with a ranking of average EC$_{50}$ titers correlated with their size (FR < E2p < I3-

01v9). Sera reacted with the SARS-CoV-1 spike similarly, albeit at a lower level (**fig. S5A**).

Lastly, we compared the bAb responses induced by three SARS-CoV-1 vaccines, S2P$_{ECTO}$ spike

(S2P$_{ECTO}$-5GS-1TD0), scaffolded RBD trimer (RBD-5GS-1TD0), and SPY-linked RBD SApNP

(RBD-5GS-SPY-5GS-FR) (**Fig. 4D** and **fig. S6**). Based on average EC$_{50}$ titers, the SARS-CoV-1

S2P$_{ECTO}$ spike appeared to have elicited a more robust bAb response than the SARS-CoV-2

S2GΔHR2 spike, whereas the SARS-CoV-1 RBD SApNP was less advantageous than its SARS-

COV-2 counterpart. Serum reactivity with the SARS-CoV-2 S2P$_{ECTO}$ spike was observed for all

three SARS-CoV-1 vaccine groups (**fig. S6A**). In summary, RBD SApNPs can elicit higher titers

of RBD-specific bAbs than the scaffolded RBD trimer and S2P$_{ECTO}$ spike, albeit at different

levels for the two SARS-CoVs. The S2GΔHR2 spike is more immunogenic than the S2P$_{ECTO}$

spike, showing on average two-fold higher bAb titers. The S2GΔHR2-presenting E2p and I3-

01v9 SApNPs are the best performers among all the spike-based vaccines, consistent with our

previous findings for these two large SApNPs (*57, 58, 60*).

SARS-CoV-1/2 vaccine-induced NAb response

One major goal in COVID-19 vaccine development is to generate a potent NAb response that

can protect against SARS-CoV-2 infection. Pseudoparticle (SARS-CoV-1/2-pp) neutralization

assays (*84*) were used to evaluate serum NAb responses elicited by different vaccines. We first

analyzed the NAb responses, as measured by 50% inhibitory dilution (ID$_{50}$) titers, in the two

SARS-CoV-2 RBD vaccine groups (**Fig. 5A** and **fig. S7**). The SPY-linked RBD SApNP elicited

a NAb response against autologous SARS-CoV-2 as early as w2, albeit with low titers, and

retained its advantage at w5 and w8, suggesting that such RBD SApNP vaccines can elicit a

rapid NAb response. The scaffolded RBD trimer group showed higher average ID$_{50}$ titers than

the S2P$_{ECTO}$ spike group at both w5 and w8. A somewhat different pattern was observed in the

SARS-CoV-1-pp assay. At w2, no vaccine group showed detectable heterologous NAb response.

At w5 and w8, the S2P$_{ECTO}$ spike elicited a more potent anti-SARS-CoV-1 NAb response than

the scaffolded RBD trimer, suggesting that non-RBD epitopes on the spike may contribute to

cross-neutralization. We then analyzed the NAb responses induced by five spike-based vaccines

(**Fig. 5B** and **fig. S8**). In terms of autologous neutralization, no spike-based vaccine elicited any

NAb response that blocks SARS-CoV-2-pps after one dose, but a consistent pattern of serum

neutralization was observed at w5 and w8 (**Fig. 5B**, upper panel). Specifically, the $S2P_{ECTO}$ spike

showed the lowest average ID_{50} titers, 879 and 2481 at w5 and w8, respectively, whereas the

S2GΔHR2 spike induced a stronger NAb response with 2.8-6.7-fold higher average ID_{50} titers,

which did not reach $P \le 0.05$ due to within-group variation. Nonetheless, this result confirmed

the beneficial effect of the 2G mutation and the HR2 stalk deletion on NAb elicitation. Among

the three SApNPs, E2p was the best performer at w5, showing an average ID_{50} titer of 8435 that

is 9.6-fold higher than $S2P_{ECTO}$ and 1.4-fold higher than S2GΔHR2, while I3-01v9 showed the

most potent NAb response at w8 with an average ID_{50} titer of 17351 that is about 7-fold and 2.5-

fold higher than $S2P_{ECTO}$ and S2GΔHR2, respectively. A similar temporal pattern was observed

in the heterologous SARS-CoV-1-pp assay (**Fig. 5B**, lower panel). It is worth noting that the

multilayered I3-01v9 SApNP elicited a SARS-CoV-1 NAb response with an average ID_{50} titer of

351 at w2, whereas all other groups showed no detectable serum neutralization. These results

suggest that a well-designed SARS-CoV-2 S2GΔHR2 SApNP vaccine may provide protection

against both SARS-CoVs. Lastly, we analyzed the NAb responses induced by three SARS-CoV-

1 vaccines (**Fig. 5C** and **fig. S9**). In the autologous SARS-CoV-1-pp assay, the $S2P_{ECTO}$ spike

and the RBD SApNP induced significantly higher NAb titers than the scaffolded RBD trimer at

w2 and w5, and all three vaccine groups showed similar ID_{50} titers at w8. However, heterologous

SARS-CoV-2 neutralization was below or at the baseline level for three SARS-CoV-1 vaccines

at w2, w5, and w8. In this study, the pseudovirus neutralization assay was validated using a panel

of known SARS-CoV-1/2 NAbs (**fig. S9C**). As a negative control, the w8 mouse sera were tested

against pseudoparticles bearing the murine leukemia virus (MLV) Env, MLV-pps, showing no

detectable reactivity (**fig. S9D**). In summary, these results demonstrate an advantage in NAb

elicitation by the SARS-CoV-2 S2GΔHR2 spike and its SApNPs compared to the widely used

S2P$_{ECTO}$ spike. Although SARS-CoV-2 RBD- and S2GΔHR2-presenting SApNPs are both

effective at eliciting an autologous NAb response, the latter can induce high NAb titers to SARS-

CoV-1 and may provide broader protection against SARS-associated CoVs.

SARS-CoV-2 vaccine-induced T-cell response

While humoral immunity is required to block host-virus interaction and prevent viral infection,

cellular immunity is essential for eliminating infected host cells to control viral infection (*85-88*).

Emerging evidence indicates that an early T-cell response (*89, 90*), as well as T-cell memory

(*91*), is critical for protection against SARS-CoV-2. However, COVID-19 vaccines must induce

a CD4$^+$ T helper 1 (Th1), but not Th2-type, T-cell response, as the latter has been implicated in

vaccine-associated enhancement of respiratory disease (VAERD) (*10*). In addition, T follicular

helper cells (Tfh) play an important role in the maturation and production of NAbs. Therefore,

understanding T-cell responses is crucial for development of an effective and safe COVID-19

vaccine.

Interferon (IFN)-γ-producing Th1 cells are important for generating an optimal antibody

response and for induction of cellular immunity (*85-87*). We first examined various SARS-CoV-

2 vaccines on induction of CD4$^+$ Th1 responses specific to the vaccine antigen at w11, two

weeks after the fourth immunization, when memory T cells had already developed in spleen (*88*).

Mouse splenocytes from the S2P$_{ECTO}$ group and two S2GΔHR2 SApNP groups (E2p and I3-

01v9) were analyzed by flow cytometry using naïve samples as a negative control. The I3-01v9

group induced about 1.5- and 2.3-fold higher frequency of IFN-γ-producing CD4$^+$ Th1 cells than

the S2P$_{ECTO}$ and E2p groups, respectively (**Fig. 6A**). Notably, after re-stimulation with the respective antigens for as few as 4 hours, both SApNP groups produced ~two-fold higher frequency of CD107a-producing cytolytic CD4$^+$ T cells than the S2P$_{ECTO}$ and naïve groups (**Fig. 6B**). IFN-γ/IL-4 (interleukin-4) double-positive cells are memory CD4$^+$ T cells that have acquired the ability to produce IL-4 while still retaining the ability to produce IFN-γ under Th1 conditions (*92*). It appeared that I3-01v9 induced three- and five-fold more IFN-γ/IL-4 double-positive memory CD4$^+$ T cells than S2P$_{ECTO}$ and E2p (**Fig. 6A**). These results suggest that I3-01v9 can induce both CD4$^+$ Th1 cells and IFN-γ/IL-4 double-positive memory CD4$^+$ T cells. In addition, I3-01v9 induced more IFN-γ/GM-CSF (granulocyte-macrophage colony-stimulating factor) double-positive CD8$^+$ effector T cells than S2P$_{ECTO}$ and E2p (**Fig. 6C**), suggesting that protective CD8$^+$ T cells were also generated in mice immunized with I3-01v9. Of note, CD8$^+$ T cells derived from mice immunized with I3-01v9, rather than those with S2P$_{ECTO}$ and E2p, acquired the ability to rapidly produce IFN-γ upon antigen re-stimulation (**Fig. 6D**), suggesting generation of I3-01v9-responsive effector/memory T cells. Our results indicate that the S2GΔHR2 I3-01v9 SApNP can induce robust T-cell responses consisting of CD4$^+$ Th1 cells, IFN-γ/IL-4 double-positive memory CD4$^+$ T cells, and effector CD8$^+$ T cells, thus providing protective cellular immunity in addition to a potent NAb response. Since T cell immunity against the SApNP backbone cannot be ruled out, a more detailed T-cell analysis using soluble spikes, spike peptides, and naked SApNPs for re-stimulation may be warranted.

DISCUSSION

COVID-19 is the first worldwide pandemic of this scale since the infamous Spanish influenza over a century ago (*93*), which caused ~50 million deaths worldwide and remains a painful reminder of our vulnerability to a new virus without a protective vaccine. The rapid spread of

SARS-CoV-2 therefore demanded rapid vaccine development (*10*). Operation Warp Speed

(OWS) was launched in May 2020 with the initial goal to deliver 300 million doses of safe and

effective vaccines by January 2021 (*94*), although that goal has not yet been realized.

Nevertheless, in December 2020, while other vaccine candidates were still being evaluated in

clinical trials, two OWS-supported mRNA vaccines were approved for EUA, marking an

important turning point in the battle against the pandemic. However, vaccine development during

a pandemic against a new virus poses unique challenges, one of which is how to balance public

health needs versus scientific rigor (*95-97*). The global vaccine campaign also provides a unique

opportunity to compare different vaccine design strategies and platforms – especially new ones –

against a common target. While mRNA and viral vector vaccines remain the front runners,

protein-based vaccines are highlighted in a recent review with the prediction that they will

eventually reach a larger fraction of the global population (*98*). As the NAb titers induced by the

first-generation nucleic acid vaccines wane over time, effective protein vaccines will be needed

to sustain long-term immunity against SARS-CoV-2.

Here, we approached SARS-CoV-2 vaccine with a rational design strategy and set out to

develop protein nanoparticle vaccines that can be used alone or as a booster vaccine. To this end,

a panel of vaccine candidates have been generated based on three SApNP platforms with

different *in vitro* and *in vivo* attributes (**table S1**). Our comparative analysis of these vaccine

constructs has offered some valuable insights. First, the choice of antigen is critical to the

success of SARS-CoV-2 vaccine irrespective of the delivery platform. Most vaccine antigens

including OWS's vaccine candidates are based on S2P, which produces a spike structure that

differs in detail from the full-length wildtype spike, e.g. in FPPR of S2 and in the relative

dispositions of the S1 domains (*52*). These differences may complicate interpretation of vaccine

outcome. S2P and other empirical spike designs (*53*) have attempted to constrain the spike

conformation and increase trimer yield. However, as we previously demonstrated for HIV-1 Env

and EBOV GP (*58, 60, 81*), it is important to identify and eliminate (if possible) the cause of

spike metastability. During antigen screening, we found that deletion of the HR2 stalk with a 2P-

to-2G substitution renders a more stable spike, which is consistent with recent reports on a highly

flexible HR2 stalk in the native spikes on SARS-CoV-2 virions (*54-56*). Thus, S2GΔHR2 would

seem to represent a major advance in spike design. Second, single-component SApNPs provide a

powerful platform for vaccine development against diverse viral pathogens (*57, 58, 60*). Here,

S2GΔHR2 was genetically fused, rather than chemically linked, to three SApNPs, including two

multilayered 60-meric carriers with enhanced stability and an embedded T-help signal. Such

protein vaccines should be more effective in eliciting a potent NAb response and less likely to

induce adverse responses (*98, 99*). An epitope-focused vaccine strategy was also explored by

designing scaffolded RBD trimers and SPY-linked RBD SApNPs. Third, to achieve high

efficacy and ensure safety, vaccine-induced NAb and T-cell responses must be evaluated in

animals prior to clinical trials. Indeed, in our mouse study, S2GΔHR2 appeared to be more

effective than S2P in NAb elicitation, both alone and displayed on SApNPs. Of note, the

multilayered S2GΔHR2 I3-01v9 SApNP elicited not only high NAb titers but also desired T-cell

responses. In addition to CD4$^+$ Th1 cells and memory CD4$^+$ T cells, this SApNP also induced

CD107a-producing cytolytic CD4$^+$ T cells, which may directly kill infected host cells, and GM-

CSF-producing CD8$^+$ effector T cells, which may promote the generation of macrophages and

functional dendritic cells (DCs) to facilitate the clearance of infected cells. It is also worth noting

that the NAb response to the large SApNPs plateaued after three doses, suggesting that the

number of injections and dosage can be reduced without diminishing vaccine-induced responses.

Lastly, expression of vaccine antigens in CHO cells followed by purification using an antibody

column, such as CR3022, would allow rapid and industrial-scale vaccine production. In

summary, our study provides promising COVID-19 vaccine candidates for evaluation in clinical

trials.

MATERIALS AND METHODS

Design, expression, and purification of SARS-CoV-1/2 RBD and spike antigens

The spike (S) genes of the SARS-CoV-1 isolate Tor2 (GenBank accession #: NC_004718) and

the SARS-CoV-2 isolate Wuhan-Hu-1 (GenBank accession #: MN908947) were used to design

all the RBD and spike constructs following codon-optimization for expression in mammalian

cells. The RBD sequence is defined as P317-D518 and P330-N532 for SARS-CoV-1 and 2,

respectively. The S_{ECTO} sequence is defined as M1-Q1190 and M1-Q1208 for SARS-CoV-1 and

2, respectively. To remove the S1/S2 cleavage site, an R667G mutation and a $^{682}GSAGSV^{687}$

modification were introduced in the SARS-CoV-1 and 2 spikes, respectively. The 2P (or 2G)

mutation was made to K968/V969 and K986/V987 in the SARS-CoV-1 and 2 spikes,

respectively. The SARS-CoV-2 C-terminal region (E1150-Q1208) containing the HR2 stalk was

removed from $S2G_{ECTO}$, resulting in an HR2-deleted spike construct, termed S2GΔHR2. The

viral capsid protein SHP (PDB: 1TD0) was used as a trimerization motif in spike constructs for

immunization, whereas the foldon domain from the bacteriophage T4 fibritin (PDB: 1RFO) was

used in coating spike antigens for ELISA to mask the 1TD0-derived antibody response. All

constructs were transiently expressed in ExpiCHO cells (Thermo Fisher). Briefly, ExpiCHO

cells were thawed and incubated with ExpiCHO^{TM} Expression Medium (Thermo Fisher) in a

shaker incubator at 37 °C, 135 rpm and 8% CO_2. When the cells reached a density of 10×10^6 ml$^-$

^{1}, ExpiCHO^{TM} Expression Medium was added to reduce cell density to 6×10^6 ml^{-1} for

transfection. The ExpiFectamine[TM] CHO/plasmid DNA complexes were prepared for 100-ml

transfection in ExpiCHO cells following the manufacturer's instructions. For a given construct,

100 μg of plasmid and 320 μl of ExpiFectamine[TM] CHO reagent were mixed in 7.7 ml of cold

OptiPRO™ medium (Thermo Fisher). After the first feed on day one, ExpiCHO cells were

cultured in a shaker incubator at 33 °C, 115 rpm and 8% CO_2 following the Max Titer protocol

with an additional feed on day five (Thermo Fisher). Culture supernatants were harvested 13 to

14 days after transfection, clarified by centrifugation at 4000 rpm for 25 min, and filtered using a

0.45 μm filter (Thermo Fisher). The CR3022 antibody column was used to extract SARS-CoV-

1/2 antigens from the supernatants, which was followed by SEC on a Superdex 200 10/300 GL

column (for scaffolded RBD trimers) or a Superose 6 10/300 GL column (for SPY-linked RBD

SApNPs, spikes, and S2GΔHR2 SApNPs). A human NAb, P2B-2F6 (*36*), was also used to pack

antibody columns for purification of the SARS-CoV-2 S2GΔHR2 spike, which was followed by

SEC on a HiLoad 16/600 Superose 6 column. For comparison, His_6-tagged S_{ECTO}-5GS-1TD0

spike protein was extracted from the supernatants using an immobilized Ni Sepharose[TM] Excel

column (GE Healthcare) and eluted with 500 mM Imidazole prior to SEC. Protein concentration

was determined using UV_{280} absorbance with theoretical extinction coefficients.

Blue native polyacrylamide gel electrophoresis

SARS-CoV-2 spikes and spike-presenting SApNPs were analyzed by blue native polyacrylamide

gel electrophoresis (*BN-PAGE*) and stained with Coomassie blue. The proteins were mixed with

sample buffer and G250 loading dye and added to a 4-12% Bis-Tris NativePAGE[TM] gel (Life

Technologies). BN-PAGE gels were run for 2 to 2.5 hours at 150 V using the NativePAGE[TM]

running buffer (Life Technologies) according to the manufacturer's instructions.

 Enzyme-linked immunosorbent assay

 Each well of a Costar[TM] 96-well assay plate (Corning) was first coated with 50 µl PBS

 containing 0.2 µg of the appropriate antigens. The plates were incubated overnight at 4 °C, and

 then washed five times with wash buffer containing PBS and 0.05% (v/v) Tween 20. Each well

 was then coated with 150 µl of a blocking buffer consisting of PBS and 40 mg ml^{-1} blotting-

 grade blocker (Bio-Rad). The plates were incubated with the blocking buffer for 1 hour at room

 temperature, and then washed five times with wash buffer. For antigen binding, antibodies (in

 the immunoglobulin G (IgG) form) were diluted in the blocking buffer to a maximum

 concentration of 5 µg ml^{-1} followed by a 10-fold dilution series. For each antibody dilution, a

 total of 50 µl volume was added to the appropriate wells. For mouse sample analysis, plasma

 was diluted by 20-fold in the blocking buffer and subjected to a 10-fold dilution series. For each

 sample dilution, a total of 50 µl volume was added to the wells. Each plate was incubated for 1

 hour at room temperature, and then washed 5 times with PBS containing 0.05% Tween 20. For

 antibody binding, a 1:5000 dilution of goat anti-human IgG antibody (Jackson ImmunoResearch

 Laboratories, Inc), or for mouse sample analysis, a 1:3000 dilution of horseradish peroxidase

 (HRP)-labeled goat anti-mouse IgG antibody (Jackson ImmunoResearch Laboratories), was then

 made in the wash buffer (PBS containing 0.05% Tween 20), with 50 µl of this diluted secondary

 antibody added to each well. The plates were incubated with the secondary antibody for 1 hour at

 room temperature, and then washed 6 times with PBS containing 0.05% Tween 20. Finally, the

 wells were developed with 50 µl of TMB (Life Sciences) for 3-5 min before stopping the

 reaction with 50 µl of 2 N sulfuric acid. The resulting plate readouts were measured at a

 wavelength of 450 nm. Of note, the w2 serum binding did not reach the plateau (or saturation) to

 allow for accurate determination of EC$_{50}$ titers. Nonetheless, the EC$_{50}$ values at w2 were derived

by setting the lower/upper constraints of OD_{450} at 0.0/3.2 to facilitate the comparison of different

vaccine groups at the first time point.

Bio-layer interferometry

The kinetics of SARS-CoV-1/2 vaccine antigens, RBD versus RBD-presenting SApNPs as well

as spike versus spike-presenting SApNPs, binding to a panel of known antibodies (in the IgG

form) was measured using an Octet RED96 instrument (FortéBio, Pall Life Sciences). All assays

were performed with agitation set to 1000 rpm in FortéBio 1× kinetic buffer. The final volume

for all the solutions was 200 μl per well. Assays were performed at 30 °C in solid black 96-well

plates (Geiger Bio-One). For all antigens except for S2GΔHR2 SApNPs, 5 μg ml^{-1} of antibody in

1× kinetic buffer was loaded onto the surface of anti-human Fc Capture Biosensors (AHC) for

300 s. For S2GΔHR2 SApNPs, anti-human Fc Quantitation Biosensors (AHQ) were used. A 60 s

biosensor baseline step was applied prior to the analysis of the association of the antibody on the

biosensor to the antigen in solution for 200 s. A two-fold concentration gradient of antigen,

starting at 950 nM for scaffolded RBD trimers, 37 nM for SPY-linked RBD FR SApNP (RBD-

5GS-SPY-5GS-FR), 150 nM for soluble spikes, and 9/3.5/3.5 nM for S2GΔHR2 presented on

FR/E2p/I3-01v9 SApNPs, was used in a titration series of six. The dissociation of the interaction

was followed for 300 s. Correction of baseline drift was performed by subtracting the mean value

of shifts recorded for a sensor loaded with antibody but not incubated with antigen, and for a

sensor without antibody but incubated with antigen. Octet data were processed by FortéBio's

data acquisition software v.8.1. Experimental data were fitted with the binding equations

describing a 2:1 interaction to achieve optimal fitting. Of note, the S2GΔHR2 spike was also

measured using AHQ biosensors to facilitate comparison of mAb binding with the S2GΔHR2-

presenting SApNPs.

Differential scanning calorimetry (DSC)

Thermal melting curves of SARS-CoV-2 S2$_{PECTO}$-5GS-1TD0 and S2GΔHR2-5GS-1TD0 spike proteins following CR3022 and SEC purification were obtained from a MicroCal PEAQ-DSC Man instrument (Malvern). The purified spike trimer protein produced from ExpiCHO cells were buffer exchanged into 1×PBS and concentrated to 0.8 μM before analysis by the instrument. Melting was probed at a scan rate of 60 °C·h^{-1} from 20 °C to 100 °C. Data processing, including buffer correction, normalization, and baseline subtraction, was conducted using the MicroCal PEAQ-DSC software. Gaussian fitting was performed using the Origin 9.0 software.

Electron microscopy (EM) assessment of nanoparticle constructs

The EM analysis of various RBD and S2GΔHR2-presenting SApNPs was performed by the Core Microscopy Facility at The Scripps Research Institute. All SApNP samples were prepared at the concentration of 0.01-0.05 mg/ml. Carbon-coated copper grids (400 mesh) were glow-discharged and 8 µL of each sample was adsorbed for 2 min. Excess sample was wicked away and grids were negatively stained with 2% uranyl formate for 2 min. Excess stain was wicked away and the grids were allowed to dry. Samples were analyzed at 80 kV with a Talos L120C transmission electron microscope (Thermo Fisher) and images were acquired with a CETA 16M CMOS camera.

Animal immunization and sample collection

Similar immunization protocols have been reported in our previous vaccine studies (*57, 58, 60*). Briefly, the Institutional Animal Care and Use Committee (IACUC) guidelines were followed with animal subjects tested in the immunization study. Eight-week-old BALB/c mice were purchased from The Jackson Laboratory and housed in ventilated cages in environmentally

controlled rooms at The Scripps Research Institute, in compliance with an approved IACUC

protocol and AAALAC (Association for Assessment and Accreditation of Laboratory Animal

Care) International guidelines. Mice were immunized at weeks 0, 3, 6, and 9 with 200 μl of

antigen/adjuvant mix containing 50 μg of vaccine antigen and 100 μl of adjuvant, AddaVax or

Adju-Phos (InvivoGen), via the intraperitoneal (i.p.) route. Blood was collected two weeks after

each immunization. All bleeds were performed through the retro-orbital sinus using heparinized

capillary tubes into EDTA-coated tubes. Samples were spun at 1200 RPM for 10 min to separate

plasma (top layer) and the rest of the whole blood layer. Upon heat inactivation at 56 °C for 30

min, the plasma was spun at 2000 RPM for 10 min to remove precipitates. The rest of the whole

blood layer was diluted with an equal volume of PBS and then overlaid on 4.5 ml of Ficoll in a

15 ml SepMate^{TM} tube (STEMCELL Technologies) and spun at 1200 RPM for 10 min at 20 °C

to separate peripheral blood mononuclear cells (PBMCs). Cells were washed once in PBS and

then resuspended in 1 ml of ACK Red Blood Cell lysis buffer (Lonza). After washing with PBS,

PBMCs were resuspended in 2 ml of Bambanker Freezing Media (Lymphotec). Spleens were

harvested at w11 and ground against a 70-μm cell strainer (BD Falcon) to release the splenocytes

into a cell suspension. Splenocytes were centrifuged, washed in PBS, treated with 5 ml of ACK

lysing buffer (Lonza), and frozen with 3ml of Bambanker freezing media. Plasma was used for

ELISA and neutralization assays to determine binding and neutralizing antibody responses in

mouse serum.

SARS-CoV-1/2 pseudovirus neutralization assay

Pseudoparticle (SARS-CoV-1/2-pp) neutralization assays were utilized to assess the neutralizing

activity of previously reported antibodies and vaccine-induced murine antibody response. SARS-

CoV-1/2-pps were generated by co-transfection of HEK293T cells with the HIV-1 pNL4-3.lucR-

E- plasmid (obtained from the NIH AIDS reagent program: https://www.aidsreagent.org/) and

the expression plasmid encoding the S gene of SARS-CoV-1 isolate Tor2 (GenBank accession #:

NC_004718) and the SARS-CoV-2 isolate Wuhan-Hu-1 (GenBank accession #: MN908947) at a

4:1 ratio by lipofectamine 3000 (Thermo Fisher Scientific). After 48 to 72 hours, SARS-CoV-

1/2-pps were collected from the supernatant by centrifugation at 4000 rpm for 10 min, aliquoted,

and stored at -80 °C before use. The mAbs at a starting concentration of 0.1-10 µg/ml, or mouse

plasma at a starting dilution of 100-fold, were mixed with the supernatant containing SARS-

CoV-1/2-pps and incubated for 1 hour at 37°C in white solid-bottom 96-well plate (Corning). A

3-fold dilution series was used in the assay. The HEK293T-hACE2 cell line (catalogue#: NR-

52511) and the vector pcDNA3.1(-) containing the SARS-CoV-2 S gene (catalogue#: NR52420)

were obtained from BEI RESOURCES (https://www.beiresources.org/) and used in pseudovirus

neutralization assays (*84*). Briefly, HEK293T-hACE2 cells at 1×10^4 were added to each well and

the plate was incubated at 37°C for 48 hours. After incubation, overlying media was removed,

and cells were lysed. The firefly luciferase signal from infected cells was determined using the

Bright-Glo Luciferase Assay System (Promega) according to the manufacturer's instructions.

Data were retrieved from a BioTek microplate reader with Gen 5 software, the average

background luminescence from a series of uninfected wells was subtracted from each well, and

neutralization curves were generated using GraphPad Prism 8.4.3, in which values from wells

were compared against a well containing SARS-CoV-1/2-pp only. The same HIV-1 vectors

pseudotyped with the murine leukemia virus (MLV) Env gene, termed MLV-pps, were produced

in HEK293T cells and included in the neutralization assays as a negative control. As the NAb

titers plateaued after w8, the w11 results were not shown in Fig. 5 but included in figs. S7-S9 for

comparison.

Dendritic cell (DC) production

Mouse bone marrow (BM) was cultured in RPMI 1640 medium containing 10% fetal bovine

serum and recombinant mouse Flt3L (50 ng/mL) and SCF (10 ng/ml) for 9 days as described

(*100*). To induce DC activation, immature DCs were incubated with lipopolysaccharide (LPS,

100 ng/mL), R848 (Resiquimod, 100 ng/mL) or CpG (ODN 1585, 1µM) overnight, which

activated Toll-like receptor (TLR)4 , TLR7/8 or TLR9 signaling, respectively. Cells were

harvested for experiments. pDCs were sorted to isolate $CD11c^+B220^+$ cells using FACS cell

sorter and magnetic beads (Miltenyi-Biotech, CA).

Antibodies and flow cytometry analysis

All antibodies used for immunofluorescence staining were purchased from eBioscience (San

Diego, CA), BioLegend (San Diego, CA) or BD Biosciences (San Jose, CA). Magnetic

microbead-conjugated Abs and streptavidin were purchased from Miltenyi-Biotech (Auburn,

CA). Recombinant human IL-2 protein was purchased from R&D Systems (Minneapolis, MN).

Recombinant mouse Flt3 ligand (Flt3L) and mouse SCF were purchased from Shenandoah

Biotech (Warwick, PA). Cells were stained with appropriate concentrations of mAbs. Dead cells

were excluded using Fixable Viability Dye from eBioscience (San Diego, CA). Flow cytometry

(FC) analyses were performed using LSRII (BD Bioscience, CA) and Canto cytometers (Becton

Dickinson, NJ). Cells were sorted on BD FACSAria II (BD Bioscience, CA).

T cell culture and activation

Splenic mononuclear cells from each group of immunized mice were cultured in the presence of

DCs pulsed with or without $S2P_{ECTO}$, multilayered S2GΔHR2-presenting E2P or I3-01v9 SApNP

$(1 \times 10^{-7} \, \mu\text{M})$ in complete IMDM medium containing IL-2 (5.0 ng/ml). Cells were collected 16

hours and 4 hours later for intracellular cytokine staining and flow cytometric analysis.

Statistics

In antibody analysis, comparison of different vaccine groups was performed in GraphPad Prism

8.4.3 using the two-tailed unpaired Student's *t* test. In T cell analysis, comparison of means was

done using the two-tailed unpaired Student's t test, ANOVA and then post-hoc *t* test. P values of

0.05 or less were considered significant.

SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at http://xxx/xxx/xxx.

fig. S1. In-vitro characterization of SARS-CoV-1/2 RBD-based immunogens.

fig. S2. In-vitro characterization of SARS-CoV-2 spike antigens.

fig. S3. In-vitro characterization of SARS-CoV-2 S2GΔHR2 SApNPs.

fig. S4. SARS-CoV-2 RBD/RBD-SApNP vaccine-induced binding antibody responses.

fig. S5. SARS-CoV-2 spike/spike-SApNP vaccine-induced binding antibody responses.

fig. S6. SARS-CoV-1 spike/RBD/RBD-SApNP vaccine-induced binding antibody responses.

fig. S7. SARS-CoV-2 RBD/RBD-SApNP vaccine-induced neutralizing antibody responses.

fig. S8. SARS-CoV-2 spike/spike-SApNP vaccine-induced neutralizing antibody responses.

fig. S9. SARS-CoV-1 spike/RBD/RBD-SApNP vaccine-induced neutralizing antibody

responses.

table. S1. Key *in vitro* and *in vivo* characteristics of RBD and spike-based SARS-CoV-2

vaccines.

677   References

1. N. Zhu *et al.*, A novel coronavirus from patients with pneumonia in China, 2019. *N. Engl.*
J. Med. **382**, 727-733 (2020).

2. P. Zhou *et al.*, A pneumonia outbreak associated with a new coronavirus of probable bat
origin. *Nature* **579**, 270-273 (2020).

3. F. Wu *et al.*, A new coronavirus associated with human respiratory disease in China.
Nature **579**, 265-226 (2020).

4. E. S. Dong, H. R. Du, L. Gardner, An interactive web-based dashboard to track COVID-
19 in real time. *Lancet Infect. Dis.* **20**, 533-534 (2020).

5. M. Hoffmann *et al.*, SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is
blocked by a clinically proven protease inhibitor. *Cell* **181**, 271-280.e278 (2020).

6. M. Z. Tay, C. M. Poh, L. Renia, P. A. MacAry, L. F. P. Ng, The trinity of COVID-19:
immunity, inflammation and intervention. *Nat. Rev. Immunol.* **20**, 363-374 (2020).

7. T. Zohar, G. Alter, Dissecting antibody-mediated protection against SARS-CoV-2. *Nat.*
Rev. Immunol. **20**, 392-394 (2020).

8. N. Vabret *et al.*, Immunology of COVID-19: Current state of the science. *Immunity* **52**,
910-941 (2020).

9. N. Lurie, M. Saville, R. Hatchett, J. Halton, Developing Covid-19 vaccines at pandemic
speed. *N. Engl. J. Med.* **382**, 1969-1973 (2020).

10. B. S. Graham, Rapid COVID-19 vaccine development. *Science* **368**, 945-946 (2020).

11. F. Krammer, SARS-CoV-2 vaccines in development. *Nature* **586**, 516-527 (2020).

12. T. T. Le, J. P. Cramer, R. Chen, S. Mayhew, Evolution of the COVID-19 vaccine
development landscape. *Nat. Rev. Drug Discov.* **19**, 667-668 (2020).

13. Q. Gao *et al.*, Development of an inactivated vaccine candidate for SARS-CoV-2.
Science **369**, 77-81 (2020).

14. H. Wang *et al.*, Development of an inactivated vaccine candidate, BBIBP-CorV, with
potent protection against SARS-CoV-2. *Cell* **182**, 713-721.e719 (2020).

15. N. van Doremalen *et al.*, ChAdOx1 nCoV-19 vaccine prevents SARS-CoV-2 pneumonia
in rhesus macaques. *Nature* **586**, 578-582 (2020).

16. P. M. Folegatti *et al.*, Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine
against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised
controlled trial. *Lancet* **396**, 467-478 (2020).

17. N. B. Mercado *et al.*, Single-shot Ad26 vaccine protects against SARS-CoV-2 in rhesus
macaques. *Nature* **586**, 583-588 (2020).

18. F.-C. Zhu *et al.*, Safety, tolerability, and immunogenicity of a recombinant adenovirus
type-5 vectored COVID-19 vaccine: a dose-escalation, open-label, non-randomised, first-
in-human trial. *Lancet* **395**, 1845-1854 (2020).

19. T. R. F. Smith *et al.*, Immunogenicity of a DNA vaccine candidate for COVID-19. *Nat.*
Commun. **11**, 2601 (2020).

20. J. Yu *et al.*, DNA vaccine protection against SARS-CoV-2 in rhesus macaques. *Science*
369, 806-811 (2020).

21. A. Patel *et al.*, Intradermal-delivered DNA vaccine provides anamnestic protection in a
rhesus macaque SARS-CoV-2 challenge model. *bioRxiv*, 2020.2007.2028.225649 (2020).

22. L. A. Jackson *et al.*, An mRNA vaccine against SARS-CoV-2 - preliminary report. *N.*
Engl. J. Med. **383**, 1920-1931 (2020).

23. E. E. Walsh *et al.*, Safety and immunogenicity of two RNA-based Covid-19 vaccine
candidates. *N. Engl. J. Med.* **383**, 2439-2450 (2020).
24. J.-H. Tian *et al.*, SARS-CoV-2 spike glycoprotein vaccine candidate NVX-CoV2373
elicits immunogenicity in baboons and protection in mice. *bioRxiv*,
2020.2006.2029.178509 (2020).
25. C. Keech *et al.*, Phase 1-2 trial of a SARS-CoV-2 recombinant spike protein nanoparticle
vaccine. *N. Engl. J. Med.* **383**, 2320-2332 (2020).
26. M. Voysey *et al.*, Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222)
against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil,
South Africa, and the UK. *Lancet*, Online ahead of print (2020).
27. F. P. Polack *et al.*, Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N.*
Engl. J. Med., Online ahead of print (2020).
28. L. R. Baden *et al.*, Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. *N.*
Engl. J. Med., Online ahead of print (2020).
29. J. Shang *et al.*, Cell entry mechanisms of SARS-CoV-2. *Proc. Natl. Acad. Sci. U. S. A.*
117, 11727-11734 (2020).
30. L. Y. Du *et al.*, The spike protein of SARS-CoV - a target for vaccine and therapeutic
development. *Nat. Rev. Microbiol.* **7**, 226-236 (2009).
31. K. Duan *et al.*, Effectiveness of convalescent plasma therapy in severe COVID-19
patients. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9490-9496 (2020).
32. S. B. Jiang, C. Hillyer, L. Y. Du, Neutralizing antibodies against SARS-CoV-2 and other
human coronaviruses. *Trends Immunol.* **41**, 355-359 (2020).
33. D. Pinto *et al.*, Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
antibody. *Nature* **583**, 290-295.
34. X. Tian *et al.*, Potent binding of 2019 novel coronavirus spike protein by a SARS
coronavirus-specific human monoclonal antibody. *Emerg. Microbes Infect.* **9**, 382-385
(2020).
35. S. J. Zost *et al.*, Rapid isolation and profiling of a diverse panel of human monoclonal
antibodies targeting the SARS-CoV-2 spike protein. *Nat. Med.* **26**, 1422-1427.
36. B. Ju *et al.*, Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature*
584, 115-119 (2020).
37. R. Shi *et al.*, A human neutralizing antibody targets the receptor-binding site of SARS-
CoV-2. *Nature* **584**, 120-124 (2020).
38. Y. Wu *et al.*, A noncompeting pair of human neutralizing antibodies block COVID-19
virus binding to its receptor ACE2. *Science* **368**, 1274-1278 (2020).
39. Y. L. Cao *et al.*, Potent neutralizing antibodies against SARS-CoV-2 identified by high-
throughput single-cell sequencing of convalescent patients' B cells. *Cell* **182**, 73-84.e16
(2020).
40. D. F. Robbiani *et al.*, Convergent antibody responses to SARS-CoV-2 in convalescent
individuals. *Nature* **584**, 437-442 (2020).
41. T. F. Rogers *et al.*, Isolation of potent SARS-CoV-2 neutralizing antibodies and
protection from disease in a small animal model. *Science* **369**, 956-963 (2020).
42. J. D. Huo *et al.*, Neutralizing nanobodies bind SARS-CoV-2 spike RBD and block
interaction with ACE2. *Nat. Struct. Mol. Biol.* **27**, 846-854.
43. D. Wrapp *et al.*, Structural basis for potent neutralization of betacoronaviruses by single-
domain camelid antibodies. *Cell* **181**, 1004-1015.e1015 (2020).

44.   D. Wrapp *et al.*, Cryo-EM structure of the 2019-nCoV spike in the prefusion
770   conformation. *Science* **367**, 1260-1263 (2020).

771   45.   A. C. Walls *et al.*, Structure, function, and antigenicity of the SARS-CoV-2 spike
772   glycoprotein. *Cell* **181**, 281-292.e286 (2020).

773   46.   R. Yan *et al.*, Structural basis for the recognition of SARS-CoV-2 by full-length human
774   ACE2. *Science* **367**, 1444-1448 (2020).

775   47.   J. Shang *et al.*, Structural basis of receptor recognition by SARS-CoV-2. *Nature* **581**,
776   221-224 (2020).

777   48.   T. Zhou *et al.*, Cryo-EM structures of SARS-CoV-2 spike without and with ACE2 reveal
778   a pH-dependent switch to mediate endosomal positioning of receptor-binding domains.
779   *Cell Host Microbe* **28**, 867-879.e865 (2020).

780   49.   C. O. Barnes *et al.*, Structures of human antibodies bound to SARS-CoV-2 spike reveal
781   common epitopes and recurrent features of antibodies. *Cell* **82**, 828-842.e816 (2020).

782   50.   M. Yuan *et al.*, A highly conserved cryptic epitope in the receptor binding domains of
783   SARS-CoV-2 and SARS-CoV. *Science* **368**, 630-633 (2020).

784   51.   M. Yuan *et al.*, Structural basis of a shared antibody response to SARS-CoV-2. *Science*
785   **369**, 1119-1123 (2020).

786   52.   Y. Cai *et al.*, Distinct conformational states of SARS-CoV-2 spike protein. *Science* **369**,
787   1586-1592 (2020).

788   53.   C.-L. Hsieh *et al.*, Structure-based design of prefusion-stabilized SARS-CoV-2 spikes.
789   *Science* **369**, 1501-1505 (2020).

790   54.   H. Yao *et al.*, Molecular architecture of the SARS-CoV-2 virus. *Cell* **183**, 730-738.e713
791   (2020).

792   55.   Z. Ke *et al.*, Structures and distributions of SARS-CoV-2 spike proteins on intact virions.
793   *Nature* **588**, 498-502 (2020).

794   56.   B. Turonova *et al.*, In situ structural analysis of SARS-CoV-2 spike reveals flexibility
795   mediated by three hinges. *Science* **370**, 203-208 (2020).

796   57.   L. L. He *et al.*, Proof of concept for rational design of hepatitis C virus E2 core
797   nanoparticle vaccines. *Sci. Adv.* **6**, eaaz6225 (2020).

798   58.   L. He *et al.*, HIV-1 vaccine design through minimizing envelope metastability. *Sci. Adv.*
799   **4**, aau6769 (2018).

800   59.   L. He *et al.*, Presenting native-like trimeric HIV-1 antigens with self-assembling
801   nanoparticles. *Nat. Commun.* **7**, 12041 (2016).

802   60.   L. He *et al.*, Single-component multilayered self-assembling nanoparticles presenting
803   rationally designed glycoprotein trimers as Ebola virus vaccines. *bioRxiv*,
804   2020.2008.2022.262634 (2020).

805   61.   B. Zakeri *et al.*, Peptide tag forming a rapid covalent bond to a protein, through
806   engineering a bacterial adhesin. *Proc. Natl. Acad. Sci. U. S. A.* **109**, E690-E697 (2012).

807   62.   J. Lan *et al.*, Structure of the SARS-CoV-2 spike receptor-binding domain bound to the
808   ACE2 receptor. *Nature* **581**, 215-220 (2020).

809   63.   R. N. Kirchdoerfer *et al.*, Stabilized coronavirus spikes are resistant to conformational
810   changes induced by receptor recognition or proteolysis. *Sci. Rep.* **8**, 15701 (2018).

811   64.   Y. Yuan *et al.*, Cryo-EM structures of MERS-CoV and SARS-CoV spike glycoproteins
812   reveal the dynamic receptor binding domains. *Nat. Commun.* **8**, 15092 (2017).

65.   R. W. Sanders *et al.*, A next-generation cleaved, soluble HIV-1 Env trimer, BG505
SOSIP.664 gp140, expresses multiple epitopes for broadly neutralizing but not non-
neutralizing antibodies. *PLoS Pathog.* **9**, e1003618 (2013).

66.   P. Pugach *et al.*, A native-like SOSIP.664 trimer based on an HIV-1 subtype B env gene.
J. Virol. **89**, 3380-3395 (2015).

67.   J. ter Meulen *et al.*, Human monoclonal antibody combination against SARS coronavirus:
Synergy and coverage of escape mutants. *PLoS Med.* **3**, 1071-1079 (2006).

68.   P. Prabakaran *et al.*, Structure of severe acute respiratory syndrome coronavirus receptor-
binding domain complexed with neutralizing antibody. *J. Biol. Chem.* **281**, 15829-15836
(2006).

69.   W. C. Hwang *et al.*, Structural basis of neutralization by a human anti-severe acute
respiratory syndrome spike protein antibody, 80R. *J. Biol. Chem.* **281**, 34610-34616
(2006).

70.   A. C. Walls *et al.*, Unexpected receptor functional mimicry elucidates activation of
coronavirus fusion. *Cell* **176**, 1026-1039.e1015 (2019).

71.   T. U. J. Bruun, A. M. C. Andersson, S. J. Draper, M. Howarth, Engineering a rugged
nanoscaffold to enhance plug-and-display vaccination. *Acs Nano* **12**, 8855-8866 (2018).

72.   L. Chu, D. K. Robinson, Industrial choices for protein production by large-scale cell
culture. *Curr. Opin. Biotechnol.* **12**, 180-187 (2001).

73.   Y. Durocher, M. Butler, Expression systems for therapeutic glycoprotein production.
Curr. Opin. Biotechnol. **20**, 700-707 (2009).

74.   J. Alexander *et al.*, Development of high potency universal DR-restricted helper epitopes
by modification of high-affinity DR-blocking peptides. *Immunity* **1**, 751-761 (1994).

75.   N. M. A. Okba *et al.*, Particulate multivalent presentation of the receptor binding domain
induces protective immune responses against MERS-CoV. *Emerg. Microbes Infect.* **9**,
1080-1091 (2020).

76.   X. Ma *et al.*, Nanoparticle vaccines based on the receptor binding domain (RBD) and
heptad repeat (HR) of SARS-CoV-2 elicit robust protective immune responses. *Immunity*
53, 1315-1330.e1319 (2020).

77.   A. C. Walls *et al.*, Elicitation of potent neutralizing antibody responses by designed
protein nanoparticle vaccines for SARS-CoV-2. *Cell* **183**, 1367-1382.e1317 (2020).

78.   J. B. Bale *et al.*, Accurate design of megadalton-scale two-component icosahedral protein
complexes. *Science* **353**, 389-394 (2016).

79.   J. Pallesen *et al.*, Immunogenicity and structures of a rationally designed prefusion
MERS-CoV spike antigen. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E7348-E7357 (2017).

80.   R. Henderson *et al.*, Controlling the SARS-CoV-2 spike glycoprotein conformation. *Nat.*
Struct. Mol. Biol. **27**, 925-933 (2020).

81.   L. Kong *et al.*, Uncleaved prefusion-optimized gp140 trimers derived from analysis of
HIV-1 envelope metastability. *Nat. Commun.* **7**, 12040 (2016).

82.   B. Zhang *et al.*, A platform incorporating trimeric antigens into self-assembling
nanoparticles reveals SARS-CoV-2-spike nanoparticles to elicit substantially higher
neutralizing responses than spike alone. *Sci. Rep.* **10**, 18149 (2020).

83.   P. J. Hotez, D. B. Corry, U. Strych, M. E. Bottazzi, COVID-19 vaccines: Neutralizing
antibodies and the alum advantage. *Nat. Rev. Immunol.* **20**, 399-400 (2020).

84.   K. H. D. Crawford *et al.*, Protocol and reagents for pseudotyping lentiviral particles with
SARS-CoV-2 spike protein for neutralization assays. *Viruses* **12**, 513 (2020).

85. E. S. Rosenberg *et al.*, Vigorous HIV-1-specific CD4(+) T cell responses associated with
control of viremia. *Science* **278**, 1447-1450 (1997).
86. L. M. Snell *et al.*, Overcoming CD4 Th1 cell fate restrictions to sustain antiviral CD8 T
cells and control persistent virus infection. *Cell Rep.* **16**, 3286-3296 (2016).
87. J. F. Zhu, H. Yamane, W. E. Paul, Differentiation of effector CD4 T cell populations.
Annu. Rev. Immunol. **28**, 445-489 (2010).
88. B. J. Laidlaw, J. E. Craft, S. M. Kaech, The multifaceted role of CD4(+) T cells in
CD8(+) T cell memory. *Nat. Rev. Immunol.* **16**, 102-111 (2016).
89. A. Grifoni *et al.*, Targets of T cell responses to SARS-CoV-2 coronavirus in humans with
COVID-19 disease and unexposed individuals. *Cell* **181**, 1489-1501.e1415 (2020).
90. M. Jeyanathan *et al.*, Immunological considerations for COVID-19 vaccine strategies
Nat. Rev. Immunol. **20**, 615-632 (2020).
91. Y. Peng *et al.*, Broad and strong memory CD4+ and CD8+ T cells induced by SARS-
CoV-2 in UK convalescent COVID-19 patients. *Nat. Immunol.* **21**, 1336-1345 (2020).
92. C. M. Krawczyk, H. Shen, E. J. Pearce, Functional plasticity in memory T helper cell
responses. *J. Immunol.* **178**, 4080-4088 (2007).
93. J. K. Taubenberger, D. M. Morens, 1918 influenza: the mother of all pandemics. *Emerg.*
Infect. Dis. **12**, 15-22 (2006).
94. K. P. O'Callaghan, A. M. Blatz, P. A. Offit, Developing a SARS-CoV-2 vaccine at warp
speed. *J. Am. Med. Assoc.* **324**, 437-438 (2020).
95. G. A. Poland, Tortoises, hares, and vaccines: A cautionary note for SARS-CoV-2 vaccine
development. *Vaccine* **38**, 4219-4220 (2020).
96. L. Corey, J. R. Mascola, A. S. Fauci, F. S. Collins, A strategic approach to COVID-19
vaccine R&D. *Science* **368**, 948-950 (2020).
97. M. S. Diamond, T. C. Pierson, The challenges of vaccine development against a new
virus during a pandemic. *Cell Host Microbe* **27**, 699-703 (2020).
98. G. Alter *et al.*, Whither COVID-19 vaccines? *Nat. Biotechnol.* **38**, 1132-1145 (2020).
99. J. P. Moore, P. J. Klasse, COVID-19 vaccines: "warp speed" needs mind melds, not
warped minds. *J. Virol.* **94**, e01083-01020 (2020).
100. K. Mochizuki *et al.*, Programming of donor T cells using allogeneic delta-like ligand 4-
positive dendritic cells to reduce GVHD in mice. *Blood* **127**, 3270-3280 (2016).
890

Acknowledgements: Funding: This work was funded by NIH Grants AI129698 and AI140844

(to J.Z.), Ufovax/SFP-2018-0416, Ufovax/SFP-2018-1013 and Ufovax/SFP-2020-0111 (to J.Z.).

Author contributions: Project design by L.H., Y.Z., I.A.W. and J.Z; construct design of RBD

and spike-based immungoens by L.H. and J.Z.; plasmid design and processing by L.H. and C.S.;

antigen production, purification and basic characterization by L.H., X.L., and T.N.; antibody

production and column packing by L.H., X.L., and T.N.; antibody-antigen ELISA and BLI by

L.H. and C.S.; DSC by L.H. and J.Z.; negative-stain EM by L.H. and J.Z.; plasma-antigen

ELISA by L.H., X.L., and C.S.; antibody and mouse plasma neutralization by L.H. and X.L.;

vaccine-induced T-cell response analysis by Y.W., C.A., and Y.Z.; Manuscript written by L.H.,

Y.Z., I.A.W. and J.Z. All authors were asked to comment on the manuscript. The TSRI

manuscript number is 30028. **Competing interests:** Authors declare no competing interests.

Data and materials availability: All data are available in the main text or in the supplementary

materials. Additional data related to this paper may be requested from the corresponding author.

Figure legends

Fig. 1. Rational design of SARS-CoV-1/2 RBD-based vaccines. (A) Structural model of RBD-

5GS-1TD0 in an extended RBD-up conformation (top view and side view). 1TD0 is a

trimerization scaffold of viral origin. **(B)** SEC profiles of SARS-CoV-1/2 scaffolded RBD

trimers following ExpiCHO expression and CR3022 purification. **(C)** ELISA binding of SARS-

CoV-1/2 scaffolded RBD trimers to a panel of mAbs. EC_{50} (μg/ml) values are labeled. **(D)** Octet

binding of the SARS-CoV-2 scaffolded RBD trimer to five mAbs. Sensorgrams were obtained

from an Octet RED96 instrument at six antigen concentrations from 950 to 29.5 nM by twofold

dilutions. **(E)** Diagram of conjugating RBD to the 24-meric FR SApNP using the

SpyTag/SpyCatcher (SPY) system. **(F)** SEC profiles of SARS-CoV-1/2 RBD-5GS-SPY-5GS-FR

SApNPs produced in ExpiCHO by co-expression (black line) and supernatant mix (red line). **(G)**

ELISA binding of SARS-CoV-1/2 RBD-FR SApNPs to a panel of mAbs. EC_{50} (μg/ml) values

are labeled. **(H)** Octet binding of the SARS-CoV-2 RBD-FR SApNP to five mAbs. Sensorgrams

were obtained from an Octet RED96 instrument at six antigen concentrations from 37 to 1.1 nM

by twofold dilutions. **(I)** EM images of SARS-CoV-1 RBD-10GS-FR (gene fusion) and RBD-

5GS-SPY-5GS-FR (co-expression). **(J)** EM images of SARS-CoV-2 RBD-5GS-SPY-5GS-FR

produced by co-expression and supernatant mix. **(K)** Diagram of conjugating RBD to the 60-

meric multilayered I3-01v9 SApNP using the SPY system. Locking domains (LD) and T-helper

epitopes within the multilayered SApNP are depicted. **(L)** and **(M)** SEC profiles and EM images

of   SARS-CoV-1/2   RBD-5GS-SPY-5GS-I3-01v9-LD7-PADRE   (or   I3-01v9-L7P)   SApNPs

produced by supernatant mix. SEC profiles of scaffolded RBD trimers in (B) and SPY-linked

RBD SApNPs in (F) and (L) were obtained from a Superdex 200 10/300 GL column and a

Superose 6 10/300 GL column, respectively.

Fig. 2. Rational design of SARS-CoV-2 spike antigens. (A) Structural model of prefusion S

spike linked to the C-terminal trimerization domain (1TD0) with a 5GS linker in transparent

molecular surface. The approximate position for the unstructured HR2 stalk, or in this case a 5-

aa G_4S linker, is highlighted with a dashed line box. **(B)** SEC profiles of SARS-CoV-1 (top) and

SARS-CoV-2 (bottom) spikes. Left to right panels: $S2P_{ECTO}$-5GS-1TD0 purified on Nickel and

CR3022 columns (1 and 2), $S2G_{ECTO}$-5GS-1TD0 on Nickel and CR3022 columns (3 and 4), and

S2GΔHR2-5GS-1TD0 on a CR3022 column (5). All spike constructs tested in (B) contain a C-

terminal His_6 tag to facilitate Nickel column purification. Results from three separate production

runs are shown for SARS-CoV-2 $S2G_{ECTO}$-5GS-1TD0. **(C)** Schematic representation of a full-

length SARS-CoV-2 spike on the virus membrane surface in the presence of host ACE2 and the

HR2 region from a neighboring spike (left), and sequence alignment of SARS-CoV-1/2 HR1 and

HR2 regions (right, top and bottom). The HR1 and HR2 segments that form a six-helix bundle in

the post-fusion state are highlighted in green and brown shade, respectively. **(D)** Left: SEC

profiles of S2GΔHR2-5GS-1TD0, showing results from four separate production runs. Right:

BN-PAGE analysis of $S2P_{ECTO}$-5GS-1TD0 and S2GΔHR2-5GS-1TD0. SEC fractions (12.5-

14.0) are shown for S2GΔHR2-5GS-1TD0 on the gel. SEC profiles in (B) and (D) were obtained

from a Superose 6 10/300 GL column. **(E)** ELISA binding of two SARS-CoV-2 spikes

(S2P$_{ECTO}$-5GS-1TD0 and S2GΔHR2-5GS-1TD0) to five mAbs. EC$_{50}$ (µg/ml) values are labeled.

(F) Octet binding of two SARS-CoV-2 spikes (S2P$_{ECTO}$-5GS-1TD0 and S2GΔHR2-5GS-1TD0)

to five mAbs. Sensorgrams were obtained from an Octet RED96 instrument at six antigen

concentrations from 150 to 4.7 nM by twofold dilutions. The spike constructs tested in (D)-(F)

do not contain a C-terminal His$_6$ tag.

Fig. 3. Rational design of SARS-CoV-2 spike-presenting nanoparticle vaccines. (A)

Schematic representation of SARS-CoV-2 virion (top) and spike-presenting SApNP vaccine

(bottom). For the SARS-CoV-2 virion, pre/post-fusion S, nucleocapsid, and RNA viral genome

are shown, while for the vaccine, stabilized spike and multilayered SApNP composition are

depicted. **(B)** Colored surface models of SApNP carriers (top) and spike-presenting SApNP

vaccines (bottom). The three SApNP carriers used here are ferritin (FR) 24-mer and multilayered

E2p and I3-01v9 60-mers (the LD layer and PADRE cluster are not shown). Particle size is

indicated by diameter (in nanometers). **(C)** SEC profiles of three SApNPs presenting the SARS-

CoV-2 S2GΔHR2 spike, from left to right, S2GΔHR2-5GS-FR, S2GΔHR2-5GS-E2p-LD4-

PADRE (or E2p-L4P), and S2GΔHR2-10GS-I3-01v9-LD7-PADRE (or I3-01v9-L7P). SEC

profiles were obtained from a Superose 6 10/300 GL column, each showing results from three

separate production runs. **(D)** EM images of three SARS-CoV-2 S2GΔHR2 SApNPs. **(E)** ELISA

binding of three SARS-CoV-2 S2GΔHR2 SApNPs to five mAbs. EC$_{50}$ (µg/ml) values are

labeled. **(F)** Antigenic profiles of SARS-CoV-2 S2GΔHR2 spike and three SApNPs against five

mAbs. Sensorgrams were obtained from an Octet RED96 using six antigen concentrations (150-

4.6 nM for spike, 9-0.27 nM for FR SApNP, and 3.5-0.1 nM for E2p and I3-01v9 SApNPs,

respectively, all by twofold dilutions) and AHQ biosensors, as shown in fig. S3B. The peak

binding signals (nm) at the highest concentration are listed. Color coding indicates the signal

strength measured by Octet (green to red: low to high).

Fig. 4. SARS-CoV-1/2 vaccine-induced binding antibody (bAb) responses in BALB/c mice.

(A) Schematic representation of the mouse immunization protocol. **(B)** SARS-CoV-2 RBD and

RBD-SApNP vaccine-induced bAb responses. Coating antigens: SARS-CoV-2 S2GΔHR2-5GS-

foldon (left) and RBD (right). **(C)** SARS-CoV-2 S2GΔHR2 spike and S2GΔHR2-SApNP

vaccine-induced bAb responses. Coating antigens: SARS-CoV-2 S2GΔHR2-5GS-foldon (top)

and RBD (bottom). In (B) and (C), the SARS-CoV-2 S2P$_{ECTO}$-5GS-1TD0 group is included for

comparison, with S2P$_{ECTO}$-5GS-foldon used as the coating antigen. **(D)** SARS-CoV-1 RBD and

RBD-SApNP vaccine-induced bAb responses. Coating antigens: SARS-CoV-1 S2P$_{ECTO}$-5GS-

foldon (left) and RBD (right). In (D), the SARS-CoV-1 S2P$_{ECTO}$-5GS-1TD0 group is included

for comparison, with S2P$_{ECTO}$-5GS-foldon used as the coating antigen. In (B) to (D), EC$_{50}$ titers

derived from the ELISA binding of mouse plasma to coating antigens are plotted, with average

EC$_{50}$ values labeled on the plots. The *P* values were determined by an unpaired *t* test in

GraphPad Prism 8.4.3 with (*) indicating the level of statistical significance. Detailed ELISA

data are shown in figs. S4-S6.

Fig. 5. SARS-CoV-1/2 vaccine-induced neutralizing antibody (NAb) responses in BALB/c

mice. (A) SARS-CoV-2 RBD and RBD-SApNP vaccine-induced NAb responses to autologous

SARS-CoV-2 (left) and heterologous SARS-CoV-1 (right). **(B)** SARS-CoV-2 S2GΔHR2 spike

and S2GΔHR2-SApNP vaccine-induced NAb responses to SARS-CoV-2 (top) and SARS-CoV-

1 (bottom). In (A) and (B), the SARS-CoV-2 S2P$_{ECTO}$-5GS-1TD0 group is included for

comparison. **(C)** SARS-CoV-1 RBD and RBD-SApNP vaccine-induced NAb responses to

autologous SARS-CoV-1 (left) and heterologous SARS-CoV-2 (right). In (C), the SARS-CoV-1

S2P$_{ECTO}$-5GS-1TD0 group is included for comparison. In (A) to (C), ID$_{50}$ titers derived from the

SARS-CoV-1/2-pp neutralization assays are plotted, with average ID$_{50}$ values labeled on the

plots. The *P* values were determined by an unpaired *t* test in GraphPad Prism 8.4.3 with (*)

indicating the level of statistical significance. Detailed SARS-CoV-1/2-pp neutralization data are

shown in figs. S7-S9.

Fig. 6. SARS-CoV-2 vaccine-induced T-cell responses in BALB/c mice. Splenocytes from

mice (n=5 in each group) immunized with S2P$_{ECTO}$ spike, S2GΔHR2 E2p SApNP, and

S2GΔHR2 I3-01v9 SApNP, were isolated at w11 and cultured in the presence of IL-2 and

dendritic cells (DC) pulsed with S2P$_{ECTO}$ (1×10^{-7} μM), S2GΔHR2 E2p (1×10^{-7} μM), and

S2GΔHR2 I3-01v9 (1×10^{-7} μM), respectively. Splenocytes from 5 naïve mice were used as

control samples and cultured in the presence of DCs without antigen pulsing. Cells were assessed

after 16 hours (A, C) and 4 hours (B, D) of culture. **(A)** and **(B)**: Vaccine-induced CD4^{+} T cell

immunity. **(C)** and **(D)**: Vaccine-induced CD8^{+} T cell immunity. Plots show the frequencies of

cell fraction. The *P* values were determined by one-way ANOVA analysis. *, $P < 0.05$; **, $P <$

0.01; ***, $P < 0.001$.

A

Postfusion S2
Prefusion spike
Nucleocapsid
RNA viral genome

SARS-CoV-2 virion

Stabilized prefusion spike
SApNP unit
Locking domain (LD)
T-helper cluster

Vaccine

B

Ferritin (FR)

12.2 nm
23.2 nm

47.9nm

E2p

55.9nm

I3-01

25.0 nm

59.3nm

C

S2GΔHR2-5GS-FR
(400 ml ExpiCHO)

S2GΔHR2-5GS- E2p-LD4-PADRE(400 ml ExpiCHO)

S2GΔHR2-10GS-I3-01v9-LD7-PADRE (400 ml ExpiCHO)

#1 #2 #3

D

S2GΔHR2-5GS-FR

S2GΔHR2-5GS-E2p-LD4-PADRE

S2GΔHR2-10GS-I3-01v9-LD7-PADRE

50nm 50nm 50nm

E

S2GΔHR2-5GS-FR

1.4E-2
6.5E-2
1.8E-3
3.2E-3
1.9E-3

S2GΔHR2-5GS-E2p-LD4-PADRE

9.4E-3
4.4E-2
1.4E-3
1.8E-3
1.4E-3

S2GΔHR2-10GS-I3-01v9-LD7-PADRE

7.2E-3
2.1E-2
1.2E-3
1.5E-3
1.4E-3

CR3022
B38
CB6
S309
P2B-2F6

F

SARS-CoV-2 S constructs	CR3022	B38	CB6	S309	P2B-2F6
S2GΔHR2-5GS-1TD0	1.17	1.07	0.83	0.94	0.77
S2GΔHR2-5GS-FR	1.12	1.41	1.27	1.06	1.68
S2GΔHR2-5GS-E2p-LD4-PADRE	1.92	2.27	1.82	1.67	1.98
S2GΔHR2-10GS-I3-01v9-LD7-PADRE	1.93	1.99	1.93	1.73	1.92

A

B SARS-CoV-2 RBD-based vaccine-induced binding antibody responses

C SARS-CoV-2 spike-based vaccine-induced binding antibody responses

D SARS-CoV-1 vaccine-induced binding antibody responses

A Induction of Th1 and memory responses

Gated on CD4$^+$ TCR-β$^+$ T cells

Naïve | S2P$_{ECTO}$ | E2p | I3-01v9

IFN-γ / IL-4 (16 hours)

IFN-γ$^+$ (Th1)
** 0.0097
*** 0.0005
* 0.0379
Percent cells (in CD4 T cells)

IFN-γ$^+$IL-4$^+$ (Memory T cell)
* 0.0187
* 0.0271

B Induction of cytolytic CD4$^+$ T cells

Gated on CD4$^+$ TCR-β$^+$ T cells

Naïve | S2P$_{ECTO}$ | E2p | I3-01v9

IFN-γ / CD107a (4 hours)

CD107a$^+$ Cytolytic CD4$^+$ T cells
** 0.0042
*** 0.00024
Percent cells (in CD4 T cells)

IFN-γ$^+$ (Th1)

C Induction of recall response of CD8$^+$ T cells

Gated on CD8$^+$ TCR-β$^+$ T cells

Naïve | S2P$_{ECTO}$ | E2p | I3-01v9

IFN-γ / GM-CSF (16 hours)

IFN-γ$^+$GM-CSF$^+$
* 0.0456
* 0.0437
* 0.0414
Percent cells (in CD8 T cells)

IFN-γ$^-$GM-CSF$^+$
* 0.0132
* 0.0157
** 0.0086

D Induction of rapid CD8$^+$ T cell recall response

Gated on CD8$^+$TCR-β$^+$ T cells

Naïve | S2P$_{ECTO}$ | E2p | I3-01v9

IFN-γ / CD107a (4 hours)

IFN-γ$^+$ CD8
* 0.0247
Percent cells (in CD8 T cells)

IFN-γ$^+$ CD107a$^+$