Dynamic gene expression and growth underlie cell-to-cell heterogeneity in *Escherichia coli* stress response

Nadia M. V. Sampaio[1,2,†,‡], Caroline M. Blassick[1,2,†], Jean-Baptiste Lugagne[1,2], Mary J. Dunlop[1,2,*]

[1] Biomedical Engineering Department, Boston University, Boston, MA 02215.
[2] Biological Design Center, Boston University, Boston, MA 02215.
[†] These authors contributed equally to this work.
[‡] Present Address: Brazilian Biorenewables National Laboratory, Brazilian Center for Research in Energy and Materials, Campinas, São Paulo, Brazil.
* Corresponding Author: mjdunlop@bu.edu

Abstract

Cell-to-cell heterogeneity in gene expression and growth can have critical functional consequences, such as determining whether individual bacteria survive or die following stress. Although phenotypic variability is well documented, the dynamics that underlie it are often unknown. This information is important because dramatically different outcomes can arise from gradual versus rapid changes in expression and growth. Using single-cell time-lapse microscopy, we measured the temporal expression of a suite of stress response reporters in *Escherichia coli*, while simultaneously monitoring growth rate. In conditions without stress, we found several examples of pulsatile expression. Single-cell growth rates were often anti-correlated with gene expression, with changes in growth preceding changes in expression. These expression and growth dynamics have functional consequences, which we demonstrate by measuring survival after challenging cells with the antibiotic ciprofloxacin. Our results suggest that fluctuations in gene expression and growth dynamics in stress response networks can have direct consequences for survival.

MAIN TEXT

Introduction

Even under otherwise constant environmental conditions, genetically identical cells can display substantial phenotypic heterogeneity, resulting from differences in gene expression and growth rate from cell to cell (*1–4*). Phenotypic heterogeneity can, in principle, arise from slow changes within the population where individual cells have different expression levels or growth rates, but maintain their state over many cell cycles. Alternatively, fast dynamics with rapid fluctuations could produce equivalent distributions. The distinction between these alternatives is significant because the timescale over which expression or growth differences persist can ultimately determine if they have functional consequences or are simply short-lived random variations that are filtered before impacting cellular outcomes. Although phenotypic heterogeneity is well documented, the timescales underlying the variation and their ultimate impact on function are often less clear.

We focused on stress response genes in *Escherichia coli* to study the dynamics of expression and growth in single cells. Genes involved in adaptation to stress are among the noisiest genome-wide

(*3*, *5*). Studies on individual pathways have revealed specific examples in which heterogeneous
expression of stress response genes can allow subpopulations of cells to survive sudden
environmental stress, such as transient exposure to antimicrobial drugs, oxidative stress, and acid
stress (*6–9*). Recent studies also indicate that heterogeneity in the expression of genes involved in
DNA repair can lead to variability in mutation rates, contributing to microbial evolution (*10–12*).
Notably, this diversity exists even in the absence of stressors. Although selected studies have
begun to reveal examples of how cell-to-cell phenotypic variation can provide important functional
capabilities to cell populations, examples of direct links are relatively sparse compared to reports
quantifying phenotypic heterogeneity. This motivated our focus on stress response pathways
because the effect of the genes involved can be directly assessed by quantifying outcomes like cell
survival versus death following stress.
58
In addition to measurements at a single time point, long-term monitoring of gene expression has
begun to uncover examples of rich dynamics in key stress response proteins (*13*). For example,
self-cleavage of the regulator LexA produces spontaneous pulses in the SOS response network
(*14*). Further, dynamic activity of transcription factors can propagate to downstream genes, with
direct consequences for stress tolerance. For instance, pulsatile expression of the transcription
factor ComK enables *Bacillus subtilis* cells to enter a transient competent state (*15*, *16*). In *E. coli*,
heterogeneous expression of RpoS, a key regulator of general stress response, originates from
pulses of activation that are inversely correlated with growth, allowing cells to survive oxidative
stress (*8*). Additionally, Kim *et al.* demonstrated that genes in the flagellar synthesis network, a
process with a pivotal role in microbial pathogenicity, are expressed with different pulsing
programs that allow cells to switch between flagellar phenotypes (*17*). Collectively, these studies
demonstrate that the temporal dynamics of gene expression play an important role in the regulation
of stress response networks.
72
Growth rate fluctuations have also been observed in single cells across many bacterial species (*2*,
8, *18–20*). They can arise due to temporal variation in the expression of metabolic enzymes (*2*,
21), the expression of burdensome proteins (*12*), and due to regulatory effects such as feedback
involved in cell size control (*19*). These single-cell differences in growth rate can play a functional
role in stress tolerance. For example, Narula *et al*. demonstrated that the growth rate of *B. subtilis*
under starvation conditions can determine whether individual cells differentiate into spores or
remain vegetative (*22*). Single-cell growth rates also play a protective role, for example correlating
with differential antibiotic susceptibility in mycobacteria (*23*) and *E. coli* (*7*, *24*).
81
Understanding the timescales associated with gene expression dynamics and growth can provide
critical insights into the strategies that cells use to hedge against environmental uncertainty. In this
study, we characterize the prevalence and dynamic properties of cell-to-cell phenotypic variation
in different branches of the stress response network in *E. coli*. Using time-lapse microscopy, we
monitored the activity of key stress response genes, as well as genes involved in biosynthesis and
metabolism, in single *E. coli* cells under precisely controlled, unstressed conditions. Our results
reveal several new examples of genes that exhibit pulses of gene expression. Furthermore,
properties of the dynamics, such as the frequency or amplitude of pulsing, are unique to each gene.
Interestingly, fluctuations in the expression of genes frequently occur following variations in
growth rate. Finally, focusing on the acid stress regulator GadX, we show that coincident
upregulation of *gadX* and reduced growth rate favors tolerance to lethal antibiotic exposure.

Together, this work reveals that non-trivial gene expression dynamics are common, even in
otherwise constant conditions, and that these dynamic patterns of gene expression can have critical
consequences for cell survival.

Results
Single-cell measurements of phenotypic heterogeneity
We began by characterizing heterogeneity in the expression of genes with a diverse range of
functions (Table S1). These include genes involved in acid resistance (*gadW*, *gadX*, *phoP*), multi-
drug resistance (*evgA*, *marA*, *rob*), heat shock (*rpoH*), oxidative stress response (*oxyR*), SOS
response (*dinB*, *recA*, *sulA*), and general stress response (*bolA*), in addition to genes involved in
biosynthesis and metabolism (*araC*, *metJ*, *purA*).

To measure heterogeneity in the expression from each promoter at the single-cell level, we used
strains containing transcriptional reporters where the promoter sequence of interest is fused to the
coding sequence of a fluorescent protein. We grew independent bulk cultures of each strain to
exponential phase and measured heterogeneity across cells in the population using fluorescence
microscopy for single cells on agarose pads. Cell-to-cell differences in gene expression resulted in
different distributions of reporter levels for each gene (Fig. 1A, Fig. S1). Measurements of the
reporters revealed many instances of wide distributions, indicative of a broad range of expression
levels across cells within a population. Distributions tended to skew to the right of the mean (skew
> 0) due to the presence of highly-expressing cells.

However, measures of static distributions do not reveal the underlying dynamics that generate
them, prompting the question: Do these distributions stem from long-lived fixed subpopulations
with slow dynamics or from conditions with fast dynamics where individual cells transition
between different expression levels over time (Fig. 1B)? The significance of this question is
particularly pertinent in the context of stress response networks, where the timescales over which
individual genes are active may have concrete implications for tolerance levels. For instance, if a
transcription factor that activates genes involved in stress response exhibits short pulses in
expression, these pulses might be insufficient to turn on expression of downstream genes, while
more sustained expression could. Further, single-cell growth rates can impact survival, thus the
interplay between expression and growth may be significant for determining tolerance.

Pulsatile gene expression can underlie cell-to-cell heterogeneity
Thus, we next aimed to quantify the temporal dynamics underlying the distributions of gene
expression and growth. We measured expression in cell lineages over many generations using a
'mother machine' microfluidic device (*25*) and time-lapse fluorescence microscopy. In this device,
'mother' cells are trapped at the top of one-ended chambers and maintained indefinitely in
exponential phase through the addition of fresh media, allowing for multi-hour imaging of cell
lineages. We used this to monitor gene expression in tens to hundreds of independent cell lineages
for each reporter for at least 15 hours. To quantify gene expression over time, we used our deep
learning based cell segmentation and tracking algorithm (*26*) to extract single-cell resolution data
from the mother cell and its progeny.

We observed heterogeneity in gene expression for cells growing in these precisely controlled
conditions, which was consistent with measurements acquired from bulk culture snapshots (Fig.

S2). Long-term monitoring of expression revealed a diversity of phenotypes, with some genes that fluctuated at a low level, likely due to variation in copy number in the reporter plasmid and inherent stochasticity in gene expression, while others were highly dynamic in their expression (Fig. 1C-E). We observed that variability within a distribution often arose from pulsatile dynamics, with single cells transitioning smoothly between different expression states. These results join other recent single-cell studies showing transcriptional pulses in expression in the absence of stress (*8*, *14*, *17*). We found that the timescales of these fluctuations were specific to each gene, and the pulses themselves ranged in intensity. For instance, *gadX*, which encodes a transcription factor that regulates ~34 genes in the acid resistance system, exhibits pulses of high amplitude and duration that persist well beyond the cell cycle length, as visible in multi-generation patterns of gene expression (Fig. 1C, Movie S1). In contrast, *recA*, which plays a central role in the processes of homologous recombination and SOS response, showed large amplitude pulses, but with shorter durations (Fig. 1D, Movie S2). Yet others, like *araC*, which encodes a transcription factor that regulates arabinose catabolism and transport, were more muted in their changes, and exhibited only mild, low amplitude fluctuations (Fig. 1E, Movie S3). Surveying 15 reporters, we observed a wide range of temporal gene expression profiles (Fig. 1F-G). In all cases, the mean fluorescence levels were consistent across the duration of the experiments despite the fluctuations in gene expression at the single-cell level (Fig. S3). Importantly, the dynamic activity observed for many reporters occurred under constant growth conditions, where parameters such as pH, temperature, and growth medium do not fluctuate.

As a control, we also included a plasmid-based constitutive reporter to quantify the baseline level of fluctuations expected from variation in plasmid copy number (Fig. 1H). The constitutive reporter exhibits mild changes in expression but does not reach the level of variation we observed in reporters like *gadX* and *recA*. As an additional control, we also chromosomally integrated the *gadX* reporter and confirmed that pulses in gene expression in cells containing the plasmid and chromosomally-integrated versions of the reporters were comparable (Fig. 1I). Low expression levels prevented us from conducting all experiments using chromosomal reporters, however the constitutive reporter serves as a control to establish baseline effects that are plasmid-based rather than specific to the stress response reporters.

Properties of temporal expression vary across the reporters
Next, we defined several metrics across which to assess dynamic behavior. We selected properties related to pulses including their frequency, duration, and amplitude (Fig. 2A). Pulse frequencies were broadly distributed, ranging from examples that exceed 0.25 pulses per hour (one pulse every four hours) for *recA* to much less frequent conditions where our peak-finding algorithm rarely identified pulses (Fig. 2B). Pulse durations typically ranged from 0 - 4 hours, and showed instances of precision (e.g. *recA*, *sulA*) in addition to examples with widely variable durations (e.g. *bolA*, *evgA, rpoH*). Pulse amplitudes were predominantly small, with average amplitudes around 75% of the mean, or 0.75x, but there were notable instances where the distributions had long tails such that amplitudes extended well above this for a subset of the pulses. For example, *gadX* and *recA* exhibit pulses that significantly deviate from the mean. To alleviate the potential concern that these relationships were a result of the specific method by which we identified pulses, we repeated the calculations under a range of different peak determination thresholds and found that our results were not sensitive to the precise threshold definition, provided the threshold was set high enough to exclude most fluctuations in the constitutive reporter control (Fig. S4).

184
We next asked what characteristic timescales the gene expression dynamics exhibited by
calculating the autocorrelation of the fluorescence signal. In all cases, we observed monotonically
decreasing autocorrelation curves, ruling out the presence of regular period signals, such as
oscillatory behavior (Fig. S5). We calculated the half life associated with the autocorrelation curve
for each gene (Fig. 2C-D). If the fluorescent reporter levels decrease solely due to dilution resulting
from growth and division, the half life will equal the cell division time. Longer half lives can
indicate the presence of memory, for example due to regulatory networks that causes signals to
persist. We found that half lives were always greater than or equal to the cell division time,
consistent with the use of stable fluorescent reporters (Fig. 2D). Notably, we observed cases where
the average exceeded the cell division time by 2-3-fold, potentially indicative of memory within
the network. We verified that these calculations performed on data from mother cells were
consistent with correlations within the lineage tree. Indeed, tracking fluorescence signals from
mother to daughter to granddaughter cells revealed a strong positive correlation that persisted
across multiple generations (Fig. S6).
199

RpoS levels influence *gadX* pulsing dynamics
One question that our results provoke is why some of the reporters exhibit pulsatile expression.
We examined this in more detail for the case of the *gadX* reporter. RpoS plays a role in *gadX*
regulation and has been shown to exhibit pulsatile dynamics during exponential growth (*8*, *27*, *28*).
Thus, we hypothesized that fluctuations in cellular RpoS levels during exponential growth could
be underlying *gadX* pulsing. We used the same *gadX* reporter and compared expression in wild
type and Δ*rpoS* strains. Deletion of *rpoS* reduced *gadX* expression and fluctuations in *gadX*
expression became significantly less noticeable (Fig. 2E, Fig. S7). The frequency, amplitude, and
duration of *gadX* pulses in the Δ*rpoS* cells were reduced relative to the characteristic profile of
wild type cells (Fig. 2F), indicating a direct impact of RpoS levels on *gadX* pulsing dynamics.
Fluctuations in the availability of the master regulator RpoS may also contribute to the profiles
observed for other downstream genes that RpoS controls. The mechanism leading to pulsing
dynamics of other genes involved the SOS response, like *recA* and *sulA*, has been shown to
originate from fluctuations in the availability of their common negative regulator LexA (*14*). Thus,
fluctuations in master regulators of gene expression might be a common mechanism leading to
heterogeneous expression of downstream genes.
216

Pulsatile expression dynamics and growth
We next asked whether growth rate contributes to the observed variability in expression levels.
While the growth rate of individual *E. coli* cells shows long-term stability during replicative aging
(*25*), it can exhibit noisy temporal behavior (*2*, *8*, *19*). Patange *et al*. demonstrated that growth
rates fluctuate, with pulses of slow growth that last for several generations and are anti-correlated
with pulses of activation of RpoS (*8*). However, it is not clear whether this feature is specific to
RpoS or if it is common to many genes. We asked whether the pulsatile dynamics we observed
coincided with dynamic patterns in growth rate. To test this, we extracted the instantaneous growth
rate from cell lineages over time and compared them to fluorescence data for the different
reporters.
227

In principle, fluorescence levels, which report underlying cellular properties such as those related
229 to metabolism, can precede changes in growth. Alternatively, growth can drive changes in

230 fluorescence since it affects dilution rates; regulatory links between growth rate and fluorescence
231 are also possible (Fig. 3A). In both cases, these relationships could be positive or negative
232 depending on the precise underlying mechanism.
233
To distinguish between these possibilities, we computed temporal cross-correlations between
fluorescence and growth rate, $R_{fluo,gr}(\tau)$ (Fig. 3B). The cross-correlation measures how well the
fluorescence and growth signals are correlated when the growth signal is shifted by a time τ relative
to the fluorescence signal. Cross-correlation curves for our reporters have characteristic shapes,
where signals are uncorrelated at large positive and negative values of the time shift, τ. For
intermediate values of τ we regularly observed valleys in the cross-correlation. We defined $R_{extreme}$
as the point on the cross-correlation with the largest absolute value, indicating the largest
correlation between the fluorescence and growth rate signals. $R_{extreme}$ can be either positive or
negative, depending on whether the signals are correlated or anti-correlated. We also defined
$\tau_{extreme}$ as the time shift associated with $R_{extreme}$, where $\tau_{extreme} < 0$ when changes in fluorescence
precede changes in growth and $\tau_{extreme} > 0$ when growth leads fluorescence.
245
Growth and expression are often anti-correlated with growth leading expression
We found examples where a strong anti-correlation between fluorescence and growth were visible,
such as with the *gadX* reporter, and other cases where the signals were uncorrelated, such as with
the *recA* reporter (Fig. 3C). To quantify this trend across many cell lineages, we calculated cross-
correlations between growth and fluorescence for all reporters (Fig. 3D). In many cases we
observed an anti-correlation with a positive time shift between growth rate and gene expression,
indicating that the pulses of expression from these promoters were preceded by a decrease in
growth rate. In addition, we confirmed that the chromosomally-integrated version of the *gadX*
reporter produces cross-correlation curves that are similar to the plasmid-based version (Fig. S8
A-B). Although the accumulation of fluorescent proteins resulting from reduced dilution due to
cell growth and division could potentially artificially generate this relationship, we observed only
a modest anti-correlation for several of the promoters, a near zero or mildly positive relationship
for the constitutive reporter (Fig. S8 C-D), and no anti-correlation for *recA* and *sulA* (Fig. 3E).
Thus, it is unlikely that these results are due solely to a reduced dilution rate.
260
Changes in growth rate never lagged changes in fluorescence for all reporters we measured ($\tau_{extreme}$
> 0) (Fig. 3F). It is possible that fluorophore maturation times could systematically introduce a lag
between actual promoter activity and read-out of the fluorescent protein. Maturation times for our
reporters are relatively rapid, with 50% of fluorescent proteins maturing within ~6 minutes (*29*),
however reporter maturation times could systematically introduce a small positive shift in $\tau_{extreme}$.
Overall, our measurements indicate that even cells growing exponentially under optimal
conditions undergo episodes of slow growth that are largely followed by the pulses of stress-related
genes, which frequently demonstrate an inverse relationship between growth rate and fluorescence
(Fig 3G).
270
We also measured correlations between fluorescence and cell length for all the reporters (Fig. S9).
These signals were not strongly correlated for most of the reporters. However, we did observe
positive correlations between expression of *recA* and *sulA* and cell length. This result is expected,
because these genes play a key role in the SOS response, which is known to induce cell

filamentation (*30–32*). Even in unstressed conditions, we find that a small subset of cells exist in
a state with high gene expression and corresponding elongated cell morphology.
277
Gene expression and growth dynamics transiently protect cells against stress
A critical question is whether these changes in gene expression and growth have concrete
implications for whether a single cell will survive or die following stress. In other words, are the
dynamics we observed sufficient to provide meaningful phenotypic differences that cause a cell to
tolerate or succumb to stress? Slow growth and induction of stress response genes have previously
been reported as mechanisms that allow survival at the populational level (*33–37*). This prompted
us to investigate whether pulses in gene expression and growth influence the chances of survival
under stress. Further, we asked whether survival was the result of a fortuitous condition of gene
expression or slow growth upon stress introduction, or if a cell's past history was important.
287
For these studies, we focused on expression of *gadX,* as our measurements demonstrate that this
reporter is expressed in large pulses and has a strong anti-correlation with growth. We previously
demonstrated that *gadX* is heterogeneously expressed in the absence of antibiotic stress and that
expression levels correlate with longer survival times under constant carbenicillin exposure (*7*).
However, it was not clear how prior history or pulsing dynamics contribute to this phenotype and
whether cells could recover normal cell division after transient drug treatment. Here, we quantified
how *gadX* fluctuations preceding sudden exposure to a lethal antibiotic dose influence survival.
To do so, we leveraged the mother machine device to rapidly switch input media and exposed
exponentially dividing cells to a short pulse of ciprofloxacin, a fluoroquinolone drug widely used
to treat bacterial infections. We treated cells with 2 µg/ml ciprofloxacin, which corresponds to
100x the minimum inhibitory concentration (MIC), for 35 minutes before switching back to
growth medium without antibiotics. This experimental set up allowed us to monitor the dynamics
of *gadX* expression and growth in single cells prior to antibiotic treatment while also recording the
outcome of each cell lineage.
302
Within the same experiment, we observed instances where single cells were able to survive
ciprofloxacin treatment (Fig. 4A, Movie S4) and cases where treatment killed the cells (Fig. 4B,
Movie S5). In addition to surviving and dying, we also observed a third category of outcomes
where cells filamented. However, it was difficult to accurately assess whether these cells survived
or died, as they were frequently swept out of the mother machine chamber and lost from the field
of view. Thus, we focused our analysis on the surviving and dying cells because we could
accurately determine their outcomes. By tracking gene expression history preceding antibiotic
treatment, we observed pulses in fluorescence from the *gadX* reporter prior to ciprofloxacin
addition in both the surviving and dying cells (Fig. 4C). Cells that exhibited expression pulses in
the past, but had low expression at the time of ciprofloxacin treatment were more likely to die. In
contrast, cells which were in the fortuitous state of having an ongoing *gadX* pulse at the time of
ciprofloxacin addition were more likely to exhibit transient stress tolerance.
315
We also looked at cellular outcome as a function of the growth rate. Consistent with our results
showing anti-correlations between fluorescence and growth for *gadX* (Fig. 3D), we observed
decreased growth rates in cells that survived, which were associated with increases in fluorescence
signals (Fig. 4D). However, reduced growth rate alone does not appear to be sufficient for survival,
as we observed cells that died which had similar growth rates to those that survived. Measurements

of the instantaneous fluorescence and growth rate at t = 0, just prior to antibiotic addition, show
that most surviving cells exhibited both elevated *gadX* expression and reduced growth rate
immediately prior to ciprofloxacin treatment (Fig. 4E).

By looking at gene expression history prior to ciprofloxacin addition, we see clear differences
emerging between the populations of cells that will survive and die ~6 hours prior to treatment
(Fig. 4F). The trend we observe does not reflect a linear increase in fluorescence in individual
cells, but rather results from *gadX* pulsing dynamics. For surviving and dying cohorts, we
quantified how long each cell had been experiencing its final pulse of *gadX* expression at the point
when ciprofloxacin was introduced (Fig. 4G). We found that cells that survived were far more
likely to be in the midst of a *gadX* expression pulse than those that died. Pulse duration did not
have a strong impact on survival; simply being within a pulse was often sufficient for survival.

Neither growth rate nor *gadX* expression alone is expected to completely determine cellular
outcome, as stress response is a complex process; instead, our results suggest that pulsing
dynamics can bias the probability that a single cell will live or die. Cell-to-cell heterogeneity in
gadX expression is the result of pulsing dynamics, which produce diverse distributions of cells.
Thus, we demonstrate that *gadX* pulsing can play a protective role against ciprofloxacin stress,
indicating a clear link between single-cell dynamics and cell fate.

Discussion
Although cells can display substantial phenotypic heterogeneity in constant environmental
conditions, the timescales over which gene expression levels and growth rate fluctuate and how
this ultimately impacts function are often less clear. To provide insight into these questions, we
focused on genes involved in stress response in *E. coli*. Our results demonstrate that distributions
of gene expression levels and growth rates can originate from rich dynamic activity, where
individual cells transition between expression levels over time. Our findings expand previously
published observations on pulsing dynamics (*8*, *13*, *14*, *17*, *38*) to a broader set of genes, suggesting
that these time-varying differences in expression and growth may be more common than
previously appreciated. These fluctuations occur under uniform, unstressed conditions and are
observed across genes with diverse stress response roles. Notably, temporal changes in gene
expression are also related to growth. In our measurements, we found many instances where pulses
of high activity followed decreases in growth rate. However, this link is not universal, which
indicates that this effect is not an unavoidable consequence of growth, and rather that the
expression and growth relationship is specific to each gene. Finally, we asked whether gene
expression dynamics have functional consequences, focusing on expression of *gadX*. We showed
that a functional outcome of *gadX* pulsing and the corresponding decrease in growth rate is that
cells are more likely to survive following sudden exposure to a lethal dose of ciprofloxacin.

Our results with *gadX* join a small number of other studies demonstrating examples of functional
phenotypic heterogeneity. For example, the stochastic activation of diverse stress response genes,
such as those encoding porins (*39*), the catalase-peroxidase KatG (*40*), and the multiple antibiotic
resistance activator MarA (*6*, *41*, *42*) result in transient tolerance to exogenous stress. Other
examples include biased partitioning of efflux pumps during cell division that results in differential

antibiotic susceptibility (*43*) and heterogeneous induction of *gad* regulon genes during antibiotic treatment that cross-protects cells against subsequent acid stress (*9*).

Reduced growth rates can also enable cells to transiently resist stress, for example, by playing a major role in the formation of persister cells (*34*, *44*). By tracking expression dynamics and growth rates simultaneously, we found many instances in which these metrics are inversely correlated over time. Interestingly, an analogous observation was reported by Patange *et al*. (*8*) that demonstrated that the stress response master regulator RpoS is expressed with pulsatile dynamics in exponentially dividing cells and is inversely correlated with growth rate. We demonstrated that RpoS availability directly contributes to *gadX* pulsing dynamics, and could potentially serve as a common mechanism underlying the expression and growth dynamics associated with many additional genes characterized in our study. Other mechanisms are also possible, as some genes in our study with strong growth rate anti-correlations (e.g. *marA*) are not known to be regulated by RpoS. In some instances other mechanisms have been demonstrated, as in a recent study that indicated that self-degradation of the SOS response repressor LexA triggers pulses in the expression of *recA* and *sulA* during exponential growth in unperturbed conditions (*14*). Fluctuations in LexA result in frequent pulses in *recA* expression, while *sulA* is less susceptible to LexA variability, corroborating our observations. It is also noteworthy that, in addition to RpoS, at least one additional sigma factor, RpoH, was activated with pulsatile dynamics during exponential growth, a behavior that can potentially be propagated to >150 downstream genes that collectively control the cellular heat-shock response (*45*, *46*). The pulsatile activity of these regulators might indicate that the mechanism by which sigma factors "time share" RNA polymerase complexes previously described in *B. subtilis* (*47*) might also be present in *E. coli*. Additionally, our results suggest that the pulses of activation of different sigma factors might be associated with different cellular growth statuses.

In future experiments, it would be interesting to conduct measurements using pairs of reporters so that expression pulses can be monitored simultaneously in the same cell. This is particularly interesting to examine in relation to the growth rate, as we observed many instances of anti-correlation, but the temporal properties of these pulses were distinct from each other. Although we measure growth rate in our analysis, our findings do not preclude the possibility that growth rate is a secondary effect that is downstream of the true coordinating signal, such as metabolic state (*48*). Experiments that use reporters to measure genes involved in metabolism alongside the stress response reporters or studies that carefully control the nutrient composition could be used to investigate this possibility. It will also be interesting to investigate how the information encoded in transcription factor dynamics is transmitted to downstream genes, and emerging technologies that enable the manipulation of gene expression over time could help to identify how these signals are propagated (*49*, *50*). Finally, although we have focused our analysis of stress survival on transient ciprofloxacin treatment in wild type cells with the *gadX* reporter, it would be interesting to test survival to other antibiotics, different stress application patterns, and the effect of different strain backgrounds.

In summary, our findings reveal that pulsatile dynamics in gene expression and growth serve as a mechanism that cells can leverage to transiently resist stress. This dynamic behavior is spread across a broad range of stress response genes, including heat-shock response, multi-drug

411 resistance, and oxidative stress and can enable subpopulations of cells to withstand temporary
412 stress.
413
414 **Materials and Methods**
415
Strains and growth media
Strains containing reporter plasmids were sourced from the collection created by Zaslaver *et al.*
(*51*), unless otherwise noted. Briefly, each strain has a low copy number plasmid (SC101 origin)
containing the promoter sequence for the gene of interest upstream of the coding sequence for
gfpmut2 green fluorescent protein (GFP). The strain reporting *purA* was sourced from Rossi *et al.*
(*7*) and the strain reporting *marA* was sourced from El Meouche *et al.* (*6*), where the *marA* reporter
has intact MarR binding sites. The *purA* and *marA* reporters use a low copy number plasmid
(SC101 origin) and the promoters of interest control the expression of cerulean cyan fluorescent
protein (CFP). The constitutive reporter plasmid was constructed by cloning a constitutive
promoter from Ref. (*52*) into the empty vector from the Zaslaver *et al.* library, pUA66.
Specifically, the promoter used here is a variant of the σ70 consensus sequence:
TTATCAAAAAGAGTA<u>TTGTCT</u>TAAAGTCTAACCTATAG<u>GAAAAT</u>TACAGCC**A**TCGAG
AGGGACACGGCGAA where the +1 transcriptional start site is shown in bold and the -10 and -
35 regions are underlined. In all cases, we used *E. coli* MG1655 as the strain background.
430
The strain containing an integrated *gadX* reporter was constructed as described by Patange *et al.*
(*8*). A cassette containing the *gadX* reporter and a kanamycin resistance gene was amplified from
the reporter plasmid using the primers
5'-ATAAACACGTTCGTGTCCCGACAGGCACACAGACGGTTAGCCACTAATTAGAGCT
CTCGAACCCCAGAGT-3'
and
5'-
GTAAGAATAAAAAAAACGGGTCACCTTCTGGCGACCCGTTTTTCTTTGCGCCTGCA
GGTCTGGACATTTA-3'
and integrated into the chromosome between the *nupG* and *speC* genes.
441
The Δ*rpoS* strain was constructed as described by Baba *et al.* (*53*). Briefly, an in-frame deletion of
rpoS was achieved through integration of a cassette containing a kanamycin resistance gene
between FLP recognition target sites and 50 base-pair homology sequences for the regions flanking
rpoS:
5'-TGAGACTGGCCTTTCTGACAGATGCTTACTTACTCGCGGAACAGCGCTTC-3'
and
5'-CTTTTGCTTGAATGTTCCGTCAAGGGATCACGGGTAGGAGCCACCTTATG-3'
After successful integration, the kanamycin resistance gene was excised using FLP recombinase.
450
451 Cells were grown in M9 medium (0.1 mM $CaCl_2$; 2 mM $MgSO_4$; 1X M9 salts) supplemented with
452 0.4% glucose, 0.2% casamino acids, and 30 µg/ml kanamycin for plasmid maintenance. Media
453 used in microfluidic experiments were supplemented with 2 g/L F-127 Pluronic to prevent cell
454 adhesion and growth outside of the mother machine growth chambers.
455
Static fluorescence microscopy snapshots

Overnight cultures were diluted 1:100 and incubated at 37C with shaking for 2.5 to 4 hours
(OD$_{600nm}$= 0.7-1.3). 1 μL of cells were placed on 1.5% MGC (0.2% glycerol, 0.01% casamino
acids, 0.15 μg/ml biotin, and 1.5 μM thiamine) low melting point agarose pads and imaged at 100X
on a Nikon Ti-E inverted fluorescence microscope. Three separate overnight cultures were
sampled and imaged for each reporter strain.
462

Mother machine microfluidic device
The mother machine microfluidic master mold used was described previously in Ref. (*26*). Briefly,
the master mold chip has 8 independent main feed channels where growth media flows in and out.
Each channel features 1,000 one-ended chambers (L x W x H = 25 μm x 1.3 μm x 1.1 μm) where
the mother cells are trapped. We made the microfluidic devices by pouring a degassed 10:1 mixture
of dimethyl-siloxane monomer and curing agent (Sylgard 184 Silicone Elastomer Kit, Dow
Corning) onto the wafer, which was then cured overnight at 65C. Individual chips were separated
from the mold and a 0.75 mm biopsy punch was used to create inlets and outlets for each flow
channel before the chip was plasma bonded to a glass slide.
472

Time-lapse microscopy movies
Overnight cultures were diluted 1:100 and incubated at 37C with shaking for ~3 hours until mid-
exponential phase (OD$_{600nm}$= 1.0-1.3). Cells were concentrated by centrifugation (6,000xg for 2
min) and loaded into the microfluidic chip. Cells were seeded in the growth chambers by
centrifugation at 6,000xg for 3 min. Media was supplied at a flow rate of 20 μL/min using a
peristaltic pump. Cells were allowed to adapt to growth in the device for 2-4 hours before imaging.
Time-lapse movies were acquired with a 100X oil objective on a Nikon Ti-E inverted fluorescence
microscope equipped with a perfect focus system and a temperature control chamber that was set
at 37C for the duration of the experiment. Images were acquired in phase contrast and
epifluorescence illumination with a GFP or CFP filter every 5 min. The total number of cell
lineages tracked for each reporter is listed in Table S2.
484

Microscopy image processing and fluorescence and growth rate analysis
For static snapshots, cell images were segmented with the SuperSegger software (*54*). For mother
machine experiments, we performed automated cell segmentation and lineage tracking using the
deep learning-based software DeLTA (*26*). In both cases, reported single-cell fluorescence values
are the average fluorescence intensities of all pixels belonging to each cell. Unless otherwise noted,
for mother machine experiments only data for the 'mother' cell trapped at the top of the growth
chambers was used in quantitative analysis. We smoothed the fluorescence data using a moving
average filter with a window of 5 frames.
493

To estimate growth rates, we calculated the difference in cell length between adjacent frames
normalized by the cell length, 1/L dL/dt. Specifically, when there is no cell division event between
frames, this is calculated as $(L_t - L_{t-1})/ (L_{t-1} \Delta t)$, where L_t is the length of the cell at time t, L_{t-1} is
the length at the preceding frame, and Δt is the time between frames (5 min). In cases where a cell
division event occurs, this calculation is modified to $(L_t + D_t - L_{t-1})/ (L_{t-1} \Delta t)$, where D_t is the length
of the daughter at time t. After calculating the growth rate, we smoothed the data using a moving
average filter with a window of 5 frames.
501

502 Cell division times are calculated as the time between division events.

Pulse identification

505 To allow comparison across all reporters, which have different expression levels and where
506 imaging exposure times are different (Table S2), we normalized each time-series by its mean.
507 Pulses of gene expression were identified using the built-in MATLAB function *findpeaks*, which
508 identifies local maxima. We set the threshold for the minimum peak prominence at 0.5, unless
509 otherwise noted. This value was determined empirically (Fig. S4).

Autocorrelation and cross-correlation calculations

We calculated the autocorrelation of the fluorescence signal—$R_{fluo,fluo}(\tau)$ where τ is the time
shift—using the MATLAB function *xcov* with the unbiased normalization method. We then
normalized the entire signal by the value of the autocorrelation at zero time lag, $R_{fluo,fluo}(0)$. The
half life was calculated by taking the mean of all autocorrelation signals from individual cell
lineages and identifying the time shift value where it crosses 0.5.

The cross-correlation between the fluorescence signal and the growth rate, $R_{fluo,gr}(\tau)$, was
calculated using the MATLAB function *xcov* with the unbiased normalization method. The signal
was then normalized by the square root of the product of the autocorrelation at zero time shift of
the fluorescence and growth rate signals: $(R_{fluo,fluo}(0)\ R_{gr,gr}(0))^{1/2}$.

$R_{extreme}$ is the value of $R_{fluo,gr}(\tau)$ with the largest absolute value. $\tau_{extreme}$ is the time lag value
associated with $R_{extreme}$.

Antibiotic survival assay

Mother machine experiments were initiated using the methods described above. During image
acquisition, cells were provided with fresh medium for 14 hours, followed by 35 minutes of
treatment with medium supplemented with 2 μg/ml ciprofloxacin, then fresh medium again for 16
hours. The outcomes of ciprofloxacin treatment for each lineage were manually scored as:
'survived,' 'died,' or 'filamented.' Cells were scored as 'survived' if cell division was observed in
the chamber at the end of the full 16 hours after the second addition of fresh medium. Cells were
scored as 'died' if growth permanently ceased in the chamber at any point after antibiotic exposure.
'Filamented' cells were excluded from the analysis, as it was difficult to accurately assess their
outcome since they were frequently swept out of the chamber and field of view.

References

1. Y. Taniguchi, P. J. Choi, G. W. Li, H. Chen, M. Babu, J. Hearn, A. Emili, X. Sunney Xie, Quantifying E. coli proteome and transcriptome with single-molecule sensitivity in single cells. *Science*. **329**, 533–538 (2010).

2. D. J. Kiviet, P. Nghe, N. Walker, S. Boulineau, V. Sunderlikova, S. J. Tans, Stochasticity of metabolism and growth at the single-cell level. *Nature*. **514**, 376–379 (2014).

3. O. K. Silander, N. Nikolic, A. Zaslaver, A. Bren, I. Kikoin, U. Alon, M. Ackermann, A genome-wide analysis of promoter-mediated phenotypic noise in Escherichia coli. *PLoS Genet.* **8** (2012), doi:10.1371/journal.pgen.1002443.

4. A. Raj, A. van Oudenaarden, Nature, Nurture, or Chance: Stochastic Gene Expression and Its Consequences. *Cell*. **135**, 216–226 (2008).

5. A. Bar-Even, J. Paulsson, N. Maheshri, M. Carmi, E. O'Shea, Y. Pilpel, N. Barkai, Noise in protein expression scales with natural protein abundance. *Nat. Genet.* **38**, 636–643 (2006).

6. I. El Meouche, Y. Siu, M. J. Dunlop, Stochastic expression of a multiple antibiotic resistance activator confers transient resistance in single cells. *Sci. Rep.* **6**, 1–9 (2016).

7. N. A. Rossi, I. El Meouche, M. J. Dunlop, Forecasting cell fate during antibiotic exposure using stochastic gene expression. *Commun. Biol.* **2**, 1–7 (2019).

8. O. Patange, C. Schwall, M. Jones, C. Villava, D. A. Griffith, A. Phillips, J. C. W. Locke, Escherichia coli can survive stress by noisy growth modulation. *Nat. Commun.* **9** (2018), doi:10.1038/s41467-018-07702-z.

9. K. Mitosch, G. Rieckh, T. Bollenbach, Noisy Response to Antibiotic Stress Predicts Subsequent Single-Cell Survival in an Acidic Environment. *Cell Syst.* **4**, 393-403.e5 (2017).

10. S. Uphoff, N. D. Lord, B. Okumus, L. Potvin-trottier, D. J. Sherratt, J. Paulsson, Stochastic activation of a DNA damage response causes cell-to-cell mutation rate variation. *Science*. **351**, 1094–1098 (2016).

11. S. Uphoff, Real-time dynamics of mutagenesis reveal the chronology of DNA repair and damage tolerance responses in single cells. *Proc. Natl. Acad. Sci.* **115**, E6516–E6525 (2018).

12. I. El Meouche, M. J. Dunlop, Heterogeneity in efflux pump expression predisposes antibiotic-resistant cells to mutation. *Science*. **362**, 686–690 (2018).

13. J. H. Levine, Y. Lin, M. B. Elowitz, Functional roles of pulsing in genetic circuits. *Science*. **342**, 1193–1200 (2013).

14. E. C. Jones, S. Uphoff, Single-molecule imaging of LexA degradation in Escherichia coli elucidates regulatory mechanisms and heterogeneity of the SOS response. *Nat. Microbiol.*, 1–10 (2021).

15. G. M. Süel, R. P. Kulkarni, J. Dworkin, G.-O. Jordi, M. B. Elowitz, Tunability and noise dependence in differentiation dynamics. *Science*. **315**, 1716–1719 (2007).

16. H. Maamar, A. Raj, D. Dubnau, Noise in gene expression determines cell fate in Bacillus subtilis. *Science*. **317** (2007), doi:10.1038/jid.2014.371.

17. J. M. Kim, M. Garcia-Alcala, E. Balleza, P. Cluzel, Stochastic transcriptional pulses orchestrate flagellar biosynthesis in Escherichia coli. *Sci. Adv.* **6**, eaax0947 (2020).

18. M. M. Logsdon, B. B. Aldridge, Stable regulation of cell cycle events in mycobacteria: Insights from inherently heterogeneous bacterial populations. *Front. Microbiol.* **9** (2018), p. 514.

19.   Y. Tanouchi, A. Pai, H. Park, S. Huang, R. Stamatov, N. E. Buchler, L. You, A noisy
linear map underlies oscillations in cell size and gene expression in bacteria. *Nature*. **523**,
357–359 (2015).

20.   E. Biselli, S. J. Schink, U. Gerland, Slower growth of *Escherichia coli* leads to longer
survival in carbon starvation due to a decrease in the maintenance rate. *Mol. Syst. Biol.* **16**
(2020), doi:10.15252/msb.20209478.

21.   D. A. F. Fuentes, P. Manfredi, U. Jenal, M. Zampieri, Pareto optimality between growth-
rate and lag-time couples metabolic noise to phenotypic heterogeneity in Escherichia coli.
Nat. Commun. 2021 121. **12**, 1–12 (2021).

22.   J. Narula, A. Kuchina, F. Zhang, M. Fujita, G. M. Süel, O. A. Igoshin, Slowdown of
growth controls cellular differentiation. *Mol. Syst. Biol.* **12**, 871 (2016).

23.   B. B. Aldridge, M. Fernandez-Suarez, D. Heller, V. Ambravaneswaran, D. Irimia, M.
Toner, S. M. Fortune, Asymmetry and aging of mycobacterial cells lead to variable
growth and antibiotic susceptibility. *Science*. **335**, 100–104 (2012).

24.   A. J. Lee, S. Wang, H. R. Meredith, B. Zhuang, Z. Dai, L. You, Robust, linear correlations
between growth rates and β-lactam–mediated lysis rates. *Proc. Natl. Acad. Sci.* **115**, 4069–
4074 (2018).

25.   P. Wang, L. Robert, J. Pelletier, W. L. Dang, F. Taddei, A. Wright, S. Jun, Robust growth
of Escherichia coli. *Curr. Biol.* **20**, 1099–1103 (2010).

26.   J. B. Lugagne, H. Lin, M. J. Dunlop, DeLTA: Automated cell segmentation, tracking, and
lineage reconstruction using deep learning. *PLoS Comput. Biol.* **16**, e1007673 (2020).

27.   A. Tramonti, P. Visca, M. De Canio, M. Falconi, D. De Biase, Functional characterization
and regulation of gadX, a gene encoding an AraC/XylS-like transcriptional activator of
the Escherichia coli glutamic acid decarboxylase system? *J. Bacteriol.* **184**, 2603–2613
(2002).

28.   S. W. Seo, D. Kim, E. J. O'Brien, R. Szubin, B. O. Palsson, Decoding genome-wide
GadEWX-transcriptional regulatory networks reveals multifaceted cellular responses to
acid stress in Escherichia coli. *Nat. Commun.* **6** (2015), doi:10.1038/ncomms8970.

29.   E. Balleza, J. M. Kim, P. Cluzel, Systematic characterization of maturation time of
fluorescent proteins in living cells. *Nat. Methods*. **15**, 47 (2018).

30.   A. Mukherjee, C. Cao, J. Lutkenhaus, Inhibition of FtsZ polymerization by SulA, an
inhibitor of septation in Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 2885–2890
(1998).

31.   O. Huisman, R. D'Ari, S. Gottesman, Cell-division control in Escherichia coli: Specific
induction of the SOS function SfiA protein is sufficient to block septation. *Proc. Natl.*
Acad. Sci. U. S. A. **81**, 4490–4494 (1984).

32.   J. Bos, R. H. Austin, E. Rogers, S. Vyawahare, Q. Zhang, S. M. Rosenberg, Emergence of
antibiotic resistance from multinucleated bacterial filaments. *Proc. Natl. Acad. Sci.* **112**,
178–183 (2014).

33.   E. Tuomanen, R. Cozens, W. Tosch, O. Zak, A. Tomasz, The rate of killing of Escherichia
coli by β-lactam antibiotics is strictly proportional to the rate of bacterial growth. *J. Gen.*
Microbiol. **132**, 1297–1304 (1986).

34.   M. H. Pontes, E. A. Groisman, Slow growth determines nonheritable antibiotic resistance
in Salmonella enterica. *Sci. Signal.* **12**, 1–11 (2019).

35.   D. J. Evans, D. G. Allison, M. R. W. Brown, P. Gilbert, Susceptibility of Pseudomonas
aeruginosa and Escherichia coli biofilms towards ciprofloxacin: Effect of specific growth

630   rate. *J. Antimicrob. Chemother.* **27**, 177–184 (1991).

36.   I. Yelin, R. Kishony, Antibiotic Resistance. *Cell.* **172** (2018),
doi:10.1016/j.cell.2018.02.018.

37.   A. C. Palmer, R. Kishony, Opposing effects of target overexpression reveal drug
mechanisms. *Nat. Commun.* (2015), doi:10.1038/ncomms5296.Opposing.

38.   J. E. Purvis, G. Lahav, Encoding and decoding cellular information through signaling
dynamics. *Cell.* **152**, 945–956 (2013).

39.   M. A. Sánchez-Romero, J. Casadesús, Contribution of phenotypic heterogeneity to
adaptive antibiotic resistance. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 355–360 (2014).

40.   Y. Wakamoto, N. Dhar, R. Chait, K. Shneider, F. Signorino-Gelo, S. Leibler, J. D. .
McKinney, Dynamic persistence of antibiotic-stressed mycobacteria. *Science.* **339**, 91–96
(2013).

41.   N. A. Rossi, M. J. Dunlop, Customized Regulation of Diverse Stress Response Genes by
the Multiple Antibiotic Resistance Activator MarA. *PLoS Comput. Biol.* **13**, e1005310
(2017).

42.   J. Garcia-Bernardo, M. J. Dunlop, Tunable stochastic pulsing in the Escherichia coli
multiple antibiotic resistance network from interlinked positive and negative feedback
loops. *PLoS Comput. Biol.* **9**, e1003229-11 (2013).

43.   T. Bergmiller, A. M. C. Andersson, K. Tomasek, E. Balleza, D. J. Kiviet, R. Hauschild, G.
Tkačik, C. C. Guet, Biased partitioning of the multidrug efflux pump AcrAB-TolC
underlies long-lived phenotypic heterogeneity. *Science.* **356**, 311–315 (2017).

44.   N. Q. Balaban, J. Merrin, R. Chait, L. Kowalik, S. Leibler, Bacterial persistence as a
phenotypic switch. *Science.* **305**, 1622–1625 (2004).

45.   F. Arsène, T. Tomoyasu, B. Bukau, The heat shock response of Escherichia coli. *Int. J.*
Food Microbiol. **55**, 3–9 (2000).

46.   T. Yura, Regulation of the heat shock response in Escherichia coli: History and
perspectives. *Genes Genet. Syst.* **94**, 103–108 (2019).

47.   J. Park, M. Dies, Y. Lin, S. Hormoz, S. E. Smith-Unna, S. Quinodoz, M. J. Hernández-
Jiménez, J. Garcia-Ojalvo, J. C. W. Locke, M. B. Elowitz, Molecular Time Sharing
through Dynamic Pulsing in Single Cells. *Cell Syst.* **6**, 216-229.e15 (2018).

48.   A. J. Lopatkin, J. M. Stokes, E. J. Zheng, J. H. Yang, M. K. Takahashi, L. You, J. J.
Collins, Bacterial metabolic state more accurately predicts antibiotic lethality than growth
rate. *Nat. Microbiol.* **4** (2019), pp. 2109–2117.

49.   J. B. Lugagne, M. J. Dunlop, Cell-machine interfaces for characterizing gene regulatory
network dynamics. *Curr. Opin. Syst. Biol.* **14**, 1–8 (2019).

50.   N. M. V. Sampaio, M. J. Dunlop, Functional roles of microbial cell-to-cell heterogeneity
and emerging technologies for analysis and control. *Curr. Opin. Microbiol.* **57**, 87–94
(2020).

51.   A. Zaslaver, A. Bren, M. Ronen, S. Itzkovitz, I. Kikoin, S. Shavit, W. Liebermeister, M.
G. Surette, U. Alon, A comprehensive library of fluorescent transcriptional reporters for
Escherichia coli. *Nat. Methods.* **3**, 623–628 (2006).

52.   Y. Siu, J. Fenno, J. M. Lindle, M. J. Dunlop, Design and Selection of a Synthetic
Feedback Loop for Optimizing Biofuel Tolerance. *ACS Synth. Biol.* **7** (2017),
doi:10.1021/acssynbio.7b00260.

53.   T. Baba, T. Ara, M. Hasegawa, Y. Takai, Y. Okumura, M. Baba, K. A. Datsenko, M.
Tomita, B. L. Wanner, H. Mori, Construction of Escherichia coli K-12 in-frame, single-

676   gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* (2006).

677 54. S. Stylianidou, C. Brennan, S. B. Nissen, N. J. Kuwada, P. A. Wiggins, SuperSegger:

678   robust image segmentation, analysis and lineage tracking of bacterial cells. *Mol.*

679   *Microbiol.* **102**, 690–700 (2016).

Acknowledgments: We thank members of the Dunlop Lab and Wilson Wong for helpful
discussions; Ahmad (Mo) Khalil and Razan Alnahhas provided input on the manuscript. Razan
Alnahhas and Eric South conducted initial experiments with the constitutive reporters. **Funding:**
This work was supported by the National Institutes of Health grants R01AI102922 and
R21AI137843. CMB received support from the National Science Foundation Graduate Research
Fellowship under Grant No. DGE-1840990. **Author contributions:** NMVS and MJD
conceptualized the study. NMVS and CMB performed experiments. NMVS, CMB, JBL, and MJD
performed data analysis. NMVS and MJD wrote the manuscript, with input from CMB and JBL.
MJD supervised the study and acquired funding. **Competing interests:** All authors declare that
they have no competing interest.
696
697

698   **Figures**

699
700
701   **Figure 1. Cell-to-cell heterogeneity and temporal dynamics of gene expression. (A)**
702   Histograms of fluorescence values show cell-to-cell variation. Values are presented normalized
703   relative to their means to allow for comparison across reporters. The same data without
704   normalization are shown in Fig. S1. Corresponding coefficient of variation (CV) values obtained
705   from snapshot images of cells grown in bulk cultures are listed in the figure. The skewness (skew)

of the distribution, which is positive for distributions that are right-skewed, is also listed. Insets show phase contrast and fluorescence images for representative strains. Functional classes are listed for each gene. Scale bar, 2 μm. **(B)** Schematic representation of fluorescence signal over time originating from single cells with slow or fast dynamics of gene expression that result in identical fluorescence distributions for the final timepoint. **(C-E)** Representative kymographs of cells containing transcriptional fluorescent reporters for **(C)** *gadX*, **(D)** *recA*, and **(E)** *araC*. In all cases, fluorescence values are normalized to the mean to allow comparisons across the reporters. Scale bar, 5 μm. **(F)** Single-cell measurements of green fluorescent protein expression (GFP) over time for the *gadX* reporter. Colored heat maps summarize the time-series data. Data from 25 cells are shown, however this represents a subset of the time-series data. **(G)** Heat maps summarizing the temporal dynamics of gene expression for all reporters. **(H)** Constitutive reporter heat map. **(I)** Chromosomally-integrated *gadX* reporter heat map. Color scale and normalization approach in (F) applies to all heat map data.

Figure 2. Pulsing dynamics vary across the reporters. (A) Schematic representation of pulse
characteristics. **(B)** Distributions of pulse frequency, duration, and amplitude for all reporters. Note
that typical movie durations are 15 - 20 hours (Table S2), thus we are limited in our ability to
measure very low frequencies accurately (1/15 = 0.067). Duration and amplitude histograms are
normalized such that the maximal height corresponds to the fraction of lineages with at least one
pulse. **(C)** Autocorrelation of the fluorescence signal for independent cell lineages with the *gadX*
reporter. Half life is defined as the value of the time shift τ when the mean of the autocorrelation
curve crosses 0.5. **(D)** Cell division time and half life for each of the reporters. Error bars show
standard deviation around the mean. **(E)** Heat maps summarizing single-cell measurements of
green fluorescent protein expression (GFP) over time for the *gadX* reporter in wild type and Δ*rpoS*

cells. Identical reporters are used in both strains, thus the unnormalized data can be compared
directly. Normalized versions are shown in Fig. S7. **(F)** Frequency, duration, and amplitude
distributions for the *gadX* reporter in the wild type and $\Delta rpoS$ backgrounds.

Figure 3. Growth and gene expression dynamics are often anti-correlated. (A) Two possible
models for the relationship between fluorescent reporter levels and growth. **(B)** Schematic of cross-
correlation function $R_{fluo,gr}(\tau)$ indicating points corresponding to $R_{extreme}$ and $\tau_{extreme}$ (top).
Temporal patterns of the fluorescence and growth rate and their impact on $R_{extreme}$ and $\tau_{extreme}$
(bottom). **(C)** Representative fluorescence (green) and growth rate (magenta) values over time for
a single mother cell with a reporter for *gadX* (left) or *recA* (right). Black dots at the top of the
figure indicate cell division events. Data are smoothed with a moving average filter with a window
of 5 frames. **(D)** Cross-correlations for all reporters. Shaded region represents standard deviation
about the mean. **(E)** $R_{extreme}$ and **(F)** $\tau_{extreme}$ values for each reporter. **(G)** $R_{extreme}$ as a function of
$\tau_{extreme}$. For signals without a peak ($|R_{extreme}| < 0.1$) we do not calculate a $\tau_{extreme}$ value.

754
755

Figure 4. Cell-to-cell differences in *gadX* expression and growth influence survival outcomes after antibiotic exposure. (A) Representative kymograph of a cell lineage that survives and **(B)** dies after antibiotic treatment. Images are from nearby chambers within the same microfluidic chip. Shaded area in red indicates 35 minute period of ciprofloxacin treatment. **(C)** Single-cell *gadX* expression and **(D)** growth rate for the period preceding ciprofloxacin treatment. Cell numbers in (C) and (D) correspond to the same cells. Time-series are sorted by the mean value of the fluorescence signal for t < 0, from high to low. **(E)** Individual cells that survived (green) or died (gray) following ciprofloxacin treatment plotted as a function of growth rate and fluorescence at t = 0. Error bars show standard deviation about the mean. **(F)** Mean fluorescence from the *gadX* reporter for the period preceding ciprofloxacin addition. Shaded regions correspond to standard deviation about the mean. Statistics are conducted at each timepoint; the horizontal bar indicates all time points where *** P < 0.001, two-sample *t* test for differences between the mean of the cells that survived and died. **(G)** Histogram of duration of the pulse that cells were in upon ciprofloxacin addition, sorted by outcome.

770
771
772

Supplementary Figures

 Figure S1. Histograms of fluorescence distributions of single cells. Microscopy exposure times vary between reporters to optimize detection of fluorescence from each reporter and are indicated in the figure.

Figure S2. Histograms displaying distributions of mean normalized fluorescence values for single cells imaged in snapshots of agarose pads (blue) or in the mother machine device (red).

Figure S3. Mean fluorescence of all mother cells over time. Shaded area represents the 25 to 75 percentile values for each reporter; center line is the mean.

Figure S4. Effect of threshold value on pulse frequency, duration, and amplitude. Note that as the threshold is increased, we exclude small pulses that result from noise in the signal. This results in decreases in the frequency and generally increases in the duration and amplitude of the pulses. The default threshold value used for analysis is 0.5.

832
833
834 **Figure S5.** Autocorrelation of the fluorescence signal for all reporters. Shaded area shows standard
835 deviation around the mean.
836
837
838

Figure S6. Correlation coefficient between mother cell's mean fluorescence signal and its offspring. "Generations from mother" = 1 corresponds to a mother-daughter comparison, 2 is a mother-granddaughter comparison, and 3 is a mother-great granddaughter comparison.

866
867 **Figure S7.** Single-cell measurements of green fluorescent protein expression (GFP) over time for
868 the *gadX* reporter shown normalized relative to their mean in the wild type and Δ*rpoS* backgrounds.
869
870
871

Figure S8. (A) Autocorrelation and **(B)** cross-correlation for the *gadX* reporter comparing plasmid-based and chromosomally-integrated versions. **(C)** Autocorrelation and **(D)** cross-correlation for the constitutive reporter. Shaded regions show standard deviation about the mean.

Figure S9. Relationship between cell length and fluorescence across reporters. (A) Distributions of cell length for all reporters. **(B)** Correlation coefficient values for cell length and fluorescence for all reporters. **(C)** As representative examples, insets show *gadX* or *recA* reporter levels plotted as a function of cell length for all time points and replicates. Correlation coefficient is listed in the figure title.

896 **Supplementary Tables**

898 **Table S1.** Summary of genes characterized in this study

Gene	Sigma Factor	Function	# of Regulatees
araC	RpoD	Arabinose catabolism	14
bolA	RpoD, RpoS	Cell morphology and protection, general stress response	5
dinB	RpoD, RpoS	SOS response, encodes DNA polymerase IV	0
evgA	RpoD, RpoS	Multi-drug resistance	19
gadW	RpoS	Acid resistance regulon	15
gadX	RpoD, RpoS	Acid resistance regulon	34
marA	RpoD	Multiple antibiotic resistance	45
metJ	RpoD	Methionine biosynthesis	15
oxyR	RpoD, RpoS	Oxidative stress	42
phoP	RpoH	Adaptation to low Mg2+, control of acid resistance genes	56
purA	RpoD	*de novo* synthesis of AMP	0
recA	RpoD	DNA recombination / repair protein, SOS response.	0
rob	Unknown	Resistance to antibiotics, organic solvents, heavy metals	26
rpoH	RpoD, RpoN, RpoE, RpoS	Sigma32. Response to heat shock during log-phase growth	174
sulA	RpoD	SOS response cell division inhibitor	0

900
901

902 **Table S2.** Number of cell lineages tracked for each reporter, where each lineage corresponds to a
903 single mother cell. Movie duration is the length of the measurement. Exposure time for the mother
904 machine time-lapse experiments, which is customized for each reporter to account for differences
905 in expression levels.
906

Gene	# cell lineages tracked	Movie duration (median, in hours)	Exposure time (msec)
araC	71	15.5	150
bolA	72	22.0	300
dinB	88	16.2	300
evgA	72	16.9	30
gadW	107	15.9	10
gadX	142	15.0	30
marA	450	16.3	50
metJ	71	15.5	150
oxyR	71	15.5	150
phoP	72	15.5	150
purA	101	16.8	50
recA	153	16.2	300
rob	108	15.9	10
rpoH	72	19.9	30
sulA	336	17.2	300
constitutive	216	16.3	350
gadX chromosomal	200	4.4	200
gadX (WT)	346	9.0	50
gadX (ΔrpoS)	221	9.0	50

907
908
909

Supplementary Movie Captions

Movie S1. Time-lapse movie of a representative mother machine chamber containing cells with the *gadX* reporter. Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.

Movie S2. Time-lapse movie of a representative mother machine chamber containing cells with the *recA* reporter. Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.

Movie S3. Time-lapse movie of a representative mother machine chamber containing cells with the *araC* reporter. Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.

Movie S4. Surviving cell in a time-lapse movie where cells containing the *gadX* reporter are exposed to ciprofloxacin (period of treatment indicated with red bar at top of the image). Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.

Movie S5. Dying cell in a time-lapse movie where cells containing the *gadX* reporter are exposed to ciprofloxacin (period of treatment indicated with red bar at top of the image). Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.