Cell-to-cell heterogeneity in *Escherichia coli* stress response originates from pulsatile expression and growth

Nadia M. V. Sampaio[1,2,†], Caroline M. Blassick[1,2], Jean-Baptiste Lugagne[1,2], Mary J. Dunlop[1,2,*]

[1] Biomedical Engineering Department, Boston University, Boston, MA 02215.
[2] Biological Design Center, Boston University, Boston, MA 02215.
[†] Present Address: Brazilian Biorenewables National Laboratory, Brazilian Center for Research in Energy and Materials, Campinas, São Paulo, Brazil.
[*] Corresponding Author: mjdunlop@bu.edu

Abstract

Cell-to-cell heterogeneity in gene expression and growth can have critical functional consequences, such as determining whether individual bacteria survive or die following stress. Although phenotypic variability is well documented, the dynamics that underlie it are often unknown. This information is critical because dramatically different outcomes can arise from gradual versus rapid changes in expression and growth. Using single-cell time-lapse microscopy, we measured the temporal expression of a suite of stress response reporters in *Escherichia coli*, while simultaneously monitoring growth rate. In conditions without stress, we found widespread examples of pulsatile expression. Single-cell growth rates were often anti-correlated with gene expression, with changes in growth preceding changes in expression. These pulsatile dynamics have functional consequences, which we demonstrate by measuring survival after challenging cells with the antibiotic ciprofloxacin. Our results suggest that pulsatile expression and growth dynamics are common in stress response networks and can have direct consequences for survival.

MAIN TEXT

Introduction

Even under otherwise constant environmental conditions, genetically identical cells can display substantial phenotypic heterogeneity, resulting from differences in gene expression and growth rate from cell to cell [1–4]. Phenotypic heterogeneity can, in principle, arise from slow changes within the population where individual cells have different expression levels or growth rates, but maintain their state over many cell cycles. Alternatively, fast dynamics with rapid fluctuations could produce equivalent distributions. The distinction between these alternatives is significant because the timescale over which expression or growth differences persist can ultimately determine if they have functional consequences or are simply short-lived random variations that are filtered before impacting cellular outcomes. Although phenotypic heterogeneity is well documented, the timescales underlying the variation and their ultimate impact on function are often less clear.

We focused on stress response genes in *Escherichia coli* to study the dynamics of expression and growth in single cells. Genes involved in adaptation to stress are among the noisiest genome-wide [3,5]. Studies on individual pathways have revealed specific examples in which heterogeneous expression of stress response genes can allow subpopulations of cells to survive sudden environmental stress, such as transient exposure to antimicrobial drugs, oxidative stress, and acid stress [6–9]. Recent studies also indicate that heterogeneity in the expression of genes involved in DNA repair can lead to variability in mutation rates, contributing to microbial evolution [10–12].

Notably, this diversity exists even in the absence of stressors. Although selected studies have begun
to reveal examples of how cell-to-cell phenotypic variation can provide important functional
capabilities to cell populations, examples of direct links are relatively sparse compared to reports
quantifying phenotypic heterogeneity. This motivated our focus on stress response pathways,
because the effect of the genes involved can be directly assessed by quantifying outcomes like cell
survival versus death following stress.

In addition to measurements at a single time point, long-term monitoring of gene expression has
begun to uncover examples of rich dynamics in key stress response proteins [13]. For example, self-
cleavage of the regulator LexA produces spontaneous pulses in the SOS response network [14].
Dynamic activity of transcription factors can propagate to downstream genes, with direct
consequences for stress tolerance. For instance, pulsatile expression of the transcription factor
ComK enables *Bacillus subtilis* cells to enter a transient competent state [15,16]. In *E. coli*,
heterogeneous expression of RpoS, a key regulator of general stress response, originates from
pulses of activation that are inversely correlated with growth, allowing cells to survive oxidative
stress [8]. Additionally, Kim *et al.* recently demonstrated that genes in the flagellar synthesis network,
a process with a pivotal role in microbial pathogenicity, are expressed with different pulsing
programs that allow cells to switch between flagellar phenotypes [17]. Collectively, these studies
demonstrate that the temporal dynamics of gene expression play an important role in the regulation
of stress response networks.

Growth rate fluctuations have also been observed in single cells across many bacterial species [2,8,18–
20]. They can arise due to temporal variation in the expression of metabolic enzymes [2], the expression
of burdensome proteins [12], and due to regulatory effects such as feedback involved in cell size
control [19]. These single-cell differences in growth rate can play a functional role in stress tolerance.
For example, Narula *et al.* demonstrated that the growth rate of *B. subtilis* under starvation
conditions can determine whether individual cells differentiate into spores or remain vegetative [21].
Single-cell growth rates also play a protective role, for example correlating with differential
antibiotic susceptibility in mycobacteria [22] and *E. coli* [7,23].

Understanding the timescales associated with gene expression dynamics and growth can provide
critical insights into the strategies that cells use to hedge against environmental uncertainty. In this
study, we characterize the prevalence and dynamic properties of cell-to-cell phenotypic variation
in different branches of the stress response network in *E. coli*. Using time-lapse microscopy, we
monitored the activity of key stress response genes, as well as genes involved in biosynthesis and
metabolism, in single *E. coli* cells under precisely controlled, unstressed conditions. Our results
reveal that pulsatile activity is common in these reporters. Furthermore, properties of the dynamics,
such as the frequency or amplitude of pulsing, are unique to each gene. In addition, fluctuations in
the expression of genes frequently occur following variations in growth rate. Finally, focusing on
the acid stress regulator GadX, we show that coincident pulses of upregulation of stress response
genes and reduced growth rates favor tolerance to lethal antibiotic exposure. Together, this work
reveals that non-trivial gene expression dynamics are common, even in otherwise constant
conditions, and that these dynamic patterns of gene expression can have critical consequences for
cell survival.

Results
Single-cell measurements of phenotypic heterogeneity
93 We began by characterizing heterogeneity in the expression of genes with a diverse range of
94 functions (Table S1). These include genes involved in acid resistance (*gadW*, *gadX*, *phoP*), multi-
95 drug resistance (*evgA*, *marA*, *rob*), heat shock (*rpoH*), oxidative stress response (*oxyR*), SOS

response (*dinB*, *recA*, *sulA*), and general stress response (*bolA*), in addition to genes involved in biosynthesis and metabolism (*araC*, *metJ*, *purA*).

In order to measure heterogeneity in the expression from each promoter at the single-cell level, we used strains containing transcriptional reporters where the promoter sequence of interest is fused to the coding sequence of a fluorescent protein. We grew independent bulk cultures of each strain to exponential phase and measured heterogeneity across cells in the population using fluorescence microscopy for single cells on agarose pads. Cell-to-cell differences in gene expression resulted in different distributions of reporter levels for each gene (Fig. 1A, Fig. S1). Measurements of the reporters revealed many instances of wide distributions, indicative of a broad range of expression levels across cells within a population. Distributions tended to skew to the right of the mean (skew > 0) due to the presence of highly-expressing cells.

However, measures of static distributions do not reveal the underlying dynamics that generate them, prompting the question: Do these distributions stem from long-lived fixed subpopulations with slow dynamics or from conditions with fast dynamics where individual cells transition between low and high expression over time (Fig. 1B)? The significance of this question is particularly pertinent in the context of stress response networks, where the timescale over which individual genes are active may have concrete implications for tolerance levels. For instance, if a transcription factor that activates genes involved in stress response exhibits short pulses in expression, these pulses might be insufficient to turn on expression of downstream genes, while more sustained expression could. Further, single-cell growth rates can impact survival, thus the interplay between expression and growth may be significant for determining tolerance.

Pulsatile gene expression underlies cell-to-cell heterogeneity
Thus, we next aimed to quantify the temporal dynamics underlying the distributions of gene expression and growth. We measured expression in cell lineages over many generations using a 'mother machine' microfluidic device [24] and time-lapse fluorescence microscopy. In this device, 'mother' cells are trapped at the top of one-ended chambers and maintained indefinitely in exponential phase through the addition of fresh media, allowing for multi-hour imaging of cell lineages. We used this to monitor gene expression in tens to hundreds of independent cell lineages for each reporter for at least 15 hours. To quantify gene expression over time, we used our deep learning based cell segmentation and tracking algorithm [25] to extract single-cell resolution data from the mother cell and its progeny.

We observed heterogeneity in gene expression for cells growing in these precisely controlled conditions, which was consistent with measurements acquired from bulk culture snapshots (Fig. S2). Long-term monitoring of expression revealed a diversity of phenotypes, with some genes that fluctuated at a low level while others were highly dynamic in their expression (Fig. 1C-E). We observed that variability within a distribution often arose from pulsatile dynamics, with single cells transitioning between low and high expression states. However, the timescales of these fluctuations were specific to each gene, and the pulses themselves ranged in intensity. For instance, *gadX*, which encodes a transcription factor that regulates ~34 genes in the acid resistance system, exhibits pulses of high amplitude and duration that persist well beyond the cell cycle length, as visible in multi-generation patterns of gene expression (Fig. 1C, Movie S1). In contrast, *recA*, which plays a central role in the processes of homologous recombination and SOS response, showed large amplitude pulses, but with shorter durations (Fig. 1D, Movie S2). Yet others, like *araC*, which encodes a transcription factor that regulates arabinose catabolism and transport, were more muted in their changes, and exhibited only mild, low amplitude fluctuations (Fig. 1E, Movie S3). Surveying 15 reporters, we observed a wide range of temporal gene expression profiles (Fig. 1F-G). Importantly,

46   the dynamic activity observed for many reporters occurred under constant growth conditions, where
47   parameters such as pH, temperature, and growth medium do not fluctuate.
48
Pulsing properties vary across the reporters
Next, we defined several metrics across which to assess dynamic behavior. We selected properties
related to pulses including their frequency, duration, and amplitude (Fig. 2A). Pulse frequencies
were broadly distributed, ranging from mean values of 0.25 pulses per hour (one pulse every four
hours) for *recA* to much less frequent conditions where our peak-finding algorithm rarely identified
pulses (Fig. 2B). The distributions themselves were also highly variable, with some genes
exhibiting tight distributions such that pulses always tended to occur at about the same frequency,
while others were broadly distributed. For instance, pulses in *bolA* tend to be fairly narrow in their
distribution while *evgA* pulses have a much less standardized frequency. Pulse durations typically
ranged from 0.5 - 3 hours, and again showed instances of precision (e.g. *recA*, *sulA*) in addition to
examples with widely variable durations (e.g. *evgA, purA, rob*). Pulse amplitudes were
predominantly small (within 25% of the mean, or 0.25x), but there were notable instances where
the distributions had long tails such that amplitudes extended well above this for a subset of the
pulses. For example, *gadX* and *recA* exhibit pulses that significantly deviate from the mean.
63
We used these data to look for general trends and relationships related to pulse properties. We found
that pulse frequency and amplitude were positively correlated for the genes we analyzed, while
frequency and duration were inversely related (Fig. 2C). To alleviate the potential concern that
these relationships were a result of the specific method by which we identified pulses, we repeated
the calculations under a range of different peak determination thresholds and found that our results
were not sensitive to the precise threshold definition (Fig. S3). Coefficients of variation from our
static snapshots were positively correlated with pulse frequency and amplitude, and inversely
correlated with duration (Fig. S4). These results on relationships between pulse properties suggest
that there may be some pulsing regimes that are uncommon in practice. For example, we did not
observe pulses with high frequency and low amplitude or long durations.
74
We next asked what characteristic timescales the gene expression dynamics exhibited by
calculating the autocorrelation of the fluorescence signal. In all cases, we observed monotonically
decreasing autocorrelation curves, ruling out the presence of regular period signals, such as
oscillatory behavior (Fig. S5). We calculated the half life associated with the autocorrelation curve
for each gene (Fig. 2D-E). If the fluorescent reporter levels decrease solely due to dilution resulting
from growth and division, the half life will equal the cell division time. Longer half lives can
indicate the presence of memory, for example due to regulatory networks that causes signals to
persist. We found that half lives were always greater than or equal to the cell division time,
consistent with the use of stable fluorescent reporters (Fig. 2E). Notably, we observed cases where
the average half life exceeded the cell division time by 2-3-fold, potentially indicative of memory
within the network. The half life of the autocorrelation curve was positively correlated with pulse
duration (Fig. S6), as longer duration pulses persist across generations. We verified that these
calculations performed on data from mother cells were consistent with correlations within the
lineage tree. Indeed, tracking fluorescence signals from mother to daughter to granddaughter cells
revealed a strong positive correlation that persisted across multiple generations (Fig. S7).
90
91   **Pulsatile expression dynamics are closely linked with growth**
92   We next asked whether growth rate contributes to the observed variability in expression levels.
93   While the growth rate of individual *E. coli* cells shows long-term stability during replicative aging
94   [24], it can exhibit noisy temporal behavior [2,8,19]. Patange *et al*. demonstrated that the growth rates
95   fluctuate, with pulses of slow growth that last for several generations and are anti-correlated with

pulses of activation of RpoS [8]. However, it is not clear whether this feature is specific to RpoS or if it is common to many genes. We asked whether the pulsatile dynamics we observed coincided with dynamic patterns in growth rate. To test this, we extracted the instantaneous growth rate, as quantified by the elongation rate, from cell lineages over time and compared them to fluorescence data for the different reporters.

In principle, fluorescence levels, which report underlying cellular properties such as those related to metabolism, can precede changes in growth. Alternatively, growth can drive changes in fluorescence since it affects dilution rates; regulatory links between growth rate and fluorescence are also possible (Fig. 3A). In both cases, these relationships could be positive or negative depending on the precise underlying mechanism.

To distinguish between these possibilities, we computed temporal cross-correlations between fluorescence and growth rate, $R_{fluo,gr}(\tau)$ (Fig. 3B). The cross-correlation measures how well the fluorescence and growth signals are correlated, when the growth signal is shifted by a time τ relative to the fluorescence signal. Cross-correlation curves for our reporters have characteristic shapes, where signals are uncorrelated at large positive and negative values of the time shift, τ. For intermediate values of τ we regularly observed peaks or valleys in the cross-correlation. We defined $R_{extreme}$ as the point on the cross-correlation with the largest absolute value, indicating the largest correlation between the fluorescence and growth rate signals. $R_{extreme}$ can be either positive or negative, depending on whether the signals are correlated or anti-correlated. We also defined $\tau_{extreme}$ as the time shift associated with $R_{extreme}$, where $\tau_{extreme} < 0$ when changes in fluorescence precede changes in growth and $\tau_{extreme} > 0$ when growth leads fluorescence.

Growth and expression are often anti-correlated with growth leading expression

We found examples where a strong anti-correlation between fluorescence and growth were visible, such as with the *gadX* reporter, and other cases where the signals were mildly positively correlated, such as with the *recA* reporter (Fig. 3C). To quantify this trend across many cell lineages, we calculated cross-correlations between growth and fluorescence for all reporters (Fig. 3D). In many cases we observed an anti-correlation with a positive time shift between growth rate and gene expression, indicating that the pulses of expression from these promoters were preceded by a decrease in growth rate. Although the accumulation of fluorescent proteins resulting from reduced dilution due to cell growth and division could potentially artificially generate this relationship, we observed only a modest anti-correlation for the majority of promoters, and no anti-correlation for *recA* and *sulA* (Fig. 3E). Thus, it is unlikely that these results are due solely to a reduced dilution rate.

Changes in growth rate never lagged changes in fluorescence for all reporters we measured ($\tau_{extreme} \geq 0$) (Fig. 3F). It is possible that fluorophore maturation times could systematically introduce a lag between actual promoter activity and read out of the fluorescent protein. Maturation times for our reporters are relatively rapid, with 50% of fluorescent proteins maturing within ~6 minutes [26], however reporter maturation times could systematically introduce a small positive shift in $\tau_{extreme}$. Overall, our measurements indicate that even cells growing exponentially under optimal conditions undergo episodes of slow growth that are largely followed by the pulses of stress-related genes, which frequently demonstrate an inverse relationship between growth rate and fluorescence (Fig 3G). In general, there was little relationship between pulse properties such as frequency, amplitude, and duration, and the properties related to growth rate and fluorescence correlations (Fig. S8), thus these features appear to be largely independent.

Gene expression and growth dynamics transiently protect cells against stress

A critical question is whether these changes in gene expression and growth have concrete implications for whether a single cell will survive or die following stress. In other words, are the dynamics we observed sufficient to provide meaningful phenotypic differences that cause a cell to evade or succumb to stress? Slow growth and induction of stress response genes have previously been reported as mechanisms that allow survival at the populational level [27–31]. This prompted us to investigate whether pulses in gene expression and growth influence the chances of survival under stress. Further, we asked whether survival was the result of a fortuitous condition of gene expression or slow growth upon stress introduction, or if a cell's past history was important.

For these studies, we focused on expression of *gadX*, as our measurements of single-cell gene expression demonstrate that this reporter is expressed in large pulses and has a strong anti-correlation with growth. We previously demonstrated that *gadX* is heterogeneously expressed in the absence of antibiotic stress and that expression levels correlate with longer survival times under constant carbenicillin exposure [7]; however, it is unclear how prior history or pulsing dynamics contribute to this phenotype. Here, we quantified how *gadX* expression preceding sudden exposure to a lethal antibiotic dose influences survival. To do so, we leveraged the mother machine device to rapidly switch input media and exposed exponentially dividing cells to a short pulse of ciprofloxacin, a fluoroquinolone drug widely used to treat bacterial infections. We treated cells with 2 µg/ml ciprofloxacin, which corresponds to 100x the minimum inhibitory concentration (MIC), for 35 minutes before switching back to growth medium without antibiotics. This experimental set up allowed us to monitor the dynamics of *gadX* expression and growth in single cells prior to antibiotic treatment while also recording the outcome of each cell lineage.

Within the same experiment, we observed instances where single cells were able to survive ciprofloxacin treatment (Fig. 4A, Movie S4) and also cases where treatment killed the cells (Fig. 4B, Movie S5). In addition to surviving and dying, we also observed a third category of outcomes where cells filamented. However, it was difficult to accurately assess whether these cells survived or died, as they were frequently swept out of the mother machine chamber and lost from the field of view. Thus, we focused our analysis on the surviving and dying cells because we could accurately assess their outcomes. By tracking gene expression history preceding antibiotic treatment, we observed pulses in fluorescence from the *gadX* reporter prior to ciprofloxacin addition in both the surviving and dying cells (Fig. 4C). However, cells that were in the midst of a pulse of expression were more likely to survive. In contrast, cells that exhibited expression pulses in the past, but had low expression at the time of ciprofloxacin treatment were more likely to die. Thus, cells which are in the fortuitous state of having an ongoing *gadX* pulse at the time of ciprofloxacin addition are more likely to exhibit transient stress tolerance.

We also looked at cellular outcome as a function of the growth rate. Consistent with our results showing anti-correlations between fluorescence and growth for *gadX* (Fig. 3D), we observed decreased growth rates in cells that survived, which were associated with increases in fluorescence signals (Fig. 4D). However, reduced growth rate alone does not appear to be sufficient for survival, as we observed cells that died which had similar growth rates to those that survived. Measurements of the instantaneous fluorescence and growth rate at t = 0, just prior to antibiotic addition, show that most surviving cells exhibited both elevated *gadX* expression and reduced growth rate immediately prior to ciprofloxacin treatment (Fig. 4E), and our findings indicate that the prior dynamics are what give rise to these conditions.

Neither growth rate nor *gadX* expression alone is expected to completely determine cellular outcome, as stress response is a complex process; instead, our results suggest that pulsing dynamics can bias the probability that a single cell will live or die. Cell-to-cell heterogeneity in *gadX*

expression is the result of pulsing dynamics, which produce diverse distributions of cells. Thus, we demonstrate that *gadX* pulsing can play a protective role against ciprofloxacin stress, demonstrating a clear link between single-cell dynamics and cell fate.

Discussion

Although cells can display substantial phenotypic heterogeneity in constant environmental conditions, the timescales over which gene expression levels and growth rate fluctuate and how this ultimately impacts function are often less clear. To provide insight into these questions, we focused on genes involved in stress response in *E. coli*. Our results demonstrate that distributions of gene expression levels and growth rates can originate from rich dynamic activity, where individual cells transition between low and high expression over time. Our findings expand previously published observations on pulsing dynamics [8,13,14,17,32] to a broader set of genes, suggesting that these time-varying differences in expression and growth may be more common than previously appreciated. These fluctuations occur under uniform, unstressed conditions and are observed across genes with diverse stress response roles. The specific properties of these fluctuations, including pulse frequency, duration, and amplitude vary significantly between individual genes. Notably, temporal changes in gene expression are also related to growth. In our measurements, we found many instances where pulses of high activity followed decreases in growth rate. However, this link is not universal, which indicates that this effect is not an unavoidable consequence of growth, and rather that the expression and growth relationship is specific to each gene. Finally, we asked whether gene expression dynamics have functional consequences, focusing on expression of *gadX*. We showed that a functional outcome of *gadX* pulsing and the corresponding decrease in growth rate is that cells are more likely to survive following sudden exposure to a lethal dose of ciprofloxacin.

Our results with *gadX* join a small number of other studies demonstrating examples of functional phenotypic heterogeneity. For example, the stochastic activation of diverse stress response genes, such as those encoding porins [33], the catalase-peroxidase KatG [34], and the multiple antibiotic resistance activator MarA [6,35,36] result in transient tolerance to exogenous stress. Other examples include biased partitioning of efflux pumps during cell division that results in differential antibiotic susceptibility [37] and heterogeneous induction of *gad* regulon genes during antibiotic treatment that cross-protects cells against subsequent acid stress [9].

Reduced growth rates can also enable cells to transiently resist stress, for example, by playing a major role in the formation of persister cells [28,38]. By tracking expression dynamics and growth rates simultaneously, we found many instances in which these metrics are inversely correlated over time. Interestingly, an analogous observation was recently reported by Patange *et al*. [8] that demonstrated that the stress response master regulator RpoS is expressed with pulsatile dynamics in exponentially dividing cells and is inversely correlated with growth rate. RpoS participates in the regulation of many genes characterized in our study including *bolA*, *dinB*, *evgA*, *gadX*, *oxyR*, and *rpoH* (Table S1), which could in principle indicate that the pulsatile RpoS expression is transmitted to several of its downstream genes. However, additional mechanisms might underlie this phenomenon, as some genes in our study with strong growth rate anti-correlations (e.g. *marA*) are not known to be regulated by RpoS. It is also noteworthy that, in addition to RpoS, at least one additional sigma factor, RpoH, was activated with pulsatile dynamics during exponential growth, a behavior that can potentially be propagated to >150 downstream genes that collectively control the cellular heat-shock response [39,40]. The pulsatile activity of these regulators might indicate that the mechanism by which sigma factors "time share" RNA polymerase complexes previously described in *B. subtilis* [41] might also be present in *E. coli*. Additionally, our results suggest that the pulses of activation of different sigma factors might be associated with different cellular growth statuses. Further

45 investigations can help to decouple the roles of growth rate and gene expression in enabling
46 survival.
47
In addition, it would be interesting to conduct measurements using pairs of reporters so that
expression pulses can be monitored simultaneously in the same cell. This is particularly interesting
to examine in relation to the growth rate, as we observed many instances of anti-correlation, but the
temporal properties of these pulses were distinct from each other. Further, future studies may be
necessary to identify the mechanism underlying the pulsing dynamics. In some instances this is
known, as in a recent study that identified LexA-induced heterogeneity as a key contributor to
pulsing in SOS response genes including *recA* [14], however other examples will require further
investigation. In addition, although we measure growth rate in our analysis, our findings do not
preclude the possibility that growth rate is a secondary effect that is downstream of the true
coordinating signal, such as metabolic state [42]. Experiments that use reporters to measure genes
involved in metabolism alongside the stress response reporters, or studies that carefully control the
nutrient composition could be used to investigate this possibility. It will also be interesting to
investigate how the information encoded in transcription factor dynamics is transmitted to
downstream genes, and emerging technologies that enable the manipulation of gene expression over
time could help to identify how these signals are propagated [43,44].
63
64 Our findings reveal that pulsatile dynamics in gene expression and growth serve as a mechanism
65 that cells can leverage to transiently resist stress. This dynamic behavior is spread across a broad
66 range of stress response genes, including heat-shock response, multi-drug resistance, and oxidative
67 stress and can enable subpopulations of cells to withstand temporary stress.
68
69 **Materials and Methods**
70
Strains and growth media
Strains containing reporter plasmids were sourced from the collection created by Zaslaver *et al.* [45],
unless otherwise noted. Briefly, each strain has a low copy number plasmid (SC101 origin)
containing the promoter sequence for the gene of interest upstream of the coding sequence for
gfpmut2 green fluorescent protein (GFP). The strain reporting *purA* was sourced from Rossi *et al.*
[7] and the strain reporting *marA* was sourced from El Meouche *et al.* [6], where the *marA* reporter has
intact MarR binding sites. The *purA* and *marA* reporters use a low copy number plasmid (SC101
origin) and the promoters of interest control the expression of cerulean cyan fluorescent protein
(CFP). In all cases, we used *E. coli* MG1655 as the strain background. Cells were grown in M9
medium (0.1 mM $CaCl_2$; 2 mM $MgSO_4$; 1X M9 salts) supplemented with 0.4% glucose, 0.2%
casamino acids, and 30 μg/ml kanamycin for plasmid maintenance. Media used in microfluidic
experiments was supplemented with 2 g/L F-127 Pluronic to prevent cell adhesion and growth
outside of the mother machine growth chambers.
84
Static fluorescence microscopy snapshots
Overnight cultures were diluted 1:100 and incubated at 37C with shaking for ~3 hours until mid-
exponential phase (OD_{600nm}= 1.0-1.3). 1 μL of cells were placed on 1.5% MGC (0.2% glycerol,
0.01% casamino acids, 0.15 μg/ml biotin, and 1.5 μM thiamine) low melting point agarose pads
and imaged at 100X on a Nikon Ti-E inverted fluorescence microscope. Three separate overnight
cultures were sampled and imaged for each reporter strain.
91
92 ***Mother machine microfluidic device***
93 The mother machine microfluidic master mold used was described previously in Ref. [25]. Briefly,
94 the master mold chip has 8 independent main feed channels where growth media flows in and out.

Each channel features 3,000 one-ended chambers (25 μm x 1.3 μm x 1.8 μm) where the mother cells are trapped. We made the microfluidic devices by pouring a degassed 10:1 mixture of dimethyl-siloxane monomer and curing agent (Sylgard 184 Silicone Elastomer Kit, Dow Corning) onto the wafer, which was then cured overnight at 65C. Individual chips were separated from the mold and a 0.75 mm biopsy punch was used to create inlets and outlets for each flow channel before the chip was plasma bonded to a glass slide.

Time-lapse microscopy movies

Overnight cultures were diluted 1:100 and incubated at 37C with shaking for ~3 hours until mid-exponential phase (OD_{600nm}= 1.0-1.3). Cells were concentrated by centrifugation (6,000xg for 2 min) and loaded into the microfluidic chip. Cells were seeded in the growth chambers by centrifugation at 6,000xg for 3 min. Media was supplied at a flow rate of 20 μL/min using a peristaltic pump. Cells were allowed to adapt to growth in the device for at least two hours before imaging. Time-lapse movies were acquired with a 100X oil objective on a Nikon Ti-E inverted fluorescence microscope equipped with a perfect focus system and a temperature control chamber that was set at 37C for the duration of the experiment. Images were acquired in phase contrast and epifluorescence illumination with a GFP or CFP filter every 5 min. At least two independent mother machine channels were imaged for each reporter; the total number of cell lineages tracked for each reporter is listed in Table S2.

Microscopy image processing and fluorescence and growth rate analysis

For static snapshots, cell images were segmented with the SuperSegger software [46]. For mother machine experiments, we performed automated cell segmentation and lineage tracking using the deep learning-based software DeLTA [25]. Unless otherwise noted, only data for the 'mother' cell trapped at the top of the growth chambers was used in quantitative analysis. We smoothed the fluorescence data using a moving average filter with a window of 5 frames.

To estimate growth rates, we calculated the numerical derivative as the difference in cell length between adjacent frames. When there is no cell division event, this derivative is calculated as (L_t – L_{t-1})/ Δt, where L_t is the length of the cell at time t and Δt is the time between frames (5 min). In cases where a cell division event occurs, this calculation is modified to (L_t + D_t – L_{t-1})/ Δt, where D_t is the length of the daughter at time t. After calculating the growth rate, we smoothed the data using a moving average filter with a window of 5 frames.

Cell division times are calculated as the time between division events.

Pulse identification

To allow comparison across all reporters, which have different expression levels and where imaging exposure times are different (Table S2), we normalized each time-series by its mean. Pulses of gene expression were identified using the built-in MATLAB function *findpeaks*, which identifies local maxima. We set the threshold for the minimum peak prominence at 0.15, unless otherwise noted.

Autocorrelation and cross-correlation calculations

We calculated the autocorrelation of the fluorescence signal—$R_{fluo,fluo}(\tau)$ where τ is the time shift—using the MATLAB function *xcov* with the unbiased normalization method. We then normalized the entire signal by the value of the autocorrelation at zero time lag, $R_{fluo,fluo}(0)$. The half life was calculated by taking the mean of all autocorrelation signals from individual cell lineages and identifying the time shift value where it crosses 0.5.

44 The cross-correlation between the fluorescence signal and the growth rate, $R_{fluo,gr}(\tau)$, was calculated
45 using the MATLAB function *xcov* with the unbiased normalization method. The signal was then
46 normalized by the square root of the product of the autocorrelation at zero time shift of the
47 fluorescence and growth rate signals: $(R_{fluo,fluo}(0)\, R_{gr,gr}(0))^{1/2}$.
48
49 $R_{extreme}$ is the value of $R_{fluo,gr}(\tau)$ with the largest absolute value. $\tau_{extreme}$ is the time lag value
50 associated with $R_{extreme}$.
51
Antibiotic survival assay
Mother machine experiments were initiated using the methods described above. During image
acquisition, cells were provided with fresh medium for 14 hours, followed by 35 minutes of
treatment with medium supplemented with 2 µg/ml ciprofloxacin, then fresh medium again for 16
hours. The outcomes of ciprofloxacin treatment for each lineage were manually scored as:
'survived,' 'died,' or 'filamented.' Cells were scored as 'surviving' if cell division was observed in
the chamber at the end of the full 16 hours after the second addition of fresh medium. Cells were
scored as 'dying' if growth ceased in the chamber at any point after antibiotic exposure.
'Filamented' cells were excluded from the analysis, as it was difficult to accurately assess their
outcome since they were frequently swept out of the chamber and field of view.
62
63 *Principal component analysis*
64 We conducted principal component analysis using six metrics for each of the 15 reporters: pulse
65 frequency, pulse duration, pulse amplitude, autocorrelation half life, $R_{extreme}$, and $\tau_{extreme}$. Principal
66 component analysis was performed using the MATLAB function *pca* with centered data and the
67 singular value decomposition algorithm.
68
69
70

References

1. Taniguchi, Y. *et al.* Quantifying E. coli proteome and transcriptome with single-molecule sensitivity in single cells. *Science* **329**, 533–538 (2010).

2. Kiviet, D. J. *et al.* Stochasticity of metabolism and growth at the single-cell level. *Nature* **514**, 376–379 (2014).

3. Silander, O. K. *et al.* A genome-wide analysis of promoter-mediated phenotypic noise in Escherichia coli. *PLoS Genet.* **8**, (2012).

4. Raj, A. & van Oudenaarden, A. Nature, Nurture, or Chance: Stochastic Gene Expression and Its Consequences. *Cell* **135**, 216–226 (2008).

5. Bar-Even, A. *et al.* Noise in protein expression scales with natural protein abundance. *Nat. Genet.* **38**, 636–643 (2006).

6. El Meouche, I., Siu, Y. & Dunlop, M. J. Stochastic expression of a multiple antibiotic resistance activator confers transient resistance in single cells. *Sci. Rep.* **6**, 1–9 (2016).

7. Rossi, N. A., El Meouche, I. & Dunlop, M. J. Forecasting cell fate during antibiotic exposure using stochastic gene expression. *Commun. Biol.* **2**, 1–7 (2019).

8. Patange, O. *et al.* Escherichia coli can survive stress by noisy growth modulation. *Nat. Commun.* **9**, (2018).

9. Mitosch, K., Rieckh, G. & Bollenbach, T. Noisy Response to Antibiotic Stress Predicts Subsequent Single-Cell Survival in an Acidic Environment. *Cell Syst.* **4**, 393-403.e5 (2017).

10. Uphoff, S. *et al.* Stochastic activation of a DNA damage response causes cell-to-cell mutation rate variation. *Science* **351**, 1094–1098 (2016).

11. Uphoff, S. Real-time dynamics of mutagenesis reveal the chronology of DNA repair and damage tolerance responses in single cells. *Proc. Natl. Acad. Sci.* **115**, E6516–E6525 (2018).

12. El Meouche, I. & Dunlop, M. J. Heterogeneity in efflux pump expression predisposes antibiotic-resistant cells to mutation. *Science* **362**, 686–690 (2018).

13. Levine, J. H., Lin, Y. & Elowitz, M. B. Functional roles of pulsing in genetic circuits. *Science* **342**, 1193–1200 (2013).

14. Jones, E. C. . & Uphoff, S. Imaging LexA degradation in cells explains regulatory mechanisms and heterogeneity of the SOS response. *bioRxiv* (2020). doi:10.1101/2020.07.07.191791

15. Süel, G. M., Kulkarni, R. P., Dworkin, J., Jordi, G.-O. & Elowitz, M. B. Tunability and noise dependence in differentiation dynamics. *Science* **315**, 1716–1719 (2007).

16. Maamar, H., Raj, A. & Dubnau, D. Noise in gene expression determines cell fate in Bacillus subtilis. *Science* **317**, (2007).

17. Kim, J. M., Garcia-Alcala, M., Balleza, E. & Cluzel, P. Stochastic transcriptional pulses orchestrate flagellar biosynthesis in Escherichia coli. *Sci. Adv.* **6**, eaax0947 (2020).

18. Logsdon, M. M. & Aldridge, B. B. Stable regulation of cell cycle events in mycobacteria: Insights from inherently heterogeneous bacterial populations. *Frontiers in Microbiology* **9**, 514 (2018).

19. Tanouchi, Y. *et al.* A noisy linear map underlies oscillations in cell size and gene expression in bacteria. *Nature* **523**, 357–359 (2015).

20. Biselli, E., Schink, S. J. & Gerland, U. Slower growth of *Escherichia coli* leads to longer survival in carbon starvation due to a decrease in the maintenance rate. *Mol. Syst. Biol.* **16**, (2020).

21. Narula, J. *et al.* Slowdown of growth controls cellular differentiation. *Mol. Syst. Biol.* **12**, 871 (2016).

22. Aldridge, B. B. *et al.* Asymmetry and aging of mycobacterial cells lead to variable growth and antibiotic susceptibility. *Science* **335**, 100–104 (2012).

23. Lee, A. J. *et al.* Robust, linear correlations between growth rates and β-lactam–mediated
lysis rates. *Proc. Natl. Acad. Sci.* **115**, 4069–4074 (2018).
24. Wang, P. *et al.* Robust growth of Escherichia coli. *Curr. Biol.* **20**, 1099–1103 (2010).
25. Lugagne, J. B., Lin, H. & Dunlop, M. J. DeLTA: Automated cell segmentation, tracking,
and lineage reconstruction using deep learning. *PLoS Comput. Biol.* **16**, e1007673 (2020).
26. Balleza, E., Kim, J. M. & Cluzel, P. Systematic characterization of maturation time of
fluorescent proteins in living cells. *Nat. Methods* **15**, 47 (2018).
27. Tuomanen, E., Cozens, R., Tosch, W., Zak, O. & Tomasz, A. The rate of killing of
Escherichia coli by β-lactam antibiotics is strictly proportional to the rate of bacterial
growth. *J. Gen. Microbiol.* **132**, 1297–1304 (1986).
28. Pontes, M. H. & Groisman, E. A. Slow growth determines nonheritable antibiotic
resistance in Salmonella enterica. *Sci. Signal.* **12**, 1–11 (2019).
29. Evans, D. J., Allison, D. G., Brown, M. R. W. & Gilbert, P. Susceptibility of Pseudomonas
aeruginosa and Escherichia coli biofilms towards ciprofloxacin: Effect of specific growth
rate. *J. Antimicrob. Chemother.* **27**, 177–184 (1991).
30. Yelin, I. & Kishony, R. Antibiotic Resistance. *Cell* **172**, (2018).
31. Palmer, A. C. & Kishony, R. Opposing effects of target overexpression reveal drug
mechanisms. *Nat. Commun.* (2015). doi:10.1038/ncomms5296.Opposing
32. Purvis, J. E. & Lahav, G. Encoding and decoding cellular information through signaling
dynamics. *Cell* **152**, 945–956 (2013).
33. Sánchez-Romero, M. A. & Casadesús, J. Contribution of phenotypic heterogeneity to
adaptive antibiotic resistance. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 355–360 (2014).
34. Wakamoto, Y. *et al.* Dynamic persistence of antibiotic-stressed mycobacteria. *Science* **339**,
91–96 (2013).
35. Rossi, N. A. & Dunlop, M. J. Customized Regulation of Diverse Stress Response Genes by
the Multiple Antibiotic Resistance Activator MarA. *PLoS Comput. Biol.* **13**, e1005310
(2017).
36. Garcia-Bernardo, J. & Dunlop, M. J. Tunable stochastic pulsing in the Escherichia coli
multiple antibiotic resistance network from interlinked positive and negative feedback
loops. *PLoS Comput. Biol.* **9**, e1003229-11 (2013).
37. Bergmiller, T. *et al.* Biased partitioning of the multidrug efflux pump AcrAB-TolC
underlies long-lived phenotypic heterogeneity. *Science* **356**, 311–315 (2017).
38. Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L. & Leibler, S. Bacterial persistence as a
phenotypic switch. *Science* **305**, 1622–1625 (2004).
39. Arsène, F., Tomoyasu, T. & Bukau, B. The heat shock response of Escherichia coli. *Int. J.*
Food Microbiol. **55**, 3–9 (2000).
40. Yura, T. Regulation of the heat shock response in Escherichia coli: History and
perspectives. *Genes Genet. Syst.* **94**, 103–108 (2019).
41. Park, J. *et al.* Molecular Time Sharing through Dynamic Pulsing in Single Cells. *Cell Syst.*
6, 216-229.e15 (2018).
42. Lopatkin, A. J. *et al.* Bacterial metabolic state more accurately predicts antibiotic lethality
than growth rate. *Nature Microbiology* **4**, 2109–2117 (2019).
43. Lugagne, J. B. & Dunlop, M. J. Cell-machine interfaces for characterizing gene regulatory
network dynamics. *Curr. Opin. Syst. Biol.* **14**, 1–8 (2019).
44. Sampaio, N. M. V. & Dunlop, M. J. Functional roles of microbial cell-to-cell heterogeneity
and emerging technologies for analysis and control. *Curr. Opin. Microbiol.* **57**, 87–94
(2020).
45. Zaslaver, A. *et al.* A comprehensive library of fluorescent transcriptional reporters for
Escherichia coli. *Nat. Methods* **3**, 623–628 (2006).
46. Stylianidou, S., Brennan, C., Nissen, S. B., Kuwada, N. J. & Wiggins, P. A. SuperSegger:

robust image segmentation, analysis and lineage tracking of bacterial cells. *Mol. Microbiol.*
102, 690–700 (2016).

Acknowledgments: We thank Imane El Meouche, Nicholas A. Rossi, Tiebin Wang, and Wilson
Wong for helpful discussions; Ahmad (Mo) Khalil and Razan Alnahhas provided input on the
manuscript. **Funding:** This work was supported by the National Institutes of Health grants
R01AI102922 and R21AI137843. **Author contributions:** NMVS and MJD conceptualized the
study. NMVS performed experiments. NMVS, CMB, JBL, and MJD performed data analysis.
NMVS and MJD wrote the manuscript, with input from CMB and JBL. MJD supervised the study
and acquired funding. **Competing interests:** All authors declare that they have no competing
interest.

86
87

Figures

89
90
Figure 1. Cell-to-cell heterogeneity and temporal dynamics of gene expression. (A) Histograms
of fluorescence values show cell-to-cell variation. Values are presented normalized relative to their
means to allow for comparison across reporters. The same data without normalization are shown in
Fig. S1. Corresponding coefficient of variation (CV) values obtained from snapshot images of cells
grown in bulk cultures are listed in the figure. The skewness (skew) of the distribution, which is
positive for distributions that are right-skewed, are also listed. Insets show phase contrast and
fluorescence images for representative strains. Functional classes are listed for each gene. **(B)**
Schematic representation of fluorescence signal over time originating from single cells with slow
or fast dynamics of gene expression that result in identical fluorescence distributions for the final
timepoint. **(C-E)** Representative kymographs of cells containing transcriptional fluorescent
reporters for **(C)** *gadX*, **(D)** *recA*, and **(E)** *araC*. In all cases, fluorescence values are normalized to
the mean to allow comparisons across the reporters. Scale bar, 5 μm. **(F)** Single-cell measurements
of green fluorescent protein expression (GFP) over time for the *gadX* reporter. Colored heat maps

summarize the time-series data. Data from 25 cells are shown, however this represents a subset of
the time-series data. **(G)** Heat maps summarizing the temporal dynamics of gene expression for all
reporters. To allow comparison across reporters, fluorescence traces are normalized by their mean
as in (F).
08
09

10
11
Figure 2. Pulsing dynamics vary across the reporters. (A) Schematic representation of pulse
characteristics. **(B)** Distributions of pulse frequency, duration, and amplitude for all reporters. **(C)**
Correlations between pulse properties. Each circle represents the median values for the reporter
analyzed. **(D)** Autocorrelation of the fluorescence signal for independent cell lineages with the *gadX*
reporter. Half life is defined as the value of the time shift τ when the mean of the autocorrelation
curve crosses 0.5. **(E)** Cell division time and half life for each of the reporters. Error bars show
standard deviation around the mean.
19
20

21
Figure 3. Growth and gene expression dynamics are often anti-correlated. (A) Two possible
models for the relationship between fluorescent reporter levels and growth. **(B)** Schematic of cross-
correlation function $R_{fluo,gr}(\tau)$ indicating points corresponding to $R_{extreme}$ and $\tau_{extreme}$ (top). Temporal
patterns of the fluorescence and growth rate and their impact on $R_{extreme}$ and $\tau_{extreme}$ (bottom). **(C)**
Representative fluorescence (green) and growth rate (magenta) values over time for a single mother
cell with a reporter for *gadX* (left) or *recA* (right). Black dots at the top of the figure indicate cell
division events. Data are smoothed with a moving average filter with window of 5 frames. **(D)**
Cross-correlations for all reporters. Shaded region represents standard deviation about the mean.
(E) $R_{extreme}$ and **(F)** $\tau_{extreme}$ values for each reporter. **(G)** $R_{extreme}$ as a function of $\tau_{extreme}$.

Figure 4. Cell-to-cell differences in *gadX* expression and growth influence survival outcomes after antibiotic exposure. (A) Representative kymograph of a cell lineage that survives and **(B)** dies after antibiotic treatment. Images are from nearby chambers within the same microfluidic chip. Shaded area in red indicates 35 minute period of ciprofloxacin treatment. **(C)** Single-cell *gadX* expression and **(D)** growth rate for the period preceding ciprofloxacin treatment. Cell numbers in (C) and (D) correspond to the same cells. Time-series are sorted by the mean value of the fluorescence signal for t < 0, from high to low. **(E)** Individual cells that survived (green) or died (black) following ciprofloxacin treatment plotted as a function of growth rate and fluorescence at t = 0.

SUPPLEMENTARY MATERIALS

Supplementary Figures

Figure S1. Histograms of fluorescence distributions of single cells. Microscopy exposure times vary between reporters to optimize detection of fluorescence from each reporter and are indicated in the figure.

Figure S2. Histograms displaying distributions of mean normalized fluorescence values for single cells imaged in snapshots of agarose pads (blue) or in the mother machine device (red).

91
92
93 **Figure S3.** R^2 correlation coefficient for pulse frequency and amplitude (left) and pulse frequency
94 and duration (right) calculated using different thresholds for the peak prominence. Calculations in
95 Fig. 2 use a threshold of 0.15, which is indicated with a dashed line.
96
97

Figure S4. Median pulse frequency, amplitude, and duration of pulses, as a function of coefficient of variation for each of the reporters.

Figure S5. Autocorrelation of the fluorescence signal for all reporters. Shaded area shows standard deviation around the mean.

34
35
36 **Figure S6.** Pulse duration versus half life for each of the reporters.
37
38

Figure S7. Correlation coefficient between mother cells mean fluorescence signal and its offspring. "Generations from mother" = 1 corresponds to a mother-daughter comparison, 2 is a mother-granddaughter comparison, and 3 is a mother-great granddaughter comparison.

Figure S8. Principal component analysis performed on traits related to gene expression dynamics for each reporter. **(A)** Percent variability explained by each principal component. **(B)** Biplot depicting first two principal components and the locations of each reporter. Vectors that are perpendicular are uncorrelated; vectors in opposite directions are anti-correlated.

Supplementary Tables

Table S1. Summary of genes characterized in this study

Gene	Sigma Factor	Function	# of Regulatees
araC	RpoD	Arabinose catabolism	14
bolA	RpoD, RpoS	Cell morphology and protection, general stress response	5
dinB	RpoD, RpoS	SOS response, encodes DNA polymerase IV	0
evgA	RpoD, RpoS	Multi-drug resistance	19
gadW	RpoS	Acid resistance regulon	15
gadX	RpoD, RpoS	Acid resistance regulon	34
marA	RpoD	Multiple antibiotic resistance	45
metJ	RpoD	Methionine biosynthesis	15
oxyR	RpoD, RpoS	Oxidative stress	42
phoP	RpoH	Adaptation to low Mg2+, control of acid resistance genes	56
purA	RpoD	*de novo* synthesis of AMP	0
recA	RpoD	DNA recombination / repair protein, SOS response.	0
rob	Unknown	Resistance to antibiotics, organic solvents, heavy metals	26
rpoH	RpoD, RpoN, RpoE, RpoS	Sigma32. Response to heat shock during log-phase growth	174
sulA	RpoD	SOS response cell division inhibitor	0

78   **Table S2.** Number of cell lineages tracked for each reporter and exposure time for the mother
79   machine time-lapse experiments
80

Gene	# cell lineages tracked	Exposure time
araC	71	150
bolA	72	300
dinB	88	300
evgA	72	30
gadW	107	10
gadX	142	30
marA	450	50
metJ	71	150
oxyR	71	150
phoP	72	150
purA	101	50
recA	153	300
rob	108	10
rpoH	72	30
sulA	336	300

81
82
83

Supplementary Movie Captions

Movie S1. Time-lapse movie of a representative mother machine chamber containing cells with the *gadX* reporter. Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.

Movie S2. Time-lapse movie of a representative mother machine chamber containing cells with the *recA* reporter. Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.

Movie S3. Time-lapse movie of a representative mother machine chamber containing cells with the *araC* reporter. Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.

Movie S4. Surviving cell in a time-lapse movie where cells containing the *gadX* reporter are exposed to ciprofloxacin (period of treatment indicated with red bar at top of the image). Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.

Movie S5. Dying cell in a time-lapse movie where cells containing the *gadX* reporter are exposed to ciprofloxacin (period of treatment indicated with red bar at top of the image). Phase contrast (gray) and fluorescence channel (green) are shown. Time is displayed as HH:MM. Scale bar, 2 μm.