# Contextual effects in sensorimotor adaptation adhere to associative learning rules

Guy Avraham*[1,2], Jordan A. Taylor[3], Assaf Breska[1,2,4], Richard B. Ivry[1,2], Samuel D. McDougle[5]

1. Department of Psychology, University of California, Berkeley, Berkeley, CA 94720, USA

2. Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, CA 94720, USA

3. Department of Psychology, Princeton University, Princeton, NJ 08544, USA

4. Max Planck Institute for Biological Cybernetics, Tübingen, Germany

5. Department of Psychology, Yale University, New Haven, CT 06520, USA

* Corresponding author

E-mail: guyavraham@berkeley.edu

Keywords: Sensorimotor adaptation; associative learning; cerebellum; forward model; context

# ABSTRACT

Traditional associative learning tasks focus on the formation of associations between salient events and arbitrary stimuli that predict those events. This is exemplified in cerebellar-dependent delay eyeblink conditioning, where arbitrary cues such as a light or tone act as conditioning stimuli that predict aversive sensations at the cornea. Here we ask if a similar framework could be applied to another type of cerebellar-dependent sensorimotor learning – sensorimotor adaptation. Models of sensorimotor adaptation posit that the introduction of an environmental perturbation results in an error signal that is used to update an internal model of a sensorimotor map for motor planning. Here we take a step towards an integrative account of these two forms of cerebellar-dependent learning, examining the relevance of core concepts from associative learning for sensorimotor adaptation. Using a visuomotor adaptation reaching task, we paired movement-related feedback with neutral auditory or visual contextual cues that served as conditioning stimuli (CSs). Trial-by trial changes in feedforward movement kinematics exhibited three key signatures of associative learning: Differential conditioning, sensitivity to the CS-US interval, and compound conditioning. Moreover, after compound conditioning, a robust negative correlation was observed between responses to the two elemental CSs of the compound (i.e., overshadowing), consistent with the additivity principle posited by theories of associative learning. The existence of associative learning effects in sensorimotor adaptation provides a proof-of-concept for linking cerebellar-dependent learning paradigms within a common theoretical framework.

## IMPACT STATEMENT

Core associative learning phenomena that are observed in studies of eyeblink conditioning are also observed in sensorimotor adaptation, pointing to a common framework for these distinct cerebellar-dependent motor learning processes.

**INTRODUCTION**

Sensorimotor adaptation refers to the gradual adjustment of movements in response to changes in the environment or body. In laboratory adaptation tasks, the introduction of perturbed sensory feedback results in a sensorimotor prediction error. This error signal is used to update a model of an internal sensorimotor mapping, thus ensuring that the sensorimotor system remains well-calibrated (Shadmehr and Krakauer, 2008; Wolpert et al., 1995; Wolpert and Ghahramani, 2000). The integrity of the cerebellum is essential for this form of error-based learning (Donchin et al., 2011; Izawa et al., 2012; Popa et al., 2016; Schlerf et al., 2012), and is thought to play a key role in generating predictions of future states given the current sensory context.

Different contexts may call for the use of different sensorimotor mappings. For instance, distinct motor memories should be retrieved when a skilled tennis player prepares to hit a service return on a clay court versus a grass court. A topic of considerable discussion in the adaptation literature centers on the constraints underlying effective contextual cues (Addou et al., 2011, 2011; Heald et al., 2021; Dawidowicz et al., 2021; Heald et al., 2018; Howard et al., 2012, 2013, 2015; Karniel and Mussa-Ivaldi, 2002; Osu et al., 2004; Schween et al., 2019; Sheahan et al., 2016, 2018; Forano and Franklin, 2020; Forano et al., 2021). Contextual cues are highly effective when the cue is directly relevant to properties of the movement. For example, people can simultaneously adapt to competing sensorimotor perturbations if the current context is established by the movement segment that precedes or follows the perturbed segment of a reach (Howard et al., 2012, 2015; Sheahan et al., 2016). In these situations, the contextual cues are hypothesized to be effective because the cues are incorporated into the motor plan (Howard et al., 2012, 2013).

Less clear is the efficacy of arbitrary contextual cues, ones that are not directly related to movement. In general, arbitrary cues do not appear to be effective for defining distinct motor memories. For example, participants show large interference between opposing perturbations that are signaled by color cues (Howard et al., 2012, 2013; Gandolfo et al., 1996; Forano et al.,

2021). This interference can be reduced with multiple days of practice, although even after 10

days of training, residual interference remains on trials immediately following a switch in context

(Addou et al., 2011; see also Osu et al., 2004).

The ineffectiveness of arbitrary contextual cues in shaping sensorimotor adaptation is

surprising when one considers another popular model task of cerebellar-dependent sensorimotor

learning – delay eyeblink conditioning. In this form of classical conditioning, a sensory cue

(conditioned stimulus, CS, e.g., a tone) is repeatedly paired with an aversive event (unconditioned

stimulus, US, e.g., an air puff to the cornea). By itself, the aversive event naturally causes an

immediate unconditioned response (UR, e.g., an eye blink in response to the air puff). After just

a short training period with a reliable CS, organisms as diverse as turtles and humans produce

an adaptive conditioned response (CR, e.g., an eye blink that anticipates the aversive air puff).

Notably, there is little constraint on the features of the predictive cues: The conditioned response

can be readily acquired in response to arbitrary conditioned stimuli, such as a tone or a light flash.

However, there is an important temporal constraint, at least with respect to cerebellar-dependent

eyeblink conditioning: The cue must appear before the aversive event and the two stimuli must

be in close temporal proximity (Schneiderman and Gormezano, 1964; Smith et al., 1969).

In the present study, we set out to re-evaluate the efficacy of arbitrary cues in sensorimotor

adaptation, drawing on key theoretical concepts and design features employed in studies of delay

eyeblink conditioning. Using a variant of a visuomotor adaptation task, we show that arbitrary

stimuli can act as effective contextual cues when a strict temporal relationship is established

between those cues and their associated movement-related sensory outcomes. We then take this

result a step further by showing that the effect of these arbitrary cues adheres to an established

core principle of associative learning – cue additivity. The parallels between adaptation and

conditioning observed in our results lay the empirical groundwork for a more parsimonious model

of cerebellar-dependent sensorimotor learning.

**RESULTS**

Our primary goal in the present study was to determine if arbitrary stimuli can serve as effective contextual cues in sensorimotor adaptation. Participants reached from a start location to a target with movement feedback provided by a cursor (Fig. 1A). In typical adaptation tasks, the onset of the target serves as the imperative for movement initiation. When considered through the lens of classical conditioning, the target appearance could be viewed as a conditioned stimulus (CS) given that it is presented just before the sensory feedback (the unconditioned stimulus, US) resulting from the movement (Fig. 1B). To test the efficacy of arbitrary cues as conditioned stimuli, the onset of the tone or light served as the imperative, with the target visible at its fixed location throughout the experimental session (Fig. 1B). For optimal delay eyeblink conditioning, the interval between the CS and the US is relatively narrow, on the order of 100-500 ms (Schneiderman and Gormezano, 1964; Smith et al., 1969). By using a neutral stimulus to cue movement initiation, we imposed a tight temporal link between the CS and the US, requiring the participant to initiate the movement within 400 ms of the onset of the imperative. Participants complied with this requirement, exhibiting rapid response times ([Mean ± STD]: 293±49.0 ms).

A common perturbation technique to elicit sensorimotor adaptation is to introduce a visuomotor rotation, where the movement path of the visual feedback cursor is rotated (about the origin) with respect to the hand's path. This drives an adaptation process in which the movement heading angle shifts across trials in the direction opposite to the rotated cursor, thus offsetting the error (Krakauer et al., 2000). To continue with our conditioning analogy, the conditioned response (CR) would be the movement heading angle elicited by a CS, expressed as a change in heading angle following the pairing of that CS with the visual error, the US (Fig. 1B). We note here that the unconditioned response (UR) in an adaptation task could correspond to an immediate movement correction in response to the perturbation; that is, an "online" correction. Given that we

required fast movements to minimize these corrections, this potential form of UR is not observed

in our design.

**Figure 1. Task design.**

(A) Reaching task. Participants reach from a start location (white circle) to a target (blue dot). Online

feedback is provided by a cursor (white dot). The target is displayed in a fixed location for the entire

experiment (location varied across participants). After a random delay, a tone or a light (white frame) is

presented, serving as the movement imperative. The participant is instructed to move directly to the target.

The cue persists until the radial position of the hand reaches the target distance. The hand is not visible

during the reach; instead, feedback is provided by a cursor. (B) Components and timing of the reaching

task described as conditioning events. The tone and light imperatives serve as conditioned stimuli (CS) and

the cursor feedback is the unconditioned stimulus (US). In our experiments, the radial position of the cursor

is aligned with the hand, but the angular position is fixed ("clamped"), appearing either 15° away from the

target (resulting in an error signal) when paired with one of the CS's (CS+) or at the target when paired with

the other CS (CS-). After repeated pairings of the CS+ and US, a conditioned response (CR) develops,

leading to an adaptive feedforward movement response in a direction opposite to the rotated feedback.

Note that we include a UR in the schematic, what we assume is an on-line, automatic corrective response

to the visual error. However, because participants are instructed to move fast, the UR is negligible.

Rather than rotate the position of the feedback cursor with respect to the actual hand

position, we opted to use a visual "clamp" where the cursor followed an invariant path with respect

to the target (Morehead et al., 2017; Shmuelof et al., 2012). In contrast to traditional movement-

contingent feedback, the clamp method eliminates confounding effects that can come about from

strategic processes (McDougle et al., 2016; Kim et al., 2020) and the size of the error can be

controlled on every trial. The participant was fully informed of this manipulation and was instructed

to ignore the task-irrelevant cursor and always reach straight to the target. Despite these

instructions, participants' reach angles gradually shift in the direction opposite the error clamp,

showing the key signatures of implicit sensorimotor adaptation (Morehead et al., 2017; Kim et al.,

2018; Tsay et al., 2020a, 2021; Avraham et al., 2021; Poh et al., 2021; Vandevoorde and Orban

de Xivry, 2019; Yin and Wei, 2020).

*Sensorimotor adaptation is modulated by arbitrary sensory cues*

In a standard differential conditioning design, one CS is paired with the US (CS+) and another CS

is presented without the US (CS-). Thus, only the CS+ should become associated with the US

and result in a CR. To implement this in Experiment 1, we used two arbitrary cues for the CSs –

a tone or a light cue (Figs. 1A, 2A). For the CS+ condition, the feedback cursor followed a clamped

path that was rotated from the target by 15°. Rather than eliminate the US on CS- trials, we used

a 0° clamp in which the feedback cursor always moved directly to the target. Thus, there was

always a US on each trial, with the 15° US creating an error signal (paired with the CS+) and the

0° US signaling the absence of error (paired with the CS-).

During a 600-trial acquisition phase, CS+ and CS- trials were randomly interleaved.

Participants exhibited a marked change in movement direction during this phase, reaching an

asymptote of ~15° (Fig. 2B). The observed rapid adaptation is consistent with previous adaptation

studies, particularly those in which the target appears at a single fixed location (Bond and Taylor,

2015; Day et al., 2016; McDougle et al., 2015, 2017; Poh et al., 2021).

**Figure 2. Experiment 1: Differential Conditioning.**

(A) Experimental protocol. During acquisition (white background), a 15° clamp (CW/CCW, counterbalanced

across participants) was associated with CS+ (e.g., a tone) and a 0° clamp with CS- (e.g., a light;

counterbalancing the associations with the tone and light across participants). During the probe phase (gray

background), the CS+ and CS- were presented without feedback. Throughout the entire experiment, CS+

and CS- trials were randomly interleaved. (B) Mean heading angle (N=16) as a function of trial number.

Clamped feedback was presented on all trials during the acquisition phase (white background) and absent

on all trials in the probe phase (gray background) phases. Shaded region represents SEM. (C, D)

Experimental results for trial-by-trial change (Δ) in heading angle (mean ± SEM) during the acquisition (*C*)

and probe (*D*) phases. Left panels present analysis results for an adaptation effect (main effect of trial *n-1*,

dark vs light blue) and a Pavlovian effect (main effect of the presented CS on the current trial *n*, filled vs

empty bars). The black outlined bar and violin plot (right panel) present the Pavlovian effect, i.e., the

subtraction of heading angle changes between CS+ and CS- trials (mean and 95% CI). (E, F) Rescorla-

Wagner model simulation results during the acquisition (*E*) and probe (*F*) phases are consistent with the

experimental results. Dots and thin lines represent individual participants.

The main analysis centered on trial-by-trial changes in heading angle. The change in

heading angle from trial *n-1* to trial *n* is normally dictated by the feedback experienced in trial *n-*

*1*. Thus, following experience with an error on CS+ trials, participants should show increased

adaptation (a positive change in heading angle), and following no error on CS- trials, decreased

adaptation (i.e., extinction). We refer to these trial-by-trial changes as the "adaptation effect", the

standard measure of learning in sensorimotor adaptation tasks. There was a robust adaptation

effect (Fig. 2C, left panel): Trial-by-trial changes in reaching direction (Δ heading angle) were

significantly affected by the CS presented on the previous trial [$F(1,15)=13.4$, $p=0.002$,

$BF_{10}=41.45$, $\eta_p^2=0.47$] such that the change in heading angle was larger after CS+ trials compared

to after CS- trials (Mean difference: 1.50°, 95% CI: [0.70° 2.30°]). That is, the error occurring on

CS+ trials resulted in learning that carried over to the next trial, whereas the absence of an error

on a previous CS- trial resulted in a relative reversion to baseline (extinction). This is the canonical

signature of incremental sensorimotor adaptation.

Importantly, the conditioning framework makes a critical prediction: The CS+ and CS-

should differentially modulate the hand's heading angle on trial $n$ itself. That is, the presentation

of the light or tone should produce a CR associated with that cue, leading to a difference in

heading angle on trials with different cues that is independent of the feedback received on the

previous trial. We refer to this as the "Pavlovian effect." As shown in Figure 2C (both panels), the

results revealed clear Pavlovian effects during the acquisition phase: The heading angle

increased in the direction of adaptation on CS+ trials (i.e., a positive change in heading angle

relative to previous trials) and decreased on CS- trials {Mean difference: 0.82°, 95% CI: [0.13°

1.51°]; $F(1,15)=5.37$, $p=0.035$, $BF_{10}=2.89$, $\eta_p^2=0.26$}. This effect provides a novel demonstration

that arbitrary sensory cues can lawfully influence implicit sensorimotor adaptation.

We also observed an interaction between CS identity on trials $n-1$ and $n$ [$F(1,15)=4.70$,

$p=0.047$, $BF_{10}=2.21$, $\eta_p^2=0.24$]. That is, the difference between CS+ and CS- was larger on trials

following a CS- (Mean: 1.05°, 95% CI: [0.20° 1.89°]) compared to trials following a CS+ (0.58°, [-

0.13° 1.30°]). This interaction effect may reflect an asymmetry between the rate of the acquisition

and extinction processes once CS-US associations are established. That is, the state following a

CS+ trial may be closer to its asymptotic limit than after a CS- trial and is thus more limited in its

potential for further change.

We note that the visual feedback was different on CS+ and CS- trials, with the cursor

deviating from the target in the former and moving in a straight line to the target in the latter. This

raises the possibility that the heading angle differences on CS+ and CS- trials could be affected

by online feedback corrections. This explanation is unlikely given that the movements were quite

rapid (mean movement time ± STD, 103±31.2 ms). To assess the online correction hypothesis

more directly, we calculated the difference in heading angle 50 ms after movement initiation and

at the radial distance of the target. There was no overall change in heading angle between the

time points (-0.32°±2.92°), and no significant difference between CS+ and CS- trials on this metric

{Mean difference: -0.04°, 95% CI: [-0.19° 0.12°]; t(15)=-0.47, p=0.640, $BF_{10}$=0.28, $d$=-0.12}. Thus,

our results appear to pertain exclusively to feedforward learning.

Following the acquisition phase, participants completed a probe phase in which CS+ and

CS- trials were randomly presented in the absence of any visual feedback (no US). This phase

provides a clean test for associative learning effects since it removes trial-by-trial effects that arise

from the differential feedback given during CS+ and CS- acquisition trials. Here too we observed

a clear Pavlovian effect: Although there was an overall decrease in heading angle across the

probe phase (i.e., a partial washout of adaptation, Fig. 2B), there was a significant main effect of

the CS presented on trial $n$ {1.06°, [0.53° 1.59°]; F(1,15)=15.45, p=0.001, $BF_{10}$=72.15, $\eta_p^2$=0.51}

(Fig. 2D), with a relative increase in heading angle on CS+ trials and a decrease on CS- trials.

Despite the absence of feedback in the probe phase, the change in heading angle on trial

n (CS+ or CS-) should also depend on the CS presented on trial n-1. After an association is

formed, the motor state, namely the actual heading angle, fluctuates as a function of the presented

CS, decreasing on CS- trials (due to extinction) and increasing on CS+ trials (due to conditioning).

As such, the heading angle on a CS+ trial should increase more if it follows a CS- trial than if it

follows a CS+ trial. Similarly, the heading angle on a CS- trial should decrease more if it follows

a CS+ trial than if it follows a CS- trial. While the data appeared to follow this pattern, there was

no main effect of the trial $n$-$1$ CS {-0.44°, [-1.11° 0.24°]; F(1,15)=1.61, p=0.224, $BF_{10}$=0.56,

$\eta_p^2$=0.10}. This may be because, statistically, the main effect of trial $n$-1 considers consecutive

trials with a repeated CS (CS+ that follows CS+, and CS- that follows CS-) where the change in

heading angle is minimal. Crucially, the Pavlovian effect (the difference between the change in

heading angle on CS+ and CS- trials on trial n) should not depend on the previous trial. Consistent

with this prediction, the trial $n$-$1$ CS x trial $n$ CS interaction was not significant [F(1,15)=2.03,

p=0.175, $BF_{10}$=0.69, $\eta_p^2$=0.12].

The Pavlovian effect did not appear to be driven by explicit awareness of the CS-US

contingency. A post-experiment survey was used to classify participants as either aware or

unaware of the CS-US associations (see *Methods*). Participants who reported being aware of the

contingencies (N=7) did not show a different Pavlovian effect compared to those who reported

being unaware (N=9) in either the acquisition [-0.34°, [-1.90° 1.22°]; t(14)=-0.470, p=0.647,

$BF_{10}$=0.46, *d*=-0.24] or probe [-0.14°, [-1.34° 1.07°]; t(14)=-0.242, p=0.813, $BF_{10}$=0.44, *d*=-0.12]

phases. A similar null effect of awareness on the strength of conditioning has also been reported

in studies of human delay eyeblink conditioning (Clark and Squire, 1998).

In summary, the observed effects of context on implicit sensorimotor adaptation in both

the acquisition and probe phases in Experiment 1 are consistent with differential conditioning

effects. Feedforward implicit sensorimotor adaptation – here operationalized as a type of CR –

was differentially modulated by arbitrary sensory CSs, with a greater response to the CS+, the

cue that was paired with a visuomotor error.

*The Rescorla-Wagner model for context-dependent sensorimotor adaptation*

One influential model that has been used to describe contextual effects in associative learning

tasks is the Rescorla-Wagner model (Rescorla and Wagner, 1972). This model formalizes

changes in conditioned responses via the modulation of learned associations. Here, the

associative strengths, *V*, of the conditioning stimuli are updated according to the learning rule (Eq.

1):

$$(1) \quad V^{[n]} = V^{[n-1]} + \alpha \cdot \beta \cdot SPE^{[n-1]}; \qquad SPE^{[n-1]} = \lambda - V^{[n-1]}$$

where *V* represents the associative strengths between the US and the CS. It is updated based on

the sensory prediction error (*SPE*) presented on trial *n*-1. The *SPE* is defined as the difference

between the maximum conditioning (asymptotic) level for the US (*λ*) and the associative strength

on the given trial. $\beta$ is the learning rate parameter of the US and $\alpha$ represents the salience of the

CS. We note that the Rescorla-Wagner model is one of many computational frameworks for

associative learning (Courville et al., 2006; Gershman, 2015), and does not provide a mechanistic

account for the error-correction process itself (e.g., the fact that the motor system "knows" to

update movements in the direction opposite of the error). For simplicity, we assume that the sign

of the change in movement direction is coded in specialized neural mechanisms for reducing

motor error (Hadjiosif et al., 2021; Herzfeld et al., 2018; Wolpert et al., 1998).

To examine whether the Rescorla-Wagner model can capture differential conditioning

behavior similar to the results observed in Experiment 1, we performed model simulations. The

simulation results demonstrated both the adaptation effect and Pavlovian effect for the acquisition

phase (Fig. 2E), as well as a clear Pavlovian effect for the Probe phase (Fig. 2F). Importantly, the

behavioral signature of differential conditioning – the larger changes in heading angle on CS+

trials compared to CS- trials – holds for essentially all combinations of parameters in the Rescorla-

Wagner model (Fig. 3, left side).

**Figure 3. Simulations of the Pavlovian effect due to differential conditioning in Experiment 1.**

Simulations of the current trial Pavlovian effect, the difference between the change in heading angle on CS+ and CS- trials, for different parameter values as predicted by the Rescorla-Wagner model (right side). The results for the acquisition phase are shown on top and probe phase on bottom. Contours represent the mean ± SEM of the Pavlovian effect from Experiment 1. All parameter values for the state-space model fail to predict a differential response on the current trial for CS+ and CS- trials (right side).

Sensorimotor adaptation is typically described by a "state-space" model in which the motor state (*x*) is updated according to the following learning rule (Eq. 2):

$$(2) \quad x^{[n]} = A \cdot x^{[n-1]} + B \cdot SPE^{[n-1]}$$

where *SPE* is the sensory prediction error – the difference between the predicted and the actual

sensory feedback – experienced on trial *n*-1, *A* is the retention factor, and *B* is the learning rate.

The state-space model is broadly similar to the Rescorla-Wagner (e.g., both models share the

Markov property, produce exponential-family learning curves, etc.). However, the basic state-

space model is unable to account for context effects: Unlike the Rescorla-Wagner model, it does

not include parameters that allow the updating of separate states that are associated with distinct

contexts. It predicts the change in behavior based on the outcome of the previous trial (i.e., the

adaptation effect); it cannot account for variation in heading angle due to the CS on the current

trial (i.e., the Pavlovian effect; Fig. 3, right side).

To formally compare the Rescorla-Wagner and state-space models in terms of the

observed results, we conducted a model comparison by fitting each participant's time course data

with the two models. The Rescorla-Wagner model provided a better fit to the data than a standard

state-space model, underscoring the significant role of the Pavlovian effect (Fig. 4; *t*-test

comparing sum of squared residuals, $-6.61 \times 10^4$, $[-1.05 \times 10^5\ -2.72 \times 10^4]$; t(15)=-3.62, p=0.003, *d*=-

0.90; *t*-test comparing Akaike Information Criterion (AIC) values, -430, [-572 -288]; t(15)=-6.44,

p<0.001, *d*=-1.61).

**Figure 4. Model comparison.**

The sum of squares residuals (SSR) and Akaike Information Criterion (AIC) difference between Rescorla-

Wagner (RW) and the state-space (SS) models for each participant of Experiment 1.

An additional analysis provided further support for an associative learning account of the

results of Experiment 1. The Rescorla-Wagner model not only provides a framework for

understanding how associations can be formed with arbitrary stimuli, but it can also capture how

the strength of these associations is constrained by the relevance of the cues. For instance,

gustatory cues are much more likely to be associated with an internal bodily state (e.g., nausea)

than a visual cue (Garcia and Koelling, 1966). In the current study, the clamped feedback (the

US) is a highly relevant stimulus for reaching; as such, we should expect it to have an immediate

strong influence on motor behavior. In contrast, the imperative cues (the tone and light CSs) have

no natural relevance for reaching; as such, their contribution to the CR should initially be quite

modest, but gradually increase with experience (Fig. 5A). To test this prediction, we examined

the time-course of the adaptation and Pavlovian effects during acquisition using a linear mixed

model analysis. As expected, at early stages of acquisition, the adaptation effect emerged rapidly

whereas the contribution of the Pavlovian effect was small. Over experience, the relative

contribution of the two effects reversed (Type X Time Bin interaction effect: $F_{(1,365)}=16.7$,

$p<0.001$): The Pavlovian effect gradually grew (mean slope: 0.14, 95% CI: [0.04 0.24]) at the

expense of the adaptation effect, which eventually exhibited a reduced contribution (-0.15, [-0.25

-0.05]) (Fig. 5B). We note that both the decrease in the adaptation effect and the increase in the

Pavlovian effect are not captured by a typical state-space model (Fig. 5A).

**Figure 5. Dynamics of adaptation and Pavlovian effects.**

(A) Time course of the mean weights (least squares regression $\beta$) of the adaptation (trial *n-1*, grey) and

Pavlovian (trial *n*, blue) effects in Experiment 1 as predicted by the Rescorla-Wagner model (solid lines)

and the state-space model (dotted lines). (B) Time course of the mean weights of the adaptation and

Pavlovian effects as derived from fits of the experimental results. Shaded regions represent SEM.

*Pavlovian effects in sensorimotor adaptation are sensitive to the CS-US interval*

The Pavlovian effects observed in Experiment 1 provide evidence that sensorimotor adaptation

can be shaped by arbitrary contextual cues. These results stand in contrast to previous work

showing that arbitrary cues are ineffective for separating competing motor memories (e.g.,

differential responses to clockwise and counterclockwise rotations). This null result has been

taken to reflect the irrelevance of such cues to the motor state (Gandolfo et al., 1996; Karniel and

Mussa-Ivaldi, 2002; Howard et al., 2012, 2013).

Why were the arbitrary cues effective in Experiment 1? Inspired by studies of eyeblink

conditioning, we opted to impose a strong temporal constraint on the CS-US interval. The

contextual cues served as the imperative signals and we required that the participants respond

quickly in response to the imperative, resulting in a mean CS-US interval of around 300 ms. The

timing of events was somewhat more relaxed in prior studies. For example, in Howard et al.

(Howard et al., 2013), the color cues (which could be seen as CSs for differential conditioning)

were presented 1,000 ms before the imperative. When adding in the reaction time, the CS-US

interval may lie beyond the optimal value for cerebellar-dependent learning (Brudner et al., 2016;

Kitazawa et al., 1995; Schneiderman and Gormezano, 1964; Smith et al., 1969). Thus, the

reported ineffectiveness of arbitrary cues in separating motor memories may be related to the

failure to meet the temporal requirements for forming associations between these cues and their

associated feedback signals.

In Experiment 2, we tested the hypothesis that extending the temporal interval between

arbitrary cues and sensorimotor feedback would attenuate or abolish the Pavlovian effect in

visuomotor adaptation. To extend the interval, we opted to have the CSs no longer serve as the

movement imperative; rather, a new cue, a color change of the target, served as the imperative

signal (Fig. 6A). To extend the CS-US interval, we imposed a delay between the onsets of the CS

and the imperative. The mean of the delay was set to 1000 ms, with the actual value selected

from a distribution ranging from 800-1,200 ms to maintain unpredictability of the imperative timing

(see *Methods* for details).

**Figure 6. Experiment 2: Pavlovian effect in differential conditioning is abolished when the delay between the CS and US is long.**

(A) Illustration of trial events and their timing. By using a separate stimulus for the CS (tone or light) and the imperative (target changed from gray to blue), a delay of ~1,000 was imposed between the CS and US. The yellow background highlights the design difference from Experiment 1 (and Experiment 3 below). (B) Mean heading angle (N=64) as a function of trial number. Shaded region represents SEM. (C, D) Trial-by-trial change in heading angle (mean ± SEM) during the acquisition (*C*) and probe (*D*) phases. Left panels present analysis results for an adaptation effect (main effect of trial *n-1*, dark vs light colors) and a Pavlovian effect (main effect of the presented CS on the current trial *n*, filled vs empty bars). The black outlined bars and violin plots (right panel) present the Pavlovian effect, i.e., the subtraction of heading angle changes between CS+ and CS- trials (mean and 95% CI). Dots and thin lines represent individual participants.

During the acquisition phase of Experiment 2, participants exhibited a marked change in

movement direction, opposite to the direction of the clamped feedback {adaptation effect:

$F(1,63)=47.3$, $p<0.001$, $BF_{10}=7.58\times10^6$, $\eta_p^2=0.43$, Mean difference, 95% CI; 1.80°, [1.29° 2.31°]},

reaching a mean asymptote of ~25° (Fig. 6B). The asymptotic values in this experiment were

higher than that seen in Experiment 1, possibly due to differences in the experimental setup (see

*Discussion*). Despite this increase in overall adaptation, a Pavlovian differential conditioning effect

was not detected {$F(1,63)=0.54$, $p=0.465$, $BF_{10}=0.16$, $\eta_p^2=0.01$, Mean difference: 0.17°, 95% CI:

[-0.29° 0.63°]} and there was no trial *n-1* x *n* interaction [$F(1,63)=0.60$, $p=0.442$, $BF_{10}=0.17$,

$\eta_p^2=0.01$] (Fig. 6C). During the probe phase, the trial *n*-1 effect disappeared with the removal of

the visual feedback {$F(1,63)=1.01$, $p=0.319$, $BF_{10}=0.21$, $\eta_p^2=0.02$, Mean difference: -0.22°, 95%

CI: [-0.65° 0.21°]}. Importantly, and consistent with the results from the acquisition phase, we

again found no reliable Pavlovian effect {$F(1,63)=1.36$, $p=0.248$, $BF_{10}=0.25$, $\eta_p^2=0.02$, Mean

difference: 0.28°, 95% CI: [-0.19° 0.75°]} and no trial *n-1* x *n* interaction [$F(1,63)=0.42$, $p=0.517$,

$BF_{10}=0.16$, $\eta_p^2=0.01$] (Fig. 6D). These results suggest that extending the CS-US interval by

approximately 1,000 ms rendered arbitrary sensory contextual cues ineffective in separating

distinct motor memories.

One possible concern with our method of increasing the CS-US interval is that, by using

a different stimulus as the imperative (the target color change), the salience of the tone and light

CSs may have been reduced since the task no longer required the participants to attend to these

cues. To address this, in Experiment 3 we used the same CSs (tone and light) and imperative

signal (color change of the target) as in Experiment 2, but now made the onsets of the CS and

imperative simultaneous on each trial (Fig. 7A). In this way, the CS-US interval is much shorter

([Mean RT ± STD]: 306±57.9 ms), similar to Experiment 1. We also increased the inter-trial

interval in Experiment 3 to roughly match that of Experiment 2 given evidence showing that the

rate of associative events impacts classical conditioning (Gallistel and Gibbon, 2000).

**Figure 7. Experiments 3: Pavlovian effect in differential conditioning is observed when the CS and imperative cue occur simultaneously.**

(A) Illustration of trial events and their timing. The green background highlights the design difference from Experiment 2, namely the simultaneous presentation of the CS and the movement imperative. (B) Mean heading angle (N=64) as a function of trial number. Shaded region represents SEM. (C, D) Trial-by-trial change in heading angle (mean ± SEM) during the acquisition (*C*) and probe (*D*) phases. Left panels present analysis results for an adaptation effect (main effect of trial *n-1*, dark vs light colors) and a Pavlovian effect (main effect of the presented CS on the current trial *n*, filled vs empty bars). The black outlined bars and violin plots (right panel) present the Pavlovian effect, i.e., the subtraction of heading angle changes

between CS+ and CS- trials (mean and 95% CI). Shaded regions and error bars represent SEM. Dots and

thin lines represent individual participants.

The time course of adaptation in Experiment 3 was similar to that observed in Experiment

2, with a mean asymptote around ~25° (Fig. 7B). During the acquisition phase, trial-by-trial

changes in heading angle showed the standard adaptation (trial $n$-1) effect {$F(1,63)=101.6$,

$p<0.001$, $BF_{10}=2.77\times10^{12}$, $\eta_p^2=0.62$, Mean difference, 95% CI; 1.46°, [1.17° 1.74°]}. However,

unlike Experiment 2, we now observed a significant Pavlovian effect {$F(1,63)=6.05$, $p=0.017$,

$BF_{10}=2.35$, $\eta_p^2=0.09$, Mean difference, 95% CI; 0.34°, [0.07° 0.61°]} (Fig. 7C). Consistent with the

Rescorla-Wanger predictions (see Fig. 2E), there was no significant interaction between trial $n$-1

and trial $n$ effects [$F(1,63)=1.30$, $p=0.258$, $BF_{10}=0.24$, $\eta_p^2=0.26$]. The Pavlovian effect was also

significant in the probe phase {$F(1,63)=25.6$, $p<0.001$, $BF_{10}=6.88\times10^3$, $\eta_p^2=0.29$, Mean difference,

95% CI; 0.77°, [0.47° 1.07°]} with neither trial $n$-1 {$F(1,63)=2.37$, $p=0.129$, $BF_{10}=0.41$, $\eta_p^2=0.04$,

Mean difference, 95% CI; -0.28°, [-0.63° 0.08°]} nor interaction effects [$F(1,63)=0.86$, $p=0.358$,

$BF_{10}=0.19$, $\eta_p^2=0.01$] (Fig. 7D).

Taken together, the results of Experiments 2 and 3 support our conjecture that the efficacy

of arbitrary stimuli in serving as contextual cues for sensorimotor conditioning is subject to strong

temporal constraints, a key feature of cerebellar-dependent learning. Even when the tone and

light no longer required attention, they proved effective for differential conditioning when the

interval between these CSs and the US was short (around 300 ms), but not when the interval was

extended (around 1,000 ms).

*Additivity principle in response to compound stimuli is observed in sensorimotor adaptation*

The results of Experiments 1-3 demonstrate that implicit sensorimotor adaptation displays

two prominent features of associative learning – the associability of sensorimotor feedback with

arbitrary sensory cues, and the key role of CS-US timing in the formation of those associations.

In Experiment 4, we tested another core feature of associative learning, the principle of additivity

(Mackintosh, 1976; Pavlov, 1927; Rescorla and Wagner, 1972). This principle is based on the

idea that there is an associative capacity for a given US- the *V* term in the Rescorla-Wagner

equation. That is, multiple CSs can become associated with a given US, but the combined

associative strength is bounded by *V*. As a result of this capacity constraint, CSs effectively

compete with one another, with the associative strength split among multiple cues.

The classic method to examine additivity is compound conditioning, where two or more

stimuli are presented simultaneously to form a "compound" CS (Eq. 3). When paired with a US,

this compound CS will come to elicit CRs. Importantly, the associative strength of the Compound

CS ($V_{comp}$) is the sum of the associative strengths of the elemental CSs ($V_i$), where *nS* in Eq. 3

represents the number of elements forming the compound. Consequently, each element of the

compound, when presented alone, elicits a proportionally weaker CR, with the degree of

attenuation being a function of the associative strength of that CS.

438
$$(3)\ V_i^{[n]} = V_i^{[n-1]} + \alpha_i \cdot \beta \cdot (\lambda - V_{comp}^{[n-1]}); \qquad V_{comp}^{[n-1]} = \sum_{i=1}^{nS} V_i^{[n-1]}$$

The additivity principle has received ample support in behavioral and neural studies of

associative learning (Giurfa, 2007; Kehoe et al., 1994; Kehoe and Schreurs, 1986; Rescorla and

Wagner, 1972; Weiss, 1972), but has not, to our knowledge, been tested in sensorimotor

adaptation. In Experiment 4, we used a compound conditioning design, pairing a 15° error clamp

stimulus with a compound CS (simultaneous presentation of the tone and light; Fig. 8A) on all

trials of the acquisition phase. As in Experiments 1-3, we again observed robust adaptation in the

acquisition phase, manifesting as a change in heading angle in the direction opposite the clamp

(Fig. 8B).

**Figure 8. Experiment 4: Compound Conditioning.**

(A) During acquisition, a tone and light were presented simultaneously (Compound CS), serving as the

imperative signal for the reaching movement. They were paired with a 15° clamp on all acquisition trials.

During the probe phase, no feedback was provided, and the CSs were presented either together

(Compound CS) or alone (Single CS; tone or light). (B) Mean heading angle (N=22) as a function of trial

number. Clamped feedback was presented on all trials during the acquisition phase (white background)

and absent on all trials in the probe phase (gray background) phases. Shaded region represents SEM. (C)

Experimental results for the Δ heading angle (mean ± SEM) during the probe phase, showing a positive

relative change for the compound CS on the current trial *n* (filled bar) and a negative relative change for

each of its elements (empty bars). (D) Predictions from the Rescorla-Wagner model for trial-by-trial Δ

heading angle during the probe phase. The two elements were assumed to have equal weight in the

simulation. (E) Scatter plot showing the between-participant trade-off in terms of the associative strength

of the two CSs (the dotted black line represents the unity line). (F) The Δ heading angle (mean ± SEM) after

pooling the two single-CS conditions to measure the effects of the previous CS type (compound versus

singleton, dark and light pink, respectively) and current CS (filled and empty bars). Thin lines represent

individual participants.

The critical test in this experiment comes from the probe phase. Here, the visual feedback

was eliminated, and the presented imperative was either the original compound CS or just the

tone or light alone. We observed a significant Pavlovian effect of CS type on these no-feedback

trials [Friedman test: $\chi^2(2)=9.82$, p=0.007, W=0.223], with larger changes in heading angles on

compound CS trials (Median: 0.39°, Interquartile range: [-0.05° 0.49°]) relative to the tone-alone

(-0.15°, [-0.47° 0.18°], p=0.020) or light-alone (-0.22°, [-0.52° 0.34°], p=0.020) trials, conforming

to the first prediction of the additivity principle (Fig. 8C, 8D).

We note that the median trial-by-trial change in heading for the tone-alone and light-alone

trials is negative. This is because this measure reflects modulations in behavior around the mean

heading angle, and this mean decreases over time due to extinction given the absence of the

error feedback. This general trend should also vary for the different CSs depending on their

relative salience (Eq.3) (Kamin, 1967). Crucially, the additivity principle posits that there should

be a negative correlation between the associative strengths of competing CSs (Rescorla and

Wagner, 1972). That is, if a strong associative bond is formed between one CS and the US, this

will come at the expense of the associative strength accrued by competing CSs (Eq. 3) given the

capacity limit on associability ($V$). This prediction was strikingly confirmed in an analysis of the

heading angle changes on tone-alone and light-alone trials: Participants who were more sensitive

to the tone stimulus were less sensitive to the light stimulus, and *vice versa* (Fig. 8E, Pearson

correlation: r=-0.72, p<0.001, $BF_{10}$=163.2).

As in Experiments 1-3, since the probe phase of Experiment 4 consisted of different types

of CSs presented randomly across trials, the behavior for a given trial should reflect not only the

CS on trial $n$ but also the motor state on trial $n$-1 (which is itself also influenced by the CS on that

trial). As a further test of compound conditioning, we pooled the two single-CS conditions to

measure the effects of the previous and current CS type (singleton versus compound) on the

observed changes in heading angle during the probe phase. We again observed the Pavlovian

effect [Mean: 0.42°, 95% CI: [0.24° 0.59°]; F(1,21)=21.2, p<0.001, $BF_{10}$=459.9, $\eta_p^2$=0.50]: There

was a relative increase in heading angle for the compound CS (0.24°, [0.13° 0.35°]) and a relative

decrease for the singleton CSs (-0.18°, [-0.28° -0.07°]) (Fig. 8F). The main effect of the CS on

trial $n$-1 was not significant [-0.01°, [-0.27° 0.24°]; F(1,21)=0.012, p=0.915, $BF_{10}$=0.21,

$\eta_p^2$=5.58×$10^{-4}$] nor was the trial $n$-1 × trial $n$ interaction [F(1,21)=0.276, p=0.605, $BF_{10}$=0.25,

$\eta_p^2$=0.01], presumably due to the elimination of adaptation effects given the absence of visual

feedback in the probe phase.

## DISCUSSION

Sensorimotor adaptation and delay eyeblink conditioning are foundational paradigms for studying error-based sensorimotor learning. They have yielded a rich empirical foundation in the development of theoretical and neural models of learning and memory, particularly with respect to cerebellar function (Albus, 1971; Ito, 1984; Marr, 1969; Wolpert et al., 1998). Adaptation has typically been modeled from a control engineering perspective, centered on the idea that changes in behavior are driven by sensory prediction errors that arise when experienced feedback deviates from that predicted by a forward model operating on an efference copy of the motor command (Krakauer et al., 2019; Shadmehr and Krakauer, 2008; Wolpert and Flanagan, 2001). In contrast, eyeblink conditioning is treated as an associative learning process that can be shaped by arbitrary sensory context cues.

The present study takes a first step toward a more parsimonious framework that might account for these two canonical forms of cerebellar-dependent motor learning. We were motivated by the apparent paradox concerning whether arbitrary stimuli can act as sufficient contextual cues for the establishment of distinct motor memories during sensorimotor adaptation. In attempting to resolve this paradox, we conceptualized adaptation as analogous to associative learning. Seen from this perspective, we recognized that prior studies failing to observe adaptation in response to arbitrary context cues had not considered the short, precise temporal relationship between these cues and sensorimotor feedback. By modifying a visuomotor adaptation task to conform to timing constraints similar to those required for delay eyeblink conditioning, we showed that adaptation exhibited the hallmarks of both differential conditioning (Experiments 1 and 3) and compound conditioning (Experiment 4). These results provide the first evidence, to our knowledge, that pairing neutral stimuli (i.e., tones and lights) with distinct visuomotor outcomes can differentially influence implicit feedforward sensorimotor adaptation in a manner consistent with core principles of associative learning.

The role of context in sensorimotor adaptation has been the subject of considerable

debate, with a large body of empirical studies asking whether different sensory cues are sufficient

to "tag" different motor memories. Postural and movement-related variables, such as different

lead-in and follow-through movements, are very effective contextual cues and allow participants

to respond to interleaved, opposing perturbations with minimal interference (Howard et al., 2012,

2013, 2015; Sheahan et al., 2016; Schween et al., 2019). In contrast, arbitrary visual features,

such as color cues, have been generally ineffective (Gandolfo et al., 1996; Karniel and Mussa-

Ivaldi, 2002; Howard et al., 2012, 2013; Krouchev and Kalaska, 2003; Osu et al., 2004; Addou et

al., 2011; Forano et al., 2021). To account for this discrepancy, it has been proposed that

movement-related cues are dynamically incorporated into the motor state, allowing for the

separation of distinct internal models, whereas arbitrary cues are not sufficiently relevant to the

motor state (Howard et al., 2012, 2013).

In the current study, we found compelling contextual effects using arbitrary sensory cues.

Importantly, like delay eyeblink conditioning, these effects were subject to strong temporal

constraint on the interval between the CS and US, similar to what is observed in other forms of

cerebellar-dependent sensorimotor learning. Robust differential conditioning behavior was

observed when the CS-US interval was short (Experiments 1 and 3); increasing the interval by

just ~1,000 ms abolished the Pavlovian effect (Experiment 2). These results are also consistent

with prior evidence showing that the efficacy of movement-related cues is highly sensitive to

timing (Howard et al., 2012).

These temporal constraints are likely built into standard adaptation tasks: The target is a

salient stimulus that defines the task goal and movement plan, and its onset usually serves as

the imperative for movement initiation. RTs in these tasks are typically below 500 ms (Kim et al.,

2018, 2019; Avraham et al., 2021). As such, a tight temporal link is established between target

appearance and movement, echoing (perhaps inadvertently) the CS-US temporal constraints

essential for cerebellar-dependent conditioning. Under these conditions, the target can be viewed

as a highly effective contextual cue.

More precisely, we propose that it is the movement plan itself that constitutes the primary,

strongest CS. While the movement plan and target appearance usually coincide spatially (i.e.,

people aim to the target), this is not always the case. For example, with a standard visuomotor

rotation, participants often deliberately aim away from the target, especially when the perturbation

is large (Hegele and Heuer, 2010; Taylor et al., 2014). Recognizing this "re-aimed" plan as

behaving like a CS may provide a parsimonious account of several phenomena. First,

generalization of adaptation appears to be centered around the direction of the movement plan,

not the target (Day et al., 2016; McDougle et al., 2017). Speculatively, this mirrors generalization

effects seen in eyeblink conditioning, where variations on the learned CS (e.g., altering a tone's

frequency) lead to parametric changes in CR probability (Siegel et al., 1968). These changes

produce a similar Gaussian generalization function as that seen in adaptation generalization.

Second, even in the absence of distinct contextual movements (e.g. follow-throughs), the mere

activation of different motor plans can serve as efficient contextual cues, negating interference

effects from opposing perturbations (Sheahan et al., 2016, 2018). Third, an emphasis on the

movement plan is in accord with theoretical models of cerebellar-dependent adaptation, where

the prediction that constitutes the basis for the sensory prediction error is computed using an

efference copy of the movement intention (Blakemore et al., 2001; Gao et al., 2016; Kawato and

Gomi, 1992; Kim et al., 2022; Wolpert et al., 1998).

Our focus in this study was on features of sensorimotor adaptation that map onto a CS,

US, and CR. The UR in this context could be defined as an automatic action that is immediately

triggered by the US, such as an online correction to the perceived error. While there is a rich

literature showing the automatic and relatively fast latency of this corrective response (e.g.,

around 150 ms, Carroll et al., 2019; Franklin and Wolpert, 2008), to isolate feedforward

adaptation, we opted to use rapid shooting movements that precluded online corrections (see

*Results*). We recognize that a feedback response potentiated by the perturbation might be utilized

for feedforward learning (Albert and Shadmehr, 2016). However, a UR is not always required for

conditioning processes arising from volitional behaviors (e.g., Cartoni et al., 2016). Future work

is needed to better understand the role of URs within our framework.

In terms of the feedforward response, there were notable differences in the extent of

adaptation across the four experiments. Overall heading angle reached asymptotic values of ~15°

in Experiments 1 and 4 and ~25° in Experiments 2 and 3. These differences are likely due to

changes in the experimental platform. Experiments 1 and 4 were conducted in our laboratory,

using a tablet and digitizing pen set-up in which the movements involve arm movements of at

least 8 cm. In contrast, Experiments 2 and 3 were conducted via a web-based platform, with each

participant using their personal computer with a mouse or a trackpad to produce wrist or finger

movements of ~2 cm (depending on the sensitivity of the device). Although basic phenomena of

visuomotor adaptation are similar between lab-based and web-based platforms (Tsay et al.,

2021), there may be mechanistic differences. For example, with our laboratory setup the monitor

is positioned just above the hand. As such, the stimuli and movement are in the same plane and

there is a 1:1 correspondence between the distance moved and the extent of the cursor

movement. For the web-based platform, the stimuli and movement occur in roughly orthogonal

planes and there is a gain factor such that the cursor movement is larger than the hand movement.

We expect that the laboratory setup may be much better suited for providing reliable information

concerning proprioception, information that may limit the extent of adaptation (Salomonczyk et

al., 2011; Tsay et al., 2022). Given that proprioceptive and visual signals are not aligned in the

web-based studies, proprioception may have a reduced effect in limiting adaptation, leading to a

higher asymptote.

To formally relate eyeblink conditioning and adaptation, we implemented the Rescorla-

Wagner model, a classic associative learning model that has been widely employed in the

classical and operant conditioning literature (Rescorla and Wagner, 1972). The success of this

model to capture the general features of the current data sets should not be surprising since the

model was developed to account for phenomena such as differential conditioning and compound

conditioning. Thus, it was somewhat overdetermined that the Rescorla-Wagner model would

provide better fits than the standard state-space model in our study, given that the latter cannot

capture contextual effects. In theory, a standard state-space model could be modified such that

different sensory cues become associated with different independent states to allow for context-

dependent learning (Heald et al., 2018). However, we emphasize that to account for compound

conditioning, and in particular the additivity principle (i.e., overshadowing), a state-space model

would essentially have to be transformed into a facsimile of a Rescorla-Wagner model, supporting

our key conclusions.

The Rescorla-Wagner model offers an account of how arbitrary sensory cues, visual

targets, and movement plans can serve as CSs for adaptation. There are, however, some notable

phenomena in the sensorimotor adaptation literature that are not accounted for by our application

of the Rescorla-Wagner model, nor by the basic state-space model. Given the prominent role of

the state-space model in the sensorimotor adaptation literature, variants of this model were

developed to capture these observations. One such effect is spontaneous recovery, the re-

manifestation of a previously adapted state in the absence of error feedback. A variant of the

state-space model, one that allows for multiple states with different learning and forgetting rates,

can capture spontaneous recovery (Smith et al., 2006). A second example is the relationship

between learning rate and environmental consistency; learning is faster in response to a

consistent versus inconsistent perturbation (Albert et al., 2021; Avraham et al., 2020;

Gonzalez Castro et al., 2014; Herzfeld et al., 2014; Hutter and Taylor, 2018). To account for this

effect, the state-space model can be modified to allow the learning rate to vary with experience

(Herzfeld et al., 2014). A third example is savings upon relearning, where faster adaptation is

observed upon a second exposure to the same perturbation (Huang et al., 2011; Krakauer et al.,

2005). Studies have modeled savings using multi-rate state-space models (Smith et al., 2006),

learning rate modulation (Herzfeld et al., 2014), or a combination of these factors (Zarahn et al.,

2008; Mawase et al., 2014).

These higher-order phenomena have also been addressed in a new model of

sensorimotor learning, one that takes a Bayesian inference approach (Heald et al., 2021). The

contextual inference (COIN) model posits that motor adaptation arises from two interacting

mechanisms: Proper learning, which involves the creation and updating of context-specific

memories, and apparent learning, a process of inferring the current context to determine which

memory to express (recall). Unlike conventional context-independent models of motor adaptation,

the COIN model stresses the importance of the environment in guiding memory retrieval. In one

sense, our implementation of the Rescorla-Wagner model follows a similar philosophy. However,

it maps to the proper learning component of the COIN model since we assume that the current

context is determined by the actual properties of the environment in an all-or-none manner (e.g.,

the CS presented by the experimenter). The COIN model suggests that spontaneous recovery,

consistency effects and savings may emerge due to contextual inferences; these inferences likely

involve cognitive systems related to executive function and nonmotor memory systems (Collins

and McDougle, 2021). Consistent with this view, growing evidence show that these phenomena

are, in large degree, the result of strategic changes (Haith et al., 2015; Morehead et al., 2015;

Leow et al., 2020; Avraham et al., 2020, 2021; Wang et al., 2022).

While the basic Rescorla-Wagner model is unable to account for higher order effects such

as spontaneous recovery or savings, we envision that complimenting it with more sophisticated,

Bayesian inference models could readily accommodate these phenomena. In fact, models that

combine associative mechanisms with some form of a Bayesian inference process that carves

the world into distinct contexts have successfully captured a range of complex phenomena in

classical conditioning and reinforcement learning (Collins and Frank, 2013; Courville et al., 2006;

Gershman, 2015; Kruschke, 2008). We believe that our results lay the foundation for adopting a

similar approach to study implicit sensorimotor adaptation and perhaps a more general account

that captures the operation and interaction of multiple learning processes.

Motivated by the cerebellar literature, we limited our focus to delay eyeblink conditioning

in considering how associative learning mechanisms can account for contextual effects in

sensorimotor adaptation. We recognize that conditioning effects can be much more complex than

the behavioral changes that arise from the pairing of two stimuli (Rescorla, 1988). These include

associations between a CS and an action (Skinner, 1938), learning that does not abide by strict

timing requirements (Gallistel and Gibbon, 2000), and other conditioning processes (e.g., trace

conditioning) that rely on brain structures outside the cerebellum (Clark et al., 2002; Clark and

Squire, 1998). Multiple learning processes allow the organism to build a comprehensive

representation of the world, with the ability to flexibly modify behavior at many levels of control in

response to changes in the contingencies between the body and environment (McDougle et al.,

2016; Kim et al., 2020). Nonetheless, by showing that core constraints of delay eyeblink

conditioning are relevant for sensorimotor adaptation, our study may inform a formal framework

that looks at these two domains in a more unified manner.

Although our results highlight principles that address context-dependent sensorimotor

adaptation, they do not speak to the error-correcting adaptation algorithm itself. Conventional

models of this algorithm focus on the role of forward models in predicting future sensory states

and updating motor commands to reduce sensory prediction errors (Wolpert and Ghahramani,

2000; but see Hadjiosif et al., 2021). Associative learning alone does not provide a mechanism

for the error correction process (e.g., the directional change of reaching movements given rotated

feedback, or the precise timing of an eyeblink response to a predicted air puff). Rather, it describes

the association between a contextual cue and a salient event (e.g., shock, reward, etc.).

Speculatively, an alternative idea is that sensorimotor adaptation may operate as a lookup table

of context-outcome associations learned by a system designed to keep the sensorimotor system

calibrated. A model-free conception like this could bring sensorimotor adaptation closer to

classical associative models of cerebellar learning and plasticity (Albus, 1971; Ito, 1984; Marr,

1969).

**METHODS**

*Participants*

A hundred and sixty-six healthy volunteers (aged 18-35 years; 98 females, 6 identified as 'other')

participated in either Experiment 1 (N=16), Experiment 2 (N=64), Experiment 3 (N=64) or

Experiment 4 (N=22). In Experiments 1 and 4, all participants were right-handed, as self-reported

and verified with the Edinburgh Handedness Inventory. The participants in Experiments 2 and 3

did not complete a handedness inventory, but 108 of them self-reported being right-handed, 17

left-handed, and 3 ambidextrous. The sample size for each experiment was set to ensure good

statistical power (details provided in the protocol for each experiment). The protocol was approved

by the Institutional Review Board at the University of California, Berkeley.

*Experimental setup and task*

In Experiments 1 and 4, the participant sat at a custom-made table that housed a horizontally

mounted LCD screen (53.2 cm by 30 cm, ASUS), positioned 27 cm above a digitizing tablet (49.3

cm by 32.7 cm, Intuos 4XL; Wacom, Vancouver, WA). The participant held in their right hand a

hockey paddle that contained an embedded digitizing stylus. The monitor occluded direct vision

of the hand, and the room lights were extinguished to minimize peripheral vision of the arm.

Reaching movements were performed by sliding the paddle across the tablet. The sampling rate

of the tablet was 200 Hz.

For Experiments 2 and 3, we used a web-based platform that was created for conducting

web-based studies in sensorimotor learning (Tsay et al., 2020b). Since participants performed the

experiment remotely with their personal computer, the test apparatus varied across participants.

Based on self-report data, 71 participants used an optical mouse, 55 used a trackpad, and 2 used

trackball. Monitor sizes varied between 11 and 30 inches. The sizes of the stimuli were adjusted

to the monitor size. Below we refer to the stimuli magnitudes in the lab-based experimental setup

(Experiments 1 and 4).

At the beginning of each trial, a white circle (0.5 cm diameter) appeared at the center of

the black screen, indicating the start location (Fig. 1A). The participant moved the stylus to the

start location. Feedback of hand position (i.e., the stylus position) was indicated by a white cursor

(0.3 cm diameter), provided only when the hand was within 1 cm of the start location. A single

blue target (0.6 cm diameter) was positioned 8 cm from the start location. In most studies of

adaptation, the appearance of the target specifies both the movement goal (where to reach) and

serves as the imperative (when to reach). From a classical conditioning perspective, the target

should constitute a very salient CS given that its onset is temporally contingent with the US, the

visual feedback associated with the movement (see below). To eliminate this temporal

contingency, the target remained visible at the same location during the entire experiment. For

each participant, the target was placed at one of four locations, either 45°, 135°, 225°, and 315°,

and this location was counterbalanced across participants.

The task included the presentation of neutral (non-spatial) CS(s). We used two different

CSs, a tone and a light, both of which have no inherent association with the US. The tone CS was

a pure sine wave tone with a frequency of 440 Hz. The light CS was a white rectangular frame

[39.4 cm X 26.2 cm] that spanned the perimeter of the visual workspace. The large frame was

selected to provide a salient visual stimulus, but one that would not be confused with the target.

Depending on the specific experimental protocol and task phase, the CSs could appear

alone or together on a trial. In Experiments 1 and 4, the onset of the CS served as the imperative

signal, with the participant instructed to rapidly reach directly towards the target, slicing through

the target. The onset of the CS occurred following a pseudo-random and predetermined delay

after the hand was positioned at the start location. This was done to mitigate predictions regarding

the timing of the CS onset. The delay ranged between 800-1,200 ms (in steps of 100 ms), drawn

from a uniform distribution. While the CSs were also presented in Experiments 2 and 3, a different

signal was used for the movement imperative (see below). In all experiments, the CS was

terminated when the hand reached the radial distance to the target (Fig. 1A). In delay eyeblink

conditioning, learning is optimal when the US is presented 100 to 500 ms after the CS

(Schneiderman and Gormezano, 1964; Smith et al., 1969). To minimize the delay between the

onset of the CS (or the other imperative cue) and the US, the auditory message "start faster" was

played whenever a reaction time (RT) exceeded 400 ms. RT was operationalized as the interval

between imperative onset and the time required for the radial distance of the hand to exceed 1

cm. Pilot work showed that RTs with the variable imperative onset time were always greater than

100 ms; as such, we did not set a minimum RT bound. Given our objective to test the link between

feedforward adaptation and classical conditioning, we sought to eliminate online feedback

corrections. Participants were instructed to make rapid movements and the auditory message

"move faster" was played whenever movement time (MT) exceeded 300 ms. The end of the

movement was operationalized as the point where the radial distance of the hand reached 8 cm.

Experiments 2 and 3 were designed to test the hypothesis that the time interval between

the arbitrary cues and the sensorimotor feedback is critical for driving Pavlovian effects. To this

end, we did not use the CS onset as the imperative signal; rather, we introduced a new cue, a

color change of the target from gray to blue, to serve as the imperative signal (Figs. 6A, 7A). The

difference between the two experiments was the interval between the onsets of the CS (tone or

light) and imperative. In Experiment 2, the imperative was delayed with respect to CS onset, with

the delay drawn from a uniform distribution ranging from 800-1,200 ms (in steps of 100 ms). As

such, we added a mean of 1,000 ms to the CS-US interval. The delay distribution that we used

added variance to the CS-US interval, allowing us to make sure that, like Experiment 1,

participants could not anticipate the onset of the imperative. This was important since

predictability of the imperative timing could have decreased reaction times, and thus effectively

decrease the delay between the CS and the feedback. We also note that previous evidence from

eyeblink conditioning shows that the conditioned response is minimally affected by the variability

in the CS-US interval (Patterson, 1970). In Experiment 3, the CS and imperative were

simultaneous (similar to the timing in Experiment 1). Trial durations were made comparable

between Experiments 2 and 3 by setting the mean interval from the trial onset to the imperative

in Experiment 3 to match that of Experiment 2.

For the unconditioned stimulus (US), we used task-irrelevant clamped feedback

(Morehead et al., 2017). With clamped feedback, the radial position of the visual cursor is matched

to the radial position of the hand. However, the angular position of the cursor is fixed. The

participant thus controlled the speed and radial distance of the cursor, but not its direction. When

designed to produce a prediction error and elicit implicit sensorimotor adaptation, the clamp

followed a path that deviated from the target by 15°, with the direction, i.e., clockwise (CW) or

counterclockwise (CCW), counterbalanced across participants. We also included no-error trials

(Experiments 1-3) by presenting clamped feedback that followed a path directly to the target (0°

clamp; Fig. 2A). The nature of the clamp manipulation was described in detail to the participant,

and they were explicitly instructed strictly to ignore the feedback, aiming their reach directly toward

the target on every trial. These instructions were designed to emphasize that the participant did

not control the cursor's angular position, and that they should always attempt to reach directly to

the target. The instructions of the task were reinforced by the presentation of short video

animations to demonstrate how the CSs would serve as imperative signals, as well as to show

the invariant direction of the clamped feedback.

For the lab-based experiments (Experiments 1 and 4), the experimental software was

custom written in Matlab (The MathWorks, Natick, MA), using the Psychtoolbox package

(Brainard, 1997). The web-based experiments (Experiments 2 and 3) were developed for a

webpage using JavaScript, HTML, and CSS. Data from the web-based experiments are

temporarily stored via the Google Firebase database and downloaded for off-line analyses.

*Experimental protocol*

All experiments included an acquisition phase and a probe phase. During the acquisition phase,

clamped feedback was presented on each trial, serving as the US. During the probe phase, the

clamped feedback was not presented. In both phases, the participants were instructed to reach

straight to the target as soon as the imperative appeared. Note that we opted to not include

baseline reaching blocks prior to the start of the acquisition phases to avoid introducing any

incidental associations between the baseline feedback and the target, movement plan, and any

other contextual variables. For lab-based Experiments 1 and 4, a break of approximately 1-minute

was provided in the middle of the experiment.

Experiments 1-3: Differential conditioning

Experiment 1 (N=16) was designed to test differential conditioning in the context of a sensorimotor

adaptation task. The target number of participants was determined based on the effect size (d =

0.67) observed for the Pavlovian effect in a pilot study, with a significance level of α = 0.05 and

power of 0.8. Experiments 2 and 3 were designed to test temporal constraints on differential

conditioning. Participants (N=64 per experiment, 128 total) were recruited using the

crowdsourcing website Prolific ([www.prolific.co](www.prolific.co)). The minimum sample size for each experiment

was determined based on the sample size of Experiment 1 (N=16) where robust Pavlovian effects

were observed. However, we have found that data collected with our web-based platform are

more variable (Tsay et al., 2021). As such, we opted to use a larger sample size, setting our target

as 64 participants for each experiment.

The session consisted of 800 trials: 600 acquisition trials followed by 200 probe trials (Fig.

2A). One of two CSs (tone or light) was presented on each trial, serving as the imperative for the

reaching response. During the acquisition phase, one CS was paired with 15° clamped error

feedback (CS+ condition) and the other CS was paired with 0° clamped feedback (CS- condition).

Each CS was presented on 50% of the trials, and the assignment of the tone and light to the CS+

and CS- was counterbalanced across participants. During the probe phase, each CS was

presented alone on half of the trials, and there was no visual feedback.

For both the acquisition and probe phases, the CS+ and CS- trials were interleaved in a

pseudo-random order that was unique to each participant. To ensure that the participant would

not be able to predict the CS type, the generated trial sequence for each participant was assessed

to verify that there was no significant lag-1 autocorrelation in the time series. At the end of the

experimental session, we assessed whether participants were aware of the contingency between

each CS and its associated feedback, asking, "Did you identify any pattern in the experiment?" in

a free-response survey.

Experiment 4: Compound conditioning

The adaptation task was modified in Experiment 2 (N=22) to provide a test of compound

conditioning. The sample size was determined based on the effect size (d = 0.56) observed for

the Pavlovian effect in Experiment 1, with a significance level of α = 0.05 and power of 0.8.

The procedure was similar to that used in Experiments 1-3 with the following changes:

First, the session consisted of 600 acquisition trials and 300 probe trials (Fig. 8A). Second, a

compound CS, consisting of both the tone and light, served as the imperative throughout the

acquisition phase, and was always paired with 15° clamped feedback. Third, the probe phase (no

visual feedback) consisted of 100 trials for each of the compound CS, tone alone CS, and light

alone CS.

*Data analysis*

The recorded position of the hand was analyzed using custom-written MATLAB scripts. Our main

analyses focused on the reach direction (heading angle) and the trial-by-trial changes in heading

angle (Δ heading angle). Heading angle was defined by two imaginary lines, one from the start

position to the target and the other from the start position to the hand position at maximum

movement velocity.

For all analyses, and to visualize the results, the sign of the heading angle was flipped for

participants who experienced a CCW clamp, such that a positive heading angle is in the direction

of expected adaptation (i.e., opposite the direction of the perturbed feedback). Moreover, the

heading angle on the first acquisition trial was treated as the baseline reaching angle and

subtracted from the heading angle on all subsequent trials. (We note that the results remain

unchanged in terms of statistical comparisons if this baseline subtraction step is omitted.)

For lab-based experiments (1 and 4), trials in which the heading angle was larger than

100° off from the target, or in which the trial-to-trial change in heading angle was larger than 25°,

were considered outliers and not included in the analyses. For web-based experiment (2 and 3),

outlier trials were considered trials in which the heading angle was beyond the range of -50°-70°

off the target, or in which the trial-to-trial change in heading angle was larger than 20°. The criteria

for considering outlier trials were determined after visual examination of the time courses of all

individuals in each experimental platform (lab-based and web-based). To eliminate biases, the

CS was not visible in the trial traces. Outliers constituted 0.03% (Experiment 1), 1.26%

(Experiment 2), 1.44% (Experiment 3) and 0.16% (Experiment 4) of the trials in each experiment.

For the change in heading angle analysis, but not for the presentation of heading angle

time courses (Figs. 2B, 6B, 7B, 8B), we excluded trials in which the reaction time exceeded 400

ms and/or movement time exceeded 300 ms (Experiment 1: 11% of acquisition trials, 4.3% of

probe trials; Experiment 2: 19% of acquisition trials, 16% of probe trials; Experiment 3: 19% of

acquisition trials, 15% of probe trials; Experiment 4: 3.4% of acquisition trials, 4.9% of probe trials).

In Experiments 1-3, the primary analyses examined how the Δ heading angle was

influenced by the CS type (CS+ vs CS-), either in terms of the previous trial (*n-1*, adaptation effect)

or current trial (*n*, Pavlovian effect). For each participant and phase, we calculated the average Δ

heading angle for four types of trials: CS+ trials that follow CS+ trials, CS- after CS+, CS+ after

CS-, and CS- after CS-. For each phase, a two-way repeated-measures ANOVA was conducted

with two within-participant independent factors, the CS on trial *n-1* and the CS on trial *n*, each with

two levels, CS+ and CS-. The Δ heading angle was the dependent variable in the two ANOVAs.

To examine the dynamics of the adaptation and Pavlovian effects (Fig. 5B), we binned the

heading angle data into epochs of 50 trials. Within each bin, we performed multiple regression

analysis to test whether trial-to-trial change in heading angle can be predicted from the previous

trial CS, the current trial CS and their interaction. Figure 5B presents the mean ± SEM regression

β weights of all simulated time courses for the previous (adaptation effect) and current (Pavlovian

effect) predictors. To evaluate statistically the changes in β weights for each type of effect across

bins, we used a linear mixed model (R statistical package: lmerTest), with Type (adaptation and

Pavlovian) and Bin as fixed effects and participants as random effects.

To assess whether the Pavlovian effect in Experiment 1 is influenced by the awareness

about the contingency between each CS and its respective feedback, we divided the participants

based on their responses to the post experiment questionnaire (see *Experimental protocol*): 7 out

of the 16 participants stated the correct contingency between the CS and the visual feedback and

were thus considered the "aware sub-group." The rest of the participants (N=9) reported that they

did not identify any pattern related to the CS-US contingency and were considered the "unaware

sub-group." Independent two-sample *t* tests were used to compare the Pavlovian effects between

these groups during the acquisition and probe phases of the experiment.

Feedback correction was operationalized as the difference between the heading angle

measured at the radial distance to the target and at 50 ms after movement initiation. We estimated

the mean and standard deviation of feedback correction across all acquisition trials in all the

participants in Experiment 1. In addition, we calculated, for each participant, the mean feedback

correction for each of the CS+ and CS- trials and used a paired-sample *t* test to examine within-

participant changes in feedback correction between the two trial types.

In Experiment 4, the analysis focused on the probe phase in which there was no visual

feedback. We compared the Δ heading angle in response to the three CSs on trial *n* (compound

CS, light alone, tone alone) regardless of the CS presented on trial *n*-1. As the assumption of

normality was not met for the distribution of one of the conditions, we used the non-parametric

Friedman test, with the Δ heading angle as the dependent variable, and the CS type as the within-

participant independent variable.

The additivity principle of the Rescorla-Wagner model states that the association strengths

of each element of the compound CS will compete for associative strength with respect to the US

(Eq. 3). This principle was tested in our data by examining the correlation between the Δ heading

angle associated with the tone and light CSs in Experiment 4.

We report effect size using Cohen's $d$ for all $t$ tests, partial eta-squared ($\eta_p^2$) for the ANOVA

and the Kendall's W for the Friedman test. We followed the assumption of normality when the

sample size was larger than N=30 based on the central limit theorem or when the Lilliefors test

indicated normality (Lilliefors, 1967). When these conditions were met, we used parametric

statistical tests. In all other cases we used non-parametric statistics (see Experiment 4). Family-

wise errors in pairwise comparisons were corrected using the Bonferroni correction. We also

report Bayes factor $BF_{10}$, the ratio of the likelihood of the alternative hypothesis ($H_1$) over the null

hypothesis ($H_0$) (Kass and Raftery, 1995).

*Model simulations*

Trial-by-trial reach angles were simulated using the Rescorla-Wagner model as a representative

model for associative learning. In this model, the motor state is updated based on the associative

strength ($V$) between the US (error signal) and all the CSs present on a given trial (Eqs. 1 and 3).

The extent of learning is determined by the maximum conditioning level ($\lambda$), and the rate of

behavioral change are determined by the learning rate of the US ($\beta$) and the salience of each

presented CS ($\alpha$). Note that in simulating the Rescorla-Wagner model for visuomotor adaptation,

we included the movement plan as a CS (*CS_plan*) on all trials in addition to the trial-specific CS

(*CS_tone* or *CS_light*). The following pseudocode describes how the heading angle, represented

by the total associative strength *V_total* between all CSs and the US, is updated in a simulation

of differential conditioning. For simplicity, we consider *CS_tone* = *CS*+ and *CS_light* =*CS*-. *US*=1

on *CS*+ trial and *US*=0 on *CS*- trials and on no feedback trials (all the trials in the probe phase).

On trial $n=1$, before experiencing the *US*, all associative strengths are initialized to 0.

**FOR** each other *trial n* in the experiment

$V\_plan_n \leftarrow V\_plan_{n-1} + \alpha\_plan \times \beta \times (\lambda \times US_{n-1} - V\_total_{n-1})$

**IF** trial $n$-1 is *CS_tone*

917            $V\_total_{n-1} \leftarrow V\_plan_{n-1} + V\_tone_{n-1}$

918            $V\_tone_n \leftarrow V\_tone_{n-1} + \alpha\_tone \times \beta \times (\lambda \times US_{n-1} - V\_total_{n-1})$

919            $V\_light_n \leftarrow V\_light_{n-1}$

920        **ELSE IF** trial $n$-1 is *CS_light*

921            $V\_total_{n-1} \leftarrow V\_plan_{n-1} + V\_light_{n-1}$

922            $V\_light_n \leftarrow V\_light_{n-1} + \alpha\_light \times \beta \times (\lambda \times US_{n-1} - V\_total_{n-1})$

923            $V\_tone_n \leftarrow V\_tone_{n-1}$

**END IF**

**END FOR**

In compound conditioning, all the trials in the acquisition phase are presented with an error ($US$=1)

and with both *CS_tone* and *CS_light* (and *CS_plan*). Therefore, both *V_tone* and *V_light* (and

*V_plan*) are updated on all trials:

**FOR** each *trial n*>1 in the acquisition phase

$V\_total_{n-1} \leftarrow V\_plan_{n-1} + V\_tone_{n-1} + V\_light_{n-1}$

$V\_plan_n \leftarrow V\_plan_{n-1} + \alpha\_plan \times \beta \times (\lambda \times US_{n-1} - V\_total_{n-1})$

$V\_tone_n \leftarrow V\_tone_{n-1} + \alpha\_tone \times \beta \times (\lambda \times US_{n-1} - V\_total_{n-1})$

$V\_light_n \leftarrow V\_light_{n-1} + \alpha\_light \times \beta \times (\lambda \times US_{n-1} - V\_total_{n-1})$

**END FOR**

During the probe phase in compound conditioning, the associative strengths are updated based

on the CS presented on each trial, either as a compound, which is simulated the same as the

above pseudocode for compound conditioning, or alone, and then the algorithm uses the same

implementation as in differential conditioning. We set $US=0$ in all the trials of the probe phase.

To illustrate the predictions of the model in Figures 2E, 2F and 8D, we chose parameters

that result in qualitatively similar effects to the experimental results. For both Experiments 1 and

2, we set $\lambda$ to 15, similar to the observed implicit adaptation asymptotes in these experiments, as

well as in other studies (Bond and Taylor, 2015; Morehead et al., 2017). The value of $\beta$ was set

to 0.12 (Exp. 1) or 0.02 (Exp. 4), the salience parameter ($\alpha$) for the movement plan CS to 0.99,

and both the tone CS and light CS to 0.002 (Exp. 1) or 0.1 (Exp. 4). These divergent salience

values are consistent with the assumption that most of the associative strength of the US would

be absorbed by the movement plan CS given its central relevance to the task of reaching to a

target. The remainder of potential associative strength to the US is thus available for the tone and

light CSs.

To demonstrate how the choice of parameters in the Rescorla-Wagner model influences

the predicted Pavlovian effects, we simulated behavior using various combinations of the free

parameters. We chose several combinations of values for $\beta$ and $\lambda$, and, for each combination, a

wide range of values for $\alpha_{goal}$, $\alpha_{tone}$ and $\alpha_{light}$. (For simplicity, we constrained $\alpha_{tone}$ and $\alpha_{light}$ to the

same value.) Figure 3 displays heatmaps resulting from these simulations, with the color of each

cell corresponding to the simulated difference in the trial-by-trial change in heading angle between

CS+ and CS- trials. In addition, we illustrated that a single-process state-space model, a standard

model of sensorimotor adaptation, cannot capture these Pavlovian effects. In this model, the

motor state is updated according to the error observed during the previous movement. The

parameters of this model determine both the trial-by-trial retention of the previous motor state and

the rate of learning from errors (see Eq. 2 in the *Results*). For the simulation presented in Figure

3, the error was fixed to one of two values during the acquisition phase, either -15° on CS+ trials

or 0° on CS- trials, and in the probe phase, it was set to 0° on all trials.

While the two models share similar features and parameters (e.g., learning rate of an error

signal), the additional parameters in the Rescorla-Wagner model allow it to capture the effects of

differential and compound conditioning, should these processes be operative in our experiments.

For example, in the differential conditioning case, modifying the parameters of the Rescorla-

Wagner model could produce changes in the magnitude of both the within-trial "Pavlovian" effects

(i.e., the modulation of behavior in response to the current CS, or "trial *n*" effects) and across-trial

adaptation effects (i.e., the modulation of behavior in response to the current state of learning, or

"trial *n-1*" effects). However, the Rescorla-Wagner model will always yield unique responses to

the CS+ and CS- given non-zero salience parameters. In contrast, no combination of the *A* and

*B* parameters in the state-space model will produce differential responses to the tone and light

CSs.

To simulate the dynamics of the adaptation and Pavlovian effects in differential

conditioning (Fig. 5A), and to compare it to the dynamics observed in the actual data (Fig. 5B),

we simulated the trial-by-trial change in heading angle during acquisition (600 trials) according to

the Rescorla-Wagner model (Eqs. 1 and 3) and the state-space model (Eq. 2), based on the

schedules of CS+ and CS- trials that were presented to the participants (16 simulated time

courses). For the Rescorla-Wagner model, we used the same parameters values that generated

the simulation results presented in Figures 2E and 2F. For the state-space model, we set the *A*

parameter to 0.9 and *B* to 0.12. We then did the same trial-by-trial regression analysis described

above for the actual heading angle data (see *Data analysis*) with the exception that the dependent

variable was now the simulated change in heading angle.

*Model fitting and comparison*

We conducted a *post-hoc* model comparison analysis (Fig. 4). In this analysis, we fit the Rescorla-

Wagner model (Eqs. 1 and 3) and the standard state-space model (Eq. 2) to participants' heading

angle time course data of Experiment 1. The two models were fit by minimizing the sum of squared

residuals between the measured and modeled movement data, using the MATLAB function

*fmincon*. To avoid local minima, 200 randomized sets of initial parameter values were used during

fitting and the best fit of each model was selected for model comparison. Models were compared

using both the sum of squared residuals and the Akaike Information Criterion approximated on

the residuals (AIC; Akaike, 1974). All free parameters were bound at [0, 1], except for $\lambda$, which

was bounded at [-30, 60]. We did not perform model fitting for Experiment 4 since all trials were

identical in the acquisition phase. As such, the salience parameters in the Rescorla-Wagner

model for the goal, tone and light are unidentifiable, and both models make indistinguishable

behavioral predictions during acquisition.

1000

1001   *Data and code availability*

1002   All raw data files and codes for data analysis and simulations are available from the GitHub

1003   repository: https://github.com/guyavr/AssociativeMotorAdaptation.git

1004

**ACKNOWLEDGEMENTS**

We thank members of the BLAM lab (Johns Hopkins) for helpful discussions. We thank Marina

Iranmanesh, Janet Hwang, Sarvenaz Pakzad and Aaron Shalf for their assistance with data

collection. JAT is funded by the NIH (NS084948), the NSF (1838462 and 1827550), ONR

N00014-18-1-2873, and J. Insley Blair Pyne Fund. RBI is funded by the NIH (NS116883 and

DC077091).

1011

**COMPETING INTERESTS**

RBI is a co-founder with equity in Magnetic Tides, Inc.

**REFERENCES**

Addou T, Krouchev N, Kalaska JF. 2011. Colored context cues can facilitate the ability to learn

and to switch between multiple dynamical force fields. *Journal of Neurophysiology*

**106**:163–183. doi:10.1152/jn.00869.2010

Akaike H. 1974. A new look at the statistical model identification. *IEEE Transactions on*

*Automatic Control* **19**:716–723. doi:10.1109/TAC.1974.1100705

Albert ST, Jang J, Sheahan HR, Teunissen L, Vandevoorde K, Herzfeld DJ, Shadmehr R. 2021.

An implicit memory of errors limits human sensorimotor adaptation. *Nature Human*

*Behaviour* 1–15. doi:10.1038/s41562-020-01036-x

Albert ST, Shadmehr R. 2016. The Neural Feedback Response to Error As a Teaching Signal

for the Motor Learning System. *J Neurosci* **36**:4832–4845.

doi:10.1523/JNEUROSCI.0159-16.2016

Albus JS. 1971. A theory of cerebellar function. *Mathematical Biosciences* **10**:25–61.

doi:10.1016/0025-5564(71)90051-4

Avraham G, Keizman M, Shmuelof L. 2020. Environmental consistency modulation of error

sensitivity during motor adaptation is explicitly controlled. *Journal of Neurophysiology*

**123**:57–69. doi:10.1152/jn.00080.2019

Avraham G, Morehead JR, Kim HE, Ivry RB. 2021. Reexposure to a sensorimotor perturbation

produces opposite effects on explicit and implicit learning processes. *PLOS Biology*

**19**:e3001147. doi:10.1371/journal.pbio.3001147

Blakemore S-J, Frith CD, Wolpert DM. 2001. The cerebellum is involved in predicting the

sensory consequences of action. *NeuroReport* **12**:1879–1884.

Bond KM, Taylor JA. 2015. Flexible explicit but rigid implicit learning in a visuomotor adaptation

task. *Journal of Neurophysiology* **113**:3836–3849. doi:10.1152/jn.00009.2015

Brainard DH. 1997. The Psychophysics Toolbox. *Spatial Vision* **10**:433–436.

doi:10.1163/156856897X00357

Brudner SN, Kethidi N, Graeupner D, Ivry RB, Taylor JA. 2016. Delayed feedback during

sensorimotor learning selectively disrupts adaptation but not strategy use. *Journal of*

*Neurophysiology* **115**:1499–1511. doi:10.1152/jn.00066.2015

Carroll TJ, McNamee D, Ingram JN, Wolpert DM. 2019. Rapid Visuomotor Responses Reflect

Value-Based Decisions. *J Neurosci* **39**:3906–3920. doi:10.1523/JNEUROSCI.1934-

18.2019

Cartoni E, Balleine B, Baldassarre G. 2016. Appetitive Pavlovian-instrumental Transfer: A

review. *Neuroscience & Biobehavioral Reviews* **71**:829–848.

doi:10.1016/j.neubiorev.2016.09.020

Clark RE, Manns JR, Squire LR. 2002. Classical conditioning, awareness, and brain systems.

*Trends in Cognitive Sciences* **6**:524–531. doi:10.1016/S1364-6613(02)02041-7

Clark RE, Squire LR. 1998. Classical Conditioning and Brain Systems: The Role of Awareness.

*Science* **280**:77–81. doi:10.1126/science.280.5360.77

Collins AGE, Frank MJ. 2013. Cognitive control over learning: Creating, clustering, and

generalizing task-set structure. *Psychological Review* **120**:190–229.

doi:10.1037/a0030852

Collins AGE, McDougle SD. 2021. Context is key for learning motor skills. *Nature*.

doi:10.1038/d41586-021-03028-x

Courville AC, Daw ND, Touretzky DS. 2006. Bayesian theories of conditioning in a changing

world. *Trends in Cognitive Sciences*, Special issue: Probabilistic models of cognition

**10**:294–300. doi:10.1016/j.tics.2006.05.004

Dawidowicz G, Shaine Y, Mawase F. 2021. Separation of multiple motor memories through

implicit and explicit processes. *Journal of Neurophysiology*. doi:10.1152/jn.00245.2021

Day KA, Roemmich RT, Taylor JA, Bastian AJ. 2016. Visuomotor Learning Generalizes Around

the Intended Movement. *eNeuro* **3**. doi:10.1523/ENEURO.0005-16.2016

Donchin O, Rabe K, Diedrichsen J, Lally N, Schoch B, Gizewski ER, Timmann D. 2011.

Cerebellar regions involved in adaptation to force field and visuomotor perturbation.

*Journal of Neurophysiology* **107**:134–147. doi:10.1152/jn.00007.2011

Forano M, Franklin DW. 2020. Timescales of motor memory formation in dual-adaptation. *PLOS*

*Computational Biology* **16**:e1008373. doi:10.1371/journal.pcbi.1008373

Forano M, Schween R, Taylor JA, Hegele M, Franklin DW. 2021. Direct and indirect cues can

enable dual adaptation, but through different learning processes. *Journal of*

*Neurophysiology* **126**:1490–1506. doi:10.1152/jn.00166.2021

Franklin DW, Wolpert DM. 2008. Specificity of Reflex Adaptation for Task-Relevant Variability. *J*

*Neurosci* **28**:14165–14175. doi:10.1523/JNEUROSCI.4406-08.2008

Gallistel CR, Gibbon J. 2000. Time, rate, and conditioning. *Psychological Review* **107**:289–344.

doi:10.1037/0033-295X.107.2.289

Gandolfo F, Mussa-Ivaldi FA, Bizzi E. 1996. Motor learning by field approximation. *PNAS*

**93**:3843–3846. doi:10.1073/pnas.93.9.3843

Gao Z, Proietti-Onori M, Lin Z, ten Brinke MM, Boele H-J, Potters J-W, Ruigrok TJH, Hoebeek

FE, De Zeeuw CI. 2016. Excitatory Cerebellar Nucleocortical Circuit Provides Internal

Amplification during Associative Conditioning. *Neuron* **89**:645–657.

doi:10.1016/j.neuron.2016.01.008

Garcia J, Koelling RA. 1966. Relation of cue to consequence in avoidance learning. *Psychon*

*Sci* **4**:123–124. doi:10.3758/BF03342209

Gershman SJ. 2015. A Unifying Probabilistic View of Associative Learning. *PLoS Comput Biol*

**11**. doi:10.1371/journal.pcbi.1004567

Giurfa M. 2007. Behavioral and neural analysis of associative learning in the honeybee: a taste

from the magic well. *J Comp Physiol A* **193**:801–824. doi:10.1007/s00359-007-0235-9

Gonzalez Castro LN, Hadjiosif AM, Hemphill MA, Smith MA. 2014. Environmental Consistency

Determines the Rate of Motor Adaptation. *Current Biology* **24**:1050–1061.

doi:10.1016/j.cub.2014.03.049

Hadjiosif AM, Krakauer JW, Haith AM. 2021. Did we get sensorimotor adaptation wrong?

Implicit adaptation as direct policy updating rather than forward-model-based learning. *J*

*Neurosci.* doi:10.1523/JNEUROSCI.2125-20.2021

Haith AM, Huberdeau DM, Krakauer JW. 2015. The Influence of Movement Preparation Time

on the Expression of Visuomotor Learning and Savings. *J Neurosci* **35**:5109–5117.

doi:10.1523/JNEUROSCI.3869-14.2015

Heald JB, Ingram JN, Flanagan JR, Wolpert DM. 2018. Multiple motor memories are learned to

control different points on a tool. *Nature Human Behaviour* **2**:300–311.

doi:10.1038/s41562-018-0324-5

Heald JB, Lengyel M, Wolpert DM. 2021. Contextual inference underlies the learning of

sensorimotor repertoires. *Nature* 1–5. doi:10.1038/s41586-021-04129-3

Hegele M, Heuer H. 2010. Implicit and explicit components of dual adaptation to visuomotor

rotations. *Consciousness and Cognition* **19**:906–917. doi:10.1016/j.concog.2010.05.005

Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R. 2018. Encoding of error and learning to

correct that error by the Purkinje cells of the cerebellum. *Nature Neuroscience* **21**:736–

743. doi:10.1038/s41593-018-0136-y

Herzfeld DJ, Vaswani PA, Marko M, Shadmehr R. 2014. A memory of errors in sensorimotor

learning. *Science* 1253138. doi:10.1126/science.1253138

Howard IS, Ingram JN, Franklin DW, Wolpert DM. 2012. Gone in 0.6 Seconds: The Encoding of

Motor Memories Depends on Recent Sensorimotor States. *J Neurosci* **32**:12756–12768.

doi:10.1523/JNEUROSCI.5909-11.2012

Howard IS, Wolpert DM, Franklin DW. 2015. The Value of the Follow-Through Derives from

Motor Learning Depending on Future Actions. *Current Biology* **25**:397–401.

doi:10.1016/j.cub.2014.12.037

Howard IS, Wolpert DM, Franklin DW. 2013. The effect of contextual cues on the encoding of

motor memories. *Journal of Neurophysiology* **109**:2632–2644.

doi:10.1152/jn.00773.2012

Huang VS, Haith A, Mazzoni P, Krakauer JW. 2011. Rethinking Motor Learning and Savings in

Adaptation Paradigms: Model-Free Memory for Successful Actions Combines with

Internal Models. *Neuron* **70**:787–801. doi:10.1016/j.neuron.2011.04.012

Hutter SA, Taylor JA. 2018. Relative sensitivity of explicit reaiming and implicit motor

adaptation. *Journal of Neurophysiology* **120**:2640–2648. doi:10.1152/jn.00283.2018

Ito M. 1984. The Cerebellum and Neural Control. Raven Press.

Izawa J, Criscimagna-Hemminger SE, Shadmehr R. 2012. Cerebellar Contributions to Reach

Adaptation and Learning Sensory Consequences of Action. *J Neurosci* **32**:4230–4239.

doi:10.1523/JNEUROSCI.6353-11.2012

Kamin LJ. 1967. Attention-like processes in classical conditioning (No. TR-5).

Karniel A, Mussa-Ivaldi FA. 2002. Does the motor control system use multiple models and

context switching to cope with a variable environment? *Exp Brain Res* **143**:520–524.

doi:10.1007/s00221-002-1054-4

Kass RE, Raftery AE. 1995. Bayes Factors. *Journal of the American Statistical Association*

**90**:773–795. doi:10.1080/01621459.1995.10476572

Kawato M, Gomi H. 1992. A computational model of four regions of the cerebellum based on

feedback-error learning. *Biol Cybern* **68**:95–103. doi:10.1007/BF00201431

Kehoe EJ, Horne AJ, Horne PS, Macrae M. 1994. Summation and configuration between and

within sensory modalities in classical conditioning of the rabbit. *Animal Learning &*

*Behavior* **22**:19–26. doi:10.3758/BF03199952

Kehoe EJ, Schreurs BG. 1986. Compound conditioning of the rabbit's nictitating membrane

response: Test trial manipulations. *Bull Psychon Soc* **24**:79–81.

doi:10.3758/BF03330509

Kim HE, Avraham G, Ivry RB. 2020. The Psychology of Reaching: Action Selection, Movement

Implementation, and Sensorimotor Learning. *Annu Rev Psychol.* doi:10.1146/annurev-

psych-010419-051053

Kim HE, Morehead JR, Parvin DE, Moazzezi R, Ivry RB. 2018. Invariant errors reveal limitations

in motor correction rather than constraints on error sensitivity. *Communications Biology*

**1**:19. doi:10.1038/s42003-018-0021-y

Kim HE, Parvin DE, Ivry RB. 2019. The influence of task outcome on implicit motor learning.

*eLife* **8**:e39882. doi:10.7554/eLife.39882

Kim OA, Forrence AD, McDougle SD. 2022. Motor learning without movement. *Proceedings of*

*the National Academy of Sciences* **119**:e2204379119. doi:10.1073/pnas.2204379119

Kitazawa S, Kohno T, Uka T. 1995. Effects of delayed visual information on the rate and amount

of prism adaptation in the human. *J Neurosci* **15**:7644–7652.

doi:10.1523/JNEUROSCI.15-11-07644.1995

Krakauer JW, Ghez C, Ghilardi MF. 2005. Adaptation to Visuomotor Transformations:

Consolidation, Interference, and Forgetting. *J Neurosci* **25**:473–478.

doi:10.1523/JNEUROSCI.4218-04.2005

Krakauer JW, Hadjiosif AM, Xu J, Wong AL, Haith AM. 2019. Motor LearningComprehensive

Physiology. American Cancer Society. pp. 613–663. doi:10.1002/cphy.c170043

Krakauer JW, Pine ZM, Ghilardi M-F, Ghez C. 2000. Learning of Visuomotor Transformations

for Vectorial Planning of Reaching Trajectories. *J Neurosci* **20**:8916–8924.

doi:10.1523/JNEUROSCI.20-23-08916.2000

Krouchev NI, Kalaska JF. 2003. Context-Dependent Anticipation of Different Task Dynamics:

Rapid Recall of Appropriate Motor Skills Using Visual Cues. *Journal of Neurophysiology*

**89**:1165–1175. doi:10.1152/jn.00779.2002

Kruschke JK. 2008. Bayesian approaches to associative learning: From passive to active

learning. *Learning & Behavior* **36**:210–226. doi:10.3758/LB.36.3.210

Leow L-A, Marinovic W, Rugy A de, Carroll TJ. 2020. Task Errors Drive Memories That Improve

Sensorimotor Adaptation. *J Neurosci* **40**:3075–3088. doi:10.1523/JNEUROSCI.1506-

19.2020

Lilliefors HW. 1967. On the Kolmogorov-Smirnov Test for Normality with Mean and Variance

Unknown. *Journal of the American Statistical Association* **62**:399–402.

doi:10.1080/01621459.1967.10482916

Mackintosh NJ. 1976. Overshadowing and stimulus intensity. *Animal Learning & Behavior*

**4**:186–192. doi:10.3758/BF03214033

Marr D. 1969. A theory of cerebellar cortex. *The Journal of Physiology* **202**:437–470.

doi:10.1113/jphysiol.1969.sp008820

Mawase F, Shmuelof L, Bar-Haim S, Karniel A. 2014. Savings in locomotor adaptation

explained by changes in learning parameters following initial adaptation. *Journal of*

*Neurophysiology* **111**:1444–1454. doi:10.1152/jn.00734.2013

McDougle SD, Bond KM, Taylor JA. 2017. Implications of plan-based generalization in

sensorimotor adaptation. *Journal of Neurophysiology* **118**:383–393.

doi:10.1152/jn.00974.2016

McDougle SD, Bond KM, Taylor JA. 2015. Explicit and Implicit Processes Constitute the Fast

and Slow Processes of Sensorimotor Learning. *J Neurosci* **35**:9568–9579.

doi:10.1523/JNEUROSCI.5061-14.2015

McDougle SD, Ivry RB, Taylor JA. 2016. Taking Aim at the Cognitive Side of Learning in

Sensorimotor Adaptation Tasks. *Trends in Cognitive Sciences* **20**:535–544.

doi:10.1016/j.tics.2016.05.002

Morehead JR, Qasim SE, Crossley MJ, Ivry R. 2015. Savings upon Re-Aiming in Visuomotor

Adaptation. *J Neurosci* **35**:14386–14396. doi:10.1523/JNEUROSCI.1046-15.2015

Morehead JR, Taylor JA, Parvin DE, Ivry RB. 2017. Characteristics of Implicit Sensorimotor

Adaptation Revealed by Task-irrelevant Clamped Feedback. *Journal of Cognitive*

*Neuroscience* **29**:1061–1074. doi:10.1162/jocn_a_01108

Osu R, Hirai S, Yoshioka T, Kawato M. 2004. Random presentation enables subjects to adapt

to two opposing forces on the hand. *Nature Neuroscience* **7**:111–112.

doi:10.1038/nn1184

Patterson MM. 1970. Classical conditioning of the rabbit's (Oryctolagus cuniculus) nictitating

membrane response with fluctuation ISI and intracranial CS. *Journal of Comparative and*

*Physiological Psychology* **72**:193–202. doi:10.1037/h0029463

Pavlov IP. 1927. Conditioned reflexes, translated by GV Anrep. *London: Oxford*.

Poh E, Al-Fawakari N, Tam R, Taylor JA, McDougle SD. 2021. Generalization of motor learning

in psychological space. *bioRxiv* 2021.02.09.430542. doi:10.1101/2021.02.09.430542

Popa LS, Streng ML, Hewitt AL, Ebner TJ. 2016. The Errors of Our Ways: Understanding Error

Representations in Cerebellar-Dependent Motor Learning. *Cerebellum* **15**:93–103.

doi:10.1007/s12311-015-0685-5

Rescorla R, Wagner A. 1972. A theory of Pavlovian conditioning: Variations in the effectiveness

of reinforcement and nonreinforcement. *Classical conditioning II: Current research and*

*theory* **2**:64–69.

Rescorla RA. 1988. Pavlovian conditioning: It's not what you think it is. *American Psychologist*

**43**:151–160. doi:10.1037/0003-066X.43.3.151

Salomonczyk D, Cressman EK, Henriques DYP. 2011. Proprioceptive recalibration following

prolonged training and increasing distortions in visuomotor adaptation.

*Neuropsychologia* **49**:3053–3062. doi:10.1016/j.neuropsychologia.2011.07.006

Schlerf JE, Ivry RB, Diedrichsen J. 2012. Encoding of Sensory Prediction Errors in the Human

Cerebellum. *J Neurosci* **32**:4913–4922. doi:10.1523/JNEUROSCI.4504-11.2012

Schneiderman N, Gormezano I. 1964. Conditioning of the nictitating membrane of the rabbit as

a function of CS-US interval. *Journal of Comparative and Physiological Psychology*

**57**:188–195. doi:10.1037/h0043419

Schween R, Langsdorf L, Taylor JA, Hegele M. 2019. How different effectors and action effects

modulate the formation of separate motor memories. *Sci Rep* **9**:17040.

doi:10.1038/s41598-019-53543-1

Shadmehr R, Krakauer JW. 2008. A computational neuroanatomy for motor control. *Exp Brain*

*Res* **185**:359–381. doi:10.1007/s00221-008-1280-5

Sheahan HR, Franklin DW, Wolpert DM. 2016. Motor Planning, Not Execution, Separates Motor

Memories. *Neuron* **92**:773–779. doi:10.1016/j.neuron.2016.10.017

Sheahan HR, Ingram JN, Žalalytė GM, Wolpert DM. 2018. Imagery of movements immediately

following performance allows learning of motor skills that interfere. *Scientific Reports*

**8**:14330. doi:10.1038/s41598-018-32606-9

Shmuelof L, Huang VS, Haith AM, Delnicki RJ, Mazzoni P, Krakauer JW. 2012. Overcoming

Motor "Forgetting" Through Reinforcement Of Learned Actions. *J Neurosci* **32**:14617–

14621a. doi:10.1523/JNEUROSCI.2184-12.2012

Siegel S, Hearst E, George N. 1968. Generalization gradients obtained from individual subjects

following classical conditioning. *Journal of Experimental Psychology* **78**:171–174.

doi:10.1037/h0026178

Skinner BF. 1938. The behavior of organisms: an experimental analysis, The behavior of

organisms: an experimental analysis. Oxford, England: Appleton-Century.

Smith MA, Ghazizadeh A, Shadmehr R. 2006. Interacting Adaptive Processes with Different

Timescales Underlie Short-Term Motor Learning. *PLOS Biology* **4**:e179.

doi:10.1371/journal.pbio.0040179

Smith MC, Coleman SR, Gormezano I. 1969. Classical conditioning of the rabbit's nictitating

membrane response at backward, simultaneous, and forward CS-US intervals. *Journal*

*of Comparative and Physiological Psychology* **69**:226–231. doi:10.1037/h0028212

Taylor JA, Krakauer JW, Ivry RB. 2014. Explicit and Implicit Contributions to Learning in a

Sensorimotor Adaptation Task. *J Neurosci* **34**:3023–3032.

doi:10.1523/JNEUROSCI.3619-13.2014

Tsay JS, Avraham G, Kim HE, Parvin DE, Wang Z, Ivry RB. 2020a. The effect of visual

uncertainty on implicit motor adaptation. *Journal of Neurophysiology* **125**:12–22.

doi:10.1152/jn.00493.2020

Tsay JS, Ivry RB, Lee A, Avraham G. 2021. Moving outside the lab: The viability of conducting

sensorimotor learning studies online. *NBDT* **5**:1–22. doi:10.51628/001c.26985

Tsay JS, Kim HE, Haith AM, Ivry RB. 2022. Understanding implicit sensorimotor adaptation as a

process of proprioceptive re-alignment. doi:10.1101/2021.12.21.473747

Tsay JS, Lee AS, Avraham G, Parvin DE, Ho J, Boggess M, Woo R, Nakayama K, Ivry RB.

2020b. OnPoint: A package for online experiments in motor control and motor learning.

PsyArXiv. doi:10.31234/osf.io/hwmpy

Vandevoorde K, Orban de Xivry J-J. 2019. Internal model recalibration does not deteriorate with

age while motor adaptation does. *Neurobiology of Aging* **80**:138–153.

doi:10.1016/j.neurobiolaging.2019.03.020

Wang T, Avraham G, Tsay JS, Ivry RB. 2022. The Effect of Perturbation Variability on

Sensorimotor Adaptation Does Not Require an Implicit Memory of Errors.

doi:10.1101/2022.05.30.493844

Weiss SJ. 1972. Stimulus compounding in free-operant and classical conditioning: A review and

analysis. *Psychological Bulletin* **78**:189–208. doi:10.1037/h0032956

Wolpert DM, Flanagan JR. 2001. Motor prediction. *Current Biology* **11**:R729–R732.

doi:10.1016/S0960-9822(01)00432-8

Wolpert DM, Ghahramani Z. 2000. Computational principles of movement neuroscience. *Nature*

*Neuroscience* **3**:1212–1217. doi:10.1038/81497

Wolpert DM, Ghahramani Z, Jordan MI. 1995. An internal model for sensorimotor integration.

*Science* **269**:1880–1882. doi:10.1126/science.7569931

Wolpert DM, Miall RC, Kawato M. 1998. Internal models in the cerebellum. *Trends in Cognitive*

*Sciences* **2**:338–347. doi:10.1016/S1364-6613(98)01221-2

Yin C, Wei K. 2020. Savings in sensorimotor adaptation without an explicit strategy. *Journal of*

*Neurophysiology* **123**:1180–1192. doi:10.1152/jn.00524.2019

Zarahn E, Weston GD, Liang J, Mazzoni P, Krakauer JW. 2008. Explaining Savings for

Visuomotor Adaptation: Linear Time-Invariant State-Space Models Are Not Sufficient.

*Journal of Neurophysiology* **100**:2537–2548. doi:10.1152/jn.90529.2008

1278

 **FIGURE CAPTIONS**

 **Figure 1. Task design.**

 (A) Reaching task. Participants reach from a start location (white circle) to a target (blue dot).

 Online feedback is provided by a cursor (white dot). The target is displayed in a fixed location for

 the entire experiment (location varied across participants). After a random delay, a tone or a light

 (white frame) is presented, serving as the movement imperative. The participant is instructed to

 move directly to the target. The cue persists until the radial position of the hand reaches the target

 distance. The hand is not visible during the reach; instead, feedback is provided by a cursor. (B)

 Components and timing of the reaching task described as conditioning events. The tone and light

 imperatives serve as conditioned stimuli (CS) and the cursor feedback is the unconditioned

 stimulus (US). In our experiments, the radial position of the cursor is aligned with the hand, but

 the angular position is fixed ("clamped"), appearing either 15° away from the target (resulting in

 an error signal) when paired with one of the CS's (CS+) or at the target when paired with the other

 CS (CS-). After repeated pairings of the CS+ and US, a conditioned response (CR) develops,

 leading to an adaptive feedforward movement response in a direction opposite to the rotated

 feedback. Note that we include a UR in the schematic, what we assume is an on-line, automatic

 corrective response to the visual error. However, because participants are instructed to move fast,

 the UR is negligible.

 **Figure 2. Experiment 1: Differential Conditioning.**

 (A) Experimental protocol. During acquisition (white background), a 15° clamp (CW/CCW,

 counterbalanced across participants) was associated with CS+ (e.g., a tone) and a 0° clamp with

 CS- (e.g., a light; counterbalancing the associations with the tone and light across participants).

 During the probe phase (gray background), the CS+ and CS- were presented without feedback.

Throughout the entire experiment, CS+ and CS- trials were randomly interleaved. (B) Mean

heading angle (N=16) as a function of trial number. Clamped feedback was presented on all trials

during the acquisition phase (white background) and absent on all trials in the probe phase (gray

background) phases. Shaded region represents SEM. (C, D) Experimental results for trial-by-trial

change (Δ) in heading angle (mean ± SEM) during the acquisition (*C*) and probe (*D*) phases. Left

panels present analysis results for an adaptation effect (main effect of trial *n-1*, dark vs light blue)

and a Pavlovian effect (main effect of the presented CS on the current trial *n*, filled vs empty bars).

The black outlined bar and violin plot (right panel) present the Pavlovian effect, i.e., the subtraction

of heading angle changes between CS+ and CS- trials (mean and 95% CI). (E, F) Rescorla-

Wagner model simulation results during the acquisition (*E*) and probe (*F*) phases are consistent

with the experimental results. Dots and thin lines represent individual participants.

1314

1315

**Figure 3. Simulations of the Pavlovian effect due to differential conditioning in Experiment**

**1.**

Simulations of the current trial Pavlovian effect, the difference between the change in heading

angle on CS+ and CS- trials, for different parameter values as predicted by the Rescorla-Wagner

model (right side). The results for the acquisition phase are shown on top and probe phase on

bottom. Contours represent the mean ± SEM of the Pavlovian effect from Experiment 1. All

parameter values for the state-space model fail to predict a differential response on the current

trial for CS+ and CS- trials (right side).

1324

1325    **Figure 4. Model comparison.**

1326      The sum of squares residuals (SSR) and Akaike Information Criterion (AIC) difference between

1327      Rescorla-Wagner (RW) and the state-space (SS) models for each participant of Experiment 1.

1328

**Figure 5. Dynamics of adaptation and Pavlovian effects.**

(A) Time course of the mean weights (least squares regression $\beta$) of the adaptation (trial $n$-$1$,

grey) and Pavlovian (trial $n$, blue) effects in Experiment 1 as predicted by the Rescorla-Wagner

model (solid lines) and the state-space model (dotted lines). (B) Time course of the mean weights

of the adaptation and Pavlovian effects as derived from fits of the experimental results. Shaded

regions represent SEM.

1335

**Figure 6. Experiment 2: Pavlovian effect in differential conditioning is abolished when the**

**delay between the CS and US is long.**

(A) Illustration of trial events and their timing. By using a separate stimulus for the CS (tone or

light) and the imperative (target changed from gray to blue), a delay of ~1,000 was imposed

between the CS and US. The yellow background highlights the design difference from Experiment

1 (and Experiment 3 below). (B) Mean heading angle (N=64) as a function of trial number. Shaded

region represents SEM. (C, D) Trial-by-trial change in heading angle (mean ± SEM) during the

acquisition (*C*) and probe (*D*) phases. Left panels present analysis results for an adaptation effect

(main effect of trial $n$-$1$, dark vs light colors) and a Pavlovian effect (main effect of the presented

CS on the current trial $n$, filled vs empty bars). The black outlined bars and violin plots (right panel)

present the Pavlovian effect, i.e., the subtraction of heading angle changes between CS+ and

CS- trials (mean and 95% CI). Dots and thin lines represent individual participants.

1348

**Figure 7. Experiments 3: Pavlovian effect in differential conditioning is observed when the**

**CS and imperative cue occur simultaneously.**

(A) Illustration of trial events and their timing. The green background highlights the design

difference from Experiment 2, namely the simultaneous presentation of the CS and the movement

imperative. (B) Mean heading angle (N=64) as a function of trial number. Shaded region

represents SEM. (C, D) Trial-by-trial change in heading angle (mean ± SEM) during the

acquisition (*C*) and probe (*D*) phases. Left panels present analysis results for an adaptation effect

(main effect of trial *n-1*, dark vs light colors) and a Pavlovian effect (main effect of the presented

CS on the current trial *n*, filled vs empty bars). The black outlined bars and violin plots (right panel)

present the Pavlovian effect, i.e., the subtraction of heading angle changes between CS+ and

CS- trials (mean and 95% CI). Shaded regions and error bars represent SEM. Dots and thin lines

represent individual participants.

1361

**Figure 8. Experiment 4: Compound Conditioning.**

(A) During acquisition, a tone and light were presented simultaneously (Compound CS), serving

as the imperative signal for the reaching movement. They were paired with a 15° clamp on all

acquisition trials. During the probe phase, no feedback was provided, and the CSs were

presented either together (Compound CS) or alone (Single CS; tone or light). (B) Mean heading

angle (N=22) as a function of trial number. Clamped feedback was presented on all trials during

the acquisition phase (white background) and absent on all trials in the probe phase (gray

background) phases. Shaded region represents SEM. (C) Experimental results for the Δ heading

angle (mean ± SEM) during the probe phase, showing a positive relative change for the compound

CS on the current trial *n* (filled bar) and a negative relative change for each of its elements (empty

bars). (D) Predictions from the Rescorla-Wagner model for trial-by-trial Δ heading angle during

the probe phase. The two elements were assumed to have equal weight in the simulation. (E)

Scatter plot showing the between-participant trade-off in terms of the associative strength of the

two CSs (the dotted black line represents the unity line). (F) The Δ heading angle (mean ± SEM)

after pooling the two single-CS conditions to measure the effects of the previous CS type

(compound versus singleton, dark and light pink, respectively) and current CS (filled and empty

bars). Thin lines represent individual participants.